Predator-prey scaling laws support a suspension-feeding lifestyle in Cambrian
luolishaniid lobopodians

AUTHORS: Jared C. Richards[1], and Javier Ortega-Hernández[1]

AFFILIATIONS:

[1]Museum of Comparative Zoology and Department of Organismic and Evolutionary Biology,

Harvard University, 26 Oxford Street, Cambridge, MA 02138, USA

*Corresponding authors: jrichards@g.harvard.edu, jortegahernandez@fas.harvard.edu

ABSTRACT

The early Paleozoic saw a dramatic diversification of shelly epibenthic metazoans adapted to

suspension and filter feeding, but the extent to which these radiations affected the evolution of

non-biomineralized suspension-feeding taxa is uncertain because these organisms are not typically

well represented in the fossil record. Luolishaniids are a highly derived and disparate clade of

(typically) armoured lobopodians, widely interpreted as suspension feeders based on the presence

of five or six anterior pairs of setulose appendages. Luolishaniids are globally widespread and

represent the only Cambrian non-biomineralized free-living epibenthic bilaterians suggested to

have a suspension-feeding mode of life, but their proposed ecology relies solely on a qualitative

interpretation of their functional morphology. Here we test the hypothesis that the setulose

appendages of luolishaniids were adapted for a suspension-feeding function. Quantitative

morphological comparisons reveal a positive and statistically significant relationship between

body length and the mesh spacing of the setulose anterior limbs of luolishaniids. Standardized

comparisons indicate that the body size disparity between luolishaniids (predators) and Cambrian

mesoplankton (prey) is consistent with patterns observed in modern suspension-feeding

organisms. Our results provide quantitative evidence for suspension feeding in luolishaniids,

which represents the first statistically supported example of modern-like predator-prey scaling

patterns observed in Cambrian soft-bodied metazoans.

Keywords: Luolishaniidae, plankton, macroecology, niche ecology, functional morphology,

comparative zoology

INTRODUCTION

Suspension feeding is a widespread life strategy in which animals capture food particles such as

phytoplankton, zooplankton, and organic detritus from the water column in aquatic environments

[1–3]. Suspension feeding is found throughout diverse animal clades, and may be further

subcategorized based on precise habitat, mode of life, and locomotory capabilities of the organisms

[1]. Given that the feeding strategy of these organisms fundamentally involves extracting organic

matter from large volumes of water, they play a key ecological role as primary consumers in

aquatic food webs where they help cycle nutrients, transport biomass from small particles into

higher trophic levels in the water column, increase light penetration, and alter the distribution of

planktonic lifeforms in space and time [1,4–7].

Suspension feeding among the earliest animal-dominated communities is well represented

during the early Paleozoic among biomineralizing organisms with high preservation potential such

as sponges, brachiopods, echinoderms, and bivalved mollusks [8–13]. By contrast, our

understanding of the evolution of non-biomineralizing suspension-feeding organisms is more

precarious due to their comparatively low preservation potential in the conventional fossil record

[14–18] (Fig. 1). Radiodonts, typically large-bodied nektonic stem-group euarthropods typified by

the presence of arthropodized frontal appendages, represent the most geographically widespread

suspension-feeding soft-bodied metazoans found in sites with exceptional preservation around the

world during the early Paleozoic [19,20]. Suspension-feeding radiodonts include the hurdiids (Fig.

1C), which become increasingly diverse after the middle Cambrian and extended into the

Devonian [21–25,25,26] and the rarer tamisiocaridids (Fig. 1B), which are restricted to the early

Cambrian [27–29]. Other Cambrian soft-bodied organisms with a likely suspension-feeding mode

of life based on comparisons with modern analogues include cnidarians, hemichordates, tunicates,

vertebrates, ctenophores, and bivalved euarthropods [30–35] (Fig. 1D, 1E). Suspension feeding

has also been proposed for luolishaniids, a diverse group of lobopodians found in several major

Cambrian sites with exceptional preservation around the world [36–43] (Fig. 1A). The general

body plan of luolishaniids consists of a tubular trunk with five or six pairs of differentiated setulose

anterior appendages, followed by a variable number of clawed walking legs on the trunk (Fig. 1).

Other aspects of their morphology vary substantially, such as extent of spinose dorsal amour

ranging from absent (*e.g., Facivermis, Ovatiovermis*) to nearly baroque (*e.g., Collinsium,*

Collinsovermis, Acinocricus), as well as their body size (roughly 10 to 100 mm in length) and

lifestyle (*e.g.,* free living or tube dwelling) [39,42]. Luolishaniids differ from other suspension-

feeding organisms because most of them had a vagile epibenthic mode of life and lack obvious

extant analogues, with phylogenetic analyses suggesting affinities as either early relatives of velvet

worms (Onychophora), water bears (Tardigrada), or even panarthropods more broadly

[39,42,44,45]. However, the view of luolishaniids as suspension feeders is exclusively based on

the qualitative interpretation of the functional morphology of their anterior setulose appendages,

and thus this ecological hypothesis remains to be critically tested, especially considering the

peculiar morphology and variable lifestyle of these organisms. Other possible functional

interpretations for the prominent setulose anterior appendages of luolishaniids include grasping

macroscopic prey through raptorial-like feeding, or possibly to sieve the sediment in search for

food items as proposed for some radiodonts [46,47].

In this study we quantitatively test the hypothesis that the functional morphology of

Cambrian luolishaniids is congruent with a suspension-feeding mode of life by examining the

relationship between body size and sieving potential of the setulose anterior appendages. Based on

comparative statistical methods, we provide the first quantitative evidence for suspension feeding

in luolishaniids and demonstrate that these lobopodians follow the same suspension feeder to prey

size scaling laws observed in modern ecosystems.

Figure 1. Diversity of exceptionally preserved suspension feeding organisms from Cambrian
deposits. A) The luolishaniid lobopodian *Acinocricus stichus* from the middle Cambrian
(Wuluian) Spence Shale, Utah (Kansas University Museum Invertebrate Paleontology collection;
KUMIP 204353). **B)** Frontal appendage of the tamisiocaridid radiodont *Echidnacaris briggsi* [29]
from the early Cambrian (Stage 4) Emu Bay shale, South Australia (South Australian Museum,
Adelaide Paleontological collection; SAMA P40180). **C)** Carapace elements and frontal
appendage of the hurdiid radiodont *Pahvantia hasta* [23] from the middle Cambrian (Drumian)
Wheeler Shale, Utah (KUMIP 314089). **D)** The vertebrate *Nuucicthys rhynchocephalus* [34] from
the mid-Cambrian (Drumian) Marjum Formation, Utah (Utah Museum of Natural History;
UMNH.IP.6084). **E)** The tunicate *Megasiphon thylakos* [31] from the mid-Cambrian (Drumian)
Marjum Formation, Utah (UMNH.IP.6079). **F.** The hexactinellid-like sponge *Polygoniella turrelli*
[13] from the mid-Cambrian (Drumian) Marjum Formation, Utah (Museum of Comparative
Zoology Invertebrate Paleontology collection, Harvard University; MCZ.IP.199049).

MATERIAL AND METHODS

Data collection

High-resolution digital images from all previously described luolishaniid taxa were loaded into Adobe Illustrator 28.2 (2024) (see Table 1). We measured the spacing between the setae (mesh spacing) on the anterior limbs for each taxon, as this is generally considered a proxy for determining the size of seston collected by suspension feeders [27,48,49]. Given the paucity of complete specimens and the widespread likelihood of morphological distortion via diagenesis, we used the maximum recorded trunk length and width in the original description (Table 1). In cases where the specimen with the best preservation of the setulose structures is also complete, we would collect the body size measurement from the same specimens.

Data Analysis

All analysis was completed in RStudio environment using the packages dplyr, plyr, magrittr, stats, lm4, lmerTest, performance, MuMIn, lmtest, ggrepel, and ggplot2 [50–60].

Given the smaller sample size of the available data (*e.g.,* multiple mesh spacing values imposed onto only one trunk length measurement per species), we used a variety of analytical tools to extract as much information as possible from our limited dataset while being methodologically conservative. We employed an Ordinary Least-Squares (OLS) linear mixed-effect model with a gaussian response distribution and an identity link function with Satterthwaite's approximation to explore the relationship between genus-level body length and unaggregated mesh spacing. We treat genus-level body length (mm) as the fixed effect and unaggregated mesh spacing as the dependent variable (μm). To account for the nested structure of the data, we included genus as a random effect

in the model. This model was used to determine whether there was a statistically significant

relationship between luolishaniid genus body length and corresponding mesh size while also

controlling for non-independent observations [48,49,54,61]. Next, we performed an OLS linear

regression examining the relationship between mean (aggregated) mesh size values and body size.

This summarized view of the mesh spacing data would be used later for further examination of the

suspension-feeding ecology of each luolishaniid species. We report the F-statistic, the coefficient

of determination (R^2), and the p-value recovered from the t-test of the regression's slope coefficient

(H_0: regression slope = 0) for both OLS models. We also ran a Harvey-Collier test on the

aggregated mesh spacing dataset to check the assumption of a linear relationship between body

and mesh size on the logarithmic scale (H_0: regression is correctly modeled as linear). Finally,

using the relationship that exists between mesh spacing and estimated prey size for a variety of

extant metazoans [27] and the mean mesh spacing values for each luolishaniid genus, we

calculated the minimum and maximum estimated prey size (μm) of all known luolishaniids to

examine the lifestyles of this rare clade even further.

150   Neontological studies frequently use the equivalent spherical distance (ESD) to represent

the size of suspensivores and their prey [49,62,63]. This measure allows the size of irregularly

shaped objects, such as organisms, to be standardized into a 1-dimensional measure, in this case

the diameter of a perfect sphere. We modelled our luolishaniids as cylinders ($V = \pi r^2 h$), where h

and $2r$ represent the trunk length and width, respectively. This cylindrical volume was then

converted into ESD using the formula ESD = $(6r^2 h)^{1/3}$ [64]. The minimum estimated prey sizes

previously calculated for luolishaniids were used as a proxy for their corresponding prey ESD, as

this represents a more conservative value and has been shown to be more strongly correlated with

mesh spacing in extant metazoans [27]. To determine if the predator-prey size relationships of

luolishaniids were in line with those of known suspension-feeding clades, the modelled data

for Cambrian luolishaniids was appended onto previously compiled data on predator and prey

ESDs for modern suspension-feeding invertebrates collected from the literature [49,62,63].

Chaetognaths, a phylum of obligate ambush predators of similar size to luolishaniids that do not

engage in suspension feeding [65,66] were included in previous studies of predator-prey

relationships on the ESD scale in marine invertebrates. Given that they can be used as benchmark

for the predator prey size ratios seen in active, marine invertebrate hunters, we included them in

our study for comparison with luolishaniids and modern suspension feeders, but do not include

them in the OLS analysis.

We performed an OLS linear regression on the logarithmic scale to determine if the

relationships between predator and prey ESD between luolishaniids and modern suspension-

feeding invertebrates was similar. We report the F-statistic, the coefficient of determination (R^2),

and the *p*-value recovered from the t-test of the regression's slope coefficient (H_0: regression slope

= 0) of the OLS model. We also ran a Harvey-Collier test on the data to check that the linearity

assumption was met.

RESULTS

Luolishaniid Allometry and Prey Size Estimate

We find a positive and statistically significant relationship between body size (mm) and mesh

spacing (μm) on the logarithmic scale for both unaggregated mesh spacing data ($F(1, 4.82) = 8.52$,

p-value $= 3.45 \times 10^{-2}$, marginal $R^2 = 0.35$, conditional $R^2 = 0.65$) and aggregated mean mesh

spacing data ($F(1,5) = 7.72$, *p*-value$= 3.91 \times 10^{-2}$, $R^2 = 0.61$; S1_OLS; Fig. 2A). The corresponding

sublinear power functions are $Log_{10}(\hat{Y}) = 0.36 \times Log_{10}(X) + 1.72$ and $Log_{10}(\hat{Y}) = 0.37 \times Log_{10}(X)$

$+ 1.72$. The linearity assumption was met in the OLS of aggregated mesh spacing data on the

logarithmic scale (Harvey-Collier statistic = 0.36, degrees of freedom = 4, p-value = 0.74).

We find that the estimated minimum particle diameters would have enabled most

luolishaniids to capture suspended prey and particles on the lower end of the mesoplanktonic scale

($\sim$230 – 460 μm) (Fig. 2B; Table 1). *Luolishania* is the only taxon estimated to have been able to

collect suspended prey and particles that were on the microplanktonic spectrum ($\sim$150 μm) (Fig.

2B; Table 1). Given that the unnamed Emu Bay luolishaniid is based on a single incomplete

specimen that obstructs our ability to measure it [38], we used the previously derived power laws

to calculate its estimated maximum body size at approximately 87.5 mm (Table 1). However, due

to the incompleteness of the only known specimen we refrain from including it in our main

analyses to minimize potential sources of error.

Figure 2. Quantitative comparisons between Cambrian suspension feeding soft-bodied metazoans. A) Allometry of mesh size of Cambrian luolishaniids. Relationship between body length (mm) and appendage mesh spacing (μm) among Cambrian luolishaniids, showing a significant positive trend (red dashed line; $Log_{10}(\hat{Y}) = 0.37 \bullet Log_{10}(X) + 1.72$, $R^2 = 0.61$). Each point represents a species with a 95% CI around the mean mesh spacing. Silhouettes illustrate representative morphologies for each taxon; note that *Acinocricus* is represented by an isolated sclerite ring. B) Comparison between mesh spacing in luolishaniids (purple) and radiodonts (red) against estimated prey size. Radiodont data based on Vinther *et al.* 2014, Lerosey-Aubril *et al.* 2018 [23,27].

Table 1. Autecological metrics for all Cambrian luolishaniid lobopodians. Note that trunk length in the Emu Bay Shale luolishaniid (asterisk) is estimated from the statistically significant scaling pattern established amongst other Cambrian luolishaniids.						
Taxa	Number of Mesh Spacings (N)	Mean Mesh Spacing (μm)	Minimum Estimated Prey Diameter (μm)	Maximum Estimated Prey Diameter (μm)	Trunk Length (mm)	Trunk Width (mm)
Acinocricus stichus [41]	21	319.3	457.1	4248.5	96.0	13.2
Collinsium ciliosum [39]	52	250.6	358.0	3421.5	85.0	4.2
Collinsovermis monstruosus [41]	25	206.2	294.1	2875.1	35.0	7.6
Enthothyreos synnaustrus [43]	14	290.4	415.4	3902.8	38.12	5.2
Facivermis yunnanicus [42]	8	202.7	289.1	2831.3	90.0	2
Luolishania longicruris [37]	8	108.4	153.8	1619.4	14.30	1
Ovatiovermis cribratus [40]	26	161.1	229.3	2305.8	18.0	3.3
Collins' monster, Emu Bay [38]	12	272.0	389.0	3681.6	87.5*	-

Luolishaniid Macroecology and Modern Suspension feeders

We find a positive and statistically significant relationship between predator and prey ESD on the

logarithmic scale for analysis of luolishaniids and modern suspension-feeding invertebrate groups

($F(1,30) = 198.1$, p-value $= 9.40 \times 10^{-15}$, $R^2 = 0.87$). The corresponding sublinear power function is

218     $Log_{10}(\hat{Y}) = 0.90 \times Log_{10}(X) - 1.13$. (Fig. 3A). The linearity assumption on the logarithmic scale

219   was met in the OLS (Harvey-Collier statistic = 0.91, degrees of freedom = 29, p-value = 0.37),

220   respectively.

221   There is a considerable range in the predator-prey ESD ratios between clades (Figure 3B).

222   The group most similar in size to their prey are chaetognaths, which are on average five times

223   larger. Copepods and rotifers are approximately 20 times larger than their prey. The ratio amongst

224   luolishaniids, trochozoans, and cladocerans ranges approximately between 25‑50, indicating that

225   most suspension-feeding clades are between one to two orders of magnitude larger than their prey

226   on the ESD scale.

Table 2. Predator: Prey ESD Ratios by Clade. Luolishaniid body sizes calculated by modelling taxa as cylinders ($V = \pi r^2 h$) to obtain their volume and converting to ESD for standardization.

Clade	N	Mean Ratio	Median Ratio	log_{10}(Mean Ratio)	log_{10}(Median Ratio)
Chaetognatha	6	4.9	5.0	0.7	0.7
Cladocera	5	45.4	47.2	1.7	1.7
Copepoda	9	20.5	17.5	1.3	1.2
Luolishaniidae	7	36.1	29.0	1.5	1.5
Rotiferia	4	17.3	17.3	1.2	1.2
Trochozoa	7	39.9	24.4	1.4	1.4

Figure 3. Quantitative comparisons between predator-prey body size scaling. A) Relationship between predator and prey size in extant and extinct marine invertebrates. Plot of estimated spherical distance (μm) of marine invertebrates and their planktonic prey, showing a significant positive trend (red dashed line; $Log_{10}(\hat{Y}) = 0.90 \bullet Log_{10}(X) - 1.13$, $R^2 = 0.87$). Note that chaetognaths represent obligate raptorial hunters, so they are shown for comparison with the remaining suspension-feeding invertebrate groups and were not included in the linear regression line. B) Predator-Prey ESD Ratios for multiple marine invertebrate clades.

DISCUSSION

Luolishaniid Macroecological Scaling and Estimated Prey Size

The positive, statistically significant relationship between trunk length and mesh spacing in the

setulose appendages of Cambrian luolishaniids (Figure 2A) on the logarithmic scale indicates that

mesh spacing does not scale uniformly with body size. For a given unit of increase in body size,

there is a proportionally smaller increase in mesh spacing. These observations recapitulate

sublinear allometric laws of body size in extant animals [67], particularly among suspension-

feeding aquatic invertebrates [48,68,69]. The strong (R^2 = 0.87, *p*-value < 0.001) relationship

between predator and prey ESD on the logarithmic scale (Figure 3A), provides quantitative support

to the hypothesis that Cambrian luolishaniids were suspension feeders. The setulose appendages

of luolishaniids would enable them to collect seston approximately 30−35 times smaller than their

body size (when expressed in terms of ESD scaling), mirroring the predator-prey size relationship

seen in predominantly filter-feeding groups such as larval mollusks and cladocerans [70–73]

(Figure 3B). In contrast, obligate small invertebrate raptorial predators such as chaetognaths use

their elongate anterior spines to actively ambush and seize relatively large prey when compared to

their own body size [65,66].

In the context of luolishaniid feeding ecology, mesh size would have had direct

implications for the planktonic prey targeted by the group. It is impossible to evaluate the precise

composition of luolishaniid diets because all their fossils to date lack evidence of gut contents or

other indications of food selectivity; thus, the mesh size provides the best available estimate of

their prey items. We find that the smallest Cambrian luolishaniids had the finest mesh size,

estimated at 108.4 *μ*m for *Luolishania* (body length = 14 mm) and 161.1 *μ*m for *Ovatiovermis*

(body length = 18 mm) (Figure 2A, Table 1), which would have allowed them to capture larger

microplankton (*e.g.*, phytoplankton), as well as the smaller cohorts of mesoplankton. All other

larger-bodied luolishaniids, ranging from *Collinsovermis* (body length = 35 mm) to *Acinocricus*

(body length = 96 mm), show a progressive increase in the mesh size on the setulose appendages,

ranging from ca. 200 μm to 450 μm. This mesh size range would have allowed medium to large

bodied luolishaniids to feed on all types of mesoplankton but would be less suitable for targeting

microplankton as it would fall below the minimum capturable size (Figure 2B). More broadly, the

range of estimated prey sizes of luolishaniids overlaps with that of suspension-feeding radiodonts

such as *Aegirocassis benmoulai, Pahvantia hastata*, and *Tamisiocaris borealis* [20,23,27]. These

comparisons further support the hypothesis of luolishaniids as suspension feeders, as both groups

follow similar patterns of mesh spacing on their feeding appendages despite the substantial

disparity in overall body size between the smaller luolishaniids (Table 1) and the typically much

larger radiodonts (*A. benmoulai*, ca. 200 cm trunk length; *P. hastana,* ca. 24 cm estimated trunk

length; *T. borealis,* ca. 34 cm estimated trunk length) [20,23,27]. These observations suggest that

mesoplankton and larger microplankton were most likely the preferred food sources of multiple

suspension-feeding clades during the Cambrian and the Early Ordovician. Critically, luolishaniids

provide a valuable and previously unavailable insight into the relationships between body size and

prey size (estimated by mesh size) of Cambrian non-biomineralized metazoans because of the

availability of complete body fossils in several members of the same clade. By contrast, this

information is absent from suspension-feeding radiodonts due to the paucity of complete body

fossils in the Cambrian record.

The microfossil record of planktonic life during the early Paleozoic offers insights into the

likely prey items that would have served as the primary food source of luolishaniids. Although the

diversity and abundance of phytoplankton (*e.g.*, cyanobacteria, acritarchs) increased rapidly

throughout the Cambrian, these microorganisms typically have body sizes below 100 μm in

diameter [74–78]. This range indicates that Cambrian phytoplankton would likely be too small for

most luolishaniids based on their mesh spacing; only the smallest luolishaniids, namely

Luolishania and *Ovatiovermis*, would have been able to feed off the largest phytoplankton (Figure

2B). Most luolishaniids would have been instead well suited for feeding on mesoplankton (ca. 0.2

to 2 mm diameter), which would include diverse organisms such as chitinozoans, radiolarians, and

various types of metazoan larvae (i.e. zooplankton). Fossil evidence for chitinozoans dates to mid-

Cambrian strata [79], although they are more common from the Ordovician to Devonian [80,81].

Chitinozoans likely extended into the early Cambrian, but it is possible that they were less

abundant and thus did not produce a robust fossil record at the time. Thus, we posit that

zooplankton likely represented the primary prey source for most Cambrian luolishaniids, which

would have included the swimming larvae of cnidarians, arthropods, and spiralians such as

brachiopods and mollusks [9,82–95].

Comparisons to Modern Analogs

Luolishaniid lobopodians are unusual among early metazoans because they are largely soft-bodied

(except for their sclerotized dorsal spines), suspension-feeding, epibenthic, vagile, and typically

free living (with the possible exception of *Facivermis* [42]), and thus occupied a specialized

ecological niche in Cambrian ecosystems. It seems plausible that other Cambrian metazoans also

fit some of these attributes, particularly when considering the substantial diversity of filter-feeding

apparatuses recovered from small carbonaceous fossils around the world [96–98]. For example,

the early Cambrian annelids *Dannychaeta* [99] and *Iotuba* [100] are likely candidates of additional

soft-bodied suspension-feeding metazoans based on comparisons with some living megaloniids

and acrociriid polychaetes. However, further discoveries of complete organisms with well-

preserved feeding structures are needed to confirm this possibility. Luolishaniids have been

directly compared with some extant suspension-feeding invertebrates, particularly caprellid

amphipod crustaceans, based on suggested similarities in their morphology including an elongate

tubular body, anterior setulose appendages with a sieving function, and clawed walking legs on

body trunk or posterior for grasping onto the substrate [40,41,101–103]. However, these taxa differ

in important aspects of their functional morphology with direct implications for their precise mode

of life and feeding strategies. Caprellid amphipods employ their cilia-laden, mucus-covered

antennae to initiate suspension feeding [40,101]. After detecting a substantial accumulation of

seston on the antennae, caprellids use their raptorial gnathopods to cleanse these appendages from

food particles, form a ball of mucus and seston, and pass it to their mouth for consumption [101].

Although luolishaniids and caprellid amphipods employ setulose anterior appendages to

capture plankton, the former differs in the absence of specialized cleaning limbs or complex

mouthparts employed in suspension feeding [39,40,42,101,102,104]. These differences suggest

that luolishaniids most likely employed a different strategy to consume seston, as there is no

evidence whatsoever that these luolishaniids could secrete or employ mucus for feeding. Instead,

suction feeding would likely be a suitable option as it is widely employed by aquatic invertebrates

where a negative pressure vacuum is produced to draw nearby water, and therefore any suspended

materials, into the buccal cavity [105,106]. This interpretation is in line with the recent hypothesis

that armored lobopodians were adapted for suction feeding based on the presence of a muscular

pharynx (e.g. *Onychodictyon ferox*) and aciculate sclerotized teeth in the foregut with a filtering

function in the hallucigeniid *Hallucgenia sparsa* and the luolishaniid *Ovatiovermis cribatus* [107].

Experimental work on the fluid dynamics of extant suction feeders has shown that no modern

invertebrates with hydrostatically supported bodies can create the same suction forces seen in

aquatic vertebrates such as fish [108]. Therefore, we hypothesize that the suction feeding capacity

of luolishaniids would have been comparatively low, just enough so that a low-pressure difference

between the buccal cavity and the environment would suffice to consume the prey captured by the

sieving apparatus on their anterior setulose appendages.

Broader Paleoecological Implications

Luolishaniids differ from other Cambrian suspension-feeding soft-bodied animals, particularly

radiodonts such as *Tamisiocaris*, *Pahvantia,* or *Hurdia*, in that the former had a slow-moving

epibenthic habitus, which means they would not be able to force water over their sieving apparatus

by simply continuously moving against the water column [20,27]. Instead, luolishaniids most

likely sought after microhabitats with low to intermediate water flow rates facing their anterior

setulose limbs perpendicular to the oncoming water current to collect suspended food items

[1,109–111]. The water flow speeds necessary for stationary suspension feeding of mesoplankton

in modern ecosystems range from 5 to 10 cm/s based on experiments involving modern suspension

feeders feeding on mesoplankton [110,112]. Assuming that luolishaniids would require a similar

flow speed necessary for effective suspension feeding, the role of the robust sclerotized terminal

claws on the posterior trunk appendages becomes critical, as it would allow these luolishaniids to

maintain a secure attachment to hard substrates during food capture without becoming dislodged

[1,39,110,112,113].

The feeding ecology of luolishaniids might also have implications for better understanding

their mode of life considering their peculiar morphology, particularly the presence of extensive

dorsal spinose defensive armature in most species [39,41,43]. Neontological behavioural work has

shown that free living animals that bear some type of external body armor (*e.g.*, epidermal spines,

plates or scales) display a higher threat tolerance or threshold before they flee or take refuge from

potential predators when compared to unarmored forms, both in aquatic and terrestrial

environments [114,115]. Applied in the context of luolishaniid paleobiology, this generalized

behavior would suggest that smaller-bodied species and/or those in which dorsal armor is relatively

light (*Luolishania*) or non-existent (*Ovatiovermis*) would be more hasty to seek refuge compared

to larger bodied and more heavily armored forms, such as *Acinocricus, Collinsium,*

Collinsovermis, and *Entothyreos* [38,39,43]. *Facivermis* is a notable exception in this regard

because it is large-bodied (ca. 90 mm long) but completely lacks dorsal armor [42]. In this case,

the unique tube-dwelling mode of life of *Facivermis* most likely represents a tradeoff that allowed

it to attain a large body size despite the absence of dorsal armor by shifting to a different defensive

strategy that instead relies on the protection of the secreted tube, facilitating a readily accessible

refuge if disturbed. It is notable that the extent of dorsal armature in luolishaniids closely follows

their overall body size, with *Acinocricus, Collinsium, Collinsovermis, Entothyreos* featuring the

heaviest coverage within the clade based on the number of dorsal spines per appendage pair. The

heavy armature of these forms would have likely made their movements cumbersome due to

excessive resistance against the water column and thus caused a tradeoff against their ability to

swiftly escape from potential predators, as also observed in modern species. Heavily armored

luolishaniids would have primarily relied on their body size and impressive arrays of dorsal spines

as primary deterrents, particularly when actively engaging in suspension feeding against the water

current [114–117].

In conclusion, our results provide novel quantitative support to the longstanding

interpretation of luolishaniids as epibenthic suspension feeders and suggest that they follow

comparable macroecologial patterns observed in modern invertebrate clades with a suspension

feeding autecology. The findings indicate that the fundamental requirements for a suspension-

feeding lifestyle among non-biomineralized metazoans were already in place by the early and

middle Cambrian based on a comparative quantification of luolishaniid morphology. This study

also provides the first estimates of planktonic prey sizes for luolishaniids and provides further

novel insights into their likely behavior based on modern analogues. Luolishaniids are among the

most eldritch organisms that evolved during the Cambrian Explosion, hence their initial moniker

as "Collin's monsters", honoring Desmond Collins who illustrated the first example of these

bizarre organisms in 1986 from a fossil collection produced by the Royal Ontario Museum field

party in 1983 [38,41]. Instead, we posit that main adaptations and mode of life of luolishaniids can

be readily explained through the lens of a modern marine suspension-feeding ecology.

ACKNOWLEDGEMENTS

We thank Rudy Lerosey-Aubril (Harvard University) for sharing digital images of *Pahvantia hastata*, and John Paterson (University of New England) for sharing digital images of *Echidnacaris briggsi*. Thanks to Bruce Lieberman (The University of Kansas) for facilitating access to the holotype of *Acinocricus stichus*. This work is funded by NSF CAREER award no. 2047192 "Ecological Turnover at the Dawn of the Great Ordovician Biodiversification Event— Quantifying the Cambro-Ordovician Transition through the Lens of Exceptional Preservation," and NSF GRFP award no. 2140743.

References

1. Hamann L, Blanke A. 2022 Suspension feeders: diversity, principles of particle separation
and biomimetic potential. *J. R. Soc. Interface.* **19**, 20210741. (doi:10.1098/rsif.2021.0741)

2. Hentschel BT, Shimeta J. 2008 Suspension Feeders. In *Encyclopedia of Ecology*, pp. 3437–
3442. Elsevier. (doi:10.1016/B978-008045405-4.00548-6)

3. Jørgensen CB. 1966 Biology of suspension feeding. *(No Title)*

4. Riisgård H, Larsen P. 2010 Particle capture mechanisms in suspension-feeding invertebrates.
Mar. Ecol. Prog. Ser. **418**, 255–293. (doi:10.3354/meps08755)

5. Ostroumov SA. 2005 Some aspects of water filtering activity of filter-feeders. *Hydrobiologia*
542, 275–286. (doi:10.1007/s10750-004-1875-1)

6. Gili J-M, Coma R. 1998 Benthic suspension feeders: their paramount role in littoral marine
food webs. *Trends in Ecology & Evolution* **13**, 316–321. (doi:10.1016/S0169-
5347(98)01365-2)

7. Lenton TM, Boyle RA, Poulton SW, Shields-Zhou GA, Butterfield NJ. 2014 Co-evolution of
eukaryotes and ocean oxygenation in the Neoproterozoic era. *Nature Geosci* **7**, 257–265.
(doi:10.1038/ngeo2108)

8. Bottjer DJ, Ausich WI. 1986 Phanerozoic development of tiering in soft substrata
suspension-feeding communities. *Paleobiology* **12**, 400–420.
(doi:10.1017/S0094837300003134)

9. Nützel A, Lehnert O, Frýda J. 2006 Origin of planktotrophy—evidence from early molluscs.
Evolution and Development **8**, 325–330. (doi:10.1111/j.1525-142X.2006.00105.x)

10. Zamora S, Deline B, Javier Álvaro J, Rahman IA. 2017 The Cambrian Substrate Revolution
and the early evolution of attachment in suspension-feeding echinoderms. *Earth-Science
Reviews* **171**, 478–491. (doi:10.1016/j.earscirev.2017.06.018)

11. Hsieh S, Bush AM, Bennington JB. 2019 Were bivalves ecologically dominant over
brachiopods in the late Paleozoic? A test using exceptionally preserved fossil assemblages.
Paleobiology **45**, 265–279. (doi:10.1017/pab.2019.3)

12. Ausich WI, Bottjer DJ. 2020 Echinoderm role in the history of Phanerozoic tiering in
suspension-feeding communities. In *Echinodermata* (eds BF Keegan, BDS O'Connor), pp.
3–11. CRC Press. (doi:10.1201/9781003079224-2)

13. Del Mouro L, Lerosey-Aubril R, Botting J, Coleman R, Gaines RR, Skabelund J, Weaver JC,
Ortega-Hernández J. 2024 A new sponge from the Marjum Formation of Utah documents the
Cambrian origin of the hexactinellid body plan. *R. Soc. Open Sci.* **11**, 231845.
(doi:10.1098/rsos.231845)

14. Orr PJ, Benton MJ, Briggs DEG. 2003 Post-Cambrian closure of the deep-water slope-basin
taphonomic window. *Geol* **31**, 769. (doi:10.1130/G19193.1)

15. Gaines RR, Hammarlund EU, Hou X, Qi C, Gabbott SE, Zhao Y, Peng J, Canfield DE. 2012
Mechanism for Burgess Shale-type preservation. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 5180–
5184. (doi:10.1073/pnas.1111784109)

16. Gaines RR. 2014 Burgess Shale-type Preservation and its Distribution in Space and Time.
Paleontol. Soc. pap. **20**, 123–146. (doi:10.1017/S1089332600002837)

17. Lefebvre B, Lerosey-Aubril R, Servais T, Van Roy P. 2016 The Fezouata Biota: An
exceptional window on the Cambro-Ordovician faunal transition. *Palaeogeography,*
Palaeoclimatology, Palaeoecology **460**, 1–6. (doi:10.1016/j.palaeo.2016.06.041)

18. Saleh F *et al.* 2021 A novel tool to untangle the ecology and fossil preservation knot in
exceptionally preserved biotas. *Earth and Planetary Science Letters* **569**, 117061.
(doi:10.1016/j.epsl.2021.117061)

19. Potin GJ-M, Daley AC. 2023 The significance of Anomalocaris and other Radiodonta for
understanding paleoecology and evolution during the Cambrian explosion. *Frontiers in*
Earth Science **11**, 1160285.

20. Potin GJ-M, Gueriau P, Daley AC. 2023 Radiodont frontal appendages from the Fezouata
Biota (Morocco) reveal high diversity and ecological adaptations to suspension-feeding
during the Early Ordovician. *Front. Ecol. Evol.* **11**, 1214109.
(doi:10.3389/fevo.2023.1214109)

21. Kühl G, Briggs DEG, Rust J. 2009 A Great-Appendage Arthropod with a Radial Mouth from
the Lower Devonian Hunsrück Slate, Germany. *Science* **323**, 771–773.
(doi:10.1126/science.1166586)

22. Van Roy P, Daley AC, Briggs DEG. 2015 Anomalocaridid trunk limb homology revealed by
a giant filter-feeder with paired flaps. *Nature* **522**, 77–80. (doi:10.1038/nature14256)

23. Lerosey-Aubril R, Pates S. 2018 New suspension-feeding radiodont suggests evolution of
microplanktivory in Cambrian macronekton. *Nat Commun* **9**, 3774. (doi:10.1038/s41467-
018-06229-7)

24. Liu Y, Lerosey-Aubril R, Audo D, Zhai D, Mai H, Ortega-Hernández J. 2020 Occurrence of
the eudemersal radiodont *Cambroraster* in the early Cambrian Chengjiang Lagerstätte and
the diversity of hurdiid ecomorphotypes. *Geol. Mag.* **157**, 1200–1206.
(doi:10.1017/S0016756820000187)

25. Pates S, Botting JP, McCobb LME, Muir LA. 2020 A miniature Ordovician hurdiid from
Wales demonstrates the adaptability of Radiodonta. *R. Soc. open sci.* **7**, 200459.
(doi:10.1098/rsos.200459)

26. Sun Z, Zeng H, Zhao F. 2020 A new middle Cambrian radiodont from North China:
Implications for morphological disparity and spatial distribution of hurdiids.
Palaeogeography, Palaeoclimatology, Palaeoecology **558**, 109947.
(doi:10.1016/j.palaeo.2020.109947)

27. Vinther J, Stein M, Longrich NR, Harper DAT. 2014 A suspension-feeding anomalocarid
from the Early Cambrian. *Nature* **507**, 496–499. (doi:10.1038/nature13010)

28. Wu Y, Fu D, Ma J, Lin W, Sun A, Zhang X. 2021 Houcaris gen. nov. from the early
Cambrian (Stage 3) Chengjiang Lagerstätte expanded the palaeogeographical distribution of
tamisiocaridids (Panarthropoda: Radiodonta). *PalZ* **95**, 209–221. (doi:10.1007/s12542-020-
00545-4)

29. Paterson JR, García-Bellido DC, Edgecombe GD. 2023 The early Cambrian Emu Bay Shale
radiodonts revisited: morphology and systematics. *Journal of Systematic Palaeontology* **21**,
2225066. (doi:10.1080/14772019.2023.2225066)

30. Nanglu K, Caron J-B, Conway Morris S, Cameron CB. 2016 Cambrian suspension-feeding
tubicolous hemichordates. *BMC Biol* **14**, 56. (doi:10.1186/s12915-016-0271-4)

31. Nanglu K, Lerosey-Aubril R, Weaver JC, Ortega-Hernández J. 2023 A mid-Cambrian
tunicate and the deep origin of the ascidiacean body plan. *Nat Commun* **14**, 3832.
(doi:10.1038/s41467-023-39012-4)

32. Zhao Y, Vinther J, Parry LA, Wei F, Green E, Pisani D, Hou X, Edgecombe GD, Cong P.
2019 Cambrian Sessile, Suspension Feeding Stem-Group Ctenophores and Evolution of the
Comb Jelly Body Plan. *Current Biology* **29**, 1112-1125.e2. (doi:10.1016/j.cub.2019.02.036)

33. Izquierdo-López A, Caron J-B. 2024 The Cambrian *Odaraia alata* and the colonization of
nektonic suspension-feeding niches by early mandibulates. *Proc. R. Soc. B.* **291**, 20240622.
(doi:10.1098/rspb.2024.0622)

34. Lerosey-Aubril R, Ortega-Hernández J. 2024 A long-headed Cambrian soft-bodied vertebrate
from the American Great Basin region. *R. Soc. Open Sci.* **11**, 240350.
(doi:10.1098/rsos.240350)

35. Ou Q, Han J, Zhang Z, Shu D, Sun G, Mayer G. 2017 Three Cambrian fossils assembled into
an extinct body plan of cnidarian affinity. *Proceedings of the National Academy of Sciences*
114, 8835–8840.

36. Hou X, Bergström J. 1995 Cambrian lobopodians-ancestors of extant onychophorans?
Zoological Journal of the Linnean Society **114**, 3–19. (doi:10.1111/j.1096-
3642.1995.tb00110.x)

37. Ma X, Hou X, Bergström J. 2009 Morphology of Luolishania longicruris (Lower Cambrian,
Chengjiang Lagerstätte, SW China) and the phylogenetic relationships within lobopodians.
Arthropod Structure & Development **38**, 271–291. (doi:10.1016/j.asd.2009.03.001)

38. García-bellido DC, Edgecombe GD, Paterson JR, Ma X. 2013 A 'Collins' monster'-type
lobopodian from the Emu Bay Shale Konservat-Lagerstätte (Cambrian), South Australia.
Alcheringa: An Australasian Journal of Palaeontology **37**, 474–478.
(doi:10.1080/03115518.2013.792456)

39. Yang J, Ortega-Hernández J, Gerber S, Butterfield NJ, Hou J, Lan T, Zhang X. 2015 A
superarmored lobopodian from the Cambrian of China and early disparity in the evolution of
Onychophora. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 8678–8683.
(doi:10.1073/pnas.1505596112)

40. Caron J-B, Aria C. 2017 Cambrian suspension-feeding lobopodians and the early radiation of
panarthropods. *BMC Evol Biol* **17**, 29. (doi:10.1186/s12862-016-0858-y)

41. Caron J, Aria C. 2020 The Collins' monster, a spinous suspension-feeding lobopodian from
the Cambrian Burgess Shale of British Columbia. *Palaeontology* **63**, 979–994.
(doi:10.1111/pala.12499)

42. Howard RJ, Hou X, Edgecombe GD, Salge T, Shi X, Ma X. 2020 A Tube-Dwelling Early
Cambrian Lobopodian. *Current Biology* **30**, 1529-1536.e2. (doi:10.1016/j.cub.2020.01.075)

43. Aria C, Caron J-B. 2024 Deep origin of articulation strategies in panarthropods: evidence
from a new luolishaniid lobopodian (Panarthropoda) from the Tulip Beds, Burgess Shale.
Journal of Systematic Palaeontology **22**, 2356090. (doi:10.1080/14772019.2024.2356090)

44. Smith MR, Ortega-Hernández J. 2014 Hallucigenia's onychophoran-like claws and the case
for Tactopoda. *Nature* **514**, 363–366. (doi:10.1038/nature13576)

45. Kihm J-H, Smith FW, Kim S, Rho HS, Zhang X, Liu J, Park T-YS. 2023 Cambrian
lobopodians shed light on the origin of the tardigrade body plan. *Proc. Natl. Acad. Sci.
U.S.A.* **120**, e2211251120. (doi:10.1073/pnas.2211251120)

46. Liu Y, Lerosey-Aubril R, Audo D, Zhai D, Mai H, Ortega-Hernández J. 2020 Occurrence of
the eudemersal radiodont *Cambroraster* in the early Cambrian Chengjiang Lagerstätte and
the diversity of hurdiid ecomorphotypes. *Geol. Mag.* **157**, 1200–1206.
(doi:10.1017/S0016756820000187)

47. Moysiuk J, Caron J-B. 2019 A new hurdiid radiodont from the Burgess Shale evinces the
exploitation of Cambrian infaunal food sources. *Proc. R. Soc. B.* **286**, 20191079.
(doi:10.1098/rspb.2019.1079)

48. Burns CW. 1968 The Relationship Between Body Size of Filter-Feeding Cladocera and the
Maximum Size of Partcile Ingested. *Limnology & Oceanography* **13**, 675–678.
(doi:10.4319/lo.1968.13.4.0675)

49. Wirtz K. 2012 Who is eating whom? Morphology and feeding type determine the size
relation between planktonic predators and their ideal prey. *Mar. Ecol. Prog. Ser.* **445**, 1–12.
(doi:10.3354/meps09502)

50. R Core Team. 2022 *R: A Language and Environment for Statistical Computing*. Vienna,
Austria: R Foundation for Statistical Computing. See https://www.R-project.org/.

51. Wickham H, François R, Henry L, Müller K, Vaughan D. 2023 dplyr: A Grammar of Data
Manipulation. R package version 1.1. 2.

52. Wickham H. 2009 *ggplot2: Elegant Graphics for Data Analysis*. New York, NY: Springer
New York. (doi:10.1007/978-0-387-98141-3)

53. Wickham H, Vaughan D, Girlich M. 2023 Tidyr: Tidy messy data [Computer software].

54. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models Using
lme4. *J. Stat. Soft.* **67**. (doi:10.18637/jss.v067.i01)

55. Kuznetsova A, Brockhoff PB, Christensen RHB. 2017 lmerTest Package: Tests in Linear
Mixed Effects Models. *J. Stat. Soft.* **82**. (doi:10.18637/jss.v082.i13)

56. Slowikowski K. 2016 ggrepel: Automatically Position Non-Overlapping Text Labels with
'ggplot2'. , 0.9.6. (doi:10.32614/CRAN.package.ggrepel)

57. Bartoń K. 2010 MuMIn: Multi-Model Inference. , 1.48.11.
(doi:10.32614/CRAN.package.MuMIn)

58. Lüdecke D, Ben-Shachar M, Patil I, Waggoner P, Makowski D. 2021 performance: An R
Package for Assessment, Comparison and Testing of Statistical Models. *JOSS* **6**, 3139.
(doi:10.21105/joss.03139)

59. Zeileis A, Hothorn T, others. 2002 Diagnostic checking in regression relationships.

60. Bache SM, Wickham H. 2025 *magrittr: A Forward-Pipe Operator for R*. See
https://magrittr.tidyverse.org.

61. Burton AL. 2021 OLS (Linear) Regression. In *The Encyclopedia of Research Methods in
Criminology and Criminal Justice* (eds JC Barnes, DR Forde), pp. 509–514. Wiley.
(doi:10.1002/9781119111931.ch104)

62. Fuchs H, Franks P. 2010 Plankton community properties determined by nutrients and size-
selective feeding. *Mar. Ecol. Prog. Ser.* **413**, 1–15. (doi:10.3354/meps08716)

63. Hansen B, Bjornsen PK, Hansen PJ. 1994 The size ratio between planktonic predators and
their prey. *Limnology & Oceanography* **39**, 395–403. (doi:10.4319/lo.1994.39.2.0395)

64. 1988 Particle size measurement: the equivalent spherical diameter. *Proc. R. Soc. Lond. A*
419, 137–149. (doi:10.1098/rspa.1988.0100)

65. Saito H. 2001 Feeding rates in the chaetognath Sagitta elegans: effects of prey size, prey
swimming behaviour and small-scale turbulence. *Journal of Plankton Research* **23**, 1385–
1398. (doi:10.1093/plankt/23.12.1385)

66. Stuart V, Verheye HM. 1991 Diel migration and feeding patterns of the chaetognath, *Sagitta friderici*, off the west coast of South Africa. *J Mar Res* **49**, 493–515. (doi:10.1357/002224091784995819)

67. Yu Y, Karbowski J, Sachdev RN, Feng J. 2014 Effect of temperature and glia in brain size enlargement and origin of allometric body-brain size scaling in vertebrates. *BMC Evol Biol* **14**, 178. (doi:10.1186/s12862-014-0178-z)

68. Burns CW. 1969 RELATION BETWEEN FILTERING RATE, TEMPERATURE, AND BODY SIZE IN FOUR SPECIES OF DAPHNIA. *Limnology & Oceanography* **14**, 693–700. (doi:10.4319/lo.1969.14.5.0693)

69. Sebens KP. 1981 THE ALLOMETRY OF FEEDING, ENERGETICS, AND BODY SIZE IN THREE SEA ANEMONE SPECIES. *The Biological Bulletin* **161**, 152–171. (doi:10.2307/1541115)

70. Ebert D. 2022 Daphnia as a versatile model system in ecology and evolution. *EvoDevo* **13**, 16. (doi:10.1186/s13227-022-00199-0)

71. Reinfelder JR, Fisher NS. 1994 The assimilation of elements ingested by marine planktonic bivalve larvae. *Limnology & Oceanography* **39**, 12–20. (doi:10.4319/lo.1994.39.1.0012)

72. Raby D, Mingelbier M, Dodson JJ, Klein B, Lagadeuc Y, Legendre L. 1997 Food-particle size and selection by bivalve larvae in a temperate embayment. *Marine Biology* **127**, 665–672. (doi:10.1007/s002270050057)

73. Gallager SM. 1988 Visual observations of particle manipulation during feeding in larvae of a bivalve mollusc. *Bulletin of marine Science* **43**, 344–365.

74. Rigby S, Milsom CV. 2000 Origins, Evolution, and Diversification of Zooplankton. *Annu. Rev. Ecol. Syst.* **31**, 293–313. (doi:10.1146/annurev.ecolsys.31.1.293)

75. Huntley J, Xiao S, Kowalewski M. 2006 1.3 Billion years of acritarch history: An empirical morphospace approach. *Precambrian Research* **144**, 52–68. (doi:10.1016/j.precamres.2005.11.003)

76. Cohen PA, Knoll AH, Kodner RB. 2009 Large spinose microfossils in Ediacaran rocks as resting stages of early animals. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 6519–6524. (doi:10.1073/pnas.0902322106)

77. Harvey THP. 2023 Colonial green algae in the Cambrian plankton. *Proc. R. Soc. B.* **290**, 20231882. (doi:10.1098/rspb.2023.1882)

78. Demoulin CF, Lara YJ, Cornet L, François C, Baurain D, Wilmotte A, Javaux EJ. 2019 Cyanobacteria evolution: Insight from the fossil record. *Free Radical Biology and Medicine* **140**, 206–223. (doi:10.1016/j.freeradbiomed.2019.05.007)

79. Shen C, Aldridge RJ, Williams M, Vandenbroucke TRA, Zhang X -g. 2013 Earliest chitinozoans discovered in the Cambrian Duyun fauna of China. *Geology* **41**, 191–194. (doi:10.1130/G33763.1)

80. Achab A, Paris F. 2007 The Ordovician chitinozoan biodiversification and its leading factors. *Palaeogeography, Palaeoclimatology, Palaeoecology* **245**, 5–19. (doi:10.1016/j.palaeo.2006.02.030)

81. Jansonius J, Jenkins WAM. 1998 Chitinozoa. In *Introduction to Marine Micropaleontology*, pp. 341–357. Elsevier. (doi:10.1016/B978-044482672-5/50015-5)

82. Rigby S, Milsom CV. 2000 Origins, Evolution, and Diversification of Zooplankton. *Annu. Rev. Ecol. Syst.* **31**, 293–313. (doi:10.1146/annurev.ecolsys.31.1.293)

83. Larsson AI, Järnegren J, Strömberg SM, Dahl MP, Lundälv T, Brooke S. 2014 Embryogenesis and Larval Biology of the Cold-Water Coral Lophelia pertusa. *PLoS ONE* **9**, e102222. (doi:10.1371/journal.pone.0102222)

84. Sandin MM, Renaudie J, Suzuki N, Not F. 2025 Extant diversity, biogeography, and evolutionary history of Radiolaria. *Current Biology* **35**, 2524-2538.e6. (doi:10.1016/j.cub.2025.04.032)

85. Chang S, Feng Q, Zhang L. 2018 New Siliceous Microfossils from the Terreneuvian Yanjiahe Formation, South China: The Possible Earliest Radiolarian Fossil Record. *J. Earth Sci.* **29**, 912–919. (doi:10.1007/s12583-017-0960-0)

86. Braun A, Chen J, Waloszek D, Maas A. 2007 First Early Cambrian Radiolaria. *SP* **286**, 143–149. (doi:10.1144/SP286.10)

87. Butler AD. 2015 Decoding the fossil record of early lophophorates: systematics and phylogeny of problematic Cambrian Lophotrochozoa. PhD Thesis, Acta Universitatis Upsaliensis.

88. Ushatinskaya GT. 2008 Origin and dispersal of the earliest brachiopods. *Paleontol. J.* **42**, 776–791. (doi:10.1134/S0031030108080029)

89. Freeman G, Lundelius J. 2008 Brachiopod larvae through time. In *Fossils and Strata* (eds DAT Harper, SL Long, C Nielsen), pp. 251–257. Wiley-Blackwell. (doi:10.18261/9781405186643-2008-27)

90. Freeman G, Lundelius JW. 2007 Variance in protegular size: a correlate of larval trophic mode in Early Palaeozoic brachiopods. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh* **98**, 233–238. (doi:10.1017/S1755691007078383)

91. Liu Y, Melzer RR, Haug JT, Haug C, Briggs DEG, Hörnig MK, He Y, Hou X. 2016 Three-dimensionally preserved minute larva of a great-appendage arthropod from the early Cambrian Chengjiang biota. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 5542–5546. (doi:10.1073/pnas.1522899113)

92. Zhang X, Maas A, Haug JT, Siveter DJ, Waloszek D. 2010 A Eucrustacean Metanauplius from the Lower Cambrian. *Current Biology* **20**, 1075–1079. (doi:10.1016/j.cub.2010.04.026)

93. Burrows M, Hawkins S, Southward A. 1999 Larval development of the intertidal barnacles *Chthamalus stellatus* and *Chthamalus montagui*. *Journal of the Marine Biological Association of the United Kingdom* **79**, 93–101.

94. Akther H, Agersted MD, Olesen J. 2015 Naupliar and Metanaupliar Development of *Thysanoessa raschii* (Malacostraca, Euphausiacea) from Godthåbsfjord, Greenland, with a Reinstatement of the Ancestral Status of the Free-Living Nauplius in Malacostracan Evolution. *PLoS ONE* **10**, e0141955. (doi:10.1371/journal.pone.0141955)

95. Speyer SE, Chatterton BDE. 1990 Trilobite Larvae, Larval Ecology and Developmental Paleobiology. *Short courses paleontol.* **3**, 137–156. (doi:10.1017/S247526300000177X)

96. Harvey TH, Butterfield NJ. 2008 Sophisticated particle-feeding in a large Early Cambrian crustacean. *Nature* **452**, 868–871.

97. Harvey TH, Vélez MI, Butterfield NJ. 2012 Exceptionally preserved crustaceans from western Canada reveal a cryptic Cambrian radiation. *Proceedings of the National Academy of Sciences* **109**, 1589–1594.

98. Mussini G, Hagadorn JW, Miller AE, Karlstrom KE, Evans R, Dehler CM, Bastien S, Butterfield NJ. 2025 Evolutionary escalation in an exceptionally preserved Cambrian biota from the Grand Canyon (Arizona, USA). *Science Advances* **11**, eadv6383.

99. Chen H, Parry LA, Vinther J, Zhai D, Hou X, Ma X. 2020 A Cambrian crown annelid reconciles phylogenomics and the fossil record. *Nature* **583**, 249–252. (doi:10.1038/s41586-020-2384-8)

100. Zhang Z, Smith MR, Ren X. 2023 The Cambrian cirratuliform *Iotuba* denotes an early annelid radiation. *Proc. R. Soc. B.* **290**, 20222014. (doi:10.1098/rspb.2022.2014)

101. Caine EA. 1974 Comparative functional morphology of feeding in three species of caprellids (Crustacea, Amphipoda) from the northwestern Florida Gulf Coast. *Journal of Experimental Marine Biology and Ecology* **15**, 81–96. (doi:10.1016/0022-0981(74)90065-3)

102. Caine EA. 1977 Feeding mechanisms and possible resource partitioning of the caprellidae (Crustacea: Amphipoda) from Puget Sound, USA. *Mar. Biol.* **42**, 331–336. (doi:10.1007/BF00402195)

103. Guerra-García JM, Tierno De Figueroa JM. 2009 What do caprellids (Crustacea: Amphipoda) feed on? *Mar Biol* **156**, 1881–1890. (doi:10.1007/s00227-009-1220-3)

104. Lustri L, Collantes L, Esteves CJP, O'Flynn RJ, Saleh F, Liu Y. 2025 A Hypothesis on Suspension Feeding in Early Chelicerates (Offacolidae). *Diversity* **17**, 412. (doi:10.3390/d17060412)

105. Wainwright P, Carroll AM, Collar DC, Day SW, Higham TE, Holzman RA. 2007 Suction
feeding mechanics, performance, and diversity in fishes. *Integrative and Comparative*
Biology **47**, 96–106. (doi:10.1093/icb/icm032)

106. Van Wassenbergh S, Aerts P. 2009 Aquatic suction feeding dynamics: insights from
computational modelling. *J. R. Soc. Interface.* **6**, 149–158. (doi:10.1098/rsif.2008.0311)

107. Ortega-Hernandez J. 2025 Hallucigenia's diet illuminates the feeding ecology of Cambrian
lobopodians. *bioRxiv* , 2025.12.28.696761. (doi:10.64898/2025.12.28.696761)

108. Wolcott TG. 1981 Inhaling without ribs: the problem of suction in soft-bodied invertebrates.
The Biological Bulletin **160**, 189–197.

109. Shimeta J. 2009 Influence of flow speed on the functional response of a passive suspension
feeder, the spionid polychaete Polydora cornuta. *Mar Biol* **156**, 2451–2460.
(doi:10.1007/s00227-009-1270-6)

110. Trager G, Achituv Y, Genin A. 1994 Effects of prey escape ability, flow speed, and predator
feeding mode on zooplankton capture by barnacles. *Marine Biology* **120**, 251–259.
(doi:10.1007/BF00349685)

111. Leonard AB, Strickler JR, Holland ND. 1988 Effects of current speed on filtration during
suspension feeding in *Oligometra serripinna* (Echinodermata: Crinoidea). *Mar. Biol.* **97**,
111–125. (doi:10.1007/BF00391251)

112. Miller DC, Bock MJ, Turner EJ. 1992 Deposit and suspension feeding in oscillatory flows
and sediment fluxes.

113. Hart DD, Merz RA, Genovese SJ, Clark BD. 1991 Feeding postures of suspension-feeding
larval black flies: the conflicting demands of drag and food acquisition. *Oecologia* **85**, 457–
463. (doi:10.1007/BF00323756)

114. Abrahams MV. 1995 The interaction between antipredator behaviour and antipredator
morphology: experiments with fathead minnows and brook sticklebacks. *Canadian Journal*
of Zoology **73**, 2209–2215.

115. Losos JB, Mouton PLFN, Bickel R, Cornelius I, Ruddock L. 2002 The effect of body
armature on escape behaviour in cordylid lizards. *Animal Behaviour* **64**, 313–321.

116. Wahle RA. 1992 Body-Size Dependent Anti-Predator Mechanisms of the American Lobster.
Oikos **65**, 52. (doi:10.2307/3544887)

117. Whitman DW, Vincent S. 2008 Large size as an antipredator defense in an insect. *Journal of*
Orthoptera Research **17**, 353–371. (doi:10.1665/1082-6467-17.2.353)