Functional divergence of Capicua isoforms explains differential tissue vulnerability in neurological disease

Hamin Lee[1,2,3], Esmeralda Villavicencio Gonzalez[1,2,3], Elias M. Rivera[2,3], Mark A. Durham[2,3,4,5],

Ronald Richman[2,3], Elizabeth H.-Y. Chu[2,3], Kailey Xia[2,3], Hu Chen[3,6], Zhandong Liu[3,6], Surabi Veeraragavan[2,3],

Binoy Shivanna[7], Huda Y. Zoghbi[1,2,3,4,6,7,8,9,10*]

[1]Program in Genetics and Genomics, Baylor College of Medicine, Houston, TX 77030, USA

[2]Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, Texas 77030, USA.

[3]Jan and Dan Duncan Neurological Research Institute at Texas Children's Hospital, Houston, Texas 77030, USA

[4]Program in Developmental Biology, Baylor College of Medicine, Houston, TX 77030, USA

[5]Medical Scientist Training Program, Baylor College of Medicine, Houston, TX 77030, USA

[6]Department of Pediatrics, Baylor College of Medicine, Houston, Texas 77030, USA

[7]Department of Pediatrics, Texas Children's Hospital, Baylor College of Medicine, Houston, TX 77030, USA

[8]Department of Neuroscience, Baylor College of Medicine, Houston, TX 77030, USA

[9]Howard Hughes Medical Institute, Baylor College of Medicine, Houston, TX 77030, USA

[10]Lead contact

*Correspondence: hzoghbi@bcm.edu

SUMMARY

Many neurological diseases attack specific brain regions despite widespread expression of the disease-related protein. Spinocerebellar ataxia type 1 (SCA1) primarily affects the cerebellum, though Ataxin-1 (ATXN1) is widely expressed. We previously showed that intensified interaction between mutant ATXN1 and Capicua (CIC) drives SCA1 pathogenesis in the cerebellum, whereas ATXN1 loss augments amyloid beta production in the hippocampus and cortex. CIC, however, forms a complex with ATXN1 and its paralog, Ataxin-1-like (ATXN1L)—yet knockout of either yields completely different phenotypes. To determine whether this could be due to CIC having two isoforms, we generated mice bearing either the long (CIC-L) or short (CIC-S) isoform. Loss of CIC-L led to cognitive deficits, whereas loss of CIC-S caused perinatal lethality, phenocopying, ATXN1 and ATXN1L knockout mice, respectively. Furthermore, CIC-L preferentially interacts with ATXN1, and CIC-S with ATXN1L. Our data underscore the importance of isoform–paralog interplay in studying regional vulnerability in neurodegenerative diseases.

Keywords

Capicua, CIC, ATXN1, ATXN1L, isoform, paralog, alloform, protein complex, neurodegeneration, neurodevelopment

INTRODUCTION

One of the puzzling features of many neurological diseases is that they seem to affect particular brain regions or cell types even though the disease-related protein is expressed widely throughout the brain and often in other tissues as well [1, 2]. Typically the disease ramifies outward from that locus of initial damage: Alzheimer's, for example, first targets the entorhinal cortex and hippocampus, whereas Parkinson's disease centers on the substantia nigra [1, 2]. The same regional specificity is seen in Mendelian disorders as well, which allow us to more readily trace their developmental origins. For example, spinocerebellar ataxia type 1 (SCA1) is caused by expansion of a polyglutamine repeat tract in the gene *ATAXIN-1* (*ATXN1*) [3, 4]. The ATXN1 protein is a transcriptional co-repressor that is expressed throughout the central nervous system (CNS) in both neurons and glia and is also expressed in organs such as the heart and liver [5]. In the CNS, it is most abundant in hippocampal pyramidal cells and cerebellar Purkinje cells, yet SCA1 pathology is largely restricted to the Purkinje cells and brainstem [6]. Interestingly, a lack of ATXN1 does not cause ataxia but rather learning and memory deficits [7-9]: copy number variation of *ATXN1* is associated with Alzheimer's disease risk [10, 11], and *Atxn1* knock-out in mice potentiates amyloid beta production [12]. If, however, we delete *Atxn1* and its paralog, *Ataxin-1*-like (*Atxn1l*), the double mutant mice develop a quite different constellation of abnormalities that includes hydrocephalus, omphalocele lung defects, and perinatal lethality [13]. Loss of *Atxn1l* by itself causes the same phenotypes but with reduced penetrance [13].

It is reasonable to assume that this combination of regional specificity independent of expression level and the very different tissues affected by loss of ATXN1L indicates that ATXN1 and ATXN1L have different native interactors. It was rather surprising, then, to find that they both form a complex with the transcriptional repressor Capicua (CIC) [13, 14], and both proteins are critical for maintaining CIC stability [13-16]. ATXN1 and ATXN1L share a conserved CIC-binding domain [16-18] and can substitute for each other in binding to CIC, exhibiting partial functional redundancy [8, 13, 19]. CIC is also broadly expressed [13, 14, 20] and interacts with ATXN1 throughout the brain [14]; the effect of the polyglutamine expansion in mutant ATXN1 is to intensify the effect of this interaction, leading to hyper-repression of CIC targets [21, 22]. The formation of the mutant ATXN1–CIC complex is necessary for Purkinje cell pathology in SCA1 [22, 23]. On the other hand, loss of ATXN1 reduces CIC levels, leading to derepression of CIC targets *Etv4* and *Etv5* [12]. These transcription factors have binding sites upstream of *Bace1* and increase its expression, which in turn promotes amyloidogenic processing of amyloid precursor protein (APP) and may be a risk factor for Alzheimer disease [12]. Notably, this upregulation occurs only in the cortex and hippocampus, not in the cerebellum [12].

CIC has broad developmental roles in both the peripheral tissues and the CNS. In mice, complete loss of CIC causes perinatal lethality, with hydrocephalus, omphalocele and impaired lung development [13, 24]— reminiscent of *Atxn1l* knockout (KO) mice. In humans, heterozygous loss-of-function variants cause CIC haploinsufficiency syndrome (CHS), a neurodevelopmental disorder characterized by intellectual disability,

autism spectrum disorder (ASD), attention deficit hyperactivity disorder (ADHD), and seizures [14]. Dual loss of both ATXN1 and ATXN1L phenocopies the complete loss of CIC [13], and conditional KO in the brain causing a CHS-like phenotype [14]. How do these two paralogs, both acting through CIC, produce such distinct loss-of-function outcomes?

We hypothesized that the answer might arise from the fact that CIC exists in two isoforms generated by alternative promoter usage, namely, CIC-L (long) and CIC-S (short) [16]. Both isoforms share the HMG box and C1 DNA-binding domains as well as the ATXN1/ATXN1L-binding domain (ATXN1/1L-BD) [16, 25, 26]. The functions of the isoform-specific N-termini remain largely undefined, leaving open the possibility that the two isoforms have distinct biological roles. To determine whether isoform-specific functions contribute to the regional specificity of ATXN1 and ATXN1L gain- or loss-of-function phenotypes, we generated CIC isoform-specific KO mice, characterized their phenotypes, and evaluated transcriptomic changes. We then performed biochemical studies to assess the formation of different complexes. We found that, surprisingly, subtle differences in the relative abundance of CIC isoforms and ATXN1 paralogs at the protein level, together with the differential complexes they assemble, can dictate regional vulnerability.

RESULTS

CIC isoforms and ATXN1/ATXN1L show different spatiotemporal expression patterns

To investigate how the expression patterns of CIC isoforms relate to that of the CIC–ATXN1/1L complex, we profiled protein levels of each CIC isoform together with ATXN1 and ATXN1L in the cortex, cerebellum, and lung, of wild-type (WT) mice at four developmental stages: birth (P0), postnatal day 6 (P6), postnatal day 21 (P21), and 12 weeks (**Figure 1A, B**). These regions and developmental timepoints were selected based on prior evidence of tissue-specific changes due to disruption of the CIC–ATXN1/1L complex: in the forebrain, *Cic* knockout (KO) or *Atxn1/Atxn1l* double KO in mice leads to selective reduction of cortical layers 2–4 beginning around postnatal day 5 (P5), with behavioral abnormalities emerging by 3 months of age [14]. In the lung, *Atxn1l* KO in mice disrupts the transcriptional program of early alveolarization at P6 and results in reduced survival by P21 [13]. The cerebellum, which is highly sensitive to ATXN1 gain-of-function toxicity, was included as a comparator to these loss-of-function–vulnerable tissues [23].

ATXN1L and both CIC isoforms showed a similar pattern of expression in the lungs, with the highest levels at P6 and decreasing at later timepoints, whereas ATXN1 expression decreases over time after P0 (**Figure 1A, B**). Notably, ATXN1L levels are considerably higher in the lungs compared to the cortex and cerebellum, with peak expression at P6. In the cortex, CIC-L and CIC-S expression patterns diverged after P6: CIC-L showed a slight increase at P21 that was maintained at 12 weeks, while CIC-S markedly decreased between P6 and P21 and remained low at 12 weeks. In the cerebellum, all four proteins showed a pronounced increase from P6 to P21, followed by a decrease in CIC-S and ATXN1L at 12 weeks. In 12-week-old brains, we also looked at the

hippocampus, striatum, and brainstem and found that both CIC isoforms were most highly expressed in the

cerebellum and least expressed in the brainstem (**Figure S1**).

Thus, CIC, ATXN1, and ATXN1L show distinct tissue- and isoform-specific expression patterns in mice

from birth through young adulthood.

CIC-S is critical for survival and for lung alveolarization

To dissect the individual roles of each CIC isoform, we used CRISPR/Cas9 to generate *Cic-L*- and *Cic-S*-

specific KO mice by targeting the unique first coding exon of each isoform. For *Cic-S*-KO, we targeted *Cic-S*

exon 1 with a single guide RNA (sgRNA) and selected founder lines with deletions that removed the in-frame

downstream ATG (**Figure 2A, S2A** and **B**). For *Cic-L*-KO, we used two sgRNAs to introduce a deletion within

exon 2, resulting in a frameshift mutation and selective disruption of the long isoform. We confirmed the deletions

by Sanger sequencing (**Figure S2C**) and used western blot to validate complete loss of either CIC-L or CIC-S in

homozygous KO animals (**Figure 2B**).

Since germline KO of *Cic* results in perinatal lethality [14, 24], we sought to determine how loss of each

isoform affects survival. We used chi-square analysis to compare observed genotype frequencies to the expected

Mendelian ratios of WT (25%), heterozygous KO (50%), and KO (25%) at postnatal days 0, 6, and 21 (P0, P6,

and P21; **Tables 1** and **2**). Both *Cic-S* and *Cic-L*-KO animals exhibited the expected Mendelian ratios at P0,

indicating no embryonic lethality. At P6, neither group showed significant deviation from expected ratios, but by

P21, approximately 70% of *Cic-S*-KO had died (chi-square analysis, $p = 0.007$; **Table 1**) while *Cic-L*-KO animals

maintained normal survival (chi-square analysis, $p = 0.74$; **Table 2**).

We measured the weight and lifespan of surviving animals beyond P21. *Cic-S*-KO and *Cic-L*-KO mice

weighed less than WT throughout the 9-week period, although *Cis-S*-KO stopped gaining weight around week 7

while *Cic-L*-KO mice continued to grow, reaching near WT weight by week 9 (**Figure 3A**). *Cic-S*-KO mice

exhibited significant early lethality after 3 weeks, with 60% (15 of 25 animals) dying before 20 weeks, whereas

Cic-L-KO animals did not display reduced survival (**Figure 3B**). Among the *Cic-S*-KOs that died, 80% developed

hydrocephalus accompanied by kyphosis, emaciation, and lethargy prior to death. One animal experienced

spontaneous and ultimately fatal home-cage seizures (**Figure 3C**). The remaining two animals that died without

clear hydrocephalus or seizures showed further weight loss beyond their already reduced baseline prior to death.

Some *Cic-S*-KO mice survived up to 40 weeks without any of these features, however, suggesting incomplete

penetrance. Neither *Cic-L*-KO nor WT animals developed any noticeable phenotype during this period.

These abnormalities of the *Cic-S*-KO mice are strongly reminiscent of those previously reported in *Atxn1l*-

KO mice (hydrocephalus with incomplete penetrance, lower body weight compared to WT controls, and reduced

survival by P21) [13]. Because *Atxn1l*-KO mice also have impaired alveolarization mediated by loss of CIC

function that dysregulates extracellular matrix (ECM) remodeling and alveolarization during late alveolarization

stage (P17–P23), which was also present in another line of mice bearing a *Cic* loss-of-function allele [13], we

performed hematoxylin and eosin (H&E) staining of lung tissue at P21 and quantified the mean linear intercepts

(MLI). Despite potential survivor bias in *Cic-S*-KO animals at this stage, the MLI values were significantly

greater in *Cic-S*-KO lungs than in WT littermates (*Cic-S*-KO, 29.73 ± 1.05 vs. WT littermate, 25.86 ± 0.71;

p=0.038, unpaired t-test; **Figure 3D**) whereas MLI values of *Cic-L*-KO lungs were comparable to WT littermate

controls (*Cic-L*-KO, 27.21 ± 1.36 vs. WT littermate, 28.14 ± 1.17; p=0.658 , unpaired t-test; **Figure 3E**).

Therefore, loss of CIC-S, but not CIC-L, impairs alveolarization, supporting functional overlap between CIC-S

and ATXN1L.

CIC-L is crucial for neurobehavioral function

Given the critical role of CIC in neurodevelopment [14, 27, 28], we performed a battery of behavioral

assays in *Cic-S*-KO, *Cic-L*-KO, and WT littermates to assess anxiety-like behavior, general locomotor activity,

motor coordination, and social interaction—phenotypes previously linked to CIC dysfunction in both people with

CHS and in mouse models [14, 29]. *Cic-L*-KO mice displayed reduced anxiety-like behavior in the elevated plus

maze assay, spending significantly more time in the open arms (**Figure 4A**), and hyperactivity in the open field

test, with increased total distance traveled and higher speed compared to controls (**Figure 4B**). They had a shorter

latency to fall from the rotarod (**Figure 4C**) and took more time to descend in the pole test (**Figure S3A**). In the

fear-conditioning assay, *Cic-L*-KO mice showed deficits in learning and memory, freezing less in both contextual

(**Figure 4D**) and cued fear tests (**Figure 4E**). In the three-chamber assay, both KO lines preferred to interact with

the conspecific than the object (**Figure S3B**).

Cic-S-KO mice exhibited minimal behavioral abnormalities (**Figure 4A, B** and **C**). The only significant

phenotype that *Cic-S*-KO mice displayed was increased freezing in the contextual test (**Figure 4D**) and pre-tone

in the cued test (**Figure 4E**), which do not signify any learning or memory deficit. These data suggest that, for the

animals that survive because of decreased penetrance, CIC-S is not critical for most neurobehavioral functions,

with only mild effects on baseline freezing behavior.

Cic-L-KO mice phenocopied *Atxn1*-KO mice, which show learning deficits [7-9]. Collectively, the survival

and neurobehavioral data point to a functional and spatio-temporal convergence between CIC-L and ATXN1 on

the one hand, and CIC-S and ATXN1L on the other.

Region-specific transcriptomic overlap between CIC isoform and ATXN1 paralog KOs

To determine whether phenotypic similarities between these two pairs of mouse lines reflect shared

transcriptional changes, we performed bulk RNA sequencing (RNA-seq) on tissues linked to the phenotype of

each CIC isoform loss (P6 lung and 12-week cortex) and compared the differentially expressed genes (DEGs)

previously identified in *Atxn1*-KO and *Atxn1l*-KO mice. We used the cerebellum as an internal control, since it

does not contribute to the CHS phenotype despite its great sensitivity to polyglutamine-expanded ATXN1 [23]

In the P6 lung, *Cic-S*-KO samples exhibited a greater number and magnitude of DEGs (*p*-adjusted < 0.05

and |log2 fold change (Log2FC)| > 0.25) than *Cic-L*-KO, with enrichment of upregulated genes consistent with

CIC's role as a transcriptional repressor (**Figure S4A** and **B**). These differences aligned with more pronounced

survival and developmental phenotypes in *Cic-S*-KO animals (**Table 1**, **Figure 3**). Gene set enrichment analysis

(GSEA) [30] revealed distinct GO term enriched for each isoform KO, with *Cic-S*-KO enriched for immune-

related terms (**Figure S4C**) and *Cic-L*-KO for chromosome segregation-related terms (**Figure S4D**).

Consistent with the phenotypic similarities, the *Cic-S*-KO DEGs showed stronger overlap with the *Atxn1l*-

KO DEGs [13] (Fisher's exact test, OR = 19.85, $p = 6.32 \times 10^{-132}$; **Figure S5A**) than *Cic-L*-KO (OR = 3.58, $p =$

1.7×10^{-12}; **Figure S5B**), with most shared genes showing concordant expression changes. Interestingly, although

Atxn1-KO lungs had only 34 DEGs, just over a third overlapped with the *Cic-S*-KO DEGs (OR = 4.74, p = 5.83 ×

10^{-5}; **Figure S5C**) or the *Cic-L*-KO DEGs (Fisher's exact test, OR = 9.86, $p = 8.54 \times 10^{-8}$; **Figure S5D**). Given

the role of the *Etv–Mmp* axis in *Atxn1l*-KO alveolarization defects [13] and the fact that *Etv1/4/5* are direct CIC

targets [13, 22, 31, 32], we examined their expression specifically. *Etv4, Mmp8, Mmp9,* and *Mmp12* were strongly

upregulated in *Cic-S*-KO lungs, while only *Etv1* and *Mmp13* showed mild upregulation in *Cic-L*-KO (**Figure**

S5E). GSEA revealed enrichment of extracellular matrix (ECM) assembly-related pathways in *Cic-S*-KO,

whereas *Cic-L*-KO showed weak or non-significant enrichment (**Figure S5F**). These results suggest that CIC-S

and ATXN1L regulate similar genes in the lung during lung development.

In the 12-week-old cortex, there were more DEGs in the *Cic-L*-KO with larger fold-changes than in *Cic-S*-

KO (**Figure S6A, B**). Whereas *Cic-S*-KO top GSEA terms were mostly related to glia or to bioenergetics (**Figure**

S6C, D), *Cic-L*-KO terms involved synapse and neurotransmission-related functions (**Figure S6E, F**).

Comparison with *Atxn1*-KO cortex DEGs (*Xhako et al., unpublished data*) revealed greater overlap and stronger

enrichment with *Cic-L*-KO than *Cic-S*-KO, with more genes changing in the same direction (**Figure S7A, B**).

Enrichment analysis of these concordant DEGs showed that only *Cic-L*-KO shared synapse- and memory-related

GO terms with *Atxn1*-KO (**Figure S7C, D**). Consistent with previous findings that the *Etv4/Etv5–Bace1* axis is

upregulated in *Atxn1*-KO cortex [12], we observed similar upregulation of this axis in *Cic-L*-KO cortex (**Figure**

S7E). These findings support the idea that CIC-L and ATXN1 regulate similar gene networks that underlie

learning and memory.

In the cerebellum, a region generally more resistant to the loss of the CIC–ATXN1/1L complex but clearly

vulnerable to the gain of CIC-ATXN1 function [22, 23], we observed fewer DEGs in *Cic-L*-KO compared to the

cortex, but more DEGs in *Cic-S*-KO relative to its cortical counterpart (**Figure S8A and B**), potentially reflecting

higher CIC-S expression in this region at 12 weeks (**Figure S1**). The two KOs also showed distinct GO term

enrichments (**Figure S8C–F**), and DEG overlap with *Atxn1*-KO was stronger for *Cic-S*-KO (Fisher's exact test,

OR = 2.41, $p = 1.69 \times 10^{-17}$; **Figure S9A**) than for *Cic-L*-KO (Fisher's exact test, OR = 1.61, $p = 2.85 \times 10^{-8}$;

Figure S9B). *Etv4/5*, which were strongly upregulated in the cortex of *Cic-L*-KO, showed less upregulation in the cerebellum, and *Bace1* remained unchanged (**Figure S9C**).

CIC-L and CIC-S, therefore, largely regulate different biological processes. Their respective transcriptional convergences with ATXN1/1L are both region- and timepoint-dependent.

CIC-S preferentially associates with ATXN1L, and CIC-L associates with ATXN1, in a region-specific manner

Given the region-specific transcriptomic overlap—between the cortex in *Cic-L*–KO and *Atxn1*–KO mice, and the cerebellum in *Cic-S*–KO and *Atxn1*–KO mice—we sought to determine whether CIC isoform–paralog pairs form distinct complexes across brain regions. We performed size exclusion chromatography (SEC) on protein extracts from cortical and cerebellar tissue harvested from 2–3-month-old *Cic-S*-KO and *Cic-L*-KO mice to examine the elution profiles of CIC, ATXN1, and ATXN1L.

In the WT cortex both ATXN1 and ATXN1L co-eluted with CIC in a high molecular weight complex (~1.7 MDa, fraction 12), as well as in distinct smaller complexes (ATXN1 peaking at fraction 15, ~300 kDa; ATXN1L at fractions 16–17, ~170–88 kDa; **Figure 5A, B**), consistent with prior reports [14]. In contrast, both *Cic-S*-KO and *Cic-L*-KO cortices showed less ATXN1 and ATXN1L in the high molecular weight fractions and a concomitant increase in smaller fractions, suggesting that CIC is essential for maintaining their incorporation into the larger complex (**Figure 5A, B**). Interestingly, in *Cic-L*-KO cortex, ATXN1 showed a stronger shift from the large to small complex (large-to-small complex ratio: 0.41± 0.05, vs. WT 1.00± 0.08; p = 0.003, one-way ANOVA) than in *Cic-S*-KO (0.67± 0.08; p =0.039) (**Figure 5C**, left), indicating preferential association of ATXN1 with CIC-L. In contrast, in *Cic-S*-KO cortex ATXN1L shifted markedly to smaller fractions (0.11± 0.03, vs. WT 1.00± 0.03; p = 0.0001) and was nearly absent from the large complex. This shift was less pronounced in *Cic-L*-KO (0.51± 0.13; p = 0.006; **Figure 5C**, right), suggesting a stronger association of ATXN1L with CIC-S. These findings confirm isoform-specific associations within the CIC–ATXN1/ATXN1L complex in the cortex.

In the *Cic-S*-KO cerebellum, ATXN1L shifted markedly toward smaller fractions (0.07± 0.03, vs. WT 1.00± 0.08; p = 0.0002). The shift was more modest in *Cic-L*-KO (0.63± 0.13; p = 0.042; **Figure 5D–F**), consistent with its preferential association with CIC-S observed in the cortex. In contrast to the cortex, ATXN1 also shifted in *Cic-S*-KO cerebellum (0.42± 0.06, vs. WT 1.00± 0.17; p = 0.015), while its distribution remained largely unchanged in *Cic-L*-KO (0.90± 0.11; p = 0.82; **Figure 5D-F**).

The cerebellum has the highest level of CIC-S among all brain regions, as well as the highest total CIC expression (**Figure S1A–C**). ATXN1 is expressed at lower levels, however, resulting in the highest CIC-to-ATXN1 ratio in the cerebellum (**Figure S1D**), which may lead to a greater proportion of ATXN1 bound to CIC. Supporting this hypothesis, comparison of WT cerebellar and cortical SEC patterns (**Figure S10**) revealed a striking difference: the cerebellum exhibits a higher proportion of ATXN1 in the large complex and lower levels

in the small complex (**Figure S10C**), a pattern not observed for ATXN1L (**Figure S10D**). The relatively high

CIC-S levels may explain why CIC-S is especially important for ATXN1 large complex formation in the

cerebellum. Moreover, the combination of a high CIC/ATXN1 ratio likely predisposes the cerebellum to gain-of-

function effects of the complex. It is worth noting that cerebellar single-nucleus RNAseq [33] show that both *Cic*

and *Atxn1* are highly expressed in Purkinje cells, which are most vulnerable to SCA1 (**Figure S11**).

Overall, these results indicate that CIC isoforms and ATXN1 paralogs favor particular interactions

depending on the context. While the CIC-S–ATXN1L interaction appears strong and consistent across regions,

ATXN1 associates with different CIC isoforms in different tissues.

The unique N-terminal of each CIC isoform mediates the ATXN1 paralog-specific interaction

To understand the mechanism of isoform-specific complex formation of CIC with either ATXN1 or

ATXN1L, we first examined the interaction of CIC-L and CIC-S with ATXN1 and ATXN1L by performing

coimmunoprecipitation (co-IP) assays in WT cortical lysates. Despite comparable expression levels of CIC

isoforms in the cortex, ATXN1 immunoprecipitation showed strong enrichment of CIC-L relative to CIC-S

(**Figure 6A**). Conversely, ATXN1L immunoprecipitation demonstrated an even more pronounced enrichment of

CIC-S over CIC-L (**Figure 6B**), which may explain the more prominent ATXN1L shift observed in SEC (**Figure**

5A-C). These results support the hypothesis that ATXN1 preferentially interacts with CIC-L, whereas ATXN1L

favors CIC-S.

Since CIC-S and CIC-L share a canonical ATXN1/ATXN1L binding domain (ATXN1/1L-BD) that can

bind both ATXN1 and ATXN1L, we hypothesized that there should be isoform-specific binding sites outside the

ATXN1/1L-BD that mediate preferential interaction between CIC-S and ATXN1L, and CIC-L and ATXN1. To

test this possibility, we performed the analysis using human CIC, ATXN1, and ATXN1L sequences. The W37A

mutation in CIC-S has been previously shown to disrupt binding with the AXH domain of ATXN1[18]. Using the

corresponding mutation in CIC-L (W946A), we found that CIC-S W37A lost interaction with ATXN1 but CIC-L

W946A retained ATXN1 binding (**Figure 6C**). This suggests that CIC-L contains an additional ATXN1-binding

region. We saw the opposite with ATXN1L, despite CIC-S containing only 22 isoform-specific amino acids at its

N-terminus: ATXN1L still bound CIC-S W37A but failed to interact with CIC-L W946A (**Figure 6D**). These data

indicate that distinct, noncanonical regions in each isoform mediate selective interactions with ATXN1 and

ATXN1L.

To test whether the N-terminal region of CIC-L can mediate the selective interaction with ATXN1, we

expressed this region alone and performed co-IP. The CIC-L N-terminus, which spans 931 amino acids and does

not contain the ATXN1/1L-BD, interacted with ATXN1 but not with ATXN1L (**Figure 6E, F**). Thus, the N-

terminal region of CIC-L is sufficient to mediate interaction with ATXN1, but not ATXN1L, and may thereby

contribute to isoform-paralog specificity.

Together, these findings reveal a previously unrecognized mechanism by which CIC isoforms interact with
different ATXN1 paralogs through isoform-specific domains, establishing a novel molecular basis for paralog-
specific complex formation.
276

CIC-L expression is more dependent on ATXN1 than CIC-S

Given the preferential interaction between ATXN1 and CIC-L, and the known role of ATXN1 in
maintaining CIC protein stability [16], we hypothesized that CIC-L would be more dependent on ATXN1. To test
this, we measured CIC-L and CIC-S protein levels in WT and *Atxn1*-KO mice across brain regions sensitive to
ATXN1 loss (cortex, hippocampus) or gain (cerebellum, brainstem). CIC-L levels were significantly lower in all
four regions in *Atxn1*-KO mice, but CIC-S levels were more modestly reduced, and only in the cerebellum
(**Figure S12A-D**). The cerebellar reduction in CIC-S was consistent with our SEC analysis showing a shift of
CIC-S in ATXN1-containing fractions (**Figure 5D-F**). IP of ATXN1 confirmed a preferential association with
CIC-L in all brain regions examined (**Figure S12E**). These findings indicate that the cortical and hippocampal
abnormalities in *Atxn1*-KO mice may result primarily from the loss of CIC-L function.

DISCUSSION

The present study uncovers functionally distinct roles for the two CIC isoforms, CIC-L and CIC-S. The
overlap of transcriptomic changes in *Cic-S*-KO and *Atxn1l*-KO mice explains why these two lines show the same
early lethality, hydrocephalus, and lung alveolarization defects, while *Cic-L*-KO and *Atxn1*-KO mice share
learning and memory deficits. These phenotypic parallels reveal an unexpected degree of specificity in CIC–
ATXN1 complex formation, given that both isoforms share the canonical ATXN1/ATXN1L binding domain
(ATXN1/1L-BD); the specificity is attributable to the unique N-terminal domains of each isoform that favor
binding between CIC-S and ATXN1L, and between CIC-L and ATXN1. The sharp distinctions between the two
sets of phenotypes also provides compelling *in vivo* support of the idea that different isoforms greatly expand
protein interaction capacities and may indeed serve as "functional alloforms" [34].
Prior studies in *Drosophila* have shown that CIC isoforms play distinct roles during development: Cic-S
is critical for early embryogenesis through its interaction with Groucho [35], while Cic-L is essential for
oogenesis [36]. These findings support the idea that the divergent N-terminal regions of CIC isoforms confer
isoform- and context-specific functions. However, *Drosophila* Cic-S has evolved on its own [35], with the species
possessing only Atx-1 (containing the AXH domain) and lacking the paralog pairs found in mammals.
Furthermore, there is no evidence for an endogenous functional complex between *Drosophila* Cic and Atx-1. By
using a mammalian system in which both CIC isoforms and ATXN1 paralogs are highly conserved, we identified
a previously unrecognized layer of regulation: isoform–paralog–specific interactions.
These isoform-specific roles are not only of evolutionary interest but also carry direct clinical relevance.

Recently, a CIC-L–specific mutation was reported in a patient with CHS [37], which is consistent with our finding that CIC-L is critical for neurological function. There may be additional patients with CIC-L variants who remain unrecognized. Furthermore, given the role of CIC-L in learning and memory and its regulation of the *Etv4/5–Bace1* axis, CIC-L variants could, like *ATXN1* [10-12, 38, 39]. contribute to susceptibility to Alzheimer's disease. More broadly, the expression of different isoforms can be dysregulated in disease, but gene-level expression measures commonly used in biology average the translational potential of each isoform and therefore may not accurately reflect protein concentration. It has also been proposed that isoform-specific therapeutics would improve efficacy and reduce adverse effects [40].

Our findings help resolve two longstanding questions: (1) What underlies the regional vulnerability to loss or gain of the CIC–ATXN1 complex despite its broad expression? (2) How does the loss of ATXN1 versus ATXN1L lead to distinct phenotypes, despite both acting through CIC? We propose that the combination of isoform–paralog–specific interactions and regional expression patterns determines the differential sensitivity to complex disruption. Consistent with this model, the similarity in learning and memory deficits, along with upregulation of the *Etv4/5–Bace1* axis, in both *Atxn1*-KO and *Cic-L*-KO mice suggests that these phenotypes arise from preferential CIC-L–ATXN1 complex formation in the cortex and possibly the hippocampus.

Although we did not assess isoform-specific contributions to the cerebellar and brainstem pathology seen in SCA1, the current study does shed light on this question. We find that total CIC levels and CIC/ATXN1 ratios are highest in the cerebellum, with a greater proportion of ATXN1 being incorporated into the large complex. Our prior work in SCA1 mice demonstrated that ~50% reduction of total CIC (affecting both isoforms) [21] or a polyQ-ATXN1 mutation that would abolish binding to both isoforms fully rescues the Purkinje cell phenotype [22, 23]. These findings indicate that total CIC levels bound to ATXN1, rather than isoform identity, are critical for Purkinje cell degeneration. By contrast, CIC expression is lowest in the brainstem, which may be why loss of the ATXN1-CIC interaction does not fully rescue brainstem SCA1 phenotypes [21, 22] and why CIC motifs are less enriched in the brainstem's disease-associated gene network [41]. In SCA1, the brainstem's ATXN1 protein complex composition is distinct from that in the cerebellum [42], indicating that brainstem pathology arises through different mechanisms.

Our findings also allow us to draw some inferences that can be tested in the future. First, ATXN1 loss does not fully recapitulate CIC-L deficiency: transcriptomic comparisons reveal only partial overlap between *Atxn1*-KO and *Cic-L*-KO datasets, indicating that ATXN1 and CIC-L have functions outside of their shared complex. Indeed, ATXN1 has interactors beyond CIC [22, 43-46], and such partners may act in compensatory transcriptional or chromatin regulatory pathways, buffering CIC-dependent gene expression programs.

What are the ATXN1/ATXN1L-independent mechanisms that distinguish CIC-L from CIC-S? If isoform differences were driven solely by interactions with ATXN1 and ATXN1L, *Cic-L*-KO and *Cic-S*-KO transcriptomes would be expected to fully overlap, differing only in the magnitude of differential expression.

Instead, the partial overlap of DEGs may reflect isoform-specific interactors that influence target gene selection.

Alternatively, the isoforms may differ in cell type expression, affecting distinct cellular populations. Divergent

GO term enrichments support this notion. Profiling isoform-specific expression across cell types and applying

single-cell transcriptomics to each isoform KO will be critical to dissect these contributions.

Finally, our findings highlight a broader principle that paralogous proteins and their isoform-specific

partners can form distinct regulatory complexes with completely different biological functions. The CIC–

ATXN1/1L complex may serve as a paradigm for understanding how paralog-isoform specific interactions shape

regional vulnerability and disease susceptibility in other diseases such as those involving amyloid precursor

protein [47-53], tau [54, 55], or α-synuclein [56], all of which exist within paralog families and have multiple

isoforms.

Limitations of the study

Because the majority of *Cic-S*-KO animals die before P21, only the healthiest underwent neurobehavioral

characterization. We therefore cannot rule out the possibility that young animals might also have neurological

deficits. To better understand the role of CIC isoforms in the adult brain, future studies employing conditional

knockout models of each isoform, as well as of *Atxn1l,* will be critical. Such models could uncover additional

functions of CIC isoforms and context-dependent effects not captured in our germline KO approach.

ACKNOWLEDGEMENTS

We thank Ji-Yoen Kim, Sameer Bajikar, Dah-eun Chloe Chung, Harini Pallikarana, Sih-Rong Wu, and Xue Deng for their critical comments on the manuscript, and Harry Orr and Vicky Brandt for editorial input. We further thank the Baylor College of Medicine (BCM) Genetically Engineered Rodent Model (GERM) Core, Jan and Dan Duncan Neurological Research Institute (NRI) Rodent Neurobehavior Core, and Neuropathology Core. Funding: This work was funded by National Institute of Aging/NIH grant F31 AG077918 (H.L.), Cockrell Family Funding (H.L.), National Institute of Neurological Disorders and Stroke/NIH grant R01 NS027699 (H.Y.Z.), Freedom Together Foundation grant MR-2023-4260 (H.Y.Z.), Howard Hughes Medical Institute (H.Y.Z.), Chao Endowment (Z.L.), Huffington Foundation (Z.L.), National Institutes of Health Cancer Center Grant /NIH grant P30 CA125123 (BCM GERM Core), Cancer Biology Research Grant /NIH grant 1R50CA283804 (BCM GERM Core), Eunice Kennedy Shriver National Institute of Child Health and Human Development Intellectual and Developmental Disabilities Research Center/NIH grant P50HD103555 (NRI Rodent Neurobehavior Core and Neovascularization Core).

AUTHOR CONTRIBUTIONS

H.L. and H.Y.Z. conceived the study, designed the experiments, and analyzed and interpreted the data. H.L. generated mouse models. H.L., E.V.G., and E.M.R. performed protein biochemistry from mouse tissues, and H.L. performed cellular experiments. H.L. and E.H.-Y.C. performed RNA extractions, and H.C. performed alignment of raw sequencing data; H.L. performed RNA-seq analysis. H.C. and Z.L. supervised RNA-seq processing and analysis. H.L., E.H.-Y.C., and K.X. performed mouse genotyping, colony management, and survival and weight tracking. M.A.D. designed and generated the hCIC-L plasmid. H.L. and M.A.D. generated mutant constructs and performed additional cloning, with technical assistance from E.H.-Y.C. R.R. performed size-exclusion chromatography. S.V. performed mouse behavior experiments. B.S. processed lung tissue for histopathology and performed morphometric analysis. H.L. wrote the manuscript and prepared the figures with input from all authors. H.Y.Z. interpreted and reviewed all data and edited the manuscript.

DECLARATION OF INTERESTS

H.Y.Z. is on the scientific advisory board of Neurogene, Lyterian, the Column Group, and Cajal Therapeutics, and is a co-founder of Cajal Therapeutics. H.Y.Z. is also a director at Regeneron. None of these activities have any relevance to the work presented in this paper.

  TABLES

390 **Table 1. Number of surviving animals from $Cic^{S\text{-}KO/+}$ intercrosses at a given time point**

Genotype	Observed (%) at P0	Observed (%) at P6	Observed (%) at P21
$Cic^{+/+}$	22 (24.4 %)	32 (28.1 %)	31 (29.8%)
$Cic^{S\text{-}KO/+}$	50 (55.6 %)	65 (57.0 %)	64 (61.5%)
$Cic^{S\text{-}KO/S\text{-}KO}$	18 (20.0 %)	17 (14.9 %)	9 (8.6 %)
Total	90	114	104
Chi-square[*]	ns (0.26)	ns (0.18)	** (0.007)

391 [*]Chi-square analysis of the observed ratio compared to the expected Mendelian ratio of $Cic^{+/+}$ (25%), $Cic^{S\text{-}KO/+}$
392 (50%), and $Cic^{S\text{-}KO/S\text{-}KO}$ (25%). Each time point is an independent cohort. ** $p<0.01$

393
394 **Table 2. Number of surviving animals from $Cic^{L\text{-}KO/+}$ intercrosses at a given time point**

Genotype	Observed (%) at P0	Observed (%) at P6	Observed (%) at P21
$Cic^{+/+}$	22 (20.9 %)	18 (25.7 %)	28 (26.7 %)
$Cic^{L\text{-}KO/+}$	53 (50.5 %)	38 (54.3 %)	54 (51.4 %)
$Cic^{L\text{-}KO/L\text{-}KO}$	30 (28.6 %)	14 (20.0 %)	23 (21.9 %)
Total	105	70	105
Chi-square[*]	ns (0.53)	ns (0.86)	ns (0.74)

396 [*]Chi-square analysis of the observed ratio compared to the expected Mendelian ratio of $Cic^{+/+}$ (25%), $Cic^{L\text{-}KO/+}$
397 (50%), and $Cic^{L\text{-}KO/L\text{-}KO}$ (25%). Each time point is an independent cohort.

398
399

FIGURES

Figure 1. CIC-L and CIC-S show different spatiotemporal expression patterns in the mouse cortex, cerebellum, and lung.

(A) Representative immunoblot of CIC-L, CIC-S, and related proteins in wild-type (WT) mouse cortex, cerebellum and lung lysates at indicated time points. P, postnatal day; w, week-old. Antibody against the C terminus of CIC recognizes both isoforms of CIC.

(B) Quantification of CIC-L and CIC-S protein expression in WT cortex, cerebellum and the lung. Each protein was normalized to total protein, and each timepoint was normalized to P0 cortex (CTX) value of

409   each protein. CIC isoforms were normalized to total CIC in P0 CTX. Data are displayed as mean ± SEM
410   from n = 4 for each time point.

Figure 2. *Cic-L*-KO and *Cic-S*-KO mice were generated using CRISPR.

(A) Gene and protein structure of human CIC-L and CIC-S. ATXN1/1L BD: ATXN1/1L binding site, HMG: HMG box, C1: C1 domain, NLS: Nuclear localization signal

414
415

(B) Immunoblot of CIC-L and CIC-S of cortex lysate of WT, *Cic-S*-KO and *Cic-L*-KO mice at 1 month of age. Vinculin (VCL) as loading control. n =3 per genotype.

416
417

419

420 **Figure 3. *Cic-S*-KO mice display early lethality and lung alveolarization defects.**

421 (A) Weekly body weights of *Cic-S*-KO, *Cic-L*-KO, and their WT littermate controls starting from week 3. n
422 =16 for each genotype, 8 males and 8 females. All data are represented as mean ± SEM. Two-way
423 ANOVA with repeated measures, with Tukey's multiple comparisons. In each case, *, **, ***, and ****
424 denote $p < 0.05$, $p < 0.01$, $p < 0.001$, and $p < 0.0001$ respectively.
425 (B) Kaplan-Meier curves for survival post-weaning (3 weeks) of WT, *Cic-S*-KO and *Cic-L*-KO mice. n = 30,
426 25, and 27 respectively). Mantel-Cox test. **** denote $p < 0.0001$.
427 (C) Left: Phenotypes observed before lethality of *Cic-S*-KO animals (n = 15). Right: Representative
428 photographs of the heads of WT, *Cic-S*-KO and *Cic-L*-KO mice at 9 weeks of age; *Cic-S*-KO mice show
429 dome-shaped heads due to hydrocephalus.

430 (D and E) Lung tissues from *Cic-S*-KO (D) and *Cic-L*-KO (E) mice and their WT littermates were collected at
431 P21 for morphometric analysis. Left: Representative hematoxylin and eosin (H&E)-stained lung sections.
432 Scale bar, 100 μm. Right: Quantification of alveolar airspace by mean linear intercept (MLI). Data are
433 presented as mean ± SEM (n = 3–5 per group). Unpaired t-test. In each case, *, and ns denote $p < 0.05$,
434 and $p > 0.05$, respectively.

Figure 4. *Cic-L*-KO mice display neurobehavioral deficits.

(A-E) Behavior characterization of *Cic-S*-KO, *Cic-L*-KO, and their WT littermate controls was performed starting at 8 weeks of age.

(A) Anxiety-like behavior measured with an elevated plus maze (EPM) at 8-10 weeks of age.

(B) Activity measured with open field assay (OFA) at 9-11 weeks of age.

(C) Motor coordination measured with rotarod assay at 9-11 weeks of age. Data points indicate male and female combined mean ± SEM. (D = Day)

(D and E) Learning and memory assessed with fear conditioning assay (FC) at 19-21 weeks of age.

For (A) and (B), n = 30 (14 males, 16 females) WT, n = 18 *Cic-S*-KO (5 males, 13 females), and n = 23 *Cic-L*-KO (12 males, 11 females) were analyzed. Due to the early lethality of *Cic-S*-KO mice, n = 17 (5 males, 12 females) and n = 15 (4 males, 11 females *Cic-S*-KO mice were analyzed for (C), (D) and (E) respectively. One-way ANOVA with Tukey's multiple comparisons test were used for (A), (B), (D) and (E), two-way ANOVA with repeated measures, with Tukey's multiple comparisons for (C). In each case, *, **, ***, **** and ns denote $p < 0.05$, $p < 0.01$, $p < 0.001$, $p < 0.0001$, and $p > 0.05$, respectively. All data are represented as mean ± SEM.

Figure 5. Differential complex formation of ATXN1 and ATXN1L in *Cic-S*-KO and *Cic-L*-KO in the cortex and the cerebellum

(A-C and D-F) Representative immunoblot of size exclusion chromatography (SEC) of WT, *Cic-L*-KO, and *Cic-S*-KO cortex (A) and cerebellum (D) lysates probed for CIC-L, CIC-S, ATXN1, and ATXN1L. The column void volume (V_0), gel-filtration standards thyroglobulin (669 kDa) and ADH (150 kDa), and

elution volume (ml) of each collected fraction are indicated. Ex, sample extract; open arrow head, CIC-L;
closed arrow head, CIC-S; *, non-specific band.

(B and E) SEC elution profile of each protein of WT, *Cic-L*-KO, and *Cic-S*-KO cortex (B) and cerebellum
(E). The percentage of each protein (mean ± SEM) in each fraction was determined from n = 3–4
independent cortical extracts. Protein abundance in each fraction was quantified by densitometry, with the
signal in each fraction normalized to the total signal across all fractions for each genotype.

(C and F) Quantification of the ratio of ATXN1 and ATXN1L in large to small molecular weight fractions in
the cortex (C) and the cerebellum (F). Left: Ratio of ATXN1 in the large (peak elution fractions 11–13) to
small (peak elution fractions 15–17) complexes, normalized to WT. Right: Ratio of ATXN1L in the large
(peak elution fractions 11–13) to small (peak elution fractions 16–17) complexes, normalized to WT.
Statistical significance was assessed by one-way ANOVA. In each case, *, **, and *** denote $p < 0.05$, p
< 0.01, $p < 0.001$, respectively. All data are represented as mean ± SEM.

Figure 6. The unique domain of CIC-L interacts exclusively with ATXN1, while the unique domain of CIC-S interacts exclusively with ATXN1L.

(A and B) Representative immunoblots of immunoprecipitation (IP) experiments from WT cortex. (A) IP of ATXN1 and co-IPed CIC-L and CIC-S. (B) IP of ATXN1L and co-IPed CIC-L and CIC-S.

(C and D) Representative immunoblots of IP experiments from HEK293T co-transfected with V5-tagged CIC-L or CIC-S, either WT or with an ATXN1/1L binding mutation (W946A or W37A), and Flag-tagged ATXN1 (C) or ATXN1L (D). IP of V5 and co-IPed Flag.

(E and F) Representative immunoblots of IP experiments from HEK293T co-transfected with V5-tagged CIC-L or CIC-L unique N-terminal (CIC-Lunq), and Flag-tagged ATXN1(C) or ATXN1L (D). IP of V5 and co-IPed Flag.

EXPERIMENTAL MODEL DETAILS

Mouse husbandry

All mice were backcrossed to C57BL/6J background. All mice were housed and maintained in the animal facilities at Baylor College of Medicine (BCM) on a 12-hour light-dark cycle. BCM Institutional Animal Care & Use Committee (IACUC) approved all animal care and experimental procedures.

HEK293T cell line

Human embryonic kidney cell T (HEK293T) (ATCC, CRL-3216) were cultured in Dulbecco's modified Eagle's medium (DMEM, Corning, MT10013CV) supplemented with 10% fetal bovine serum (Gibco A4736201) and 1X antibiotic-antimycotic (Gibco, 15240096). Cell cultures were maintained at 37 °C in a humidified 5% CO_2 incubator.

METHOD DETAILS

Generation of *Cic-L*-KO and *Cic-S*-KO mice

Cic-L-KO and *Cic-S*-KO alleles were generated using CRISPR/Cas9 genome editing. For *Cic-L*-KO, two crRNAs targeting exon 2 of *Cic-L*, the first coding exon unique to the CIC-L isoform, were designed (5′-CGCACCTGGCGCACGACTGC-3′ and 5′-TCCAGCCGCACCGACTGCTG-3′; IDT) to generate a large deletion. For Cic-S-KO, a single crRNA targeting exon 1 of Cic-S was designed (5′-ACGCGGGTATCAGGGGCCTG-3′; IDT).

crRNA, tracrRNA, and Cas9 protein were prepared as described previously [22] with modifications to total reaction volume, concentration and delivery method. Final concentrations were 6.6 μM crRNA, 6.6 μM tracrRNA, and 6 μM Cas9 in $T_{10}E_{0.1}$ buffer. For *Cic-L*-KO, each crRNA was diluted to 3.3 μM. The mixture was electroporated into zygotes by the Genetically Engineered Rodent Model (GERM) Core at Baylor College of Medicine (BCM).

Founders were screened by PCR using primers flanking the expected deletion sites (*Cic-L*-KO: 5′-TCCTCCTCTACTGACACGGC-3′, 5′-TAGGACAAGCAGCTCATCACC-3′; *Cic-S*-KO: 5′-GGCCAAGGAGCGGGTTAC-3′, 5′-TCCTAGTGCAGTACCGTCCA-3′). PCR products were TOPO cloned (Invitrogen) and analyzed by Sanger sequencing for frameshift deletions. *Cic-S*-KO founders were selected for

510 deletions eliminating the downstream ATG in exon 1. Founder lines were backcrossed to C57BL/6J WT mice

511 (Jackson Laboratory), and germline transmission was confirmed by sequencing of F1 progeny.

512 Immunoblotting was performed on homozygous KO animals to confirm the loss of the corresponding CIC

513 isoform. One independent founder line was established for each isoform KO and maintained by successive

514 backcrossing (>5 generations) to C57BL/6J prior to experimental use.

Genotyping of *Cic-L*-KO and *Cic-S*-KO mice

Genotyping was performed by PCR using allele-specific primers. For *Cic-L*, WT and KO alleles were

distinguished using two primer sets: *Cic-L*-WT (5′-ACTGGAGAAGGGAGCTGCAC-3′ and 5′-

CTGTCCCTCGAAGTCTCATCG-3′) and *Cic-L*-KO (5′-TCCTCCTCTACTGACACGGC-3′ and 5′-

TAGGACAAGCAGCTCATCACC-3′). For *Cic-S*, a single primer pair (5′-GGCCAAGGAGCGGGTTAC-3′ and

5′-TCCTAGTGCAGTACCGTCCA-3′) was used, and WT versus KO alleles were distinguished based on product

size.

Lung Histopathology

The lungs were harvested for morphometry studies on postnatal day (P) 21, as described before [57]. Both lungs

of the experimental mouse (n=3-5/group) were inflated at 25 cm pressure for at least 10 minutes via tracheal

cannulation with 4% paraformaldehyde. The inflated lungs were then fixed overnight with 4% paraformaldehyde

at room temperature with gentle agitation. The fixed lungs were stored in 70% ethanol in a refrigerator until

further processing for lung histopathology. The lungs were then paraffin-embedded, sliced into 5 µm thin sections,

and stained with H&E. We examined at least five random non-overlapping fields of 10X H&E stained images

from both lungs of each animal to quantify alveolarization by mean linear intercepts (MLIs). To calculate the

MLI, grids of horizontal and vertical lines were placed on the H&E stained lung images, and the number of times

the lines intersected with the lung tissue was counted. The MLI was determined by dividing the total length of the

grid lines by the number of intersects.

Behavioral Assays

All behavioral analyses were conducted during the light phase of a 12-hour light-dark cycle by an experimenter

blinded to mouse genotype. Mice had *ad libitum* access to food and water except during testing. Age-matched

mice were used in all experiments, and wild-type (WT) littermates from both C*ic-S*-KO and *Cic-L*-KO breeding

were pooled as controls. Mice were habituated in the test room for at least 30 min prior to each assay. Tests were
performed under room lighting of 150 lux with background white noise at 60 dB.

Elevated plus maze: Mice (8-10 weeks of age) were placed in the center of an elevated plus maze consisting of
two open arms (25 × 8 cm, with 0.5 cm raised edges) and two closed arms (25 × 8 cm, enclosed by 15 cm high
walls), elevated 50 cm above the floor. Behavior was recorded for 10 min using a camera and ANY-maze video
tracking system (Stoelting, Wood Dale, IL, USA). Measures included time spent, distance travelled and number of
entries in open and closed arms.

Open field assay: Mice (9-11 weeks of age) were placed in the center of a 40 × 40 × 30 cm chamber equipped
with photobeams (AccuScan, Columbus, OH, USA) and allowed to explore for 30 min testing period. Locomotor
activity was automatically analyzed using AccuScan Fusion software (Omnitech). Parameters included total
distance traveled and mean motor speed.

Rotarod: This assay was performed as described previously [22]. Mice (9-11 weeks of age) were placed on a
rotating rod (3 cm diameter, 30 cm length; Ugo Basile) and tested for four consecutive days, with four trials per
day and a minimum 30-min inter-trial interval. Each trial lasted a maximum of 10 min. The rod accelerated
linearly from 5 to 40 rpm over the first 5 min, and then remained at 40 rpm for the final 5 min. Latency to fall was
recorded, with two consecutive passive rotations scored as a fall.

Pole test: Mice (15-17 weeks of age) were placed head-upward at the top of a vertical threaded metal pole. Time
to descend to the base was recorded with a 60 s cutoff. If a mouse turned but fell before reaching the floor, the
time to fall was recorded. Mice unable to turn downward were scored as 60 s.

Conditioned fear: On day 1, mice (19-21 weeks of age) were placed in a conditioning chamber (Actimetrics,
Coulbourn instruments, Wilmette, IL) and presented with two conditioned stimulus (CS)- unconditioned stimulus
(US) pairings, programmed using the FreezeFrame software (Actimetrics). The trial comprised of 2 min of no
sound, followed by 30 s of sound at 80 dB (CS), which was followed by a mild 0.8 mA foot shock for 2 s (US).
The tone-shock pairing was repeated after 2 minutes. Twenty-four hours later, contextual memory was assessed
by returning the mice to the same chamber for 5 min and quantifying freezing behavior. One hour later, cued fear
memory was assessed in a novel chamber by monitoring freezing for 3 min without stimulus, followed by 3 min
with the tone. Freezing was recorded and quantified using FreezeFrame software (Actimetrics).

Three-chamber sociability assay: Partner mice (matched for age, sex, and genetic background) were habituated
to the three-chamber apparatus (length: 54 cm; width: 40 cm; height: 23 cm) divided into three chambers by
removable partitions, for 1 h per day on two consecutive days prior to testing, while placed inside wire cups. On
the test day, following habituation, experimental mice (12-15 weeks of age) were placed in the central chamber

and allowed to freely explore for 10 min. A novel partner mouse was then placed inside a wire cup in either the

left or right chamber, while a novel object was placed inside a wire cup in the opposite chamber. Placement of the

partner mouse was randomized across trials. Experimental mice were allowed to explore for 10 min, and

interaction time with each wire cup was manually scored.

RNA extraction

Cortical and cerebellar tissues from 12-week-old mice, and lung tissues from P6 mice, were dissected, flash-

frozen in liquid nitrogen, and stored at −80 °C until further processing. Total RNA was isolated using the

miRNeasy Mini Kit (Qiagen, 217004) according to the manufacturer's instructions. RNA concentration and purity

were measured using a NanoDrop 1000 (Thermo Fisher).

RNA sequencing and analysis

RNA samples were submitted to Azenta Life Sciences for RNA integrity assessment, library preparation, and

sequencing. Libraries were generated following polyA selection for mRNA and sequenced on Illumina NovaSeq

or HiSeq platforms, yielding ~ 30 million 150-bp paired-end reads per sample.

For each sample, the raw reads from multiple lanes were merged, and then were aligned to the Mus musculus

reference transcriptome (GRCm38.p6, GENCODE M18) using STAR v2.7 with "-quantMode GeneCounts" to

generate raw read counts [58].

Differential gene expression analysis was performed using DESeq2 v1.44.0 [59]. Principal component analysis

(PCA) was performed on the top 500 most variable genes. Genes were considered differentially expressed if

adjusted $p < 0.05$ and |log2 fold change| > 0.25.

Previously published lung P6 microarray data of *Atxn1*-KO and *Atxn1l*-KO were obtained from *Lee et al., 2011*.

Differentially expressed genes (DEGs) reported in that dataset were used for comparison; the original publication

defined DEGs using thresholds of $p < 0.05$ and |fold change| > 1.2. For *Atxn1*-KO RNA-seq data from cortex and

cerebellum (*Xhako et al., unpublished data*), read count data were processed with the same DESeq2 pipeline, and

limma v3.60.6 [60] was used for batch correction prior to PCA.

Gene set enrichment analysis (GSEA) for Gene Ontology (GO) terms was conducted on all ranked genes using

fgsea v1.30.0, with neuro/glia-related GO terms selected based on keywords including "neuron," "neuro,"

"nervous," "synapse," "axon," "glia," "glial," "astrocyte," "oligodendrocyte," "microglia," "myelin," and "brain."

[61] Overlapping differentially expressed genes were analyzed for GO term enrichment using clusterProfiler

v4.12.6 [62], with the background universe defined as all genes detected in the RNA-seq dataset for the
corresponding tissue.

Protein extraction for size exclusion chromatography

Cortical or cerebellar tissues were collected from two 2–3-month-old mice per genotype and pooled per biological
replicate. Tissues were homogenized in TX buffer (50 mM Tris, pH 8.0; 75 mM NaCl; 0.5% Triton X-100;
Sigma) supplemented with freshly added protease inhibitor (GenDEPOT, P3100-020) and phosphatase inhibitors
(GenDEPOT, P3200-020) using a Dounce homogenizer. Homogenates were extracted at 4 °C for 20 min with
rotation and clarified by centrifugation at 20,000 × g for 10 min. Supernatants were re-centrifuged, filtered
through 0.45 μm centrifugal filters (Millipore), and protein concentrations were determined by BCA assay
(Pierce).

Size exclusion chromatography

Gel filtration chromatography was performed on a Superose 6 10/300 GL column (Cytiva) using an AKTA Pure
system at 4 °C. The column was equilibrated in 50 mM Tris (pH 8.0), 50 mM NaCl, 0.1% Triton X-100. Column
calibration was performed with thyroglobulin (667 kDa), apoferritin (440 kDa), alcohol dehydrogenase (150
kDa), BSA (66 kDa), and cytochrome C (14.3 kDa). Fractions (1 mL) were collected, supplemented with protease
inhibitor (GenDEPOT, P3100-020) and phosphatase inhibitors (GenDEPOT, P3200-020), and analyzed by SDS-
PAGE on NuPAGE 3–8% Tris-Acetate gels (Invitrogen).

Cloning

The hCIC-S cDNA was obtained from a previously described plasmid [16]. The unique N-terminal region of
hCIC-L was synthesized (GenScript) and cloned into a PU57 backbone. Because direct amplification of the ~4.8
kb CIC-Shared region was hindered by high GC content at the 3′ end, a two-step cloning strategy was employed:
the 5′ and 3′ CIC fragments were first assembled into pENTR223 containing a synthetic multiple cloning site with
an AvrII site, followed by insertion of the ~4 kb central CIC fragment by PCR to generate full-length CIC-Shared.

A W37A point mutation (W37A in CIC-S, corresponding to W946A in CIC-L) was introduced into hCIC-Shared
in pENTR223 using gene blocks and Gibson assembly (New England Biolabs (NEB), E2621L) after XhoI (5′
end) and StuI (N-terminal) digestion. CIC-L and CIC-S entry clones were generated by Gibson assembly of the
respective unique 5′ regions into the CIC-Shared backbone, followed by addition of C-terminal V5 tags. The

hCIC-L–unique sequence was cloned into pENTR223 with an N-terminal V5 tag and a stop codon after amino

acid 931. All entry clones were recombined into pcDNA3.1-DEST (derived from pcDNA3.1-3×FLAG-DEST,

with stop codons preventing FLAG expression) using Gateway LR recombination (Thermo Fisher).

The hATXN1L cDNA (Horizon Discovery) was cloned with an added N-terminal FLAG tag into pLV-EF1α-

IRES-Puro [63] using PCR amplification and Gibson assembly. The FLAG-hATXN1 coding sequence was

obtained from the previously described pcDNA1-ATXN1 plasmid [18] and cloned into pcDNA3.1 by Gibson

assembly.

Transfection and co-immunoprecipitation in HEK293T cells

HEK293T cells were transfected with the indicated plasmids using TransIT-293 transfection reagent (Mirus Bio)

according to the manufacturer's instructions and incubated for 48–72 h. Cells were washed with PBS and either

processed immediately or stored at –80 °C. Frozen cells were thawed on ice and lysed in cold lysis buffer (75 mM

NaCl, 5 mM MgCl$_2$, 50 mM Tris-HCl (pH 8.0), 0.5% Triton X-100, with protease and phosphatase inhibitors) for

30–60 min at 4 °C with rotation. Lysates were cleared by centrifugation at 20,000 × g for 20 min at 4 °C.

For co-immunoprecipitation, lysates from one well of a 6-well plate were incubated with 1 µg V5 antibody

(Invitrogen) pre-bound to 15 µL Protein G Dynabeads (Fisher Scientific, Cat. #10-009-D) for 2–4 h at 4 °C with

rotation. Beads were washed four times with lysis buffer, and bound proteins were eluted in 1× NuPAGE Sample

Reducing Agent and NuPAGE LDS Sample Buffer (Invitrogen), boiled for 10 min, and analyzed by SDS-PAGE

on NuPAGE 3–8% Tris-Acetate gels (Invitrogen).

Immunoprecipitation in mouse brain tissue

Cortices from 12-week-old WT mice were homogenized in either cold NEMT buffer (50 mM Tris-HCl pH 7.5,

150 mM NaCl, 1 mM EDTA, 0.5% NP-40) or lysis buffer (75 mM NaCl, 5 mM MgCl$_2$, 50 mM Tris-HCl pH 8.0,

0.5% Triton X-100), each supplemented with freshly added protease inhibitor (GenDEPOT, P3100-020) and

phosphatase inhibitors (GenDEPOT, P3200-020). Tissue homogenization was performed using an electric pestle

(handheld polytron, WPR, 47747-370), followed by mechanical shearing through a 27G syringe. Lysates were

incubated on ice for 20 min and clarified by centrifugation at 20,000 × g for 20 min at 4 °C. Ten percent of the

supernatant was reserved as input, and the remaining lysate was precleared with Dynabeads Protein G (Fisher

Scientific, 10-009-D) for 30 min at 4 °C. Precleared lysates were incubated for 2–3 h at 4 °C with 1–2 µg of

antibody—anti-ATXN1 (534, in-house [64]), anti-ATXN1L (881, in-house), or mouse IgG control—conjugated to

30 µL Dynabeads Protein G. Beads were washed four times with the corresponding lysis buffer and final wash

with PBS. Bound proteins were eluted in 1× NuPAGE Sample Reducing Agent and NuPAGE LDS Sample Buffer
(Invitrogen), boiled for 10 min, and analyzed by SDS-PAGE on NuPAGE 3–8% Tris-Acetate gels (Invitrogen).

For *Atxn1*-KO experiments, cortex, hippocampus, cerebellum, and brainstem were dissected from 4-week-old WT
and *Atxn1*-KO mice and processed using the ATXN1 immunoprecipitation protocol described above.

Protein extraction for immunoblotting

Brain and lung tissues were collected at postnatal day zero (P0), P6, P21, and 12 weeks and snap-frozen in liquid
nitrogen and kept in -80°C until processing. For protein extraction, frozen tissues were homogenized in cold lysis
buffer (75 mM NaCl, 5 mM MgCl$_2$, 50 mM Tris pH 8.0, 0.5% Triton X-100, supplemented with freshly added
protease inhibitor (GenDEPOT, P3100-020) and phosphatase inhibitors (GenDEPOT, P3200-020) using an
electric pestle (handheld polytron, WPR, 47747-370), followed by mechanical shearing through a 27G syringe.
Lysates were incubated on ice for 20 min and centrifuged at 20,000 × g for 20 min at 4 °C. Protein concentrations
in the supernatant were measured using the Pierce BCA Protein Assay Kit (Thermo Fisher Scientific), and
samples were normalized to equal concentrations.

Western blot analysis

For immunoblotting, samples were mixed with NuPAGE Sample Reducing Agent and NuPAGE LDS Sample
Buffer (Invitrogen), boiled for 10 min, and run on NuPAGE 3–8% Tris-Acetate gels (Invitrogen). Proteins were
transferred to nitrocellulose membranes (Amersham, #10600004) using transfer buffer containing 10% methanol.
Membranes were blocked with either 5% skim milk in Tris-buffered saline with 0.1% Tween® 20 (TBST) or
Intercept (TBS) Blocking Buffer (LI-COR Biosciences) for 1 h at room temperature and incubated overnight at
4 °C with primary antibodies diluted in 3% BSA in TBST. The following primary antibodies were used: anti-CIC
(rabbit, in-house, 1:1,000), anti-ATXN1[5] (rabbit, in-house, 11750VII, 1:2,000), anti-ATXN1L (mouse, 674, in-
house, 1:1,000), anti-Vinculin (mouse, Sigma-Aldrich, 1:5,000), anti-Flag (Sigma-Aldrich, 1:4000), and anti-V5
(Invitrogen, 1:2,000).

After washing in TBST, membranes were incubated for 1 h at room temperature with either HRP-conjugated
secondary antibodies (Jackson ImmunoResearch, 1:10,000 in 5% milk/TBST) or fluorophore-conjugated
secondary antibodies (LI-COR Biosciences, 1:10,000 in Intercept Blocking Buffer). HRP blots were developed
using ECL (Cytiva, RPN2236) or SuperSignal Femto (Thermo Fisher Scientific, 34095) and imaged with a GE
Amersham Imager 680 or ChemiDoc MP Imaging System or LI-COR Odyssey M system. Fluorescent blots were
imaged directly using the LI-COR Odyssey M system.

Statistical analysis

All statistical analyses were performed using GraphPad Prism 9.0, except for RNA-seq analyses. Survival of animals was compared to the expected Mendelian ratio using a chi-square test. Comparisons between two groups were performed using unpaired t-tests. Comparisons involving more than two groups were analyzed using one-way or two-way ANOVA, followed by Tukey's multiple comparisons test. Repeated-measures data were analyzed using two-way repeated-measures ANOVA with Tukey's multiple comparisons test. Survival curves were analyzed using the Mantel–Cox test. All reported p values are from post hoc comparisons. Data are presented as mean ± SEM. Statistical significance was defined as $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***), and $p < 0.0001$ (****).

RNA-seq analyses were performed in R using DESeq2, as described above. Fisher's exact test was used to calculate odds ratios and p-values for gene set overlaps, and pairwise comparisons between genotypes for box plots were performed using two-tailed t-tests with Benjamini–Hochberg correction for multiple comparisons.

Sample sizes were based on prior studies and standard practice in the field. Biological replicates refer to independent animals. Experimental analyses were performed in a blinded manner whenever possible.

SUPPLEMENTARY FIGURES

Figure S1. CIC-L, CIC-S, ATXN1, and ATXN1L expression in different brain regions.

(A) Representative immunoblot of CIC-L, CIC-S, ATXN1, and ATXN1L in 12-week-old WT mouse brain region lysates at the indicated brain regions (CTX: cortex, HP: hippocampus, STR: striatum, CB: cerebellum, BS: brainstem).

(B and C) Quantification of each protein's expression. (C) Total CIC combining CIC-L and CIC-S. Protein levels were normalized to vinculin (VCL), and values at each region were normalized to the cortex value for the respective protein.

(D) Total CIC was normalized to ATXN1 for each region. Mean of each region compared to the normalized cortical expression of each protein. One-way ANOVA. In each case, *, **, ***, and **** denote $p < 0.05$, $p < 0.01$, $p < 0.001$, and $p < 0.0001$ respectively. All data are represented as mean $\pm$ SEM from n = 4 per region.

Figure S2. *Cic-S*-KO and *Cic-L*-KO were generated by frameshift indels.

(A) Schematic of *Cic-L*-KO and *Cic-S*-KO allele generation. *Cic-L*-KO was generated by a 1975bp deletion of the first coding exon of *Cic-L* (*Cic-L* exon2). The deletion is downstream of the start codon and leads to a frameshift of *Cic-L*. *Cic-S*-KO was generated by an indel that spans the *Cic-S* unique exon1 and the downstream intron. The deletion is downstream of the start codon and spans 50bp of the exon including the downstream start codon and 14bp of the intron with a one bp insertion. Overall, this leads to a frameshift of *Cic-S*. The start codon is marked with red and the downstream ATG is marked with black. The PCR primers for genotyping are marked as arrows.

(B) Representative gel image of genotyping PCR reaction. *Cic-L* KO allele (383bp), *Cic-L* KO allele (403bp), *Cic-S* WT allele (633bp), and *Cic-S* KO allele (570bp).

(C) Sanger sequencing confirming a frameshift indel of *Cic-S*-KO (top), and *Cic-L*-KO (bottom) genomic DNA.

Figure S3. *Cic-L*-KO shows motor deficits in the pole test and more time spent with partner during a social interaction assay.

(A) Fine motor behavior measured with pole test at 15-17 weeks of age.

(B) Social behavior measured with the three-chamber assay at 12-15 weeks of age.

For (A) n = 17 (8 males, 9 females) WT, n = 8 *Cic-S*-KO (1 males, 7 females), and n =16 *Cic-L*-KO (9 males, 7 females) were analyzed with one-way ANOVA with Tukey's multiple comparisons. For (B) n = 27 (12 males, 15 females) WT, n = 10 *Cic-S*-KO (3 males, 7 females), and n =14 *Cic-L*-KO (5 males, 9 females) were analyzed with two-way ANOVA. In each case, ***, **** and ns denote $p < 0.001$, $p < 0.0001$, and $p > 0.05$, respectively. All data are represented as mean ± SEM.

Figure S4. *Cic-S*-KO shows greater number and magnitude of DEGs in the lungs compared to *Cic-L*-KO
749
DEGs.
750

(A) Principal component analysis (PCA) of RNA-seq data from P6 lungs (WT, *Cic-S*-KO [SKO], *Cic-L*-KO
751
[LKO]) using the top 500 most variable genes.
752

(B) Left: Venn diagram showing overlap of differentially expressed genes (DEGs; padj < 0.05 & |log2 fold
753
change (log2FC)| >0.25) between *Cic-S*-KO and *Cic-L*-KO. Odds ratio (OR) and Fisher's exact *p*-value
754

755   are indicated. A total of 15473 genes were analyzed. Right: Scatterplot comparing log2FC of DEGs *Cic-*
756   *S*-KO versus *Cic-L*-KO. Pearson correlation coefficient (r) and R^2 are shown. Overlapping DEGs marked
757   in purple.
758 (C and D) Top gene ontology (GO) enrichment terms for biological processes of *Cic-S*-KO (C) and *Cic-L*-KO
759   (D).

Figure S5. *Cic-S*-KO DEGs show greater overlap with *Atxn1l*-KO DEGs than *Cic-L*-KO DEGs in the lungs.

(A and B) Left: Venn diagrams showing overlap between DEGs in *Cic-S*-KO [SKO] (A) or *Cic-L*-KO [LKO] (B) and DEGs from *Atxn1l*-KO lungs reported in *Lee et al., 2011*. Odds ratio (OR) and Fisher's exact test p-value are indicated. A total of 15473 genes were analyzed. Right: Scatterplot comparing log2FC of DEGs of *Cic-S*-KO (A) or *Cic-L*-KO (B) versus *Atxn1l*-KO. Pearson correlation coefficient (r) and R^2 are shown. Overlapping DEGs marked in blue (A) or red (B).

(C and D) Left: Venn diagrams showing overlap between DEGs in *Cic-S*-KO (A) or *Cic-L*-KO (B) and DEGs from *Atxn1*-KO lungs reported in *Lee et al., 2011*. OR and Fisher's exact *p*-value are indicated. A total of

15473 genes were analyzed. Right: Scatterplot comparing log2FC of DEGs of *Cic-S*-KO (C) or *Cic-L*-KO

(D) versus *Atxn1*-KO. Pearson correlation coefficient (r) and R^2 are shown. Overlapping DEGs marked in

brown (C) or navy (D).

(E) Boxplot of *Etv-Mmp* genes. Statistical analysis was performed using pairwise two-tailed t-tests with

Benjamini–Hochberg correction for multiple comparisons. In each case, *, **, ***, and **** denote $p <$

0.05, $p < 0.01$, $p < 0.001$, and $p < 0.0001$, respectively. Non-significant comparisons ($p > 0.05$) are not

shown.

(F) Enrichment of GO of extracellular matrix assembly and disassembly terms of *Cic-S*-KO (Left) or *Cic-L*-

KO (Right) gene set enrichment analysis (GSEA).

Figure S6. *Cic-L*-KO shows greater number and magnitude of DEGs than *Cic-S*-KO DEGs in the cortex.

(A) Principal component analysis (PCA) of RNA-seq data from 12-week-old cortex of WT, *Cic-S*-KO [SKO],
Cic-L-KO [LKO] of this study and 10-week-old cortex from *Xhako et al., in preparation. Atxn1*-KO
[AKO] and WT using the top 500 most variable genes.

(B) Left: Venn diagram showing overlap of DEGs between *Cic-S*-KO and *Cic-L*-KO. OR and Fisher's exact
p-value are indicated. A total of 15354 genes were analyzed. Right: Scatterplot comparing log2FC of
DEGs *Cic-S*-KO versus *Cic-L*-KO. Pearson correlation coefficient (r) and R² are shown. Overlapping
DEGs marked in purple.

790   (C and D) Top gene ontology (GO) terms of all biological processes (C) and neuro- or glia- related terms (D)

791   of *Cic-S*-KO gene set enrichment analysis (GSEA).

792   (E and F) Top enriched GO terms of all biological process (E) and neuro- or glia- related -terms (F) of *Cic-L*-

793   KO GSEA.

Figure S7. *Cic-L*-KO DEGs show greater overlap with *Atxn1*-KO DEGs than *Cic-S*-KO in the cortex.

(A and B) Left: Venn diagrams showing overlap between DEGs in *Cic-S*-KO [SKO] (A) or *Cic-L*-KO [LKO] (B) and DEGs (padj < 0.05 & |log2FC| >0.25) from *Atxn1*-KO cortex from *Xhako et al., in preparation*. OR and Fisher's exact *p*-value are indicated. A total of 15290 genes were analyzed. Right: Scatterplot comparing log2FC of DEGs of *Cic-S*-KO (A) or *Cic-L*-KO (B) versus *Atxn1*-KO. Pearson correlation coefficient (r) and R^2 are shown. Overlapping DEGs marked in brown (A) or navy (B).

(C and D) Top enriched gene ontology terms of biological process of the DEGs that overlap and change in the same direction between *Atxn1*-KO and *Cic-S*-KO (C), or *Atxn1*-KO and *Cic-L*-KO (D).

(E) Boxplot of *Etv1/4/5-Bace1*. Two-tailed t-tests with Benjamini–Hochberg correction for multiple comparisons. In each case, **** denotes p < 0.0001. Non-significant comparisons (p > 0.05) are not shown.

Figure S8. *Cic-L*-**KO shows greater number and magnitude of DEGs in the cerebellum than** *Cic-S*-**KO.**

(A) Principal component analysis (PCA) of RNA-seq data from 12-week-old cerebellum of WT, *Cic-S*-KO
[SKO], *Cic-L*-KO [LKO] of this study and 10-week-old cerebellum from *Xhako et al., in preparation*.
Atxn1-KO [AKO] and WT using the top 500 most variable genes.
(B) Left: Venn diagram showing overlap of DEGs between *Cic-S*-KO and *Cic-L*-KO. OR and Fisher's exact
p-value are indicated. A total of 15354 genes were analyzed. Right: Scatterplot comparing log2FC of
DEGs *Cic-S*-KO versus *Cic-L*-KO. Pearson correlation coefficient (r) and R^2 are shown. Overlapping
DEGs marked in purple.
(C and D) Top gene ontology (GO) terms of all biological process (C) and neuro- or glia- related terms (D) of
Cic-S-KO gene set enrichment analysis (GSEA).
(E and F) Top enriched GO terms of all biological process (E) and neuro- or glia- related -terms (F) of *Cic-L*-
KO GSEA.

Figure S9. *Cic-S*-KO DEGs show greater overlap in the cerebellum with *Atxn1l*-KO than *Cic-L*-KO DEGs.

(A and B) Left: Venn diagrams showing overlap between DEGs in *Cic-S*-KO [SKO] (A) or *Cic-L*-KO [LKO] (B) and DEGs (padj < 0.05 & |log2FC| >0.25) from *Atxn1*-KO cerebellum from *Xhako et al., in preparation*. OR and Fisher's exact *p*-value are indicated. A total of 15290 genes were analyzed. Right: Scatterplot comparing log2FC of DEGs of *Cic-S*-KO (A) or *Cic-L*-KO (B) versus *Atxn1*-KO. Pearson correlation coefficient (r) and R² are shown. Overlapping DEGs marked in brown (A) or navy (B).

(C) Boxplot of *Etv1/4/5-Bace1*. Two-tailed t-tests with Benjamini–Hochberg correction for multiple comparisons. In each case, *, **, and ***, denote p < 0.05, p < 0.01, and p < 0.001, respectively. Non-significant comparisons (p > 0.05) are not shown.

Figure S10. CIC isoforms form different complexes with ATXN1 and ATXN1L in the WT cortex and cerebellum (related to Figure 5).

(A-D) SEC elution profile of each protein of WT cortex and cerebellum. The percentage of each protein (mean ± SEM) in each fraction was determined from n = 3–4 independent cortical extracts. Protein abundance in each fraction was quantified by densitometry, with the signal in each fraction normalized to the total signal across all fractions for both regions. Statistical significance was assessed by two-way ANOVA. In each case, *, **, ***, and **** denote $p < 0.05$, $p < 0.01$, $p < 0.001$, $p < 0.0001$ respectively. All data are represented as mean ± SEM.

Figure S11. *Cic, Atxn1,* and *Atxn1l* expression in mouse cerebellum at P60.

Figure S12. CIC-L is more dependent on ATXN1 than CIC-S in all brain regions

(A-D) Representative western blot images (top) and quantification of input (bottom) of ATXN1, ATXN1L and CIC expression in WT mice and Atxn1-KO mice of four different brain regions. Protein levels were normalized to VCL, then relative levels were normalized to WT expression.

850   (E) Binding preference of CIC-L/CIC-S in ATXN1 IP normalized to the ratio in input.

851   CTX, cortex; HP, hippocampus; CB, cerebellum; BS, brainstem; Data=mean $\pm$ SEM; n = 4 per region. One-
852   way ANOVA; *$p<0.05$; **$p<0.01$; ***$p<0.001$; ***$p<0.0001$.

REFERENCES

1. Fu, H., et al., *Selective vulnerability in neurodegenerative diseases.* Nature Neuroscience 2018 21:10, 2018. **21**(10).
2. Kampmann, M., *Molecular and cellular mechanisms of selective vulnerability in neurodegenerative diseases.* Nat Rev Neurosci, 2024. **25**(5): p. 351-371.
3. Orr, H.T., et al., *Expansion of an unstable trinucleotide CAG repeat in spinocerebellar ataxia type 1.* Nat Genet, 1993. **4**(3): p. 221-6.
4. Banfi, S., et al., *Identification and characterization of the gene causing type 1 spinocerebellar ataxia.* Nat Genet, 1994. **7**(4): p. 513-20.
5. Servadio, A., et al., *Expression analysis of the ataxin-1 protein in tissues from normal and spinocerebellar ataxia type 1 individuals.* Nat Genet, 1995. **10**(1): p. 94-8.
6. Zoghbi, H.Y. and H.T. Orr, *Spinocerebellar ataxia type 1.* Semin Cell Biol, 1995. **6**(1): p. 29-35.
7. Matilla, A., et al., *Mice lacking ataxin-1 display learning deficits and decreased hippocampal paired-pulse facilitation.* J Neurosci, 1998. **18**(14): p. 5508-16.
8. Crespo-Barreto, J., et al., *Partial loss of ataxin-1 function contributes to transcriptional dysregulation in spinocerebellar ataxia type 1 pathogenesis.* PLoS Genet, 2010. **6**(7): p. e1001021.
9. Asher, M., et al., *Cerebellar contribution to the cognitive alterations in SCA1: evidence from mouse models.* Hum Mol Genet, 2020. **29**(1): p. 117-131.
10. Swaminathan, S., et al., *Genomic Copy Number Analysis in Alzheimer's Disease and Mild Cognitive Impairment: An ADNI Study.* Int J Alzheimers Dis, 2011. **2011**: p. 729478.
11. Swaminathan, S., et al., *Analysis of copy number variation in Alzheimer's disease: the NIALOAD/NCRAD Family Study.* Curr Alzheimer Res, 2012. **9**(7): p. 801-14.
12. Suh, J., et al., *Loss of Ataxin-1 Potentiates Alzheimer's Pathogenesis by Elevating Cerebral BACE1 Transcription.* Cell, 2019. **178**(5): p. 1159-1175 e17.
13. Lee, Y., et al., *ATXN1 protein family and CIC regulate extracellular matrix remodeling and lung alveolarization.* Dev Cell, 2011. **21**(4): p. 746-57.
14. Lu, H.C., et al., *Disruption of the ATXN1-CIC complex causes a spectrum of neurobehavioral phenotypes in mice and humans.* Nat Genet, 2017. **49**(4): p. 527-536.
15. Wong, D., et al., *TRIM25 promotes Capicua degradation independently of ERK in the absence of ATXN1L.* BMC Biol, 2020. **18**(1): p. 154.
16. Lam, Y.C., et al., *ATAXIN-1 interacts with the repressor Capicua in its native complex to cause SCA1 neuropathology.* Cell, 2006. **127**(7): p. 1335-47.
17. Mizutani, A., et al., *Boat, an AXH domain protein, suppresses the cytotoxicity of mutant ataxin-1.* EMBO J, 2005. **24**(18): p. 3339-51.
18. Kim, E., et al., *Structural basis of protein complex formation and reconfiguration by polyglutamine disease protein Ataxin-1 and Capicua.* Genes Dev, 2013. **27**(6): p. 590-5.
19. Bowman, A.B., et al., *Duplication of Atxn1l suppresses SCA1 neuropathology by decreasing incorporation of polyglutamine-expanded ataxin-1 into native complexes.* Nat Genet, 2007. **39**(3): p. 373-9.
20. Kim, E., et al., *Deficiency of Capicua disrupts bile acid homeostasis.* Sci Rep, 2015. **5**: p. 8272.
21. Fryer, J.D., et al., *Exercise and genetic rescue of SCA1 via the transcriptional repressor Capicua.* Science, 2011. **334**(6056): p. 690-3.
22. Coffin, S.L., et al., *Disruption of the ATXN1-CIC complex reveals the role of additional nuclear ATXN1 interactors in spinocerebellar ataxia type 1.* Neuron, 2023. **111**(4): p. 481-492 e8.
23. Rousseaux, M.W.C., et al., *ATXN1-CIC Complex Is the Primary Driver of Cerebellar Pathology in Spinocerebellar Ataxia Type 1 through a Gain-of-Function Mechanism.* Neuron, 2018. **97**(6): p. 1235-1243 e5.
24. Simon-Carrasco, L., et al., *Inactivation of Capicua in adult mice causes T-cell lymphoblastic lymphoma.* Genes Dev, 2017. **31**(14): p. 1456-1468.
25. Lee, Y., *Regulation and function of capicua in mammals.* Exp Mol Med, 2020. **52**(4): p. 531-537.

26. Jimenez, G., S.Y. Shvartsman, and Z. Paroush, *The Capicua repressor--a general sensor of RTK signaling*
in development and disease. J Cell Sci, 2012. **125**(Pt 6): p. 1383-91.
27. Hwang, I., et al., *CIC is a critical regulator of neuronal differentiation.* JCI Insight, 2020. **5**(9).
28. Ahmad, S.T., et al., *Capicua regulates neural stem cell proliferation and lineage specification through*
control of Ets factors. Nat Commun, 2019. **10**(1): p. 2000.
29. Tan, Q. and H.Y. Zoghbi, *Mouse models as a tool for discovering new neurological diseases.* Neurobiol
Learn Mem, 2019. **165**: p. 106902.
30. Subramanian, A., et al., *Gene set enrichment analysis: A knowledge-based approach for interpreting*
genome-wide expression profiles. Proceedings of the National Academy of Sciences, 2005. **102**(43).
31. Dissanayake, K., et al., *ERK/p90(RSK)/14-3-3 signalling has an impact on expression of PEA3 Ets*
transcription factors via the transcriptional repressor capicua. Biochem J, 2011. **433**(3): p. 515-25.
32. Weissmann, S., et al., *The Tumor Suppressor CIC Directly Regulates MAPK Pathway Genes via Histone*
Deacetylation. Cancer Res, 2018. **78**(15): p. 4114-4125.
33. Kozareva, V., et al., *A transcriptomic atlas of mouse cerebellar cortex comprehensively defines cell types.*
Nature, 2021. **598**(7879): p. 214-219.
34. Yang, X., et al., *Widespread Expansion of Protein Interaction Capabilities by Alternative Splicing.* Cell,
2016. **164**(4): p. 805-17.
35. Fores, M., et al., *Origins of context-dependent gene repression by capicua.* PLoS Genet, 2015. **11**(1): p.
e1004902.
36. Rodriguez-Munoz, L., et al., *Noncanonical function of Capicua as a growth termination signal in*
Drosophila oogenesis. Proc Natl Acad Sci U S A, 2022. **119**(31): p. e2123467119.
37. Sharma, S., et al., *Novel CIC variants identified in individuals with neurodevelopmental phenotypes.* Hum
Mutat, 2022. **43**(7): p. 889-899.
38. Bertram, L., et al., *Genome-wide association analysis reveals putative Alzheimer's disease susceptibility*
loci in addition to APOE. Am J Hum Genet, 2008. **83**(5): p. 623-32.
39. Bettens, K., et al., *Follow-up study of susceptibility loci for Alzheimer's disease and onset age identified*
by genome-wide association. J Alzheimers Dis, 2010. **19**(4): p. 1169-75.
40. Kjer-Hansen, P., T.G. Phan, and R.J. Weatheritt, *Protein isoform-centric therapeutics: expanding targets*
and increasing specificity. Nat Rev Drug Discov, 2024. **23**(10): p. 759-779.
41. Friedrich, J., et al., *Antisense oligonucleotide-mediated ataxin-1 reduction prolongs survival in SCA1*
mice and reveals disease-associated transcriptome profiles. JCI Insight, 2018. **3**(21).
42. Jafar-Nejad, P., et al., *Regional rescue of spinocerebellar ataxia type 1 phenotypes by 14-3-3epsilon*
haploinsufficiency in mice underscores complex pathogenicity in neurodegeneration. Proc Natl Acad Sci
U S A, 2011. **108**(5): p. 2142-7.
43. Tsuda, H., et al., *The AXH domain of Ataxin-1 mediates neurodegeneration through its interaction with*
Gfi-1/Senseless proteins. Cell, 2005. **122**(4): p. 633-44.
44. Lim, J., et al., *Opposing effects of polyglutamine expansion on native protein complexes contribute to*
SCA1. Nature, 2008. **452**(7188): p. 713-8.
45. Serra, H.G., et al., *RORalpha-mediated Purkinje cell development determines disease severity in adult*
SCA1 mice. Cell, 2006. **127**(4): p. 697-708.
46. Chen, H.K., et al., *Interaction of Akt-phosphorylated ataxin-1 with 14-3-3 mediates neurodegeneration in*
spinocerebellar ataxia type 1. Cell, 2003. **113**(4): p. 457-68.
47. Kang, J., et al., *The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface*
receptor. Nature, 1987. **325**(6106): p. 733-6.
48. Ponte, P., et al., *A new A4 amyloid mRNA contains a domain homologous to serine proteinase inhibitors.*
Nature, 1988. **331**(6156): p. 525-7.
49. Tanzi, R.E., et al., *Protease inhibitor domain encoded by an amyloid protein precursor mRNA associated*
with Alzheimer's disease. Nature, 1988. **331**(6156): p. 528-30.
50. Kitaguchi, N., et al., *Novel precursor of Alzheimer's disease amyloid protein shows protease inhibitory*
activity. Nature, 1988. **331**(6156): p. 530-2.

51.　Wasco, W., et al., *Identification of a mouse brain cDNA that encodes a protein related to the Alzheimer*
disease-associated amyloid beta protein precursor. Proc Natl Acad Sci U S A, 1992. **89**(22): p. 10758-62.
52.　Wasco, W., et al., *Isolation and characterization of APLP2 encoding a homologue of the Alzheimer's*
associated amyloid beta protein precursor. Nat Genet, 1993. **5**(1): p. 95-100.
53.　Sprecher, C.A., et al., *Molecular cloning of the cDNA for a human amyloid precursor protein homolog:*
evidence for a multigene family. Biochemistry, 1993. **32**(17): p. 4481-6.
54.　Chung, D.C., et al., *Cellular and pathological heterogeneity of primary tauopathies.* Mol Neurodegener,
2021. **16**(1): p. 57.
55.　Wang, Y. and E. Mandelkow, *Tau in physiology and pathology.* Nature Reviews Neuroscience, 2016.
17(1): p. 22-35.
56.　Rhinn, H., et al., *Alternative α-synuclein transcript usage as a convergent mechanism in Parkinson's*
disease pathology. Nat Commun, 2012. **3**: p. 1084.
57.　Shrestha, A.K., et al., *Consequences of early postnatal lipopolysaccharide exposure on developing lungs*
in mice. Am J Physiol Lung Cell Mol Physiol, 2019. **316**(1): p. L229-l244.
58.　Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1): p. 15-21.
59.　Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for RNA-seq*
data with DESeq2. Genome Biol, 2014. **15**(12): p. 550.
60.　Ritchie, M.E., et al., *limma powers differential expression analyses for RNA-sequencing and microarray*
studies. Nucleic Acids Res, 2015. **43**(7): p. e47.
61.　Korotkevich, G., V. Sukhov, and A. Sergushichev, *Fast gene set enrichment analysis.* bioRxiv, 2019: p.
060012.
62.　Xu, S., et al., *Using clusterProfiler to characterize multiomics data.* Nat Protoc, 2024. **19**(11): p. 3292-
3320.
63.　Hayer, A., et al., *Engulfed cadherin fingers are polarized junctional structures between collectively*
migrating endothelial cells. Nat Cell Biol, 2016. **18**(12): p. 1311-1323.
64.　Gennarino, V.A., et al., *A Mild PUM1 Mutation Is Associated with Adult-Onset Ataxia, whereas*
Haploinsufficiency Causes Developmental Delay and Seizures. Cell, 2018. **172**(5): p. 924-936 e11.