Atomistic TCR-ligand interactions instruct memory T-cell differentiation

Aoi Akitsu[1,2,3], Kemin Tan[4], Robert J. Mallis[1,2,3,5], Matthew A. Booker[6], Jonathan S. Duke-Cohan[1,2,3], Kristine N. Brazin[1,2,3], Evan H. Kirkpatrick[7], Andrew N. Parkins[1,2,3], Sarita Aryal[1,2,3], Vincenzo Cinella[1,2], Jonathan J. Lee[1,2], Kaveri I. Uberoy[1,2], Jenna K. Koenig[1,2], Matthew Biddle[1,2], Cameron M. Messier[8], Patrick H. Lizotte[8], Michael Y. Tolstorukov[6], Wonmuk Hwang[9,10], Matthew J. Lang[7,11]*, Ellis L. Reinherz[1,2,3]*

[1]Laboratory of Immunobiology, Dana-Farber Cancer Institute; Boston, MA 02115, USA.

[2]Department of Medical Oncology, Dana-Farber Cancer Institute; Boston, MA 02115, USA.

[3]Department of Medicine, Harvard Medical School; Boston; MA 02115, USA.

[4]Structural Biology Center, X-Ray Science Division, Advanced Photon Source, Argonne National Laboratory; Lemont, IL 60439, USA.

[5]Department of Dermatology, Harvard Medical School; Boston, MA 02115, USA.

[6]Department of Informatics and Analytics, Dana-Farber Cancer Institute; Boston, MA 02115, USA.

[7]Department of Chemical and Biomolecular Engineering, Vanderbilt University; Nashville, TN 37212, USA

[8]Belfer Center for Applied Cancer Science, Dana-Farber Cancer Institute; Boston, MA, USA.

[9]Departments of Biomedical Engineering, Materials Science & Engineering, Physics & Astronomy, Texas A&M University; College Station, TX 77843, USA

[10]Center for AI and Natural Sciences, Korea Institute for Advanced Study; Seoul 02455, Republic of Korea

[11]Department of Molecular Physiology and Biophysics; Vanderbilt University School of Medicine; Nashville, TN 37232, USA

*Corresponding authors. Email: ellis_reinherz@dfci.harvard.edu, matt.lang@vanderbilt.edu

34   **Summary**
35
36
Memory T cells endow the mammalian host with an essential form of adaptive immunity
that generates durable and rapid protection following re-exposure to an infectious pathogen
or a cancerous transformation[1-4]. How naive CD8[+] T cells develop into memory T cells that
bifurcate into long-lived central (T_{CM}) versus effector (T_{EM}) subsets after such an antigen
encounter remains ill-defined at the molecular level. To address this gap, here we dissect the
functional T-cell receptor (TCR) repertoire of 242 murine CD8[+] TCR$\alpha\beta$ clonotypes specific
for the peptide-major histocompatibility complex molecule (pMHC) comprising an
immunodominant influenza A virus (IAV) oligopeptide epitope $NP_{366-374}$ presented by H-2D[b]
[5,6]. Using single-cell transcriptomics with TCR sequencing, biophysical measurements of
force-dependent TCR-pMHC interactions, *in vivo* memory development, and structural
analyses, we integrate these data to define relationships between signaling polarity and
memory fate. We find that T_{CM} and T_{EM} polarities are reflected in distinct subunit-related
mechanotransduction biases: T_{CM}-associated TCRs favor TCRβ-driven pMHC engagement,
whereas T_{EM}-associated TCRs preferentially favor TCRα-driven pMHC engagement.
Moreover, "bipolar" clonotypes (T_{BP}) capable of generating both memory subsets exhibit
more balanced signaling, the largest average clonal expansions, and broadest heterosubtypic
crossreactivities. By contrast, variegated T_{CM} TCR sequences foster mostly clonal singlets
with the greatest IAV mutant recognition potential within the overall memory repertoire,
thus revealing two complementary memory niche crossreactivity strategies. Weak, force-
sensitive TCR bonds are pervasive for this entire repertoire, being most conspicuous amongst
T_{CM} clonotypes. At the atomic level, highly nuanced variation in TCR$\alpha\beta$ bonding to either
peptide and/or MHC modulates mechanical load transmission across the holoreceptor's
ectodomains, transmembrane segments, and CD3 cytoplasmic tails to differentially influence
downstream tyrosine phosphorylation and memory development. Collectively, TCR
diversity anticipates epitope-specific pathogen evolution, whereas divergent TCR signaling
regulates memory fate with implications for adoptive T-cell immunotherapies seeking to
maximize persistence and minimize functional exhaustion.

Introduction

Immunological memory is a hallmark of the adaptive immune system, enabling rapid and robust targeted responses upon re-exposure to pathogens or tumors. CD8$^+$ T cells play a central role in this process by first recognizing threats through their clone-specific (clonotypic) TCRs and then eliminating infected or malignant cells by elaborating cytotoxic molecules and cytokines. Following acute viral infection, a small population of effector CD8$^+$ T cells that survive the predominant contraction phase differentiates into long-lived memory cells to provide durable protection [1,2]. Memory CD8$^+$ T cells are functionally and phenotypically heterogeneous but broadly classified into two circulating lymphocyte subsets, termed central memory (T$_{CM}$) and effector memory (T$_{EM}$) cells [3]. A non-circulating subset of tissue-resident memory T cells (T$_{RM}$) also forms [4]. T$_{CM}$ and T$_{EM}$ cells differ in surface markers, migratory behavior, functional potential, and transcriptional and epigenetic landscapes [4]. T$_{CM}$ cells express lymphoid-homing receptors such as CD62L and CCR7, enabling their lodging in secondary lymphoid organs. They also express transcription factors including Tcf1 [7], Lef1 [8], and Id3 [9] which support T$_{CM}$-like properties such as self-renewal, long-term persistence, and a robust proliferative response upon antigen re-encounter [10,11]. In contrast, T$_{EM}$ cells lack CD62L and CCR7, allowing them to circulate through the blood and access both lymphoid and non-lymphoid tissues. They preferentially express Id2 [12,13], which promotes effector differentiation, and are enriched for cytotoxic mediators such as granzyme B (Gzmb) and granzyme K (Gzmk), positioning them as a frontline defense capable of exerting immediate effector function [1,3].

The generation of memory subsets during viral infection is believed to be shaped by both extrinsic factors such as antigen abundance, inflammatory cytokines, and the tissue microenvironment as well as intrinsic transcriptional regulators and metabolic programs [1,14,15]. How memory fate is ultimately determined, however, remains unresolved. Notwithstanding, three conceptual models have been proposed [1]. The first, the intrinsic memory T cell fate model, suggests that fate is imprinted in naïve T cells prior to activation, i.e. the TCR *per se* directs memory T cell development. The second, the linear differentiation model, suggests that memory precursor cells (T$_{MP}$) give rise to memory subsets sequentially from T effector cells (T$_{EFF}$) to T$_{EM}$ followed by T$_{CM}$, with the latter especially pronounced in the context of high precursor frequencies [10,16-18]. The third, so-called signal strength model, alternatively posits that the intensity of TCR and co-stimulatory signaling during priming governs fate decisions, with stronger signals promoting T$_{EM}$ differentiation and weaker signals favoring T$_{CM}$ [19-21]. These models are largely based on several TCR transgenic systems, with or without altered peptide ligands (APLs) utilized to tune TCR signal strength. While elegant, these systems cannot recapitulate the diversity of TCRs and T-cell competition operative in the mammalian repertoire.

As recent studies have shown that TCR ligand recognition specificity and sensitivity are regulated by T-cell sensing of physical load placed on a TCR-pMHC bond during immune surveillance [5,22-32], we examined the biophysical and structural contributions of this mechanosensing to intrinsic CD8 memory fate within a single TCR repertoire. Our results indicate that memory T$_{EM}$ and T$_{CM}$ differentiation is unrelated to TCR signaling threshold performance metrics. Instead, differentiation "polarity" is dependent upon disparate signaling linked to the structural features of the TCR-pMHC interaction, including bonding and surface topology

120   involving dynamic TCR behaviors under force that impact mechanotransduction. These attributes
121   define an atomistic basis for divergence amongst TCRs differing by as few as 1-3 amino acids.
122
123
124
125   **Results**
126
127   **TCR clonotypes of varying T-cell burst sizes and polarities**
128
To investigate whether αβTCRs *per se* impact memory T cell subset fate, we performed single-
cell RNA Sequence (scRNA-seq) on $NP_{336-374}/D^b$-specific CD8 T cells isolated from pooled
mediastinal lymph nodes (mLNs) of 10 B6 mice at 34 days post-IAV infection (dpi 34) using the
mouse-adapted A/Puerto Rico/8/1934 (PR8) strain (Extended Data Fig. 1a). A total of 5,429 CD8
T cells comprising 242 clonotypes with paired TCRα and TCRβ V domain sequences were
identified. Uniform Manifold Approximation and Projection (UMAP) analysis revealed two major
clusters corresponding to central memory CD8 T cells (T_{CM}) and effector memory CD8 T cells
(T_{EM}), appearing as left- and right-sided lobulated aggregates, respectively, (Fig. 1a). Furthermore,
differentially expressed genes (DEGs) and hierarchical clustering (Supplementary Data 1 and
Extended Data Fig. 1b) discriminated three distinct T_{CM} clusters and three T_{EM} clusters. Of the
former, $T_{CM-CONV}$ (T_{CMC}) expressed conventional T_{CM} genes (*Klf2, Id3, Tcf7, Ccr7*), $T_{CM-XCL1}$
(T_{CMX}) specifically expressed *Xcl1, Cxcl10, and Mif*, and T_{CM-PRO} (T_{CMP}) was enriched for
proliferative genes (histone-related genes, *Stmn1, Pclaf, Birc5, and Ube2x*). With respect to the
T_{EM}, $T_{EM-CONV}$ (T_{EMC}) cluster expressed conventional T_{EM} genes (*Gzmb, Gzmk, Id2, Ccl5*), T_{EM-}
$_{KLR}$ (T_{EMK}) was enriched for KLR genes (*Klrd1 and Klrc1*), and T_{EM-EXH} (T_{EME}) expressed
exhaustion-associated genes (*Pdcd1*, *Lag3*, *Havcr2*). Selected top genes are shown in Fig. 1b.
Independent gene signatures derived from T_{CM} and T_{EM} clusters, identified by separate scRNASeq
analysis on $CD8\beta^+CD44^+$ memory cells isolated by FACS from pooled mLNs of four B6 mice at
dpi 33 (Methods) validated the transcriptional identity of these subsets specifically in the IAV
context (Fig. 1c and Extended Data Fig. 2a). Interestingly, *Sell*-encoded CD62L, a canonical T_{CM}
cell surface marker, was weakly expressed even within the T_{CM} cluster, as reflected by the low
frequency on $NP_{366-374}/D^b$-specific memory $CD8^+$ T cells (Extended Data Fig. 2b,c), compared to
the level on $PA_{224-233}/D^b$-, and $PB1_{703-711}/K^b$-specific $CD8^+$ T cells. Hence, the expression of
classical memory subset markers is apparently modulated in an epitope-specific manner.
153
154
We analyzed the distribution of clonotypes across the UMAP clusters to determine whether
TCRs significantly influence memory fate (Fig. 1d), identifying those with single TCRα and TCRβ
chains after filtration (Supplementary Data 2 and Methods). Among the top 40 clonotypes, 25 were
distributed comparably amongst subsets of T_{CM} vs T_{EM} clusters. In contrast, the remaining 15
clonotypes were highly skewed toward one or the other subsets, suggesting that TCRs *per se*
impact memory T cell fate as indicated by the low adjusted p-values (Fig. 1e and Supplementary
Table 1). Accordingly, clonotypes were categorized as belonging to the T_{CM} vs T_{EM} type when
>70% of the cells in each clone's progeny were polarized toward one or more of the three
respective subclusters therein, as defined in Fig. 1a. Other clonotypes with a more equivalent
distribution not achieving the 70% threshold were classified as bipolar (T_{BP}). Representative
clonotypes listed in Fig. 1e underwent further analysis. The NP41, as well as NP34 and NP63,

TCRs identified in a previous study of the IAV memory recall response [5] were also observed here
in the primary memory pool, as one T_{CM} and two T_{BP} clonotypes, respectively. Moreover, since
both analogue NP34 and digital NP63 are categorized as T_{BP}, their different performance under
pN forces (i.e. digital TCRs need only a few pMHC for activation while analogue TCRs require
dozens to hundreds), is not linked to the T memory subset determination. The cell projection on
the UMAP (Fig. 1f) and the Euclidean distance of principal component-1 (PC1) for NP201, NP204,
and NP207 (Fig. 1g) confirmed their locations in $T_{CM,}$ $T_{EM,}$ or both clusters, highlighting gene
expression similarity between individual clonotypes and the overall T_{CM} and T_{EM} populations as
well as total T cells.

We further examined the localization of all cell types belonging to each memory clonotype
category (CT) and projected their distribution on the UMAP of the entire CD8 repertoire (Fig. 1h).
"CT-matched" cells (i.e., T_{CM} cells in T_{CM} CTs and T_{EM} cells in T_{EM} CTs) are localized primarily
toward the periphery of the respective T_{CM} and T_{EM} regions, whereas T_{CM} and T_{EM} cells in T_{BP} CTs
are diffusely spread but with more concentrated toward the center of UMAP. This differential
distribution suggests that CT-matched cells express their respective signature genes more strongly
than T_{BP} clonotypes or those "CT-mismatched" cells (i.e., T_{CM} cells in T_{EM} CTs and T_{EM} cells in
T_{CM} CTs), thereby enhancing their fate determination. Consistent with this, RNA velocity analysis
supports the notion that T_{CM} and T_{EM} clonotypes represent more terminally differentiated states,
as directional flow arrows from CT-matched cells are either absent or pointed toward the outer
UMAP edges (Extended Data Fig. 3a). In contrast, T_{BP} clonotypes appear more interchangeable,
as the arrows from T_{CM} and T_{EM} cells in T_{BP} CTs indicate some bidirectional flow.

To determine whether the expression of all signature memory genes or only a specific
subset underlie the memory polarization, we analyzed differentially expressed genes in T_{EM} and
T_{CM} clusters (Supplementary Data 3) across CT-matched cells, CT-mismatched cells, and T_{CM} and
T_{EM} cells in T_{BP} CTs. Both aggregate T_{EM} and T_{CM} signature gene expressions increased
progressively with clonotype polarization (Extended Data Fig. 3b,c). Of the 126 T_{EM} signature
genes elevated in the T_{EM} cluster, only 45 genes, including *Ccl5*, *Gzmb*, *Gzmk*, and *Fasl* showed a
gradual increase in expression corresponding to the degree of clonotype polarization, suggesting
that these genes may contribute directly to clonotype fate commitment (Extended Data Fig. 3b).
Similarly, of 140 T_{CM} genes, only 65 genes, such as *Lef1*, *Tcf7*, and *Tnfsf8* exhibited graded
expression aligned with clonotype polarity, implying they are key regulators of T_{CM} fate (Extended
Data Fig. 3c).

The T_{CM} repertoire is most variegated

Having established that a TCR clonotype influences memory fate, we examined whether TCR
gene segment features are differentially associated with the T_{CM}, T_{EM}, and/or T_{BP} transcriptional
phenotype. V and J gene usage analysis revealed the preferences for TRAV16N/TRAV16, TRBJ2-
2/TRBJ2-7, and TRBV13-1 among all 242 NP-specific clonotypes (Fig. 2a), consistent with a
previous study [33]. Notably, the 140 T_{CM} clonotypes exhibited greater diversity in V and J gene
usage, while 44 T_{BP} clonotypes showed a more restricted repertoire, with a higher frequency of the
dominant VJ combinations found across the NP-specific TCRs. The 58 T_{EM} clonotypes displayed

an intermediate pattern. Entropy analysis confirmed the differences in VJ diversity among the
memory subsets (Extended Data Fig. 4a).

Next, we compared clonal burst sizes amongst memory populations. T_{CM} clonotypes were
the largest fraction in lower cell number categories, particularly in the single-cell group, compared
to T_{EM} clonotypes (Fig. 2b). In contrast, T_{BP} clonotypes were significantly more expanded than
either T_{CM} or T_{EM} clonotypes (Fig. 2c). Since clonotypes with only one cell cannot be classified as
T_{BP} by definition, we repeated the analysis excluding clonotypes with single-cell clonotypes and
observed consistent result (Extended Data Fig. 4b). These findings suggest that T_{CM} tend to persist
as a small number of cells, whereas T_{BP} are more clonally expanded.

To distinguish clonotypes with similar versus divergent TCR sequences, we categorized
branches of a TCR sequence tree into "short" and "long", respectively, based on paired TCR
sequence distances metrics (Fig. 2d). For each clonotype, we calculated the distance to all other
clonotypes and computed the mean of the top 10 shortest pairwise distances. For example, CT_25,
categorized as a short-branch clonotype, had a mean top 10 distance of 7.8, whereas CT_50,
categorized as a long-branch clonotype, had a corresponding value of 196.4 (Extended Data Fig.
5a,b). A histogram of these mean top 10 distances across all clonotypes revealed a bimodal
distribution, and we defined the valley between the two peaks (mean distance = 109) as the
threshold distinguishing short- and long-branch clonotypes (Extended Data Fig. 5b). A TCR
distance heatmap further supported the validity of this classification (Extended Data Fig. 5c).
Moreover, analysis of short- and long-branch clonotypes showed that short-branch clonotypes
exhibited more restricted V and J segments commonly used in NP-specific TCRs, whereas long-
branch clonotypes displayed greater diversity (Extended Data Fig. 5d).

We then examined DEGs between short vs. long branch clonotypes. Short-branch
clonotypes showed higher expression of T_{EM} genes such as *Gzmk* and *Ccl4*, whereas long-branch
clonotypes expressed higher levels of T_{CM} genes, including *Sell*, *Ccr7*, and *Satb1* (Fig. 2e and
Supplementary Data 4). Gene enrichment analysis confirmed a significant enrichment of T_{EM} gene
signatures in short branches and T_{CM} gene signatures in long branches (Fig. 2f). We further
examined the distribution of representative memory clonotypes on the TCR tree, including NP101
and NP105, identified from an independent scRNA-Seq experiment (Methods) with NP101
showing identical amino acid sequence to CT_5 and CT_56 (Supplementary Table 1) and NP105
bearing a significant relationship to NP204 described below. Representative T_{EM} clonotypes such
as NP201 and NP202, as well as T_{BP} clonotypes including NP34, NP63, NP101, NP206, and
NP207, fell within the short branch (Fig. 2d). In contrast, T_{CM} clonotypes NP41, NP204, and
NP205 were found in long branch, supporting the notion that TCR sequence similarity is associated
with T_{EM} or T_{BP} phenotypes, while greater sequence divergence is linked to the T_{CM} phenotype.

Deviations to this pattern include T_{CM} clonotypes NP105 and NP203 belonging to the short branch
(Fig. 2d). In the case of NP203, this is probably explained by the TCR sequence differing by only
1-3 amino acids from short-branch T_{BP} clonotypes NP101 and NP207 and T_{EM} clonotype NP202.
Hence, minor sequence differences can override the general pattern to influence fate decisions
despite overall amino acid similarity. This observation prompted us to pursue structural analyses
described in a later section to reveal "the exception that proves the rule". In sum, the T_{CM} repertoire

is more variegated than that of the T_{EM} repertoire but interclonal TCR distances *per se* do not define
transcriptome polarity.
253
254
255
256 **Weak TCR-NP$_{336-374}$/D^b bonding profiles**
257
Single molecule (SM) TCR$\alpha\beta$-pMHC analysis was used to evaluate the performance and bonding
characteristics of clonotypes belonging to each memory category (Extended Data Fig. 6a,b). T_{CM}
clonotypes NP203, NP204, NP205, and NP105 were all weak binders with profiles consistent with
slip-bonds. T_{EM} clonotypes NP202 and NP201 were also weak binders but exhibited more typical
catch-bond profiles with peaks around 15 piconewtons (pN). T_{BP} clonotypes NP101 and NP207
exhibited mixed profiles, with higher lifetime amplitudes at both the typical slip and catch force
regions (5 and 15 pN, respectively), while NP63 was a catch bond as in previous work[5].
Additionally, overall lifetime amplitudes were weak relative to verified digital clonotypes such as
NP63[5] (Fig. 2g and Extended Data Fig. 6c). The ability to form a stable single-molecule TCR$\alpha\beta$-
pMHC interaction irrespective of load was determined by comparing the total number of binding
opportunities allowed to the total number of events measured. Stable tethering formation varied
substantially among the clonotypes, with T_{CM} clonotypes showing a notably lower tendency to
form stable bonds, which also tended to bind in the slip-type manner.
271
Excluding NP63, ranking relative performance based on lifetime amplitude around 15 pN in
addition to the tendency to form stable tethers, suggest that T_{EM} clonotype NP202 and T_{BP}
clonotypes NP207 and NP101 are a tier above T_{CM} clonotypes NP203, NP205, and NP105. NP204
and NP201 are the lowest performing clonotypes, exhibiting low lifetime amplitudes across all
force ranges measured, coupled with a poor ability to form stable bonds measured by binding
probability (Fig. 2h). Binding probability was strongest for NP202 (and NP63) compared to the
other clonotypes profiled, where multiple tethers were observed.
279
Retrogenic profiling reprises polarization
To determine whether $\alpha\beta$TCRs are linked to CD8 differentiative fate following antigen encounter
and memory phenotypes identified by scRNA-seq reflect clonotype-specific polarization *in vivo*,
we generated retrogenic chimera (RgC) mice. This was achieved by adoptively transferring naïve
CD8$^+$ retrogenic (Rg) T cells from Rg donor mice into B6 [CD45 (2/2) Thy1.2] mice, followed by
infection with the PR8 one day later (Fig. 2i). At 30dpi in mixed NP101/NP105 RgC mice, NP105
Rg T cells exhibited significantly higher expression of T_{CM} markers, CD62L and Tcf-1, compared
to NP101 Rg T cells (Fig. 2j and Extended Data Fig. 7), indicating that the RgC system faithfully
recapitulates clonotype polarization patterns identified by scRNA-seq.

Memory IAV crossreactivities

The role of T cells in providing heterosubtypic IAV protection to largely conserved epitopes is
well established [34] despite limited structural analysis [35]. To ascertain if such crossreactivity could
be extended more broadly to various memory clonotypes recognizing NP$_{366-374}$/D^b, we selected 16
TCRs that were transduced into mCD8$\alpha\beta^+$BW5147.3 (BW) cells and sorted for comparable TCR

expression. BW transductants were then stimulated with WT-PR8 $NP_{366-374}$ peptide or 9 naturally
occurring variant peptides derived from 11 other IAV strains [36] on APCs. IL-2 production was
assayed as a readout of TCR activation (Fig. 2k). The importance of the structurally featured
residues of the peptide (p4E, p6M and p7E) were also quantified (Extended Data Fig. 8a,b) and
referenced to the IAV variants (Extended Data Fig. 8c). Crossreactivity was observed in 14 of 16
TCRs, ranging from 1-6 of the 9 variants, with all TCRs responding to the isologous PR8 strain
(Fig. 2k and Extended Data Fig. 8b,d). As 0/6 T_{EM}, 1/5 T_{CM} and 3/5 T_{BP} recognized 5 or more viral
epitopes, T_{BP} recognition trended as more crossreactive compared to the polar clonotypes (p =
0.0632 using a two-tailed Fisher's exact test). The greatest crossreactivity was exclusive to the T_{BP}
(NP101, NP206 and NP207). As an example, detailed IL-2 production from NP207 is further
quantified by peptide titration (Extended Data Fig. 8e).

305
306 **Protein expression, crystallization, and analyses**
307
To obtain diffraction quality crystals of TCRαβ-$NP_{366-374}$/D^b complexes for structural studies, we
engineered mouse TCRs and TCR chimeras with murine V domains fused to human C domains
(sequences in Supplementary Data 5). All TCRαβ heterodimers were generated in eukaryotic
Expi293F N-acetylglucoseaminotransferase I-deficient (GnTI⁻) cells to facilitate deglycosylation
under native conditions. H-2D^b and β2m proteins were produced in *E. coli* and refolded *in vitro*
with chemically synthesized $NP_{336-374}$ peptide to generate pMHC complexes. Procedures for
purifications, crystallization screening, and data collection are detailed in Methods. A series of
crystals of unligated TCRs and their complexes were obtained and structurally determined.
Crystallographic statistics, complex interface buried surface areas (BSA) and surface
complementarity (Sc values) along with H-bonding and van der Waals interactions are given in
Supplementary Table 2-4. Supplementary Discussion Item 1-4 offer specific information on
TCRαβ apoproteins, both wild-type (wt) and chimeras, and delineate incidence, twist, and shift
docking parameters on the same $NP_{336-374}$/D^b ligand.

321
Fig. 3a tabulates information on the six TCRs from B6 mice that recognize the $NP_{366-374}$
PR8 IAV epitope used for structural analysis. Four TCRs are representative of T_{EM} (NP202), T_{BP}
(NP101, NP207), and T_{CM} (NP203). It is striking that the TCRs utilize identical TRAV and TRBV
gene segments differing by only 1-3 CDR3 amino acid residues and yet belong to functionally
distinct memory CD8 subsets (Fig. 3a). We also selected the NP204 and NP105 T_{CM} TCRs
comprising distinct V gene segments but with similar CDR3β loops given a shared TRBJ1-6
differing but just 4 of 17 residues. Note that CT_29 (Supplementary Table 1) is a T_{CM} like NP105
and NP204. CT_29 is most similar to NP105 with utilization of the same TRAV, TRBV, TRBJ,
but a different TRAJ.

331
Each of the six TCRs adopts a canonical binding mode in their respective complex such
that Vβ overlies the C-terminal half of the peptide and the MHC α1 helix while Vα overlies the
N-terminal half of the peptide and MHCα2 (Fig. 3b). Amongst the various TCR-$NP_{336-374}$/D^b
complexes, the docking modes of the four closely related clonotypes, NP202, NP101, NP207 and
NP203 appear quite similar and with comparable footprints on $NP_{366-374}$/D^b (Fig. 3c), complex
interfaces (BSA and Sc) (Supplementary Table 3), and docking parameters (incidence angle, twist
and shift) (Supplementary Discussion Item 4). Those interactions are certainly distinct from the

two footprints of the "Vβ-centric" NP105 and NP204 TCRs analyzed using the same set of
parameters. We refer to NP105 and NP204 as Vβ-centric since the TCR-pMHC complex in each
is dominated by structural interactions involving its Vβ domain. For these two Vβ centric TCRs,
the ratio of TCRα/TCRβ BSA in the TCR-pMHC complex is 0.26-0.59. In contrast, for the four
NP 202, NP101, NP207 and NP203, these parameters are 1.23-1.33, due to considerably greater
Vα interactions. Amongst those four highly related NP TCRs, their distinct interactions with NP_{336-}
$_{374}$/D^b are only appreciated following detailed structural examination.
346
347
348   **Atomistic basis of CD8 memory fates**
349
The primary crystallographic features of the NP202 T_{EM} TCR clonotype's interaction with NP_{336-}
$_{374}$/D^b are shared with NP101, NP207 and NP203 interactions. The most exposed $NP_{336-374}$ peptide
residues extending upward from the MHC groove are p4E and p6M, and to a lesser extent, p8T
(Extended Data Fig. 8a), in agreement with a prior pMHC structure [37]. When the NP202 TCR
docks on $NP_{366-374}$/D^b, the negatively charged carboxylic group of p4E is positioned below the
center of a glycine-rich loop from CDR3α, forming 2-3 H-bonds to the amide groups (Fig. 3d).
The central p6M sidechain is sandwiched by CDR3α and CDR3β and constrained in a
hydrophobic pocket involving aliphatic sidechains of K98α and R99β. An additional specific
interaction is a H-bond between side chains of p3N and CDR3 K98α.
359
The NP101 T_{BP} TCR differs from the NP202 T_{EM} TCR by a single CDR3β residue (D95β
vs. E95β), as highlighted in Fig. 4a. Consequently, the two TCR docking angles and binding sites
on $NP_{366-374}$/D^b are very similar (Supplementary Discussion Item 4). Structural alignment of their
H-2D^b α1α2 domains demonstrates excellent overlay of the peptide binding grooves and
corresponding $NP_{366-374}$ peptides except at the central p6M residue differing by a slight shift and
side chain rotation. Although the single-residue substitution resides in CDR3β, the Vβ domain
including CDR3β exhibit limited deviation (shortest red arrow). However, a positional shift is
evident in the Vα domain comparison as a combined result of slightly different TCR docking
angles onto the pMHC and deviations at the Vα-Vβ interface between the two TCR V modules
(longer two red arrows). This shift brings important pocket-forming residues, including F33α and
Y54α (in green), as well as the aliphatic part of R99β (in cyan), closer to p6M, which significantly
reduces the size of the hydrophobic pocket. For example, the Cα-Cα distance between D52α and
R99β, a representative distance between CDR2α and CDR3β, is reduced from 7.78Å in NP202 to
6.80Å in NP101. The apposition of the Vα C' strand edge of to the Vβ G strand edge is reflected
in the formation of two salt bridges, R50α to E101β and D52α to R99β in NP101. Shrinking of
the p6M hydrophobic pocket strengthens hydrophobic and van der Waals interactions with NP101.
Meanwhile, the distant interaction between p4E to G100α is lost in NP101 (Supplementary Table
4). Together, the overall structural alterations foster the repositioning of the TCR-pMHC
interaction center from the p4E to the p6M residue, thereby doubling the number of p6M van der
Waals contacts made by NP101 relative to NP202 (3 vs 6, Supplementary Table 4). This subtle
difference is not revealed by TCR docking angles or footprints onto pMHC (Supplementary
Discussion Item 4 or Fig. 3c). A second T_{BP} clonotype, namely NP207, manifests a similar shift
akin to NP101 associated with TCR crossreactivity against variant $NP_{366-374}$ heterotypic IAV
strains, as noted in Fig. 2k and Extended Data Fig. 8.

385    NP207 (T_{BP}) and NP203 (T_{CM}) differ at two residues, K98α vs R98α in CDR3α and V99β

386 vs L99β in CDR3β, respectively, as highlighted in Fig. 4b. In the NP207-NP$_{366\text{-}374}$/D^b complex,

387 K98α forms a H-bond with p3N. A well-defined water molecule (labeled as W) also bridges the

388 interactions between K98α and p3N to the ND1 atom of H155 in the D^b α2-helix. Meanwhile, p7E

389 and NE2 of H155 form a salt bridge. However, in the NP203 complex, the longer R98α sidechain

390 both pushes p3N slightly downward and displaces the water molecule found in the NP207 complex

391 to form a salt bridge directly with the ND1 atom of H155. The ND1 atom of H155 is likely

392 deprotonated to neutralize the extra positive charge introduced by the K$\rightarrow$R substitution, while the

393 NE2 of H155 is protonated and neutralized by p7E, thereby becoming indispensable in the NP203

394 TCR recognition of pMHC, in contrast to NP207, as elaborated below. The CDR3β L99 in NP203

395 (T_{CM}) relative to CDR3β V94 in NP207 (T_{BP}) augments van der Waals contacts with p6M, whereas

396 the H-bonds of p4E to CDR3α are weakened with the bond length (p4E to G101α) increase from

397 2.97Å to 3.34Å (Supplementary Table 4). These changes collectively contribute to the apparent

398 shift of the NP203 TCR mechanical load from α chain to β chain.

400    Fig. 4c reveals two additional differences between the NP207 (T_{BP}) and NP203 (T_{CM})

401 complexes related to their respective Vα domains and the single N-glycosylation site adduct. In

402 NP207, as well as in NP101 and NP202, the Vα C" edge strand of shifts from the GFCC' to the

403 ABED face as commonly observed in TCR Vα domains [38]. In NP207, the conserved N-glycan

404 binding residue N58α is a part of CDR2α. The first N-acetylglucosamine (GlcNAc) attached to

405 N58α is visible in the structure, pointing away from the ABEDC" sheet of Vα domain. In this

406 orientation, even a mature N-linked glycan attached to N58α is unlikely to extend to the TCR-

407 pMHC interface. In contrast, in the NP203 complex, the C" of Vα domain is absent with no

408 secondary structural element between C'-strand and D-strand. Instead, the extended loop from

409 C'- to D-strands is stabilized by two side chain-main chain H-bonds between the loop and the D-

410 strand (boxed insert in Fig. 4c). Because E57α points to the D-strand, N58α pivots from the D-

411 strand to the interface edge between the Vα C'-stand and the Vβ G-strand, repositioning the

412 attached GlcNAc to above the α2-helix apex. The N-glycan with its additional sugar adducts

413 attached to N58α (absent here due to EndoH cleavage) presumably contacts the pMHC, impacting

414 TCR ligation, mechanotransduction involving the TCR-pMHC interface, and NP203 T_{CM} polarity.

416    Considering the striking divergence in the transcriptomes of the NP202, NP101, NP207

417 and NP203 clonotypes, these structural distinctions must impact gene activation. Molecular

418 dynamics simulations under force show that interdomain motion, molecular compliance,

419 fluctuating forces, and interfacial contacts collectively determine the mechanical response and

420 fine peptide TCR discrimination [39,40]. To measure differential physical stress responses of

421 individual clonotypes we counted the number of TCRα and TCRβ residues that form 6 or more

422 intradomain contacts with occupancy greater than 80% during a 700ns simulation. The greater the

423 number of these 'hub' residues, the more tightly packed and rigid the corresponding domain is,

424 which indicates that the domain bears greater physical load. Although these TCRs differ by only a

425 few amino acids (Fig. 3a), under identical loading conditions the numbers of the hub residues

426 varied considerably among the 4 subdomains (Supplementary Discussion Item 5), which comports

427 with exquisite biomechanical nuancing.

The two NP204 and NP105 T_{CM} clonotypes distinguish themselves from NP202, NP101,
NP207 and NP203 above by their Vβ-dominant docking footprints on $NP_{336-374}/D^b$ (Fig. 3b,c), and
as further visualized by the comparison of NP202 versus NP105 side views (Fig. 4d, left panel).
The NP105 TCR-pMHC interaction is dominated by Vβ, notably involving its
CDR3β motif D96RRxS100β (Fig. 4d, right panel). The NP105 CDR3β D96 forms a hydrogen
bond to p8T, akin to D95β or E95β found in other NP TCRs. Meanwhile, R97β forms a salt bridge
to p4E, which lacks interaction with the Vα domain in contrast to the four TRBV5-containing NP
TCRs (Fig. 4a,b). R98β forms a H-bond to the carbonyl group of p5N. The side chain and amide
group of S100β form a pair of H-bonds with the p7E sidechain. Furthermore, p6M is no longer
sandwiched between Vα and Vβ. Instead, it is surrounded by a part of CDR3β (from R98β to
S100β) and MHC residues H155 and Y156 (not shown) and capped by Y94α of CDR3α.
Strikingly, the Vα contribution to peptide recognition is minimal. The primary features of the Vβ-
dominant docking in NP105 are largely conserved in the other T_{CM} NP204 that utilizes distinct Vα
and Vβ genes but shares the CDR3β motif with NP105 (Extended Data Fig. 8g).

Differential tyrosine phosphorylation and V-centric TCR binding

The distinct transcriptional programs elicited by pMHC ligation of TCRs, correlating with Vα- vs
Vβ-centric interactions , suggest that different downstream signaling pathways should be apparent,
especially considering the known compartmentalization of CD3 dimers in the holoreceptor
(Extended Data Fig. 9a,b). As CD3ζ tyrosine phosphorylation is maximal at 1 minute following
pMHC ligation of TCRs in triggered T cells[41], we used this as a proxy to investigate NP202 versus
NP105. As shown in Extended Data Fig. 9c, the NP202-associated CD3ζ subunit is significantly
more phosphorylated following $NP_{366-374}/D^b$ tetramer stimulation compared to that of NP105, as
measured by phospho-Tyr 142 detection. This difference in three independent experiments is not
due to reduced NP105 signaling ability, as higher molecular weight proteins in the 50-75 kDa and
100-120 kDa ranges are as or more heavily phosphorylated in NP105 compared to NP202
(Extended Data Fig. 9d). Yet, within the ~20-25 kDa range, where the CD3 subunits are found to
migrate, the NP202 appears to result in a greater level of phosphorylation. Thus, while the NP202
α-dominant TCR results in higher levels of CD3 phosphorylation, the NP105 β-dominant TCR
shows phosphorylation of alternative signaling proteins.

Structural basis of heterosubtypic immunity

Structural analysis also delineates the basis for differential TCR heterosubtypic immunity. For
example, NP105 tolerates a peptide E7D viral mutation, whereas NP204 does not. The weaker H-
bonds between pE7 and CDR3β residues in NP204 (approximated as 3.10 Å and 3.49 Å) would
have a greater impact than those between pE7 and CDR3β residues in NP105, which are shorter
(2.77 Å and 2.82 Å) (Supplementary Table 4). Pointedly, the E7D mutation results in a 1.54Å
sidechain foreshortening given a loss of one carbon-carbon bond. Distances between hydrogen
bonding atoms would likely become too weak and break for NP204. Likewise, the tilt observed in
NP101 (Fig. 4a) and NP207 (Extended Data Fig. 8f) relative to the T_{EM} NP202, makes both T_{BP}
clonotypes more crossreactive to p6M variants given that smaller sidechains such as I, L and V

(Fig 2k) readily accommodate the reduction in volume of the hydrophobic pocket hosting the
central p6 residue.
475
476
Activated T$_{CM}$ performance

Because high numbers of transferred T cells modulate memory phenotypes during antigen
competition [16-18], we asked whether TCRs would continue to dictate intrinsic differentiation bias
under such challenging conditions. 100,000 Rg T cells bearing NP101 or NP105 TCRs were
adoptively transferred to B6 hosts and infected with PR8 24 hrs later. Then, at 35 dpi mLNs were
isolated and Rg T cells were sorted based on congenic markers. PCA of gene expression following
RNA-Seq for the 14 individual lymph node samples (7 for NP101 [C101 series], 7 for NP105
[C105 series]) delineated 3 clusters, 2 consisting only of NP101 (Cluster 1) or NP105 (Cluster 3),
and an intermediate cluster (Cluster 2) of mixed clonotypes (Fig. 5a). Differential expression of
the top 500 genes identified 2 clear sets of genes partitioning with high probability between the
NP101 and NP105 clusters (Fig. 5b). Manual curation of the gene list for those reported to have T
cell functionality reduced the set to 59 expressed genes segregating with high probability between
the NP101 cluster and the NP105 cluster (Fig. 5c). Application of this gene list to all 3 clusters
identified the NP101 cluster 1 as displaying an exhaustion signature, as generally found in T$_{EM}$,
especially in chronic infection and tumor environments [42,43]. In contrast, the NP105 cluster
expressing higher *Sell, Il7r* and *Bcl2* transcripts together with cytotoxic transcripts (*Gzma/Gzmb)*
and active metabolic programming (*Myc*) as well as stem cell proliferative genes (*Placd8, Pim1)*
suggested an activated T$_{CM}$ state with effector functionality (Fig. 5d). Meanwhile, Cluster 2 with
mixed NP101/NP105 samples revealed a representation of both activated T$_{CM}$ and exhaustion
signatures. To extend these findings, we performed a similar independent RNA-Seq for NP202,
NP204, and NP207 Rg T cells (C202/C204/C207 series). Clustering of this dataset revealed a
graded activation signature in addition to the T$_{CM}$ and exhaustion signature (Extended Data Fig.
10). Aggregation of the data (Extended Data Fig. 10e) identified differential partitioning by cluster
of the clonotypic retrogenic cells implying this was a property of each unique NP-specific TCR
(Fig. 5e). By χ^2 analysis, C101 and C105 samples partition differentially between clusters 1-3 (χ^2
= 19.14, dof = 6, P < 0.004), and the C202, C204, and C207 dataset also partition differentially
between clusters 1-4 (χ^2 = 23.33, dof = 9, P <0.006). Notably, only NP105 and NP204 manifest as
activated T$_{CM}$. Collectively, the deep bulk RNA-seq analyses identified TCR-linked CD8 IAV
memory responses of adoptively transferred Rg T cells to manifest greater exhaustion of T$_{BP}$ and
T$_{EM}$ counterparts relative to that of T$_{CM}$ under adverse conditions.

507
Discussion

TCR-pMHC interaction *per se,* primarily dictates fate commitment as revealed through single cell
transcriptomic/paired TCR analysis of B6 mice as well as TCR retrogenic chimeric animals. While
clonal plasticity, as noted herein (Fig. 5 and Extended Data Fig. 10) and reported elsewhere [44],
exists in relation to T-cell functionality states responsive to environmental queuing, TCR-linked
transcriptome bias is unquestionable. In this context, *Abdullah et al.* [44] recently explored
recombinant vesicular stomatitis virus (VSV), as well as the more inflammatory bacterial *Listeria*
monocytogenes systems, to study the naive SIINFEKL/K^b T cell response during those infections.

For both systems, TCR intrinsic bias among clonotypes was observed by day 7 of the acute
response, when short-lived effectors and memory precursor effector CD8 T-cells were generated,
several weeks in advance of memory development. Given that Abdullah *et al.* and our current
study utilized different pathogens and kinetics to affirm that TCR-pMHC interaction is critical for
naïve CD8 T cell differentiation, the collective conclusions are broad and complementary.

The structural mechanisms underlying fate determination were identified here through
structural analyses of six distinct TCR-NP$_{366-374}$/D^b complexes. Two tiers of differential
interactions were observed. In the <u>first tier,</u> a pronounced positional docking variance as to TCR
footprints of Vα versus Vβ domains, as well as BSA of each's TCR-pMHC interaction, pointed to
differential binding of T$_{EM}$ and T$_{CM}$ for NP202 vs NP105 and NP204, respectively. The former
TCR was dominated by Vα interactions focused over the α1 MHC helix, whereas the latter two
TCRs were Vβ binding centered around the α2 MHC helix. In these examples, the distinct TCR
footprint topologies at the ligand interface are termed Vα-centric for NP202 and Vβ-centric for
NP105 and NP204. Within the αβTCR holoreceptor complex, eight of the 10 cytoplasmic tail
ITAMs (CD3εδ, CD3ζζ) are attached to the inner leaflet of the membrane on the TCRα side, but
just 2 (CD3εγ) are located on the TCRβ side and further compartmentalized by transmembrane
segments separated by intercalated cholesterol molecules [[45] and references therein] (Extended
Data Fig. 9b). Such partitioning and binding asymmetry likely foster divergent signaling following
ITAM release (reviewed in [31]), impacting the magnitude and quality of downstream responses as
reflected in differential gene signatures noted after one week [44] and one month (current study), and
is consistent with the differential tyrosine phosphorylation observed herein for CD3ζζ. Signaling
differences probably also relate to the greater memory expansion mediated by the T$_{BP}$ clonotypes.

As the preTCR lacks a Vα domain in its pTα surrogate subunit that pairs with a canonical
β subunit preceding αβTCR development in the thymus [46], we posit that the preTCR pTα-β
heterodimer is an extreme example of a Vβ-centric interaction linked to asymmetric
mechanotransduction [47-49]. Given that the preTCR functions to foster survival [50], promoting the
incorporation and persistence of β chains into the evolving αβ lineage of thymocytes, there is an
analogy to Vβ-centricity involving peripheral T cells associated with long-term T$_{CM}$ development
and renewal, i.e. stemness. Consistent with this notion, five key genes involved in transcriptional
programming and/or survival that are upregulated in common during the preTCR-dependent β-
selection checkpoint and CD8 T$_{CM}$ memory generation are *Tcf7* [7,51], *Lef1* [8,51], *Il7r* [52,53], *Id3* [9,54] and
Bcl2 [55,56]. By contrast, T$_{EM}$ promote rapid effector functions but generally displays more limited
survival and cellular plasticity characteristics [1-3].

In the <u>second tier</u> of docking interactions, there is pronounced subtlety as footprints of
NP202, NP101, NP207 and NP203 are largely indistinguishable. Nonetheless, NP203 is less
capable of transducing TCRα-associated signals for two reasons: first, Vα structural rigidity is
attenuated by loss of the NP203 C" strand, and second, the attendant glycan adduct intercalating
between TCR and pMHC interaction surfaces may dampen applied physical load. We postulate
that this diminished TCRα mechanotransduction fosters T$_{CM}$ development. In contrast, NP101 and
NP207 T$_{BP}$ TCRs are without such structural impediments, but both are tilted to the Vβ side
relative to NP202 T$_{EM}$. We posit that tilt results in a differential force vector to transmit load,
creating a more comparable Vα/Vβ balance centered on the smaller TCR pocket hosting p6M,

fostering T_{BP} development unlike that of the more Vα dominant NP202 programming T_{EM}
differentiation.
562
Fig. 6 provides an overview of our observations. Panel a shows that CD3 signaling
molecules are asymmetrically positioned relative to the TCRα and TCRβ subunits, while panel b
illustrates that antigenic peptide residues and MHC helices interact in various patterns at the TCR-
pMHC interface, demonstrated by comparison of NP202, NP207, and NP105 as examples. Panel
c uses a seesaw analogy to explain how mechanical forces are transmitted through the TCR
subunits with ball size representing the relative mechanical load from the pMHC. Just as gravity
tilts a seesaw, uneven loading on the TCR influences which signaling pathways are activated.
When the load is balanced, i.e. for T_{BP} clonotypes, it allows for stochastic outcomes in how T cells
differentiate into T_{EM} or T_{CM} progeny during repeated cycles of attachment and detachment from
antigen-presenting cells (APCs) throughout clonal expansion [57-59] This balanced distribution of
T_{EM} and T_{CM} likely results from chance mechanical interactions at the contact points between T
cell microvilli and APCs, which generate different shear forces. In contrast, when the load is
biased, as regards polar T_{EM} and T_{CM} clonotypes, impact from this randomness is reduced and the
T cell is more likely to follow a specific signaling pathway.
577
Shrewdness in TCR mechanotransduction biology can dictate digital vs analog TCR
performance based on as little as a single CDR3α amino acid variation among two TCRs (NP63
vs. NP34) binding to the same NP$_{366-374}$/D^b ligand [5]. Conversely, a single amino acid change in a
peptide ligand (i.e , APL) bound to one MHC molecule can generate differential functional T cell
activation through the same TCR [60]. For the latter, molecular dynamics simulations under force
revealed a combined role of interdomain motion, molecular compliance, fluctuating forces, and
interfacial contacts that collectively determine the mechanical response and fine peptide TCR
discrimination [39,40]. The differential physical stress response observed here by molecular dynamics
amongst closely related TCR$\alpha\beta$ sequences underscores this exquisite nuancing (Supplementary
Discussion Item 5).
588
Given different clonotypic TCR sequences comprising a single T-cell repertoire, those
TCRs may experience varying levels of stress and strain distributions upon mechanical loading.
Thus, their load propagation pathways through the ectodomains of the $\alpha\beta$TCR holoreceptor into
the membrane and cytoplasmic tail elements may also deviate. These differences (Extended Data
Fig. 9) likely instruct distinguishable release of the ten ITAMs per $\alpha\beta$TCR complex and engage
divergent downstream signaling, comporting with our distinct tyrosine phosphorylation data and
transcriptomes. It is important to highlight that bipolar (balanced) vs polar (unbalanced) signaling
is evident for G-protein coupled receptors (GPCRs), where activating both G-protein and β-
arrestin routes collectively versus the latter alone dramatically alters functional cellular outcome
in μ-opioid and chemokine receptors (reviewed in [61]). For GPCRs, however, dissimilar ligands
acting on the same receptors mediate the differential outcomes, whereas in the T-cell memory
system explored here, the TCRs are distinct receptors that bind the same pMHC. While nearly all
of the ~800 various GPCRs are not mechanosensors, unlike TCRs, our findings hold significant
biological relevance and, as noted, are not without precedence[62 63].
603
The NP$_{366-374}$/D^b T-cell repertoire is generated against a ligand expressed at ~2000
copies/antigen-presenting cell [64] utilizing mostly analog TCRs [5]. The crossreactivities found in

~20% of the TCRs offer a teleological explanation for the highly divergent TCRs incorporated into
the memory niche, namely to foster recognition of natural viral strain diversification. The T_{BP}
clonotypes in our analysis manifest the greatest crossreactivity and largest burst sizes with rapid
cytotoxicity derived from their T_{EM} progeny, implying superior attributes. However, the T_{CM}
revealed the least exhaustion and greatest TCR repertoire diversity comprising single-cell
clonotypes occupying compact memory niche "real estate" to recognize viral variants. We predict
that a comparable memory niche strategy is also found in other CD8 repertoires, including $PA_{224\text{-}233}/D^b$,
where digital performers are elicited exclusively due to the sparseness of antigen display [5]
and shall extend to CD4 repertoires directed at pMHC class II epitopes as well. Harnessing this
multidimensional information presents a compelling opportunity to create immunotherapies
involving both TCR and synthetic receptor designs, incorporating necessary survival and persistent
effector functionalities. Consideration of receptor-pMHC binding centricities can likely improve
upon clever AI-based designs [65-67] or other cellular therapies to foster effective clinical translation.
619
620

Methods

Mice and IAV infection

C57BL/6N (B6), B6.129S6-Rag2^{tm1Fwa}N12 (*Rag2$^{-/-}$*), and B6.SJL-*Ptprc^a*/BoyAiTac (CD45.1/1) mice were purchased from Taconic Biosciences, Inc. B6.PL-*Thy1^a*/CyJ (Thy1.1) were obtained from The Jackson Laboratory. CD45.1/1-*Rag2$^{-/-}$* and CD45.1/2-*Rag2$^{-/-}$* were generated by intercrossing CD45.1/1 and *Rag2$^{-/-}$* strains. Thy1.1-*Rag2$^{-/-}$* mice were generated by intercrossing Thy1.1 and *Rag2$^{-/-}$* mice. All mice were housed and bred under specific pathogen-free conditions at the Dana-Farber Cancer Institute (DFCI) Animal Facility, accredited by the Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC). Euthanasia was performed by CO_2 inhalation followed by cervical dislocation. Mice aged 6-10 weeks were used for experiments. Sex-matched mice were used for each experiment. No gender preference was applied, and the sex in each experiment was not deliberately selected.
Mice were infected intranasally with 20 EID_{50} of Influenza A/PR/8/34 virus (PR8, H1N1, Charles River Laboratories) as a primary infection under anesthesia with intraperitoneal injection of ketamine/xylazine (120 mg/kg ketamine, 10 mg/kg xylazine) PR8-infected mice were rechallenged intranasally with $1x10^7$ EID_{50} of serologically distinct X:31, A/Aichi/68 (X31, H3N2, Charles River Laboratories). These viral doses were used consistently across all experiments. All mouse maintenance, breeding, and experimental procedures were approved under Dana-Farber Cancer Institute Institutional Animal Care and Use Committee (IACUC) protocol 04-113.

Cell lines and cell culture

BW5147.3 cells and CD3$\delta\gamma\epsilon\zeta$ pMIY vector were a gift from the Vignali lab (St. Jude Children's Research Hospital, Memphis, Tennessee), and mCD8$\alpha\beta^+$ BW5147.3 cells were generated as previously described [22]. The cells were maintained in D10 [DMEM medium (Gibco), 10% fetal bovine serum (FBS) (Sigma-Aldrich), 100 IU Penicillin and 100μg/mL Streptomycin (Corning), 2 mM L-glutamine (Corning) and 55 μM 2-mercaptoethanol (Gibco)] with 400 μg/mL Hygromycin B (Gibco) and 400 μg/mL Geneticin (Gibco). R8 cells were maintained in R10 (RPMI-1640 (Gibco), 10% FBS, 100 IU Penicillin and 100 μg/mL Streptomycin, and 55 μM 2-mercaptoethanol). All cells were grown in a 5% CO_2 incubator at 37°C.

Cell isolation, flow cytometry and cell sorting

Mediastinal lymph nodes (mLNs) were harvested, mashed with a 3 mL syringe on a 24-well plate with RPMI-1640, filtered with 80 μm mesh, washed with RPMI-1640, and resuspended with FACS buffer (2% FBS/0.05% NaN$_3$/PBS) or appropriate buffer as indicated for the following experiments. Cell suspensions from mLN were first treated with anti-mouse CD16/CD32 mAbs in FACS buffer to block FcR binding for 10 minutes at 4°C and then stained with antibodies indicated in each figure caption in FACS buffer for 20 minutes at 4°C. For intracellular staining, cells were fixed with 4% PFA/PBS and permeabilized using permeabilization buffer following the manufacturer's protocol (Invitrogen). After washing, cells were incubated with anti-mouse TCF-1 mAb for 30 minutes at 4 °C. For scRNA-seq of NP$_{366-374}$ /H-2D^b-specific memory T cells, cell

suspensions from pooled mLNs from 10 mice was stained with APC-conjugated $NP_{366-374}$/H-2D^b
tetramer (MBL International Corporation) for 30 min at RT followed by staining with Zombie
Aqua, and CD8β and CD44 mAb. $NP_{366-374}$-tetramer$^+$ cells were sorted after gating on Zombie
Aqua$^-$ CD8β$^+$ CD44$^+$ cells. For scRNA-seq of non-tetramer sorted T cells at memory and recall
phase, Zombie Aqua$^-$ CD8b$^+$ and CD44$^+$ cells were sorted from pooled mLNs from four and two
mice, respectively. Rg T cells for bulk RNA-Seq were isolated from individual RgC mouse mLN
as Zombie$^-$ CD8β$^+$ cells in combination with congenic markers CD45.1, CD45.2, and Thy1.1. All
antibodies and the concentration used are listed in Supplementary Data 6. All cells were analyzed
on a BD LSR Fortessa™ Cell Analyzer (BD Biosciences) or sorted on a FACS Aria II (BD
Biosciences). Data were analyzed with FlowJo software (FlowJo, LLC).
677
678
679 **Single-cell RNASeq library construction and sequencing**
680
$NP_{366-374}$-specific memory CD8$^+$ cells were sorted into 500 mL R10 in 1.5 mL tube. Cell density
was adjusted to analyze 10,000 cells per sample. Single-cell gene expression (scRNA-seq) was
performed using Chromium Single Cell 5' Library and Gel Bead Kit with V(D)J enrichment kit
(10x Genomics). Gel Bead-in-Emulsion reverse transcription (GEM-RT) reaction, clean-up, and
PCR amplification were performed to obtain cDNA for library generation. Single cell sequencing
libraries were generated using 5'GEM-X v1 according to manufacturer's protocol. Libraries were
uniquely indexed for multiplexed sequencing. Libraries were single i7-indexed using the
Chromium i7 Multiplex kit (indices listed in table). Following isolation and clean-up of library
DNA, concentration and fragment size were assessed using an Agilent Bioanalyzer and
quantification by Qubit analysis (ThermoFisher). 150bp paired-end sequencing was performed on
a Hiseq 3000 utilizing the following allocation parameters per lane: two 5'GEM-X libraries (2 x
40% of reads) and two V(D)J libraries (2 x 10% of reads).
693
694
695 **Single-cell RNA-seq data analysis**
696
Gene expression and TCR FASTQ files were processed using CellRanger version 3.1.0 [68] using
the GRCm38 mouse genome reference. The R package Seurat (version 5.1.0) [69] was used to filter
out cells, cluster the data using 7 principal components, and perform downstream analyses. Cells
were required to have less than 10% of mitochondrial content, between 5 and 7,000 features, and
a minimum of 1,000 UMIs. Potential doublets were removed using the Python package "scrublet"
with default parameters [70]. Additionally, only those cell with exactly one alpha TCR chain and
exactly one beta TCR chain were retained. Gene signature scores were calculated using the
package AUCell (version 1.26) [71]. The first principal component was used to generate a cell density
ridge plot. UMAP approach was used for data visualization [72].
706
707
708 **Verification of memory clusters**
709
scRNA-seq was performed on non-tetramer sorted (NS) memory CD8 T cells from mLNs, and
these were processed as described above with cluster resolution set to "0.2". Previously defined
gene sets associated with T_{CM} and T_{EM} cells [73] and AUCell were used to determine which clusters

in the NS data constituted T_{CM} and T_{EM} clusters. Differential gene expression between NS T_{EM}
clusters and T_{CM} clusters was performed using "FindMarkers" from the "Seurat" package with
"min.pct" set to 0.3. Final T_{EM} and T_{CM} signatures were those with an adjusted p-value <= 0.05,
and, for T_{EM}, (pct.1 – pct.2) >= 0.3 and pct.2 < 0.2 and, for T_{CM}, (pct.1 – pct.2) <= -0.3 and pct.1
< 0.2. The final T_{EM} gene signature includes *"Gzmk", "Itga1", "S100a4", "Id2", "S100a6", "Mdfic",*
"Rgs1", "Fasl", "Pdcd1", "Ly6a", and *"AU020206"*. The final T_{CM} gene signature includes *"Sell",*
"Satb1", "Tmem108", "Txk", "Nsg2", "S1pr1", "Cmah", "Pde2a", "Ccr7", "Treml2", "Rras2",
"Dapl1", "Il6st", "Pdlim1", "Id3", and *"Dennd2d"*. These gene signatures were used to verify T_{CM}
and T_{EM} clusters on scRNA-seq of $NP_{366-374}/D^b$-specific memory CD8 T cells using "AUCell".

Memory clonotype definition

726 T_{CM} and T_{EM} clonotypes were broadly defined as those in which more than 70% of cells localized
727 to the polarized T_{CM} or T_{EM} cluster, respectively. T_{BP} clonotypes were those in which fewer than
728 70% of cells localized to either cluster. Polarized clonotypes manifest statistically distinct and non-
729 random distributions in contrast to bipolar clonotypes.

Clonotype origin

Clonotypes NP201-207 were isolated from pooled mLNs of 10 B6 mice at day 34 following
primary infection with the PR8 strain of IAV. Clonotypes NP101 and NP105 were derived from
pooled mLNs of 2 B6 mice undergoing a memory recall response, 4 days after heterologous
challenge with X31. Clonotypes NP34, NP41, and NP63 were previously identified from lungs 5
days after secondary X31 infection [5]. Clonotypes NP1-3 were isolated from the T_{EM} cluster of
CD44$^+$ memory CD8β$^+$ T cells sorted from pooled mLNs of four B6 mice at day 33 post-infection.

Gene enrichment analysis

744 Gene Set Enrichment analysis was performed on $\log_2$ fold changes using the "GSEA" function of
745 the R package "clusterProfiler" (version 4.12.6) [74] with parameters minGSSize = 6 and
746 pAdjustMethod = "fdr".

RNA velocity analysis

751 Intronic read counts were calculated using "velocyto" (version 0.17.17) [75]. The Python package
752 "scVelo" (version 0.2.5) [76] was used to perform velocity analysis using parameter mode set to
753 'stochastic'.

Generation of TCR sequence tree and analysis
Va and Vb sequence similarities (TCR distances) were calculated using the package "TCRDist3"
(version 0.2.2) [77] with the distance matrices for the two summed. Clonotypes with the TRBV gene

"TRBV12-2+TRBV13-2" were excluded from the analysis due to the absence of this V-gene from
the TCRDist3 database. A distance cladogram was generated using the R package "hclust" using
complete linkage. The R package "pheatmap" (version 1.0.12) was used to create a heatmap of
TCR distances.
763
764
765 **Definition of short and long branch**
766
Short and long branches of the TCR tree are defined based on pairwise TCR sequence distance
metrics. First, the mean of the top 10 shortest pairwise distances was calculated for each clonotype.
Then the threshold value was determined based on the bimodal distribution of the mean distances
for all clonotypes (mean distance = 109.0, Extended Data Fig. 5b). This threshold value was used
to define 'short' and 'long' branches. For example, CT_25, categorized as a short-branch
clonotype, had a mean top 10 distance of 7.8 < 109.0, whereas CT_50, categorized as a long-
branch clonotype, had a corresponding value of 196.4 > 109.0.
774
775
776 **Entropy analysis**
777
778 Shannon entropy of V and J gene usage was calculated using the "Entropy" function of R package
779 DescTools (version 0.99.58). Entropy values were normalized by calculating the maximum
780 possible entropy for given numbers of V and J genes and independent clonotypes using a custom
781 script. The numbers of possible V and J genes were retrieved from CellRanger 3.1.0 VDJ database.
782
783
784 **Retrovirus production and transduction**
785
To generate non-fluorescent TCRab-expressing vectors, the IRES-EGFP site was cut out from
retroviral vector pMSCV-IRES-GFP II (pMIG II, Addgene, #52107) with restriction enzymes and
replaced with synthesized DNA encoding TCRβ-P2A-TCRα. The vector was transfected into Plat-
E packaging cells using the calcium phosphate precipitation method. A transfection mixture was
prepared by combining 35 μg of plasmid DNA, 125 μl of 2.5 M CaCl$_2$, and 1090 μl of 0.1× TE
buffer. After incubation at room temperature for 15 minutes, 1250 μl of 2× HBS buffer (100 mM
HEPES, 281 mM NaCl, 1.5 mM Na$_2$HPO$_4$) was added dropwise to the DNA/CaCl$_2$ solution while
vortexing. The resulting DNA/calcium phosphate precipitate was immediately added to Plat-E
cells. Retrovirus in the cultured supernatant was collected 48 hours post-transfection and frozen at
-80°C until use. Thawed retrovirus was transduced with Retronectin (Takara Bio) using
Retronectin-bound virus infection methods according to the manufacturer's protocol. Retrovirus
was transduced into mCD8αβ$^+$ BW5147.3 cells to generate TCRαβ-expressing cell lines or into
mouse bone marrow (BM) cells to generate Rg mice. To establish TCRαβ-expressing BW cell
lines, the transduced cells were sorted by FACS Aria II (BD Biosciences) to match the TCR
expression based on the cell surface expression of CD3ε. Before all assays were performed, CD3ε
or TCRβ surface level was confirmed to be matched in a group.
802
803
804

Generation of Rg mice

Rg mice were generated as previously described [5,78]. Briefly, BM cells were harvested from CD45.1/1-*Rag2*[-/-], CD45.1/2-*Rag2*[-/-], or Thy1.1-*Rag2*[-/-] mice as indicated in figure legends, and hematopoietic stem cells (HSC) were enriched by EasySep Mouse Hematopoietic Progenitor Cell Isolation Kit (STEMCELL Technologies) followed by expansion in Stem cell medium (StemPro™-34 SFM (Gibco), 5% FBS, 100 IU Penicillin and 100 μg/mL Streptomycin (Corning), 2 mM L-glutamine, with 50 ng/mL mIL-3 (STEMCELL Technologies), 50 ng/mL hIL-6 (STEMCELL Technologies) and 50 ng/mL mSCF (STEMCELL Technologies)) for 3 days. Subsequently, HSC were transduced with retrovirus encoding TCRαβ, cultured in Stem cell medium for 3 days, and then transferred into sublethally irradiated (600 rad) *Rag2*[-/-] mice using a Gamma Cell 40 Cs[137] Irradiator (Thratonics) one day before. Rg mouse blood was analyzed for CD8 development and TCRβ expression by flow cytometry 6 weeks after transplantation, and mice expressing more than 10% of CD8 T cells in CD8[+] cells were used for the generation of RgC mice.

Generation of RgC mice

RgC mice were generated by adoptive transfer of naïve CD8β[+]CD44[−] Rg T cells, isolated from peripheral lymph nodes and spleen of Rg donor mice, into B6 (CD45.2/2, Thy1.2) recipient mice. The number of transferred cells ranged from 0.5 to 10×10^4 per mouse, as indicated in figure captions. To avoid potential immunogenicity associated with fluorescent proteins such as EGFP [79], we used congenic surface markers, such as CD45.1/1, CD45.1/2, and Thy1.1, as identifiers for donor cells, allowing for reliable tracking of individual clones. Mixed RgC mice were generated by co-transferring equal numbers of Rg T cells expressing different TCRs to minimize variability across individual mice. The equal input ratios of each clone were verified by flow cytometry using congenic marker expression. Single RgC mice received Rg T cells expressing a single TCR to eliminate interclonal competition amongst different Rg T cell clonotypes. All RgC mice were intranasally infected with PR8 one day after adoptive transfer. At >30 days post-infection, transferred Rg T cells were analyzed as memory T cells by flow cytometry and RNA-Seq to assess phenotype and gene expression profiles.

TCR Ectodomain Constructs for X-ray Crystallography

The design and expression of TCRαβ heterodimers has been described previously [5,23,80,81]. Briefly, DNA vectors containing TCRα or TCRβ (i.e. V(D)J-C) ectodomains, a HRV3C protease cleavage site [82], a base or acid leucine zipper (LZ) motif to facilitate heterodimer formation upon expression [83], and a stop codon, were designed utilizing human cytomegalovirus promoter. TCRα and β vectors were constructed as separate eukaryotic expression vectors using the pGDom vector from prior work with TCRβ expression in mammalian cells [84,85], retaining the vector signal sequence and encoding the murine Cα-HRV3C-basic-LZ or Cβ-HRV3C-acid-LZ, named as pGDomCα-LZ or pGDomCβ-LZ, respectively. A novel cloning region was inserted between the signal sequence and the start of the C domain utilizing two BsaI type IIS restriction endonuclease sites. This allowed in-frame insertion, using Golden Gate Cloning strategies [86], of fragments encoding only

VJ-α or VDJ-β. These were isolated by PCR amplification from plasmids of full-length constructs
used in retroviral transductions or purchased as double stranded DNA fragments (Azenta) and
subsequently ligated into the respective pGDomCα-LZ or pGDomCβ-LZ vector. TCR clonotypes
with insufficient protein yields were alternately cloned into similar vectors encoding human C-
domains along with the addition of a non-native disulfide between the Cα and Cβ, resulting in a
chimeric construct [87-89]. Annotated DNA and translated amino acid sequences of all constructs are
included in a Supplementary Data 5.
858
859

860 **TCR Protein Expression and Purification**
861

The plasmids containing TCRα and TCRβ constructs were transfected together into a culture
containing $2x10^6$ cells/mL Expi293F glucoseaminotransferase deficient (GnTi⁻) cells
(ThermoFisher) in fresh Expi293 Expression Medium (ThermoFisher) at a final concentration of
0.5 µg vector/mL culture each, using ExpiFectamine transfection reagent (ThermoFisher) and
following the manufacturer's protocol. The Expi293F GnTi⁻ variant produces only high mannose-
modified N-linked glycans, allowing the removal of glycans under native conditions using
endoglycosidase H (EndoH) [85,90]. Transfected cells were cultured at 37°C with rotary shaking at
125 rpm, per manufacturer recommendation, for 7 days for expression of protein. The supernatant
was then collected and clarified by centrifugation and filtration before being passed at least twice
over protein G resin (GammaBind Pl; Cytiva) coupled with the anti-LZ mAb 2H11, utilizing a
gravity flow apparatus at a flow rate of approximately 1 mL/min or slower [81]. Following washing
with 10 column volumes of TBS pH 8.0, bound protein was eluted with 50 mM Citrate, 500 mM
NaCl buffer, pH 2.9 (adjusted with Tris base), in six 9 mL fractions, and neutralized immediately
upon collection with 1 mL (i.e., 1/10 vol) 1M Tris-HCl, pH 8.0. The fractions containing eluted
protein were concentrated and buffer exchanged into TBS and purified via size exclusion
chromatography (SEC) utilizing a HiLoad 16/600 Superdex 200 pg column (S200P; Cytiva).
Fractions containing the TCRαβ heterodimer were concentrated to 1 mg/mL, calculated via
absorbance at 280 nm (A_{280}), prior to digestion with HRV3C protease (1 µg enzyme/50 µg TCR)
at 25°C for 2 hours for complete removal of the LZ. Glycans were then digested with EndoH at
1U enzyme/1 µg TCR with incubation at 25°C for up to 24 hours. Completeness of the digestions
was confirmed with SDS-PAGE. The fully digested protein was then purified with anion-exchange
chromatography utilizing a MonoQ 5/50 column (MonoQ; Cytiva) and eluted as a single peak.
Final protein purity was assessed by non-reducing (NR) and reducing SDS-PAGE prior to
crystallization trials. For crystallization screening, the purified protein was buffer exchanged into
10mM Tris-HCl, pH 8.0, 100 mM NaCl and concentrated to 10 mg/mL, per A_{280} measurements.
For SM use, all constructs were the murine C domain variants with the single C-terminal
interchain disulfide only. Expression and purification mirrored that for structural work with the
following exceptions. Expression utilized the wildtype Expi293F cells (ThermoFisher), retaining
complex glycan modifications. Following elution from 2H11 affinity columns, the proteins were
buffer exchanged into PBS pH 7.4 using SEC chromatography with a 21 mL analytical-scale S200
column (Cytiva). The sample was then concentrated to approximately 1 mg/mL concentration and
flash frozen with LZ regions intact for SM analysis.
894
895
896

pMHC expression, refolding and complex formation, and purification

NP$_{366-374}$ (NP366) peptide (ASNENMETM) was chemically synthesized (United Biosystems, Inc) and NP366/H-2D^b production was executed similarly to that previously described [48,91,92]. Plasmids containing H-2D^b or murine β2m were transformed into BL21-DE3 *E.Coli* for expression. Transformed cells were grown to an OD$_{600}$ of 0.6-0.8 in Luria broth (LB) at 37°C under agitation at which point expression was induced with the addition of 0.5mM IPTG. After 3 hours, cells were harvested by centrifugation, supernatant was discarded and the cell pellet was resuspended in 30 mL of TBS, pH 8.0, 1 mg/mL lysozyme was added, and the resuspension was frozen at -80°C for at least 2 hr. The lysate was then thawed and 0.3 mg/mL of DNAse and EDTA free protease inhibitor (Roche) were added before being frozen again at -80°C > 2 hr or overnight. The lysate was thawed, and inclusion bodies were purified via centrifugation and resuspension cycles, washing twice with TBS, twice with TBS + 0.1% TritonX-100, and once with TBS. The final product was a pure inclusion body pellet, which was dissolved in 10 mL of 5.4M Guanidine HCl, 0.1M Tris, pH 8.0. The solution was centrifuged and the supernatant was collected before the addition of dithiothreitol to 2 mM and storage at -80°C. H-2D^b, β2M, and NP366 peptide were added to 300 mL 20 mM Tris, 8M Urea at a 60:20:10 w/w ratio before serial dialysis (6-8 KDa nominal molecular weight cutoff; Spectrum Laboratories, Inc.) against 4 L of 20 mM Tris + 2 M, 1M, 0.5M and 0M Urea, with each step greater than 2 h or overnight for a total of at least 48 hours dialysis/refold time. The dialyzed refolded protein was then concentrated for purification via SEC (S200P; Cytiva) followed by anion-exchange chromatography (MonoQ; Cytiva). The final purification yielded a mono-dispersed peak and subsequent NR SDS-PAGE revealed two bands, corresponding to H-2D^b and β2M. For crystallization screening, the purified protein was buffer exchanged into 10 mM Tris-HCl, pH 8.0, 100 mM NaCl and concentrated to 10 mg/mL, per A$_{280}$ measurements.

SM assay

Tether geometry and optical tweezers measurements were carried out essentially as detailed in [23], and as shown schematically in Figure (Extended Data Fig. 6a). Streptavidin-coated polystyrene beads (1.29 μm diameter, Spherotech Inc) were connected to biotin terminated 1010-bp double-stranded DNA containing covalently bound 2H11 antibody that binds to the TCRαβ construct paired LZ region. Incubation time and DNA concentration were varied to control tether density around the bead, which was optimized for the binding probability of each individual TCRαβ-pMHC pair. After incubation at RT with DNA tethers, DNA-labeled beads were washed then resuspended in 15 μL of TCRαβ (~0.1 mg/mL) and incubated at RT for an additional 1 hr. The tethered beads were then washed a final time before resuspension in PBS buffer.

A 10-μL flow cell was prepared using two pieces of double-sided sticky tape. Cover-glass surfaces were coated with functionalized PEG silane 99% mixed with 1% biotin-PEG silane like previously described work [23]. Casein blocker (0.1 mg/mL) was introduced to the flow cell for 10 minutes then washed out with PBS buffer. Streptavidin (0.5 mg/mL) was introduced into the flow cell, incubated for 30 min, and washed with PBS buffer. Biotinylated pMHC (~0.05 mg/mL) was introduced and incubated for 30 min and washed again with PBS buffer. Finally, tethered bead solution was introduced to the flow cell for bond lifetime measurements.

Tethers were formed by scanning a bead near the cover-glass surface using 300 nm steps
of the piezo stage in the x-y plane, pausing for several seconds between stage movements to allow
for TCRαβ-pMHC bond formation (Extended Data Fig. 6b). Successful tether formation was
indicated by bead displacement relative to the trap center. After rupture, beads were mapped for
position sensing using automated procedures followed by stiffness calibration. Control
experiments were TCRαβ was omitted from the assay resulted in minimal, atypical tether
formation. Binding probability can be assessed by comparing the number of stage movements
which resulted in a binding event to the total number of stage movements.
951
952

953 **Peptide stimulation and IL-2 measurement**
954

1×10^5 NP-specific TCR-transduced BW cells were cultured in D10 with 1μg/mL WT-NP$_{366-374}$,
1mg/mL NP$_{366-374}$ peptide variants, or titrated theses peptides (from 1×10^{-5} to 1×10^4 ng/mL) at
37°C for 16-18 hours overnight. 1×10^5 R8 cells used as APC were treated with Mitomycin C
(Sigma-Aldrich) and washed with PBS three times prior to use. IL-2 concentration in culture
supernatant was measured by ELISA assay according to the manufacturer's instruction. EC$_{50}$ of
peptide response to each BW cells was calculated by Prism 10 (GraphPad Software).
961
962

963 **Tetramer activation assay and pTyr analysis**
964

Detection of CD3 subunits, pTyr, and CD3ζ pTyr142 pre- and post-tetramer activation was carried
out using 5×10^6 cells/ml in DMEM. WT NP$_{366-374}$/D^b tetramer (0.75 μg/ml final concentration)
was added to the NP202 or NP105 cell samples matched for cell-surface TCR level expression
prior to use. Control samples were simultaneously prepared for each cell line without the addition
of tetramer at time zero. All cell samples were incubated on ice for 20 min and then washed with
DMEM to remove unbound tetramer. The cells were resuspended at a final concentration of 2 x
10^6 cells per 0.1 mL and incubated at 37∘C for 1 min. Then, 1 mL of ice-cold PBS was added to
the cells and the cells were pelleted at 1200 rpm for 5 min. The PBS wash was removed and lysis
buffer (1% Triton X-100, 0.1% SDS, 50 mM Tris (pH 7.4), 150 mM NaCl, 2 mM NaVO3, 1 mM
N-ethylmaleimide, and Roche cØmplete protease cocktail) was added to the cell pellet. The
resuspended samples were immediately placed on dry ice to await further processing.
For analysis of CD3 and phosphorylated proteins, cell samples were then thawed for 10
min on ice and centrifuged at 13,000 rpm, 4∘C for 15 min. The clarified lysate was transferred to
a clean tube. Aliquots were separated non-reducing with 4-12% bis-tris NuPAGE gels, transferred
to polyvinylidene difluoride (PVDF) membrane, and detected with anti-CD3ζ pTyr142 (clone EM-
54), and anti-CD3ζ (clone H146-968). The western blots were imaged using a Bio-Rad ChemiDoc
imaging system. Band density was measured using the Image Lab software, with band densities
normalized to the median within each experimental set to mitigate antibody staining artifacts The
level of phosphorylated CD3ζ Tyr142 was determined in proportion to CD3ζ for each sample.
Three biological replicates were prepared for each activation assay with tetramer activation being
carried out on three separate days with three separate cell culture preparations. To examine overall
phosphotyrosine levels, the cell lysates were separated on a reducing gel, transferred to PVDF
membrane, and assayed for pTyr (clone 4G10).
988

 Bulk RNA-seq analyses

991 Following phenotyping and gating for lymph node cells of retrogenic origin (see table below),
992 cells were sorted into 1.5 mL tubes containing TCL buffer x2 (350 µL; Qiagen) supplemented with
993 2-mercaptoethanol (1%).

995 Gating parameters and cell yields for retrogenic cells:

997 C101   Gating: viable $CD8\beta^+CD45.1^+/CD45.2^+$;   host: B6 $(CD45.2^+)$

Mouse:	1	2	3	4	5	6	7
Cell yield:	5,517	2,852	7,515	8,758	16,051	992	2,658

1000

1001 C105   Gating: viable $CD8\beta^+CD45.1^+$;   host: B6 $(CD45.2^+)$

Mouse:	1A	2	4	5	6	7	1B $(CD45.1^{int})$
Cell yield:	1,222	275	227	504	5,517	7,165	7,305

1004

1005 C202   Gating: viable $CD8\beta^+Thy1.1^+$;   host: B6 $(Thy1.2^+)$

Mouse:	1	2	3	5	6	7
Cell yield:	6,695	1,183	2,172	3,108	7,011	5,065

1008

1009 C204   Gating: viable $CD8\beta^+CD45.1^+$;   host: B6 $(CD45.2^+)$

Mouse:	2	3	5	6
Cell yield:	3.088	347	5,203	5,418

1012

1013 C207   Gating: viable $CD8\beta^+CD45.1^+/CD45.2^+$;   host: B6 $(CD45.2^+)$

Mouse:	2	3	4	5
Cell yield:	2,306	11,467	10,620	7,549

1016

Total RNA was extracted from each sample (n = 28) using the Arcturus Pico Pure RNA kit
(ThermoFisher) following the manufacturer's protocol including on-column RNase-free DNase
treatment (27 Kunitz units/sample; Qiagen), and eluting into 11 µL elution buffer, snap freezing at
-80ºC and storage at the same temperature until library preparation and sequencing (Admera
Health, LLC). Following initial RNA preparation quality control, libraries were prepared using the
SMART-Seq v4 Ultra Low Input kit with Tagmentation (Takara Bio USA, Inc.), Illumina adapters
added, and sequenced on the NovaSeq X Plus platform (2 x 250 bp paired end sequencing, 40 x
10^6 reads/sample). The paired output fasta files for each sample were checked for quality using
FastQC, and adapters trimmed using FastqMcf and Cutadapt [93]. Contaminating non-polyA-tailed
RNA sequence, mitochondrial genome sequences, ribosomal RNAs and transfer RNAs were then
removed using Bowtie2 [94] and the resulting filtered reads aligned to the mouse reference genome
assembly GRCm39 using STAR (v2.7.3a) [95]. HTSeq-count was then used to count the reads
aligned to each gene. Gene expression in transcripts per million (tpm) normalizing for transcript
length and library size, was obtained using the DESeq2 package under Bioconductor version 3.21
running on the R version 4.5 platform. All further processing for hierarchical clustering and heat
maps, and differential gene expression (DESeq2), was performed on the same platform. Note that
for Extended Data Fig. 10e, the heat maps represent transcripts per million (tpm) for a reduced
panel of transcripts representing genes validated as upregulated on T cell activation. In this

instance, tpm can be used as a broad measure for comparison of the two data sets (C101/C105 and
C202/C204/C207) since all samples were of similar origin (mLN CD8$^+$ retrogenic T cells), and
identical RNA preparation, sequencing, and quality control statistics (e.g. mitochondrial transcript
representation) [96]. All data was organized/plotted using MS-Excel or plotted using PSI-Plot version
9 (Poly Software International).
1040
1041
1042 **Crystallization**
1043
Each NP TCR was mixed with NP366-H-2D^b (NP$_{366-374}$/D^b) in a 1:1 molar ratio. The mixed TCR-
pMHC solution was concentrated to about 10-20mg/ml. Sitting drop vapor diffusion technique
in 96-well CrystalQuick plates (Greiner) was used for crystallization trials with the help of a
Mosquito nanoliter liquid handler (TTP LabTech) at 16°C. For each condition, 0.4 mL of
crystallization screening formulation was added to 0.4 ml of concentrated TCR-pMHC solution;
the mixture was equilibrated against 140 mL of the same screening buffer in each reservoir well.
The crystallization screens used were MCSG-1-4 (Anatrace), Peg/Ion (Hampton Research) and
Top96 (Molecular Dimensions). Crystals appeared under the conditions 1-8 as listed for each
construct:
1053
1054 1) NP101-NP366/H-2D^b (mouse/human hybrid TCR) = NP101-NP$_{336-374}$/D^b
1055   0.2 M Ammonium iodide pH 6.2, 20% (w/v) PEG 3,350.
1056
1057 2) NP105-NP366/H-2D^b (mouse/human hybrid TCR) = NP105-NP$_{336-374}$/D^b
1058   0.1 M Sodium citrate, 5% (v/v) 2-Propanol, 20% (w/v) PEG 4,000,
1059
1060 3) NP202 apo form (mouse TCR)
1061   0.2 M Ammonium nitrate pH 6.2, 20% (w/v) PEG 3,350.
1062
1063 4) NP202-NP366/H-2D^b (mouse TCR) = NP202-NP$_{336-374}$/D^b
1064   0.1 M Tris hydrochloride pH 8.5, 8% (w/v) PEG 8,000.
1065
1066 5) NP203-NP366/H-2D^b (mouse/human hybrid TCR) = NP203-NP$_{336-374}$/D^b
1067   0.2 M Ammonium fluoride pH 6.2, 20% (w/v) PEG 3,350.
1068
1069 6) NP204-NP366/H-2D^b (mouse TCR) = NP204-NP$_{336-374}$/D^b
1070   1% (w/v) Tryptone, 0.05 M HEPES sodium pH 7.0, 12% (w/v) PEG 3,350.
1071
1072 7) NP207 apo form (mouse TCR)
1073   0.2 M Potassium Nitrate: HCl pH 6.9, 20% (w/v) PEG 3,350.
1074
1075 8) NP207-NP366/H-2D^b (mouse/human hybrid TCR) = NP207-NP$_{336-374}$/D^b
1076   0.6 M Sodium Chloride, 0.1 M MES:NaOH pH 6.5, 20% (w/v) PEG 4,000.
1077
1078
1079
1080

X-ray diffraction data collection and processing

All crystals were harvested and treated with a cryoprotectant solution (25% glycerol or 25% ethylene glycol in mother liquor) and then flash-frozen in liquid nitrogen before X-ray diffraction data collection. Those data were collected at 100 K from the cryocooled crystals at the 19-ID at the 19-ID (SBC) beamline of the Advanced Photon Source at Argonne National Laboratory or 19-ID (NYX) beamline or 17-ID-2 (FMX) beamline of National Synchrotron Light Source II at the Brookhaven National Laboratory. The intensities of each data set were integrated, scaled, and merged with the HKL-3000 program suite [97]. The structure of NP202-NP366-H-2D^b (NP202-NP$_{366-374}$/D^b) was first determined using the molecular replacement (MR) method [98], in which the murine P14 TCR and the H-2D^b from their complex (PDB code: 5M01) were used as search templates (Supplementary Table 2). The following structural determination of NP TCR apo forms and other complexes were performed with the same MR method using NP202 TCR and H-2D^b obtained in NP202-NP366-H-2D^b complex (NP202-NP$_{366-374}$/D^b) that was determined and refined in the first place. All the models from structural determination were rebuilt using the program Coot [99]. The final models was refined using the program phenix.refine [100]. Structural validation of each model was performed using the program MolProbity [101].

Molecular dynamics simulation

Molecular dynamics simulation under force was performed based on the method described in [40]. Structure preparation, preparatory simulation, and trajectory analysis were performed by using CHARMM [102] and production runs were performed by using OpenMM [103]. Briefly, crystal structures of NP202, NP101, NP207, and NP203 complexed with NP$_{366-374}$/H-2D^b were prepared by converting the Cα, and Cβ sequences from human (used for x-ray crystallography) to mouse (used for other experiments). The structure of NP202 had the full mouse sequence except for S184 of Cβ, which was mutated to C184 for mouse. The resulting structure of the C-module was aligned and copied to NP101, NP207, and NP203 to generate the mouse versions. 5-residue long linkers were added to the C-terminal ends of MHC and the TCR based on their amino acid sequences so that the last residues were Y283 for MHC, T214 for Cα, and A245 for Cβ, respectively. C$_\alpha$ atoms of the terminal residues were harmonically restrained at 168-Å extensions to apply fluctuating loads that are comparable in magnitude to those used in single-molecule experiments. We modified the contact analysis method developed in [26] to examine intradomain contacts as shown in Supplementary Discussion Item 5. Additional simulations and more extensive analyses will be published elsewhere.

Statistics

Statistical tests used throughout this study are indicated in figure captions or the main text. Paired t-tests were used for RgC mouse and p-tyrosine experiments. Chi-squared tests were employed for analysis of T_{CM} vs. T_{EM} clonotype distributions, Shannon entropy, and RNA-seq clustering. Mann–Whitney U-tests evaluated differences in cell counts. Trend tests assessed graded gene expression across cell–clonotype categories. Two-sided Fisher's exact tests were used for cross-reactive

1126 clonotype comparisons. Gene Set Enrichment Analysis (GSEA) was evaluated by a permutation
1127 test.
1128
1129

Data and materials availability

Sequence files for scRNA-Seq of $NP_{366-374}/D^b$-specific memory CD8 T cells, non-tetramer sorted
CD8 T cells at memory phase, and at recall response have been deposited in NCBI Gene
Expression Omnibus (GEO) under accession GSE302680. Bulk RNAseq data are accession
GSE302856. PDB accession code will be available at the time of publication for all X-ray
crystallographic structures. Materials and protocols are available on request to scientists through
the corresponding authors.

1137
1138 **References**
1139

1   Kaech, S. M. & Cui, W. Transcriptional control of effector and memory CD8+ T cell
differentiation. *Nat Rev Immunol* **12**, 749-761 (2012).
https://doi.org/10.1038/nri3307
2   Wherry, E. J. & Ahmed, R. Memory CD8 T-cell differentiation during viral infection. *J*
Virol **78**, 5535-5545 (2004). https://doi.org/10.1128/JVI.78.11.5535-5545.2004
3   Sallusto, F., Geginat, J. & Lanzavecchia, A. Central memory and effector memory T
cell subsets: function, generation, and maintenance. *Annu Rev Immunol* **22**, 745-
763 (2004). https://doi.org/10.1146/annurev.immunol.22.012703.104702
4   Jameson, S. C. & Masopust, D. Understanding Subset Diversity in T Cell Memory.
Immunity **48**, 214-226 (2018). https://doi.org/10.1016/j.immuni.2018.02.010
5   Akitsu, A. *et al.* Parsing digital or analog TCR performance through piconewton
forces. *Sci Adv* **10**, eado4313 (2024). https://doi.org/10.1126/sciadv.ado4313
6   La Gruta, N. L. *et al.* A virus-specific CD8+ T cell immunodominance hierarchy
determined by antigen dose and precursor frequencies. *Proc Natl Acad Sci U S A*
103, 994-999 (2006). https://doi.org/10.1073/pnas.0510429103
7   Zhou, X. *et al.* Differentiation and persistence of memory CD8(+) T cells depend on T
cell factor 1. *Immunity* **33**, 229-240 (2010).
https://doi.org/10.1016/j.immuni.2010.08.002
8   Zhou, X. & Xue, H. H. Cutting edge: generation of memory precursors and functional
memory CD8+ T cells depends on T cell factor-1 and lymphoid enhancer-binding
factor-1. *J Immunol* **189**, 2722-2726 (2012).
https://doi.org/10.4049/jimmunol.1201150
9   Yang, C. Y. *et al.* The transcriptional regulators Id2 and Id3 control the formation of
distinct memory CD8+ T cell subsets. *Nat Immunol* **12**, 1221-1229 (2011).
https://doi.org/10.1038/ni.2158
10   Wherry, E. J. *et al.* Lineage relationship and protective immunity of memory CD8 T
cell subsets. *Nat Immunol* **4**, 225-234 (2003). https://doi.org/10.1038/ni889

11 Jeannet, G. *et al*. Essential role of the Wnt pathway effector Tcf-1 for the
establishment of functional CD8 T cell memory. *Proc Natl Acad Sci U S A* **107**, 9777-
9782 (2010). https://doi.org/10.1073/pnas.0914127107
12 Cannarile, M. A. *et al*. Transcriptional regulator Id2 mediates CD8+ T cell immunity.
Nat Immunol **7**, 1317-1325 (2006). https://doi.org/10.1038/ni1403
13 Omilusik, K. D. *et al*. Sustained Id2 regulation of E proteins is required for terminal
differentiation of effector CD8(+) T cells. *J Exp Med* **215**, 773-783 (2018).
https://doi.org/10.1084/jem.20171584
14 Hamilton, S. E. & Jameson, S. C. CD8 T cell memory: it takes all kinds. *Front
Immunol* **3**, 353 (2012). https://doi.org/10.3389/fimmu.2012.00353
15 van der Windt, G. J. & Pearce, E. L. Metabolic switching and fuel choice during T-cell
differentiation and memory development. *Immunol Rev* **249**, 27-42 (2012).
https://doi.org/10.1111/j.1600-065X.2012.01150.x
16 Marzo, A. L. *et al*. Initial T cell frequency dictates memory CD8+ T cell lineage
commitment. *Nat Immunol* **6**, 793-799 (2005). https://doi.org/10.1038/ni1227
17 Obar, J. J. & Lefrancois, L. Early signals during CD8 T cell priming regulate the
generation of central memory cells. *J Immunol* **185**, 263-272 (2010).
https://doi.org/10.4049/jimmunol.1000492
18 Badovinac, V. P., Haring, J. S. & Harty, J. T. Initial T cell receptor transgenic cell
precursor frequency dictates critical aspects of the CD8(+) T cell response to
infection. *Immunity* **26**, 827-841 (2007).
https://doi.org/10.1016/j.immuni.2007.04.013
19 Daniels, M. A. & Teixeiro, E. TCR Signaling in T Cell Memory. *Front Immunol* **6**, 617
(2015). https://doi.org/10.3389/fimmu.2015.00617
20 Smith-Garvin, J. E. *et al*. T-cell receptor signals direct the composition and function
of the memory CD8+ T-cell pool. *Blood* **116**, 5548-5559 (2010).
https://doi.org/10.1182/blood-2010-06-292748
21 Solouki, S. *et al*. TCR Signal Strength and Antigen Affinity Regulate CD8(+) Memory T
Cells. *J Immunol* **205**, 1217-1227 (2020). https://doi.org/10.4049/jimmunol.1901167
22 Brazin, K. N. *et al*. The T Cell Antigen Receptor alpha Transmembrane Domain
Coordinates Triggering through Regulation of Bilayer Immersion and CD3 Subunit
Associations. *Immunity* **49**, 829-841 e826 (2018).
https://doi.org/10.1016/j.immuni.2018.09.007
23 Das, D. K. *et al*. Force-dependent transition in the T-cell receptor beta-subunit
allosterically regulates peptide discrimination and pMHC bond lifetime. *Proc Natl
Acad Sci U S A* **112**, 1517-1522 (2015). https://doi.org/10.1073/pnas.1424829112
24 Feng, Y. *et al*. Mechanosensing drives acuity of alphabeta T-cell recognition. *Proc
Natl Acad Sci U S A* **114**, E8204-E8213 (2017).
https://doi.org/10.1073/pnas.1703559114
25 Feng, Y., Reinherz, E. L. & Lang, M. J. alphabeta T Cell Receptor Mechanosensing
Forces out Serial Engagement. *Trends Immunol* **39**, 596-609 (2018).
https://doi.org/10.1016/j.it.2018.05.005
26 Hwang, W., Mallis, R. J., Lang, M. J. & Reinherz, E. L. The alphabetaTCR
mechanosensor exploits dynamic ectodomain allostery to optimize its ligand

recognition site. *Proc Natl Acad Sci U S A* **117**, 21336-21345 (2020).
https://doi.org/10.1073/pnas.2005899117

27 Kim, S. T. *et al.* The alphabeta T cell receptor is an anisotropic mechanosensor. *J Biol*
Chem **284**, 31028-31037 (2009). https://doi.org/10.1074/jbc.M109.052712

28 Liu, B., Chen, W., Evavold, B. D. & Zhu, C. Accumulation of dynamic catch bonds
between TCR and agonist peptide-MHC triggers T cell signaling. *Cell* **157**, 357-368
(2014). https://doi.org/10.1016/j.cell.2014.02.053

29 Wu, P. *et al.* Mechano-regulation of Peptide-MHC Class I Conformations
Determines TCR Antigen Recognition. *Mol Cell* **73**, 1015-1027 e1017 (2019).
https://doi.org/10.1016/j.molcel.2018.12.018

30 Zhao, X. *et al.* Tuning T cell receptor sensitivity through catch bond engineering.
Science **376**, eabl5282 (2022). https://doi.org/10.1126/science.abl5282

31 Mallis, R. J. *et al.* Biophysical and Structural Features of alphabetaT-Cell Receptor
Mechanosensing: A Paradigmatic Shift in Understanding T-Cell Activation. *Immunol*
Rev **329**, e13432 (2025). https://doi.org/10.1111/imr.13432

32 Hu, Y., Duan, Y. & Salaita, K. DNA Nanotechnology for Investigating Mechanical
Signaling in the Immune System. *Angew Chem Int Ed Engl* **62**, e202302967 (2023).
https://doi.org/10.1002/anie.202302967

33 Dash, P. *et al.* Quantifiable predictive features define epitope-specific T cell
receptor repertoires. *Nature* **547**, 89-93 (2017). https://doi.org/10.1038/nature22383

34 Sridhar, S. *et al.* Cellular immune correlates of protection against symptomatic
pandemic influenza. *Nat Med* **19**, 1305-1312 (2013).
https://doi.org/10.1038/nm.3350

35 Valkenburg, S. A. *et al.* Molecular basis for universal HLA-A*0201-restricted CD8+ T-
cell immunity against influenza viruses. *Proc Natl Acad Sci U S A* **113**, 4440-4445
(2016). https://doi.org/10.1073/pnas.1603106113

36 Zhong, W. *et al.* Significant impact of sequence variations in the nucleoprotein on
CD8 T cell-mediated cross-protection against influenza A virus infections. *PLoS One*
5, e10583 (2010). https://doi.org/10.1371/journal.pone.0010583

37 Young, A. C., Zhang, W., Sacchettini, J. C. & Nathenson, S. G. The three-dimensional
structure of H-2Db at 2.4 A resolution: implications for antigen-determinant
selection. *Cell* **76**, 39-50 (1994). https://doi.org/10.1016/0092-8674(94)90171-6

38 Garcia, K. C. *et al.* An alphabeta T cell receptor structure at 2.5 A and its orientation
in the TCR-MHC complex. *Science* **274**, 209-219 (1996).
https://doi.org/10.1126/science.274.5285.209

39 Chang-Gonzalez, A. C. *et al.* Load-based divergence in the dynamic allostery of two
TCRs recognizing the same pMHC. *Elife* **13** (2025).
https://doi.org/10.7554/eLife.104280

40 Chang-Gonzalez, A. C., Mallis, R. J., Lang, M. J., Reinherz, E. L. & Hwang, W.
Asymmetric framework motion of TCRalphabeta controls load-dependent peptide
discrimination. *Elife* **13** (2024). https://doi.org/10.7554/eLife.91881

41 Chan, A. C., Irving, B. A., Fraser, J. D. & Weiss, A. The zeta chain is associated with a
tyrosine kinase and upon T-cell antigen receptor stimulation associates with ZAP-

70, a 70-kDa tyrosine phosphoprotein. *Proc Natl Acad Sci U S A* **88**, 9166-9170
(1991). https://doi.org/10.1073/pnas.88.20.9166

42 Sen, D. R. *et al.* The epigenetic landscape of T cell exhaustion. *Science* **354**, 1165-
1169 (2016). https://doi.org/10.1126/science.aae0491

43 Kurtulus, S. *et al.* Checkpoint Blockade Immunotherapy Induces Dynamic Changes
in PD-1(-)CD8(+) Tumor-Infiltrating T Cells. *Immunity* **50**, 181-194 e186 (2019).
https://doi.org/10.1016/j.immuni.2018.11.014

44 Abdullah, L. *et al.* The endogenous antigen-specific CD8(+) T cell repertoire is
composed of unbiased and biased clonotypes with differential fate commitments.
Immunity **58**, 601-615 e609 (2025). https://doi.org/10.1016/j.immuni.2025.02.001

45 Notti, R. Q. *et al.* The resting and ligand-bound states of the membrane-embedded
human T-cell receptor-CD3 complex. *bioRxiv* (2024).
https://doi.org/10.1101/2023.08.22.554360

46 von Boehmer, H. Unique features of the pre-T-cell receptor alpha-chain: not just a
surrogate. *Nat Rev Immunol* **5**, 571-577 (2005). https://doi.org/10.1038/nri1636

47 Das, D. K. *et al.* Pre-T Cell Receptors (Pre-TCRs) Leverage Vbeta Complementarity
Determining Regions (CDRs) and Hydrophobic Patch in Mechanosensing Thymic
Self-ligands. *J Biol Chem* **291**, 25292-25305 (2016).
https://doi.org/10.1074/jbc.M116.752865

48 Mallis, R. J., Arthanari, H., Lang, M. J., Reinherz, E. L. & Wagner, G. NMR-directed
design of pre-TCRbeta and pMHC molecules implies a distinct geometry for pre-TCR
relative to alphabetaTCR recognition of pMHC. *J Biol Chem* **293**, 754-766 (2018).
https://doi.org/10.1074/jbc.M117.813493

49 Li, X. *et al.* Pre-T cell receptors topologically sample self-ligands during thymocyte
beta-selection. *Science* **371**, 181-185 (2021).
https://doi.org/10.1126/science.abe0918

50 Borowski, C., Li, X., Aifantis, I., Gounari, F. & von Boehmer, H. Pre-TCRalpha and
TCRalpha are not interchangeable partners of TCRbeta during T lymphocyte
development. *J Exp Med* **199**, 607-615 (2004).
https://doi.org/10.1084/jem.20031973

51 Gounari, F. & Khazaie, K. TCF-1: a maverick in T cell development and function. *Nat
Immunol* **23**, 671-678 (2022). https://doi.org/10.1038/s41590-022-01194-2

52 Hong, C., Luckey, M. A. & Park, J. H. Intrathymic IL-7: the where, when, and why of IL-
7 signaling during T cell development. *Semin Immunol* **24**, 151-158 (2012).
https://doi.org/10.1016/j.smim.2012.02.002

53 Kaech, S. M. *et al.* Selective expression of the interleukin 7 receptor identifies
effector CD8 T cells that give rise to long-lived memory cells. *Nat Immunol* **4**, 1191-
1198 (2003). https://doi.org/10.1038/ni1009

54 Miyazaki, M. *et al.* The opposing roles of the transcription factor E2A and its
antagonist Id3 that orchestrate and enforce the naive fate of T cells. *Nat Immunol*
12, 992-1001 (2011). https://doi.org/10.1038/ni.2086

55 Zhang, N., Hartig, H., Dzhagalov, I., Draper, D. & He, Y. W. The role of apoptosis in the
development and function of T lymphocytes. *Cell Res* **15**, 749-769 (2005).
https://doi.org/10.1038/sj.cr.7290345

56 Dunkle, A., Dzhagalov, I., Gordy, C. & He, Y. W. Transfer of CD8+ T cell memory using
Bcl-2 as a marker. *J Immunol* **190**, 940-947 (2013).
https://doi.org/10.4049/jimmunol.1103481

57 Bohineust, A., Garcia, Z., Beuneu, H., Lemaitre, F. & Bousso, P. Termination of T cell
priming relies on a phase of unresponsiveness promoting disengagement from
APCs and T cell division. *J Exp Med* **215**, 1481-1492 (2018).
https://doi.org/10.1084/jem.20171708

58 Mempel, T. R., Henrickson, S. E. & Von Andrian, U. H. T-cell priming by dendritic
cells in lymph nodes occurs in three distinct phases. *Nature* **427**, 154-159 (2004).
https://doi.org/10.1038/nature02238

59 Miller, M. J., Wei, S. H., Parker, I. & Cahalan, M. D. Two-photon imaging of
lymphocyte motility and antigen response in intact lymph node. *Science* **296**, 1869-
1873 (2002). https://doi.org/10.1126/science.1070051

60 Ding, Y. H., Baker, B. M., Garboczi, D. N., Biddison, W. E. & Wiley, D. C. Four A6-
TCR/peptide/HLA-A2 structures that generate very different T cell signals are nearly
identical. *Immunity* **11**, 45-56 (1999). https://doi.org/10.1016/s1074-7613(00)80080-
1

61 Hodavance, S. Y., Gareri, C., Torok, R. D. & Rockman, H. A. G Protein-coupled
Receptor Biased Agonism. *J Cardiovasc Pharmacol* **67**, 193-202 (2016).
https://doi.org/10.1097/FJC.0000000000000356

62 Erdogmus, S. *et al*. Helix 8 is the essential structural motif of mechanosensitive
GPCRs. *Nat Commun* **10**, 5784 (2019). https://doi.org/10.1038/s41467-019-13722-0

63 Furness, S. G. & Sexton, P. M. Coding GPCR-G protein specificity. *Cell Res* **27**, 1193-
1194 (2017). https://doi.org/10.1038/cr.2017.92

64 Wu, T. *et al*. Quantification of epitope abundance reveals the effect of direct and
cross-presentation on influenza CTL responses. *Nat Commun* **10**, 2846 (2019).
https://doi.org/10.1038/s41467-019-10661-8

65 Householder, K. D. *et al*. De novo design and structure of a peptide-centric TCR
mimic binding module. *Science* **389**, 375-379 (2025).
https://doi.org/10.1126/science.adv3813

66 Johansen, K. H. *et al*. De novo-designed pMHC binders facilitate T cell-mediated
cytotoxicity toward cancer cells. *Science* **389**, 380-385 (2025).
https://doi.org/10.1126/science.adv0422

67 Liu, B. *et al*. Design of high-specificity binders for peptide-MHC-I complexes.
Science **389**, 386-391 (2025). https://doi.org/10.1126/science.adv0185

68 Zheng, G. X. *et al*. Massively parallel digital transcriptional profiling of single cells.
Nat Commun **8**, 14049 (2017). https://doi.org/10.1038/ncomms14049

69 Hao, Y. *et al*. Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587
e3529 (2021). https://doi.org/10.1016/j.cell.2021.04.048

70 Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational Identification of Cell
Doublets in Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291 e289 (2019).
https://doi.org/10.1016/j.cels.2018.11.005

71 Aibar, S. *et al*. SCENIC: single-cell regulatory network inference and clustering. *Nat
Methods* **14**, 1083-1086 (2017). https://doi.org/10.1038/nmeth.4463

72 McInnes, L., Healy, J., Saul, N. & Großberger, L. UMAP: Uniform Manifold
Approximation and Projection. *Journal of Open Source Software* **3 (29)**, 861 (2018).
https://doi.org/10.21105/joss.00861
73 Hudson, W. H. *et al.* Expression of novel long noncoding RNAs defines virus-specific
effector and memory CD8(+) T cells. *Nat Commun* **10**, 196 (2019).
https://doi.org/10.1038/s41467-018-07956-7
74 Xu, S. *et al.* Using clusterProfiler to characterize multiomics data. *Nat Protoc* **19**,
3292-3320 (2024). https://doi.org/10.1038/s41596-024-01020-z
75 La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494-498 (2018).
https://doi.org/10.1038/s41586-018-0414-6
76 Weiler, P., Lange, M., Klein, M., Pe'er, D. & Theis, F. CellRank 2: unified fate mapping
in multiview single-cell data. *Nat Methods* **21**, 1196-1205 (2024).
https://doi.org/10.1038/s41592-024-02303-9
77 Mayer-Blackwell, K. *et al.* TCR meta-clonotypes for biomarker discovery with
tcrdist3 enabled identification of public, HLA-restricted clusters of SARS-CoV-2
TCRs. *Elife* **10** (2021). https://doi.org/10.7554/eLife.68605
78 Holst, J. *et al.* Generation of T-cell receptor retrogenic mice. *Nat Protoc* **1**, 406-417
(2006). https://doi.org/10.1038/nprot.2006.61
79 Agudo, J. *et al.* GFP-specific CD8 T cells enable targeted cell depletion and
visualization of T-cell interactions. *Nat Biotechnol* **33**, 1287-1292 (2015).
https://doi.org/10.1038/nbt.3386
80 Chang, H. C. *et al.* A general method for facilitating heterodimeric pairing between
two proteins: application to expression of alpha and beta T-cell receptor
extracellular segments. *Proc Natl Acad Sci U S A* **91**, 11408-11412 (1994).
https://doi.org/10.1073/pnas.91.24.11408
81 Liu, J. *et al.* Crystallization of a deglycosylated T cell receptor (TCR) complexed with
an anti-TCR Fab fragment. *J Biol Chem* **271**, 33639-33646 (1996).
https://doi.org/10.1074/jbc.271.52.33639
82 Waugh, D. S. An overview of enzymatic reagents for the removal of affinity tags.
Protein Expr Purif **80**, 283-293 (2011). https://doi.org/10.1016/j.pep.2011.08.005
83 O'Shea, E. K., Lumb, K. J. & Kim, P. S. Peptide 'Velcro': design of a heterodimeric
coiled coil. *Curr Biol* **3**, 658-667 (1993). https://doi.org/10.1016/0960-
9822(93)90063-t
84 Tiller, T., Busse, C. E. & Wardemann, H. Cloning and expression of murine Ig genes
from single B cells. *J Immunol Methods* **350**, 183-193 (2009).
https://doi.org/10.1016/j.jim.2009.08.009
85 Mallis, R. J. *et al.* Efficient and economic protein labeling for NMR in mammalian
expression systems: Application to a preT-cell and T-cell receptor protein. *Protein
Sci* **33**, e4950 (2024). https://doi.org/10.1002/pro.4950
86 Engler, C., Kandzia, R. & Marillonnet, S. A one pot, one step, precision cloning
method with high throughput capability. *PLoS One* **3**, e3647 (2008).
https://doi.org/10.1371/journal.pone.0003647

87   Boulter, J. M. *et al.* Stable, soluble T-cell receptor molecules for crystallization and
therapeutics. *Protein Eng* **16**, 707-711 (2003).
https://doi.org/10.1093/protein/gzg087
88   Yoshida, K. *et al.* The diabetogenic mouse MHC class II molecule I-Ag7 is endowed
with a switch that modulates TCR affinity. *J Clin Invest* **120**, 1578-1590 (2010).
https://doi.org/10.1172/JCI41502
89   Day, E. B. *et al.* Structural basis for enabling T-cell receptor diversity within biased
virus-specific CD8+ T-cell responses. *Proc Natl Acad Sci U S A* **108**, 9536-9541
(2011). https://doi.org/10.1073/pnas.1106851108
90   Reeves, P. J., Callewaert, N., Contreras, R. & Khorana, H. G. Structure and function
in rhodopsin: high-level expression of rhodopsin with restricted and homogeneous
N-glycosylation by a tetracycline-inducible N-acetylglucosaminyltransferase I-
negative HEK293S stable mammalian cell line. *Proc Natl Acad Sci U S A* **99**, 13419-
13424 (2002). https://doi.org/10.1073/pnas.212519299
91   Moody, A. M., Xiong, Y., Chang, H. C. & Reinherz, E. L. The CD8alphabeta co-
receptor on double-positive thymocytes binds with differing affinities to the
products of distinct class I MHC loci. *Eur J Immunol* **31**, 2791-2799 (2001).
https://doi.org/10.1002/1521-4141(200109)31:9<2791::aid-immu2791>3.0.co;2-x
92   Mallis, R. J. *et al.* Pre-TCR ligand binding impacts thymocyte development before
alphabetaTCR expression. *Proc Natl Acad Sci U S A* **112**, 8373-8378 (2015).
https://doi.org/10.1073/pnas.1504971112
93   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *2011* **17**, 3 (2011). https://doi.org/10.14806/ej.17.1.200
94   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat
Methods* **9**, 357-359 (2012). https://doi.org/10.1038/nmeth.1923
95   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013). https://doi.org/10.1093/bioinformatics/bts635
96   Zhao, S., Ye, Z. & Stanton, R. Misuse of RPKM or TPM normalization when comparing
across samples and sequencing protocols. *RNA* **26**, 903-909 (2020).
https://doi.org/10.1261/rna.074922.120
97   Minor, W., Cymborowski, M., Otwinowski, Z. & Chruszcz, M. HKL-3000: the
integration of data reduction and structure solution--from diffraction images to an
initial model in minutes. *Acta Crystallogr D Biol Crystallogr* **62**, 859-866 (2006).
https://doi.org/10.1107/S0907444906019949
98   Vagin, A. & Teplyakov, A. Molecular replacement with MOLREP. *Acta Crystallogr D
Biol Crystallogr* **66**, 22-25 (2010). https://doi.org/10.1107/S0907444909042589
99   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of
Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501 (2010).
https://doi.org/10.1107/S0907444910007493
100   Afonine, P. V. *et al.* Towards automated crystallographic structure refinement with
phenix.refine. *Acta Crystallogr D Biol Crystallogr* **68**, 352-367 (2012).
https://doi.org/10.1107/S0907444912001308

101 Chen, V. B. *et al.* MolProbity: all-atom structure validation for macromolecular
crystallography. *Acta Crystallogr D Biol Crystallogr* **66**, 12-21 (2010).
https://doi.org/10.1107/S0907444909042073
102 Hwang, W. *et al.* CHARMM at 45: Enhancements in Accessibility, Functionality, and
Speed. *J Phys Chem B* **128**, 9976-10042 (2024).
https://doi.org/10.1021/acs.jpcb.4c04100
103 Eastman, P. *et al.* OpenMM 7: Rapid development of high performance algorithms
for molecular dynamics. *PLoS Comput Biol* **13**, e1005659 (2017).
https://doi.org/10.1371/journal.pcbi.1005659

1435
1436
1437
1438

Acknowledgments

We gratefully acknowledge Dr. Jia-huai Wang for carefully reviewing the manuscript. We thank Amelia Seabury for assistance with figure preparation. The use of SBC 19-ID at Argonne National Laboratory and the use of NYX 19-ID and FMX 17ID-2 at Brookhaven National Laboratory were supported by DOE contract no. DE-AC02-06CH11357 and DE-SC0012704, respectively. Monomers and tetramers listed in Supplementary Data 6 were obtained through the NIH Tetramer Core Facility. This work was supported by National Institutes of Health grant 1P01AI143565 (M.J.L, R.J.M, W.H, and E.L.R). Simulations were performed by using computers at the Texas A&M High Performance Research Computing facility.

Author contributions

A.A and K.T contributed to conceptualization, methodology, investigation, formal analysis, visualization, data curation, writing the original draft, and reviewing and editing the manuscript. R.J.M contributed to methodology, investigation, reviewing and editing the manuscript. M.A.B contributed to formal analysis, visualization, data curation, reviewing and editing the manuscript. J.S.D-C, contributed to methodology, investigation, formal analysis, visualization, data curation, reviewing and editing the manuscript. K.N.B contributed to methodology, investigation, formal analysis, visualization, reviewing and editing the manuscript. E.H.K contributed to methodology, investigation, formal analysis, and visualization. A.N.P contributed to methodology, formal analysis, visualization, reviewing and editing the manuscript. S.A contributed to investigation. V.C contributed to investigation and formal analysis. J.J.L, K.I.U, J.K.K, M.B, and C.M.M contributed to investigation. P.H.L contributed to data curation. M.Y.T contributed to data curation, supervision, reviewing and editing the manuscript. W.H contributed to formal analysis, Visualization, Funding acquisition, reviewing, and editing the manuscript. M.J.L contributed to conceptualization, visualization, funding acquisition, supervision, reviewing and editing the manuscript. E.L.R contributed to conceptualization, methodology, visualization, funding acquisition, supervision, writing the original draft, and reviewing and editing the manuscript.

Competing interests

Authors declare that they have no competing interests.

1468

Materials & Correspondence

1472

Fig. 1. TCRαβ clonotypes intrinsically influence memory CD8$^+$ T cell fate. a, UMAP clustering of NP$_{366-374}$-specific memory CD8$^+$ T cells. **b,** Dotplot of select immune-related genes. Circles are proportional to the fraction of cells with expression of the indicated gene. Red indicates high expression relative to other clusters while blue indicates low expression. **c,** Violin plots of T$_{CM}$ and T$_{EM}$ gene signatures generated from an independent experiment and applied to the six clusters. The black dots show the medians. **d,** Correlation matrix of the 40 most abundant clonotypes. Pie slice sizes are proportional to the fraction of the clonotype in each cluster. The barplot above the matrix indicates the number of cells with the given clonotype while the barplot to the left indicates the number of cells in each cluster. **e,** Table of select clonotypes and their distribution across T$_{CM}$ and T$_{EM}$ clusters. T$_{CM}$ and T$_{EM}$ clonotypes are those for which more than 70% of cells are in T$_{CM}$ or T$_{EM}$ cluster, respectively. T$_{BP}$ clonotypes are those for which fewer than 70% of cells are in a T$_{CM}$ or T$_{EM}$ cluster. P-values were generated using a binomial test (with adjustment by the Benjamini-Hochberg procedure) relative to the overall distribution T$_{CM}$ or T$_{EM}$ cells. The prime (') dictates the usage of TRAV16 for CT_33, which differs at two residues from TRAV16N used by NP63 (V$\rightarrow$ M and V$\rightarrow$ E) at sites distal to the pMHC interface. **f,** Blue dots on UMAPs showing which cells bear the indicated clonotypes. **g,** Ridge plot indicating the position of cells across the first principal component. Sets of cells include all cells, cells from T$_{CM}$ clusters, T$_{EM}$ clusters, and select clonotypes. **h,** UMAP with cells colored by the indicated CT category and memory cell types comprising each. T$_{CM}$ and T$_{EM}$ cells are cells in a T$_{CM}$ or T$_{EM}$ cluster, respectively.

Fig. 2. Distinct features of memory T cells according to their polarities. a, Sankey plot of V and J-gene usage by clonotype category. **b,** Stacked barplot of clonotype type distribution binned by cell number. The p-values by chi-squared test for comparison between T_{CM} and T_{EM} are: 1 cell vs overall distribution of cells: ****; p = 7.58 x 10^{-7}, 2-10 cells: **; p = 0.0093, and 11+ cells: p = 0.29. **c,** Boxplot of cell number by clonotype type. P-values for comparisons were calculated by Mann-Whitney-U test. **d,** Phylogram of clonotypes generated by hierarchical clustering of TCR distances with complete linkage. Clonotypes that are sequence similar to multiple other clonotypes ("short branch") are shaded in purple while the remaining clonotypes ("long branch") are in black. Representative T_{EM} clonotypes are highlighted in red, T_{CM} in green, and T_{BP} in blue. **e,** Volcano plot of differential expression of clonotypes in the long branch category versus short branch category. **f,** Gene Set Enrichment Analysis of the differential expression in E using the T_{CM} and T_{EM} gene signatures. **g,** Force vs. lifetime plot for T_{EM} clones NP201 (red curve, N=56), NP202 (magenta curve, N=136); T_{CM} clones NP203 (blue curve, N=58), NP204 (yellow curve, N=110), NP205 (green curve, N=79), NP105 (black curve, N=101); and T_{BP} clones NP101 (orange curve, N=103), NP207 (cyan curve, N=100) And NP63 (pink curve, N=40) with its cognate ligand NP366-374/Db. Error bars indicate SEM. **h,** Tether forming probability for the SM optical tweezers assay under identical conditions of tethers per bead and surface ligand density. Error bars indicate SEM. **i,** Experimental scheme for generating single and mixed retrogenic (RgC) mice using congenic markers (i.e., CD45.1/1, CD45.1/2 or Thy1.1) is detailed in Methods. **j,** Tcf-1 and CD62L expression on Rg T cells in mLN of mixed NP101/105 RgC mice that were recipients of 5,000 Rg T cells bearing each clonotype (i.e. 10,000 cells total). **k,** Crossreactivity profiles of 16 NP-specific TCRs against a panel of naturally occurring IAV variants carrying mutated NP peptides. Reactivity was assessed by measuring IL-2 production from TCR-transduced BW cell lines following stimulation with peptide-loaded APC. Memory category color coding matches that in **d**. Data is the representative of two or three independent experiments.

Fig. 3. TCR sequences and structures of six NP TCR-NP$_{366-374}$/D^b complexes derived from T$_{EM}$, T$_{CM}$ and T$_{BP}$ memory CD8 T cells. a, The genes and CDR3 sequences of six NP TCRs. In the first four NP TCRs, the three residues that vary among their CDR3s are highlighted in green or cyan. Between NP105 and NP204 key CDR3β residues are in common as a result of a shared TRBJ1-6. Only varied residues between two CDR3β are highlighted in yellow. **b,** Ribbon diagrams of six NP TCR-NP$_{366-374}$ /D^b complexes arranged in the order from T$_{EM}$ (α-dominant) to T$_{CM}$ (β-dominant) from left to right. Their α1α2 domains of MHC are positioned in the same orientation. The NP$_{366-374}$ peptide is displayed in sticks with the central residue p6M highlighted in red. TCR Vα, TCR Vβ, NP$_{366-374}$ peptide, MHC α chain and MHC β$_{2m}$ are colored in green, blue, yellow, grey and light orange, respectively. The Cα domain of NP 204 TCR is absent due to structural disorder. **c,** TCR CDR loops and their footprints on pMHC. Magenta and pink surface areas

represent the MHC and peptide residues that interact with TCR Vα, respectively. Orange and
bright orange surface represent the MHC and peptide residues that interact with TCR Vβ,
respectively. **d,** A zoomed-in view of the interaction between TCR and peptide in the NP202-
$NP_{366-374}$ complex. The three varied CDR3 residues noted in **a** are highlighted in the same color
schemes to show their positions and roles. All peptide residues are labeled, with p4E and p6M
highlighted in red for their prominent roles in TCR's recognition.
1542

1543
1544
Fig. 4. Fine structure details of TCR-ligand interactions in a single CD8 T cell repertoire
linked to memory differentiation. **a,** TCR-pMHC interface area of NP101 and NP202 complexes
superimposed based on their D^b $\alpha 1 \alpha 2$ MHC domains. The NP202 complex is colored in grey as a
reference. The TCR α-chain, TCR β-chain, peptide and MHC of NP101 complex are colored in
green, cyan, orange and pink, respectively. A few relevant residues are displayed in sticks for
comparison. **b,** Side-by-side comparison between NP207 and NP203 complexes. NP207 complex

is colored in grey on the left. The TCR α-chain, TCR β-chain, peptide and MHC of NP203
complex are colored in green, cyan, pink and magenta, respectively. Some key residues described
in text are shown in sticks to display hydrogen bonds and salt bridges. The ND1 and NE2 atoms
of H2-D^b H155 are labeled to show their potential pronation/deprotonation status. **c,** Side-by-side
comparison between NP207 and NP203 complexes in a view focusing on the C'(C")-D edge of
Vα domain. The NAG moiety attached to N58α is shown in sticks. In NP203 complex, the
extended loop between C' strand and D strand is stabilized by two hydrogen bonds between
sidechain and mainchain as shown in the insert. **d,** A docking comparison of β-dominated NP105
(in grey) with α-dominated NP202 (colored) on the left and a zoomed-in display of NP105
interaction with NP$_{366-374}$/D^b on the right, in which a part of α2-helix in front of peptide is omitted
for clarity.
1562

1563
1564
Fig. 5. Functional characterization by transcriptome of Rg T cells isolated from mediastinal
lymph nodes of IAV-infected host animals. a, PCA for gene expression of Rg mLN-derived T
cells from B6 mice adoptively transferred with either 10^5 C101 (green) or C105 (blue) retrogenic
CD8[+] T cells expressing NP101 or NP105 TCRs, respectively, 30 d after IAV infection. Each data
point represents results from an individual mouse (n = 14 mice). **b,** Volcano plot of significant
differential gene expression (962 genes) between Cluster 1 Rg cells (all C101, green) and Cluster
3 (all C105, blue). **c,** Volcano plot of a 59 differentially expressed gene subset flagged for known
T cell functionality (T cell panel) where Cluster 1 (C101.n) exhibits an "exhausted" signature and
Cluster 3 (C105.n) exhibits a functionally activated T_{CM} transitioning to T_{EM} signature. **d,**
Hierarchical clustering/heat map of 59 gene T cell panel reclusters samples to 5 exhibiting an
"exhausted" signature (all C101), 5 with activated T_{CM} signature (all C105), and a mixed
C101/C105 cluster of 4 with an intermediate signature. Each row scaled where 1 (white) = base
gene mean, red = fold-change above, and blue = fold-change below, the base mean. **e,** Schematic
of merged C100.n [expressing NP101 and NP105 TCRs] and C200.n [expressing NP202, NP204,

and NP207 TCRs] samples (see Extended Data Fig. 10), shows skewed distribution of clonotypic
Rg T cells by exhaustion signature, activated T_{CM} functionality, and activation state (n = 28 RNA-
seq analyses). The asterisk for C204.3 indicates quiescent signature by activation state (Extended
Data Fig. 10c) but T_{CM} signature by differential T cell panel (Extended Data Fig. 10d).
1583

1584
1585
Fig. 6. Schematic representation of memory T-cell fate and relationships to signaling balance
or asymmetry. a, Differential signaling related to TCR sidedness as a cartoon representation with
CD3 elements and their ITAMs exposed to yield activation as shown. **b,** Graphic depicting key
peptide residues involved in TCR interaction illustrated as magenta balls on the peptide backbone
(simple cylinder) that sits in the antigen presenting platform between the labeled MHC helices in
the top row. The bottom row shows graphics based upon X-ray crystallographic structures of those
residues in magenta ball and stick display with CDR3α and CDR3β (green and cyan) at the top.

Both rows are based on the TCRs NP202, NP207, and NP105 from left to right. Other neighboring
residues are not labeled for clarity. In both rows, the MHC helix closest to the mass center of the
TCR is indicated by the dark color in each complex. **c,** Force on the TCRα and TCRβ subunits are
balanced for T_{BP}, in contrast to the indicated imbalances associated with T_{EM} and T_{CM} represented
as balls of equivalent or differing sizes on a seesaw in gravitation pull to the ground. With respect
to TCR mechanobiology, however, the seesaws would be flipped as the balls represent sizes of
pMHC-related forces impacting the indicated TCR subunits.
1600