RNA polymerase II-TFIIE-TFIIH interface functions in transcription start site selection in
Saccharomyces cerevisiae

Pratik Basnet[1], Yunye Zhu[1†], Irina O. Vvedenskaya[2], Payal Arora[1], Staci Hammer[3††], Brittany McVicar[1], Shawn Aji Alex[4], Bryce E. Nickels[2], and Craig D. Kaplan[1]

[1]Department of Biological Sciences, University of Pittsburgh, Pittsburgh, United States; [2]Department of Genetics and Waksman Institute, Rutgers University, Piscataway, United States; [3]Department of Chemistry, Loras College, Dubuque, Iowa; [4]Department of Biochemistry & Biophysics, Texas A&M University, College Station, Texas.

*For correspondence: craig.kaplan@pitt.edu
[1]Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA 15260, United States

[†]Present address: Mingma Technologies Co., Ltd., Shanghai, China.
[††]Present address: Baylor College of Medicine, Houston, Texas

ABSTRACT

During transcription initiation in *Saccharomyces cerevisiae*, RNA polymerase II (Pol II) and the general transcription factors (GTFs) assemble upstream of transcription start sites (TSSs) to form the pre-initiation complex (PIC). In this model organism, yeast, the PIC selects TSSs through a unidirectional scanning mechanism referred to as promoter scanning. Previous studies have shown that the TFIIH subunit Tfb3 connects TFIIH to the rest of the PIC via interactions with (i) Pol II, (ii) TFIIE, and (iii) other TFIIH components. Activities within the PIC that influence TSS selection can do so by control of initiation efficiency at individual TSSs and by control of TSS scanning (either rate of scanning or scanning processivity). Here, we identified classes of *tfb3* and *tfa1* mutants within the TFIIH-Pol II-TFIIE interface able to alter promoter scanning in either upstream or downstream directions, suggesting that changes to this interface can fine-tune scanning behavior through evolution. Mutants were isolated based on transcriptional phenotypes and were validated for changes to promoter-specific TSS usage at *ADH1*. Subsets of alleles were analyzed using TSS sequencing approaches, showing that tested *tfb3* and *tfa1* alleles shift TSS distributions across most genomic promoters. Genetic interaction and genomic analysis revealed that the Tfb3 interfaces with Rpb7 and Tfa1 both contribute to promoter scanning, and that *tfb3* alleles exhibit additive effects with *ssl2* and *sub1Δ*, consistent with Tfb3-PIC interactions likely modulating scanning processivity.

INTRODUCTION

In eukaryotes, RNA polymerase II (Pol II) transcribes all protein-coding genes and numerous noncoding RNAs in the first step of gene expression. This process proceeds through three consecutive steps: initiation, elongation, and termination (ROEDER 2019). The first step in transcription, initiation, requires promoter opening, insertion of the template DNA strand into the Pol II active site, and transcription start site (TSS) selection (HAHN 2004; CRAMER *et al.* 2008; HABERLE AND STARK 2018). Most promoters have multiple TSSs, and the precise choice of TSS can influence mRNA content and gene expression (CARNINCI *et al.* 2005; ZHANG AND DIETRICH 2005; KAPLAN 2013; NEPAL *et al.* 2013; HABERLE *et al.* 2014). Initiation is a highly regulated process and requires the activities of RNA Pol II along with several general transcription factors (GTFs) (CONAWAY AND CONAWAY 1993; KORNBERG 2007; GRÜNBERG AND HAHN 2013). Pol II and GTFs are recruited and assembled upstream of the TSS on promoter DNA to form a pre-initiation complex (PIC).

Although PIC components and their structures are highly conserved (BURATOWSKI *et al.* 1988; TOMKO *et al.* 2021), TSS selection in most eukaryotes is distinct from that in budding yeast. In higher eukaryotes, transcription initiation for promoters containing a TATA-box core promoter element occurs at ~30-31 bp downstream of the TATA-box (KADONAGA 2012; DANINO *et al.* 2015; DUDNYK *et al.* 2024). In contrast, initiation in *S. cerevisiae* occurs at multiple TSSs ranging from ~40 to 120 bp downstream of a TATA-box when present (STRUHL 1989; ZHANG AND DIETRICH 2005). The mechanism of TSS selection in yeast involves a unidirectional scanning mechanism wherein the PIC surveys downstream positions for suitable TSSs to initiate transcription. This mechanism of initiation in yeast is known as "promoter scanning" (GIARDINA AND LIS 1993; HAMPSEY 2006; KUEHNER AND BROW 2006; FAZAL *et al.* 2015; MURAKAMI *et al.* 2015). Genetic and genomic studies have established promoter scanning as a general feature of yeast transcription initiation(QIU *et al.* 2020; LU AND LIN 2021; YANG *et al.* 2022; ZHU *et al.* 2024).

The promoter scanning model posits that individual TSSs possess intrinsic differences in their likelihood of being used, and genetic and molecular data indicate that changes in the activity of Pol II and GTFs can shift TSS usage upstream or downstream, as reflected by changes to the overall distribution of TSS usage at promoters (PINTO *et al.* 1992; SUN 1996; BANGUR *et al.* 1997; WU *et al.* 1999; FAITAR *et al.* 2001; GHAZY *et al.* 2004; FREIRE-PICOS 2005; KAPLAN *et al.* 2012; BRABERG *et al.* 2013; JIN AND KAPLAN 2015; ZHAO *et al.* 2021). Promoter scanning has been proposed to be regulated through two distinct networks, the first relating to the innate initiation efficiency of individual TSSs and Pol II catalytic activity, and the second relating to TFIIH DNA translocase activity as the proposed engine for promoter scanning (ZHAO *et al.* 2021; ZHU *et al.* 2024; ARORA *et al.* 2025). While TSS

selection is primarily dictated by the activities of TFIIH and Pol II, there are multiple interactions among
PIC components during initiation. How each PIC component and their interactions influence the TSS
selection mechanism, either through communication to different enzymatic activities in the PIC or
through structural means, is not clear at the molecular level.
TFIIH, a multiprotein complex of 11 subunits in yeast, functions to drive promoter opening for Pol II
promoters in eukaryotes (SVEJSTRUP *et al.* 1994; RANISH *et al.* 2004; MURAKAMI *et al.* 2012; GRÜNBERG
AND HAHN 2013). To do this, TFIIH translocates downstream DNA towards the PIC using the ATPase
activity of its Ssl2 subunit (the yeast homolog of human XPB) (FAZAL *et al.* 2015; FISHBURN *et al.* 2015;
FISHBURN *et al.* 2016; TOMKO *et al.* 2021). During promoter scanning initiation, TFIIH has been
proposed to continue translocating on downstream DNA subsequent to promoter opening, allowing
downstream TSSs to be accessed by the PIC (FAZAL *et al.* 2015; TOMKO *et al.* 2021; ZHAO *et al.* 2021;
BASSETT *et al.* 2022). Structural studies have shown that the TFIIH component Tfb3, the yeast homolog
of human MAT1, links TFIIH to the rest of the PIC via interactions with the Pol II subunit Rpb7 and
TFIIE, through Tfa1 (CHEN *et al.* 2021; SCHILBACH *et al.* 2021). Intriguingly, yeast PICs lacking the
TFIIK module (Tfb3, Kin28, Ccl1) or just the Tfb3 RING domain initiate far upstream in vitro near the
position observed for TSS selection in other eukaryotes (MURAKAMI *et al.* 2015; YANG *et al.* 2022). This
work highlighted the potential importance of these interactions in the evolution of promoter scanning.
Genetic and molecular data suggest that Tfb3-Rpb7 interactions are functionally important for
promoter scanning in vivo, as mutations on either side of this interface cause upstream shifts to TSS
usage (BRABERG *et al.* 2013; YANG *et al.* 2022; YANG *et al.* 2024). Given the strong effects of Tfb3 on
initiation in vitro and in vivo, the interface between Pol II and TFIIH has been suggested as one facet for
how promoter scanning has evolved in yeast, with residues differing between yeast and human located
within Tfb3 shown to affect TSS selection in vivo (YANG *et al.* 2024). One residue in particular, Tfb3 P51
appears to work by supporting adjacent Tfb3 residues in electrostatic interaction with Rpb7 (YANG *et al.*
2024). Interestingly, the MAT1 (human Tfb3)-Rpb7 interface may also undergo remodeling during
human promoter opening (AIBARA *et al.* 2021). In reconstituted closed complex PIC structures, MAT1-
Rpb7 interactions are intact, including through conserved residues that can support salt bridges in both
human and yeast PICs. Putatively, upon XPB translocation this interface breaks and subsequently
density for the MAT1 RING domain is not observed, suggesting these interactions may control coupling
of Ssl2/XPB to the rest of the PIC during promoter opening/promoter scanning (AIBARA *et al.* 2021).
Here, we have undertaken genetic and genomic analysis of the Pol II-TFIIH-TFIIE interface
functions in TSS selection. We have identified novel *tfb3* and *tfa1* alleles by genetic screens for
transcription-related phenotypes. For both Tfb3 and Tfa1, we identified two classes of alleles that
produce distinct polarity of shifts in TSS usage in vivo. In addition, genetic interactions of *tfb3* alleles
with either *ssl2* alleles or *sub1Δ* were primarily additive, suggesting that Tfb3 defects can alter TFIIH
processivity. We have probed the contributions of electrostatic interactions between Rpb7 and Tfb3 in
promoter scanning, providing evidence for the contribution of two conserved salt bridges at this
interface for TSS selection. Intramolecular and intermolecular genetic interactions support that contacts
between both Tfb3-Rpb7 and Tfb3-Tfa1 additively contribute to promoter scanning. Importantly,
intramolecular suppression of putative interface disrupting mutations that shift TSSs upstream by
opposite polarity, downstream-shifting mutations are consistent with the latter class enhancing
interactions at the interface. In parallel, we mapped TSSs genome-wide for representative *tfb3* and *tfa1*
alleles, which demonstrated corresponding global upstream or downstream shifts in TSS usage.
Collectively, our findings indicate that the Pol II-TFIIH-TFIIE interface, mediated through Tfb3-Rpb7 and
Tfb3-Tfa1 interactions, plays a critical role in the promoter scanning mechanism and can be tuned by
mutation.
133
RESULTS

Genetic screens for *tfb3* and *tfa1* alleles show transcription-related phenotypes

To dissect how promoter scanning governs TSSs selection during transcription initiation in yeast
and to probe the role of an RNA Pol II-GTF interface, we performed genetic screens for *tfb3* and *tfa1*
alleles with transcription-initiation-related phenotypes (see Materials and methods). We applied a
battery of phenotypes during our genetic screening. Two of these phenotypes are related to
transcription initiation at the *IMD2* promoter (**Figure 1a**) (KAPLAN *et al.* 2012; BRABERG *et al.* 2013).
One initiation phenotype reflects the sensitivity of yeast cells to Inosine-5′-monophosphate
dehydrogenase (IMPDH) inhibitors such as the drug mycophenolic acid (MPA) (SHAW AND REINES
2000; HYLE *et al.* 2003). IMDPH is the rate limiting step in GTP synthesis, and *IMD2* encodes an MPA-
resistant form of IMDPH and is necessary for cell viability in the presence of MPA. When MPA is
present, the reduction in GTP levels leads to usage of a downstream TSS that allows for production of
IMD2 mRNA (SHAW AND REINES 2000; HYLE *et al.* 2003; JENKS *et al.* 2008; KUEHNER AND BROW 2008;
KWAPISZ *et al.* 2008). Mutants that shift TSSs upstream reduce the usage of the downstream TSS
required for MPA resistance and are therefore sensitive to MPA (**Figure 1a**, **top**) (KAPLAN *et al.* 2012;
JIN AND KAPLAN 2015; ZHAO *et al.* 2021). We screened for downstream-shifting mutants using the
converse *IMD2* phenotype: the constitutive usage of the downstream *IMD2* TSS seen for a wide range
of initiation mutants. We detect constitutive *IMD2* expression using the reporter allele *imd2Δ::HIS3*,
where *HIS3* replaces the *IMD2* ORF as it can easily be observed as a His$^+$ phenotype (**Figure 1a**,
bottom) (MALIK *et al.* 2017; ZHAO *et al.* 2021). These phenotypes have been used as genetic tools to
probe initiation defects in previous studies relating to Pol II and other GTF mutants.
We focused on *TFA1*, encoding a component of TFIIE, and *TFB3*, encoding a component of
TFIIH because of the structurally described Rpb7-Tfb3-Tfa1 interface in the yeast PIC (SCHILBACH *et al.*
2017; SCHILBACH *et al.* 2021). Prior functional work implicated the RING finger domain of Tfb3 as
critical for proper TSS selection during scanning. Tfb3 binds in a pocket formed by the Tfa1 E-linker
helices and the OB domain of Rpb7 (SCHILBACH *et al.* 2017; YANG *et al.* 2022). We performed *TFB3*
and *TFA1* genetic screens by a gap-repair strategy using random PCR mutagenesis followed by
plasmid shuffling (**Figure 1b**) (**Materials and methods**). We identified 31 *tfb3* and 26 *tfa1* alleles each
containing a unique single amino acid substitution (**Table 1 and Table 2**). Mutant growth phenotypes
are shown in spot assays. We identified alleles for both *TFA1* and *TFB3* with presumptive TSS-shifting
phenotypes of each polarity, MPA^s presumptive upstream shifting and His$^+$ presumptive downstream
shifting alleles (**Figures 1c, 1d, S1a, S1b**). We failed to identify Spt$^-$ mutants for either *TFB3* or *TFA1*
screen, which is distinct from a previous genetic screen where these were identified as a subset of *ssl2*
His$^+$ alleles in Zhao et al (ZHAO *et al.* 2021). Additionally, a subset of these novel *tfb3 and tfa1* mutants
showed Gal^R phenotypes from the suppression of *gal10Δ56*, a *GAL10* allele that is sensitive to
transcription defects (GREGER 1998; GREGER *et al.* 2000; KAPLAN *et al.* 2005). These Gal^R phenotypes
were only exhibited by a fraction of MPA^s alleles, for both *tfb3* (*C13S*, *C16R*, *S23P* and *I183F*) and *tfa1*
(*K187E, K187N* and *I183F*) (**Figures S1a, S1b**).
172
***tfb3* and *tfa1* MPA^S alleles map to the TFIIE-TFIIH interface**
Mapping substituted residues of *tfb3* and *tfa1* alleles onto the structure of the yeast Pol II PIC
revealed that MPA^s mutants were clustered phenotype-selectively at the Tfb3-Tfa1 interface, the Tfb3-
Rpb7 interface, and in residues important for structural integrity (Zn binding) in the Tfb3 RING domain
(SCHILBACH *et al.* 2017; SCHILBACH *et al.* 2021) (**Figures 1e, 1f**). These results are in line with
collaborative studies from our lab and the Murakami lab showing that Tfb3 interactions with Rpb7 are
critical for promoter scanning and usage of downstream TSSs (YANG *et al.* 2022). Our screen extends
these results and is consistent with disruptions of structural integrity across Tfb3 PIC-interfaces causing
MPA sensitivity. *tfb3* alleles at either the Tfb3-Rpb7 interface (*e.g. tfb3-R64K, tfb3-N66D*) or Tfb3-Tfa1
interface (e.g. *tfb3-K17E*, *tfb3-Y21H*, *tfb3-L22P*, and *tfb3-S23P*) were the strongest MPA^S mutants with
a mix of strong and medium MPAs alleles in Tfa1 on the Tfa1 side of the Tfa1-Tfb3 interface (**Figures
1e, 1f**). Consistent with this interface being important, we identified alleles on both sides of specific
interaction between Tfa1 K99 and Tfb3 Y21 (*tfa1-K99I* and *tfb3-Y21H*) and Tfa1 M176 and Tfb3 K17
(*tfa1-M176T* and *tfb3-K17E*).

His$^+$ alleles are represented by more subtle changes and may reflect altered stability or protein dynamics

In contrast to MPA^S alleles clearly disrupting protein interactions or key structural components, His$^+$ alleles were more distributed with most located at intramolecular or core interactions, though a few alleles mapped to protein interfaces. A general trend for these His$^+$ mutants in both Tfa1 and Tfb3 is that substitutions within the protein core typically involved hydrophobic residues replaced by amino acids of similar nonpolar character, preserving overall hydrophobicity. In contrast, His$^+$ surface substitutions tended to occur at polar residues, where changes in side-chain length or charge may help maintain or remodel intermolecular interactions.

For Tfb3, most His$^+$ alleles were clustered within the RING domain. A recent structural model of the human PIC proposed that early in initiation, remodeling of MAT1/Tfb3 disrupts Pol II-TFIIH coupling by displacing the RING domain, thereby limiting further promoter opening in that system as opposed to the scanning that can occur in yeast (AIBARA *et al.* 2021). The 16 substitutions at 13 residues conferring the His$^+$ phenotype in Tfb3 were mostly located at internal positions within the RING domain rather than at intermolecular interfaces or Zn^{2+}-binding residues, suggesting that these changes may subtly stabilize or alter the dynamics of the RING fold. For example, residues I38 (I38F) and F28 (F28L) reside in the β1 and β2 sheets that contribute to the characteristic αββαβ architecture of the RING finger domain (Gervais et al., 2001). Likewise, I15 (I15N) and M12 (M12K) lie adjacent to the zinc-binding residues C16 and C13 respectively and confer the opposite phenotype to mutations in zinc-binding cysteines. For Tfa1, nine substitutions across eight residues confer the His$^+$ phenotype. Only K99I and H103R are near the Tfa1-Tfb3 interface, while the remaining substitutions are concentrated internally (**Figure 1f**). This distribution suggests that most of these His$^+$ alleles primarily affect intramolecular stability rather than disrupting intermolecular contacts.

For both Tfa1 and Tfb3, we identified residues where distinct substitutions at the same position confer either the His$^+$ or MPA^S phenotypes, arguing for their distinct characters. In Tfb3, the K17 and S23 residues illustrate this dual behavior. The K17E substitution produces a severe MPA^S phenotype, likely by disrupting (i) intramolecular interactions involving nearby polar residues within the α-helix or (ii) intermolecular contacts with the Tfa1 interface. Consistent with this, K17 in Tfb3 is positioned to interact with M176 of Tfa1, and the M176T substitution in Tfa1 also yields an MPA^S phenotype. In contrast, the K17R substitution, which retains the positive charge but alters side chain character, results in a moderate His$^+$ phenotype, suggesting that preservation of electrostatic interactions at this site contribute to the altered phenotype. Similarly, for Tfb3 S23, which is near the Tfa1-K99 and Tfb3 Y21 interaction (described below), we identified *tfb3* S23P which shows strong MPA^S along with Gal^R phenotype, likely by disturbing intermolecular contacts between Tfa1-Tfb3 interface. In contrast, Arora et al. 2025 isolated *tfb3* S23F alleles with the His$^+$ phenotype, supporting idea that this residue is also placed uniquely at Tfb3-Tfa1 interface where depending upon the substation it can either reinforce or disturb the Tfa1-Tfb3 interaction, which is crucial for promoter scanning (ARORA *et al.* 2025).

A parallel pattern was observed for Tfa1 at K99, which forms a hydrogen bond with Y21 of Tfb3 (SCHILBACH *et al.* 2021). The K99R substitution leads to an MPA^S phenotype (**Figure S1b**), likely for reasons similar to those observed for Tfb3 K17E, where perturbation of interfacial interactions compromises functional coupling (discussed below). Consistently, the Y21H substitution in Tfb3 also causes an MPA^S phenotype (**Figure S1a**). Whereas the K99I substitution in Tfa1 produces a strong His$^+$ phenotype (**Figures 1d, S1b**). Given that the K99 residue occupies a unique position at the boundary of alleles associated with MPA^S and His$^+$ phenotypes, implicating K99 in Tfb3-Tfa1 interface interactions and potentially intramolecular stabilization within Tfa1.

Novel *tfb3* and *tfa1* alleles display unique patterns of TSS usage, differing from the previously defined 'efficiency' and 'processivity' mutants

The core promoter of *ADH1* has been used as a model to quantitatively determine TSS usage defects. Previously, we found a perfect concordance between PIC mutants conferring initiation phenotypes at *IMD2* with those that also show quantitative defects at *ADH1* (KAPLAN *et al.* 2012; JIN

AND KAPLAN 2015; MALIK *et al.* 2017; ZHAO *et al.* 2021). Furthermore, mutations showing growth
phenotypes at *IMD2* and TSS phenotypes at *ADH1* also showed global TSS selection effects across
yeast promoters for the subset that have been evaluated (QIU *et al.* 2020; ZHAO *et al.* 2021). We
employ primer extension (PE) to measure the TSS usage at *ADH1*, which contains two major TSSs at -
37 and -27 positions relative to the first base of the translation start codon (+1) along with multiple
minor TSSs (**Figure 2a**) (HULL *et al.* 1995; SUN 1996). The quantification scheme for observed *ADH1*
TSS usage is described in (JIN AND KAPLAN 2015) (**Figure 2b**). Quantifications of PE for *tfb3-L218S* (a
putative upstream shifting mutant) and *tfb3-F28L* (a putative downstream shifting mutant) are shown as
examples. Consistent with previously studied Pol II and other GTF mutants' examined by primer
extension, all tested *tfb3* and *tfa1* MPA^S alleles shifted TSS usage upstream at *ADH1*, whereas tested
His$^+$ alleles shifted TSS usage downstream (**Figures 2c, 2d**) (KAPLAN *et al.* 2012; JIN AND KAPLAN
2015; MALIK *et al.* 2017; QIU *et al.* 2020; ZHAO *et al.* 2021). The effects of strong MPA^S *tfa1* and *tfb3*
alleles appeared intermediate between those previously observed for tested Pol II MPA^S alleles and
tested *ssl2* MPA^S alleles. Pol II alleles have been interpreted as solely acting through increased
catalysis, which allows increased efficiency and therefore increases in absolute as well as relative
usage of upstream TSSs, whereas *ssl2* alleles were interpreted as being defective in scanning
processivity, meaning that primarily downstream TSS usage was truncated, with generally only a
relative increase in usage of upstream TSSs (KAPLAN *et al.* 2012; MALIK *et al.* 2017; QIU *et al.* 2020;
ZHAO *et al.* 2021). Interestingly, both *tfb3* and *tfa1* MPA^S alleles, consistent with their relative
phenotypic strengths, simultaneously increase the usage of upstream TSSs while reducing usage of
downstream TSSs. As expected, and consistent with their phenotypic strengths, both *tfb3* and *tfa1* His$^+$
alleles shifted the TSS distribution downstream by increasing usage of downstream TSSs (see increase
in quantification bins 5 and 6) (**Figures 2c, 2d**).
261
***tfb3* alleles show mostly additive interactions with *ssl2* mutants and *sub1*Δ**
To investigate the relationships between putative *ssl2* processivity mutants and *tfb3* alleles, we
generated *tfb3/ssl2* double mutants that combined both upstream- and downstream-shifting variants of
the two genes. When combining upstream-shifting *tfb3* alleles with upstream-shifting *ssl2* alleles we
observed extensive synthetic sickness or lethality, consistent with each allele conferring defects
independently and the double mutants having stronger phenotypes (**Figures 3a, S2a**). When
combining *tfb3* and *ssl2* alleles of opposing phenotypes and TSS shifts, we observe that there is partial
mutual suppression between the upstream *tfb3-K17E* allele and downstream shifting *ssl2* alleles. The
tfb3-K17E allele shows moderate to strong suppression of the *ssl2* allele His$^+$ phenotype but there is
weaker suppression of the *tfb3-K17E* MPA^S phenotype in the doubles. This result is consistent with
some additivity and that *tfb3-K17E* is either a stronger upstream-shifting allele than the *ssl2* alleles are
downstream, or there is partial epistasis of *tfb3-K17E* to the tested *ssl2* alleles (ssl2 *N230I* and *R636C*)
(**Figure 3b top**). Additionally, in a patch phenotyping assay for double mutants combining *ssl2* His$^+$
alleles (N230I and R636C) with *tfb3* alleles exhibiting varying strengths of MPA^S phenotype, we
observed a spectrum of suppression for transcriptional phenotypes. Specifically, double mutants
carrying the strong *tfb3* MPA^S alleles (C16R, S23P, L22P, and C39R) showed suppression of both the
MPA^S phenotype of the *tfb3* single mutants and the His$^+$ phenotype of the *ssl2* single mutants (**Figure
S2a**).
When combining upstream shifting *ssl2* alleles with downstream shifting *tfb3-F28L* we observed
moderate suppression of the *ssl2* alleles' MPA^S, consistent with additivity between these alleles. This
additive interaction was further supported by patch phenotyping of double mutants combining *ssl2*
MPA^S alleles with *tfb3* His$^+$ alleles exhibiting different strengths of the corresponding phenotypes.
Specifically, double mutants carrying strong or moderate *tfb3* His$^+$ alleles together with *ssl2* MPA^S
alleles displayed weaker MPA^S phenotypes compared to the corresponding *ssl2* single mutants
(**Figures 3b, S2b**). Finally, we also observe stronger His$^+$ phenotypes when His$^+$ alleles of *ssl2* are
combined with *tfb3-F28L*, also consistent with additivity (**Figure 3b bottom**). Similarly, additive
interaction of *ssl2* His$^+$ with *tfb3* His$^+$ was further supported by patch phenotyping where combination of

the *ssl2-R636C* with other *tfb3* His[+] alleles showed stronger His[+] phenotypes than *ssl2-R636C* single
(**Figure S2c**). These double mutant analysis results are in contrast with Pol II allele genetic interactions
with *ssl2* alleles, where epistasis was much more prominent and there was very little synthetic lethality
(ZHAO *et al.* 2021). In other words, we interpret effects of *tfb3* alleles as potentially altering scanning
processivity independently or in parallel with *ssl2* allele effects on processivity, but we can't rule out
additional effects on initiation efficiency or other inputs to scanning (See Discussion).
We also asked about genetic interactions between *sub1Δ* and *tfb3* alleles. Sub1, the yeast
homolog of PC4 (HENRY *et al.* 1996; KNAUS *et al.* 1996), is a PIC-associated protein (TIMOTHY *et al.*
2011) that can shows extensive genetic interactions with PIC components (KNAUS *et al.* 1996; WU *et al.*
1999; CALVO AND MANLEY 2005; JIN AND KAPLAN 2015; GARAVÍS *et al.* 2017; ZHAO *et al.* 2021). Deletion
of *SUB1* (*sub1Δ*) causes a significant downstream shift in TSS distributions in yeast at *ADH1* (KOYAMA
et al. 2008; BRABERG *et al.* 2013). Moreover, its distinct genetic interactions with Pol II or *sua7-1* alleles
compared with *ssl2* alleles suggested that Sub1 might function in controlling scanning processivity, with
loss of Sub1 leading to an increase in processivity (BRABERG *et al.* 2013; JIN AND KAPLAN 2015; ZHAO *et*
al. 2021). When *sub1Δ* is combined with upstream-shifting *tfb3* alleles, *tfb3* alleles' MPA[S] phenotypes
appeared to be mostly unaffected and the *sub1Δ* His[+] phenotype was suppressed. These phenotypes
report on *IMD2* TSS usage and interpretation from these phenotypes alone would suggest *tfb3* alleles
were epistatic to *sub1Δ* (**Figure 3c**)*. However, examination of primer extension for *ADH1* TSS
distributions suggests that in fact *sub1Δ* and *tfb3* upstream-shifting alleles can be additive. When
combining *sub1Δ* with downstream shifting *tfb3* alleles, His[+] plate phenotypes of double mutants were
difficult to differentiate from *sub1Δ* alone, because it's His[+] phenotype is already so strong. However,
examination of *ADH1* TSS distributions in double mutants shows that double mutants have slightly
greater effects than *sub1Δ* alone, consistent with additivity between *sub1Δ* and *tfb3* downstream-
shifting alleles (**Figures 3d, S2d**).
313
***rpb7* and *tfb3* alleles at the Tfb3-Rpb7 interface show strong initiation phenotypes**
To further study the mechanism of Tfb3 function in TSS selection, we focused our attention on
the Tfb3-Rpb7 interface. The Cryo-EM structure of the budding yeast PIC from Schilbach *et al.* 2021
revealed the TFIIE-TFIIH-Pol II interactions within the PIC (**Figures 1e, 4a**) (SCHILBACH *et al.* 2021).
Our genetic screens suggest that Tfb3-Tfa1 interfaces play a critical role in TSS selection during
promoter scanning, likely by contributing to scanning processivity. In addition, structural studies and
limited mutational analysis previously pointed to potential salt bridges between Tfb3 and Rpb7 as being
critical for normal initiation by promoter scanning (BRABERG *et al.* 2013; SCHILBACH *et al.* 2017;
SCHILBACH *et al.* 2021; YANG *et al.* 2022). A salt bridge connects Tfb3 R64 to Rpb7 D166 (which we will
refer to as salt bridge 1 (SB-1), and a second salt bridge connects Tfb3 K65 to Rpb7 E165 (SB-2)
(SCHILBACH *et al.* 2017) (**Figure 4b, left**). However, in a more recent cryo-EM structure only SB-1
between Tfb3 R64 and Rpb7 D166 was observed (**Figure 4b, right**) (SCHILBACH *et al.* 2021). To further
dissect this interface, we performed systematic genetic analyses to determine whether one or both salt
bridges functionally contribute to scanning.
To determine the contributions for putative salt bridge forming residues at the Rpb7-Tfb3
interface to TSS selection, we generated a battery of *rpb7* and *tfb3* single mutants and *rpb7/tfb3* double
mutants (Materials and Method). To enable ease of double mutant generation, we employed a plasmid-
shuffle approach where *tfb3* mutants were generated using site-directed mutagenesis in *TFB3 LEU2*
plasmids and *rpb7* mutants were generated in the genome by CRISPR. *TFB3* or *tfb3* plasmids were
introduced into relevant *RPB7* or *rpb7* strains deleted for *TFB3* by transformation, followed by selection
against a WT *TFB3 URA3* plasmid (**Figure S3a**).
Alleles with substitutions at either one or both salt bridge-forming residues at this interface were
screened for initiation phenotypes using those employed for genetic screening. We found that
mutations in all individual residues within the putative Rpb7-Tfb3 salt bridges conferred MPA sensitivity,
which is predictive of upstream TSS shifts relative to WT. Mutations in SB-1 residues were generally
more MPA sensitive and exhibited stronger overall growth defects than those in SB-2. Similarly, for

both *tfb3* and *rpb7* mutants, charge-altering substitutions caused more severe phenotypes (both in
growth and MPA sensitivity) than alanine substitutions (**Figure 4c**). We interpret this as consistent with
charge incompatibility arising when only one partner of the putative salt bridge is altered, resulting in
the introduction of like charges in proximity. We next sought to test this model more rigorously through
an extensive series of double-mutant analyses.
345
Genetic dissection of mutants at the Tfb3-Rpb7 interface supports functional interactions
between Tfb3-Rpb7 via two pairs of amino residues observed as salt bridges in different cryo-
EM structures
To determine if Tfb3 and Rpb7 residues work together through physical interactions to facilitate
promoter scanning in *S. cerevisiae*, we evaluated the genetic interactions of *tfb3/rpb7* double mutants
and the potential restoration of growth when one or both salt bridges were restored but in novel
configurations through charge reversal mutations. First, to assess the nature of the genetic
interactions—additive, suppressive, or epistatic—at the predicted salt bridge–forming sites, we
performed phenotypic analyses of double mutants harboring alanine-substituted *rpb7* alleles in
combination with alanine-substituted *tfb3* alleles (MANI *et al.* 2008). We observed that combining either
single or double alanine-substituted *rpb7* alleles at putative SB-1 or SB-12 or both with either single or
double alanine-substituted *tfb3* alleles show epistatic interactions (**Figure 4d**). These results are
consistent with these residues requiring each other for function, and mutating one side of the interaction
means that mutation of the other side does not confer a further defect.
Next, to determine whether the Rpb7-Tfb3 interface relies on one or both salt bridges, we
performed charge identity reversal experiments at SB-1, SB-2, or both simultaneously. We
hypothesized that only complete charge reversal at both putative salt bridges would restore the
transcriptional defects caused by mutations that compromise both sites. As a starting point, we
characterized the growth and MPA phenotypes of the individual *rpb7* and *tfb3* alleles used to construct
the combined mutants (**Figure 4e, top**). The *rpb7-D166R/E165K* mutant, in which both salt bridge–
forming residues (D166 and E165) are converted to positively charged residues, is in direct opposition
to the positively charged partners of Tfb3 (R64 and K65). This mutant exhibits a strong MPA^S
phenotype and a pronounced growth defect. We therefore used *rpb7-D166R/E165K* as the reference
"sick" allele in plasmid shuffle experiments designed to evaluate whether restoring charge
complementarity at the Tfb3 positions could suppress its MPA^S and growth phenotype.
The charge reversal at a single putative salt bridge provided only limited suppression of the
initiation defect of the *rpb7-D166R/E165K* mutant. For example, the combinations *rpb7-D166R/E165K*
with *tfb3-R64D* (putatively restoring SB-1) and *rpb7-D166R/E165K* with t*fb3-K65E* (putatively restoring
SB-2) yielded only modest suppression in growth and MPA sensitivity compared to *rpb7-D166R/E165K*
alone. To account for potential charge incompatibility from having two adjacent positively charged
residues, we generated quadruple mutants designed for charge reversal of a single salt bridge but with
an alanine substitution in one of the pair of residues supporting the other salt bridge: *rpb7-*
D166R/E165K + *tfb3-R64D/K65A* (restoring SB-1 but removing charge incompatibility at SB-2) and
rpb7-D166R/E165K + *tfb3-R64A/K65E* (restoring SB-2 but removing charge incompatibility at SB-1).
These mutant combinations displayed improved growth phenotypes relative to their charge-
incompatible counterparts were still not wild type (**Figure 4e, bottom**). Additionally, to remove charge
or side chain effects from each of the salt bridge pairs and examine the relative contributions of SB-1
and SB-2 more carefully, we constructed charge reversal mutations for each salt bridge while alanine
substituting both residues of the other salt bridge. These combinations included *rpb7-D166R/E165A* or
rpb7-D166A/E165K, which were combined with various *tfb3* alleles. Quadruple mutants such as r*pb7-*
D166R/E165A + *tfb3-R64D/K65A* (restoring SB-1 only) and *rpb7-D166A/E165K* + *tfb3-R64A/K65E*
(restoring SB-2 only) showed partial suppression of the double-salt-bridge defect (**Figure S3b**),
consistent with the results obtained from *rpb7-D166R/E165K* + *tfb3-R64D/K65A and rpb7-*
D166R/E165K + *tfb3-R64A/K65E*. Finally, the complete charge reversal at both salt bridges achieved
by combining *rpb7-D166R/E165K* with *tfb3-R64D/K65E*, suppressed the severe growth defect and

suppressed the MPA^S phenotype of the *rpb7-D166R/E165K* mutant (**Figure 4e**). In this case,
restoration of charge complementarity at both interfaces was sufficient to suppress the most growth
defects.
To further confirm the charge-based nature of the Tfb3-Rpb7 interaction, we generated an
alternative mutant, *rpb7-D166K/E165R,* which reverses the charge pattern with different substitutions.
When combined with the corresponding charge-reversal *tfb3* alleles, the combined mutant exhibited
suppression comparable to *rpb7-D166R/E165K*, reinforcing the functional relevance of charge
complementarity at this interface (**Figure S3c**). Collectively, these genetic data indicate that both salt
bridges contribute to a stable and functional Tfb3-Rpb7 interaction, and that disruption of either alone is
sufficient to perturb normal initiation.

rpb7 alleles at Tfb3-Rpb7 salt bridge forming interface show genetic interactions with additional
tfb3 alleles
To further investigate the nature of the Tfb3-Rpb7 interaction, we examined the genetic
interactions of *rpb7* alleles at interface with novel *tfb3* alleles outside the Tfb3-Rpb7 interface. The
strong His$^+$ *tfb3-M12K* shows slight suppression of MPA^S alleles at Rpb7-Tfb3 interface. The *tfb3-M12K*
suppresses the growth defect and MPA^S phenotype of *rpb7-D166A* whereas it slightly suppresses
growth defect of *rpb7-D166A/E165A* (**Figure 5**). This suppression of MPA^S interface mutants by His$^+$
mutants support the scanning defect at Rpb7-Tfb3 is plastic in nature and that His$^+$ mutants may indeed
be acting by increasing interactions. The MPA^S mutants disrupting salt bridge interactions at Tfb3-Rpb7
interface such as *rpb7-D166R/E165K*, *rpb7-D166A/E165A* and *tfb3-R64A/K65A* are strong MPA^S and
had growth defects (**Figure 4**) but they are still viable. This observation suggests that the defect at
Tfb3-Rpb7 interface still supports some scanning in vivo. However, when we combine the *rpb7* alleles
at Tfb3-Rpb7 interface with *tfb3* MPA^S alleles elsewhere, we observed additive effects. When the sick
MPA^S *tfb3-C39R* allele was combined with *rpb7-D166A* or *rpb7- D166A/E165A,* we observed lethality
for both combinations. Additionally, with *tfb3-Y21H*, when combined with *rpb7-D166A*, we observed
strong additive effects in MPA^S phenotype and growth defects. When we combined *tfb3-Y21H* with
rpb7-D166A/E165A, lethality was observed, consistent with an additive interaction (**Figure 5**).

Genome-wide sequencing (TSS-seq and STRIPE-seq) identifies global *tfa1*, *tfb3*, and *rpb7*
initiation effects in *S. cerevisiae*
We examined the 5′ ends of RNA transcripts for select *tfb3*, *tfa1* and *rpb7* alleles on a genome
wide level. For a subset of representative mutants isolated through genetic screens we performed TSS-
Seq (VVEDENSKAYA *et al.* 2015), and a subset of those mutants were additionally validated by STRIPE-
Seq (POLICASTRO *et al.* 2020) (a CAGE-related protocol for TSS mapping) in parallel (**Figures S4a**,
S4b, **Materials and Methods**). We also analyzed salt bridge alleles by STRIPE-Seq (see below). In
total, six *tfb3* alleles, three MPA^S (K17E, C39R, and R64K) and three His$^+$ (K17R, F28L, and I62A)),
four *tfa1* alleles, two MPA^S (K187E, and I183T) and two His$^+$ (F14L, and K99I) and one *rpb7* (MPA^S,
D166G) allele were analyzed along with their strain-specific WT controls by TSS-Seq. The STRIPE-Seq
analyzed strains included two *tfb3* (MPA^S; R64K and His$^+$; F28L) and two *tfa1* (MPA^S; K187E and His$^+$;
K99I) along with their strain-specific WT controls. The positions and counts of the 5′ ends of uniquely
mapped reads were extracted to assess reproducibility between biological replicates. We observed
strong correlations among replicates for both TSS-Seq and STRIPE-Seq datasets (**Figures 6a**, **S4c,**
S5a, S5b). As described in Qiu et al. 2020 and Zhao et al. 2021, we investigated 5979 promoters
across the genome and after stringent data filtering and read depth thresholding (**Materials and**
Methods), we retained 3226 promoters for TSS-Seq and 2883 for STRIPE-Seq (QIU *et al.* 2020; ZHAO
et al. 2021).
To examine how these *tfb3*, *tfa1*, and *rpb7* alleles affect transcription start site (TSS)
distributions, we applied two main metrics for TSS usage as described in Qiu et al. 2020 and Zhao et
al. 2021 (QIU *et al.* 2020; ZHAO *et al.* 2021). First, we quantified TSS distribution differences between
WT and mutants within defined 401 bp promoter windows (**Figure 6b, left**). We observed that all MPA^S
alleles, such as *tfb3-R64K*, *tfa1-K187E*, and *rpb7-D166G*, exhibited upstream shifts in the median

position of the TSS distribution on average across analyzed promoters. In contrast, His$^+$ alleles for both *tfa1* and *tfb3* displayed increased TSS usage at downstream sites, consistent with downstream shifts in initiation (**Figures 6b, 6c**). Next, to quantify changes in TSS usage, we used a metric defined as the "median TSS shift", calculated by subtracting the WT median TSS position within each promoter window from that of the mutant (**Figure 6b, right**). For each mutant, we show the median TSS shifts across promoters (**Figures 6d, 6e**). Consistent with genetic and primer extension observations; *tfb3*, *tfa1* and *rpb7* MPA^S alleles shift the median TSS positions upstream whereas His$^+$ alleles shift median TSS positions downstream at the majority of promoters (**Figures 6d, 6e, S5a**). As both MPA^S and His$^+$ alleles show a gradient of phenotypic strength in genetic tests, the same holds true for median TSS shift. For example, Stronger MPA^S mutants like *tfb3-C39R* and *tfa1-K187E* shows stronger median TSS shift in upstream direction, with the same pattern holding true for His$^+$ alleles (**Figures 1c, 1d, 6e**). These results are consistent with phenotypes of mutants being predictive of global TSS defects. TSS-Seq and STRIPE-Seq techniques both recapitulate the observed effect (**Figures 6d, 6e**).

Genome-wide analysis of TSS selection via STRIPE-seq corroborates interaction between Tfb3-Rpb7 via two salt bridges

The observed growth suppression of the sick *rpb7-D166R/E165K* allele by charge reversal *tfb3-R64D/K65E* suggests potential widespread effects on the *rpb7-D166R/E165K* phenotype. However, the MPA phenotype assessed derives from defects at a single promoter, *IMD2*. To assess potential suppression of defects genome wide, we analyzed a series of mutants by STRIPE-seq (**Figure 7**). Tested alleles included *tfb3-R64K* (SB-1 impaired), *tfb3-R64A/K65A* (both salt bridges impaired from the Tfb3 side), *rpb7-D166A/K165A* (both salt bridges impaired from Rpb7 side), *rpb7-D166R/E165K* (a slow growth allele with both salt bridges impaired along with charge incompatibility) and *rpb7-D166R/E165K+tfb3-R64D/K65E* (quadruple mutant with charge reversal). As expected, these alleles shift the median TSS positions upstream across the majority of promoters, consistent with their MPA^S phenotypic strength (**Figures 7a, 7b**). The alleles where both salt bridge interactions were impaired showed strong shift in median TSS upstream whereas the quadruple mutant with the potential for salt bridge restoration was much closer to wild type with a median shift near zero. This observation validates our salt bridge interaction, where charge reversal allele strongly suppresses the upstream shift in TSSs for the sick *rpb7-D166R/E165K* allele at the majority of promoters in genome wide level.

Qualitative similarity between TSS sequencing techniques but technique-specific differences

Our experiments assessed two types of TSS mapping techniques. TSS-seq as employed by our group, starts with rRNA-depleted RNA and uses a relatively large amount, but is highly reproducible and generates deep promoter coverage (CHO *et al.* 2009; MITSCHKE *et al.* 2011; VVEDENSKAYA *et al.* 2015; QIU *et al.* 2020; ZHAO *et al.* 2021). STRIPE-seq is a CAGE-related protocol that utilizes template switching for addition of a 5^I adapter and second strand synthesis (POLICASTRO *et al.* 2020). An advantage of STRIPE-seq is that it uses a much smaller amount of starting material. STRIPE-seq showed strong agreement with TSS-seq in capturing the expected direction of TSS shifts for tested alleles, although biological replicates showed some variability in median shift (**Figure S5a, left**), and fewer promoters were above threshold for consideration compared to TSS-seq. To directly compare the two methods, we analyzed 2,564 promoters detected by both. Within this shared set, mutants exhibited consistent directional TSS shifts, but the magnitude of shifts differed between techniques (**Figures S6b, S6d**). Clustering analysis of the median TSS positions revealed that data points derived from each technique grouped separately (**Figure S6a**). Similarly, for median TSS shift, although upstream- and downstream-shifting alleles formed distinct clusters, the separation appeared to be driven primarily by the sequencing technique rather than by allele-specific effects (**Figures S6b**). Principal component analysis of the median TSS shifts further confirmed that variation was largely technique-dependent (**Figure S6c**). Together, these results demonstrate that STRIPE-seq reliably captures qualitative TSS-shift behavior of mutants, but differences due to each technique suggest they can be compared qualitatively and not quantitatively.

DISCUSSION

Our genetic and genomic results reveal a notable plasticity in how tested GTFs can contribute to promoter scanning. For both TFIIE subunit Tfa1 and the architectural TFIIH component Tfb3, we were able to obtain mutants that altered TSS distributions in either the upstream or downstream direction. For previously examined Pol II and *SSL2* alleles, the enzymatic activity of each provided a straightforward explanation for finding alleles of different classes – that they increased or decreased activity relative to WT. Here, our studies show that additional GTFs may also increase or decrease the outcomes of scanning as evidenced by shifts in TSS distributions. Consistent with our targeted genetic screens revealing mutants in GTF components able to alter TSS distributions, our other work using unbiased genetic selections across the genome have also revealed mutants across GTFs and in multiple components of TFIIH able to shift TSSs (ARORA *et al.* 2025).

Previous genetic studies identified Rpb7 and Tfb3 alleles at the Tfb3-Rpb7 interface that conferred MPA^S (BRABERG *et al.* 2013; YANG *et al.* 2022; YANG *et al.* 2024). The location and nature of those alleles suggest that defects in protein interactions confer the MPA^S phenotype, and we interpret this phenotype as, minimally, a reduction in TFIIH scanning processivity leading to inability for scanning to reach downstream TSSs. These results would be consistent with this conserved interface being important for the coupling of TFIIH translocation to continued TFIIH scanning in yeast. However, our genetic screens have isolated His$^+$ alleles able to shift TSS usage downstream for both Tfa1 and Tfb3. As noted above, Arora et al. 2025 also identified His$^+$ alleles for the TFIIH components Rad3, Tfb2, Tfb4 and Ssl1 (ARORA *et al.* 2025).

How do these novel alleles alter scanning? Given that we interpret the converse phenotype (MPA^S) for alleles of these genes as compromising the physical integrity of the Tfa1-Tfb3-Rpb7 interface, we interpret His$^+$ alleles as strengthening the interface or adding plasticity to it. This interpretation suggests roles for the architectural components of TFIIH in the PIC that are not enzymatic components themselves but can feedback or modulate the biochemical activities involved in promoter scanning, which are Pol II catalysis and TFIIH translocation. Our genetic interactions between interface components and *ssl2* alleles or *sub1Δ* suggest that *tfb3* alleles at least are additive with *ssl2* and *sub1Δ*, suggesting that some of their effects alter TFIIH processivity. However, upstream shifting *rpb7, tfa1,* and *tfb3* alleles tested at *ADH1* all appear stronger than tested *ssl2* alleles (**Figure 2**), though weaker than strong upstream shifting Pol II alleles (KAPLAN *et al.* 2012; BRABERG *et al.* 2013; QIU *et al.* 2016; ZHAO *et al.* 2021). A simple explanation is that these observations reflect quantitative differences in allele strength. This explanation is somewhat unappealing because an allele that only affects scanning processivity would truncate downstream scanning, therefore leading to relative increases in upstream TSSs. Processivity mutants would not be predicted to alter actual initiation efficiencies at any position within the scanning window, just the length of the scanning window. In contrast, Pol II and TFIIF alleles that shift TSSs upstream appear to strongly activate the use of upstream TSSs and this is consistent with their ability to alter efficiency of each TSS within the scanning window. An intriguing alternative is that Pol II-TFIIH interface mutants might communicate with the Pol II active site (affecting innate initiation efficiency), the TFIIH active site (affecting scanning rate), or an additional PIC activity during scanning. Such an additional activity could reflect the unknown mechanism that restricts initiation close to the core promoter/TATA element. Relief of this restriction would also be predicted to activate TSS usage at upstream positions without needing to directly communicate to the Pol II active site.

Together, these findings suggest that multiple components of the PIC, both enzymatic and structural, participate in shaping the dynamics of promoter scanning. The ability of non-enzymatic factors and interfaces to influence scanning underscores the plasticity of the scanning process and that fine-tuning of these interactions may have contributed to the for evolution of promoter scanning within otherwise highly conserved and essential components of the eukaryotic PIC.

MATERIALS AND METHODS

Yeast strains and media

Yeast strains are derived from a GAL$^+$ of S288C (FY2) (WINSTON *et al.* 1995). Yeast strains used in this study for *the tfb3* mutant screen, *tfb3/ssl2* double mutants, *tfb3/sub1Δ double mutants*, *tfa1* mutant screen, *rpb7* and *tfb3* mutants at the Tfb3-Rpb7 interacting surface, and related experiments are listed in **Supplementary Table 1.** Yeast media are prepared following standard protocols (AMBERG 2005). YP solid medium is made of yeast extract (1% w/v; BD), peptone (2% w/v; BD, 211677) and Bacto agar (2% w/v; BD, 214010), supplemented with adenine (0.15 mM; Sigma-Aldrich, A9126) and L-tryptophan (0.4 mM; Sigma-Aldrich T0254). YPD plates contained dextrose (2% w/v; VWR, VWRBK876), YPRaf plates contained raffinose (2% w/v; Amresco, J392), and YPRaf/Gal plates contained raffinose (2% w/v; Amresco, J392) and galactose (1% w/v; Amresco, 0637) as carbon sources. YPRaf and YPRaf/Gal plates also contain antimycin A (1 µg/ml; Sigma, A8674-100mg). Minimal media plates are synthetic complete ("SC") with amino-acids dropped out as appropriate as described in (AMBERG 2005) with minor alterations as described in (KAPLAN *et al.* 2012): per standard batch formulation, adenine hemisulfate (Sigma-Aldrich, A9126) was 2 g, L-Leucine (Sigma-Aldrich, L8000) was 4 g, Myo-inositol was 0.1 g, para-aminobenzoic acid (PABA) was 0.2 g. SC-Leu+5FOA plates contained 1 mg/ml final concentration of 5-fluoroorotic acid monohydrate (5-FOA, GoldBio, F-230). SC-Leu+MPA plates contained either 20 ug/ml, 5 ug/ml, or 1ug/ml final concentration of mycophenolic acid (MPA, Sigma, M3536-250MG). SC-His+3AT plates contained different final concentrations of 3-Amino-1,2,4-triazole (3-AT, Sigma-Aldrich, A8056).

Plasmids, bacterial strains, and oligonucleotides

Bacterial strains, plasmids, and oligonucleotides used in this study are listed in **Supplementary Tables 2 and 3.**

tfb3 and *tfa1* genetic screening

Novel *tfb3* and *tfa1* mutants were created by PCR-based random mutagenesis coupled with a gap repair. Briefly, to generate randomly mutagenized *tfb3* (*tfb3**), standard PCR was performed using Taq polymerase (NEB). *tfb3** PCR products were then transformed into yeast along with a linearized pRS315-*TFB3* (pCK1664) plasmid with most of the WT TFB3 sequence removed by restriction digest using *Bgl*II and *Nru*I (Thermo Scientific). Leu$^+$ transformants were selected. Homologous sequences on each end of the *tfb3** PCR products and the gapped *tfb3* vector allowed homologous recombination, resulting in a library of gap-repaired plasmids containing potential *tfb3** mutants. Because *TFB3* is an essential gene, these yeast cells are pre-transformed with a pRSII316-*TFB3* (pCK1632) *URA3* plasmid to support growth, while the genomic *TFB3* was deleted to allow plasmid *tfb3* alleles to exhibit phenotypes. After gap repair, cells retaining pCK1632 (pRSII316-*TFB3 URA3*) plasmids were killed when transformants were replica-plated to medium containing 5-FOA (GoldBio). Yeast cells were then replica-plated to various media to screen for mutants with transcription-related or conditional phenotypes. Plasmids from yeast mutants were recovered and transformed into *Escherichia coli* for amplification, followed by sequencing to identify mutations. Additional phenotyping was performed to determine if the mutant phenotype was plasmid-linked or not. All mutants described here were verified by retransformation into a clean genetic background. A similar approach was taken for the *tfa1* genetic screen.

Site-directed mutagenesis at Tfb3-Rpb7 interface: CRISPR Cas9 to generate *rpb7* alleles

Single or double amino acids substituted mutants were created targeting Rpb7 residues; D166 and E165, using the CRISPR-Cas9 single plasmid pML107 (LAUGHERY *et al.* 2015). This Leu auxotrophic marker-based pML107 (gift from Francois Robert Lab) allows the expression of a guide RNA (gRNA) and Cas9 gene together. First, the forward (5′ portion contains a sequence that allows gRNA to bind to Cas9) and reverse strands of gRNA targeting the region containing the PAM site located near the site

of interest for mutation were designed. Similarly, the forward repair oligo containing the mutated PAM
site, and the reverse oligo harboring either single or double substituted amino acids for D166 and E165
were designed. After annealing gRNA oligos, they were cloned into pML107 at *Swa*I and *Bcl*I restriction
enzyme digested sites. Finally, 400ng of pML107 expressing Cas9 and gRNA insertion was
transformed into yeast cells along with double strand repair oligo (up to 1ug) following the yeast high-
efficiency transformation protocol described in (GIETZ AND SCHIESTL 2007). Transformants were grown
on selective SC-Leu plates. Several colonies were picked and initially selected for initiation phenotypes
at SC-Leu+MPA (20ug/ml) plates. Several MPA^S and some MPA^R (control) colonies were picked. PCR
fragments harboring the PAM site with mutation site, and homology sites were sent for sequencing and
verification for desired *rpb7* mutations.
600
Site-directed mutagenesis at Tfb3-Rpb7 interface: PCR sewing (Overlap PCR) to generate *tfb3*
alleles
tfb3 mutants at the Tfb3-Rpb7 interface were created by overlap PCR. Two PCR products overlapping
at the location of a desired mutation introduced using mutant oligos were used in an additional PCR
reaction to generate mutant DNA amplicons with otherwise WT *TFB3* sequence for the insertion into
the full length *TFB3* plasmid. Desired *tfb3** PCR products were ligated into *Bcl*I and *Smi*I (NEB),
digested pRS315 *TFB3 LEU2* plasmid. *tfb3** harboring pRS315 *tfb3* LEU2* plasmids were verified by
sequencing. Finally, 200ng of each pRS315 *tfb3* LEU2* plasmid was transformed into yeast cells
containing pRSII316 *TFB3 URA3* plasmid, while the genomic *TFB3* was removed to allow plasmid
TFB3 alleles to exhibit phenotypes. Transformants were grown on selective SC-Leu plates. Six
candidate colonies from SC-Leu plates were patched on SC-Leu and then replicated to a medium
containing 5-FOA (GoldBio), to kill cells retaining pRSII316 *TFB3 URA3* plasmids. Clones that have lost
URA3 containing plasmid and harbored *tfb3** were then saved to the lab collection.
614
Primer extension assay
Primer extension (PE) assays to detect putative usage of TSSs in *ADH1* promoter, were performed on
30ug total RNA extracted by using a phenol-chloroform method (SCHMITT *et al.* 1990) per reaction as
described in (RANISH AND HAHN 1991) with modifications included in (KAPLAN *et al.* 2012). Primer
CKO401 complementary to *ADH1* mRNA was end-labeled with 32P γ-ATP (PerkinElmer) and T4
polynucleotide kinase (Thermo Scientific). A reverse transcription reaction was performed by mixing M-
MuLV Reverse Transcriptase (NEB, RNase inhibitor (NEB), dNTPs (GE), DTT, and labeled primer with
total RNA. After cDNA formation, RNase A was added to remove excess RNA. Then, the product was
run into a sequencing gel that contained 8% acrylamide gel (19:1 acrylamide: bisacrylamide) (Bio-Rad),
1× TBE, and 7 M urea. Finally, the gels were visualized by Molecular Imager PharosFX™ Plus System
(Bio-Rad) and quantified by Image Lab (5.2).**TSS-seq library preparation**
Yeast cell cultures were grown in liquid YPD media and cells were harvested at the mid-log phase at a
density of 1-1.5 × 10^7 cells/ml. Cell counts per culture were determined by counting cells with a
hemocytometer. We performed library construction for TSS-seq essentially as described by
(VVEDENSKAYA *et al.* 2015); the steps are briefly described as follows; 100 μg of the extracted total RNA
(phenol-chloroform extraction) was treated with 30 U of DNase I (QIAGEN) and purified using RNeasy
Mini Kit (QIAGEN). A riboPOOL probe (siTOOLS Biotech) was used to deplete rRNAs from DNase-
treated RNAs. rRNA depletion was done following published riboPOOL Protocol_v1-6 from siTOOLs
Biotech which was modified by us to deplete 25ug of DNAse-treated RNAs at once. The rRNA-depleted
RNA was resuspended in ~15 μl of nuclease-free water. From rRNA-depleted RNA, to remove RNA
transcripts carrying a 5′ monophosphate moiety (5′-P), ~1 μg of rRNA-depleted RNA were treated with
1 U Terminator 5′-Phosphate-Dependent Exonuclease (Epicentre) in the 1× Buffer A in the presence of
40 U RNaseOUT in a 50 μl reaction at 30°C for 1 hr. Then, RNA was extracted using the phenol-
chloroform method. Next, to get rid of 5′-terminal phosphates, RNAs were treated with 1.5 U CIP (NEB)
in 1× NEBuffer three in the presence of 40 U RNaseOUT in a 50 μl reaction at 37°C for 30 min. After
the reaction was completed, RNAs were again extracted using the phenol-chloroform method. Now to

convert enriched 5ʹ-capped RNA transcripts to 5ʹ-monophosphate RNAs, CIP-treated RNAs were mixed
with 12.5 U CapClip (Cellscript) and 40 U RNaseOUT in 1× CapClip reaction buffer in a 40 µl total
volume and incubated at 37°C for 1 hr. Then, the reaction went through the phenol-chloroform
extraction method. To ligate the 5ʹ adapter, these CapClip-treated RNA products were mixed with 1 µM
5ʹ adapter oligonucleotide s1086, 1× T4 RNA ligase buffer, 40 U RNaseOUT, 1 mM ATP, 10% PEG
8000, and 10 U T4 RNA ligase 1 in a 30 µl reaction volume. The reaction was incubated at 16°C for 16
hr and was halted by adding 30 µl of 2× RNA loading dye. The mixtures were then separated by using
electrophoresis on 10% 7 M urea slab gels in 1× TBE buffer. Then the gel was incubated with SYBR
Gold nucleic acid gel stain. RNA products migrating above the 5ʹ adapter oligo were recovered from the
gel as described (PINTO *et al.* 1994) and were purified by ethanol precipitation. To generate the first
strand cDNA, 5ʹ-adaptor-ligated products were mixed with 0.3 µl of 100 µM s1082 oligonucleotide
containing a randomized 9 nt sequence, incubated at 65°C for 5 min, and cooled to 4°C. A mixture of 4
µl of 5× First-Strand buffer, 1 µl (40 U) RNaseOUT, 1 µl of 10 mM dNTP mix, 1 µl of 100mM DTT, 1 µl
(200 U) of SuperScript III Reverse Transcriptase and 1.7 µl of nuclease-free water was added to the
reaction. The reaction was incubated at 25°C for 5 min, 55°C for 60 min, 70°C for 15 min, and then
cooled to 25°C. 10 U RNase H was added, the mixtures were again incubated for 20 min at 37°C, and
20 µl of 2× DNA loading solution (PippinPrep Reagent Kit, Sage Science) was added. Then, the DNAs
were separated by electrophoresis on 2% agarose gel (PippinPrep Reagent Kit, external Marker B) to
collect a library of ~90 to ~ 550 nt range which is recovered by ethanol precipitation. To amplify cDNA,
9 µl of gel-isolated cDNA was added to the mixture containing 1× Phusion HF reaction buffer, 0.2 mM
dNTPs, 0.25 µM of one of an i5 index containing lab-designed custom primer (CKO3414-3423), 0.25
µM of one of an Illumina TruSeq small RNA indexing primer (RPI1-32) and 0.02 U/µl Phusion HF
polymerase in 30 µl reaction. Then, PCR was performed with an initial denaturation step of 10 s at
98°C, amplification for 12 cycles (denaturation for 5 s at 98°C, annealing for 15 s at 62°C and extension
for 15 s at 72°C), and a final extension of 5 min at 72°C. Finally, PCR-amplified cDNA libraries were
isolated by electrophoresis on 2% agarose gel and products in the range of ~180 to ~ 550 nt were
collected following ethanol precipitation. Thus, generated dual indexed TSS-seq libraries which were
sequenced together with STRIPE-seq libraries. TSS-seq and STRIPE-seq libraries were pooled and
sequenced on an Illumina NovaSeq-S1-200 cycles sequencing flow cell using custom sequencing
primer (s1115) and custom i5-index primer (CKO3424). **Supplement figures S4a and S4b** show the
library design schemes for both TSS-seq and STRIPE-seq.
672
STRIPE-seq library preparation
Yeast cell cultures were grown in liquid YPD media and cells were harvested at the mid-log phase at a
density of 1-1.5 × 107 cells/ml. STRIPE-seq library construction was performed as described by
(POLICASTRO *et al.* 2020; POLICASTRO *et al.* 2021) with modifications on the oligos used, to make it
compatible with the TSS-seq sequencing scheme. The steps are briefly described as follows;
Total RNA was extracted, DNase I digested, and rRNA depleted as described above in the TSS-Seq
method section. For a given library, two reactions apiece with 100ng rRNA-depleted RNA were set up
in parallel. Terminator exonuclease (TEX) reaction, and template switching reverse (TSRT) reaction
were performed as described in (POLICASTRO *et al.* 2020). For all the libraries, the same Reverse
Transcription Oligo (RTO, CKO3425) with random nonamer was used instead of a unique RTO per
library. See **Supplemental Table 3** for RTO and TSO sequences. Bead-purified TSRT products for two
reactions per library were pooled into a single reaction before the final PCR. Final library amplification
was performed on cleaned TSRT products using one of the i5 index containing lab-designed custom
primers (CKO3414-3423), and one of the Illumina TruSeq small RNA index primers (RPI1-32) to
generate dual indexed libraries. Amplified libraries were initially purified with SPRI select beads to
remove small fragments and then gel extracted from 1.8% agarose gel to size select library products
from 200-800 bp using NucleoSpin® Gel and PCR Clean-up (Macherey Nagel, NC0389463) kit. Hence,
the generated STRIPE-seq libraries were sequenced with TSS-seq in the same Nova-Seq flow cell.

TSS-seq data analysis
TSS-Seq data analysis followed the pipeline outlined in (Zнао *et al.* 2021) (Arora *et al.* 2025). Quality
control, read trimming, and mapping were performed as described by (Qiu *et al.* 2020; Zнао *et al.*
2021), resulting in a TSS count table. Thus, the generated count table contains TSS-seq reads at each
position within defined promoter windows (median TSS ± 250 nt upstream and 150 nt downstream) for
5,979 selected promoters. The data were then filtered to retain counts from n = X promoters with an
average of ≥20 reads per promoter, which were used for downstream analysis. Key TSS metrics, such
as 'TSS spread' and 'TSS shift (or Median TSS shift), were calculated using scripts from (Zнао *et al.*
2021). Differences in TSS spread and shift between WT and mutant for selected promoter classes were
visualized as heatmaps (Morpheus) and/or boxplots.
701
STRIPE-seq data analysis:
The STRIPE-Seq data analysis pipeline was similar to TSS-Seq, generating metrics like 'TSS spread'
and 'TSS shift (or Median TSS shift).' However, after alignment to the reference genome, the Picard
toolkit removed reads with more than three soft-clipped bases. Downstream analysis from here was the
same as described above in the TSS-seq data analysis section and in (Arora *et al.* 2025).

DATA AVAILABILITY
The sequencing data generated in this study for TSS-Seq and STRIPE-seq have been deposited in the
NCBI BioProject database under accession number **PRJNA1346453**.

ACKNOWLEDGEMENTS
We acknowledge funding from National Institutes of Health (NIH) grants R01GM097260,
R01GM120450, and R35GM144116 awarded to CDK, and for R35GM118059 awarded to BEN. We
acknowledge funding from the National Science Foundation (US) to Texas A&M University for REU
support for SEH (award DBI-0851611). For computing resources, we acknowledge funding to the
University of Pittsburgh Center for Research Computing and Data (RRID:SCR_022735). Specifically,
this award used the HTC cluster supported by NIH grant S10OD028483. We acknowledge Gabe
Zentner and Robert Policastro for their STRIPE-seq protocol and Chhabi Govind for his valuable input
on implementation of STRIPE-seq. We would like to dedicate this paper to the memory of Gabe
Zentner.

REFERENCES

Aibara, S., S. Schilbach and P. Cramer, 2021 Structures of mammalian RNA polymerase II pre-initiation complexes. Nature 594**:** 124-128.

Amberg, D. C., D. J. Burke and J. N. Strathern, 2005 Methods in Yeast Genetics: A Cold Spring Harbor Laboratory Course Manual. Cold Spring Harbor Press.

Arora, P., K. F. Brennan, M. H. Warner and C. D. Kaplan, 2025 Genetic dissection of transcription start site selection by RNA Polymerase II in *Saccharomyces cerevisiae*, pp. Cold Spring Harbor Laboratory.

Bangur, C. S., T. S. Pardee and A. S. Ponticelli, 1997 Mutational Analysis of the D1/E1 Core Helices and the Conserved N-Terminal Region of Yeast Transcription Factor IIB (TFIIB): Identification of an N-terminal Mutant That Stabilizes TATA-Binding Protein-TFIIB-DNA Complexes. Molecular and Cellular Biology 17**:** 6784-6793.

Bassett, J., J. K. Rimel, S. Basu, P. Basnet, J. Luo *et al.*, 2022 Systematic mutagenesis of TFIIH subunit p52/Tfb2 identifies residues required for XPB/Ssl2 subunit function and genetic interactions with TFB6. Journal of Biological Chemistry 298**:** 102433.

Braberg, H., H. Jin, Erica, Yujia, S. Wang *et al.*, 2013 From Structure to Systems: High-Resolution, Quantitative Genetic Analysis of RNA Polymerase II. Cell 154**:** 775-788.

Buratowski, S., S. Hahn, P. A. Sharp and L. Guarente, 1988 Function of a yeast TATA element-binding protein in a mammalian transcription system. Nature 334**:** 37-42.

Calvo, O., and J. L. Manley, 2005 The transcriptional coactivator PC4/Sub1 has multiple functions in RNA polymerase II transcription. The EMBO Journal 24**:** 1009-1020.

Carninci, P., T. Kasukawa, S. Katayama, J. Gough, M. C. Frith *et al.*, 2005 The Transcriptional Landscape of the Mammalian Genome. Science 309**:** 1559-1563.

Chen, X., Y. Qi, Z. Wu, X. Wang, J. Li *et al.*, 2021 Structural insights into preinitiation complex assembly on core promoters. Science 372**:** eaba8490.

Cho, B.-K., K. Zengler, Y. Qiu, Y. S. Park, E. M. Knight *et al.*, 2009 The transcription unit architecture of the Escherichia coli genome. Nature Biotechnology 27**:** 1043-1049.

Conaway, R. C., and J. W. Conaway, 1993 GENERAL INITIATION FACTORS FOR RNA POLYMERASE II. Annual Review of Biochemistry 62**:** 161-190.

Cramer, P., K. J. Armache, S. Baumli, S. Benkert, F. Brueckner *et al.*, 2008 Structure of Eukaryotic RNA Polymerases. Annual Review of Biophysics 37**:** 337-352.

Danino, Y. M., D. Even, D. Ideses and T. Juven-Gershon, 2015 The core promoter: At the heart of gene expression. Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms 1849**:** 1116-1131.

Dudnyk, K., D. Cai, C. Shi, J. Xu and J. Zhou, 2024 Sequence basis of transcription initiation in the human genome. Science 384.

Faitar, S. L., S. A. Brodie and A. S. Ponticelli, 2001 Promoter-Specific Shifts in Transcription Initiation Conferred by Yeast TFIIB Mutations Are Determined by the Sequence in the Immediate Vicinity of the Start Sites. Molecular and Cellular Biology 21**:** 4427-4440.

Fazal, F. M., C. A. Meng, K. Murakami, R. D. Kornberg and S. M. Block, 2015 Real-time observation of the initiation of RNA polymerase II transcription. Nature 525**:** 274-277.

Fishburn, J., E. Galburt and S. Hahn, 2016 Transcription Start Site Scanning and the Requirement for ATP during Transcription Initiation by RNA Polymerase II. Journal of Biological Chemistry 291**:** 13040-13047.

Fishburn, J., E. Tomko, E. Galburt and S. Hahn, 2015 Double-stranded DNA translocase activity of transcription factor TFIIH and the mechanism of RNA polymerase II open complex formation. Proceedings of the National Academy of Sciences 112**:** 3961-3966.

Freire-Picos, M. A., 2005 Evidence that the Tfg1/Tfg2 dimer interface of TFIIF lies near the active center of the RNA polymerase II initiation complex. Nucleic Acids Research 33**:** 5045-5052.

Garavís, M., N. González-Polo, P. Allepuz-Fuster, J. A. Louro, C. Fernández-Tornero *et al.*, 2017 Sub1
contacts the RNA polymerase II stalk to modulate mRNA synthesis. Nucleic Acids Research 45**:**
2458-2471.
Ghazy, M. A., S. A. Brodie, M. L. Ammerman, L. M. Ziegler and A. S. Ponticelli, 2004 Amino Acid
Substitutions in Yeast TFIIF Confer Upstream Shifts in Transcription Initiation and Altered
Interaction with RNA Polymerase II. Molecular and Cellular Biology 24**:** 10975-10985.
Giardina, C., and J. T. Lis, 1993 DNA Melting on Yeast RNA Polymerase II Promoters. Science 261**:**
759-762.
Gietz, R. D., and R. H. Schiestl, 2007 High-efficiency yeast transformation using the LiAc/SS carrier
DNA/PEG method. Nature Protocols 2**:** 31-34.
Greger, I. H., 1998 Poly(A) signals control both transcriptional termination and initiation between the
tandem GAL10 and GAL7 genes of Saccharomyces cerevisiae. The EMBO Journal 17**:** 4771-
4779.
Greger, I. H., A. Aranda and N. Proudfoot, 2000 Balancing transcriptional interference and initiation on
the *GAL7* promoter of *Saccharomyces cerevisiae*. Proceedings of the National Academy of
Sciences 97**:** 8415-8420.
Grünberg, S., and S. Hahn, 2013 Structural insights into transcription initiation by RNA polymerase II.
Trends in Biochemical Sciences 38**:** 603-611.
Haberle, V., N. Li, Y. Hadzhiev, C. Plessy, C. Previti *et al.*, 2014 Two independent transcription initiation
codes overlap on vertebrate core promoters. Nature 507**:** 381-385.
Haberle, V., and A. Stark, 2018 Eukaryotic core promoters and the functional basis of transcription
initiation. Nature Reviews Molecular Cell Biology 19**:** 621-637.
Hahn, S., 2004 Structure and mechanism of the RNA polymerase II transcription machinery. Nature
Structural & Molecular Biology 11**:** 394-403.
Hampsey, M., 2006 The Pol II initiation complex: finding a place to start. Nature Structural &
Molecular Biology 13**:** 564-566.
Henry, N. L., D. A. Bushnell and R. D. Kornberg, 1996 A Yeast Transcriptional Stimulatory Protein
Similar to Human PC4. Journal of Biological Chemistry 271**:** 21842-21847.
Hull, M. W., K. McKune and N. A. Woychik, 1995 RNA polymerase II subunit RPB9 is required for
accurate start site selection. Genes & Development 9**:** 481-490.
Hyle, J. W., R. J. Shaw and D. Reines, 2003 Functional Distinctions between IMP Dehydrogenase
Genes in Providing Mycophenolate Resistance and Guanine Prototrophy to Yeast. Journal of
Biological Chemistry 278**:** 28470-28478.
Jenks, M. H., T. W. O'Rourke and D. Reines, 2008 Properties of an Intergenic Terminator and Start
Site Switch That Regulate *IMD2* Transcription in Yeast. Molecular and Cellular Biology 28**:**
3883-3893.
Jin, H., and C. D. Kaplan, 2015 Relationships of RNA Polymerase II Genetic Interactors to
Transcription Start Site Usage Defects and Growth in *Saccharomyces cerevisiae*. G3
Genes|Genomes|Genetics 5**:** 21-33.
Kadonaga, J. T., 2012 Perspectives on the RNA polymerase II core promoter. WIREs Developmental
Biology 1**:** 40-51.
Kaplan, C. D., 2013 Basic mechanisms of RNA polymerase II activity and alteration of gene expression
in Saccharomyces cerevisiae. Biochimica et Biophysica Acta (BBA) - Gene Regulatory
Mechanisms 1829**:** 39-54.
Kaplan, C. D., M. J. Holland and F. Winston, 2005 Interaction between Transcription Elongation Factors
and mRNA 3′ -End Formation at the Saccharomyces cerevisiae GAL10-GAL7 Locus. Journal
of Biological Chemistry 280**:** 913-922.
Kaplan, C. D., H. Jin, I. L. Zhang and A. Belyanin, 2012 Dissection of Pol II Trigger Loop Function and
Pol II Activity–Dependent Control of Start Site Selection In Vivo. PLoS Genetics 8**:** e1002627.
Knaus, R., R. Pollock and L. Guarente, 1996 Yeast SUB1 is a suppressor of TFIIB mutations and has
homology to the human co-activator PC4. EMBO J. 15**:** 1933-1940.

Kornberg, R. D., 2007 The molecular basis of eukaryotic transcription. Proceedings of the National
824   Academy of Sciences 104**:** 12955-12961.
825   Koyama, H., E. Sumiya, M. Nagata, T. Ito and K. Sekimizu, 2008 Transcriptional repression of the
826   *IMD2* gene mediated by the transcriptional co‐activator Sub1. Genes to Cells 13**:** 1113-1126.
827   Kuehner, J. N., and D. A. Brow, 2006 Quantitative Analysis of in Vivo Initiator Selection by Yeast RNA
828   Polymerase II Supports a Scanning Model. Journal of Biological Chemistry 281**:** 14119-14128.
829   Kuehner, J. N., and D. A. Brow, 2008 Regulation of a Eukaryotic Gene by GTP-Dependent Start Site
830   Selection and Transcription Attenuation. Molecular Cell 31**:** 201-211.
831   Kwapisz, M., M. Wery, D. Després, Y. Ghavi-Helm, J. Soutourina *et al.*, 2008 Mutations of RNA
832   polymerase II activate key genes of the nucleoside triphosphate biosynthetic pathways. The
833   EMBO Journal 27**:** 2411-2421.
834   Laughery, M. F., T. Hunter, A. Brown, J. Hoopes, T. Ostbye *et al.*, 2015 New vectors for simple and
835   streamlined CRISPR-Cas9 genome editing in*Saccharomyces cerevisiae*. Yeast 32**:** 711-720.
836   Lu, Z., and Z. Lin, 2021 The origin and evolution of a distinct mechanism of transcription initiation in
837   yeasts. Genome Research 31**:** 51-63.
838   Malik, I., C. Qiu, T. Snavely and C. D. Kaplan, 2017 Wide-ranging and unexpected consequences of
839   altered Pol II catalytic activityin vivo. Nucleic Acids Research**:** gkx037.
840   Mani, R., R. P. St.Onge, J. L. Hartman, G. Giaever and F. P. Roth, 2008 Defining genetic interaction.
841   Proceedings of the National Academy of Sciences 105**:** 3461-3466.
842   Mitschke, J., J. Georg, I. Scholz, C. M. Sharma, D. Dienst *et al.*, 2011 An experimentally anchored map
843   of transcriptional start sites in the model cyanobacterium *Synechocystis* sp. PCC6803.
844   Proceedings of the National Academy of Sciences 108**:** 2124-2129.
845   Murakami, K., B. J. Gibbons, R. E. Davis, S. Nagai, X. Liu *et al.*, 2012 Tfb6, a previously unidentified
846   subunit of the general transcription factor TFIIH, facilitates dissociation of Ssl2 helicase after
847   transcription initiation. Proceedings of the National Academy of Sciences 109**:** 4816-4821.
848   Murakami, K., P.-J. Mattei, Ralph, H. Jin, Craig *et al.*, 2015 Uncoupling Promoter Opening from Start-
849   Site Scanning. Molecular Cell 59**:** 133-138.
850   Nepal, C., Y. Hadzhiev, C. Previti, V. Haberle, N. Li *et al.*, 2013 Dynamic regulation of the transcription
851   initiation landscape at single nucleotide resolution during vertebrate embryogenesis. Genome
852   Research 23**:** 1938-1950.
853   Pinto, I., D. E. Ware and M. Hampsey, 1992 The yeast SUA7 gene encodes a homolog of human
854   transcription factor TFIIB and is required for normal start site selection in vivo. Cell 68**:** 977-988.
855   Pinto, I., W. H. Wu, J. G. Na and M. Hampsey, 1994 Characterization of sua7 mutations defines a
856   domain of TFIIB involved in transcription start site selection in yeast. Journal of Biological
857   Chemistry 269**:** 30569-30573.
858   Policastro, R. A., D. J. McDonald, V. P. Brendel and G. E. Zentner, 2021 Flexible analysis of TSS
859   mapping data and detection of TSS shifts with TSRexploreR. NAR Genomics and
860   Bioinformatics 3.
861   Policastro, R. A., R. T. Raborn, V. P. Brendel and G. E. Zentner, 2020 Simple and efficient profiling of
862   transcription initiation and transcript levels with STRIPE-seq. Genome Research 30**:** 910-923.
863   Qiu, C., O. C. Erinne, J. M. Dave, P. Cui, H. Jin *et al.*, 2016 High-Resolution Phenotypic Landscape of
864   the RNA Polymerase II Trigger Loop. PLOS Genetics 12**:** e1006321.
865   Qiu, C., H. Jin, I. Vvedenskaya, J. A. Llenas, T. Zhao *et al.*, 2020 Universal promoter scanning by Pol II
866   during transcription initiation in Saccharomyces cerevisiae. Genome Biol 21**:** 132.
867   Ranish, J. A., and S. Hahn, 1991 The yeast general transcription factor TFIIA is composed of two
868   polypeptide subunits. Journal of Biological Chemistry 266**:** 19320-19327.
869   Ranish, J. A., S. Hahn, Y. Lu, E. C. Yi, X.-J. Li *et al.*, 2004 Identification of TFB5, a new component of
870   general transcription and DNA repair factor IIH. Nature Genetics 36**:** 707-713.
871   Roeder, R. G., 2019 50+ years of eukaryotic transcription: an expanding universe of factors and
872   mechanisms. Nature Structural & Molecular Biology 26**:** 783-791.

873 Schilbach, S., S. Aibara, C. Dienemann, F. Grabbe and P. Cramer, 2021 Structure of RNA polymerase
874   II pre-initiation complex at 2.9 Å defines initial DNA opening. Cell 184**:** 4064-4072.e4028.
875 Schilbach, S., M. Hantsche, D. Tegunov, C. Dienemann, C. Wigge *et al.*, 2017 Structures of
876   transcription pre-initiation complex with TFIIH and Mediator. Nature 551**:** 204-209.
877 Schmitt, M. E., T. A. Brown and B. L. Trumpower, 1990 A rapid and simple method for preparation of
878   RNA from*Saccharomyces cerevisiae*. Nucleic Acids Research 18**:** 3091-3092.
879 Shaw, R. J., and D. Reines, 2000 *Saccharomyces cerevisiae* Transcription Elongation Mutants Are
880   Defective in *PUR5* Induction in Response to Nucleotide Depletion. Molecular and Cellular
881   Biology 20**:** 7427-7437.
882 Struhl, K., 1989 MOLECULAR MECHANISMS OF TRANSCRIPTIONAL REGULATION IN YEAST.
883   Annual Review of Biochemistry 58**:** 1051-1077.
884 Sun, Z., 1996 Functional interaction between TFIIB and the Rpb9 (Ssu73) subunit of RNA polymerase
885   II in Saccharomyces cerevisiae. Nucleic Acids Research 24**:** 2560-2566.
886 Svejstrup, J. Q., W. J. Feaver, J. Lapointe and R. D. Kornberg, 1994 RNA polymerase transcription
887   factor IIH holoenzyme from yeast. Journal of Biological Chemistry 269**:** 28044-28048.
888 Timothy, Scott, J. Holik, T. Kim, Oliver *et al.*, 2011 Sub1 and RPA Associate with RNA Polymerase II at
889   Different Stages of Transcription. Molecular Cell 44**:** 397-409.
890 Tomko, E. J., O. Luyties, J. K. Rimel, C.-L. Tsai, J. O. Fuss *et al.*, 2021 The Role of XPB/Ssl2 dsDNA
891   Translocase Processivity in Transcription Start-site Scanning. Journal of Molecular Biology 433**:**
892   166813.
893 Vvedenskaya, I. O., S. R. Goldman and B. E. Nickels, 2015 Preparation of cDNA Libraries for High-
894   Throughput RNA Sequencing Analysis of RNA 5′ Ends, pp. 211-228. Springer New York.
895 Winston, F., C. Dollard and S. L. Ricupero‐Hovasse, 1995 Construction of a set of convenient
896   *saccharomyces cerevisiae* strains that are isogenic to S288C. Yeast 11**:** 53-55.
897 Wu, W.-H., I. Pinto, B.-S. Chen and M. Hampsey, 1999 Mutational Analysis of Yeast TFIIB: A
898   Functional Relationship Between Ssu72 and Sub1/Tsp1 Defined by Allele-Specific Interactions
899   With TFIIB. Genetics 153**:** 643-652.
900 Yang, C., P. Basnet, S. Sharmin, H. Shen, Craig *et al.*, 2024 Transcription start site scanning requires
901   the fungi-specific hydrophobic loop of Tfb3. Nucleic Acids Research 52**:** 11602-11611.
902 Yang, C., R. Fujiwara, H. J. Kim, P. Basnet, Y. Zhu *et al.*, 2022 Structural visualization of de novo
903   transcription initiation by Saccharomyces cerevisiae RNA polymerase II. Mol Cell 82**:** 660-676
904   e669.
905 Zhang, Z., and F. S. Dietrich, 2005 Mapping of transcription start sites in *Saccharomyces cerevisiae*
906   using 5′ SAGE. Nucleic Acids Research 33**:** 2838-2851.
907 Zhao, T., I. O. Vvedenskaya, W. K. Lai, S. Basu, B. F. Pugh *et al.*, 2021 Ssl2/TFIIH function in
908   transcription start site scanning by RNA polymerase II in Saccharomyces cerevisiae. eLife 10.
909 Zhu, Y., I. O. Vvedenskaya, S. H. Sze, B. E. Nickels and C. D. Kaplan, 2024 Quantitative analysis of
910   transcription start site selection reveals control by DNA sequence, RNA polymerase II activity
911   and NTP levels. Nat Struct Mol Biol 31**:** 190-202.

914 **Table 1:** *tfb3* alleles with single amino acid substitutions (Ts = temperature sensitive and, Cs = cold
915 sensitive phenotypes).

#	Substitution position	Amino acid change	Phenotypes
1	7	E7K	His$^+$
2	12	M12K	His$^+$
3	13	C13S	MPA^S/Gal^R/Ts
4	15	I15N	His$^+$
5	16	C16R	MPA^S/Gal^R/Ts/Cs
6	17	K17E	MPA^S/Ts
7	17	K17R	His$^+$
8	21	Y21H	MPA^S
9	22	L22P	MPA^S/Gal^R/Ts/Cs
10	23	S23P	MPA^S/Gal^R/Ts/Cs
11	28	F28L	His$^+$
12	38	I38F	His$^+$
13	39	C39R	MPA^S/Ts/Cs
14	44	D44G	His$^+$
15	46	I46T	His$^+$
16	56	Y56C	His$^+$
17	56	Y56F	His$^+$
18	62	I62A	His$^+$
19	62	I62T	His$^+$
20	64	R64K	MPA^S
21	66	N66D	MPA^S
22	68	F68L	MPA^S
23	68	F68S	Gal^R
24	68	F68V	MPA^S
25	73	F73S	MPA^S/Gal^R
26	88	V88E	Gal^R
27	112	E112K	His$^+$
28	112	E112V	His$^+$
29	131	E131V	His$^+$
30	204	D204N	His$^+$
31	218	L218S	MPA^S

Table 2: *tfa1* alleles with single amino acid substitutions (Ts = temperature sensitive and, Cs = cold sensitive phenotypes).

#	Substitution position	Amino acid change	Phenotypes
1	12	L12H	MPA^S
2	14	F14L	His$^+$
3	17	V17D	His$^+$/Ts
4	20	F20S	His$^+$
5	29	L29P	MPA^S
6	33	L33P	MPA^S
7	35	H35R	MPA^S
8	57	L57P	His$^+$
9	64	D64G	MPA^S
10	80	K80E	MPA^S
11	99	K99I	His$^+$
12	99	K99R	MPA^S
13	103	H103R	His$^+$
14	106	V106L	MPA^S/Ts
15	124	C124S	MPA^S/Ts
16	176	M176T	MPA^S
17	183	I183T	MPA^S
18	183	I183F	MPA^S
19	183	I183V	MPA^S
20	186	L186S	MPA^S
21	187	K187E	MPA^S/Gal^R
22	187	K187N	MPA^S/Gal^R
23	187	K187Q	MPA^S/Gal^R
24	190	D190G	His$^+$
25	190	D190N	His$^+$
26	250	R250G	His$^+$

FIGURES AND CAPTIONS

931
Figure 1: Genetic screens identify novel mutants with initiation phenotypes at the interface
between Pol II/TFIIE and TFIIH. **a.** Schematic representation of two transcriptional phenotypes used in
this study. Mycophenolic acid (MPA) sensitivity: MPA inhibits the yeast inosine monophosphate

dehydrogenase (IMPDH) enzymes encoded by *IMD3* and *IMD4*, leading to GTP depletion. In response,
wild-type cells induce *IMD2* expression by a TSS shift from an upstream "G" TSSs, which produces
nonfunctional cryptic transcripts, to a downstream "A" TSS that yields a functional *IMD2* mRNA.
Mutants that are unable to shift the usage to the downstream TSS show MPA^S (top)**.** Suppression of
imd2Δ::HIS3: The *IMD2* ORF was replaced with the *HIS3* to generate the reporter construct. In the
absence of MPA, wild-type cells initiate transcription from upstream TSSs of the *IMD2* promoter,
producing nonfunctional transcripts and failing to grow on medium lacking histidine (SC-His). Mutants
that shift transcription constitutively to the downstream TSS express *HIS3* and show a His$^+$ phenotype
(bottom). **b.** Schematic of genetic screening by PCR-based random mutagenesis coupled with gap
repair/plasmid shuffle to examine *tfb3* or *tfa1* alleles. **c.** Initiation phenotypes of select representative
novel *tfb3* mutants obtained from genetic screening. **d.** Initiation phenotypes of select representative
novel *tfa1* mutants obtained from genetic screening. 10-fold serial dilutions of saturated cultures of WT
and mutant strains were plated on different phenotyping media. **e.** Positions of Tfb3 and Tfa1 relative to
TFIIH and Pol II in the yeast PIC (PDB 7O4J) (SCHILBACH *et al.* 2021). **f.** Mapping novel *tfb3 and tfa1*
mutant residues shown as spheres on the cartoon representation of Tfb3 and Tfa1 protein structures
from (**e**). Residues are colored according to phenotype: shades of purple indicate increasing MPA^S
mutant strength; shades of orange indicate increasing His$^+$ mutant strength; exhibiting with distinct
substitutions conferring MPA^S or His$^+$ phenotypes are shown in red.

953
Figure 2: *tfb3* and *tfa1* alleles show TSS distribution alterations at *ADH1*. a. Schematic of the TSS
region at *ADH1*. The *ADH1* promoter contains two major TSSs at −37 and −27 nucleotides upstream of
the *ADH1* start codon (designated as −37 and −27). **b.** Quantification scheme for *ADH1* TSS usage for
tfb3 MPA^S and His$^+$ mutants. *ADH1* TSS signals were divided into six bins, quantified, and normalized
to the total signal from all bins, with the background removed based on "no RNA" reaction signals. The

change in the TSS usage fraction for mutants relative to WT was determined by subtracting the TSS
usage of each bin in WT from the corresponding bin in mutants. The upstream and downstream shifters
are colored purple and orange, respectively. **c**. Primer extension and quantification showing analyzed
tfb3 MPA^S (Left) and His$^+$ (Right) mutant effects on *ADH1* TSS distribution. **d.** Primer extension and
quantification showing analyzed *tfa1* MPA^S and His$^+$ mutant effects on *ADH1* TSS distribution. Color
coding of the *tfb3* and *tfa1* allele class in this graph is used throughout the figures. Shades of purple
represent upstream shifting *tfa1* or *tfb3* alleles when used to annotate figures. Shades of orange
represent downstream shifting *tfa1* or *tfb3* alleles. Averages of ≥3 biological replicates ± standard
deviation are shown.

969 **Figure 3: Genetic interactions of *tfb3* alleles with *ssl2* alleles or the PIC cofactor *sub1*Δ show**
970 **additive interactions**. **a.** Patch assay showing viability of *tfb3/ssl2* double mutants assessed by
971 plasmid shuffling. The left panel shows growth on SC-Leu plate which is the control state where WT
972 *TFB3* is present. Middle and right panels show growth on SC-Leu+5FOA plate on day 1 and day 5.
973 Upward and downward arrows denote previously detected or predicted single-mutant effects on TSS
974 distribution (up arrow indicating upstream shifts, down arrow indicating downstream shifts), with greater
975 color intensity indicating stronger phenotypic effects. Patches in black boxes are double mutants that

combine *tfb3* MPA^S mutants and *ssl2* MPA^S mutants. **b.** Spot assay results for double mutants
combining *tfb3 K17E* (MPA^S allele) and *ssl2* His$^+$ alleles (Top), *tfb3 F28L* (His$^+$ allele) and *ssl2* MPA^S
alleles (middle) and *tfb3 F28L* (His$^+$ allele) and *ssl2* His$^+$ alleles (bottom). **c.** Spot assay results for
double mutants combining *tfb3* alleles (both MPA^S and His$^+$) and *sub1Δ.* **d.** Quantification of primer
extension result showing TSS usage changes of single and double mutants from WT. Signals for *tfb3*
MPA^S, *tfb3* His$^+$, *sub1Δ*, and *tfb3/sub1Δ* mutants are shown in purple, orange, brown, and gray,
respectively.

a

b

Salt bridges interaction, 5OQJ   Salt bridge interaction, 7O4J

c

d

e

 Figure 4: Genetic dissection of Rpb7-Tfb3-Tfa1 interface. a. Position of Rpb7-Tfb3-Tfa1 interface to
TFIIH and Pol II in the yeast PIC (PDB 7O4J)(SCHILBACH *et al.* 2021). **b**. Interaction between Rpb7 and
Tfb3 via two putative salt bridges (left, PDB 5OQJ) (SCHILBACH *et al.* 2017) vs single putative salt
bridge (right, PDB 7O4J) (SCHILBACH *et al.* 2021) in distinct cryo-EM structures of yeast PIC. Rpb7 and
Tfb3 are colored yellow and blue, respectively. Residue numbers are color-coded according to their
parent protein color. **c.** Mutants at Rpb7-Tfb3 interacting surfaces show strong MPA^S phenotypes. Spot
assay results for *tfb3* (left) and *rpb7* (right) alleles on YPD, SC-Leu and SC-Leu+20 µg/ml MPA. WT is
labeled in black, mutants present at putative SB-1 are labeled in brown, mutants at putative SB-2 are
labeled in orange, with previously described mutants underlined. **d.** Double mutant combination of
alanine-substituted *rpb7* and *tfb3* alleles at either putative salt bridges or both, are epistatic. **e**. Charge
reversal at both putative salt bridges at the Tfb3-Rpb7 interface suppresses individual mutant initiation
defects. Lower MPA concentrations (1 and 5 µg/ml) were used in panels **e–f** to reveal subtle
differences in phenotype strength that are masked at 20 µg/ml MPA.

997
998 **Figure 5:** ***rpb7* mutants in a salt bridge-forming residue of Rpb7-Tfb3 show additive interactions**
999 **with upstream and downstream shifting *tfb3* alleles outside the Rpb7-Tfb3 interface.** Spot assay
1000 results for double mutant combination of *rpb7-D166A* or *rpb7-D166A/E165A* with upstream shifting *tfb3*
1001 alleles (purple) or a downstream shifting *tfb3* allele (orange).

Figure 6: The *tfb3*, *tfa1*, and *rpb7* allele shift transcription start site (TSS) distributions genome wide. a. Scatter plot showing the correlation of $\log_2$ transformed reads at individual genome positions for all positions ≥ 3 reads for biological replicates shown for representative alleles. **b.** Schematic depicting the TSS distribution within a promoter window centered on the median of the wild-type (WT) TSS distribution (Left). Schematic showing shift of the median TSS, either upstream or downstream, in a mutant relative to the WT (Right). **c.** Heatmaps of TSS-normalized read differences between WT and two *tfb3* (*tfb3-K17E*; upstream shifting and *tfb3-I62A*; downstream shifting) mutants within defined promoter windows. Promoter classes were rank ordered according to total reads in WT. The promoter-normalized read differences between the mutant and WT are represented using a color gradient from cyan (negative) to orange (positive). **d.** Heatmap with hierarchical clustering of median TSS shifts in *tfa1*, *tfb3*, or *rpb7* mutants generated by TSS-Seq (Left) and a subset of mutants examined by STRIPE-Seq (Right). **e.** Box plots showing median TSS shifts across promoters in *tfa1*, *tfb3*, or *rpb7* mutants, TSS-Seq (Left) and a subset of mutants with STRIPE-Seq (Right).

1016
1017 **Figure 7: STRIPE-seq validates salt bridge interaction as charge reversal partially restores**
1018 **upstream shift of TSSs in sick *rpb7* mutant. a.** Heatmap with hierarchical clustering of median TSS
1019 shifts in *tfb3*, or *rpb7* salt bridge mutants. **b.** Box plot showing median TSS shifts across promoters in
1020 *tfb3*, or *rpb7* salt bridge mutants. Charge reversal mutant (grey) suppresses the median TSS shift.
1021

Figure S1: Initiation phenotypes of all *tfb3* and *tfa1* mutants obtained from genetic screening. a. Initiation phenotypes of all *tfb3* mutants obtained from screening. **b**. Initiation phenotypes of all *tfa1*

1025 mutants obtained from screening. Spot assays show the 10-fold serial dilutions of saturated cultures of
1026 wild type and mutants plated on different phenotyping media.

1027
1028 **Figure S2: Genetic interactions of *tfb3* alleges with *ssl2* and *sub1* show additive interactions**. **a.**
1029 The patch phenotyping assay of *tfb3* MPA^S/*ssl2* His$^+$ double mutants. **b.** Patch phenotyping assay of

1030   *tfb3* His$^+$/*ssl2* MPA^S double mutants. **c.** Patch phenotyping assay of *tfb3* His$^+$/*ssl2* His$^+$ double mutants.

1031   **d.** TSSs at *ADH1* detected by primer extension for *tfb3, sub1Δ, and tfb3/sub1Δ* mutants.

a

b

c

1032

1033 **Figure S3: Additional double mutant analyses provide evidence for the functional coupling**
1034 **between both salt bridge-forming residue pairs. a**. Schematic representation of CRISPR editing and
1035 PCR sewing followed by plasmid shuffle to generate mutants at salt bridge-forming residues. **b.** The
1036 charge reversal at individual salt bridges partially suppresses double SB mutants. **c.** An alternative *rpb7*
1037 mutant i.e. *rpb7-D166K/E165R* + *tfb3-R64D/K65E* also shows suppression through charge reversal.
1038

Figure S4: The transcription start site sequencing (TSS-seq) and STRIPE-Seq identify global
initiation effects. a. TSS-seq library construction is based on the protocol described by Vvedenskaya
et al., 2015 and Qiu et al. 2020, with dual-index oligos replacing the single-indexed approach during the
final library amplification step (VVEDENSKAYA *et al.* 2015; QIU *et al.* 2020), see Materials and Methods.
b. STRIPE-seq library construction is based on the protocol described by Policastro et al., 2020
(POLICASTRO *et al.* 2020), implementing a single unique reverse transcription oligo (RTO) strategy
instead of a unique RTO per library, see Materials and Methods. **c.** Scatter plot showing the correlation
of $\log_2$ transformed reads at individual genome positions for all positions ≥ 3 reads for biological
replicates of TSS-Seq and STRIPE-Seq libraries.
1049

1051 **Figure S5: Analysis of transcription start site (TSS) shifts in mutants via TSS-Seq and STRIPE-**
1052 **Seq. a.** Heatmap showing hierarchical clustering of median TSS shifts for *tfb3*, *tfa1*, and *rpb7* mutants
1053 across individual replicates (TSS-Seq; left and STRIPE-Seq; right). Data are presented as in **Figure**
1054 **6d**, but for individual replicate libraries. **b.** The principal component analysis (PCA) for median TSS shift
1055 *tfb3*, *tfa1*, and *rpb7* mutants in individual replicates (TSS-Seq; left and STRIPE-Seq; right).

1056
Figure S6: Mutant behavior seems consistent between TSS vs STRIPE-seq techniques with
1057
technique driven variability. a. Hierarchical clustering of Pearson correlation coefficients for *tfb3*, and
1058
tfa1 median shift position in 2654 promoters common to both techniques. **b**. Heatmap showing
1059

hierarchical clustering of median TSS shifts for *tfb3*, and *tfa1* mutants (promoters as in (a)). **c.** The
principal component analysis (PCA) for median TSS shift in *tfb3*, and *tfa1* mutants shows variation
among mutants is technique driven. **d.** Box plot showing median TSS shifts across promoters in *tfb3*,
and *tfa1* mutants. Charge reversal mutants (grey) suppress the median TSS shift. Techniques differ in
magnitude at promoters but consistently capture median TSS shift at the individual promoter level. Blue
label for libraries that are sequenced by STRIPE-Seq and black for TSS-Seq.
1066
1067
1068
1069