1

MS#25-7271

West Nile virus dissemination among individual wild mosquito vectors in Northern

Colorado, 2023

Running head: WEST NILE INFECTIONS IN WILD MOSQUITOES

Authors: Jackson R. DeCook,[1] Molly E. Ring,[1], Claire Stewart,[1] Hana M. Pavelko,[1] Timothy A.

Burton,[1] Gregory D. Ebel,[1] Joseph R. Fauver[2] and Brian D. Foy[1,3]

[1]*Center for Vector-borne and Infectious Diseases, Department of Microbiology, Immunology,*
and Pathology, Colorado State University*, 1685 Campus Delivery, Fort Collins, CO, 80523-
1685

[2]*Department of Epidemiology, University of Nebraska Medical Center, Omaha, Nebraska, USA.*

*The institution at which the work was done

[3]Corresponding author: Jackson DeCook. (307) 315-3442, decookjackson@gmail.com

ABSTRACT. West Nile virus (WNV) transmission risk is typically estimated from pooled whole-mosquito infection data, which may overestimate the proportion of mosquitoes capable of transmission. To assess natural viral dissemination in field-collected *Culex tarsalis,* we tested infection rates in tissues of 1,793 individual mosquitoes collected from Northern Colorado in August 2023. Abdomens were screened for WNV RNA, and corresponding thorax and head tissues from positive mosquitoes were tested. Fifteen mosquitoes had detectable abdominal infections, but WNV RNA was detected in only 53% (8/15) of both the thorax and head tissues, while another 27% (4/15) had WNV RNA detected in either the thorax or the head alone. Logistic regression suggests an inconsistent relationship between abdominal viral ribonucleic acid (RNA) load and virus dissemination, whereas receiver operating characteristic analysis identifies a threshold of ~59,000 RNA copies in the abdomen predictive of dissemination (AUC 0.80, 95% CI: 0.545, 1). These results suggest whole-body RNA detection may overestimate transmission potential from field-captured mosquitoes, and that incorporating infection data could refine surveillance-based risk indices for WNV.

INTRODUCTION

West Nile virus (WNV) is the most widespread arthropod-borne virus in the United

States and poses a persistent public health threat and economic burden (Barrett 2014; Ronca et al.

2019; Snyder et al. 2021; Padda 2025). Since its introduction to the country in 1999, thousands

of cases of West Nile disease (WND) have been diagnosed in humans annually (Patel et al.

2015) despite evidence of significant underreporting of infections and disease (CDC 2024a).

Some estimates put the true number of human infections in the US since the virus emerged at

roughly 7 million (Reimann et al. 2008; Ronca et al. 2019). Mosquitoes from the genus *Culex* are

the primary vectors for WNV, and *Culex tarsalis* (Coquillett) is one of the most significant

enzootic and epidemic vectors in the western U.S., playing a key role in virus amplification via

the enzootic cycle among birds and transmission to humans (Colpitts et al. 2012; Rhodes et al.

2023). Because there is no human vaccine or specific antiviral treatment for WND, disease

prevention relies heavily on entomological surveillance, vector control, and personal protective

measures (CDC 2024b). These Integrated Vector Management (IVM) strategies include WNV

surveillance through mosquito trapping and WNV testing, vector control operations (i.e.

larvicide, adulticide, and water drainage applications), recommendations for personal protection

measures (i.e. changing of behaviors and repellent use), and public communications of WNV

transmission risk (US EPA 2013; Nasci and Mutebi 2019). This approach, although crucial to

reducing transmission risk, may incur significant costs to taxpayers where local and state

governments pay to implement it. Furthermore, the efficacy of some of these approaches can be

hampered by weather and insecticide resistance in the mosquito populations (Tiffin et al. 2025),

and there may be concern from individuals or groups about the effect of widespread insecticide

use on pollinators, other non-target insect populations, and the environment (Peterson et al. 2006;

Rampold et al. 2020; Duval et al. 2023). Accordingly, IVM-connected WNV surveillance

strategies should be employed in at-risk areas using the most precise WNV detection methods, as

these provide critical insight for mosquito control districts/programs, public health authorities

and elected officials about contemporary and temporal risks of WNV transmission risk in

humans and helps to guide decision making around public communications and control efforts.

Current WNV surveillance programs rely on pooled mosquito testing, whereby whole

female mosquitoes in pools (usually up to 50 per pool) are homogenized in a buffer solution. The

buffer is then tested for either WNV antigen with a rapid test or WNV RNA through reverse-

transcription quantitative PCR (RT-qPCR) to estimate pooled infection rates and the Vector

Index (VI), an estimate of the abundance of infected mosquitoes in an area, which can be used to

guide public health interventions (Lanciotti et al. 2000; CDC 2024c). The RT-qPCR method to

quantify WNV RNA is considered to be the most sensitive and accurate method but is more

resource-intensive (Ho-Pun-Cheung et al. 2009; Tomar et al. 2021). West Nile virus positive

pools are determined from amplification of WNV RNA in pools of mosquitoes captured in

standard surveillance traps – meeting a predetermined threshold cycle threshold (CT) value or,

ideally, a threshold CT value determined by comparing amplification to a standard curve of

known viral RNA quantities (Burkhalter et al. 2014). This method assumes that one or more

mosquitoes from a pool determined to be below this CT threshold is infected with WNV and has

the potential to transmit WNV to humans. Data from tested mosquito pools over an area are then

used to calculate the mosquito infection rate via either the minimum infection rate (MIR), which

assumes only one infected mosquito per positive pool, or via the maximum likelihood estimation

(MLE), which probabilistically (via a binomial distribution of infected individuals in a positive

pool) estimates the mosquito infection rate given the observed number of positive mosquito

pools and their pool sizes (Gu et al. 2003). The mosquito infection rate multiplied with the

species capture rate forms the VI for each captured and tested species (Ward et al. 2023). These

mathematical estimates have been shown to be acceptable measures of true mosquito infection

rates, with VI correlating with human WNV infections in the following weeks. Therefore, they

are used as action thresholds to initiate specific vector control interventions to mitigate human

risk (Condotta et al. 2004; Kilpatrick and Pape 2013). Importantly, the mosquito infection rate

variable used in the VI equation naturally varies among mosquito species due to differences in

species vector competence for WNV.

89   Following ingestion of an infectious bloodmeal, the vector competence of a mosquito

species or strain for WNV depends on the ability of the virus to infect the midgut and foregut

tissues, replicate and spread in those tissues, disseminate into hemocoelic tissues, infect the

salivary gland tissue in the thorax, and finally disseminate into the saliva. The vector competence

is variable depending on the virus titer ingested in the blood meal (Richards et al. 2011), and the

ingested and newly produced progeny virions must pass through infection/escape barriers in the

midgut and salivary gland tissues before the mosquito can become infectious (Fitzmeyer et al.

2023). The time that this process takes is known as the extrinsic incubation period (EIP) and has

been characterized rigorously under laboratory conditions (Girard et al. 2004; Reisen et al. 2006).

This period is temperature-dependent but generally lasts between 8-18 days under summer

conditions in temperate regions (Danforth et al. 2015).

Much less is known about WNV dissemination rates in individual wild-caught

mosquitoes, as testing individual field-derived mosquitoes is often not economical due to low

prevalence within mosquito populations, which typically necessitates pooled testing (Burkhalter

et al. 2014; Balingit et al. 2020; Bigeard et al. 2024). Condotta et al. tested individual and pooled

mosquitoes to evaluate the actual infection rate in WNV positive mosquitoes for *Cx. pipiens*

(Linnaeus) and *Cx. restuans* (Theobald) as well as to evaluate if the MIR and MLE overestimate

or underestimate WNV infection, but the analysis did not account for the possibility of non-

disseminated infections. If a substantial proportion of these mosquitoes harbor only localized

midgut infections, risk assessments based solely on infection prevalence may overestimate actual

transmission potential. Improved understanding of viral dissemination in wild mosquitoes is

therefore essential to refine entomological risk indices and enhance the accuracy of outbreak

predictions.

The summer 2023 in Northern Colorado had unusually high mosquito abundance and

infection prevalence, which allowed for a meaningful sample size of individual WNV-positive

mosquitoes to be found over a short two-week time frame. Using these naturally infected field-

derived mosquitoes, we tested the hypothesis that traditional approaches to assessing WNV risk

based on testing of mosquito pools overestimates actual risk. Specifically, we sought to

determine the extent to which WNV-positive, wild *Cx. tarsalis* mosquitoes lack viral RNA in the

head and thorax, indicating a lower-than-expected potential for transmission. By dissecting

individual mosquitoes and estimating viral loads in specific tissues, we aimed to quantify the

proportion of infections that are non-disseminated and assess the implications for surveillance-

based risk assessment.

MATERIALS AND METHODS

Adult *Cx. tarsalis* from Larimer, Weld, and Morgan counties, Colorado, USA were

trapped using the Centers for Disease Control and Prevention (CDC) miniature traps, using dry

ice but no light as the attractant, during the weeks of August 7[th] and 14[th], 2023. In total, 1791

mosquitoes were captured and stored at –80°C until dissection and testing began in 2024.

Mosquito samples were dissected into abdomen, thorax, and head anatomical regions for

tissue-specific analysis, with the thorax samples retaining the legs and wings. Each mosquito and

dissection forceps were surface sterilized by dipping it into 70% EtOH and then rinsed twice in

PBS before dissection. Each mosquito was dissected separately in depression wells of a glass

depression well plate and the plate was sterilized between dissections to avoid contamination

among samples. For most samples, dissected abdomen and head tissues were stored individually

in microfuge tubes without media at –80°C before testing, while thorax tissues were pooled

(n=11 per pool) for testing. These strategies were implemented to increase cost-effectiveness of

the experiment through a reduction of diluent, probe, primer, and extraction media usage.

Samples 19, 76, 124, and 172 were dissected and stored under a slightly different protocol that

stored and tested the thoraces individually because they contain the salivary glands. The heads

were pooled in groups of 11 similar to the thoraces of the other samples, and the dissection and

storage methods were identical.

Mosquito diluent was made in 1-liter batches using 779 mL of 1X sterile phosphate

buffered saline (PBS), 200 mL heat-inactivated EquaFetal fetal bovine serum (FBS), 10 mL

100X penicillin streptomycin (penicillin - 10,000 IU/mL, streptomycin - 10,000 µg/ml), 10 mL

amphotericin B (250 µg/ml), and 1 mL gentamicin (50 mg/mL), and was stored at 4°C. One

sterile ball bearing and 300 µL of mosquito diluent was added to each abdominal tissue sample.

Individual abdominal tissues were homogenized at 24 Hz frequency for 60 seconds and

centrifuged. Subsequently, 15 µL of supernatant from each abdomen homogenate was pooled in

groups of 11 into a microcentrifuge tube, the RNA was extracted from the homogenate pool and

was finally tested using RT-qPCR to identify the presence of WNV RNA in each homogenate

pool.

Mosquito RNA was extracted using a KingFisher Flex instrument and an Omega Bio-Tek

Mag-Bind Viral DNA/RNA 96 kit. Sample RNA was loaded onto a plate with a master mix

consisting of 60 µL TNA lysis buffer, 1 µL linear acrylamide, 70 µL isopropanol, 5µL MagBind

beads, 5 µL proteinase K, and 50 µL of the mosquito homogenate per well. The samples were

then washed twice with 200 µL of SPR buffer and once with 200 µL of VHB buffer and

unloaded into 50 µL elution of nuclease free water. A primer and probe mix containing WNV

forward primer, 5'-TCA GCG ATC TCT CCA CCA AAG-3', WNV reverse primer, 5'-GGG

TCA GCA CGT TTG TCA TTG-3', and WNV FAM probe, 5'-FAM-TGC CCG ACC ATG

GGA GAA GCTC-BHQ-3' was created and 4 µL per sample was added to a master mix

containing 1 µL EXPRESS One-Step SuperScript™ RT enzyme and 10 µL 2X qPCR SuperMix

Universal. After the addition of 15 µL of this master mix per sample, 5 µL of sample RNA was

added per well. PCR testing included positive, negative, and water controls. Quantification of

viral RNA was then done using Applied Biosystems QuantStudio 3 Real-Time PCR system.

Each cycle consisted of a hold stage and PCR stage. Step 1 in the hold stage increased the

temperature by 1.6°C/s up to 50.0°C where it was held for 15 minutes. For step 2, temperature

was increased by 1.6°C/s up to 95.0°C where it was held for 3 minutes. Then in step 1 of the

PCR stage, temperature was held at 95.0°C for 5 seconds, then in step 2 of the PCR stage it

decreased by 1.6°C/s to 60°C, where it was held for 30 seconds. This was repeated for 40 cycles.

Infection of the midgut is a prerequisite for viral dissemination of the entire mosquito.

Therefore, only individuals with RNA presence in the abdomen were included in this study. If a

pooled abdomen supernatant was positive, the 11 individual abdomen homogenates were

subsequently tested for the presence of WNV. Once an individual mosquito was identified as

having an abdomen positive for WNV, the corresponding thorax pool and individual head and

thorax tissues were tested. CT values were obtained from raw PCR output at a consistent

fluorescence threshold of 34,000. Aliquots of varying concentrations of WNV RNA standards

were cloned from viral RNA isolated from stocks and stored at –80°C until their use in a

standard curve run alongside the samples. RNA concentrations in these aliquots were 100,

10,000, and 1,000,000 copies, and the number of copies in each positive sample was estimated

from a linear regression of the standard curve CT values to account for fluctuations in CT

estimates between runs. Whole-body CT estimates were created by summing the RNA estimates

from the standard curves of the three separate tissues and converting them back into a CT value.

Based on previous studies that showed WNV-infected mosquito cells consistently produced

greater than 10^2 RNA copies for each infectious plaque forming unit (PFU) (Fay et al. 2021;

Hamel et al. 2024), a conservative binary infection threshold of 200 RNA copies was created to

distinguish weakly infected tissue. This cutoff minimizes false positives from non-infectious

viral RNA while maintaining sensitivity for biologically relevant infections which are likely to

contain an infectious unit.

Statistical analysis was performed in R version 4.5.1 using packages ggplot2 (Wickham

2016) for visualizations, patchwork (Pederson 2025) for organizing plots, corrplot (Wei and

Simko 2024) for correlation analysis, and pROC (Robin et al. 2011) for receiver operator

characteristics analysis. Logistic regression was performed using a generalized linear model with

a logit link function, and statistical significance was determined as a p value below 0.05.

RESULTS

To provide historical context for the 2023 dissections, the combined *Cx. pipiens* and *Cx.*

tarsalis average weekly mosquito abundance as well as the vector index (VI) in Northern

Colorado regions from long-term (2012–2024) surveillance data are displayed relative to 2023

data (Fig. 1). These elevated VI and mosquito abundance numbers are consistent with high

human West Nile virus (WNV) cases from the same surveillance period (Fig. 2).

Sampled mosquito RNA values varied widely from nearly undetectable to millions of

copies in each tissue compartment but were generally higher in the abdominal tissues (Table 1).

Corresponding thorax and head CT values suggest variable dissemination efficiency, with

several mosquitoes exhibiting high abdominal RNA but limited or no evidence of dissemination

to secondary tissues. Overall, the abdominal infection rate was approximately 1 per 120

mosquitoes, or 0.84%. Of those infected, 8/15 (53.33%) had viral RNA in both the thorax and

head tissue.

Mosquitoes with disseminated infections exhibited consistently low CT values across all

compartments, reflecting high RNA loads (Fig. 3). In contrast, those with failed or incomplete

dissemination showed sharp CT increases (lower RNA) between abdomen and head/thorax

tissues. These differ from weak infections, which exhibit disseminated infections with uniformly

high and/or undetectable CT values across all tissue compartments.

To assess whether RNA concentrations in one tissue could predict viral presence in

others, we conducted correlation analyses using two approaches: raw CT values and log□□-

transformed RNA values (Fig. 4). Raw CT values (Panel A) showed a strong positive correlation

between thorax and head tissues (r = 0.70), moderate correlation between abdomen and head (r =

0.49), and weak correlation between abdomen and thorax (r = 0.23). RNA values that were

estimated from CT outputs were then log-transformed (Panel B) and yielded similar weak to

strong correlations: r = 0.63 for thorax–head, r = 0.50 for abdomen-head, and r = 0.21 for

abdomen–thorax.

A logistic regression was then conducted to evaluate the relationship between presence or

absence of a disseminated WNV infection in mosquitoes and the log-transformed abdominal

RNA levels (Fig. 5). A binary infection threshold of 200 RNA copies in both the head/thorax

was selected to classify individuals as "infected" or "uninfected" in the target tissue. For every

log increase in abdominal RNA, the odds of WNV successfully disseminating into a mosquito's

head and thorax increased by approximately 2.5 times, but the association did not reach

statistical significance (OR = 2.45, 95% CI: [0.81–7.36], p = 0.11).

Lastly, we evaluated whether RNA concentration in the abdomen could serve as a

predictor for the presence of virus in the head and thorax, which is a key indicator of

dissemination and potential transmission competence. The same threshold of 200 copies of RNA

from the previous analysis was used. The receiver operating characteristic (ROC) curve analysis

revealed good overall discriminative ability, with an area under the curve (AUC) of 0.80 (95%

CI: [0.545–1.000]; Fig. 6). This suggests that abdomen RNA levels can moderately to strongly

differentiate mosquitoes with disseminated infections from those without. The optimal abdomen

RNA threshold for predicting head/thorax infection was estimated from the AUC using the same

RNA copy threshold and measured ~59,000 RNA copies. At this threshold, sensitivity was 1.00,

indicating that all mosquitoes with ≥200 copies in the head and thorax were correctly identified.

Specificity was 0.625, meaning that 62.5% of mosquitoes without complete dissemination were

correctly classified as negative.

237   **DISCUSSION**

238   Our study provides novel insights into WNV infection dynamics in wild, field-collected

239   *Cx. tarsalis*, highlighting that detection of WNV in pools of mosquitoes does not explicitly

240   equate to transmission potential in the wild as many of these positive mosquitoes do not have

completely disseminated infections. Of the 15 mosquitoes that were positive in abdominal tissue, nearly half failed to demonstrate complete viral dissemination into the thorax and head. This indicates that a substantial proportion of naturally infected mosquitoes may not contribute to transmission at the time of capture, despite being classified as "infected" under current surveillance standard methods. The addition of whole-body estimates to our analysis provides a comparison to what likely would have been observed had the whole mosquito been tested by itself, or in a pool of non-infected mosquitoes, mimicking the standard approach used in pool-based arbovirus surveillance.

Importantly, this study used mosquitoes that were collected during the unique conditions of 2023, when Northern Colorado experienced unusually high mosquito abundance, WNV activity, and both mild and severe neuroinvasive human cases. These coincided with elevated mosquito abundance and vector index values during mid-August, when field collections were performed. Reported human cases are most often neuroinvasive, which are the cases most likely to be captured by health systems due to underreporting of more mild symptoms. Relative to the preceding decade, both mosquito abundance and VI in 2023 ranked among the highest observed. These circumstances enabled the collection of enough naturally infected individuals (0.83%) to allow for relatively cost-effective screening and testing of individual mosquitoes for tissue-specific analyses, which is rarely economically feasible under typical field conditions (Ward et al. 2023).

Our correlation analysis suggests that viral burden in head and thorax tissues generally increases with abdominal RNA, but the relationship is not strictly proportional. Mosquitoes with failed or incomplete dissemination showed sharp CT increases between abdomen and head/thorax, suggesting potential midgut escape or salivary gland infection barriers or a

mosquito that was captured early in the extrinsic incubation period for the virus. Overall,

correlations between WNV infection in abdomen and thorax tissues were lower than abdomen

and head tissues. Correlation statistics were relatively similar between CT value and log-RNA

estimates.

The relationship between abdominal viral RNA burden and dissemination was not

statistically significant in a logistic regression, though the directionality and magnitude of the

estimated effect suggest that higher abdominal titers increased the odds of dissemination.

According to our model, for every log increase in abdominal RNA, the odds of WNV

successfully disseminating in a mosquito increased by approximately 2.5 times. The lack of

precision from our estimate is reflected in the wide confidence interval and is consistent with the

small sample size (n = 15) and limited statistical power, which likely constrained our ability to

detect moderate effects. Although not statistically significant, the observed directionality of the

effect is consistent with prior laboratory-based studies that suggest higher viral titer in the

infectious bloodmeal, and therefore a lower CT value, leads to an increased chance of a fully

disseminated infection (Condotta et al. 2004). Larger studies with greater statistical power will

be necessary to more precisely estimate the strength of the association and to determine whether

abdominal viral load can serve as a reliable predictor of dissemination.

The ROC curve analysis further explored the predictive utility of abdominal RNA levels

for dissemination. We identified an abdominal RNA threshold of approximately ~59,000 copies

that discriminated between disseminated and non-disseminated infections with high sensitivity

(1.0) but moderate specificity (0.625), yielding an AUC of 0.80 (95% CI: 0.55–1.0). This finding

suggests that increased abdominal RNA values strengthen predictions of a disseminated infection

in mosquitoes. The relatively low specificity indicates that a subset of mosquitoes with high

abdominal titers may still fail to disseminate or could have recently taken an infectious

bloodmeal and are early in the EIP. These results reinforce the biological complexity of WNV

dissemination and the limitations of using whole-body RNA measurements as proxies for vector

competence.

291   Several methodological considerations of our data warrant further discussion. First, the

rarity of naturally infected mosquitoes necessitated pooling strategies in our initial screening, and

while this improved efficiency, it introduced challenges in interpreting thorax RNA values when

multiple individuals were combined. In two cases, pooled thorax homogenates might have

masked heterogeneity between individuals, diluting strong infections or inflating weak ones.

Second, viral RNA quantification by RT-qPCR is inherently variable at low copy numbers, such

as the binary cutoff of 200 RNA copies that we applied (He et al. 2022). Although the threshold

approach provides a consistent framework, different cutoffs could shift classification,

underscoring the need for standardized dissemination definitions for field caught mosquitoes.

Finally, our sample size, though substantial for field collections, yielded only 15 positives. This

limits our ability to detect subtle trends and necessitates cautious interpretation of the statistical

associations we performed.

303   Nonetheless, our results align with and extend prior laboratory studies of *Culex spp.*

vector competence. Experimental infections in colony mosquitoes often report higher

dissemination rates than observed here, possibly reflecting circumstances such as differences in

mosquito microbiome, viral dose, mosquito age, or environmental stressors between controlled

and field conditions (Girard et al. 2004; Reisen et al. 2006). The lower dissemination rates in

wild mosquitoes may most obviously reflect the timing of their capture before the EIP was

complete. Alternatively, they could also reflect a combination of midgut escape barriers, immune

priming, or ecological pressures not captured in laboratory systems (Fitzmeyer et al. 2023).

These findings suggest that extrapolating laboratory-based competence data to field populations

may overestimate true transmission risk at the time of mosquito capture and testing.

From a surveillance perspective, these findings have important implications. Current

arboviral monitoring programs typically classify any WNV-positive mosquito sample below a

cycle threshold value of ~36-39 as potentially infectious, regardless of dissemination status

(Burkhalter et al. 2014; Herman 2015). Our data suggests this approach may inflate estimates of

vectorial capacity by failing to account for the proportion of mosquitoes with unsuccessfully

disseminated infections restricted to the midgut, or weak disseminated infections that may be

mosquitoes captured early in the EIP or that can only transmit few infectious virions. Similarly, a

previous modeling study that incorporated WNV infection dynamics in mosquitoes into

transmission models found that a potential explanation for the variability of CT values observed

in WNV surveillance pools is due to a substantial proportion of mosquitoes that are exposed to

WNV and do not develop a productive infection (Alahakoon et al. 2025). Incorporating tissue-

specific viral dissemination data into predictive risk indices could improve their accuracy and

reduce overuse in areas where disease control resources may be scarce. While the logistical

challenges of dissecting large numbers of mosquitoes are arduous, statistical adjustment factors

that incorporate CT values as a proxy for predicting the potential dissemination and

infectiousness of a mosquito informed by this study and future studies like it may offer a solution.

Future research should focus on validating dissemination thresholds in larger sample sets

and across temporal and ecological contexts. Our sampling occurred within a limited two-week

period in mid-August, representing only a narrow temporal window of mosquito activity.

Seasonal or age-related changes in mosquito physiology may alter dissemination efficiency, and

longitudinal studies with broader sampling could illustrate these dynamics to better capture

seasonal variation in infection prevalence or dissemination efficiency. An important future step

will be integrating dissection-based dissemination data from laboratory studies with salivary

secretion of infectious virions, which would directly establish the link between tissue-level

infection and actual transmission capability (Marín-López et al. 2023). Finally, exploring viral

genetic strain variation in dissemination success would provide valuable insight into the

evolutionary pressures shaping WNV transmission.

In summary, our findings demonstrate that a substantial fraction of field-collected *Cx.*

tarsalis harbor WNV infections restricted to the abdomen, without evidence of dissemination to

the thorax and head. Abdominal viral RNA levels showed predictive value for dissemination, but

with limited specificity. These results demonstrate the importance of distinguishing between

infection and transmissibility in surveillance-based risk models. Refining entomological risk

indices to include CT values as a proxy for transmission likelihood to account for incomplete

dissemination will improve the accuracy of WNV outbreak predictions and guide more efficient

public health responses.

ACKNOWLEDGEMENTS

The Centers for Disease Control and Prevention, Department of Health and Human Services provided financial support for this project. The award provided 100% of total costs and totaled $5,675,000.00. The contents are those of the authors. They may not reflect the policies of the Department of Health and Human Services or the U.S. government.

REFERENCES CITED

Alahakoon P, Marchinton I, Fauver JR, Hay JA. 2025. Tracking West Nile virus dynamics using viral loads from trapped mosquitoes. bioRxiv. 2025.07.02.662782. https://doi.org/10.1101/2025.07.02.662782

Balingit JC, Carvajal TM, Saito-Obata M, Gamboa M, Nicolasora AD, Sy AK, Oshitani H, Watanabe K. 2020. Surveillance of dengue virus in individual *Aedes aegypti* mosquitoes collected concurrently with suspected human cases in Tarlac City, Philippines. *Parasit Vectors*. 13:594. https://doi.org/10.1186/s13071-020-04470-y

Barrett ADT. 2014. Economic burden of West Nile virus in the United States. Am J Trop Med Hy. 90:389–390. https://doi.org/10.4269/ajtmh.14-0009

Bigeard C, Pezzi L, Klitting R, Ayhan N, L'Ambert G, Gomez N, Piorkowski G, Amaral R, Durand GA, Colmant AMG, Giraud C, Ramiara K, Migné C, Grard G, Touzet T, Zientara S, Charrel R, Gonzalez G, Duvignaud A, Malvy D, Lamballerie X De, Fontaine A. 2024. Molecular Xenomonitoring (MX) allows real-time surveillance of West Nile and Usutu virus in mosquito populations. *PLoS Negl Trop Dis*. 18(12):e0012754. https://doi.org/10.1371/journal.pntd.0012754

Burkhalter KL, Horiuchi K, Biggerstaff BJ, Savage HM, Nasci RS. 2014. Evaluation of a Rapid Analyte Measurement Platform and real-time reverse transcriptase polymerase chain reaction assay West Nile virus detection system in mosquito pools. *J Am Mosq Control Assoc*. 30:21–30. https://doi.org/10.2987/13-6386.1

Centers for Disease Control and Prevention. 2024a. Mosquitoes; [accessed 2025 Aug 19].

https://www.cdc.gov/mosquitoes/php/arbonet/index.html

Centers for Disease Control and Prevention. 2024b. West Nile Virus; [accessed 2025 Aug 19].

https://www.cdc.gov/west-nile-virus/hcp/treatment-prevention/index.html

Centers for Disease Control and Prevention. 2024c. West Nile Virus; [accessed 2025 Aug 20].

https://www.cdc.gov/west-nile-virus/php/surveillance-and-control-guidelines/index.html

Colpitts TM, Conway MJ, Montgomery RR, Fikrig E. 2012. West Nile Virus: Biology,

Transmission, and Human Infection. *Clin Microbiol Rev* 25:635–648.

https://doi.org/10.1128/CMR.00045-12

Condotta SA, Hunter FF, Bidochka MJ. 2004. West Nile virus infection rates in pooled and

individual mosquito samples. *Vector Borne Zoonotic Di*s 4:198–203.

https://doi.org/10.1089/vbz.2004.4.198

Danforth ME, Reisen WK, Barker CM. 2015. Extrinsic incubation rate is not accelerated in

recent California strains of West Nile Virus in *Culex tarsalis* (Diptera: Culicidae). *J Med*

Entomol. 52:1083–1089. https://doi.org/10.1093/jme/tjv082

Duval P, Aschan-Leygonie C, Valiente Moro C. 2023. A review of knowledge, attitudes and

practices regarding mosquitoes and mosquito-borne infectious diseases in nonendemic regions.

Front Public Health 11:1239874. https://doi.org/10.3389/fpubh.2023.1239874

Fay RL, Ngo KA, Kuo L, Willsey GC, Kramer LD, Ciota AT. 2021. Experimental evolution of

West Nile Virus at higher Temperatures facilitates broad adaptation and increased genetic

diversity. *Viruses* 13 [accessed 2025 Nov 25]. https://www.mdpi.com/1999-4915/13/10/1889.

https://doi.org/10.3390/v13101889

Fitzmeyer EA, Gallichotte EN, Weger-Lucarelli J, Kapuscinski ML, Adbo Z, Pyron K, Young

MC, Ebel GD. 2023. Loss of West Nile virus genetic diversity during mosquito infection due

to species-dependent population bottlenecks. *iScience* 26:107711.

https://doi.org/10.1016/j.isci.2023.107711

Girard YA, Klingler KA, Higgs S. 2004. West Nile virus dissemination and tissue tropisms in

orally infected *Culex pipiens quinquefasciatus*. *Vector Borne Zoonotic Dis* 4:109–122.

https://doi.org/10.1089/1530366041210729

Gu W, Lampman R, Novak RJ. 2003. Problems in estimating Mosquito infection rates using

Minimum Infection Rate. J Med Entomol. 40(5):595–596. https://doi.org/10.1603/0022-2585-

40.5.595

Hamel R, Narpon Q, Serrato-Pomar I, Gauliard C, Berthomieu A, Wichit S, Missé D, Sofonea

MT, Pompon J.. 2024. West Nile virus can be transmitted within mosquito populations

through infectious mosquito excreta. *iScience* 27 [accessed 2025 Nov 25].

https://www.cell.com/iscience/abstract/S2589-0042(24)02324-1.

https://doi.org/10.1016/j.isci.2024.111099

He Y, Xie T, Tu Q, Tong Y. 2022. Importance of sample input volume for accurate SARS-CoV-

2 qPCR testing. *Anal Chim Acta* 1199:339585. https://doi.org/10.1016/j.aca.2022.339585

Herman NM. 2015. Real-time RT-PCR assays to detect West Nile virus in mosquito pools.

Public Health Theses. 1124. https://elischolar.library.yale.edu/ysphtdl/1124

Ho-Pun-Cheung A, Bascoul-Mollevi C, Assenat E, Boissiére-Michot F, Bibeau F, Cellier D,

Ychou M, Lopez-Crapez E. 2009. Reverse transcription-quantitative polymerase chain

reaction: description of a RIN-based algorithm for accurate data normalization. *BMC Mol Biol*

10:31. https://doi.org/10.1186/1471-2199-10-31

Kilpatrick AM, Pape WJ. 2013. Predicting human West Nile virus infections with mosquito

surveillance data. *Am J Epidemiol* 178:829–835. https://doi.org/10.1093/aje/kwt046

Lanciotti RS, Kerst AJ, Nasci RS, Godsey MS, Mitchell CJ, Savage MS, Komar N, Panella NS,

Allen BC, Volpe KE, Davis BS, Roehrig JR. 2000. Rapid detection of West Nile virus from

human clinical specimens, field-collected mosquitoes, and avian samples by a TaqMan

reverse transcriptase-PCR assay. *J Clin Microbiol* 38:4066–4071.

https://doi.org/10.1128/JCM.38.11.4066-4071.2000

Marín-López A, Raduwan H, Chen TY, Utrilla-Trigo S, Wolfhard DP, Fikrig E. 2023. Mosquito

salivary proteins and arbovirus infection: from viral enhancers to potential targets for vaccines.

Pathogens 12:371. https://doi.org/10.3390/pathogens12030371

Nasci RS, Mutebi J-P. 2019. Reducing West Nile virus risk through vector management. *J Med*

Entomol 56:1516–1521. https://doi.org/10.1093/jme/tjz083

Padda H. 2025. West Nile Virus and Other Nationally Notifiable Arboviral Diseases, United

States, 2023. *MMWR Morb Mortal Wkly Re*p. 74 [accessed 2025 Aug 19].

https://www.cdc.gov/mmwr/volumes/74/wr/mm7421a1.htm.

https://doi.org/10.15585/mmwr.mm7421a1

Patel H, Sander B, Nelder MP. 2015. Long-term sequelae of West Nile virus-related illness: a

systematic review. *Lancet Infect Dis* 15):951–959. https://doi.org/10.1016/S1473-

3099(15)00134-6

Pederson T. 2025. *patchwork: The Composer of Plots.* R package version 1.3.2.9000

https://patchwork.data-imaginist.com

Peterson RKD, Macedo PA, Davis RS. 2006. A human-health risk assessment for West Nile

virus and insecticides used in mosquito management. *Environ Health Perspect* 114:366–372.

https://doi.org/10.1289/ehp.8667

Rampold S, Telg R, Poulin A, Anderson S, Lindsey AB, McLeod-Morin A, Stokes P. 2020.

Public perceptions of mosquitoes and mosquito Control: AEC694/WC357, 5/2020. *EDIS*

2020 [accessed 2025 Nov 25]. https://journals.flvc.org/edis/article/view/121357.

https://doi.org/10.32473/edis-wc357-2020

Reimann CA, Hayes EB, DiGuiseppi C, Hoffman R, Lehman JA, Lindsey NP, Campbell GL,

Fischer M. 2008. Epidemiology of neuroinvasive arboviral disease in the United States, 1999-

2007. *Am J Trop Med Hy*. 79:974–979

Reisen WK, Fang Y, Martinez VM. 2006. Effects of temperature on the transmission of West

Nile virus by *Culex tarsalis* (Diptera: Culicidae). *J Med Entomol* 43:309–317.

https://doi.org/10.1603/0022-2585(2006)043%255B0309:EOTOTT%255D2.0.CO;2

Rhodes CG, Chaves LF, Bergmann LR, Hamer GL. 2023. Ensemble species distribution

modeling of *Culex tarsalis* (Diptera: Culicidae) in the continental United States. *J Med*

Entomol 60:664–679. https://doi.org/10.1093/jme/tjad027

Richards SL, Anderson SL, Lord CC, Tabachnick WJ. 2011. Impact of West Nile Virus dose and

incubation period on vector competence of *Culex nigripalpus* (Diptera: Culicidae). *Vector-*

Borne Zoonotic Dis 11:1487–1491. https://doi.org/10.1089/vbz.2010.0229

Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez J, Müller M. 2011. pROC: an open-

source package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics,*12,

77.

Ronca SE, Murray KO, Nolan MS. 2019. Cumulative incidence of West Nile Virus infection,

continental United States, 1999–2016. *Emerg Infect Dis* 25:325–327.

https://doi.org/10.3201/eid2502.180765

Snyder RE, Cooksey GS, Kramer V, Jain S, Vugia DJ. 2021. West Nile virus–associated

hospitalizations, California, 2004–2017. *Clin Infect Dis* 73:441–447.

https://doi.org/10.1093/cid/ciaa749

Tiffin HS, Gordon JR, Poh KC. 2025. One Health, many approaches: integrated vector

management strategies support One Health goals. *Front Insect Sci* 5 [accessed 2025 Nov 25].

https://www.frontiersin.org/journals/insect-science/articles/10.3389/finsc.2025.1549348/full.

https://doi.org/10.3389/finsc.2025.1549348

Tomar PS, Kumar S, Patel S, Kumar JS. 2021. Development and evaluation of real-time reverse

transcription recombinase polymerase amplification assay for rapid and sensitive detection of

West Nile Virus in human clinical samples. *Front Cell Infec* Microbiol. 10 [accessed 2025

Nov 25]. https://www.frontiersin.org/journals/cellular-and-infection-

microbiology/articles/10.3389/fcimb.2020.619071/full.

https://doi.org/10.3389/fcimb.2020.619071

US EPA [United States Environmental Protection Agency] 2013. [accessed 2025 Aug 19].

https://www.epa.gov/insect-repellents/deet

Ward MJ, Sorek-Hamer M, Vemuri KK, DeFelice NB. 2023. Statistical tools for West Nile virus

disease analysis. In: Bai F, editor. *West Nile Virus: Methods and Protocols*. Springer US; p

171–191 [accessed 2025 Nov 25]. https://doi.org/10.1007/978-1-0716-2760-0_16.

https://doi.org/10.1007/978-1-0716-2760-0_16

Wei T, Simko V. 2024. R package "corrplot": Visualization of a Correlation Matrix.

https://github.com/taiyun/corrplot

Wickham H. 2016. ggplot2: *Elegant Graphics for Data Analysis*. Springer-Verlag New York.

https://ggplot2.tidyverse.org

Table 1

Table. Infection data from all wild *Culex tarsalis* mosquitoes that tested positive in their abdomens for West Nile virus RNA						
Mosquito ID	**Abdomen CT**	**Abdomen RNA (copies)**	**Thorax CT**	**Thorax RNA (copies)**	**Head CT**	**Head RNA (copies)**
A3	15.86	5,985,545.67	25.24	8,461.57	23.39	30,850.83
A4	32.81	42.27	25.24	8,461.57	33.84	20.52
D8	17.54	1,845,606.78	23.85	22,353.08	24.37	15,554.08
D10	16.28	4,471,412.28	23.85	22,353.08	28.74	732.22
19	22.54	68,236.08	21.76	118,408.02	24.41	18,157.96
76	22.99	49,615.82	Negative	Negative	Negative	Negative
124	23.37	37,916.02	Negative	Negative	Negative	Negative
182	37.00	2.46	Negative	Negative	Negative	Negative
259	33.42	137.19	28.14	4,593.69	Negative	Negative
703	21.71	334,350.00	Negative	Negative	33.71	112.79
928	16.98	7,788,455.58	32.01	349.40	Negative	Negative
1103	19.20	1,767,845.07	Negative	Negative	33.87	101.73
1150	18.54	2,752,642.96	30.14	1,212.83	30.59	898.81
1186	23.50	100,984.11	27.16	8,863.06	24.67	46,551.64
1341	18.17	3,358,847.22	25.08	109,088.94	31.48	2,683.82
A total of 1,793 mosquitoes were tested in pools, and 143 mosquitoes from positive pools were tested as individuals. The RNA numbers refer to estimated RNA copies estimated from a linear regression of CT values from a standard curve. Tissues with less than 200 copies of RNA were considered negative. A prerequisite for inclusion in the analysis and further testing was the presence of viral RNA in the abdomen tissue.						

                                    FIGURE LEGENDS

Fig. 1. Historical West Nile virus mosquito vector abundance and VI in Northern Colorado

regions during the 2023 season compared to other seasons. The years 2012-2024 (excluding

2023) are highlighted in grey, while the 2023 season is highlighted in red. BC: Boulder County,

CO; BE: Berthoud, CO; FC: Fort Collins, CO; LV: Loveland, CO.

Fig. 2. Confirmed human West Nile virus cases in Colorado during 2023 compared to other

years. The 2023 season, highlighted in yellow, showed a large spike in human cases compared to

other seasons from 2012-2024 highlighted in blue.

Fig. 3. Paired CT values for abdomen, thorax, and head compartments of WNV-positive

mosquitoes revealed distinct patterns by dissemination category. WNV-infected mosquitoes were

grouped into four phenotypes: disseminated infection (n = 8), failed dissemination (n = 2),

incomplete dissemination (n = 4), and weak midgut infection (n = 1). Plots include

corresponding whole body mosquito CT estimates determined by adding together and combining

RNA estimates from given tissue CT values, then converting them back into one whole body

estimate CT estimate. *Denotes multiple positive samples in one thorax pool, which artificially

deflated the CT estimate. **Denotes a weaker midgut infection that had disseminated into the

thorax but not the head.

Fig. 4. Correlation analyses of RNA values in paired tissue compartments of mosquito samples.

Correlation matrices for raw CT values (Panel A) and log-transformed RNA estimates (Panel B).

Fig. 5. Logistic regression model predicting complete dissemination of viral RNA via log-

transformed RNA estimates. Blue dots represent WNV-positive mosquitoes. The red line

represents the regression line and the shaded red area the 95% CI of the regression line. For

every log increase in abdominal RNA, the odds of WNV successfully disseminating to the head

and thorax increased by approximately 2.5 times (OR = 2.45, 95% CI: 0.81, 7.36, p = 0.11)

Fig. 6: Receiver Operating Characteristic (ROC) curve analysis for abdomen RNA value that

predicts binary threshold for complete dissemination. The binary infection threshold was set at

200 copies of viral RNA. The area under the curve (AUC) for prediction of infection was 0.8

(95% CI: 0.545, 1). The optimal threshold for prediction of complete dissemination at 200 copies

of RNA by abdomen RNA was ~59,000 copies with a sensitivity of 1 and a specificity of 0.625.

Raw CT values

Log RNA values