IRIS, a tool for the in-silico evaluation of mosquito control trial designs based on inundative releases

Jagadeesh Chitturi[1,$], Paulo C. Ventura[1,$], Allisandra G. Kummer[1], Chalmers Vasquez[2], Ethan SeRine[1], Megan D. Hill[1], Mattia Manica[3], Piero Poletti[3], John C. Beier[4], Keisuke Ejima[5], Michael Johansson[6], Stefano Merler[3], Hongjie Yu[7], John-Paul Mutebi[2], Maria Litvinova[8,#], André B.B. Wilke[8,#], Marco Ajelli[1,#,*]

[1]Laboratory for Computational Epidemiology and Public Health, Department of Epidemiology and Biostatistics, Indiana University School of Public Health, Bloomington, IN, USA

[2]Miami-Dade County Mosquito Control Division, Miami, FL, USA

[3]Center for Health Emergencies, Bruno Kessler Foundation, Trento, Italy

[4]Department of Public Health Sciences, Miller School of Medicine, University of Miami, Miami, FL, USA

[5]Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore, Singapore

[6]Network Science Institute, Northeastern University, Boston, MA, USA

[7]School of Public Health, Key Laboratory of Public Health Safety, Ministry of Education, Fudan University, Shanghai, China

[8]Department of Epidemiology and Biostatistics, Indiana University School of Public Health, Bloomington, IN, USA

[$]Co-first author

[#]Co-senior author

[*]Corresponding author: majelli@iu.edu

Abstract

Mosquito control strategies based on the mass release of modified males, such as
genetically modified mosquitoes (GMMs), aim to suppress wild populations by impairing
reproduction. Evaluating these interventions requires resource-intensive field trials, but a
lack of standardized implementation practices, particularly regarding release ratios of
modified males to wild female mosquitoes and trial timing, has led to variable outcomes.
This study's objective is to propose a modeling tool for the "in-silico" simulation of trial
designs before field implementation. To this aim, we developed an agent-based model of
mosquito population dynamics. As a case study, we calibrated the model using 2019-
2023 *Aedes aegypti* surveillance data from Miami-Dade County, Florida, and compared
two GMM trial designs as illustrative examples. Our results show that depending on the
implementation choices (e.g., trial start date and duration, release ratio), trials yield highly
variable outcomes. For example, changing the start date while fixing all other
implementation details can lead to effectiveness between 50% and 90%. Our findings
suggest that "in-silico" simulation is a valuable tool for improving trial protocol design,
allowing stakeholders to test strategies and reduce outcome uncertainty before
committing to a fieldwork experiment.

Keywords: Genetically Modified Mosquitoes; *Aedes aegypti*; Agent-Based Modeling;
Population Dynamics

Introduction

Traditional insecticide-based strategies are the cornerstone of mosquito control [1–4]. However, they have limitations such as insecticide resistance [5–7] and potential environmental impacts and effects on biodiversity, ecosystems, and human health, regardless of the effectiveness on mosquito control [8]. Therefore, in past decades, a new generation of interventions has emerged, which are based on mass releases of irradiated sterile mosquitoes, *Wolbachia*-infected mosquitoes, or genetically modified mosquitoes (GMM) [9–18]. These strategies, here referred to as "inundative" mosquito control strategies, rely on mass releases of adult male mosquitoes that are either sterile or carrying a modified gene or endosymbiont, which impair reproduction [12,19]. However, implementing inundative strategies still requires rigorous testing and evaluation to assess their effectiveness, ecological impact, and safety before large-scale implementation.

Field trials represent the gold standard in evaluating mosquito control strategies. However, they require substantial financial resources, logistical support, and expert coordination to ensure proper implementation [20,21]. To date, trial results of inundative mosquito control strategies have shown high variability, highlighting a major knowledge gap in the underlying causes for this heterogeneity [9–18]. While pilot studies in Kentucky, Panama, Brazil, China, Cuba, and the Cayman Islands have yielded promising results in decreasing mosquito populations [10,12–15,17], other trials in Florida, Mexico, Italy, and Malaysia have had more limited success [9,11,16,22]. The lack of standardized guidelines on the design and implementation of field trials for inundative mosquito control strategies has possibly contributed to these results. For instance, the release ratio of sterilized, *Wolbachia*-infected, or genetically modified male mosquitoes to wild-type males varied widely [9,10,14], including trials that released mosquitoes without assessing the ratio with the wild-type population [13,15]. As another example, the trial start and end dates were variable between studies, with some trials being conducted during the low mosquito season [22], others during the high season [16], and some spanning most of the year [13,15].

Without standardized trial designs, it is challenging to compare findings across different studies and settings. Mathematical models have been used to inform and guide the implementation of field trials [23], including those for Ebola [24] and dengue [25] vaccines, and SARS-CoV-2 antiviral therapies [26,27]. Thus, our study introduces IRIS (Inundative mosquito Release Intervention Simulator), an agent-based model for the "in-silico" simulation of trial designs for inundative mosquito releases. As a case study, we focus on GMM-based strategies, evaluating two alternative trial designs to control the *Aedes aegypti* population, a primary vector for dengue, Zika, and chikungunya [28] and target species for previous trials [11,12,15]. As a study location, we selected Miami-Dade County, Florida, as it has an abundant *Ae. aegypti* population, an archive of historical mosquito surveillance data, and a history of local transmission of arboviruses [29–31].

Methods

Data

We used mosquito surveillance data from January 1, 2019, to December 31, 2023, for Miami-Dade County, FL, which is collected and managed by the Miami-Dade County Mosquito Control Division. The surveillance system consists of about 300 BG-Sentinel-2 traps baited with dry ice that are deployed once per week for 24 hours. Collected mosquitoes are identified using taxonomic keys. Details on the data collection are reported in our previous publication [32]. Daily mean temperature data for Miami-Dade County, Florida, was retrieved from publicly available records from the Miami International Airport [33].

Model Formulation

As an illustrative example of our modeling tool, in this study, we simulate GMM-based trials that aim at reducing the mosquito population. GMM population suppression strategies involve the release of homozygous adult male mosquitoes carrying a gene that impairs the fertility of female offspring [34]. Specifically, when a genetically modified male mosquito mates with a wild adult female mosquito, it will produce a heterozygous offspring carrying that gene. Heterozygous females are infertile, while heterozygous males can mate with wild females, further propagating the gene in the wild mosquito population [19,34–37].

IRIS is an agent-based model that produces stochastic simulations of *Ae. aegypti* population dynamics considering four stages of mosquito development: egg, larva, pupa, and adult (Fig. 1A). The model estimates the number of *Ae. aegypti* in each stage of their life cycle for each calendar date at a time step of 0.1 days. Each mosquito is represented by an agent with the following attributes: sex (male or female), genetic makeup (genetically modified homozygous, genetically modified heterozygous, or wild type), and current stage of development (egg, larva, pupa, or adult). We assumed a 1:1 female-to-male ratio for eggs. At each time step of the simulation, each agent's transition into the next development stage or mortality and adult female oviposition are sampled from a Bernoulli distribution using the appropriate temperature-dependent rate [38–41]. Wild-type adult female mosquitoes lay eggs at a given temperature-dependent oviposition rate [39,41] with an average of 100 eggs laid per oviposition [42]. Newly laid eggs then become new agents in the model. Eggs' mortality is both temperature- and density-dependent to account for the carrying capacity of the environment (see section Model Calibration).

To simulate GMM-based trials, we periodically introduce a given number of new adult male agents that are genetically modified (homozygous) into the population. These mosquitoes continue through the life cycle in the same way as wild-type mosquitoes. When a wild-type female mates with a homozygous male, the offspring is heterozygous, and heterozygous females are infertile (i.e., their oviposition rate is set to 0). When a wild-type female mates with a heterozygous male, half of the offspring will be heterozygous for the transgene, and half will not carry the transgene. Moreover, we assume that female mosquitoes mate only once in their lifetime [43]. Finally, mosquitoes carrying the transgene have a different (generally lower) fitness than wild-type males, which reduces

Figure 1. A. Conceptual representation of the modeled mosquito life cycle. **B.** Conceptual representation of the GMM mechanisms of action and list of trial design variables that are investigated in this study.

128 the likelihood of successful mating [44]. A conceptual representation of the process is
129 shown in Fig. 1B.

Model Calibration

Temperature-dependent development and mortality rates of mosquito lifecycle stages
were taken from the literature [38–40]. The model also considers seasonal changes in
the carrying capacity to account for variations throughout the year [41]. We used a Markov
chain Monte Carlo (MCMC) procedure to obtain estimates of the posterior distributions of
the carrying capacity for two periods of the year. MCMC explored the likelihood of
observing the actual number of collected adult female *Ae. aegypti* specimens from Miami-
Dade County, Florida, from a negative binomial distribution of the mean given by the
model and the over-dispersion parameter, which is estimated as well [45]. Details are
reported in the Supplementary Material.

Simulation Analysis of Trial Implementation

The calibrated model simulates *Ae. aegypti* population dynamics from January 2019
through December 2023 and provides estimates of the daily number of *Ae. aegypti*
specimens for each life stage of development, sex, and genetic makeup. This allows us
to simulate different trial designs and trial implementation variables "in silico". In this
analysis, we simulated 2-arm trials where the control arm consists of the model
realizations that are run without any GMM-based intervention, while the intervention arm
consists of the model realizations simulating a GMM-based intervention.

We consider two trial designs:

- **Constant-release trial design.** The number of genetically modified (homozygous) male mosquitoes released per release in the intervention is set to obtain the desired ratio between the number of released male mosquitoes and the number of female mosquitoes per trap night collected during a baseline entomological survey. This baseline entomological survey is conducted before the start of the trial to assess the relative abundance of the mosquito population by monitoring traps for adult female mosquitoes (e.g., BG-Sentinel-2 traps). In this design, the number of released male mosquitoes per release is constant for the entire duration of the trial; thus, in populations following seasonal trends and where control is effective, the actual ratio of released male mosquitoes as compared to the actual female mosquito population changes over time.
- **Adaptive-release trial design.** The number of genetically modified (homozygous) male mosquitoes released per release in the intervention is dependent on the number of mosquitoes estimated in the control arm. The idea of the adaptive-release trial design is to rely on the control arm to obtain an estimate of the actual target female population relative abundance at any time point. Then, this estimate is used at the time of each release as the denominator to calculate the number of male mosquitoes to be released to obtain a constant release ratio as compared to the actual female mosquito population at any given point in time. Specifically, the number of male mosquitoes released at time t, $G(t)$ is given by: $G(t) = r \cdot s \cdot \bar{C}(t-7)$, where r is the target release ratio, s is the trap collection ratio estimated in the literature (which corresponds to the ratio between the collected adult female mosquitoes and the total number of adult female mosquitoes), and $\bar{C}(t-7)$ is the mean estimated number of mosquitoes in the control arm over the past week. The one-week lag was incorporated to account for the time required for trap collection, mosquito identification, data entry, and release planning.

In this study, the effectiveness of a simulated trial (trial outcome) is defined as the reduction in the *Ae. aegypti* female population within the monitored area during the trial period as:

$$E_{ff} = 1 - \frac{T}{C}$$

where T is the sum of estimated adult female mosquitoes in the treatment arm from 2 months after the start of the trial until the end of the trial, and C is the same quantity but for the control arm.

Results for each simulated trial are based on 100 stochastic model realizations obtained by uniformly sampling 100 indexes of MCMC corresponding to the joint posterior distributions of the carrying capacity for the two periods. Then, for each sampled index, we run the control and the treatment arm for both trial designs.

Standardized guidelines for trial implementation of GMM mosquito control strategies encompass a wide range of aspects and cover different steps of the implementation of the trial, from laboratory protocols to field practices. Among them, we investigate four variables: i) release ratio, ii) start date of the trial, iii) the duration of the baseline

entomological survey (which is relevant only for the constant-release trial design), iv)
release frequency, and v) trial duration. For the main analysis, we simulated 6-month trials
with a release frequency of once per week, a 2-month baseline entomological survey, and
that genetically modified mosquitoes have the same fitness as wildtype ones. We
investigated release ratios from 0.1 to 9.0 by 0.1 steps, and we consider starting dates of
the trial on the first day of the month for every month of the year in 2022. To estimate
release ratios, we considered a 24.6% trap collection ratio within the trap catchment area
[46]. Thus, an n:1 release ratio is relative to the collected female mosquitoes, which
represents an underestimate of all female mosquitoes within the area.

We conducted several sensitivity analyses varying the release frequency (every 3, 7, and
14 days), trial duration (90, 180, and 365 days), duration of the baseline entomological
survey (2, 6, and 12 months), and the fitness of the genetically-modified mosquitoes (set
at 50% and 75% relative to the fitness of wild-type male mosquitoes). Moreover, we
conducted a sensitivity analysis considering two alternative values for the collection ratio,
12.3% and 49.2%, encompassing the uncertainty reported in the reference study [46]. To
assess possible differences between years, we conducted a sensitivity analysis where
the trials were carried out in 2020, 2021, and 2022. Finally, to assess the impact of
temperature and carrying capacity on the effectiveness of the GMM-based trial, we
performed an analysis where the temperature was set to a constant 25°C and the carrying
capacity was kept constant throughout the year, while all other parameters remained the
same as in the main analysis.

Code Availability

212 The codebase for IRIS is openly available on GitHub at https://github.com/jagadeesh-
213 chitturi/iris-model.

Results

The weekly number of adult female *Ae. aegypti* per trap night estimated by IRIS
compares well with the observed data for Miami-Dade County over the entire study period
(Fig. 2A). The data show a year-round presence of *Ae. aegypti* with a clear seasonal trend
in its abundance, with peak activity taking place in July (about 10 collected adult female
Ae. aegypti per trap night) and lower activity during the winter months, with about 2
collected specimens per trap night.

Figure 2B shows an illustrative example of a trial simulated by IRIS: a constant-release
GMM trial of a 6-month duration, weekly releases, a 4:1 release ratio, starting on March
1, and with a 2-month baseline entomological survey. Figure 2C shows the adaptive-
release GMM trial based on the same implementation details as the constant-release trial
design. The constant-release and adaptive-release trial designs follow a similar trajectory
to the control arm for the first few weeks of the trial before diverging. Overall, we estimated
an effectiveness of 47.3% (IQR: 37.5%-58.4%) and 81.3% (IQR: 71.3%-92.-0%) for the
constant-release ratio and adaptive-release ratio, respectively. These results are not
indicative of which trial design has a better suppression effectiveness; they merely
represent the result of the considered implementation details and should not be
interpreted as a comparative analysis of the two designs to identify an optimal one. For

Figure 2. A. Number of collected adult female *Ae. aegypti* per trap night by month of the year as reported in the Miami-Dade mosquito surveillance system and as estimated by the model. **B.** Daily number of collected adult female *Ae. aegypti* per trap night estimated by the model in the control arm and in the treatment arm of a constant-release trial. Trial implementation details are: start date May 1, 6-month trial duration, weekly releases, 4:1 release ratio, and 2 months baseline entomological survey. **C.** As B, but for an adaptive-release trial design. Trial implementation details are the same as in panel B, except for the baseline entomological survey, which is not needed for this trial design.

example, this simulated implementation of the adaptive-release trial requires a larger number of released mosquitoes over the entire trial period, which yields better effectiveness.

We used IRIS to explore different trial implementation details and their impact on the estimated effectiveness. First, we fixed weekly release frequency for a 6-month trial duration. For the constant-release design, we also fixed the duration of the entomological survey at 2 months. Regardless of the trial design, we found that the effectiveness is highly variable (Fig. 3A and B). For any trial start date, the effectiveness is highly dependent on the release ratio, which can lead to the outcome spanning the entire 0 to ~100% range. If the release ratio is 12:1 or higher, the effectiveness is over 90% regardless of the trial start date. However, for lower values of the release ratio, the effectiveness is highly dependent on the trial start date (Fig. 3C). The estimated dependencies of trial effectiveness on the mosquito release ratio and trial start date may

Figure 3. A. Estimated effectiveness of a constant-release trial as a function of the start date of the trial and release ratio. The other trial variables are set as follows: 6-month trial duration, weekly release, and 2-month baseline entomological survey. **B.** As A, but for an adaptive-release trial. **C.** Estimated median and IQR of the effectiveness of a constant-release and adaptive-release trial designs given a release ratio, for any possible trial start date. The other trial variables are set as in panel A. **D.** Estimated trial effectiveness as a function of the total number of genetically modified adult male mosquitoes released per trap over the entire duration of the trial. Each dot represents a single simulation; dashed lines represent the mean. The panel shows the results for 6-month trials with and weekly releases, any starting date, and any release ratio from 1:0.1 to 9:1. For the constant-release trial design, the duration of the baseline entomological survey is set to 2 months.

help explain the variability observed in field trials, which were based on different designs
and implementations.

A more stable indicator we found to determine the effectiveness is the overall number of
released mosquitoes during the entire duration of the trial (Fig. 3D). This holds true for
any trial start date, release ratio, and trial design. In other words, a trial design with a
given release ratio and start date requires a given number of mosquitoes to be released
over the entire duration of the trial; this number essentially determines the final
effectiveness of the trial.

Figure 4. A. Estimated median and IQR of the effectiveness of a constant-release and adaptive-release trial designs given the release frequency, for any possible trial start date. The release ratio is set to 4 and the trial duration to 6 months; for the constant-release design, the duration of the baseline entomological survey is set to 2 months. **B.** Estimated median and IQR of the effectiveness of a constant-release and adaptive-release trial designs given the trial duration, for any possible trial start date. The release ratio is set to 4 and the release frequency to once per week; for the constant-release design, the duration of the baseline entomological survey is set to 2 months. **C.** Estimated median and IQR of the effectiveness of a constant-release and adaptive-release trial designs given the fitness of genetically modified male mosquitoes, for any possible trial start date. The release ratio is set to 4, the trial duration is set to 6 months, and the release frequency to once per week; for the constant-release design, the duration of the baseline entomological survey is set to 2 months. **D.** Estimated median and IQR of the effectiveness of a constant-release trial design given the duration of the baseline entomological survey, for any possible trial start date. The release ratio is set to 4, the trial duration is set to 6 months, and the release frequency to once per week.

Our results show that more frequent releases, longer trial durations, and greater fitness
of transgenic mosquitoes increase the effectiveness of the trial and decrease the
variability in the estimated effectiveness (Fig. 4A-C). For the constant-release design, a
longer baseline entomological survey decreases variability (Fig. 4D).

We found that all our findings remain unchanged if trials are simulated using data from
different years (see Supplementary Material). However, in a hypothetical scenario where
the temperature and carrying capacity do not change over the course of the year, we
estimated that a much larger release ratio is needed to obtain the same effectiveness
estimated for the simulated trials using data for Miami-Dade County, FL (see
Supplementary Material). Given that no changes take place over time and thus the
mosquito population is constant year-round, the variability in the estimated effectiveness
drastically decreases, suggesting that in areas where the mosquito population is more
constant throughout the year, trial results may be more stable. Finally, we estimated a
higher effectiveness of the trials for higher collection ratios of the traps, but our main
finding about the lower variability in the effectiveness estimated using the adaptive-
release design remains unchanged (see Supplementary Material).

Discussion

In this study, we propose IRIS, an agent-based model designed to be a decision-support
tool for public health authorities and researchers to in-silico test different trial designs and
implementation details. This model builds on the wide body of existing literature on
mathematical modeling for the evaluation of mosquito control strategies based on
inundative mosquito releases [19,47–62], adapting it for trial simulation.

This analysis focuses on a specific study case: GMM-based trials for *Ae. aegypti*
suppression in Miami-Dade County, Florida. Our results show that the effectiveness of
GMM trials is highly sensitive to implementation choices, particularly the release ratio of
modified males and the trial start date. This finding provides a possible explanation for
the high variability observed in field trials across different settings [9–18]. Other trial
implementation details, such as differences in release frequency, trial duration, and length
of baseline entomological survey (for the constant-release trial design), may have further
contributed to the observed variability in field estimates. On the other hand, our analysis
suggests that the total number of modified mosquitoes released over the entire trial period
may be a more stable indicator of trial effectiveness. This metric appears to transcend
specific trial designs and implementation choices. Future research that considers a wider
range of designs would be needed to further support/challenge this preliminary finding.

Our study is not designed to provide specific recommendations about field trial designs
and their implementation details. Thus, we are not endorsing the implementation of a
specific design. Instead, our work is intended to showcase a modeling approach for the
in-silico simulation of trial designs and provide general insights to better interpret results
from previous field studies. Our primary contribution is to provide an inexpensive and
flexible modeling tool to test trial designs before committing the significant monetary and
logistic resources needed for field implementation. This is especially relevant for proof-of-
concept trials, which are the crucial first step in a pipeline toward full-scale public health
adoption. The outcomes of such initial trials are critical, as they determine whether
stakeholders may decide to commit resources to subsequent, larger-scale trials focused
on optimizing long-term effectiveness (e.g., year-round release schedules). This
optimization phase is, in turn, the prerequisite for the eventual, sustainable
implementation of these strategies in real-world public health programs. IRIS directly
supports the first step by informing the design of field trials with higher likelihood of

success. Furthermore, IRIS codebase can be readily adapted to support both the design
of larger-scale trials and the optimization of a wide range of other mosquito control
interventions.

Our study is subject to the following limitations. First, our model simplifies certain
biological, ecological, and environmental aspects. For example, it assumes a 1:1 sex
ratio, female mosquitoes mating only once in their lifetime, uniform mixing between male
and female mosquitoes, and relies on specific temperature-dependent parameters
derived from the literature on *Ae. aegypti* rather than on the specifics of the population in
the study site. While these assumptions are standard in many mosquito population
dynamic models, they may not fully capture the nuances of real-world dynamics. Second,
our analysis did not consider several logistical constraints and practical considerations
associated with the implementation of field trials (e.g., implementation costs, required
manpower, number of deployed traps per arm, number of clusters per arm, and the
constant availability of a sufficient number of genetically modified male mosquitoes to
release). However, although these constraints and considerations are crucial for the
successful implementation of a trial, these should be considered by stakeholders to
determine which designs and implementation details represent feasible options for them.
Then, our tool can be used to compare these options to inform their decision among them.
Third, our model was validated against historical mosquito surveillance data from Miami-
Dade County, Florida, providing a good fit to observed population trends. However, it has
not yet been validated against data from actual GMM field trials. Such validation would
be crucial to confirm the model's predictive power regarding the impact of GMM releases.
Fourth, the current model does not account for spatial heterogeneity in breeding site
availability, microclimatic conditions, etc., which may affect the outcome of a trial.
Incorporating spatial factors into future model iterations would enhance the model's
capability to test more nuanced factors affecting the outcome of a trial. For example, the
model currently considers a fitness parameter that represents the (potentially) reduced
ability of genetically modified male mosquitoes to successfully mate. A spatial model
would allow for the evaluation of different mechanisms driving such a reduction (e.g.,
shortened flight range, lower energy level) as well as boundary and spillover effects that
are not considered here. Other differences between genetically-modified and wild-type
male mosquitoes (e.g., lifespan, which is currently embedded into the fitness parameter)
could be explicitly modeled and investigated as well. Fifth, our analysis focuses solely on
GMM-based interventions and does not consider the potential for integrating other
mosquito control methods. In practice, GMM releases might be combined with other
interventions, such as the application of *Bacillus thuringiensis israelensis* (*Bti*) or other
larvicides before the start of a trial.

In conclusion, this study presents a modeling tool designed for the "in-silico" simulation
of field trials for mosquito control strategies based on inundative mosquito releases.
Grounded in empirical mosquito surveillance data and incorporating key environmental
factors, such as temperature-dependent development and mortality rates and temporal
variations in carrying capacity, IRIS offers a platform for exploring different trial designs.
While our analysis focused on *Ae. aegypti* in Miami-Dade County, our modeling approach
allows for its adaptation to other mosquito species and geographic locations, making it an
asset in the broader field of vector control. By enabling the identification of trial

implementations with the highest likelihood of success, our model can contribute to the
development of standardized guidelines for field trial design.

Funding

349 M.L., A.B.B.W., and M.A. were supported by the National Science Foundation (DMS-
350 2526926). The funder had no role in the design of the study and collection, analysis, and
351 interpretation of data, and in writing the manuscript.

Competing Interests

353 H.Y. has received research funding from Sanofi Pasteur, Shenzhen Sanofi Pasteur
354 Biological Products Co., Ltd, Shanghai Roche Pharmaceutical Company, and SINOVAC
355 Biotech Ltd. None of these funding is related to this work. The other authors have declared
356 that no competing interests exist.

References

1.   Wilke ABB, Vasquez C, Carvajal A, Ramirez M, Cardenas G, Petrie WD, Beier JC. 2021
Effectiveness of adulticide and larvicide in controlling high densities of Aedes aegypti in
urban environments. *PLoS One* **16**, e0246046. (doi:10.1371/journal.pone.0246046)

2.   Juri MJD, Zaidenberg M, Santana M. 2013 The efficacy of a combined larvicide-adulticide
in ultralow volume and fumigant canister formulations in controlling the dengue vector
Aedes aegypti (Diptera: Culicidae) in Northwest of Argentina. *Parasitol. Res.* **112**, 1237–
1246. (doi:10.1007/s00436-012-3256-4)

3.   Harburguer L, Seccacini E, Licastro S, Zerba E, Masuh H. 2012 Droplet size and efficacy of
an adulticide-larvicide ultralow-volume formulation on Aedes aegypti using different
solvents and spray application methods. *Pest Manag. Sci.* **68**, 137–141.
(doi:10.1002/ps.2237)

4.   McAllister JC *et al.* 2020 Mosquito Control Activities during Local Transmission of Zika Virus,
Miami-Dade County, Florida, USA, 2016. *Emerg. Infect. Dis.* **26**, 881–890.
(doi:10.3201/eid2605.191606)

5.   Hirata K, Komagata O, Itokawa K, Yamamoto A, Tomita T, Kasai S. 2014 A Single Crossing-
Over Event in Voltage-Sensitive Na+ Channel Genes May Cause Critical Failure of Dengue
Mosquito Control by Insecticides. *PLoS Negl. Trop. Dis.* **8**, e3085.
(doi:10.1371/journal.pntd.0003085)

6.   Moyes CL *et al.* 2017 Contemporary status of insecticide resistance in the major Aedes
vectors of arboviruses infecting humans. *PLoS Negl Trop Dis.* **11**, 1–20.
(doi:10.1371/journal.pntd.0005625)

7.   Pocquet N *et al.* 2013 Multiple Insecticide Resistances in the Disease Vector Culex p.
Quinquefasciatus from Western Indian Ocean. *PLoS One* **8**, 1–14.
(doi:10.1371/journal.pone.0077855)

8.   Silver MK, Shao J, Zhu B, Chen M, Xia Y, Kaciroti N, Lozoff B, Meeker JD. 2017 Prenatal naled
and chlorpyrifos exposure is associated with deficits in infant motor function in a cohort of
Chinese infants. *Environ. Int.* **106**, 248–256. (doi:10.1016/j.envint.2017.05.015)

9.   Bellini R, Medici A, Puggioli A, Balestrino F, Carrieri M. 2013 Pilot field trials with aedes
albopictus irradiated sterile males in Italian Urban Areas. *J. Med. Entomol.* **50**, 317–325.
(doi:10.1603/ME12048)

10.   Harris AF *et al.* 2012 Successful suppression of a field mosquito population by sustained
release of engineered male mosquitoes. *Nat. Biotechnol.* **30**, 828–830.
(doi:10.1038/nbt.2350)

11.   Lacroix R *et al.* 2012 Open field release of genetically engineered sterile male Aedes aegypti
in Malaysia. *PLoS One* **7**, 1–9. (doi:10.1371/journal.pone.0042771)

12.   Carvalho DO, McKemey AR, Garziera L, Lacroix R, Donnelly CA, Alphey L, Malavasi A,
Capurro ML. 2015 Suppression of a field population of Aedes aegypti in Brazil by sustained
release of transgenic male mosquitoes. *PLoS Negl. Trop. Dis.* **9**, 2016–2018.
(doi:10.1371/journal.pntd.0003864)

13.   Gorman K *et al.* 2016 Short-term suppression of Aedes aegypti using genetic control does
not facilitate Aedes albopictus. *Pest Manag. Sci.* **72**, 618–628. (doi:10.1002/ps.4151)

14.   Zheng X *et al.* 2019 Incompatible and sterile insect techniques combined eliminate
mosquitoes. *Nature* **572**, 56–61. (doi:10.1038/s41586-019-1407-9)

15.   Gato R *et al.* 2021 Sterile insect technique: Successful suppression of an Aedes aegypti field
population in Cuba. *Insects* **12**. (doi:10.3390/insects12050469)

16.   Martín-Park A *et al.* 2022 Pilot trial using mass field-releases of sterile males produced with
the incompatible and sterile insect techniques as part of integrated Aedes aegypti control
in Mexico. *PLoS Negl. Trop. Dis.* **16**. (doi:10.1371/journal.pntd.0010324)

17.   Mains JW, Brelsfoard CL, Rose RI, Dobson SL. 2016 Female adult aedes Albopictus
suppression by Wolbachia-infected male mosquitoes. *Sci. Rep.* **6**. (doi:10.1038/srep33846)

18.   Bellini R, Carrieri M, Balestrino F, Puggioli A, Malfacini M, Bouyer J. 2021 Field
Competitiveness of *Aedes albopictus* (Diptera: Culicidae) Irradiated Males in Pilot Sterile
Insect Technique Trials in Northern Italy. *J. Med. Entomol.* **58**, 807–813.
(doi:10.1093/jme/tjaa235)

19.   Phuc HK *et al.* 2007 Late-acting dominant lethal genetic systems and mosquito control.
BMC Biol. **5**, 11. (doi:10.1186/1741-7007-5-11)

20.   WHO. 2014 Third Meeting of the Vector Control Advisory Group. , 1–52.

21.   Wilke ABB, Beier JC, Benelli G. 2018 Transgenic Mosquitoes – Fact or Fiction? *Trends
Parasitol.* **34**, 456–465. (doi:10.1016/j.pt.2018.02.003)

22.   Mains JW, Kelly PH, Dobson KL, Petrie WD, Dobson SL. 2019 Localized Control of Aedes
aegypti (Diptera: Culicidae) in Miami, FL, via Inundative Releases of Wolbachia-Infected
Male Mosquitoes. *J. Med. Entomol.* **56**, 1296–1303. (doi:10.1093/jme/tjz051)

420 23. Halloran ME *et al.* 2017 Simulations for designing and interpreting intervention trials in
421   infectious diseases. *BMC Med.* **15**, 223. (doi:10.1186/s12916-017-0985-3)

422 24. Dean NE *et al.* 2019 Design of vaccine efficacy trials during public health emergencies. *Sci.*
423   *Transl. Med.* **11**. (doi:10.1126/scitranslmed.aat0360)

424 25. Odio CD *et al.* 2023 Phase 1 trial to model primary, secondary, and tertiary dengue using a
425   monovalent vaccine. *BMC Infect. Dis.* **23**, 345. (doi:10.1186/s12879-023-08299-5)

426 26. Zheng Q, Bao C, Ji Y, Li P, Ma Z, Wang X, Meng Q, Pan Q. 2023 Treating SARS-CoV-2 Omicron
427   variant infection by molnupiravir for pandemic mitigation and living with the virus: a
428   mathematical modeling study. *Sci. Rep.* **13**, 5474. (doi:10.1038/s41598-023-32619-z)

429 27. Lingas G *et al.* 2022 Effect of remdesivir on viral dynamics in COVID-19 hospitalized
430   patients: a modelling analysis of the randomized, controlled, open-label DisCoVeRy trial.
431   *Journal of Antimicrobial Chemotherapy* **77**, 1404–1412. (doi:10.1093/jac/dkac048)

432 28. Aryaprema VS, Steck MR, Peper ST, Xue R De, Qualls WA. 2023 A systematic review of
433   published literature on mosquito control action thresholds across the world. *PLoS Negl.*
434   *Trop. Dis.* **17**, 1–22. (doi:10.1371/journal.pntd.0011173)

435 29. Ventura PC *et al.* 2024 Forecasting the relative abundance of Aedes vector populations to
436   enhance situational awareness for mosquito control operations. *PLoS Negl. Trop. Dis.* **18**,
437   e0012671. (doi:10.1371/journal.pntd.0012671)

438 30. Coatsworth H *et al.* 2022 A molecular surveillance-guided vector control response to
439   concurrent dengue and West Nile virus outbreaks in a COVID-19 hotspot of Florida. *The*
440   *Lancet Regional Health - Americas* **11**, 100231. (doi:10.1016/j.lana.2022.100231)

441 31. Likos A *et al.* 2016 Local Mosquito-Borne Transmission of Zika Virus — Miami-Dade and
442   Broward Counties, Florida, June–August 2016. *MMWR Morb. Mortal. Wkly. Rep.* **65**, 1032–
443   1038. (doi:10.15585/mmwr.mm6538e1)

444 32. Vasquez C, Multini LC, Mutebi J-P, SeRine E, Hill MD, Litvinova M, Ajelli M, Wilke ABB. 2025
445   Spatiotemporal Dynamics of *Aedes aegypti* and *Culex quinquefasciatus* populations in
446   Miami-Dade County, Florida. (doi:10.1101/2025.10.01.679884)

447 33. https://www.ncei.noaa.gov/cdo-web/. In press. NOAA - Climate Data Online.

448 34. Alphey L. 2002 Re-engineering the sterile insect technique. *Insect Biochem. Mol. Biol.* **32**,
449   1243–1247. (doi:10.1016/S0965-1748(02)00087-5)

450 35. Fu G, Condon KC, Epton MJ, Gong P, Jin L, Condon GC, Morrison NI, Dafa'alla TH, Alphey L.
451   2007 Female-specific insect lethality engineered using alternative splicing. *Nat. Biotechnol.*
452   **25**, 353–357. (doi:10.1038/nbt1283)

453 36. World Health Organization, Who. 2009 Progress and prospects for the use of genetically
454   modified mosquitoes to inhibit disease transmission. *WHO*. (doi:10.978-924-1599238)

455 37. Wilke ABB, Nimmo DD, St John O, Kojin BB, Capurro ML, Marrelli MT. 2009 Mini-review:
456   Genetic enhancements to the sterile insect technique to control mosquito populations.
457   *Asia. Pac. J. Mol. Biol. Biotechnol.* **17**, 65–74.

38. Cristina Farnesi L, Jesus Martins A, Valle D, Lazzaro Rezende G. 2009 Embryonic
development of Aedes aegypti (Diptera: Culicidae): influence of different constant
temperatures. *Mem Inst Oswaldo Cruz, Rio de Janeiro*. **104**.

39. Yang HM, Macoris MLG, Galvani KC, Andrighetti MTM, Wanderley DMV. 2009 Assessing the
effects of temperature on the population of Aedes aegypti, the vector of dengue.
Epidemiol. Infect. **137**, 1188–1202. (doi:10.1017/S0950268809002040)

40. Yang HM, Macoris M de L da G, Galvani KC, Andrighetti MTM. 2011 Follow up estimation
of Aedes aegypti entomological parameters and mathematical modellings. *Biosystems* **103**,
360–371. (doi:10.1016/j.biosystems.2010.11.002)

41. Kummer AG, Wilke ABB, Ventura PC, Vasquez C, Medina J, Ejima K, Ajelli M. 2023
Comparing the effectiveness of adulticide application interventions on mitigating local
transmission of dengue virus. *J. Pest Sci. (2004).* (doi:10.1007/s10340-024-01787-w)

42. Clemons A *et al.* 2010 *Aedes aegypti* : An Emerging Model for Vector Mosquito
Development: Figure 1. *Cold Spring Harb. Protoc.* **2010**, pdb.emo141.
(doi:10.1101/pdb.emo141)

43. Degner EC, Harrington LC. 2016 Polyandry depends on postmating time interval in the
dengue vector Aedes aegypti. *American Journal of Tropical Medicine and Hygiene* **94**, 780–
785. (doi:10.4269/ajtmh.15-0893)

44. Massonnet-bruneel B, Corre-catelin N, Lacroix R, Lees RS, Hoang KP, Nimmo D, Alphey L,
Reiter P. 2013 Fitness of Transgenic Mosquito Aedes aegypti Males Carrying a Dominant
Lethal Genetic System. **8**, 1–7. (doi:10.1371/journal.pone.0062711)

45. Gilks WR, Richardson S, Spiegelhalter D, editors. 1995 *Markov Chain Monte Carlo in*
Practice. Chapman and Hall/CRC. (doi:10.1201/b14835)

46. Johnson PH, Spitzauer V, Ritchie SA. 2012 Field sampling rate of BG-sentinel traps for Aedes
aegypti (Diptera: Culicidae) in Suburban Cairns, Australia. *J. Med. Entomol.* **49**, 29–34.
(doi:10.1603/ME11116)

47. Xia S, Ury J, Powell JR. 2021 Increasing effectiveness of genetically modifying mosquito
populations: Risk assessment of releasing blood-fed females. *American Journal of Tropical*
Medicine and Hygiene **104**, 1895–1906. (doi:10.4269/ajtmh.19-0729)

48. O'Reilly KM *et al.* 2019 Estimating the burden of dengue and the impact of release of wMel
Wolbachia-infected mosquitoes in Indonesia: A modelling study. *BMC Med.* **17**, 1–14.
(doi:10.1186/s12916-019-1396-4)

49. Patinvoh RJ, Susu AA. 2014 Mathematical Modelling of Sterile Insect Technology for
Mosquito Control. *Advances in Entomology* **02**, 180–193. (doi:10.4236/ae.2014.24027)

50. Esteva L, Mo Yang H. 2005 Mathematical model to assess the control of Aedes aegypti
mosquitoes by the sterile insect technique. *Math. Biosci.* **198**, 132–147.
(doi:10.1016/j.mbs.2005.06.004)

51. Cai L, Huang J, Song X, Zhang Y. 2019 Bifurcation analysis of a mosquito population model
for proportional releasing sterile mosquitoes. *Discrete Continuous Dyn. Syst. Ser. B* **24**,
6279–6295. (doi:10.3934/dcdsb.2019139)

52. Li J. 2008 Differential equations models for interacting wild and transgenic mosquito
populations. *J. Biol. Dyn.* **2**, 241–258. (doi:10.1080/17513750701779633)

53. Li Y, Li J. 2019 Discrete-time model for malaria transmission with constant releases of sterile
mosquitoes. *J. Biol. Dyn.* **13**, 225–246. (doi:10.1080/17513758.2018.1551580)

54. Iboi EA, Gumel AB, Taylor JE. 2020 Mathematical modeling of the impact of periodic release
of sterile male mosquitoes and seasonality on the population abundance of malaria
mosquitoes.

55. Li J. 2010 Modeling of mosquitoes with dominant or recessive transgenes and allee effects.
Mathematical Biosciences and Engineering **7**, 99–121. (doi:10.3934/mbe.2010.7.99)

56. Li J, Yuan Z. 2014 Modelling releases of sterile mosquitoes with different strategies. *J. Biol.*
Dyn. **9**, 1–14. (doi:10.1080/17513758.2014.977971)

57. Sacker RJ, Von Bremen HF. In press. Some Stability Results in a Model for Genetically
Altered Mosquitoes.

58. Robert MA, Legros M, Facchinelli L, Valerio L, Ramsey JM, Scott TW, Gould F, Lloyd AL. 2012
Mathematical Models as Aids for Design and Development of Experiments: The Case of
Transgenic Mosquitoes. *J. Med. Entomol.* **49**, 1177–1188. (doi:10.1603/ME11205)

59. Sacker RJ, Von Bremen HF. In press. Global Asymptotic Stability in the Jia Li Model for
Genetically Altered mosquitoes.

60. Sacker2 RJ, Von Bremen HF. In press. Stability in a Periodic Model for Genetically Altered
Mosquitos.

61. White SM, Rohani P, Sait SM. 2010 Modelling pulsed releases for sterile insect techniques:
Fitness costs of sterile and transgenic males and the effects on mosquito dynamics. *Journal*
of Applied Ecology **47**, 1329–1339. (doi:10.1111/j.1365-2664.2010.01880.x)

62. Dumont Y, Tchuenche JM. 2012 Mathematical studies on the sterile insect technique for
the Chikungunya disease and Aedes albopictus. *J. Math. Biol.* **65**, 809–854.
(doi:10.1007/s00285-011-0477-6)

524