Conkazal-M1 from the MKAVA family of conotoxins – a dual-function protease inhibitor and neuroactive peptide

Celeste M. Hackney[1], Thomas Lund Koch[2,3], Nicklas Lund Ryding[1], Aymeric Rogalski[2], Kevin Chase[4], Matías Leonel Giglio[4], Samuel S. Espino[4], Zildjian G. Acyatan[2], Maren Watkins[4], Baldomero M. Olivera[4], Helena Safavi-Hemami[2,4], Kaare Teilum[1]*, and Lars Ellgaard[1]*

Affiliations:

[1] Department of Biology, Linderstrøm-Lang Centre for Protein Science, University of Copenhagen, Copenhagen, Denmark

[2] Department of Biochemistry, University of Utah, Salt Lake City, Utah, United States of America

[3] Department of Biomedical Sciences, University of Copenhagen, Denmark

[4] School of Biological Sciences, University of Utah, Salt Lake City, Utah, United States of America

*Correspondence: Lars Ellgaard, University of Copenhagen, Department of Biology, Ole Maaløes Vej 5, DK-2200 Copenhagen N., Denmark. Email: lellgaard@bio.ku.dk

Kaare Teilum, University of Copenhagen, Department of Biology, Ole Maaløes Vej 5, DK-2200 Copenhagen N., Denmark. Email: kaare.teilum@bio.ku.dk

20   Keywords: Cone snails, conkazal, conotoxin, Kazal-type protease inhibitor, Kunitz domain,

21   molecular evolution, schistosomin

ABSTRACT

Marine cone snails produce a diverse array of bioactive peptides, known as conotoxins, in their venom. Given their high target potency and specificity, conotoxins are attractive compounds for the development of precision research tools and pharmacological agents. Here, we provide the first experimental characterization of a conotoxin from the MKAVA superfamily, conkazal-M1, from *Conus magus*. Using NMR spectroscopy, we show that conkazal-M1 adopts a fold characteristic of the Kazal-type protease inhibitor family, featuring a Glu residue at the inhibitory P1 position. Recombinantly expressed conkazal-M1 inhibits the proteolytic activity of Subtilisin A with an apparent Ki of 1.1 μM. In addition, conkazal-M1 partially inhibits calcium transients in mouse sensory neurons, suggesting a potential role in modulating ion-channel activity as seen for many other toxins. The dual function of conkazal-M1 in protease inhibition and neuroactivity is analogous to the dual function of several toxins harboring a Kunitz-type fold. The well-conserved sequence of the MKAVAs indicates an evolutionary trajectory where these proteins face an adaptive conflict when mutations that enhance one activity compromise the other. Collectively, this work provides new structural and functional insight into a previously uncharacterized toxin superfamily in cone snails and expands our knowledge about the toxin arsenal available to venomous animals.

INTRODUCTION

Peptide toxins from animal venoms target a broad range of biological processes to subdue prey or deter predators (1). To exert these effects, toxins act by disrupting the cellular ion balance or by interfering with signaling pathways mediated by transmembrane proteins such as ion channels, G protein–coupled receptors (GPCRs), transporters, and tyrosine kinases (2). Other major classes of toxins target hemostatic processes, such as blood coagulation (3). Due to the evolutionary conservation of such key processes and pathways, many peptide toxins hold great potential as research tools and potential pharmaceutical compounds.

The diverse functions of peptide toxins are enabled by a variety of molecular structures. Many of these toxins fold into small, compact structures stabilized by multiple disulfide bonds. While the cysteine residues involved in disulfide-bond formation are typically conserved, inter-cysteine residues normally undergo rapid evolution that generate diversity (4, 5). Consequently, toxin sequences – even within the same evolutionary family – often exhibit a high degree of sequence variation.

Several prevalent toxins function as protease inhibitors, reflecting the critical role of protease activity in the physiology of target animals. For example, venom protease inhibitors target proteases in the blood-coagulation system (causing internal bleeding and hypotension), enzymes of the cholinergic system (disrupting neurotransmitter metabolism), and the immune system (modulating immune and inflammatory responses). Common protease-inhibitor folds present in venoms include Kazal-, Kunitz- and cystatin-like domains, Snake Venom Serine Protease Inhibitors, and Cysteine-Rich Secretory Proteins. As is common for other venom peptides, such folds are often recruited into the venom gland from endogenous genes and are subsequently "weaponized" (6). Through this process, these compact and stable structural scaffolds can be

repurposed to perform diverse functions that differ from the original, while retaining their overall structure.

Well-characterized Kunitz-type toxins are known to have acquired new functions different from the original protease-inhibitor function through evolution. For instance, the conkunitzins from venomous marine cone snails (7, 8), as well as mambaquaretin-1 (9) and calcicludine (10) from the Eastern green mamba, have evolved to inhibit ion channels or receptors. Moreover, certain Kunitz-like toxins display a dual function as both protease inhibitors and ion channel blockers (11-14). At the molecular level, the dual function of the tarantula Kunitz-type toxin HWTX-X is achieved through two distinct binding interfaces (12). It has also been shown that Kunitz-type protease inhibitors from venomous snakes undergo positive selection at sites that cover 30-50% of the domain (15). These examples provide textbook illustrations of the sequence plasticity often observed in toxins that allows them to adopt new functions using a preexisting structural scaffold.

A prominent example of a protease-inhibitor fold found in several venoms is the Kazal-type domain that inhibits a broad range of serine proteases. Kazal-type venom peptides are found in various venomous animals including assassin bugs, shrew, honeybees, and wasps, where they have been associated with functions like anti-coagulation (16-18), pro-coagulation (19), antimicrobial activity (18), melanization in insects (20), and analgesia (19). In humans, the SPINK family of Kazal-type protease inhibitors (KPIs) play important functions in regulating cellular protease activity, as exemplified by the pancreatic trypsin inhibitor SPINK1, where sequence variants are associated with pancreatitis and pancreatic cancer (21). Another example of a well-characterized KPI is greglin from the locust *Schistocerca gregaria*, which inhibits both elastase and subtilisin with low nanomolar Ki-values (22). In general, KPIs function as substrate analogs.

They act by inserting their inhibitory P1 residue, the residue positioned immediately adjacent to

the scissile bond, into the specificity pocket of the target protease (23). This residue is typically

located two positions C-terminally to the second cysteine in the sequence on the so-called reactive-

site loop (23). The nature of the P1 residue can vary considerably, in turn influencing substrate

specificity – for instance, trypsin-inhibiting KPIs often contain a lysine or arginine residue at the

P1 position to mimic physiological trypsin substrates (23). Classical KPIs harbor 40-60 amino acid

residues and fold into a structure that comprises an α-helix and a three-stranded β-sheet and is held

together by three disulfide bonds (23). KPIs that deviate from this definition (e.g. more or fewer

disulfides or varying disulfide patterns) are classified as non-classical KPIs (see Results for more

details).

The conotoxins produced by venomous marine cone snails are peptides encoded as precursor

sequences that comprise a well-conserved N-terminal signal sequence that directs the peptide to

the endoplasmic reticulum followed in some cases by a propeptide of intermediate sequence

conservation, and, finally, the mature conotoxin sequence. Approximately 70 different conotoxin

gene families have presently been identified (24-26). While most mature conotoxins are 15-45

residues in length, many larger peptides also exist (25). Many families of large conotoxins – also

known as macro-conotoxins (27) – remain uncharacterized due to the difficulty in producing large

disulfide-bonded peptides. These larger peptides can be expected to perform functions unavailable

to shorter conotoxins, e.g., as enzymes or as a consequence of interacting with their targets through

a larger binding interface.

Here, we surveyed venom gland transcriptomes to identify new macro-conotoxin families.

We describe the previously uncharacterized MKAVA family of conotoxins and show by NMR

spectroscopy that an MKAVA-family toxin from *Conus magus*, conkazal-M1, comprises a single

non-classical Kazal-type domain with both protease inhibitor activity and the ability to inhibit

KCl-induced calcium influx in a subset of dorsal root ganglia (DRG), as also seen for several

toxins harboring a Kunitz-type fold.

RESULTS

The MKAVA conotoxin superfamily is widely distributed among venomous cone snails

To characterize the MKAVA conotoxin superfamily, we mined publicly available venom gland transcriptomes from 49 cone snail species and identified 89 transcripts sharing the MKAVA consensus signal sequence (MKAVAVFLVVALAVAYG) with slight variations (Figure S1; all sequences provided in Supplementary File 1). In addition to their wide distribution across *Conus* species, we found that closely related MKAVA-like peptides are also found in other members of Conoidea, the larger superfamily of marine snails to which cone snails belong. Searches against the NCBI Sequence Read Archive (SRA) database identified similar sequences in species belonging to Drillidae, Turridae, Profundiconus, and Terebridae (see Materials and Methods section for SRA entry numbers). It should be noted that these sequences are incomplete, as they derive from unassembled short-read datasets.

In *Conus*, all identified sequences lack an apparent propeptide with the predicted mature toxin sequence (~73 residues) following immediately after the signal sequence. The toxins exhibit a high level of conservation (Figure 1a) – within the mature toxin sequence, 32 residues are >95% conserved (including all eight cysteines) and approximately 44 positions in an alignment of all identified full-length sequences show ≥75% identity (Figure S1). Selection analyses show that approximately 95% of sites evolved under negative to neutral selection (from comparisons of the M2 and M1a models and the M8 and M7 models in the Phylogenetic Analysis by Maximum Likelihood (PAML) package (28)), while only three sites (T38, K46 and K48) evolved under weak positive selection with >90% likelihood based on dN/dS ratios (see Supplementary File 2).

Generally, venom gland transcripts that are highly expressed across multiple species likely encode peptides that function as toxins. Analysis of the expression levels of MKAVA transcripts

revealed that most cone snail species express two transcripts: one with a high level of expression

(100–several thousand transcripts per million (tpm)) and the other with a low level of expression

(<100 tpm) (Figure 1b). Phylogenetic analysis of transcripts with high and low expression revealed

a well-resolved clustering pattern consistent with their relative expression (Figure 1c). The

bipartite nature of the gene tree is reflected at the sequence level, where specific positions

distinguish toxins encoded by either high- or low-expressing transcripts (Figure 1a). The high

expression of many MKAVA conotoxin-encoding transcripts support a function of these as toxins.

This is further corroborated by their presence in the venom. Using publicly available tandem mass

spectrometry data of the venom of *Conus geographus* and *Conus marmoreus*, we identify several

tryptic peptides corresponding to MKAVA sequences from these species (Figure S2).

The MKAVAs are evolutionarily related to a schistosomin

To identify possible homologs of MKAVAs outside of Conoidea, we used a high-expressing

precursor sequence from *C. magus* to perform homology searches against the NCBI non-redundant

protein database and transcriptome shotgun assemblies. Given that we find this toxin from *C.*

magus to display a KPI structure (see below), and that it is the first characterized protein from its

family, we name this toxin "conkazal-M1" (where M stands for *magus*). Likewise, we will use the

term "conkazal", rather than MKAVA, for this family of conotoxins from here on.

The searches revealed sequence similarity, including conservation of all eight cysteines,

with schistosomin-like proteins from several molluscs (Figures 2a and S3), but seemingly absent

in other phyla. Schistosomin was first described in the freshwater pond snail, *Lymnaea stagnalis*,

as a peptide upregulated in response to infection with the schistosome parasite *Trichobilharzia*

ocellata (29) (see also Discussion). Overall, modest sequence conservation is observed across the

conkazals and schistosomins. However, in addition to the cysteines, three residues show high

conservation. These comprise a glutamate residue located two residues after the second cysteine

in both groups of proteins, i.e. at the P1 position of KPIs, at position 34 in the alignment, a proline

immediately following the first Cys residue, and a positively charged residue at position 62 in the

alignment (Figure 2a and S3).

A CLANS (CLuster ANalysis of Sequences) clustering analysis (30), which groups protein

families based on all-against-all pairwise protein sequence similarities, revealed a clear similarity

of the *Conus*-derived conkazal sequences with molluscan schistosomins from gastropods (Figure

2b; all sequences used in this analysis are provided in Supplementary File 3). Moreover, gene

structure analysis revealed a common gene structure of conkazals and molluscan schistosomins.

The genes have a common phase-1 intron located immediately downstream of the signal sequence

and a conserved phase-2 intron located between the sixth and seventh cysteine residues (Figure

2c). The common gene structures further support an evolutionary relationship between conkazal

conotoxins and schistosomins.

While molluscan species exhibit a high degree of intrachromosomal gene translocation,

macrosynteny tends to be conserved. We therefore compared the genomic location of conkazal

genes from *C. ventricosus* with schistosomins from representative gastropods (Figure S4). The two

conkazal genes in *C. ventricosus* are adjacent to each another on scaffold 23, which represents one

of the chromosome-scale scaffolds in the cone snail genome. This chromosome shows homology

to chromosome NC_037594.1 of the gastropod *Pomacea canaliculata* and chromosome

NC_090294.1/ NC_090290.1 of another gastropod *Biomphalaria glabrata* (Figure S4). However,

neither of these chromosomes contain the schistosomin genes, which are located on NC_037595.1

and NC_074716.1, respectively. Surprisingly, both *P. canaliculata* and *B. glabrata* have multiple schistosomin gene copies located in tandem though the neighboring genes show no syntenic relationship (Figure S4b). Overall, while conkazals show a high level of similarity to schistosomins, strongly suggesting an evolutionary relationship, the genes are not part of a common gene linkage group.

Conkazal-M1 can be produced in the fully oxidized form with the csDisCoTune *E. coli* expression platform

To gain insight into the structure and function of the conkazal superfamily, conkazal-M1 from the magician cone, *C. magus*, was selected for recombinant expression and characterization. Conkazal-M1 was recombinantly expressed in *E. coli* using the csDisCoTune system (31). This expression platform, a recent further development of the original CyDisCo system (32) and our own csCyDisCo system (33), is tailored specifically for the expression of conotoxins and allows for the production of correctly folded, fully oxidized disulfide-rich proteins in the bacterial cytosol by co-expression of three foldases encoded by an auxiliary plasmid – the Erv1p oxidase, human protein disulfide isomerase (hPDI), and a conotoxin-specific protein disulfide isomerase (csPDI) (34). Conkazal-M1 was expressed as a fusion protein with an N-terminal ubiquitin (Ub) tag containing an internal loop modified to contain ten consecutive histidine residues (His_{10}) and a tobacco etch virus protease (TEVp) recognition site, allowing for release of the mature toxin (35).

SDS-PAGE analysis of small-scale test expressions of Ub-His_{10}-conkazal-M1 alone, Ub-His_{10}-conkazal-M1 with the csDisCoTune-encoded foldases, or with the foldases encoded by the csCyDisCo expression platform clearly showed the beneficial effect of both expression systems

on Ub-His$_{10}$-conkazal-M1 production. Both systems resulted in the expression of soluble Ub-

His$_{10}$-conkazal-M1 fusion protein, in contrast to expression without the foldases where equivalent

total protein levels were observed, but the fusion protein was predominantly found in the insoluble

fraction in a reduced and/or partially oxidized state (Figure S5a). Because the csDisCoTune

showed higher levels of oxidized Ub-His$_{10}$-conkazal-M1 compared to the csCyDisCo system,

large-scale expression was carried out with csDisCoTune. Cleavage of affinity-purified Ub-His$_{10}$-

conkazal-M1 by TEVp resulted in liberation of the 73-residue mature region of conkazal-M1,

which was purified (>95% purity) to a final yield of just over 5 mg per liter culture (Figure S5b).

Disulfide-bond formation in the purified toxin was verified by SDS-PAGE analysis, where

a clear downward mobility shift was observed upon addition of a reducing agent (Figure S5c), as

well as Q-TOF mass spectrometry, which revealed the presence of two species in the purified

preparation. The primary species displayed a monoisotopic mass of 7990.54 Da in accordance with

the theoretical mass (7990.55 Da) of the fully oxidized protein (Figure S5d). The secondary

component exhibited a monoisotopic mass of 7973.49 Da, i.e., 17 Da lower than the theoretical

mass. We attribute this reduction in mass to modification of the N-terminal Gln to a pyroglutamate,

a post-translational modification commonly found in conotoxins (36). This modification can be

enzymatically catalyzed by glutaminyl cyclase but is also known to occur spontaneously (37).

222 The NMR structure of conkazal-M1 reveals a Kazal-type protease inhibitor fold

The three-dimensional structure of ^{15}N and ^{13}C stable isotope-labelled conkazal-M1 was solved in

solution using distance and torsion angle restraints obtained by NMR spectroscopy. We assigned

the chemical shifts of 79% of the backbone ^{1}H, ^{13}C and ^{15}N resonances and 66% of sidechain

resonances. In total 69% of the chemical shifts were assigned. We observed no resonances from

the first seven residues. 602 short distances assigned in NOESY spectra were combined with 100

dihedral angles calculated from chemical shifts and used as restraints in structure calculations

(Table 1). The structure is well ordered from residue 10 to the C-terminus with an RMSD value of

0.31Å for the positions of the backbone atoms in the 20 models that represent the structure (Figure

S6).

The structure consists of a central three-stranded antiparallel β-sheet and two α-helices; α-

helix 1 that is parallel to the β-strands and α-helix 2 that is almost perpendicular to α-helix 1

(Figure 3a and b). The N- and C-termini are tethered to the core structure via two disulfide bridges.

Located at each end of α-helix 1, Cys40 and Cys48 form disulfide bonds with the N-terminal

Cys14 and Cys4, respectively. The C-terminus is affixed to the short loop connecting β-strand 2

and α-helix 1 by the bridge connecting Cys37 and Cys68. The last disulfide bond is formed by

Cys24 and Cys59, thus connecting β-strand 1 and α-helix 2.

As structural similarity often correlates with conserved functional properties, we used

FoldSeek (38) and PDB eFold (39) to identify proteins structurally similar to conkazal-M1. Both

search engines identified several Kazal-type protease inhibitors among the most similar structures

to conkazal-M1, with the AlphaFold-predicted structure of a schistosomin from *Lymnaea stagnalis*

found as the top hit in the FoldSeek search with conkazal-M1. We also identified an unpublished

crystal structure of schistosomin from another freshwater snail, *Biomphalaria glabrata* (PDB ID:

9FDO), in the PDB. Conkazal-M1 and this schistosomin (Figure 3a) superpose with an RMSD

value of 1.9 Å over 38 Cα-atoms (Figure 3b). Both structures harbor the KPI core structure as well

as an additional C-terminal α-helix not found in classical KPIs. The three characteristic disulfide

bridges of the classical KPIs are present in both conkazal-M1 and schistosomin and are similarly

positioned in the two structures (blue arrowheads, Figure 3b). Both proteins harbor a fourth

disulfide bridge that tethers the C-terminal regions to the core structure of the proteins (Figure 3a).

To illustrate the structural similarity between conkazal-M1 and a verified KPI, Figure S7 shows

an overlay of conkazal-M1 with the crystal structure of greglin (40). Similar to the other structures,

greglin contains a fourth disulfide bond, but its position differs from that in conkazal-M1 and

schistosomin (Figure S7; orange).

Conkazal-M1 inhibits the proteolytic activity of Subtilisin A

Because conkazal-M1 adopts a Kazal fold, we speculated that the toxin may exhibit inhibitory

activity against serine proteases. Consistent with this hypothesis, Glu15 at the P1 position in

conkazal-M1 is solvent exposed in all 20 of the lowest energy structures (Figure S6). We therefore

investigated the proteolytic activities of chymotrypsin, Endoproteinase GluC and Subtilisin A

(SubA) in the presence of increasing amounts (0.25 μM – 3 μM) of conkazal-M1. GluC was chosen

as a possible candidate due to its preferential cleavage of peptide bonds C-terminal to glutamic

acid residues. SubA is a non-specific protease that favors cleavage after hydrophobic residues but

is not restricted to them. For instance, a Glu residue at the P1 position in the second domain of the

CrSPI two-domain Kazal-type protein from horseshoe crab is the primary residue for mediating

the contacts with subtilisin in the co-crystal of the complex between the two proteins (41).

SubA activity was monitored using the chromogenic substrate N-succinyl-Ala-Ala-Pro-

Phe-pNA over a range of substrate concentrations in the presence and absence of conkazal-M1.

Initial rates were estimated from changes in absorbance at 405 nm over time and were converted

to reaction rates (µM/min/mg enzyme). Nonlinear regression of the initial rate data was performed

using Michaelis-Menten and competitive, non-competitive and uncompetitive inhibition models.

Model comparison by Akaike Information Criterion indicated that competitive inhibition best

described the data – as expected for a KPI – with estimated kinetic parameters of Km=201 µM

(Confidence Interval (C.I.): 173‑234 µM), Vmax= 12,888 µM/min/mg (C.I.: 12,357-13,456

µM/min/mg) and Ki=1.1 µM (C.I.: 0.92-1.5 µM) (Figure 4). We performed similar investigations

with chymotrypsin and Endoproteinase GluC in the absence and presence (0.5 µM-2 µM) of

conkazal-M1 where no effect of the toxin on enzyme activity was discerned (Figure S8).

279 Conkazal-M1 modulates calcium transients in mouse sensory neurons

A wide variety of venom peptides function as neurotoxins that target neuronal ion channels.

Therefore, we next assessed the bioactivity of conkazal-M1 using the calcium imaging-based

constellation pharmacology assay (Teichert et al, 2012). This assay enables the monitoring of Ca^{2+}

influx in primary cell cultures obtained from mouse DRGs, which comprise different types of

nociceptors, mechanoreceptors, and glia. During each experiment, intracellular calcium levels

were simultaneously monitored in 500-1,000 cells using the Fura-2-AM dye. Cells were assigned

to seven major classes based on their size, isolectin B4 staining (non-peptidergic nociceptors) and

calcitonin gene-related peptide (CGRP) expression (peptidergic nociceptors) (Fang et al, 2010).

To further characterize neuronal subtypes, a series of pharmacological agents (conotoxin κM-

RIIIJ, allyl isothiocyanate [AITC], menthol, and capsaicin) were applied at the end of each

experiment (Giacobassi et al, 2020).

Dissociated DRG neurons were depolarized using 15 s pulses of high-potassium

extracellular solution (30 mM KCl), resulting in intracellular Ca^{2+} elevation, as indicated by the

rise in Fura-2 signal. At the lowest concentration tested (1 μM), conkazal-M1 did not produce

significant effects on any neuronal subclasses. In contrast, application of 10 μM conkazal-M1

reduced the KCl–evoked calcium signals in a subset of cells (12.2% of the 1,494 cells analyzed)

(Figure 5a). The blockade of calcium transients was predominantly observed in small (<400 $μm^2$)

peptidergic nociceptive neurons (34.1% of affected cells) (identified by the expression of CGRP

genetically labeled with green fluorescent protein) (Figure 5b). Other neuronal subclasses affected

by 10 μM conkazal-M1 included large-diameter (>400 $μm^2$) mechanoreceptors (18.7%), a subset

of non-peptidergic nociceptors (15.9%), and C-low threshold mechanoreceptors (C-LTMR;

14.3%). Thus, conkazal-M1 inhibits KCl-induced calcium influx in a subset of neuronal cells,

albeit with low affinity.

Lastly, because conkazal-M1 originates from the venom of *C. magus*, a fish-hunting cone

snail, we hypothesized that the toxin might elicit measurable effects in a fish model. To test this,

we injected goldfish (*Carassius auratus*) with 5 or 10 nmol of toxin via both intraperitoneal and

intramuscular routes (N=3 for each group) and monitored them closely for a period of three hours

post-injection. No overt phenotypic effects were observed in either treatment group compared with

the controls (data not shown).

DISCUSSION

Here, we provide the first characterization of a toxin from the conkazal conotoxin family, conkazal-M1. The conkazals constitute a large and widespread toxin family found across phylogenetically diverse *Conus* species and several other species in the Conoidean superfamily, indicating an important functional role of these peptides. The common evolutionary history with the schistosomins indicates that an ancestral gene was recruited into the venom in early Conoidea and then acquired neuroactive functions. However, given that the function of schistosomins remains unclear, the exact process of recruitment is presently unknown.

Many conkazal transcripts are highly expressed in the venom gland, and we confirmed their presence in venom by proteomic analysis, supporting a role as toxins. We show that conkazal-M1 can perform dual functions as a protease inhibitor and as a partial inhibitor of KCl-stimulated calcium influx in sensory neurons. Such dual activities have also been reported in Kunitz-type venom proteins (11-14). In the context of constellation pharmacology, the partial suppression of KCl-evoked calcium transients by conkazal-M1 provides preliminary evidence that this peptide can modulate ion channel activity in sensory neurons. Although the effect was observed in only a subset of cells, the preferential targeting of small-diameter, CGRP-positive nociceptors suggests that conkazal-M1 may interact with molecular components enriched in these neurons, such as high-voltage-activated calcium channels (*e.g.*, Cav2.2 or Cav2.3) or auxiliary subunits that shape their kinetics (42-44). The modest inhibition at 10 µM indicates either a low-affinity or indirect mechanism, potentially consistent with dual-function peptides in which ion-channel modulation is a secondary activity to protease inhibition, as further discussed below. Future studies are needed to identify the target and mode of action conkazal-M1 in neuronal cells.

In the context of protease inhibitor function, subtilisin-targeting KPIs with a Glu at the P1

position have previously been characterized (see e.g., (41, 45-47)). It is therefore unsurprising that

conkazal-M1 with its Glu15 at the P1 position was able to inhibit subtilisin (although additional

residues in the reactive-site loop are likely also to contribute to substrate specificity). While certain

venoms appear to harbor microbiomes (48), bacterial subtilisin is an unlikely physiological target

of the protease-inhibitory function of conkazal-M1. This is also reflected in the relatively high K_i

value of conkazal-M1 for subtilisin when compared to many other subtilisin inhibitors, such as

greglin. In principle, in its capacity as a protease inhibitor, conkazal-M1 could perform a function

either in the venom (to regulate the activity of venom proteases to maintain venom stability before

delivery) or in the prey to exert a deleterious effect upon delivery. At present, we cannot distinguish

between these possibilities although we note that we have not been able to find evidence in the

literature for an endogenous regulatory function of protease inhibitors in other venomous

organisms.

When comparing conkazal sequences with schistosomins from other molluscs, Glu15 at

the P1 position is one of the few residues conserved across species (Figure 2). This observation

supports to the hypothesis that the conkazals and mollusk schistosomins function as KPIs. In

addition to the eight cysteine residues, only two other residues are as highly conserved as Glu15:

a proline residue immediately following the first Cys residue, likely playing a structural role, and

a surface-exposed lysine residue at a position corresponding to Lys43 in the mature conkazal-M1

sequence, which is most likely to have a functional role.

Schistosomiasis, a parasitic infection, is caused by flatworms known as schistosomes. In

the schistosomiasis life cycle, freshwater snails serve as intermediate hosts that release infectious

larvae into the water. These larvae can penetrate human skin, gaining access to the host, where the

subsequent release of eggs can cause a range of symptoms, including serious damage to organs

such as the liver and bladder. In the snail-schistosome system *L. stagnalis-T. ocellata*, schistosomin

was initially proposed to be a neuropeptide induced by parasite infection, but subsequent studies

showed that it is produced in connective tissue cells and in hemocytes (reviewed in (49)). In the

hemolymph, schistosomin was proposed to act as an antagonist of gonadotropic hormones thereby

inhibiting reproductive activity in snails. However, in the *Biomphalaria glabrata-Schistosoma*

mansoni system, no correlation was found between parasite infection and schistosomin levels and

infection did not result in parasitic castration (50). Thus, the physiological function of schistosomin

remains unclear, and further investigation is warranted, particularly with its potential function as

a KPI in mind.

An ancestral protease inhibitor that acquires the ability to modulate neuronal activity while

retaining protease inhibition represents a functional change in which purifying selection preserves

the original function, while weak-to-moderate selection can favor the emergence of a novel activity

(6, 51). In contrast, a complete shift toward modulation of neuronal activity accompanied by partial

or total loss of protease activity can be explained by strong positive selection for the new function,

coupled with relaxed constraint on the ancestral activity. Because we detect no evidence of positive

selection, we favor the first evolutionary model. In this scenario, conkazals would be

"moonlighting" proteins that face an adaptive conflict when mutations that enhance one activity

compromise the other. Such conflicts are typically resolved through gene duplication followed by

subfunctionalization that lead to partitioning of the ancestral roles between paralogs or to

neofunctionalization of one copy. Our data do not support a neofunctionalization event. If such an

event had occurred, we would expect to detect both an ancestral, schistosomin-like gene and a

duplicated conkazal gene. Instead, we only observe conkazal genes when interrogating available

cone snail genome data, consistent with a moonlighting role. This interpretation is further supported by the expression of conkazals in tissues other than the venom gland, including the venom bulb, nerve ring, and foot (see Materials and Methods section for SRA entry numbers), where these peptides likely retain an endophysiological, ancestral function.

We note that multi-domain Kazal-type proteins are found in many venomous animals, including the two-domain Kazal-type toxins BQTX from shrews (19) and Rhodniin from the kissing bug *Rhodnius prolixus* (17, 52). In general, secreted cysteine-rich repeat-proteins (53) – including a broad range of venom peptides – are common in nature and have been cataloged in the ScrepYard database (54). Their multi-domain structure enables multivalent substrate interactions, allowing them to bind targets strongly through an avidity effect. Indeed, Rhodniin binds its substrate thrombin through multivalent interactions involving the reactive-site loop of one domain and electrostatic interactions of the second domain, together conferring a picomolar affinity (17, 52). Analogous to the two-domain Kazal-type toxins described above, we have recently identified a family of such conotoxins, the MKIAV family. As shown in Figure S9a-b, both Kazal domains of these conotoxins, with few exceptions, contain a Glu or Asp residue three positions C-terminal to the second Cys in the mature sequence. Moreover, each MKIAV protein has a Glu/Asp at this position in one of the two kazal-type domains. Although not located at the canonical P1 position, we speculate that this residue could still perform the inhibitory function of a P1 residue. The AlphaFold 3 predictions of the MKIAVs reveal two very similar Kazal-type domains connected by a short linker, with each domain closely resembling the structure of conkazal-M1 (Figure S9c-e).

In this study, we identify two previously unrecognized families of conotoxins, conkazal and MKIAV, which encode single- and two-domain Kazal-type domain structures, respectively.

402   We uncover the dual function of conkazal-M1 as a neuroactive peptide with the ability to inhibit

403   calcium transients in a subset of DRGs and as a protease inhibitor. Together, these observations

404   highlight the structural and functional diversity of Kazal-type toxins in *Conus*, extend the current

405   knowledge about the toxin complement available to venomous animals, and illustrate the adaptive

406   conflict that can shape the evolutionary trajectories of dual function toxins.

MATERIALS AND METHODS

Sequence mining and analysis

Multiple sequence alignments were generated with the MAFFT version 7 (55) online interface using the G-INSi method (56). Alignments were visualized in JalView (57). Schistosomin sequences were obtained using web version of blast to query against the non-redundant (nr) protein database and transcriptome shotgun assemblies with standard settings. Conotoxin sequences were obtained from (25). These sequences were used as input for CLANS (30), which is a sequence-based clustering program that assigns attractive values between sequences based on their BLOSUM62 blastp p-value. In a uniformly repelling force field this causes similar sequences to cluster together. Gene structure analyses were performed with using the protein2genome (p2g) model in Exonerate (v.2.76.2) (58). Macrosynteny analysis was performed with macrosyntR using OrthoFinder to assign orthologous proteins across *C. ventricosus* (GCA_018398815.2)*, P. canaliculata* (GCF_003073045.1)*,* and *B. glabrata* (GCF_947242115.1). The local gene neighborhoods were assessed and visualized in Figure 2C. Maximum likelihood gene trees were constructed using alignments generated with MAFFT v.7.526 and the tree was constructed using IQ-TREE v. 2.2.2.3 (59) with UF bootstrap (60) calculation for 1000 replicates. The best model according to the Bayesian Information Criterion was calculated to be TMP3 + G4. The consensus tree was visualized using FigTree (61). Conkazal-like sequences were identified in the NCBI SRA database using the conkazal-M1 amino acid sequence as a query sequence in tblastn searches that retrieved the following SRA entries: SRX8002177, SRR11423812 (Drillidae), SRR26897535, SRR26897526 (Turridae), SRR10192879, SRR10192877 (Profundiconus), SRR2060989 (Terebridae), SRR16493587 (venom bulb), SRR16493599, SRR16493598 (nerve ring), and SRR16493591 (foot).

Proteomics-based identification of MKAVA peptides in venom

Proteomics files of reduced, alkylated and trypsin digested venom of *Conus geographus* (accession: PXD052869) and *Conus marmoreus* (PXD038992) were retrieved from the PRIDE ProteomeXChange repository and searched for MKAVA sequences using the Byonic software (version 3.2.0 Protein metrics) with the following settings: cleavage site: RK, missed cleavage: 1, Modifications: carbamidomethyl (C, +57.02146), Oxidation (M,P, +15.994915), pyro-Glu (Q, +17.026549), Carboxy (E, +43.989829). High scoring peptide hits (score> 400) were manually verified.

Plasmid generation

The codon-optimized sequence encoding mature conkazal-M1 was synthesized and inserted (TWIST Bioscience) into the pET39_Ub19 expression vector (35). The resulting plasmid (pLE657) encodes ubiquitin (Ub)-His$_{10}$-tagged conkazal-M1 (Ub-His$_{10}$-conkazal-M1) with a recognition site for TEV protease located immediately N-terminally to the mature toxin sequence.

Protein expression and preparation of lysate

Chemically competent BL21(DE3) were co-transformed with pLE657 and the csDisCoTune expression plasmid (pLE900) or, for comparison, alone or together with the csCyDisCo plasmid (pLE577). Cells were plated on lysogeny broth (LB) agar supplemented with kanamycin (50 µg/mL) and – when co-transforming with pLE900/pLE577 – chloramphenicol (30 µg/mL). LB medium containing the same type and concentration of antibiotic as in the transformation plates

was inoculated with a single colony and incubated for at 37° C for 18 hours at 190 rpm on an orbital shaker.

Initial small-scale expression tests were performed in 50 mL LB medium containing appropriate antibiotics and supplemented with 0.05% glucose. Test cultures were inoculated with 2% overnight culture and grown at 37° C, 190 rpm until OD_{600} reached ~0.8. Expression was induced by addition of isopropyl -D-1-thiogalactopyranoside (IPTG) to a final concentration of 1 mM and the cultures grown for 18 hours at 25° C with shaking at 190 rpm to allow protein expression. Large-scale expression was performed similarly in 1 L LB medium. Expression performed in 0.5 L M9 minimal medium for stable isotope labelling was performed as described in (62).

Induced cultures were harvested by centrifugation at 4,000 x g for 20 minutes at 4° C. The cell pellets were resuspended in 20 mL lysis buffer (50 mM Tris (pH 8.0), 300 mM NaCl, 20 mM imidazole) per liter culture. Cell lysis was performed using a UP200S ultrasonic processor (Hielscher) while keeping the cells on ice. The cells were lysed with 8 x 30-second pulses at 90% power with 30-second rests between each pulse. Cell debris was subsequently pelleted by centrifugation at 30,000 x g for 45 minutes at 4° C. The cleared supernatant was reserved, and cell pellets were resuspended in an equal volume lysis buffer containing 8 M urea for SDS-PAGE analysis.

Protein purification

Ub-His$_{10}$-conkazal-M1 was affinity purified from the clarified lysate on an ÄKTA START system equipped with a 5-mL prepacked HisTrap HP (Cytiva) column equilibrated in lysis buffer. The

lysate was applied to the column and washed with 30 column volumes (CVs) of lysis buffer before

elution of Ub-His$_{10}$-conkazal-M1 with a gradient of 0% to 100% elution buffer (50 mM Tris (pH

8.0), 300 mM NaCl, 400 mM imidazole) developed over 20 CVs. Pooled fractions were dialyzed

twice against 2 L cleavage buffer (50 mM Tris (pH 8.0), 300 mM NaCl).

The Ub-His10-conkazal-M1 fusion protein was cleaved using His$_6$-tagged TEVp,

expressed, and purified essentially as described previously (27). A molar ratio Ub-His$_{10}$-conkazal-

M1: His$_6$-TEVp of 1:2 was used. To avoid reducing the disulfides in conkazal-M1, His$_6$-TEVp –

preactivated with 2 mM dithiothreitol (DTT) for 30 minutes at room temperature – was diluted to

approximately 0.002 mM DTT by 3 rounds of dilution/concentration in an Amicon Ultra 15 mL

10K Centrifugal Filter (Merck Millipore). TEVp cleavage was performed overnight at room

temperature.

To remove uncleaved Ub-His$_{10}$-conkazal-M1, the Ub-His$_{10}$ tag, and His$_6$-TEVp, the

cleavage mixture was applied to a 5 mL prepacked HisTrapHP (Cytiva) affixed an ÄKTA START

equilibrated in cleavage buffer. The flow-through and the first wash fraction were collected. The

presence of cleaved conkazal-M1 in the flow-through and wash fractions was investigated by

analysis on a 15% SDS-PAGE gel, and the protein-containing fractions pooled. Due to minor

remaining impurities, the cleaved conkazal-M1 was subjected to size exclusion chromatography

on a Superdex 75 Increase 10/300 GL column (Cytiva) equilibrated in 50 mM Tris (pH 7.8), 150

mM NaCl. Fractions containing purified conkazal-M1 were pooled and stored at 4 C.

SDS-PAGE analysis

Samples from bacterial expression and subsequent purification steps were separated on 15% glycine SDS-PAGE gels and where indicated samples were treated with 40 mM DTT. Protein bands were visualized with Coomassie brilliant blue stain and images were captured with a BioRad Chemidoc Imaging System.

Protein concentration determination

Concentration was determined by measuring absorbance at 280 nm and using the theoretical extinction coefficient provided by the Expasy ProtParam tool available through the Expasy bioinformatics resource web portal (63).

Mass spectrometry

Mass spectrometry was performed on a Waters Xevo G3 QTof system equipped with a ACQUITY UPLC C4 reversed-phase liquid chromatography column (300 Å, 1.7 μm, 2.1 mm × 50 mm) using a sample of purified conkazal-M1 at 100 μM adjusted to pH 2 with trifluoroacetic acid as input.

NMR spectroscopy

A sample of 500 μM ^{13}C,^{15}N-labelled conkazal-M1 was prepared in 10 mM 2-(N-morpholino)ethanesulfonic acid (MES), 200 mM NaCl, 5% D_2O, pH 6.0. For backbone chemical shift assignments ^{15}N-HSQC, HNCA, HN(CO)CA, CBCA(CO)NH, CBCANH, CBCA(CO)NH, HNCO, and HN(CA)CO spectra were recorded on a Bruker Avance III HD spectrometer operating at 750 MHz with a triple resonance cryoprobe. Side chain assignments were achieved from

HCCONH and HCCH-TOCSY spectra. Distance restraints were obtained from ^{15}N-NOESY-

HSQC and ^{13}C-NOESY-HSQC experiments recorded using a mixing time of 120 ms. The triple

resonance spectra were recorded with non-uniform sampling at 25% and reconstructed with

qMDD (64). All spectra were processed with nmrPipe (65) and analyzed in CCPNMR analysis

(66).

NMR structure calculations

Automated NOE assignment was performed using Cyana (67). The NOE list was manually refined.

XPLOR-NIH (68) was used for the final structural refinement including a torsion angle database

potential (69) and an implicit solvent mode (70). For the final ensemble representing the solution

structure of conkazal-M1, the 20 structures with the lowest energy were chosen from 100

calculated structures. The Ramachandran-plot statistics was calculated using PROCHECK (71)

and the coordinate precisions were calculated by PyMOL (DeLano Scientific). Structure

visualization was also performed with PyMOL. The structure has been deposited in the Protein

Data Bank (PDB, wwpdb.org) with PDB code 9SLR and chemical shifts and peaks lists from the

NOESY spectra have been deposited in the Biological Magnetic Resonance Bank

(http://www.bmrb.wisc.edu/) with ID 53313.

Structural homology search

Structures similar to conkazal-M1 were identified using FoldSeek (38), and structural

superpositions were executed manually in PyMOL (DeLano Scientific) utilizing the CEalign

command.

Protease assays

Endoproteinase GluC, bovine pancreas α-chymotrypsin and subtilisin A and substrates for chymotrypsin (N-Succinyl-Gly-Gly-Phe-*p*-nitroanilide) and subtilisin A (N-succinyl-Ala-Ala-Pro-Phe-p-nitroanilide) were all purchased from Merck. Substrate for GluC (Z-Phe-Leu-Glu-pNA) was purchased from Bachem.

Inhibition assays with chymotrypsin and GluC were performed in assay buffer (50 mM Tris, 2 mM CaCl$_2$, pH 7.8). Subtilisin A inhibition experiments were performed in assay buffer, pH 8 supplemented with 5 mM CaCl$_2$. The proteases (3.6 nM SubA, 100 nM GluC and 400 nM chymotrypsin) were incubated with varying concentrations of conkazal-M1 for 10 min at 29° C. Proteolytic reactions were initiated by addition of the substrates and absorbance at 405 nm was recorded every 30 seconds for 30 minutes. The 96-well plate was shaken orbitally for 10 seconds immediately prior to measurement. Initial reaction rates were estimated with the web-based tool, IceKat (72). The absorbance-based rates for Subtilisin A were converted to reaction rates in µM/min/mg using the extinction coefficient for AAPF-p-nitroaniline ($\varepsilon = 8800$ M^{-1}cm^{-1}) and an estimated path length of 0.2 cm for 100 µL reaction volumes in 96-well plates. The absorbance-based rates (Abs/min) for Chymotrypsin and GluC were plotted without conversion. The data were fit to four models of enzyme inhibition: Michaelis-Menten (no inhibitor), competitive, uncompetitive, and noncompetitive inhibition.

$$v = \frac{V_{max}[S]}{K_m + [S]}$$

561
$$v = \frac{V_{max}[S]}{K_m\left(1 + \frac{[I]}{K_i}\right) + [S]}$$

563
$$v = \frac{V_{max}[S]}{K_m + [S]\left(1 + \frac{[I]}{K_i}\right)}$$

565
$$v = \frac{V_{max}[S]}{K_m\left(1 + \frac{[I]}{K_i}\right) + [S]\left(1 + \frac{[I]}{K_i}\right)}$$

566   Model comparison using Akaike Information Criterion identified the best fit to the data.

Cell culture and constellation pharmacology

Experiments were performed as previously described (73). Briefly, all lumbar DRG neurons from

C57BL/6 mice 30-40 days old of both genders were dissociated by trypsin digestion and

mechanical trituration and plated on polylysine-coated plates. The cells were incubated overnight

in a 5% CO_2 incubator at 37 °C in neuronal culture medium [minimal essential media

supplemented with 10% (vol/vol) fetal bovine serum, penicillin (100 U/mL), streptomycin (100

µg/mL), 1× Glutamax, 10 mM HEPES, and 0.4% (wt/vol) glucose, adjusted to pH 7.4]. One hour

before calcium imaging experiments, cells were loaded with 2.5 µM fura-2-acetoxymethyl ester

(Fura-2-AM, Sigma-Aldrich) and incubated at 37 °C. Cells loaded with Fura-2-AM dye were

excited intermittently with 340- and 380-nm light; fluorescence emission was monitored at 510

nm. An image was captured at each excitation wavelength, and the ratio of fluorescence intensities

(340/380 nm) was acquired every 2 s to monitor the relative changes in intracellular calcium

concentration in each cell as a function of time. Typically, 500-1,000 neurons were imaged for

each experiment. The calcium transients were elicited by 15 s applications of depolarizing stimulus

(30 mM KCl), followed by four washes with extracellular solution [145 mM NaCl, 5 mM KCl, 2

mM CaCl2, 1 mM MgCl2, 1 mM sodium citrate, 10 mM HEPES, and 10 mM glucose, adjusted to

pH 7.4]. For activity screening, conkazal-M1 was applied in extracellular solution at 1 and 10 μM

between KCl pulses and activity was assessed by significant change in the KCl response after

incubation with conkazal-M1. Four different pharmacological agents were used for cell

classification: conotoxin κM-RIIIJ (1 μM), AITC (100 μM), menthol (400 μM) and capsaicin (300

nM). At the end of the experiment, cells were incubated with 0.7 mL of Hoechst stain (1000 μg/mL)

and Alexa-Fluor 647 Isolectin B4 (2.5 μg/mL) for 5 min. The data was acquired using NIS-

Elements and further processed with CellProfiler v.3.1.9 (Jones et al, 2008). Custom-built scripts

in Python v.3.7.2 and R were used for further data analysis and visualization. The experimental

methods were approved by the Institutional Animal Care and Use Committee (IACUC) of the

University of Utah (Protocol number: 17–05017).

Statistical analysis of calcium imaging data

Each experiment had a series of KCl-induced responses. Their magnitude was measured as the

maximum area under the curve for any 15 s interval. The deviation of this response after incubation

with conkazal-M1 was used to estimate the effect of conkazal-M1. We estimated this effect on

each neuron using the linear model (lm function in R) with the following:

lm(Kauc15 ~ linear + toxin)

where Kauc15 was the maximum area under the curve for any 15 s interval for each KCl pulse,

linear was the sequential trend coded as sequential integers, toxin was an indicator variable for the

pulse immediately following incubation with conkazal-M1.

To control for false positives and set appropriate thresholds for significance we used Monte

Carlo simulations. Each simulation generated random numbers drawn from a normal distribution

with mean and standard deviations of the actual data. The T-stats were estimated for all cells and

recorded. This is considered a null distribution. The T-stat estimates from 100 Monte Carlo

simulations were used to establish the thresholds for single test cell significance as well as per-

experiment significance.

We used the relative residual of the affected KCl pulse in the model as a ratio of the

observed to the model predicted magnitude of effect. This was calculated as:

(KPp-KPobs)/KPp,

where KPp was the predicted peak height of the pulse immediately after conkazal-M1 application

and KPobs was the observed KCl pulse.

Fish behavioral assays

All animal assays were conducted in accordance with the *Guide for the Care and Use of*

Laboratory Animals and were approved by the Institutional Animal Care and Use Committee

(IACUC) of the University of Utah (protocol #001629). Goldfish (*Carassius auratus*) of both

sexes (weight= 0.85 ± 0.11 g, mean $\pm$ SD) were obtained from a local supplier and used for the

experiments. An in-house prepared E3 Buffer (4.96 mM NaCl, 0.18 mM KCl, 0.33 mM

$CaCl_2 \cdot 2H_2O$, 0.4 mM $MgCl_2 \cdot 6H_2O$; pH 7.2) served as the vehicle. Groups of three fish each

623 received injections of either 13.5 µL E3 buffer alone (control) or E3 buffer containing the desired

624 concentration of conkazal-M1.

625  All injections were performed with a 1 mL insulin syringe (31 G). For intraperitoneal

626 injections, fish were positioned ventral side up on a chilled sponge embedded with E3 Buffer under

627 a dissecting microscope, and the solution was delivered into the abdominal cavity. For

628 intramuscular injections, fish were positioned laterally on a hard surface moistened with E3 buffer

629 under a dissecting microscope, and the solution was administered into the lateral muscle below the

630 posterior end of the dorsal fin. Following injection, all fish were maintained under observation for

631 three hours.

 TABLES

 Table 1. Statistics for the ensemble of 20 conformers representing conkazal-M1

Experimental restraints	
NOE based restraints	
Intraresidual (|i - j| = 0)	130
Sequential (|i - j| = 1)	183
Medium range (2 ≤ |i - j| ≤ 4)	92
Long range (|i - j| ≥ 5)	197
Total	602
Dihedral angles	100
Disulfide bonds	4
Restraint violation statistics[a]	
NOE-based distances (Å)	0.046 ± 0.004
Dihedral angles (°)	1.04 ± 0.09
Ramachandran plot statistics[b]	
Most favored regions	83.0%
Allowed regions	15.3%
Generously allowed regions	1.5%
Disallowed regions	0.3%
Coordinate precision (RMSD to the average)[c]	
Residues 10 - 73	
Backbone N, C$^\alpha$ and C'	0.31 ± 0.07 Å
All heavy atoms	2.4 ± 0.6 Å

 [a]XPLOR-NIH, [b]PROCHECK, [c]PyMOL

ABBREVIATIONS

CLANS, CLuster ANalysis of Sequences); csPDI, conotoxin-specific PDI; DRG: dorsal root ganglion, DTT, dithiothreitol; ER, endoplasmic reticulum; HEPES, 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid; hPDI, human PDI; IMAC, immobilized metal affinity chromatography; IPTG, isopropyl ß-D-1-thiogalactopyranoside; KPI: Kazal-type protease inhibitor, LB, lysogeny broth: MSA, multiple sequence alignment; NCBI, National Center for Biotechnology Information; PDB, Protein Data Bank; PDI, protein disulfide isomerase; RMSD, root mean square deviation; SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis; SRA, sequence read archive; TEVp, tobacco etch virus protease; TPM, transcripts per million; Ub, ubiquitin; Ub-His$_{10}$, Ub containing 10 consecutive histidines.

ACKNOWLEDGMENTS

We thank Signe A. Sjørup, SBiNLab, University of Copenhagen, for recording mass spectra.

FUNDING

This work was supported by the Independent Research Fund Denmark (10.46540/3103-00126B to L.E.), the Lundbeck Foundation (R500-2024-1904 to L.E.), the Novo Nordisk Foundation (NNF18OC0032996 to K.T.), and a National Institutes of Health grant (GM144719 to B.M.O. and H.S.-H). T.L.K. was supported by an international postdoc fellowship from the Independent Research Fund Denmark (3102-00006B).

AUTHOR CONTRIBUTIONS

Conceptualization, C.M.H., T.L.K., H.S.-H., and L.E.; Data curation, C.M.H., T.L.K., A.R., K.C.,

M.W., K.T., and L.E.; Formal Analysis, C.M.H., T.L.K., A.R., K.C., and K.T.; Funding acquisition,

T.L.K., B.O., H.S.-H., K.T., and L.E.; Investigation, C.M.H., T.L.K., N.L.R., M.W., A.R., K.T. and

L.E.; Project administration, L.E.; Supervision, B.O., H.S.-H., and L.E.; Visualization, C.M.H.,

T.L.K., A.R., K.T., and L.E.; writing—original draft preparation, C.M.H., T.L.K., A.R., H.S.-H.,

K.T. and L.E.; writing—review and editing, all authors. All authors have read and agreed to the

published version of the manuscript.

COMPETING INTERESTS

The authors declare that no competing interests exist.

FOOTNOTES

The atomic coordinates for conkazal-M1 are available with the Protein Data Bank under accession

number 9SLR and in the Biological Magnetic Resonance Data Bank under ID number 53313.

FIGURES

Figure 1. The MKAVAs are encoded by either low or high expressing transcripts. (a) Multiple

sequence alignment of select *Conus* precursor sequences from the MKAVA superfamily.

Sequences encoded by high or low expressing transcripts are indicated in dark teal and orange,

respectively. The signal sequence and mature toxin regions are identified by light and dark teal

bars above the alignment. The prey preference of each species (i.e. piscivorous, molluscivorous or

vermivorous) is indicated by the icons on the left. **(b)** Gene expression levels of 85 MKAVA

transcripts from 49 cone snail species. Dark teal and orange circles indicate high (>100 tpm) and

low (<100 tpm) expressors, respectively, with light teal bars showing the average tpm-value of the

transcripts for each species. **(c)** Phylogenetic reconstruction of the MKAVA transcripts from (b).

Maximum-likelihood was performed using IQ-Tree's UF bootstrap with 1000 replicates with best

687   fit model TMP3 + G4 selected by the Bayesian Information Criterion. High-expressing transcripts

688   are shown in dark teal and low-expressing in orange. Some bootstrap values were removed for

689   visualization purposes but can be seen in Figure S10. The tree was rooted to separate the two

690   paralogous genes.

692

Figure 2. The MKAVAs are evolutionarily related to schistosomin. (a) Multiple sequence alignment of MKAVAs and schistosomin-like sequences from *Aplysia californica* (XP_005098364.1)*, Pomacea cataticulata* (PVD28780.1)*, Biomphilaria glabrata* (NP_001298209.1)*, Rapana venosa* (GDIA01100366.1) and *Crassostrea gigas* (GHAT01224603.1). **(b)** CLANS clustering analysis of conotoxins and schistosomins. Deep teal: conotoxins from MKAVA superfamily; pink: molluscan schistosomin. Nodes depict individual precursor sequences, and the edges correspond to BLOSUM62 blastp p-values. **(c)** Gene structure analysis of toxins with high or low expression from *Conus ventricosus*, and of schistosomins from *A. californica* and *P. cataticulata*. The exons are depicted as boxes proportional to the length of the sequences, while introns are represented as thin lines (with sequence length labelled). The intron phases (0, 1, or 2) are indicated prior to the introns. The yellow lines indicate cysteine positions within the exons.

Figure 3. Conkazal-M1 harbors a similar fold to schistosomin. (a). Graphic representations of Conkazal-M1 (teal) and schistosomin from *Biomphalaria glabrata* (pink). α-helices and β-strands are represented by cylinders and block arrows, respectively, and disulfide bridges are represented as black brackets. **(b).** Cartoon representation of the conkazal-M1 NMR structure (PDB: 9SLR) (left), cartoon representation of the *B. glabrata* schistosomin structure solved by X-ray crystallography (PDB: 9FDO) (middle), and superposition of conkazal-M1 and schistosomin (right). Coloration as described in Panel (a). Disulfide bridges are colored yellow and the amino acid residue at the P1 position is shown in red. The RMSD value is derived from 38 α-carbons and determined using the PyMOL super command.

Figure 4. Conkazal-M1 inhibits proteolytic activity of Subtilisin A. Kinetic analysis of

Subtilisin A in the presence of increasing concentrations Conkazal-M1. Reaction rates were

measured across a range of substrate concentrations and were normalized to enzyme mass. The

plots display experimental data points (colored according to inhibitor concentration: 0 μM

(orange), 0.25 μM (pink), 0.5 μM (purple), 1 μM (light teal), 2 μM (cyan) and 3 μM (dark teal))

overlaid with lines showing showing the best-fit model (competitive inhibition).

Figure 5. Effects of conkazal-M1 on mouse sensory neurons by constellation pharmacology.

(**A**) Representative calcium-imaging traces of individual neurons responding with a partial block to applications of 10 µM conkazal-M1 (top 7 traces, conkazal-M1 application highlighted by blue box), and neurons that did not respond (bottom 4 traces). Light grey shading indicates KCl (30 mM) depolarization pulses. Each trace represents the calcium signal (ΔF/F) of the neuron pictured on the left (GFP: CGRP expression; IB4: isolectin B4 staining; BF: brightfield image; 10x magnification). Pharmacological agents used for cell classification: RIIIJ (1 µM, burgundy box); AITC (100 µM), menthol (400 µM, Men), and capsaicin (300 nM, Cap). A pulse of 40 mM KCl (K40) was used to assess cell viability at the end of the experiment. Cell size is indicated on the left side of each trace. (**B**) Bar graph showing the percentage of each class of conkazal-M1-sensitive DRG neurons. The pie chart represents the population of neurons analyzed (n = 1,494), highlighting the 12.2% of conkazal-M1 (10 µM)-sensitive sensory neurons. AITC: allyl isothiocyanate; Aδ-LTMRs: Aδ–low-threshold mechanoreceptors; CGRP: calcitonin-gene related peptide; C-LTMR: C-low threshold mechanoreceptors; GFP: green fluorescent protein; N16: capsaicin-positive small-diameter neurons; RIIIJ: conotoxin κM-RIIIJ.

REFERENCES

1.   Casewell, N. R., Wuster, W., Vonk, F. J., Harrison, R. A., and Fry, B. G. (2013) Complex cocktails: the evolutionary novelty of venoms *Trends Ecol Evol* **28**, 219-229

2.   Lewis, R. J., and Garcia, M. L. (2003) Therapeutic potential of venom peptides *Nat Rev Drug Discov* **2**, 790-802

3.   Sachetto, A. T. A., and Mackman, N. (2019) Modulation of the mammalian coagulation system by venoms and other proteins from snakes, arthropods, nematodes and insects *Thromb Res* **178**, 145-154

4.   Woodward, S. R., Cruz, L. J., Olivera, B. M., and Hillyard, D. R. (1990) Constant and hypervariable regions in conotoxin propeptides *The EMBO journal* **9**, 1015-1020

5.   Sollod, B. L., Wilson, D., Zhaxybayeva, O., Gogarten, J. P., Drinkwater, R., and King, G. F. (2005) Were arachnids the first to use combinatorial peptide libraries? *Peptides* **26**, 131-139

6.   Jackson, T. N. W., and Koludarov, I. (2020) How the Toxin got its Toxicity *Front Pharmacol* **11**, 574925

7.   Bayrhuber, M., Vijayan, V., Ferber, M., Graf, R., Korukottu, J., Imperial, J. *et al.* (2005) Conkunitzin-S1 is the first member of a new Kunitz-type neurotoxin family: structural and functional characterization *J Biol Chem* **280**, 23766-23770

8.   Finol-Urdaneta, R. K., Remedi, M. S., Raasch, W., Becker, S., Clark, R. B., Struver, N. *et al.* (2012) Block of Kv1.7 potassium currents increases glucose-stimulated insulin secretion *EMBO molecular medicine* **4**, 424-434

9.   Droctove, L., Ciolek, J., Mendre, C., Chorfa, A., Huerta, P., Carvalho, C. *et al.* (2022) A

new Kunitz-type snake toxin family associated with an original mode of interaction with

the vasopressin 2 receptor *Br J Pharmacol* **179**, 3470-3481

10.   Schweitz, H., Heurteaux, C., Bois, P., Moinier, D., Romey, G., and Lazdunski, M. (1994)

Calcicludine, a venom peptide of the Kunitz-type protease inhibitor family, is a potent

blocker of high-threshold Ca2+ channels with a high affinity for L-type channels in

cerebellar granule neurons *Proc Natl Acad Sci USA* **91**, 878-882

11.   Mishra, M. (2020) Evolutionary Aspects of the Structural Convergence and Functional

Diversification of Kunitz-Domain Inhibitors *J Mol Evol* **88**, 537-548

12.   Yuan, C. H., He, Q. Y., Peng, K., Diao, J. B., Jiang, L. P., Tang, X. *et al.* (2008) Discovery

of a distinct superfamily of Kunitz-type toxin (KTT) from tarantulas *PloS one* **3**, e3414

13.   Gladkikh, I., Peigneur, S., Sintsova, O., Lopes Pinheiro-Junior, E., Klimovich, A.,

Menshov, A. *et al.* (2020) Kunitz-Type Peptides from the Sea Anemone Heteractis crispa

Demonstrate Potassium Channel Blocking and Anti-Inflammatory Activities *Biomedicines*

8,

14.   Peigneur, S., Billen, B., Derua, R., Waelkens, E., Debaveye, S., Beress, L. *et al.* (2011) A

bifunctional sea anemone peptide with Kunitz type protease and potassium channel

inhibiting properties *Biochem Pharmacol* **82**, 81-90

15.   Zupunski, V., and Kordis, D. (2016) Strong and widespread action of site-specific positive

selection in the snake venom Kunitz/BPTI protein family *Sci Rep* **6**, 37054

16.   Zdenek, C. N., Cardoso, F. C., Robinson, S. D., Mercedes, R. S., Raidjoe, E. R., Hernandez-

Vargas, M. J. *et al.* (2024) Venom exaptation and adaptation during the trophic switch to

blood-feeding by kissing bugs *iScience* **27**, 110723

17.   Friedrich, T., Kroger, B., Bialojan, S., Lemaire, H. G., Hoffken, H. W., Reuschenbach, P.

790   *et al.* (1993) A Kazal-type inhibitor with thrombin specificity from Rhodnius prolixus *J*

791   *Biol Chem* **268**, 16216-16222

792   18.   Kim, B. Y., Lee, K. S., Zou, F. M., Wan, H., Choi, Y. S., Yoon, H. J. *et al.* (2013)

793   Antimicrobial activity of a honeybee (Apis cerana) venom Kazal-type serine protease

794   inhibitor *Toxicon* **76**, 110-117

795   19.   Liao, Z., Tang, X., Chen, W., Jiang, X., Chen, Z., He, K. *et al.* (2022) Shrew's venom

796   quickly causes circulation disorder, analgesia and hypokinesia *Cell Mol Life Sci* **79**, 35

797   20.   Qian, C., Fang, Q., Wang, L., and Ye, G. Y. (2015) Molecular Cloning and Functional

798   Studies of Two Kazal-Type Serine Protease Inhibitors Specifically Expressed by Nasonia

799   vitripennis Venom Apparatus *Toxins* **7**, 2888-2905

800   21.   Wang, Q. W., Zou, W. B., Masson, E., Ferec, C., Liao, Z., and Chen, J. M. (2025) Genetics

801   and clinical implications of SPINK1 in the pancreatitis continuum and pancreatic cancer

802   *Hum Genomics* **19**, 32

803   22.   Brillard-Bourdet, M., Hamdaoui, A., Hajjar, E., Boudier, C., Reuter, N., Ehret-Sabatier, L.

804   *et al.* (2006) A novel locust (Schistocerca gregaria) serine protease inhibitor with a high

805   affinity for neutrophil elastase *Biochem J* **400**, 467-476

806   23.   Rimphanitchayakit, V., and Tassanakajon, A. (2010) Structure and function of invertebrate

807   Kazal-type serine proteinase inhibitors *Dev Comp Immunol* **34**, 377-386

808   24.   Safavi-Hemami, H., Foged, M. M., and Ellgaard, L. (2018) Evolutionary Adaptations to

809   Cysteine-rich Peptide Folding In *Oxidative Folding of Proteins: Basic Principles, Cellular*

810   *Regulation and Engineering*, Feige M, ed. Royal Society of Chemistry, 99-128

811   25.   Koch, T. L., Robinson, S. D., Salcedo, P. F., Chase, K., Biggs, J., Fedosov, A. E. *et al.*

812   (2024) Prey Shifts Drive Venom Evolution in Cone Snails *Mol Biol Evol* **41**,

813   26.   Robinson, S. D., Safavi-Hemami, H., McIntosh, L. D., Purcell, A. W., Norton, R. S., and

814   Papenfuss, A. T. (2014) Diversity of conotoxin gene superfamilies in the venomous snail,

815   Conus victoriae *PloS one* **9**, e87648

816   27.   Hackney, C. M., Florez Salcedo, P., Müller, E., Koch, T. L., Kjelgaard, L. D., Watkins, M.

817   *et al.* (2023) A previously unrecognized superfamily of macro-conotoxins includes an

818   inhibitor of the sensory neuron calcium channel Cav2.3 *PLoS Biol* **21**, e3002217

819   28.   Yang, Z. (1997) PAML: a program package for phylogenetic analysis by maximum

820   likelihood *Comput Appl Biosci* **13**, 555-556

821   29.   Hordijk, P. L., de Jong-Brink, M., ter Maat, A., Pieneman, A. W., Lodder, J. C., and Kits,

822   K. S. (1992) The neuropeptide schistosomin and haemolymph from parasitized snails

823   induce similar changes in excitability in neuroendocrine cells controlling reproduction and

824   growth in a freshwater snail *Neurosci Lett* **136**, 193-197

825   30.   Frickey, T., and Lupas, A. (2004) CLANS: a Java application for visualizing protein

826   families based on pairwise similarity *Bioinformatics* **20**, 3702-3704

827   31.   Bertelsen, A. B., Hackney, C. M., Bayer, C. N., Kjelgaard, L. D., Rennig, M., Christensen,

828   B. *et al.* (2021) DisCoTune: versatile auxiliary plasmids for the production of disulphide-

829   containing proteins and peptides in the E. coli T7 system *Microb Biotechnol* 10.1111/1751-

830   7915.13895

831   32.   Gaciarz, A., Khatri, N. K., Velez-Suberbie, M. L., Saaranen, M. J., Uchida, Y., Keshavarz-

832   Moore, E. *et al.* (2017) Efficient soluble expression of disulfide bonded proteins in the

cytoplasm of Escherichia coli in fed-batch fermentations on chemically defined minimal media *Microb Cell Fact* **16**, 108

33. Nielsen, L. D., Foged, M. M., Albert, A., Bertelsen, A. B., Soltoft, C. L., Robinson, S. D. *et al.* (2019) The three-dimensional structure of an H-superfamily conotoxin reveals a granulin fold arising from a common ICK cysteine framework *J Biol Chem* **294**, 8745-8759

34. Safavi-Hemami, H., Li, Q., Jackson, R. L., Song, A. S., Boomsma, W., Bandyopadhyay, P. K. *et al.* (2016) Rapid expansion of the protein disulfide isomerase gene family facilitates the folding of venom peptides *Proc Natl Acad Sci USA* **113**, 3227-3232

35. Rogov, V. V., Rozenknop, A., Rogova, N. Y., Lohr, F., Tikole, S., Jaravine, V. *et al.* (2012) A universal expression tag for structural and functional studies of proteins *Chembiochem* **13**, 959-963

36. Buczek, O., Bulaj, G., and Olivera, B. M. (2005) Conotoxins and the posttranslational modification of secreted gene products *Cell Mol Life Sci* **62**, 3067-3079

37. Fischer, W. H., and Spiess, J. (1987) Identification of a mammalian glutaminyl cyclase converting glutaminyl into pyroglutamyl peptides *Proc Natl Acad Sci USA* **84**, 3628-3632

38. van Kempen, M., Kim, S. S., Tumescheit, C., Mirdita, M., Lee, J., Gilchrist, C. L. M. *et al.* (2024) Fast and accurate protein structure search with Foldseek *Nat Biotechnol* **42**, 243-246

39. Krissinel, E., and Henrick, K. (2004) Secondary-structure matching (SSM), a new tool for fast protein structure alignment in three dimensions *Acta Crystallogr D Biol Crystallogr* **60**, 2256-2268

40.   Derache, C., Epinette, C., Roussel, A., Gabant, G., Cadene, M., Korkmaz, B. *et al.* (2012)

Crystal structure of greglin, a novel non-classical Kazal inhibitor, in complex with

subtilisin *FEBS J* **279**, 4466-4478

41.   Shenoy, R. T., Thangamani, S., Velazquez-Campoy, A., Ho, B., Ding, J. L., and Sivaraman,

J. (2011) Structural basis for dual-inhibition mechanism of a non-classical Kazal-type

serine protease inhibitor from horseshoe crab in complex with subtilisin *PloS one* **6**, e18838

42.   Rivier, J., Galyean, R., Gray, W. R., Azimi-Zonooz, A., McIntosh, J. M., Cruz, L. J. *et al.*

(1987) Neuronal calcium channel inhibitors. Synthesis of omega-conotoxin GVIA and

effects on 45Ca uptake by synaptosomes *J Biol Chem* **262**, 1194-1198

43.   Qi, L., Iskols, M., Shi, D., Reddy, P., Walker, C., Lezgiyeva, K. *et al.* (2024) A mouse DRG

genetic toolkit reveals morphological and physiological diversity of somatosensory neuron

subtypes *Cell* **187**, 1508-1526 e1516

44.   Bhuiyan, S. A., Xu, M., Yang, L., Semizoglou, E., Bhatia, P., Pantaleo, K. I. *et al.* (2024)

Harmonized cross-species cell atlases of trigeminal and dorsal root ganglia *Sci Adv* **10**,

eadj9173

45.   Tian, M., Benedetti, B., and Kamoun, S. (2005) A Second Kazal-like protease inhibitor

from Phytophthora infestans inhibits and interacts with the apoplastic pathogenesis-related

protease P69B of tomato *Plant Physiol* **138**, 1785-1793

46.   Tian, M., and Kamoun, S. (2005) A two disulfide bridge Kazal domain from Phytophthora

exhibits stable inhibitory activity against serine proteases of the subtilisin family *BMC*

Biochem **6**, 15

47.   Li, X. C., Wang, X. W., Wang, Z. H., Zhao, X. F., and Wang, J. X. (2009) A three-domain

Kazal-type serine proteinase inhibitor exhibiting domain inhibitory and bacteriostatic

activities from freshwater crayfish Procambarus clarkii *Dev Comp Immunol* **33**, 1229-1238

48.   Esmaeilishirazifard, E., Usher, L., Trim, C., Denise, H., Sangal, V., Tyson, G. H. *et al.*

(2022) Bacterial Adaptation to Venom in Snakes and Arachnida *Microbiol Spectr* **10**,

e0240821

49.   De Jong-Brink, M. (1995) How schistosomes profit from the stress responses they elicit in

their hosts *Adv Parasitol* **35**, 177-256

50.   Zhang, S. M., Nian, H., Wang, B., Loker, E. S., and Adema, C. M. (2009) Schistosomin

from the snail Biomphalaria glabrata: expression studies suggest no involvement in

trematode-mediated castration *Mol Biochem Parasitol* **165**, 79-86

51.   Force, A., Lynch, M., Pickett, F. B., Amores, A., Yan, Y. L., and Postlethwait, J. (1999)

Preservation of duplicate genes by complementary, degenerative mutations *Genetics* **151**,

1531-1545

52.   van de Locht, A., Lamba, D., Bauer, M., Huber, R., Friedrich, T., Kroger, B. *et al.* (1995)

Two heads are better than one: crystal structure of the insect derived double domain Kazal

inhibitor rhodniin in complex with thrombin *The EMBO journal* **14**, 5149-5157

53.   Maxwell, M., Undheim, E. A. B., and Mobli, M. (2018) Secreted Cysteine-Rich Repeat

Proteins "SCREPs": A Novel Multi-Domain Architecture *Front Pharmacol* **9**, 1333

54.   Liu, J., Maxwell, M., Cuddihy, T., Crawford, T., Bassetti, M., Hyde, C. *et al.* (2023)

ScrepYard: An online resource for disulfide-stabilized tandem repeat peptides *Protein Sci*

32, e4566

55.   Katoh, K., Misawa, K., Kuma, K., and Miyata, T. (2002) MAFFT: a novel method for rapid

multiple sequence alignment based on fast Fourier transform *Nucleic Acids Res* **30**, 3059-

3066

56.   Katoh, K., and Standley, D. M. (2013) MAFFT multiple sequence alignment software

version 7: improvements in performance and usability *Mol Biol Evol* **30**, 772-780

57.   Waterhouse, A. M., Procter, J. B., Martin, D. M., Clamp, M., and Barton, G. J. (2009)

Jalview Version 2--a multiple sequence alignment editor and analysis workbench

Bioinformatics **25**, 1189-1191

58.   Slater, G. S., and Birney, E. (2005) Automated generation of heuristics for biological

sequence comparison *BMC Bioinformatics* **6**, 31

59.   Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von

Haeseler, A. *et al.* (2020) IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era *Mol Biol Evol* **37**, 1530-1534

60.   Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q., and Vinh, L. S. (2018)

UFBoot2: Improving the Ultrafast Bootstrap Approximation *Mol Biol Evol* **35**, 518-522

61.   Rambaut, A. (2018) FigTree (Version 1.4.4) Institute of Evolutionary Biology, University

of Edinburgh,

62.   Khilji, M. S., Hackney, C. M., Koch, T. L., Hone, A. J., Rogalski, A., Watkins, M. *et al.*

(2025) Structural similarities reveal an expansive conotoxin family with a two-finger toxin

fold *Protein Sci* **34**, e70333

63.   Wilkins, M. R., Gasteiger, E., Bairoch, A., Sanchez, J. C., Williams, K. L., Appel, R. D. *et*

al. (1999) Protein identification and analysis tools in the ExPASy server *Methods Mol Biol*

112, 531-552

64.   Kazimierczuk, K., and Orekhov, V. Y. (2011) Accelerated NMR spectroscopy by using

922   compressed sensing *Angew Chem Int Ed Engl* **50**, 5556-5559

923   65.   Delaglio, F., Grzesiek, S., Vuister, G. W., Zhu, G., Pfeifer, J., and Bax, A. (1995) NMRPipe:

924   a multidimensional spectral processing system based on UNIX pipes *J Biomol NMR* **6**,

925   277-293

926   66.   Skinner, S. P., Fogh, R. H., Boucher, W., Ragan, T. J., Mureddu, L. G., and Vuister, G. W.

927   (2016) CcpNmr AnalysisAssign: a flexible platform for integrated NMR analysis *J Biomol*

928   *NMR* **66**, 111-124

929   67.   Güntert, P., and Buchner, L. (2015) Combined automated NOE assignment and structure

930   calculation with CYANA *J Biomol NMR* **62**, 453-471

931   68.   Schwieters, C. D., Kuszewski, J. J., Tjandra, N., and Clore, G. M. (2003) The Xplor-NIH

932   NMR molecular structure determination package *J Magn Reson* **160**, 65-73

933   69.   Bermejo, G. A., Clore, G. M., and Schwieters, C. D. (2012) Smooth statistical torsion angle

934   potential derived from a large conformational database via adaptive kernel density

935   estimation improves the quality of NMR protein structures *Protein Sci* **21**, 1824-1836

936   70.   Tian, Y., Schwieters, C. D., Opella, S. J., and Marassi, F. M. (2017) High quality NMR

937   structures: a new force field with implicit water and membrane solvation for Xplor-NIH *J*

938   *Biomol NMR* **67**, 35-49

939   71.   Laskowski, R. A., Rullmannn, J. A., MacArthur, M. W., Kaptein, R., and Thornton, J. M.

940   (1996) AQUA and PROCHECK-NMR: programs for checking the quality of protein

941   structures solved by NMR *J Biomol NMR* **8**, 477-486

942 72. Olp, M. D., Kalous, K. S., and Smith, B. C. (2020) ICEKAT: an interactive online tool for

943   calculating initial rates from continuous enzyme kinetic traces *BMC Bioinformatics* **21**,

944   186

945 73. Teichert, R. W., Smith, N. J., Raghuraman, S., Yoshikami, D., Light, A. R., and Olivera, B.

946   M. (2012) Functional profiling of neurons through cellular neuropharmacology *Proc Natl*

947   *Acad Sci USA* **109**, 1388-1395

948