Title: **Genomic resources of *Ascidiella aspersa* and comparative analysis across tunicates reveal**
conserved class-level features and evolutionary diversification

Takumi T. Shito[1], Vasanthan Jayakumar[2], Koki Nishitsuji[3,5], Yoshie Nishitsuji[3,5], Shimon
Kawai[4], Shunsuke O. Miyasaka[4], Kotaro Oka[2], Yasubumi Sakakibara[2]* and Kohji Hotta[4]*

[1] Department of Chemistry, Biology and Marine Science, Faculty of Science, University of the
Ryukyus, Okinawa, Japan

[2] School of Frontier Engineering, Kitasato University, Sagamihara, Japan

[3] Department of Advanced Aquaculture Science, Faculty of Marine Science and Technology, Fukui
Prefectural University, Obama, Fukui, 917-0116, Japan.

[4] Department of Bioscience and Informatics, Faculty of Science and Technology, Keio University,
Yokohama, 223-8522, Japan

[5] Marine Genomics Unit, Okinawa Institute of Science and Technology Graduate University,
Okinawa, Japan.

*Correspondence: Kohji Hotta khotta@bio.keio.ac.jp,

Yasubumi Sakakibara sakakibara.yasubumi@kitasato-u.ac.jp

Keywords: *De novo* assembly, tunicate, ascidian, Ascidiidae, *Ascidiella aspersa*, codon usage
bias, GC content, genome evolution, phylogenomics, evo-devo

Abstract (349 / 350 words)

Background

Ascidiella aspersa is an invasive tunicate, one of the closest relatives of vertebrates. Despite its
ecological nuisance, *A. aspersa* is anticipated to be a valuable model organism for developmental
studies due to its remarkably transparent embryos. However, annotated genome assemblies and
transcriptomic resources have not yet been fully established. Although several tunicate genomes have
been sequenced, most lack annotations, and a comprehensive analysis across tunicates has not yet
been conducted.

Results

We performed *de novo* genome assembly and transcriptome analysis of *A. aspersa,* producing a high-
quality 306.5 Mb genome assembly. The transcriptome was derived from nine different organs of
adults and embryos at six developmental stages. Ab initio and homology-based gene predictions
identified 24,504 genes as part of the KAS25 gene model with a BUSCO score of 92.2%. Functional
annotation was added for 18,636 genes in the model. To understand the relative features of this
species among tunicates, we conducted genome-wide comparative analysis using publicly available
data from 35 other tunicate genomes across three classes, five orders, and 12 families, constructing
gene models for 27 species with BUSCO scores >80%. Overall, Phylogenetic analysis revealed a
new hypothesis regarding the relationships among Phlebobranchia and Aplousobranchia families.
Gene duplication analysis showed distinct contractions of gene families in some taxa, with losses of

specific DNA repair related genes that were shared among Thaliacea, which may have contributed to
their evolutionary diversification. Tunicate genomes exhibited a high level of variation in genomic
GC content (28.0%–42.7%). *A. aspersa* has the highest GC content in coding regions and the third
position of codons among tunicate species, with changes in codon usage bias differing from other
Ascidiidae species. We also constructed an online comparative tunicate genome database
(TUNOME), that provide functional annotations of gene models and ortholog analyses based on
these genomic and transcriptomic data.

Conclusions

46   We constructed genomic resources for *A. aspersa* and another 35 tunicate gene models. Comparative
47   analysis reveals a variety in tunicate species genomes and characterizes class-level features. Our
48   resources are expected to serve as a foundation for experimental studies involving non-model
49   tunicates and for comparative analysis among tunicate species.

Background

Tunicates belong to a sister group of vertebrates and consist of approximately 3,000 species
traditionally divided into three classes: Appendicularia, Thaliacea, and Ascidiacea [1]. Their unique
evolutionary position makes them important clades in evo-devo studies, particularly in Ascidiacea
species, which have sessile adults and planktonic embryos suitable for developmental experiments
due to their many unique merits, including a short period of embryogenesis, invariant cleavage
patterns, and compact genomes [2]. For studying initial development, *Ciona intestinalis* type A
(*Ciona robusta*) and *Halocynthia roretzi* are primarily used for experiments and have led to
significant discoveries in a variety of scientific fields [2, 3].

Recently, *Phallusia mammillata* has emerged as a new model ascidian due to its remarkably
transparent embryos, which enable whole embryo live imaging and thereby greatly facilitate the
understanding of the developmental mechanisms in chordate [4]. The family Ascidiidae, which
includes *P. mammillata*, is characterized by distinctive biological traits such as transparent eggs and
embryos [5]. However, as *P. mammillata* is geographically restricted to Europe, alternative model
organisms with similar optical transparency are needed in other regions.

One promising candidate is *Ascidiella aspersa,* a cosmopolitan ascidian species belonging to the
family Ascidiidae. Although *A. aspersa* is an invasive species that causes biofouling in shellfish
aquaculture, particularly in Japan [6–8], its biological characteristic make it an exceptionally valuable
experimental organism. The eggs and embryos of *A. aspersa* are highly transparent—transmitting
approximately 90% of visible light—and its unfertilized eggs retain mRNA translation capability [5,
9]. Moreover, a standardized developmental table and a three dimensional embryological image
resource (RAMNE) have already been established for this species [9].

These features, together with its worldwide distribution and ease of collection, highlight *A. aspersa*
as a practical and powerful model system for developmental and molecular studies. Although low
fertilization efficiency has been limitation, an improved fertilization protocol has recently been
developed to overcome this issue [10]. Despite these advances, a high-quality, annotated genome
assembly—an essential foundation for molecular and genetic analyses—has not yet been available
for *A. aspersa*.

Since the draft genome of *Ciona intestinalis* [11], many ascidian genomes have been reported. The
ANISEED genome database has been developed, which includes model ascidians like *H. roretzi* and
P. mammillata [3]. ANISEED currently hosts 12 tunicate genomes, and additional tunicate genomes
are available in NCBI; however, some lack gene annotation or contain low-quality gene annotations.
Furthermore, comprehensive comparative analysis across tunicate genomes has not yet been
conducted.

Here, we present a *de novo* genome assembly and transcriptome analysis of *A. aspersa*, along with
updated gene models for 38 tunicate species. We also compared genomic statistics among species,
such as genome size and GC content, characterizing specific tunicate taxa. The newly built genomic
resources for tunicates will facilitate comparative analysis across tunicate families and expand the use
of non-model tunicates in evo-devo studies.

Methods

Genome sequencing and assembly of *A. aspersa*.

A. aspersa specimens were collected from fixed scallop ropes at Onagawa (Onagawa Field Center,
Tohoku University) in October 2019. Sperm from a single adult was isolated for DNA extraction.
DNA was extracted using the Qiagen DNeasy Kit and quantified using a spectrophotometer and
bioanalyzer; subsequently, it was stored at -30°C until sequencing. The genome sequence was
obtained using Pacific Bioscience (PacBio) Sequel and Illumina NovaSeq. PacBio generated over
46.7 Gb of long reads (2.9 million reads), while Illumina paired-end sequencing produced 70 Gb,
which was used for three rounds of error correction of long-read data (Table S2).

For genome assembly, wtdbg2 [12], SMARTdenovo [13], Flye [14], and Canu [15] were employed.
Three rounds of polishing were performed. Repetitive sequences in the genome were identified and
masked using Repeat Modeler v2.0.4 [16] and RepeatMasker v4.1.5 [17]. Completeness of the
genome assembly was assessed using BUSCO v5 [18] on the gVolante platform [19], utilizing the
eukaryota database.

Transcriptome sequencing and assembly of *A. aspersa*

Samples for the transcriptome analysis were also collected from Onagawa, Hakodate (Hokkaido
Research Organization, Hakodate Fisheries Research Institute), and Yoichi (Hokkaido Research
Organization, Central Fisheries Research Institute) in Japan. Adult organs were dissected and
preserved in RNAlater at -80°C until RNA isolation. Matured eggs isolated from gonoducts were
dechorionated using a solution of 0.05% actinase-E and 1% mercaptoacetic acid sodium salt in
seawater. Dechorionated eggs and intact embryos were fixed with liquid nitrogen and stored at -80°C.
Samples were disrupted by manipulation and repeated freezing with liquid nitrogen, followed by
treatment with the RNeasy kit (QIAGEN).

Nine adult organs (gill cage, atrial siphon, oral siphon, fascia, granule, intestinal tract, orange organs,
anus, and gonad) were dissected, and eggs or embryos at five developmental stages (gastrula, neurula,
early tailbud, mid tailbud, and hatching larva at 25 hpf) were also collected (Fig. S1). For
insemination, eggs and sperm were isolated from gonoducts and spermducts, respectively. The sperm
were pre-activated with pH 9.5 ASW for 10–15 minutes, following a method used in experiments
with *Phallusia* species [20]. Pre-activated sperm were added to eggs in seawater, then diluted

fivefold 20 minutes after fertilization. Embryos were incubated at 20°C, and four developmental
stages were fixed with liquid nitrogen according to the standard developmental table (Funakoshi et al.
2021). Fixed embryos were stored at -80°C. Embryos with chorions were treated using the TransZol
Plant kit (TransGen Biotech).

From the extracted RNA, stranded library was prepared with polyA selection and paired-end
sequencing was performed using Illumina MiSeq and Illumina NovaSeq 6000, producing an average
of 62.0 million reads, comprising 8.67 GBases per sample (Table S4). The acquired transcriptome
data were quality-checked using FastQC v0.11.9 [21] and SeqKit v2.3.0 [22]. Adapter sequences
were trimmed using fastp v0.20.1 [23]. Adapter-trimmed reads were assembled using Trinity v2.14.0
[24]. After adapter trimming, paired-end RNA sequences were aligned to the Trinity assembly using
Hisat2 [25], achieving nearly 90% mapping efficiency. Reads that failed to align concordantly were
discarded, and the remaining clean reads were subsequently aligned to the genome assembly for
downstream analyses (Table S5).

Gene model construction of *A. aspersa*

Gene prediction was performed using two software packages according to Ito et al. [26]. The soft-
masked assembly was initially annotated by BRAKER2 pipeline [27] with metazoan proteins of
orthoDB10 [28]. To improve prediction accuracy, we also applied the GeMoMa v1.9 pipeline [29],
which incorporates RNA-seq evidence and existing gene models of phylogenetically related species
for homology-based gene prediction. Gene models from *Ciona robusta* and *Ascidia mentula,*
produced with BRAKER2 and having high BUSCO scores, were selected as references for GeMoMa.
For exhaustive prediction, genes without start/stop codons were accepted in the GeMoMa step. The
two predicted gene models were integrated using GffCompare [30]. We relied on the GeMoMa
model, and additional annotations from BRAKER2 were adopted.

Translated protein sequences were obtained from the merged gene model using GffRead [30], then
processed with R scripts to omit alternative isoforms by retaining the longest splicing variant.
Functional annotation was added using EnTAP [31] with default settings and the following protein
resources: Universal Protein Resource (Swissprot, Trembl) and RefSeq (Invertebrate). BRAKER-
predicted genes were filtered by the GeMoMa filter.

Gene model construction of other tunicate genomes

Genome sequences were downloaded from repositories such as NCBI or ANISEED. The genome of
Ciona robusta [32, 33] was downloaded from the Kyoto Ghost Database (https://ghost.zool.kyoto-
u.ac.jp/download_ht.html). Genomes of *Ciona savignyi* [34], *Botrylloides schlosseri* [35],
Botrylloides leachii [36], and *Molgura* spp. [37], as well as *Oikopleura dioica* [38] have been
reported and downloaded from ANISEED. *Phallusia mammillata, Phallusia fumigata*, *Halocynthia
roretzi*, and *Halocynthia aurantium* have not been published but are available on ANISEED [39].
Oikopleura dioica [40], five Oikopleuridae spp. [41], *Clavelina lepadiformis* [42], *Aplidium
turbinatum* [43], *Trididemnum clinides* [44], *Corella eumyota* [45], *Corella inflata* [46], *Ascidia
mentula* [47], *Styela plicata* [48], and *Styela clava* [49] have been published and are available on
NCBI. *Pegea* sp., *Salpa aspera*, *Salpa thompsoni*, *Thalia democratica*, *Trididemnum miniatum*,
Trididemnum nubilum, *Diplosoma virens*, *Perophora annectens*, *Halocynthia papillosa*, and *Boltenia
villosa* have not been published but are available on NCBI. First chromosome-level genome

assembly of *Botryllus schlosseri* [50] and *Didemnum vexillum* [51] have been published but are
unavailable and hence not used in this study. In total, tunicate genomes from 37 species are available
in NCBI genome resources, although many of them lack gene models (Table S1).

All available genomes were downloaded. Before repeat masking step, the headers of FASTA files
were processed to be shorter and characters were converted to uppercase. For existing gene models,
the numbers of genes and transcripts were checked using GeMoMa scripts. BUSCO completeness of
protein sequences was assessed using BUSCO v5 on gVolante with the eukaryota dataset. Available
gene models with 95% BUSCO value were processed with the GeMoMa filter and finalized. Braker2
was performed on other tunicate genomes, while BRAKER3 [52] was used for some genomes due to
the failure of BRAKER2 in highly fragmented genomes. Gene models predicted by BRAKER2 with
>95% completeness were processed with the GeMoMa filter and finalized. Other genomes with gene
models <95% were subsequently processed with GeMoMa gene prediction using high quality gene
models from relatively close species. Combinations of species were tested, and predictions with
higher BUSCO completeness were selected. Alternative splicing was permitted in all prediction steps
and processed by R scripts to select the representative longest splicing variants for subsequent
analysis. Protein sequences with stop codons were omitted from downloaded models.

180
Phylogenetic analysis

OrthoFinder v2.5.4 [53] was used to analyze tunicate amino acid sequences from multiple species'
genomes. One of cephalochordate *Branchiostoma floridae* was selected as an outgroup and genomic
data were downloaded from the NCBI database via accession number GCF_000003815.2.
Phylogenetic analysis was performed according to Maeda et al [54]. In each orthogroup, gene trees
were constructed using MAFFT and FastTree [55] in OrthoFinder. A species tree with all species was
drawn from gene trees in orthogroups with all species present using STAG in OrthoFinder [56]. For
more precise analysis without the effect of paralogues, single-copy orthogroups were used in the
following analysis. Protein sequences were trimmed using trimAl v1.4.rev15 [57] and aligned using
MAFFT v7.505 [58]. Aligned sequences were analyzed using IQ-TREE2 v2.1.4-beta [59] for
maximum likelihood analysis. Bootstrap resampling with 1,000 pseudo-replicates was performed to
obtain statistical support for the nodes. The ML tree was converted to an ultrametric tree based on
divergence times between Cephalochordata and the clade comprising Vertebrata and Tunicata (578
Myr), between Stolidobranchia and the clade comprising Thaliacea, Phlebobranchia, and
Aplousobranchia (389 Myr), between Thaliacea and the clade comprising Phlebobranchia and
Aplousobranchia (296 Myr), and between Molgulidae and the clade comprising Pyulidae and
Styelidae (350 Myr) [1]. The conversion was conducted using r8s v1.81 [60] for subsequent analysis.

Gene duplication analysis

The percentage of duplicated genes was calculated by the ratio of genes assigned to any orthogroup
to the total gene number. Rapidly expanded or contracted families were identified using CAFÉ v4.2.1
[61]. For Gene Ontology (GO) analysis, all protein sequences were used as a query to search human
protein sequences using BLASTP. Homologies were filtered by BLASTP e-value < 0.01. Genes with
the lowest e-value in each orthogroup were used as representative genes. Enrichment analysis was
performed using the enrichGO function in the R package clusterProfiler v4.2.1 [62].

GC content analysis of tunicate genomes

GC content in tunicate genomes was collected from the NCBI database or calculated using SeqKit. Coding sequences (CDS) were extracted using gffread. CDS sequences, including alternative splices, were converted into unique sequences for each locus, and sequences without start/stop codons were omitted using R scripts. GC content in each position of codons was calculated using R scripts. Statistics were presented using the ggplot2 package in R [63]. Kernel density estimation was performed using the geom_density function. Codon usage of each CDS was calculated using cusp script in EMBOSS [64]. Dimension reduction of codon usage fractions was carried out via PCA utilizing the built-in R packages. A parameter evaluating codon usage bias was effective number of codons (ENC), ranging from 20 (only one synonymous codon for encoding amino acids) to 61 (every synonymous codon is used equally), calculated using CodonW v1.4.4 (http://codonw.sourceforge.net/). A smaller ENC value indicates stronger codon usage bias. ENC-GC3 plots and theoretical ENC curves were drawn using the formula (1) [65]. If codon usage bias is affected only by mutation pressure, the genes will be located on or near the theoretical curve, while natural selection will position them below the curve [66].

$$ENC = 2 + GC3s + \frac{29}{GC3s^2 + (1 - GC3s^2)} \ (1)$$

Results

Genome assembly and transcriptome profiling of *Ascidiella aspersa*

To establish a reference genome for *A. aspersa*, we performed hybrid assembly using long-read (PacBio Sequel) and short-read (Illumina NovaSeq 6000) sequencing, achieving approximately 46,8 Gb and 141.4 Gb, respectively. After comparing the results of all 4 assemblies (Table S3), wtdbg2 assembly, refined through multiple rounds of error correction, was selected for downstream analyses because it yielded the longest contig and the highest N50 value (Table 1). The final assembly had 92.2% BUSCO completeness.

Table1. Statistics for the assembly and annotation

Genome assembly		
Total assembled length		306,513,209 bp
Number of contigs		1,411
Mean Length		217,231 bp
Largest contig		12,818,253 bp
Contig N50		2,058,548 bp
Repeat sequences (%)		41.3%
GC (%)		36.9%
BUSCO	complete	94.9%
	partial	3.1%
	fragment	2.0%
Gene Model (KAS25 model)		
Genes		24,504
Transcripts		30,856
BUSCO	complete	92.2%

partial	5.9%
fragment	1.9%

Transcriptome sequencing across nine adult organs and six embryonic stages generated over 60
million reads per sample. The assembled RNA sequences yielded 101,713 transcripts, comprising 97
MBases (Table S5).

Braker2 produced a gene model comprising 20,886 genes and 21,623 transcripts for the *A. aspersa*
genome, assessed with a relatively low BUSCO score of 69.8% (Table 2). GeMoMa integrating
RNA-seq evidence and homology-based refinement predicted 19,302 genes and 25,645 transcripts,
achieving a higher BUSCO completeness of 92.2% (Table 2; Table S6). Two models were integrated,
yielding the final gene model comprises 24,504 genes and 30,856 transcripts, with BUSCO
completeness of 92.2% (Table 1). This gene model has been designated as the Keio *Ascidiella*
aspersa 2025 version (KAS25) gene model.

In total, 18,636 of the 24,504 predicted protein sequences were successfully annotated (Table 2).
Homology-based searches identified similarities for 18,636 sequences across the following
databases: Uniprot_sprot (9,134 sequences), Uniprot_trembl (14,642 sequences), and Refseq_inV
(9,494 sequences). Family-based annotation assigned 18,098 sequences, including 15,049 with GO
terms and 5,492 with KEGG terms. In total, 5,868 sequences remained without functional annotation.

Table 2. Summary of Functional annotation of KAS25 model

	Gene/Transcripts (BUSCO)
Braker2 gene model	20886/21623 (69.8%)
GeMoMa gene model	19302/25645 (92.2%)
Total genes number of KAS25 model	24504/30856 (92.2%)
Total functional annotated gene numbers	18,636
Total unique sequences with an alignment by similarity search:	16,872
Total unique transcripts with an alignment: uniprot_sprot	9,134
Total unique transcripts with an alignment: uniprot_trembl	14,642
Total unique transcripts with an alignment: refseq_inV	9,494
Total unique sequences with family assignment:	18,098
Total unique sequences with at least one GO term:	15,049
Total unique sequences with at least one pathway (KEGG) assignment:	5,492
Total unannotated genes	5,868

Gene model construction of other tunicate genomes

In total, 37 tunicate genomes spanning three classes, five orders, and 12 families were downloaded
from online databases (Table S1). Genome sizes ranged from 64.3 Mb (*Oikopleura dioica*) to 1.1 Gb
(*Trididemnum nubilum*). However, BUSCO completeness scores varied widely, from 17.3% to
98.9%. Three species (*Clavelina lepadiformis*, C*iona robusta*, *Styela clava*) exhibited completeness
>95% and were used for downstream analysis without modification (Table S6). For the remaining 34

species, gene models were reconstructed using BRAKER2. In two cases (*B. villosa* and *M.*
erythrocephalus), gene predictions using BRAKER2 failed, possibly due to highly fragmented
genome sequences, and BRAKER3 was used instead. Five species (*Pegea* sp., *Thalia democratica*,
Ascidia mentula, *Corella eumyota*, and *Botrylloides violaceus*) produced gene models with
completeness greater than 95% using BRAKER2 and were finalized.

For the remaining 27 genomes, we applied the GeMoMa pipeline, using reference gene models of
closely related species with higher BUSCO values. Multiple combination of one to three reference
genomes was tested in each GeMoMa prediction step, and gene models with higher BUSCO values
were selected (Table S6). BRAKER2 and GeMoMa models were then integrated for each genome to
maximize completeness, except for three species (*Oikopleura albicans*, *Oikopleura longicauda*, and
Oikopleura vanhoeffeni), where BRAKER2 alone provided more accurate models (Table S6).

The number of transcripts, including alternatively spliced isoforms, as well as the BUSCO
duplication rate, increased with the number of reference species used. However, the duplication value
was generally lower than 10% when alternative isoforms were removed (Table S6).

Overall, our newly constructed gene models showed an average improvement of 15.2% in BUSCO
scores compared to 12 existing online models. Finally, 27 of the 34 models we constructed achieved
more than 80% BUSCO completeness, with the exception of seven species (*Bathochordaeus stygius*,
Mesochordaeus erythrocephalus, *Oikopleura albicans*, *Oikopleura longicauda*, *Oikopleura*
vanhoeffeni, *Boltenia villosa*, and *Botryllus schlosseri*) (Table S7; Table 3).

Table 3 Gene models of all 38 tunicate genomes

Class	Order	Family	Species name	genes	BUSCO completeness
Appendicularia Copelata Oikopleuridae			*Bathochordaeus stygius*	16,907	16.1%
			Mesochordaeus erythrocephalus	22,295	19.6%
			Oikopleura albicans	37,978	77.7%
			Oikopleura dioica	16,744	83.1%
			Oikopleura longicauda	62,461	76.5%
			Oikopleura vanhoeffeni	32,793	57.7%
Thaliacea Salpida Salpidae			*Pegea* sp.	13,826	96.5%
			Salpa aspera	32,638	93.7%
			Salpa thompsoni	29,086	94.5%
			Thalia democratica	37,001	96.9%
Ascidiacea Aplousobranchia		Polyclinidae	*Aplidium turbinatum*	24,461	98.1%
		Clavelinidae	*Clavelina lepadiformis*	25,119	95.7%
		Didemnidae	*Diplosoma virens*	33,042	96.8%
			Trididemnum clinides	34,563	94.9%
			Trididemnum miniatum	34,498	96.8%
			Trididemnum nubilum	21,480	96.5%

Ascidiacea Phlebobranchia	Ascidiidae	*Ascidiella aspersa*	24,504	92.2%
		Ascidia mentula	20,419	97.3%
		Phallusia fumigata	39,443	89.4%
		Phallusia mammillata	28,259	96.5%
	Cionidae	*Ciona robusta*	18,788	99.2%
		Ciona savignyi	21,684	93.3%
	Corellidae	*Corella eumyota*	18,534	98.4%
		Corella inflata	29,590	80.8%
	Perophoridae	*Perofora annectens*	23,788	89.8%
Ascidiacea Stollidobranchia	Styelidae	*Styela plicata*	23,705	98.8%
		Styela clava	20,018	96.5%
		Polycarpa mamillaris	34,763	94.5%
		Botrylloides leachii	17,311	98.9%
		Botryllus schlosseri	54,922	80.4%
		Botrylloides violaceus	14,677	95.7%
	Pyuridae	*Boltenia villosa*	32,157	51.4%
		Halocynthia aurantium	20,018	91.4%
		Halocynthia papillosa	14,723	98.4%
		Halocynthia roretzi	15,643	97.7%
	Molgulidae	*Molgula occidentalis*	27,511	92.2%
		Molgula occulta	28,185	87.5%
		Molgula oculata	23,318	93.3%

Comparison of gene numbers in constructed gene models

Species with larger genomes generally exhibited a higher number of predicted genes (Fig. S2a–b). *B.*
schlosseri and *O. longicauda* exhibit an exceptionally high number of predicted genes relative to
their moderate genome sizes, compared to other tunicates. Notably, these species show relatively low
BUSCO completeness scores (≤80%). Excluding gene models with low BUSCO values, a clearer
positive correlation between genome size and gene number was observed.

Pegea sp., *B. violaceus*, *H. papillosa*, *H. roretzi*, *O. dioica*, and *B. leachii* have relatively small
genomes (<200 Mb) and correspondingly low gene counts (<18,000). There was no strong tendency
for closely related species to cluster together in terms of gene number or genome size.

The ratio of duplicated genes to total gene number was also calculated and showed a positive
correlation with total gene number (Fig. S2c–d). Species such as *H. papillosa*, *H. roretzi*, *Pegea sp.*,
B. violaceus, and *O. dioica* exhibited small genomes, low gene counts, and low gene duplication
ratios. Since gene duplication rates can be affected by genome completeness, species with BUSCO
completeness ≤90% were excluded from further analysis. This exclusion clarified the trend: species

with larger genomes tended to have higher numbers of duplicated genes (Fig. 1a).

Phylogenetic analysis

We first conducted ortholog analysis using 39 species (38 tunicate species and *Branchiostoma*
floridae as an outgroup) with OrthoFinder. Of the total 1,053,122 genes, 959,536 genes were
assigned to 60,568 orthogroups. Because the 39 species did not share any single copy orthologs,
shared 1,518 orthogroups containing paralogs derived from all 39 species were used to construct the
species tree using STAG (Fig. S3a). The resulting topology supported recent class-level analyses,
such as the paraphyly in Ascidiacea, including Thaliacea [1]. The monophyly of Appendicularia and
its internal relationships were also supported in recent phylogenetic studies [67] and the monophyly
of Phlebobranchia and Aplousobranchia was consistent with [68], while STAG analysis contains the
effect of paralogs.

To minimize the impact of paralogs, species were ranked by the BUSCO completeness of their gene
models, and those with lower scores were progressively excluded to maximize the number of single
copy orthologs (Table 8). We reduced the number of species from 39 to 32 (BUSCO > around 80%),
30 (BUSCO > 85%), 29, and 28 (BUSCO > around 90%), which yielded 44,324, 42,405, 42,405, and
39,311 orthogroups, respectively (Table 8). The number of single copy orthogroups progressively
increased to 94, 217, 240, and 385 as the number of species decreased.

Consequently, 28 tunicate species were chosen, and 385 single-copy orthologs were used for
phylogenetic analysis based on maximum likelihood analysis using IQ-TREE (Fig. 1b; Fig. S3b).
Bootstrap values supported the tree with nearly 100%, except for the nodes of Corellidae and
Cionidae, which had 94% support (Fig. S3b). ML Analysis with other species subsets (Table S8)
produced the same topology.

In the resulting tree, Stolidobranchia was the earliest branching group, followed by Thaliacea, then
the clade comprising Ascidiidae and Perophoridae, and finally the group of Corellidae, Cionidae, and
Aplousobranchia. This tree suggests that Aplousobranchia was nested within the paraphyletic
Phlebobranchia, which is inconsistent with STAG phylogeny (Fig. S4). This paraphyly of
Phlebobranchia is consistent with recent studies, although the relationships of families differ [1, 46,
69] (Fig. S4). Delsuc et al. [1] and Shenkar et al. [69] positioned Ascidiidae as the sister group of
Cionidae and the Aplousobranchia clade, consistent with our tree, although [69] placed Corellidae as
a sister group of Ascidiidae, in contrast to our tree, which suggests the sisterhood of Corellidae and
Cionidae. On the other hand, [46] placed Cionidae as a sister group of Aplousobranchia, Ascidiidae,
and Corellidae clade, while [68] positioned Aplousobranchia as a sister group of Ascidiidae,
Corellidae, and Cionidae (Fig. S4). These four studies did not include Perophoridae, and our study
suggests a sister group relationship between Ascidiidae and Perophoridae.

Gene duplication analysis

We used 28 species with BUSCO scores >90%, as gene model completeness is important for
duplication analysis. The percentage of duplicated genes was calculated by the results from
OrthoFinder and plotted against genome size (Fig. 1a). Higher genome sizes corresponded to higher
numbers of duplicated genes. Two species in *Halocynthia* and two species in *Botrylloides* exhibited
low duplication rate (44%–56%). In contrast, *T. democratica* showed an approximately 85%

duplication rate. These tendencies were preserved in the scatter plot of duplication rate and total gene
number (Fig. 1a).

The numbers of rapidly expanded or contracted gene families were investigated by CAFÉ and
mapped onto the phylogenetic tree of the 28 species (Fig. 1b). Interestingly, most gene families
showed rapid expansion rather than contraction. Exceptions were observed in *Botrylloides* and
Thaliacea, which exhibited rapid contraction. *Botrylloides* species also show a low duplication rate in
overall gene sets (Fig. 1a). By contrast, *Halocynthia* species, despite their low duplication rates, did
not exhibit substantial gene family contraction. Gene ontology enrichment analysis suggested that
rapidly contracted orthogroups in *Botrylloides* and Thaliacea contain some groups related to
development, morphogenesis, and metabolism (Table S9).

Rapidly expanded gene families in *A. aspersa* and gene families unique to this species were also
investigated. A total of 51 gene families were identified as rapidly expanded, among which 21
showed significant similarity to human genes (BLASTP, e-value < 0.01). These families are mainly
associated with DNA binding, hormone biosynthetic process, transcriptional regulation, and
extracellular matrix organization (Table S9). In addition, 315 gene families were found to be unique
to *A. aspersa*. Gene Ontology analysis indicated that these genes are enriched in functions related to
the extracellular matrix, endoplasmic reticulum, and the omega-hydroxylase P450 pathway.

346
Comparison based on Genome size and GC content
Firstly, genome size and GC content from 38 tunicate species genomes were mapped (Fig. 2a–b).
Genome size varied 17-fold, ranging from 64.3 Mb in *Oikopleura dioica* to 1,100 Mb in
Trididemnum nubilum. In Appendicularia, *Oikopleura vanhoeffeni* and *Mesochordaeus*
erythrocephalus show relatively high genome sizes (643.7 Mb and 874 Mb, respectively). Thaliacean
species, including *Thalia democratica* (790.7 Mb), *Salpa aspera* (903.3 Mb), and *Salpa thompsoni*
(742.8 Mb), also exhibit large genome sizes. *Pegea* sp. is an exception with a small genome size of
105.8 Mb. Among ascidians, seven of the nine species with genomes larger than 400 Mb were
colonial.

Genome GC content varied from 25.3% in *T. democratica* to 42.7% in *Polycarpa mamillaris.*
Thaliacean species, including *T. democratica* and *Pegea* sp., show very low GC content values
(25.3% and 28.0%). In contrast, six species (*P. mamillaris*, *Botrylloides violaceus*, *Botrylloides*
leachii, *Oikopleura dioica*, *Perophora annectens*, and *Botryllus schlosseri*) exhibited relatively high
GC content above 40%, four of which belong to the family Styelidae.

A dense cluster of species was observed in the region of genome sizes between 120 Mb and 400 Mb
and GC content between 35% and 37% (Fig. 2b). This cluster included two appendicularians
(*Oikopleura* albicans and *Bathochordaeus stygius*) and 11 ascidian species. Of these 11 ascidians,
except for the colonial *Clavelina lepadiformis*, all were solitary. Genera represented by more than
two species generally clustered together (Fig. 2a–b). Species within the same genus were mapped
relatively close to each other. Notable exceptions were *Molgula occidentalis*, *Oikopleura* species, and
Trididemnum species, which were dispersed among genera; three species in the same genus
(*Trididemnum* and *Oikopleura*) were arranged linearly. In general, genus-level clusters exhibited a
downward-right slope, except in closely related genera such as *Ciona*, *Styela*, and *Botrylloides*.

GC content in coding regions (CDS) shows different trends compared to whole-genome GC content.
GC content in CDS ranged from 37% to 47%, typically 5% to 10% higher than genomic GC within
the same species (e.g., 28.0% to 37.2% in *Pegea* sp.) (Fig. 2c–d; Table S9). While the relative

positions among species were mostly conserved, several species (*T. democratica*, *T. clinides*, *A.*
aspersa, *O. dioica*) exhibited marked shifts. Despite its extremely low genomic GC content, CDS GC
content in *T. democratica* showed an intermediate value, with an increase of almost 17% (Fig. 2c). *T.*
clinides and *O. dioica* also shifted in relative position within their genera. Interestingly, *A. aspersa*
exhibited the highest CDS GC content among all 38 genomes (47.1%), whereas its genomic GC
content was intermediate (Fig. 2).

380   GC content of tunicate genomes in different codon positions

To further investigate GC variation, we calculated GC content at the first, second, and third codon
positions (GC1, GC2, GC3, respectively) for each species (Table S9). On average, GC1 showed a
relatively high value of 48.3%, while GC2 was relatively low at 38% (Fig. 3a). Interestingly,
compared to GC1 and GC2, GC3 exhibited a wide distribution, ranging from 29.9% (*Pegea* sp.) to
50.2% (*O. dioica*). GC contents across all regions were positively correlated, especially GC3, which
showed the strongest correlation with CDS GC content (correlation coefficient = 0.975; Fig. 3b).
Kernel density estimation of genome GC content revealed a major peak around 35% in both
Appendicularia and Ascidiacea. In these two classes, GC peaks were also evident in CDS and GC3,
whereas Appendicularia displayed a bimodal peak in GC1 and a downward shift in GC2. In contrast,
Thaliaceans exhibited different positions, particularly with low values in *Pegea* sp. (Fig. 3b).

We next examined GC content distributions for each gene in individual species (Fig. S5). In most
species, GC1–3 density profiles formed sharp, unimodal peaks, while *O. longicauda*, *T. clinides*, *C.*
lepadiformis, and *C. eumyota* displayed bimodal distributions. *O. longicauda* exhibited two peaks at
around 30% and 52%, while *C. eumyota* showed a small second peak at about 50% across GC1–3. *T.*
clinides displayed a second peak at about 60% only in GC1 and GC3. *C. lepadiformis* exhibited
multiple minor peaks on the upper side of the main peak. Among Appendicularia, *O. dioica* showed a
GC3 peak shift upwards. In Thaliacea, GC3 in *T. democratica* displayed a relatively low, rounded
peak shape. In Phlebobranchia, *A. aspersa* showed a gradually shifted peak compared to other species
(Fig. 3c; Fig. S5d). In Stolidobranchia, the shapes of the curves were almost identical, except for a
lower shift in GC3 of *M. occidentalis*.

Overall, GC1 tended to be relatively high, GC2 relatively low, and GC3 showed the widest variation
among species. Notably, GC3 most strongly reflected interspecific differences and correlated closely
with the overall CDS GC content.

405   Codon usage bias in tunicate genomes

Codon usages of all 38 tunicates were calculated (Fig. S6). By combining usage fractions in all
codons, we performed principal component analysis (PCA) and extracted PC1 and PC2 (Fig. 4a).
These components accounted for 58.8% and 13.3% of the overall variance, respectively.
Appendicularians were located distantly from other tunicates, except for *O. vanhoeffeni* and *O. dioica*.
Similarly, Pegea was positioned separately from other Salpidae species. Three *Trididemnum* species
in Didemnidae, three molgulidae species, and four Pyulidae species exhibited clustering in the PCA
map (Fig. 4a; Table S11). Interestingly, *A. aspersa* was located in isolation from other ascidiids (Fig.
4a).

Compared to other Ascidiidae, *A. aspersa* exhibited some changes in codon usage fractions (Fig. 4b).
For example, in codons for cysteine (C), *A. aspersa* prefers TGC than TGT, while other Ascidiidae
species favor TGT. Similarly, *A. aspersa* prefers GAC rather than GAT for aspartic acid (D), while
other species show the opposite tendency. Additionally, *A. aspersa* favors CAC was preferred than
CAT for histidine (H) codons. Fractional changes were also notable in alanine (A), lysine (K),
arginine (R), and threonine (T). Many of these changes involved a conversion from A/T to G/C at the
third codon. These tendency switches were also observed in comparisons with other phlebobranchs.

Codon usage variation analysis in related species is also remarkable in *O. dioica,* contrary to other
Appendicularia species (Fig. S6a). Tendency changes were observed in cysteine (C), phenylalanine
(F), Histidine (H), isoleucine (I), Asparagine (N), and tyrosine (Y), sharing C and H with *A. aspersa.*
In the AT-rich *Pegea* sp., significant switches in usage tendency were only observed in Asparagine
(N), but the usage bias was enhanced in many codons (Fig. S6b). In Phlebobranchia, Corella species
shared codon bias switches compared to other phlebobranchs, particularly in asparagine (N).
Aplousobranchia, Stolidobranchia species exhibited fewer changes in codon usage bias.

For codon usage variation analysis, ENC-GC3 plots were drawn (Fig. 5; Fig. S4). A larger ENC
value indicates a stronger bias in codon usage and a smaller ENC value indicates equal usage among
codons. Mutational effects may be the dominant factor in the codon preference of genes on the
expected curve [65]. A position below the curve indicates that the preference is under selection
pressure. Distributions of genes in most Appendicularia is comparatively below the left of the peak
(Fig. S7a). In contrast, *O. dioica* shows a clear tendency of extending down to the right, indicating
some popularity of genes under the influence of natural selection. *O. longicauda* shows high GC
genes but is relatively positioned higher compared to *O. dioica*, which may be driven by sequence.

Among Thaliacea, *Pegea* sp. is distinctly positioned on the left side of the standard curve (Fig. S7b).
Plots of the other three thaliaceans had similar shapes, extending to the right side along the standard
curve. Aplousobranchia species show almost left-side distributions along the curve, but *T. clinides*
displays a right distribution similar to the left. The tendency of right distribution is weakly observed
in other *Trididemnum* species, *T. miniatum* and *T. nubilum* (Fig. S7c).

Phlebobranchians show distributions at the top of the peak and extending left, except for *Ascidiella*
aspersa, which has a right distribution (Fig. S7d). Stolidobranchians show almost AT-side
distributions along the standard line distribution, except for *B. schlosseri* (Fig. S7e).

Overall, codon usage in ascidians reflects patterns characteristic of each family, with some
exceptions, such as *A. aspersa*, *O. dioica,* and *Pegea* sp., which shows clear preferences diverging
from closely related species (Fig. 4; Fig. 5).

Discussion

Variation in genome size and gene numbers in tunicates

Consistent with our results, some reports indicate a positive relationship between genome size and
predicted gene size, although protein-coding regions contribute only a small fraction to overall
genome size [70]. Genome size and organism complexity are not necessarily related, but cell size and
cell division rate are associated; larger genome sizes are observed in nuclei of larger, slowly dividing
cells [70]. Compound species *Botryllus schlosseri*, which has a relatively large genome in tunicates

(about 580 Mb), exhibits slow embryo development compared to *Ciona robusta*, which has a smaller
genome (about 123 Mb) [71]. To verify the existence of a constraint on developmental speed and
genome size, further investigation of embryogenesis in other compound ascidian species with larger
genome sizes is required, as this remains unclear in most species.

O. dioica has extreme genome scrambling caused by thousands of chromosomal rearrangements,
which may be promoted by the loss of the canonical nonhomologous end-joining (NHEJ) DNA
repair pathway among appendicularians [67, 72]. Our dataset also suggests a lack of DNA repair
pathways containing NHEJ in appendicularians (Fig. S8). Interestingly, thaliacean species also share
a lack of genes associated with DNA repair (smug1, MUTYH, MPG, BRCA1, XRCC4), although
Appendicularia and Thaliacea are not sister groups. Base Excision Repair (BER), which includes
smug1, MUTYH, and MPG, is thought to increase GC to AT transitions when impaired [73, 74]. As
suggested in appendicularians, potential losses in DNA repair pathways may have occurred in the
common ancestor of thaliaceans, affecting the high variability in thaliacean genomes, which is
represented by genome size, GC value, and the rapid evolutionary rate observed in the maximum
likelihood tree (Fig. S3b). Thaliaceans have a large number of rapidly contracted gene families
containing genes associated with development and metabolism (Fig. S1b; Table S9). These drastic
genome changes and the loss of specific gene sets may be associated with the remarkable lifestyle
transition from a sessile form to a motile form in Thaliaceans compared to other ascidian species.

473
The phylogeny of the Tunicata

Phylogenetic analysis in tunicates has been complicated by accelerated rates of evolution in their
nuclear and mitochondrial genomes, which could result in long-branch attraction (LBA) [1]. The
long branch observed in *Oikopleura dioica* reflects the high evolutionary rate of this species,
complicating the debate regarding its basal position as the sister group of all other tunicates [75].
Additionally, 18S rRNA sequences of Aplousobranchia species seem to evolve at high rates, causing
long branches [76]. In our STAG tree, long branches were not observed in Appendicularians,
suggesting that transcriptomic-level analysis might resolve the long branch effects in tunicates,
although this analysis contains the effects of paralogs (Fig. S3a). In ML tree, long branches were not
seen in Aplousobranchia, while Thaliacea contained a slightly long branch (Fig. S3b).

Two main unresolved phylogenetic questions in tunicates are: 1) the relationships among the three
classes of tunicates, and 2) the relationships between Phlebobranchia and Aplousobranchia. For the
first question, reliable appendicularian genomes are necessary, as there is still no reliable
phylogenetic analysis based on ML or Bayesian. Salpidae in Thaliacea were placed within
Ascidiacea with 100% bootstrap support in our analysis (Fig. 1b; Fig. S3b), consistent with recent
literature, while the monophyly of Thaliacea should be proven through phylogenomics approaches.
Regarding the second question, there is no consensus on the relationships among families in
Phlebobranchia and Aplousobranchia, with completely reliable bootstrap values (Fig. S4). To address
this problem, comprehensive sampling of various taxa is required, as there is a substantial lack of
representatives from five families in Phlebobranchia, including Agneziidae and Plurelidae, many of
which inhabit deep-sea environment. The problem is even more serious in Aplousobranchia, which
has the highest number of species and families in tunicates, making species identification relatively
difficult [77]. For phylogenomic approaches, several alternative strategies may be helpful: using
genes that are not necessarily single-copy [78] and accurately identifying orthologs [79].

498
Variety of GC Content in tunicate genomes

T. clinides exhibits bimodal GC content, and several other species have relatively high GC values in
their CDS. High GC content in genomes can complicate gene prediction. In our analysis, BRAKER2
proved to be a powerful tool for various species, including *A. mentula*, *P. mammillata*, and *B.*
violaceus, generating gene models that included nearly all BUSCO genes predicted across the whole
genome (Table S12). However, BRAKER2 produced less accurate gene models for some species, as
indicated by a significant decline in BUSCO scores from the whole genome to the predicted gene
models (Table S12). In these cases, GeMoMa performed well, producing BUSCO scores around 90%
(Table S12). Our run in GeMoMa ignored missing start/stop codons, which almost explained the
gaps observed in some species (*S. thompsoni*, *S. aspera*, and *P. annectens*) (Table S12). There still
remain gaps between BRAKER2 and GeMoMa in *T. clinides* and *A. aspersa*. Gene prediction may
be challenging due to high GC content, as observed in the gene prediction of the western honey bee
genome, which also exhibits bimodal GC content [80].

A study reported a mean GC content of 39.9% for sampled metazoan genomes, although some
species show unusually high or low GC content, such as the hydroid *Hydra magnipapillata* (29%
GC) and the sea lamprey *Petromyzon marinus* (46% GC) [81]. Interestingly, tunicates exhibit a wide
range of genomic GC content within a single subphylum, ranging from 25.3% in *T. democratica* to
42.7% in *P. mamillaris*. In the scatter plot of genome size versus genomic GC content, species within
the same genus were either clustered closely or dispersed along a downward-sloping trend line in
genera with more than two species, such as *Trididemnum* and *Oikopleura* (Fig. 2a). A negative
correlation between genome size and GC content has also been reported in monocot genomes,
possibly due to the higher energetic cost of nucleotide synthesis, which causes a deficiency of dGTP
and dCTP, leading to the misincorporation of the less costly dATP and dTTP, and an overall
mutation bias toward AT-rich DNA [82].

Regarding coding regions, GC content was generally more than 5% higher than that of the whole
genome, ranging from 29.9% to 50.2% (Fig. 2c–d). These values of CDS GC content were relatively
low compared to a dataset of 33 mammals, which ranged from 43.9% to 57.9% [83]. Tunicate
genomes exhibited higher GC content at the first codon position (GC1) than at the second codon
position (GC2), while the third codon position (GC3) showed a wider range among tunicate species
compared to GC1 and GC2 (Fig. 3a). GC indices have positive correlations with each other,
especially GC3, which has a high correlation with CDS GC content (Fig. 3b). This tendency is also
observed in the prokaryotes and viral genes [84].

Codon usage bias in tunicates

Synonymous codon composition affects translation efficiency, and natural selection acts on codon
usage [85, 86]. Our data revealed some change in codon usage bias among related species in
tunicates (Fig. 4a). Notably, *A. aspersa* exhibited changes in codon usage tendencies compared to
other ascidiids, particularly for cysteine, aspartic acid, histidine, and others (Fig. 4b). Codon usage
transfer of these amino acids was also observed in *O. dioica*. The ENC-GC3 plot was used to
determine the effect of mutation bias versus natural selection on GC3 content bias. Most ascidian
genomes show a left-down distribution on the theoretical line, indicating that mutational bias is
dominant (Fig. 5). Three species in Thaliacea show an upper distribution of GC compared to
ascidians. *T. clinides* have a clearly right-down shape, while other *Trididemnum* species show a weak
tendency in that direction, which may indicate phylogenetic closeness. Conversely, *O. dioica* exhibits
a right-down distribution relative to the theoretical line, suggesting that some preferred codons were
selected for translational efficiency under natural selection [86].

Another important force contributing to nucleotide heterogeneity is thought to be GC-biased gene
conversion (gBGC), a bias in the DNA repair machinery that results in meiotic distortion favoring G
and C over A and T alleles in highly recombining regions [81, 83]. *O. dioica* exhibits high GC
content and has an extremely short life cycle of only five days at 20°C [87], which may accelerate the
effects of gBGC by increasing recombination rates. In this scenario, the right-down shape of the
ENC-GC3 plot in this species may reflect not natural selection on translational efficiency but rather
the effects of gBGC, as gBGC mechanically restricts the number of codons by shaping local base
compositions with increasing GC content, including surrounding non-coding regions, which also
reduces the ENC [86]. gBGC is thought to be primarily caused by the mismatch repair system [88],
which genes were conserved in *O. dioica*, except for POLD3 (Fig. S8c).

A. aspersa was characterized by normal genomic GC and remarkably high GC3, along with a
different codon usage bias compared to related species. *A. aspersa* also shows a right-down
distribution along the theoretical line in the ENC-GC3 plot, which may indicate that it is not
subjected to natural selection or gBGC. The actual cause of the high mutation bias toward GC and
the effect of this GC bias on its ecology is still unknown. Further investigations into the
characteristics of associated genes, such as those involved in DNA repair and environmental factors,
across tunicate genomes, will reveal the distinct evolution of this species.

The genomic basis of tunicates

We constructed a genome assembly of *A. aspersa* and gene models across 38 tunicates. We
integrated all constructed genome sequences, gene models, functional annotations, and ortholog
analyses into a new tunicate genome database, TUNOME
(https://ciona.bpni.bio.keio.ac.jp/Tunome/Latest/). For an initial overview, functional annotations of
all 38 tunicates can be easily searched using various queries, such as gene ID, common names, or
domain names (Fig. S9a). All annotations corresponding to a selected gene can be displayed,
Expresion level represented with transcripts per million (TPM) are graphically showed, and both
nucleotide and protein sequences are available for download (Fig. S9b). The phylogram and all
relative sequences of the orthogroup containing the selected gene can also be viewed (Fig. S9c). A
genome browser is provided, enabling visualization of RNA-seq alignments derived from different
organs or embryonic stages (Fig. S9d). Additionally, BLAST searches can be performed against all
38 genomes and their gene models (Fig. S9e). The "Blast to all species" mode allows simultaneous
searches across all species, and both summary tables and hit sequences can be downloaded (Fig. S9f).

Using our resources, 38 tunicate genomes, including non-model animals, can be easily compared.
Our resources also bridge comparative evo-devo: orthogroup analyses anchored by our gene models
can aid investigations into shared features, such as bio-transparency in ascidian embryos [89]. These
resources can also be helpful for developmental experiments. Our database sheds light on
experiments in non-model animals and facilitates species comparison among tunicate traits.

Conclusions

We constructed genomic and transcriptomic resources for *A. aspersa*. Other tunicate species'
genomes were also annotated, and gene models were rebuilt, achieving high-quality gene models
containing 27 species with BUSCO scores greater than 80%. Phylogenetic analysis produced a new

hypothesis in tunicate phylogeny, and gene duplication analysis characterized multiple taxa within
tunicates. Genome statistics of tunicates exhibited high variability in genome size and GC content,
including high GC content in the CDS of *A. aspersa* genome and codon usage bias. This analysis and
our constructed genome database, TUNOME, can accelerate evo-devo research through species
comparisons, including non-model organisms.

Data availability

594 Genome sequencing data and assembly were deposited to BioProject accession PRJNA1345923.
595 Gene models and protein sequences for other species were deposited in the TUNOME database:
596 https://ciona.bpni.bio.keio.ac.jp/Tunome/Latest/

Abbreviations

GC1: GC content in the first codon

GC2: GC content in the second codon

GC3: GC content in the third codon

gBGC: GC-biased gene conversion

ENC: Effective number of codons

PCA: Principal component analysis

Author Contributions

607 KH conceived the project. KH and TS designed the experiments. TS and SK performed experiments
608 with living samples. KN, YN performed the sequencing. VJ and YS conducted the assembly. TS, SK,
609 SM, and KN carried out the gene model construction. TS performed subsequent analysis and
610 constructed the website. TS and KH drafted the manuscript. KO, YS, KN, and KH critically revised
611 the manuscript and supervised the research. All authors reviewed the manuscript.

Funding

This study was supported by JSPS KAKENHI (21H00440, 23H04717, 24K02038, and 25H01799),
the Keio University Research and Education Center for Natural Sciences Budget, and the KLL Keio
Leading Program to KH. The Research Institute of Marine Invertebrates (IKU2021-02) supported
TTS. The Keio University Doctorate Student Grant-in-Aid Program from the Ushioda Memorial
Fund supported TTS. JSPS KAKENHI Grant Numbers JP 22J22628 and 25KJ0268 supported TTS.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

Acknowledgements

We thank Dr. Minoru Ikeda and Captain Toyokazu Hiratsuka (Onagawa Field Center of Tohoku
University); Dr. Makoto Kanamori, Dr. Masafumi Natsuike, and Takuya Mizukami (Hokkaido
Research Organization, Hakodate Fisheries Research Institute); Dr. Gaku Kumano (Graduate School
of Life Sciences, Tohoku University); and Akio Takiya, Kazuya Sakai, Dr. Tatsunari Mori, Dr.
Takaaki Kayaba, and Dr. Motohito Yamaguchi (Hokkaido Research Organization, Central Fisheries
Research Institute) for their assistance in collecting samples. We also thank Dr. Kazuho Ikeo and Dr.
Sonoko Kinjo (National Institute of Genetics) and Dr. Takashi Makino (Tohoku University) and
Naohiro Hasegawa (Hiroshima Shudo University) and Euichi Hirose (University of Ryukyus) for
their helpful comments. We thank Mr. Wu Jiayang for building the computer for the analysis. We
thank Ms. Haruka M. Funakoshi and Ms. Yuki S. Kogure for handling *Ascidiella aspersa* samples.
We are grateful to Dr. Ken Naito (The University of Tokyo), Dr. Taro Maeda (Keio University), and
Dr. Jun Inoue (The University of Tokyo) for sharing valuable articles about genome analysis
methods on their website. We would like to express our sincere gratitude to Dr. Nori Satoh for
sequencing support.

Figure Legends

Table 1 Statistics for the assembly of *A. aspersa*

Statistics of the adopted assembly of the *A. aspersa* genome and constructed gene model (KAS25
model) summary. BUSCO values were assessed on the eukaryote gene sets.

Table 2 Gene prediction and functional annotation of *A. aspersa*

Summary of functional annotations attached to the KAS25 gene model. Gene numbers and transcript
numbers containing alternative splicing (BUSCO value) are shown.

Table 3 Gene models of all 38 tunicate genomes

Gene numbers, transcript numbers, and BUSCO values of constructed gene models of 38 tunicate
genomes. Classes, orders, and families to which they belong are also shown.

Fig. 1 Phylogeny of tunicates and duplicated gene families

a) Scatter plot of duplicated gene ratio versus genome size in 28 tunicate species with gene models
showing about 90% or higher BUSCO completeness. b) Phylogenetic tree constructed using the ML
method with 27 tunicates and *Branchiostoma floridae* as the outgroup. Families are separated by
different colors, and body plan or life cycles (solitary swimming, solitary fixed, compound

swimming, compound fixed) of clades are illustrated at right. The numbers of highly expanded or
contracted gene families are shown at each clade in red or blue.

Fig. 2 Comparison of genome size and GC contents

a) Scatter plot of genome size versus GC content in whole genomes among all 38 tunicate species.
Species in the same genus were clustered. b) Close-up view of the crowded area within the orange
rectangle in a). c) Scatter plot of genome size versus GC content in coding regions (CDS). d) Close-
up view of the crowded area within the orange rectangle in c). Species positions that drastically
changed are exhibited as red or orange colors.

Fig. 3 GC content in different positions of codons

a) Box plot of GC content in different regions of genomes. GC1–3 refers to GC content at the first,
second, and third positions of codons. One dot represents the average value of one species. b) Scatter
plot, density plot, and correlations of each combination of GC contents in different genome regions.
Red, green, and blue colors correspond to Appendicularia, Ascidiacea, and Thaliacea, respectively. c)
Density plot of GC content in all genes from different positions in the *A. aspersa* gene model
(KAS25 model).

Fig. 4 Codon usage analysis in tunicate species

a) Principal component analysis was performed on codon usage data of 38 tunicate species. Red,
green, and blue colors correspond to Appendicularia, Ascidiacea, and Thaliacea, respectively.
Different shapes of dots correspond to all families of tunicates. PC1 and PC2 accounted for 58.8%
and 13.3% of the overall variance, respectively. b) Codon usages of amino acids in four ascidiids.
The vertical axis show the fractions of each codon. Black pins indicate the reversal of codon usage
compared to other Ascididae species. White pins indicate codon usage changes that are not as
significant as reversals.

Fig. 5 ENC-GC3 plot of 38 tunicates

Scatter plot of effective codon numbers (ENC) and GC content in the third codon position (GC3) in
38 tunicates. Axes were omitted almost and the example in *Oikopleura vanhoeffeni* is shown (Right
Upper). One dot corresponds to one gene. The theoretical line was plotted according to the
description in the methods section.

References

1. Delsuc F, Philippe H, Tsagkogeorga G, Simion P, Tilak M-K, Turon X, et al. A phylogenomic
framework and timescale for comparative studies of tunicates. BMC Biol. 2018;16:39.

2. Satoh N. Developmental Genomics of Ascidians. Nashville, TN: John Wiley & Sons; 2014.

3. Dardaillon J, Dauga D, Simion P, Faure E, Onuma TA, DeBiasse MB, et al. ANISEED 2019: 4D
exploration of genetic data for an extended range of tunicates. Nucleic Acids Res. 2020;48:D668–75.

4. Yasuo H, McDougall A. Practical Guide for Ascidian Microinjection: Phallusia mammillata. Adv
Exp Med Biol. 2018;1029:15–24.

5. Shito TT, Hasegawa N, Oka K, Hotta K. Phylogenetic comparison of egg transparency in ascidians
by hyperspectral imaging. Sci Rep. 2020;10:20829.

6. Lynch SA, Darmody G, O'Dwyer K, Gallagher MC, Nolan S, McAllen R, et al. Biology of the
invasive ascidian Ascidiella aspersa in its native habitat: Reproductive patterns and parasite load.
Estuar Coast Shelf Sci. 2016;181:249–55.

7. Palanisamy SK, Thomas OP, P McCormack G. Bio-invasive ascidians in Ireland: A threat for the
shellfish industry but also a source of high added value products. Bioengineered. 2018;9:55–60.

8. Nishikawa T, Oohara I, Saitoh K, Shigenobu Y, Hasegawa N, Kanamori M, et al. Molecular and
morphological discrimination between an invasive ascidian, Ascidiella aspersa, and its congener A.
scabra (Urochordata: Ascidiacea). Zoolog Sci. 2014;31:180–5.

9. Funakoshi HM, Shito TT, Oka K, Hotta K. Developmental Table and Three-Dimensional
Embryological Image Resource of the Ascidian Ascidiella aspersa. Frontiers in Cell and
Developmental Biology. 2021;9. https://doi.org/10.3389/fcell.2021.789046.

10. Kogure YS, Hotta K. An improved fertilization protocol for*Ascidiella aspersa*: Promoting a new
chordate model in developmental biology. bioRxiv. 2025;:2025.03.10.642315.
https://doi.org/10.1101/2025.03.10.642315.

11. Dehal P, Satou Y, Campbell RK, Chapman J, Degnan B, De Tomaso A, et al. The draft genome
of Ciona intestinalis: insights into chordate and vertebrate origins. Science. 2002;298:2157–67.

12. Ruan J, Li H. Fast and accurate long-read assembly with wtdbg2. Nat Methods. 2020;17:155–8.

13. Liu H, Wu S, Li A, Ruan J. SMARTdenovo: a de novo assembler using long noisy reads.
Gigabyte. 2021;2021:1–9.

14. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat
graphs. Nat Biotechnol. 2019;37:540–6.

15. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable and
accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res.
2017;27:722–36.

16. Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, et al. RepeatModeler2 for
automated genomic discovery of transposable element families. Proc Natl Acad Sci U S A.
2020;117:9451–7.

17. Smit, AFA, Hubley, R & Green, P. RepeatMasker Open-4.0. 2013-2015.

18. Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO update: Novel and
streamlined workflows along with broader and deeper phylogenetic coverage for scoring of

eukaryotic, prokaryotic, and viral genomes. Mol Biol Evol. 2021;38:4647–54.

19. Nishimura O, Hara Y, Kuraku S. gVolante for standardizing completeness assessment of genome
and transcriptome assemblies. Bioinformatics. 2017;33:3635–7.

20. Sensui N, Itoh Y, Okura N, Shiba K, Baba SA, Inaba K, et al. Spawning-Induced pH Increase
Activates Sperm Attraction and Fertilization Abilities in Eggs of the Ascidian, Phallusia
philippinensis and Ciona intestinalis. Int J Mol Sci. 2023;24:2666.

21. Andrews S. FastQC: a quality control tool for high throughput sequence data. 2010.

22. Shen W, Le S, Li Y, Hu F. SeqKit: A cross-platform and ultrafast toolkit for FASTA/Q file
manipulation. PLoS One. 2016;11:e0163962.

23. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics. 2018;34:i884–90.

24. Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, et al. De novo
transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation
and analysis. Nat Protoc. 2013;8:1494–512.

25. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol. 2019;37:907–15.

26. Ito M, Ohashi H, Takemoto M, Muto C, Seiko T, Noda Y, et al. Single candidate gene for salt
tolerance of Vigna nakashimae (Ohwi) Ohwi & Ohashi identified by QTL mapping, whole genome
sequencing and triplicated RNA-seq analyses. Breed Sci. 2024;74:93–102.

27. Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M. BRAKER2: automatic eukaryotic
genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database. NAR
Genom Bioinform. 2021;3:lqaa108.

28. Waterhouse RM, Tegenfeldt F, Li J, Zdobnov EM, Kriventseva EV. OrthoDB: a hierarchical
catalog of animal, fungal and bacterial orthologs. Nucleic Acids Res. 2013;41 Database issue:D358-
65.

29. Keilwagen J, Hartung F, Grau J. GeMoMa: Homology-Based Gene Prediction Utilizing Intron
Position Conservation and RNA-seq Data. Methods Mol Biol. 2019;1962:161–77.

30. Pertea G, Pertea M. GFF Utilities: GffRead and GffCompare. F1000Res. 2020;9:304.

31. Hart AJ, Ginzburg S, Xu MS, Fisher CR, Rahmatpour N, Mitton JB, et al. EnTAP: Bringing
faster and smarter functional annotation to non-model eukaryotic transcriptomes. Mol Ecol Resour.
2020;20:591–604.

32. Satou Y, Nakamura R, Yu D, Yoshida R, Hamada M, Fujie M, et al. A Nearly Complete Genome
of Ciona intestinalis Type A (C. robusta) Reveals the Contribution of Inversion to Chromosomal
Evolution in the Genus Ciona. Genome Biol Evol. 2019;11:3144–57.

33. Satou Y, Tokuoka M, Oda-Ishii I, Tokuhiro S, Ishida T, Liu B, et al. A Manually Curated Gene
Model Set for an Ascidian, Ciona robusta (Ciona intestinalis Type A). Zoolog Sci. 2022;39:253–60.

34. Small KS, Brudno M, Hill MM, Sidow A. A haplome alignment and reference sequence of the
highly polymorphic Ciona savignyi genome. Genome Biol. 2007;8:R41.

35. Voskoboynik A, Neff NF, Sahoo D, Newman AM, Pushkarev D, Koh W, et al. The genome
sequence of the colonial chordate, Botryllus schlosseri. Elife. 2013;2:e00569.

36. Blanchoud S, Rutherford K, Zondag L, Gemmell NJ, Wilson MJ. De novo draft assembly of the
Botrylloides leachii genome provides further insight into tunicate evolution. Sci Rep. 2018;8:5518.

37. Stolfi A, Lowe EK, Racioppi C, Ristoratore F, Titus Brown C, Swalla BJ, et al. Divergent
mechanisms regulate conserved cardiopharyngeal development and gene expression in distantly
related ascidians. Elife. 2014;3. https://doi.org/10.7554/eLife.03728.

38. Wang K, Tomura R, Chen W, Kiyooka M, Ishizaki H, Aizu T, et al. A genome database for a
Japanese population of the larvacean Oikopleura dioica. Dev Growth Differ. 2020;62.
https://doi.org/10.1111/dgd.12689.

39. Brozovic M, Dantec C, Dardaillon J, Dauga D, Faure E, Gineste M, et al. ANISEED 2017:
extending the integrated ascidian database to the exploration and evolutionary comparison of
genome-scale datasets. Nucleic Acids Res. 2018;46:D718–25.

40. Bliznina A, Masunaga A, Mansfield MJ, Tan Y, Liu AW, West C, et al. Telomere-to-telomere
assembly of the genome of an individual Oikopleura dioica from Okinawa using Nanopore-based
sequencing. BMC Genomics. 2021;22:222.

41. Naville M, Henriet S, Warren I, Sumic S, Reeve M, Volff J-N, et al. Massive changes of genome
size driven by expansions of non-autonomous transposable elements. Curr Biol. 2019;29:1161-
1168.e6.

42. Daric V, Lanoizelet M, Mayeur H, Leblond C, Darras S. Genomic Resources and Annotations for
a Colonial Ascidian, the Light-Bulb Sea Squirt Clavelina lepadiformis. Genome Biol Evol. 2024;16.
https://doi.org/10.1093/gbe/evae038.

43. Bishop J, Harley J, Mrowicki R, Marine Biological Association Genome Acquisition Lab,
Darwin Tree of Life Barcoding collective, Wellcome Sanger Institute Tree of Life programme, et al.
The genome sequence of *Aplidium turbinatum* (Savigny 1816), a colonial sea squirt. Wellcome Open
Res. 2022;7:106.

44. Hirose E, Lopez JV, Oatley G, Sinclair E, Aunin E, Gettle N, et al. The chromosomal genome
sequence of the photosymbiotic ascidian, Trididemnum clinides Kott, 1977 and its associated
microbial metagenome sequences. Wellcome Open Res. 2025;10:357.

45. Bishop J, Wood C, Mrowicki RJ, Harley J, Marine Biological Association Genome Acquisition
Lab, Darwin Tree of Life Barcoding collective, et al. The genome sequence of the Orange-tipped sea
squirt, Corella eumyota Traustedt, 1882. Wellcome Open Res. 2024;9:146.

46. DeBiasse MB, Colgan WN, Harris L, Davidson B, Ryan JF. Inferring Tunicate Relationships and
the Evolution of the Tunicate Hox Cluster with the Genome of Corella inflata. Genome Biol Evol.
2020;12:948–64.

47. Bishop J, Wood C, Marine Biological Association Genome Acquisition Lab, Darwin Tree of Life
Barcoding collective, Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory
team, Wellcome Sanger Institute Scientific Operations: Sequencing Operations, et al. The genome
sequence of a solitary sea squirt, Ascidia mentula (Müller, 1776). Wellcome Open Res. 2023;8:583.

48. Galià-Camps C, Schell T, Pegueroles C, Baranski D, Hamadou AB, Turon X, et al. Genomic
richness enables worldwide invasive success. Research Square. 2024.
https://doi.org/10.21203/rs.3.rs-3902873/v1.

49. Wei J, Zhang J, Lu Q, Ren P, Guo X, Wang J, et al. Genomic basis of environmental adaptation
in the leathery sea squirt (Styela clava). Mol Ecol Resour. 2020;20:1414–31.

50. De Thier O, Tawfeeq MM, Faure R, Lebel M, Dru P, Blanchoud S, et al. First chromosome-level
genome assembly of the colonial tunicate Botryllus schlosseri. bioRxiv. 2024.
https://doi.org/10.1101/2024.05.29.594498.

51. Parra-Rincón E, Velandia-Huerto CA, Gittenberger A, Fallmann J, Gatter T, Brown FD, et al.
The Genome of the "Sea Vomit" Didemnum vexillum. Life. 2021;11.
https://doi.org/10.3390/life11121377.

52. Gabriel L, Brůna T, Hoff KJ, Ebel M, Lomsadze A, Borodovsky M, et al. BRAKER3: Fully
automated genome annotation using RNA-seq and protein evidence with GeneMark-ETP,
AUGUSTUS, and TSEBRA. Genome Res. 2024;34:769–77.

53. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative genomics.
Genome Biol. 2019;20:238.

54. Maeda T, Takahashi S, Yoshida T, Shimamura S, Takaki Y, Nagai Y, et al. Chloroplast
acquisition without the gene transfer in kleptoplastic sea slugs, Plakobranchus ocellatus. Elife.
2021;10. https://doi.org/10.7554/eLife.60176.

55. Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood trees for large
alignments. PLoS One. 2010;5:e9490.

56. Emms DM, Kelly S. STAG: Species tree inference from all genes. bioRxiv. 2018;:267914.
https://doi.org/10.1101/267914.

57. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment
trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25:1972–3.

58. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements
in performance and usability. Mol Biol Evol. 2013;30:772–80.

59. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-
TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. Mol Biol
Evol. 2020;37:1530–4.

60. Sanderson MJ. R8s: Inferring absolute rates of molecular evolution and divergence times in the
absence of a molecular clock. Bioinformatics. 2003;19:301–2.

61. Han MV, Thomas GWC, Lugo-Martinez J, Hahn MW. Estimating gene gain and loss rates in the
presence of error in genome assembly and annotation using CAFE 3. Mol Biol Evol. 2013;30:1987–
97.

62. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrichment tool
for interpreting omics data. Innovation (Camb). 2021;2:100141.

63. Wickham H. Ggplot2: Elegant graphics for data analysis. 1st edition. New York, NY: Springer;
2009.

64. Rice P, Longden I, Bleasby A. EMBOSS: The European molecular biology open software suite.
Trends Genet. 2000;16:276–7.

65. Wright F. The "effective number of codons" used in a gene. Gene. 1990;87:23–9.

66. Wang Y, Jiang D, Guo K, Zhao L, Meng F, Xiao J, et al. Comparative analysis of codon usage
patterns in chloroplast genomes of ten Epimedium species. BMC Genom Data. 2023;24:3.

67. Plessy C, Mansfield MJ, Bliznina A, Masunaga A, West C, Tan Y, et al. Extreme genome
scrambling in marine planktonic Oikopleura dioica cryptic species. Genome Res. 2024;34:426–40.

68. Kocot KM, Tassia MG, Halanych KM, Swalla BJ. Phylogenomics offers resolution of major
tunicate relationships. Mol Phylogenet Evol. 2018;121:166–73.

69. Shenkar N, Koplovitz G, Dray L, Gissi C, Huchon D. Back to solitude: Solving the phylogenetic
position of the Diazonidae using molecular and developmental characters. Mol Phylogenet Evol.
2016;100:51–6.

70. Elliott TA, Gregory TR. What's in a genome? The C-value enigma and the evolution of
eukaryotic genome content. Philos Trans R Soc Lond B Biol Sci. 2015;370:20140331.

71. Anselmi C, Ishizuka KJ, Palmeri KJ, Burighel P, Voskoboynik A, Hotta K, et al. Speed vs
completeness: a comparative study of solitary and colonial tunicate embryogenesis. Front Cell Dev
Biol. 2025;13:1540212.

72. Deng W, Henriet S, Chourrout D. Prevalence of mutation-prone microhomology-mediated end
joining in a chordate lacking the c-NHEJ DNA repair pathway. Curr Biol. 2018;28:3337-3341.e4.

73. Moggioli G, Panossian B, Sun Y, Thiel D, Martín-Zamora FM, Tran M, et al. Distinct genomic
routes underlie transitions to specialised symbiotic lifestyles in deep-sea annelid worms. Nat
Commun. 2023;14:2814.

74. Krokan HE, Bjørås M. Base excision repair. Cold Spring Harb Perspect Biol. 2013;5:a012583.

75. Delsuc F, Brinkmann H, Chourrout D, Philippe H. Tunicates and not cephalochordates are the
closest living relatives of vertebrates. Nature. 2006;439:965–8.

76. Tsagkogeorga G, Turon X, Hopcroft RR, Tilak M-K, Feldstein T, Shenkar N, et al. An updated
18S rRNA phylogeny of tunicates based on mixture and secondary structure models. BMC Evol Biol.
2009;9:187.

77. Shenkar N, Swalla BJ. Global diversity of Ascidiacea. PLoS One. 2011;6:e20657.

78. Smith ML, Hahn MW. New approaches for inferring phylogenies in the presence of paralogs.
Trends Genet. 2021;37:174–87.

79. Inoue J, Satoh N. ORTHOSCOPE: An automatic web tool for phylogenetically inferring
bilaterian orthogroups with user-selected taxa. Mol Biol Evol. 2019;36:621–31.

80. Elsik CG, Worley KC, Bennett AK, Beye M, Camara F, Childers CP, et al. Finding the missing
honey bee genes: lessons learned from a genome upgrade. BMC Genomics. 2014;15:86.

81. Kyriacou RG, Mulhair PO, Holland PWH. GC content across insect genomes: Phylogenetic
patterns, causes and consequences. J Mol Evol. 2024;92:138–52.

82. Šmarda P, Bureš P, Horová L, Leitch IJ, Mucina L, Pacini E, et al. Ecological and evolutionary
significance of genomic GC content diversity in monocots. Proc Natl Acad Sci U S A.
2014;111:E4096-102.

83. Romiguier J, Ranwez V, Douzery EJP, Galtier N. Contrasting GC-content dynamics across 33
mammalian genomes: relationship with life-history traits and chromosome sizes. Genome Res.
2010;20:1001–9.

84. Bernardi G, Bernardi G. Compositional constraints and genome evolution. J Mol Evol.
1986;24:1–11.

85. Clessin A, Joseph J, Lartillot N. Evolution of GC-biased gene conversion by natural selection.
Genetics. 2025;230:iyaf111.

86. Romiguier J, Roux C. Analytical biases associated with GC-content in molecular evolution. Front
Genet. 2017;8:16.

87. Onuma TA, Nishida H. Developmental biology of the larvacean Oikopleura dioica: Genome
resources, functional screening, and imaging. Dev Growth Differ. 2022;64:67–82.

88. Glémin S, Arndt PF, Messer PW, Petrov D, Galtier N, Duret L. Quantification of GC-biased gene
conversion in the human genome. Genome Res. 2015;25:1215–28.

89. Hotta K, Miyasaka SO, Oka K, Shito TT. Staring into a crystal ball: understanding evolution and
development of in vivo aquatic organismal transparency. Front Ecol Evol. 2024;12:1428976.

Supplementary Material 1

SupplementaryMaterial_1.xlsx. Table S1–S12.

Supplementary Material 2

SupplementaryMaterial_2.docx. Fig. S1–S7.

Supplementary Material 3

SupplementaryMaterial_3.docx. Fig. S8–S9.

Fig. 1

a

b

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Appendicularia

B. stygius *M. eryth.* *O. albicans* *O. dioica* *O. longi.* *O. vanho.*

Effective Number of Codons (ENC)

0.0   0.5   1.0

GC3 / %

Thaliacea

Pegea sp. *S. aspera* *S. thomp.* *T. demo.*

Aplousobranchia

A. turbi. *C. lepa.* *D. virens* *T. clinides* *T. miniatum* *T. nubilum*

Phlebobranchia

A. aspersa *A. mentula* *P. fumigata* *P. mamillata* *C. eumyota* *C. inflata*

C. savignyi *C. robusta* *P. annectens*

B. villosa

Stolidobranchia

M. occi. *M. occulta* *M. oculata* *H. aura.* *H. papillosa* *H. roretzi*

S. plicata *S. clava* *P. mamillaris* *B. leachii* *B. violaceus* *B. shclosseri*