Patient-Specific Midbrain Organoids with CRISPR Correction Reveal Disease Mechanisms and Enable Therapeutic Evaluation in Neuronopathic Gaucher Disease

Yi Lin[1], Benjamin Liou[1], Venette Fannin[1], Stuart Adler[1], Christopher N. Mayhew[2,3], Jason E. Hammonds[2,4], Yueh-Chiang Hu[2,5], Jason Tchieu[2,6], Wujuan Zhang[7], Xueheng Zhao[2,7], Rebecca L. Beres[7], Kenneth DR. Setchell[2,7], Ahmet Kaynak[8], Xiaoyang Qi[2,8], Ricardo A. Feldman[9], Ying Sun[1,2*].

[1] Division of Human Genetics, Cincinnati Children's Hospital Medical Center, Cincinnati, OH

[2] Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH

[3] Pluripotent Stem Cell Facility & Developmental Biology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH

[4] Division of Infectious Diseases, Cincinnati Children's Hospital Medical Center, Cincinnati, OH

[5] Transgenic Animal and Genome Editing Facility, Cincinnati Children's Hospital Medical Center, Cincinnati, OH

[6] Division of Developmental Biology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH

[7] Division of Pathology and Laboratory Medicine, Cincinnati Children's Hospital Medical Center, Cincinnati, OH

[8] Division of Hematology/Oncology, Department of Internal Medicine, University of Cincinnati, College of Medicine, Cincinnati, OH

[9] Department of Microbiology and Immunology, University of Maryland School of Medicine, Baltimore, MD

25   *Corresponding author. Division of Human Genetics, Cincinnati Children's Hospital Medical

26   Center, S Building Rm. 4.216, 3333 Burnet Ave, Cincinnati, OH 45229, United States. E-mail:

27   Ying.Sun@cchmc.org

Abstract

Neuronopathic Gaucher disease (nGD) is a lysosomal storage disorder caused by *GBA1* mutations, leading to defective acid β-glucosidase (GCase) and accumulation of glycosphingolipid substrates, causing inflammation and neurodegeneration. Patients with nGD manifests severe neurological symptoms, but current animal models fail to fully recapitulate human condition, posing a major barrier to the development of effective therapies targeting the brain. To bridge this gap, we have developed midbrain-like organoids (MLOs) from human induced pluripotent stem cells (hiPSCs) of nGD patients with $GBA1^{L444P/P415R}$ and $GBA1^{L444P/RecNcil}$ mutations to model nGD brain pathogenesis. These nGD MLOs exhibited GCase deficiency, resulting in diminished enzymatic function, accumulation of lipid substrates, widespread transcriptomic changes, and impaired dopaminergic neuron differentiation, mirroring nGD pathology. *GBA1* mutation correction mediated by CRISPR/Cas9 restored GCase activity, normalized lipid substrate levels, and rescued dopaminergic neuron function, confirming the causal role of *GBA1* mutations during early brain development. Using this novel platform, we further evaluated therapeutic strategies, including SapC-DOPS nanovesicles delivering GCase, AAV9-GBA1 gene therapy, and substrate reduction therapy with GZ452, a glucosylceramide synthase inhibitor currently under clinical investigation. These treatments either restored GCase activity, reduced lipid substrate accumulation, improved autophagic and lysosomal abnormalities, or ameliorated dysregulated genes involved in neural development. These patient-specific, 3D neural models offer a transformative, physiologically relevant platform for unravelling disease mechanisms and accelerating the discovery of therapies for patients with nGD.

Introduction

Gaucher disease (GD) is a lysosomal storage disorder caused by *GBA1* mutations, which impair acid β-glucosidase (GCase). Defective GCase leads to accumulation of its glycosphingolipid substrates, glucosylceramide (GluCer) and glucosylsphingosine (GluSph), triggering inflammation and neurodegeneration. GD affects 1/450 in Ashkenazi individuals and 1/50,000 globally [1]. There are three GD types: Type 1 present mainly visceral symptoms and is not life-threatening [1]. while neuronopathic GD (nGD, Types 2 and 3) involves severe neurological manifestations. Type 2 patients typically die by age 2, indicating early central nervous system (CNS) defects [2,3].

Current knowledge of brain pathogenesis in nGD is limited to a few postmortem brain analyses showing prominent neuronal loss and astrogliosis in the midbrain [4-6]. GD mouse models, including GCase knockout, irreversible GCase inhibitor conduritol B-epoxide (CBE) induced mouse models, double transgenic mutant model harboring both *Gba1* and Saposin C mutations (Saposin C is the activator of GCase), have been widely used to study disease pathogenesis and therapies [7-11]. These models replicate neuronal loss and astrogliosis [4-6,8,12], but have limitations. The knockout and CBE-induced models lack *GBA1* mutations, and the double transgenic model has a second gene that may change the GD phenotype, and knock-in models fail to mimic human phenotypes [10,13]. For example, mice with the *Gba1*[L444P/L444P] or *Gba1*[D409H/D409H] mutations show no neuronopathic disease despite <10% of residual GCase activity [10,13-15]. These constraints highlight the need for physiologically relevant human nGD models.

Brain organoids derived from human induced pluripotent stem cells (hiPSC) emerged as a

powerful 3D model to study brain development and CNS diseases [16]. Unlike 2D monolayer

neural cells, brain organoids contain diverse neural and glial cell types arranged in 3D

comprising subventricular zone and multiple organized cellular layers that more closely

represents the spatial architecture of the human brain [17]. They have been used to study

neurodegenerative and lysosomal diseases, including Parkinson's disease (PD), Alzheimer's

disease, and Sandhoff disease [18-22]. A brain organoid derived from hiPSCs of a healthy individual

with *GBA1* knockout and α-synuclein overexpression exhibited some PD features [23]. However,

this model lacks patient-specific *GBA1* mutations needed for clinical relevance. To develop a

relevant model, we developed novel midbrain organoids from nGD patients with *GBA1*

mutations, a model not yet well studied.

The midbrain region is prominently affected in nGD [4-6]. To model this, we derived midbrain-like

organoids (MLOs) from Type 2 patient-derived hiPSCs carrying either $GBA1^{L444P/P415R}$ or

$GBA1^{L444P/RecNcil}$ mutations. These GD MLOs showed reduced GCase activity, lipid substrate

accumulation, transcriptomic alterations, and impaired dopaminergic neuron differentiation.

CRISPR/Cas9-mediated correction of *GBA1* mutation confirmed the causal role of *GBA1*

mutations in these phenotypes and mitigated disease phenotypes. To explore therapeutic potential,

we tested three emerging treatments in GD MLOs: CNS accessible enzyme therapy

(SapC-DOPS-GCase), gene therapy (AAV9-GBA1) and substrate reduction therapy (GZ452).

Our results demonstrate the utility of patient-derived MLOs for studying nGD pathogenesis and

advance drug development.

Materials and Methods

Maintenance of hiPSCs and generation of MLOs

Three hiPSCs lines (WT-75.1, GD2-1260 and GD2-10-257) generated previously [24,25] (Supplementary Table 1) were maintained in mTeSR1 complete medium (StemCell) on six-well plates coated with Vitronectin (1:100 diluted in DPBS, 1 mL/well, incubated at RT for 1 hour). HiPSCs at ~70% confluency were treated and replated at 1:20 to 1:60 ratios, with optional 10 µM ROCK inhibitor (Y-27632, StemCell) for survival. All hiPSC lines were authenticated using short tandem repeat (STR) profiling and were verified as mycoplasma-free.

MLO generation method was modified based on published protocol [26,27]. Briefly, hiPSCs were pretreated with 50 µM Y-27632 for 1 hour, dissociated with Accutase (Millipore-Sigma) into singlets, centrifuged ($200 \times$ g, 5 minutes), and resuspended in embryoid body (EB) medium (EBM; consisting of DMEM/F12, 20% KnockOut Serum Replacement (KSR), 1% penicillin/streptomycin, GlutaMAX, NEAA, 55 µM β-mercaptoethanol, 1 µg/mL heparin, 3% FBS, 4 ng/mL bFGF, $1\times$ CEPT). For each organoid, $1.2\sim1.5 \times 10^4$ hiPSC cells were seeded in 100 µL EBM per well in ultra-low attachment (ULA) U-bottom 96-well plates, centrifuged at $500 \times$ g for 3 minutes, and incubated at 37°C, 5% CO_2; on day 2, 150 µL EBM with 4 ng/mL bFGF (no ROCK inhibitor) was added. Once the EBs grow to 400 to 475 µm in size, they were transferred to a new ULA 96-well plate with 125 µL brain organoid generation medium (BGM, consisting of 50% DMEM/F12, 50% Neurobasal, $1\times$ N2, $1\times$ B27 without vitamin A, 1% penicillin/streptomycin, 1% GlutaMAX, 1% NEAA, 55 µM β-mercaptoethanol, 1 µg/mL heparin) supplemented with dual SMAD inhibitors (SMADi: 2 µM dorsomorphin, 2 µM A83-01, 3 µM CHIR99021, 1 µM IWP2), refreshed every other day until day 11. On day 7 or 8, mesencephalic

floor plate induction was initiated by adding 125 μL BGM containing 1× SMADi, 100 ng/mL

FGF8, and 2 μM SAG until day 16. On day 11, four EBs were transferred to one well of

ultra-low attachment 24-well plates with 500 μL BGM-MLOs induction medium, consisting of

BGM with 100 ng/mL FGF8, 2 μM SAG, 200 ng/mL laminin, 2.5 μg/mL insulin, 2% Growth

Factor Reduced (GFR) Matrigel, and refreshed with Matrigel on day 14. On day 16, three MLOs

per well were transferred to ultra-low attachment 6-well plates in brain organoid maturation

medium (BGM-BMM), consisting of BGM with 10 ng/mL BDNF, 10 ng/mL GDNF, 200 μM

ascorbic acid, 125 μM db-cAMP, 1% GFR Matrigel, cultured on an orbital shaker (100 rpm) to

reduce spontaneous fusion with medium changes every 3 days. Starting at day 30, GFR Matrigel

was removed, and brain organoids were cultured solely in BGM medium till the date of analysis.

Immunostaining and image analysis of sectioned MLOs

Organoids were washed with 1xDPBS and fixed in 4% paraformaldehyde (PFA) overnight at

4°C. After washing with PBS-T (1 x DPBS with 0.1% Tween 20), organoids were cryoprotected

in 30% sucrose solution until fully equilibrated. Samples were then embedded in gelatin solution

(7.5% gelatin, 10% sucrose in 1xDPBS) at 37°C for 1 hour, snap-frozen in a dry ice/ethanol

slurry, and stored at -80°C. Cryosectioning was performed to obtain 16-μm-thick sections. For

immunofluorescence, sections were air-dried and outlined with a hydrophobic PAP pen before

blocked with 5% normal goat serum in PBS-T for 1 hour, incubated overnight with primary

antibodies (Supplementary Table 1) in primary antibody dilution buffer (PBS-T with 5% BSA

and 0.05% sodium azide), and followed by secondary antibodies (Alexa Fluor® conjugates) for 1

hour at room temperature. Slides were washed, stained with DAPI solution (0.2 μg/mL),

mounted with Antifade Mounting Medium (VECTASHIELD®, H-1000-10), and imaged at 10×

magnification for whole organoid scans or 60× magnification for detailed cellular analysis using

a fluorescence confocal microscope (Nikon Eclipse *Ti*) (Nikon, Tokyo, Japan) to assess neuronal

organization of MLOs. The morphologies of neurons, astrocytes, and colocalization analysis

were performed by taking multi-layer stacking (Z-stack) images under 60× oil immersion

objective lens. For quantification of immunoreactive cells, six images for each group of MLOs

were binarized by applying the automatic threshold and counted using ImageJ (NIH).

Genome editing patient iPSC clones

The iPSC line (GD2-1260) used for gene correction was derived from fibroblasts obtained from

a GD Type 2 patient carrying compound heterozygous *GBA1* mutations (P415R/L444P) [24]. A

guide (g) RNA was designed to target SpCas9-mediated double strand break introduction

proximal to the L444P mutation (*GBA1* mutation at nt14446, T>C) in GD2-1260.

Oligonucleotides   (caccGAAGAACGACCCGGACGCAG   and

aaacCTGCGTCCGGGTCGTTCTTC; overhangs in lower case; target mutation site underlined;

IDT) transcribing the sgRNA target sequence were annealed and cloned via BbsI restriction

digest into plasmid pX458M that contains a U6 promoter-driven sgRNA and a SpCas9-2A-EGFP

expression cassette. The pX458M plasmid is modified from the pX458 plasmid (addgene #48138)

[28] and carries an optimized sgRNA scaffold [29]. The targeting activity of this plasmid was

validated by T7E1 assay using 293 cells. GD2-1260 cells were transfected with pX458M and a

phosphorothioate-modified single stranded oligonucleotide (ssODN; IDT) donor template using

TransIT-LT1 (Mirus). The ssODN (Supplementary Figure 1A) was designed to introduce the

desired wildtype *GBA1* sequence flanked by homology arms to the targeted genomic region. The

ssODN was also designed to contain silent mutations to prevent retargeting by SpCas9 and to

introduce a BtgI restriction site to facilitate identification of targeted clones. Forty-eight hours

post-transfection, GFP-positive cells were isolated by FACS and replated at cloning density

(250-500 cells/well of a 6 well matrigel-coated plate). Replated cells were cultured for 4 days in

mTeSR1 containing 10% CloneR (StemCell Technologies), using the manufacturer's

recommended protocol. Cells were subsequently fed daily with mTeSR1 for an additional 9 days

before colonies were manually harvested and expanded for genotyping. Primers VS4247

(gtgcgtaactttgtcgacagtcc) and VS4249 (ctgagagtgtgatcctgccaag) were used to PCR amplify the

targeted *GBA1* genomic region and products were subjected to BtgI digestion to identify putative

edited clones. Selected clones were subsequently confirmed by Sanger sequencing.

SapC-DOPS-fGCase treatment of MLOs

As previously described [30,31], SapC-DOPS nanovesicles were formulated by combining saposin

C (SapC) with dioleoylphosphatidylserine (DOPS) to encapsulate either recombinant human acid

β-glucosidase (fGCase, Freeline Therapeutics) or fluorescent CellVue Maroon (CVM) for uptake

studies. The SapC-DOPS-fGCase complex was prepared at a final concentration of 0.6 μg/mL

fGCase and added to the BGM culture medium of MLOs derived from WT-75.1, GD2-1260, and

GD2-10-257 hiPSC lines. To assess nanovesicle uptake, WT-75.1 MLOs at week 13 were

co-cultured with SapC-DOPS-CVM for 48 hours, followed by confocal imaging to confirm

internalization of the fluorescent CVM within the organoids.

For enzyme delivery and therapeutic evaluation, MLOs were treated with SapC-DOPS-fGCase

or SapC-DOPS alone (control) for a 48-hour period to assess uptake and enzyme activity. For

long-term treatment efficacy, MLOs were treated for two weeks before evaluating therapeutic

efficacy. During treatment, MLOs were maintained in BGM media containing nanovesicles and

were replaced every 3 days to ensure consistent exposure. Post-treatment, MLOs were harvested

for confocal imaging, enzymatic assays, and biochemical analyses to assess GCase activity,

protein expression, and correction of GD phenotypes.

AAV injection into MLOs

Transfer vector of AAV9-GBA1 virus containing GBA1/GFP expressing cassette

(CB-GBA1-IRES-GFP) was packaged in AAV9 capsid at AAVnerGene (Rockville, MD, USA).

For the delivery of AAV9-GBA1 gene therapy to MLOs, a precise injection protocol was

employed using a nanoliter injector system (World Precision Instruments). MLOs derived from

WT-75.1, GD2-1260, and GD2-10-257 hiPSC lines were placed in a dish containing sterile 1 x

PBS. A glass pipette connected to a gas injector was lowered into the center of each MLO at a

desired depth of 200 to 300 μm, as determined by microscopic visualization. A volume of 500 nL

(0.5μL) of AAV9-GBA1 vector, at a concentration of 3.6×10^{13} vg/mL, was injected into ten

sites of MLOs at a volume of 50 nL per injection site. The injection glass needle was retracted

slowly after completion of the injection to minimize damage and ensure vector distribution.

Morphology and viability were assessed post-injection, and MLOs were then maintained in

culture for 3 weeks prior to subsequent analyses.

Substrate Reduction Therapy treatment of MLOs

For evaluation of the tolerated dose of Substrate Reduction Therapy (SRT), WT-75.1 MLOs were

treated with SRT compound GZ452 (AstaTech, P14969), also named GZ-682452, an analogue of

venglustat [32], at concentrations of 0.3, 1, 2, and 3 μM starting on day 0 for a period of 6 weeks.

MLO size was measured weekly using bright-field microscopy and analyzed with ImageJ

software to assess morphological changes. For short-term SRT treatment, 13-week-old

GD2-1260 MLO were treated with 300nM GZ452 for 3 weeks, with medium refreshed every 3

days to maintain consistent drug exposure. For long-term SRT treatment, GZ452 treatment at the

concentration of 300nM was started at the beginning of MLO generation (day 1) and continued

for 28 weeks before collection for analysis. Untreated MLOs served as controls.

Glycosphingolipids analyses

MLOs were washed with ice-cold PBS and homogenized and sonicated in sterile ddH_2O.

Glycosphingolipids were then extracted from tissue homogenates using chloroform/methanol

following protocol outlined previously [33]. Aliquots of lipids extracts were processed for GluCer

and GluSph quantification by ultra-high performance chromatography coupled to tandem mass

spectrometry (UHPLC-MS/MS) using a Waters Xevo TQ-S Micro triple quadrupole mass

spectrometer (Waters, Milford, MA) at CCHMC Clinical Mass Spectrometry Laboratory.

Chromatographic separation for GluCer and GluSph was achieved using a XSelect CSH C18 XP

Column (100 x 2.1 mm, 2.5 µm, Waters) column. Quantification by LC-MS/MS was operated in

the multiple reaction monitoring (MRM) mode, with detection of the transition pair of the

individual protonated parent ions of GluCers and daughter ion m/z 264.2. GluSph was measured

by monitoring the mass transition m/z 462.3> 282.4. Calibration curves were prepared for C16

GluCer, C18 GluCer, C24 GluCer, and C24:1 GluCer using C18 glucosyl(ß) ceramide-D5 as the

internal standard. Quantification of GluCer species with various fatty acid chain lengths was

realized by the calibration curve of each species or with the closest fatty acyl chain length. The

quantification of GluSph was based on the calibration curve using glucosyl(ß) sphingosine-D5 as

internal standard. The calibration curve for GluCer and GluSph was 25 pg–10 ng on column. The

GluCer and GluSph levels in MLO were normalized to mg of cell protein that was determined by

BCA assay and expressed as pmol/mg protein. Additionally, GluSph levels in the culture medium

were quantified and normalized to the medium volume (pmol/mL) [34].

Measuring Dopamine levels by ELISA

Dopamine levels in MLO culture medium were quantified using the Dopamine ELISA Kit

(Abnova) that involves dopamine extraction, acylation and enzymatic conversion before assay.

Briefly, MLO culture medium was collected from 4 MLOs cultured in 3 mL BGM medium for

72 hours. Dopamine extraction was performed by pipetting 10 µL of standards, controls, and 750

µL of the sample into the extraction plate wells, filling each to 750 µL with deionized water,

followed by 25 µL of TE buffer. Shake the covered plate for 60 minutes at room temperature (RT,

20-25°C) at 600 rpm, wash with 1 mL wash buffer twice, then acylate with 150 µL acylation

buffer and 25 µL acylation reagent for 20 minutes. After washing again, elute with 100 µL

hydrochloric acid, and transfer 90 µL of supernatant to the microtiter plate for enzymatic

conversion with 25 µL freshly prepared enzyme solution, incubating at 37°C for 2 hours. Then,

100 µL of the converted samples and 50 µL dopamine antiserum were added to the dopamine

microtiter strips. The mixture was then incubated overnight at 2-8°C. The strips were then

washed and incubated with 100 µL enzyme conjugate for 30 minutes, followed by 100 µL of

substrate for 20-30 minutes, and finally, 100 µL of stop solution was added. Absorbance was

measured at 450 nm within 10 minutes, and dopamine concentrations were calculated using a

calibration curve, applying the correction factor (10/500 = 0.02) to adjust and normalize by

sample volume.

GCase activity assay

GCase enzyme activities in hiPSCs and MLOs were determined fluorometrically with 4-methylumberlliferyl-β-D-glucopyranoside (4MU-Glc) in the presence of the GCase irreversible inhibitor, Conduritol B epoxide (CBE, 2 mM, Millipore, Bedford, MD) as previously described [35]. Briefly, hiPSC cell pellet or MLO tissues were homogenized using Precellys Evo Homogenizer/CK Mix beads (Bertin Technologies, France) in 1% sodium taurocholate/1%Triton X-100 (Tc/Tx) solution. GCase enzyme activity was then determined by 4MU-Glc as substrate in 0.25% Tc/Tx diluted in 0.1 M citrate phosphate buffer (pH 5.6). Protein concentrations were determined by BCA assay for normalization. The GCase-specific activity was calculated by subtracting non-specific activity (with CBE) from total activity (without CBE) and normalized to total protein mass.

Transcriptome analysis of MLOs

Total RNAs were extracted from week 8 MLOs using the RNeasy Mini Kit (Qiagen). Total RNAs (150 to 300 ng), quantified by Qubit (Invitrogen) high-sensitivity spectrofluorometric assay, were poly-A selected and reverse transcribed using Illumina's TruSeq® stranded mRNA library preparation kit (Illumina). Each sample was fitted with one of 96 adapters containing different 8-base molecular barcodes for high-level multiplexing. After 15 cycles of PCR amplification, completed libraries were sequenced on NovaSeq 6000 (Illumina), generating 30 million high-quality 100-base-long paired-end reads per sample. A quality control check on the fastq files was performed using FastQC. Upon passing basic quality metrics, the reads were trimmed to remove adapters and low-quality reads using default parameters in Trimmomatic

(Version 0.33). The trimmed reads were then mapped to human reference genome GRCh38 using

default parameters with strandness option in Hisat2 (Version 2.0.5). Transcript or gene

abundance was determined using kallisto (Version 0.43.1). A transcriptome index in kallisto was

created using Ensembl cDNA sequences for the reference genome. This index was then used to

quantify transcript abundance in raw counts and transcript per million (TPM). Differential

expression analysis was performed using the R package DESeq2 [36]. Genes with BaseMean $\geq$ 50,

|fold change| $\geq$ 2, p-adj $\leq$ 0.05 were identified as differentially expressed genes (DEGs).

Heatmaps were made using the R package ggplot2. Gene enrichment analysis using cellular

component, KEGG pathways, molecular function, and biological process (GO analysis) was

performed using DAVID 6.8 [37]. Three MLOs were pooled for each sample and three samples

were profiled for each genotype. MLO Sequencing datasets and partially processed results have

been deposited to the National Center for Biotechnology Information (NCBI)'s Gene Expression

Omnibus (GEO) database (GSE303993).

Gene expression analysis by qRT-PCR

Total RNA was isolated from MLO tissue using the RNeasy Kit (Qiagen). Quantitative RT-PCR

was carried out as described previously using primers listed in Supplementary Table 1 [38].

Housekeeping gene *GAPDH* or *ACTB* was used as internal controls for mRNA quantification.

Relative expression of mRNAs was determined by the $2^{-\Delta\Delta CT}$ method.

Immunoblotting

Immunoblotting was used to detect protein expression in iPSCs and MLOs. Briefly, brain

organoids or cells were homogenized in freshly prepared RIPA buffer with protease inhibitors

and anti-phosphatase inhibitors using a Precellys Evo Homogenizer (Bertin). The samples were

then centrifuged at 12,000 g for 10 min to produce lysate for electrophoresis, membrane

transferring, and detection using primary and secondary antibodies indicated in Supplementary

Table 1. Semi-quantification of proteins was performed using ImageJ (NIH).

Statistical analysis

For comparison of two or more groups, analysis was performed by unpaired Student's t-tests

with normal distribution or one-way ANOVA followed by Tukey's or Bonferroni's multiple

comparison, according to the sample size. Exclusion of outliers was made based on cutoffs of the

mean ± 2 standard deviations. All statistical analyses were performed using GraphPad Prism 10

Software. $P < 0.05$ was considered statistically significant.

Results

Generation and characterization of MLOs derived from hiPSCs

To generate MLOs from iPSCs, healthy hiPSCs were differentiated following a stepwise protocol,

neural induction, neuroepithelial expansion, and maturation, adopted with modifications from

previous studies [26,27] (Figure 1A). Midbrain specification during development requires

appropriate posterior differentiation signals [39,40]. Our protocol used sequential

activation/inhibition of critical signaling pathways involved in midbrain specification: Wnt

signaling (activated by CHIR99021), BMP signaling (inhibited by dual-SMAD), and FGF

signaling (activated by FGF8) (Figure 1A). MLOs were matured and characterized using

immunofluorescence, gene expression, and immunoblotting. Week 8 MLOs showed diverse

neural cell types, including pan-neurons (Tuj1/NeuN), astrocytes (GFAP), dopaminergic (DA)

neurons (FOXA2/TH), and neural progenitor cells (SOX2/Ki67) (Figure 1B). Comparative

analysis of forebrain marker FOXG1 expression between MLOs and cerebral organoids (COs)

showed barely detectable FOXG1 in MLOs (Figure 1C). Quantitative RT-PCR analysis

demonstrated a significant increase in midbrain/DA neuron-specific genes (*FOXA2*, *ASCL1*,

LMX1A, *PLZF*, and *TH*) from week 3 and week 8 MLOs, demonstrating the midbrain identity of

these organoids. Expression of glial-specific genes (*GLAST*, *S100B*) increased over time, while

pluripotency markers (*SOX2*, *NANOG*, *OCT4*) were downregulated, suggesting progressive

neural differentiation (Figure 1D). Immunoblot further validated hiPSCs differentiation into

MLOs by expression of neuronal (Tuj1, MAP2), DA neuronal (TH), and glial (GFAP, S100B)

markers at week 8, with reduced Sox2 expression in MLOs versus hiPSCs (Figure 1E). Together,

these results confirm the successful generation of MLOs with midbrain-like cellular composition

and molecular identities.

GCase deficiency in GD MLOs leads to glycosphingolipids accumulation and altered transcriptomic profiling

To model nGD and investigate the impact of GCase deficiency on MLOs, we generated GD MLO using the GD2-1260 hiPSC line derived from GD Type 2 patient harboring $GBA1^{L444P/P415R}$, compound heterozygous mutations, using same protocol in Figure 1 (Figure 2) [24]. Midbrain identity of GD MLOs was confirmed by TH/FOXA2 expression (Figure 3). Immunoblot showed an approximately 85% reduction in GCase protein levels in week 8 GD2-1260 MLOs versus WT-75.1 MLOs (Figure 2A). Consistent with reduced GCase protein, GCase activity decreased to 14.2% in GD2-1260 hiPSCs and 15.0% in GD2-1260 MLOs vs. WT-75.1 ($p < 0.001$; Figure 2B). Bright-field imaging at weeks 4, 8, and 15 showed no difference on overall MLOs appearance between WT-75.1 and GD2-1260 MLOs (Figure 2C). MLO size measurements revealed a trend toward smaller GD2-1260 MLOs versus WT-75.1 at week 8, though not statistically significant (Figure 2D).

Loss of GCase function in GD disrupts glycosphingolipids metabolism, a primary pathogenic factor in nGD [41]. Mass spectrometry analysis demonstrated significant glycosphingolipid substrate accumulation in GD2-1260 MLOs. At week 15, total glucosylceramide (GluCer) level was elevated approximately 1.39-fold in GD2-1260 MLOs compared to WT-75.1 MLOs ($p <$ 0.05; Figure 2E). GluCer analysis revealed marked accumulation of brain enrich species (18:0 and 16:0) in GD2-1260 MLOs, while other species remained largely unchanged (Figure 2F). Similarly, glucosylsphingosine (GluSph), a toxic lipid associated with nGD [4,32,42], was significantly elevated approximately 3.3-fold and 5.6-fold in GD2-1260 MLOs at weeks 15 and 28, respectively, versus WT-75.1 controls (Figure 2G). These findings indicate profound lipid

substrates dysregulation in GCase-deficient GD MLOs, consistent with the biochemical hallmark

of GD.

To explore the transcriptomic consequences of GCase deficiency, we performed bulk RNA

sequencing on week 8 MLOs, when DA neurons maturation and GD phenotypes were evident.

Principal component analysis (PCA) revealed distinct clustering of WT-75.1 and GD2-1260

MLOs, with the first principal component accounting for 53% of the variance, indicating

significant transcriptomic differences between two genotype groups (Figure 2H). An MA plot

identified 1,429 differentially expressed genes (DEGs) in GD2-1260 MLOs compared to

WT-75.1, with 664 genes upregulated and 765 genes downregulated. GO analysis of these DEGs

revealed significant enrichment in pathways including cAMP, PI3K-AKT, and WNT signaling

that control the nervous system development, axon guidance, and neuron differentiation (Figure

2J). KEGG analysis similarly identified dysregulation in neural signaling pathways, synaptic

transmission, and lysosomal function (Figure 2K). The heatmap of these DEGs elucidated

specific gene expression changes within key pathways (Figure 2L-2O), including WNT signaling

(Figure 2L) [25], anterior-posterior brain specification (Figure 2M), neuronal function (Figure 2N),

and lysosome-phagosome (Figure 2O) in GD2-1260 MLOs, with many genes exhibiting

upregulation or downregulation consistent with nGD pathology.

Collectively, these results demonstrate that *GBA1* mutation in GD MLOs leads to reduced

enzyme activity, causes lipid substrate accumulation, and alters transcriptomic profiles,

particularly in neural development, signaling, and lysosomal function. These findings provide

insights into nGD mechanism and support MLOs as a model for studying nGD.

Skewed specification of midbrain patterning and dopaminergic neuron differentiation in GD MLOs

RNA sequencing data revealed elevated FOXG1 and PAX6, and reduced FOXP1 mRNAs in

GD2-1260 MLOs versus WT-75.1 MLOs at week 8 (Figure 2M, Figure 3A). These genes are

fate-determining regulators in controlling proper anterior-posterior neural patterning [43-45].

Immunofluorescence confirmed reduced FOXP1+ and increased FOXG1+ cell counts in

GD2-1260 MLOs (Figure 3B, 3C). About 11.6% cells were FOXP1+ FOXG1+ cells in

GD2-1260 MLOs, but they were barely seen in WT-75.1 MLOs. These results indicate a

significant disruption in expression of genes involved in forebrain and midbrain patterning in GD

MLOs.

The skewed specification of midbrain patterning may contribute to dysregulated dopaminergic

neuron development in GCase-deficient GD MLOs (Figure 3). There were clear differences in

the expression of midbrain-specific markers between WT-75.1 and GD2-1260 organoids at week

6 (Figure 3D). In WT-75.1 MLOs, FOXA2 (a midbrain progenitor marker) and TH

(dopaminergic neuron marker) were strongly expressed and co-localized (Figure 3D, 3E),

indicating normal midbrain patterning. In contrast, GD2-1260 MLOs showed markedly reduced

FOXA2+, TH+ and FOXA2+TH+ cells (Figure 3D, 3E). Quantitative analysis confirmed the

significant reduction of midbrain progenitor markers (*ASCL1*, *TH*, *FOXA2*, and *PLZF*) in

GD2-1260 MLOs versus WT-75.1 (Figure 3F). At week 8, expression levels of *ASCL1*, *TH*,

FOXA2, and *PLZF* were significantly reduced by approximately 71.0%, 44.4%, 90.0% and

69.8%, respectively, indicating a profound impairment in midbrain progenitor specification due

to GCase deficiency. Immunoblot showed ~77.5% and ~66.5% reductions in TH and FOXA2,

respectively, and a 27.7% increase in MAP2 in GD2-1260 versus WT-75.1 MLOs (Figure 3H).

Dopaminergic function was determined by measuring dopamine levels in the culture medium

(Figure 3I). Dopamine levels in week 12 MLOs culture medium (72 hours culture) were 76.0%

(p < 0.001) lower in GD2-1260 MLOs versus WT-75.1 MLOs, confirming impaired

dopaminergic function in GD MLOs.

These results indicate that GCase deficiency leads to skewed midbrain patterning and

dopaminergic neuron differentiation, underscoring the critical role of GCase in midbrain

development, as revealed for the first time, using patient iPSC-derived brain organoids.

CRISPR/Cas9-mediated *GBA1* mutation correction rescues disease phenotypes in GD

MLOs

GD is an autosomal recessive disorder. Individuals who are carriers or heterozygous for *GBA1*

mutations typically do not exhibit disease phenotypes [1]. To determine whether correction of the

GBA1 mutation could rescue GD phenotypes in GD MLOs, we used CRISPR/Cas9 technology

to correct the L444P mutation in GD2-1260 hiPSCs (*GBA1*$^{L444P/P415R}$) and generated isogenic

iso-GD2-1260 hiPSCs with a heterozygous genotype of *GBA1*$^{WT/P415R}$ (Figure 4A;

Supplementary Figure 1). Karyotype analysis of GD2-1260 and iso-GD2-1260 demonstrated no

detectable chromosomal abnormalities, confirming the genetic stability of both cell lines

(Supplementary Figure 1D). Furthermore, genetic correction did not alter the expression of

hiPSC related pluripotent genes (Supplementary Figure 1E). The MLOs derived from WT-75.1,

GD2-1260, and iso-GD2-1260 hiPSCs were then compared to evaluating the impact of mutation

correction on GD phenotypes.

At week 8, GCase protein in iso-GD2-1260 MLOs was restored to 45.8% of WT-75.1 levels

(Figure 4B, 4C), consistent with correction of one *GBA1* allele [24]. GCase activity also improved,

reaching 43.9% and 46.4% of WT-75.1 levels in GD2-1260 hiPSCs and MLOs, respectively,

versus approximate 15% in GD2-1260 hiPSCs and MLOs (Figure 4D). Despite reduced GCase,

early neural rosette formation in GD2-1260 MLOs remained largely unaffected in GD2-1260 and

iso-GD2-1260, with similar counts of SOX2+Ki67+ proliferating neural progenitor cells in

neural rosettes (Supplementary Figure 2A, 2B), suggesting early neural development is not

impacted by *GBA1* L444P mutation.

GBA1 L444P mutation correction ameliorated lipid substrate accumulation in iso-GD2-1260

MLOs. GluSph levels which were elevated in GD2-1260 MLOs at weeks 15 and 28

(approximately 3.3-fold and 5.6-fold higher than WT-75.1, respectively; $p < 0.01$) was

normalized to WT-75.1 levels (Figure 4E). Organoid size of iso-GD2-1260 MLOs was similar to

GD2-1260 MLOs, except slightly decrease at week 8 (Figure 4F, 4G).

Correction of the L444P mutation restored midbrain and dopaminergic neuron differentiation.

TH and FOXA2 levels, which were reduced in week 16 GD2-1260 MLOs by approximately 76.4%

and 72.4%, respectively, were restored in iso-GD2-1260 MLOs to WT-75.1 levels for TH, and

approximately 53% of WT-75.1 levels for FOXA2 (Figure 4H, 4I). Dopamine levels in

iso-GD2-1260 MLOs culture matched those in WT-75.1, unlike the reduced levels in GD2-1260

MLOs (Figure 4J), indicating restored dopaminergic function. The autophagy-lysosomal pathway,

which is dysregulated in nGD, was partially corrected by *GBA1* mutation restoration (Figure 4K,

4L). Immunoblot analysis of lysosomal proteins showed that LAMP1 levels, which were

significantly decreased in GD2-1260 MLOs by approximately 42.5% versus WT-75.1, were

partially restored in iso-GD2-1260 MLOs (Figure 4K, 4L), however, Cathepsin D levels

remained unchanged (Figure 4K, 4L). Dysregulated autophagy flux demonstrated by the elevated

LC3-II and LC3-I was not significantly improved in isogenic MLOs compared to GD2-1260

MLOs (Figure 4M, 4N).

GCase deficiency leads to mTOR hyperactivation in nGD [12,46,47]. In GD2-1260 MLOs, levels of

4E-BP1 and its phosphorylated form P-4E-BP1 (Thr37/46) were significantly elevated (Figure

4O). These levels were normalized or significantly reduced in iso-GD2-1260 MLOs. While S6

and its phosphorylated form (P-Ser235/236) remain unchanged (Figure 4O, 4P).

These results demonstrate that CRISPR/Cas9-mediated correction of *GBA1* mutation in

GD2-1260 hiPSCs effectively rescues key nGD phenotypes and downstream effects, highlighting

gene correction as a therapeutic strategy and validating MLOs as a preclinical disease model.

SapC-DOPS nanoparticle-mediated GCase enzyme therapy corrects GD phenotypes in GD

MLOs

SapC-DOPS nanoparticles, composed of saposin C (SapC) and dioleoylphosphatidylserine

(DOPS), showed promise as a CNS-targeted delivery system for lysosomal disorders like nGD

[30,31]. The effectiveness of this approach in rescuing GD phenotypes was evaluated in MLOs

derived from GD2-1260 hiPSCs and another GD2 hiPSC line, GD2-10-257, which carries the

$GBA1^{L444P/RecNcil}$ mutation [48]. MLOs were derived from these hiPSCs following the protocol in

Figure 1 and described in the Methods section. SapC-DOPS nanoparticle was formulated with a

fGCase, a recombinant GCase variant with over 21-fold longer active half-life at lysosomal pH

than wild-type GCase [49,50] (Figure 5A).

To test SapC-DOPS uptake, WT-75.1 MLOs were co-cultured with CVM (CellVue Maroon)

loaded SapC-DOPS nanoparticles for 48 hours. CVM signals were detected throughout the

organoids, confirming successful cargo delivery (Figure 5B). SapC-DOPS-fGCase was then

prepared with 0.6 μg/mL fGCase [30] and cocultured with MLOs. After 48 hours, treatment

significantly increased GCase activity in GD2-1260 MLOs, restoring it to approximately 66.7%

of WT-75.1 levels, confirming efficient delivery (Figure 5C).

To evaluate efficacy, GD2-1260 and GD2-10-257 MLOs were treated for two weeks with

SapC-DOPS or SapC-DOPS-fGCase. Confocal imaging confirmed restored GCase expression,

with SapC-DOPS-fGCase-treated GD2-1260 and GD2-10-257 MLOs showing increased GCase

approaching the level in WT-75.1 MLO (Figure 5D). Additionally, SapC-DOPS-fGCase

treatment significantly elevated GCase load in both dopaminergic neurons (TH+) and astrocytes

(GFAP+), as shown by colocalization of GCase with TH and GFAP in GD2-1260 and

GD2-10-257 MLOs, respectively (Supplementary Figure 3A-D). Corresponding to increased

GCase protein, SapC-DOPS-fGCase treatment elevated GCase activity to 2.3-fold of untreated

WT-75.1 MLO (Figure 5E). In GD2-1260 MLOs, SapC-DOPS-fGCase fully restored GCase

activity to WT levels, while SapC-DOPS alone had no effect (Figure 5F). Similar GCase

restoration was observed in the additional patient organoids, GD2-10-257 MLOs, indicating the

effectiveness of this approach across different GD MLO models with various *GBA1* mutations

(Figure 5G). Importantly, SapC-DOPS-fGCase significantly reduced elevated GluSph levels in

GD2-1260 and GD2-10-257 MLOs, matching WT-75.1 levels (Figure 5H, 5I), confirming

effective lipid substrate clearance by delivered fGCase.

Furthermore, we evaluated the impact of SapC-DOPS-fGCase on the autophagy and lysosomal

pathway and mTOR signaling. fGCase was found effectively delivered to lysosomal

compartments, evidenced by colocalization of LAMP1 and LC3-II in treated GD2-1260 and

GD2-10-257 MLOs (Supplementary Figure 4A-D), indicating restoration of GCase in lysosomal

and autophagosome compartments. LAMP1 level was decreased in GD2-1260 MLOs and

partially restored following SapC-DOPS-fGCase treatment (Figure 5J). High levels of LC3-II,

indicator of autophagy flux, were reduced in treated GD2-1260 MLOs, suggesting enhanced

autophagic activity (Figure 5J). Levels of 4E-BP1 and p-4E-BP1 (Thr37/46), markers of

mTOR signaling, were improved in SapC-DOPS-fGCase-treated MLOs, indicating a reduction

in mTOR hyperactivation (Figure 5J).

In conclusion, the effective restoration of GCase activity and correction of GD-related disease

phenotypes highlight both the utility of the MLO platform for studying nGD and the therapeutic

potential of this nanoparticle-based approach for treating nGD.

AAV-mediated gene transfer in GD MLOs enhances GCase activity and mitigates GD

phenotypes

AAV-based gene therapies have shown clinical safety and efficacy in neurodegenerative diseases

[51,52]. Trials for GD Type 1 (NCT05487599) and Type 2 (NCT04411654) using AAV9 have

initiated. While animal models support AAV's effectiveness in GD, its cellular impact on human

or patient derived models remains unknown. To investigate the therapeutic effect of AAV gene

therapy in human GD organoids, AAV9-GBA1 was injected at dose of 1.8×10^{10} vg per organoid

at week 13 and evaluated 3 weeks after the injection (Figure 6A). In WT-75.1 MLOs, GCase

activity was significantly increased in AAV9-GBA1-treated MLOs compared to untreated

controls at week 15 (Figure 6B, left panel), suggesting effective transgene expression in normal

organoids. AAV9-GBA1 restored GCase activity in GD2-1260 and GD2-10-257 MLOs to 47.8%

and 37.7% of WT-75.1 levels, respectively (p < 0.001), compared to 6.0% and 8.8% in untreated

controls (Figure 6B, middle and right panel). Immunoblot results further confirmed the

restoration of GCase protein in AAV9-GBA1 treated GD MLOs (Figure 6B, bottom panel). Mass

spectrometry analysis showed that AAV9-GBA1 normalized GluSph levels in GD2-1260 and

GD2-10-257 MLOs, contrasting with elevated levels in untreated MLOs versus WT-75.1

controls (Figure 6C), indicating effective clearance of toxic lipid substrates in MLO by AAV

gene therapy. Of note, LAMP1 levels were elevated after AAV9-GBA1 treatment, reflecting

correction of lysosomal dysfunction (Figure 6D). However, the protein level of DA neuron

marker (TH) was not significantly increased after AAV gene therapy compared to untreated

group, suggesting longer treatment or earlier gene therapy intervention might be needed to

recover dopaminergic neurons. Confocal imaging corroborated that AAV9-GBA1-treated

GD2-1260 MLOs exhibited GFP signals and restored GCase expression in pan-neurons, DA

neurons and astrocytes (Figure 6E).

These results collectively demonstrate that AAV9-GBA1 gene therapy effectively corrected

GCase deficiency, reduced GluSph accumulation, and ameliorates lysosomal pathology in GD

MLOs, highlighting its therapeutic potential for nGD.

Substrate Reduction Therapy reduces lipid accumulation and improves lysosomal function

in GD MLOs

Current Substrate Reduction Therapy (SRT) in the standard clinical treatments for GD [53,54] has

limited efficacy in addressing the neurological manifestations of nGD due to their inability to

cross the blood-brain barrier efficiently. GZ-682452 (termed herein as GZ452) is an analogue of

venglustat that is presently under clinical evaluation for treating nGD [55]. To evaluate the human

MLO as a preclinical model for SRT drug development, we assessed GZ452's effects on MLO

growth, lipid accumulation, midbrain markers, and lysosomal function.

GZ452 toxicity was first evaluated in WT-75.1 MLOs. Treatment at 0.3, 1, 2, 3 µM over 6 weeks

modestly reduced MLO size at 1 µM, with the 2 µM and 3 µM doses significantly decreasing

size versus untreated controls (Figure 7A), indicating high dose of GZ452 notably suppressed

MLO growth. We next tested GZ452 at doses (0.01, 0.05, 0.3 µM) over 6 weeks in WT-75.1

MLOs. GZ452 dose-dependently reduced total GluCer levels (Figure 7B) and GluCer species,

with the 0.3 µM dose showing the most pronounced effect (Figure 7C).

Expression of midbrain markers *ASCL1*, *TH*, and *PLZF* in WT-75.1 MLOs was largely

unaffected by GZ452 (0.3 µM) treatment started at week 6, with no significant changes in

relative mRNA levels compared to untreated controls (Supplementary Figure 5). We therefore

applied this optimal tolerated GZ452 dose of 0.3 μM (300nM) in short-term (2 weeks) and

long-term (up to 28 weeks) treatment to test its therapeutic effects in GD MLOs. After two weeks

of treatment starting in week 13, GZ452 significantly ameliorated GluCer and GluSph

accumulation in GD2-1260 MLOs (Figure 7E, 7F). Long-term GZ452 treatment from day 2 to

week 28 robustly cleared GluSph storage and normalized it to the WT level (Figure 7G, 7H).

Immunoblot analysis showed that long-term GZ452 treatment improved lysosomal function,

partially restoring LAMP1 levels (0.8 ± 0.1; $p < 0.05$) and reducing LC3-II by 21.4% ($p < 0.01$)

(Figure 7I, 7J). These findings demonstrate that GZ452 effectively reduced lipid accumulation

and improves lysosomal function in GD MLOs, supporting its therapeutic potential for

ameliorating key disease phenotypes.

Discussion

We developed MLOs from GD Type 2 patient-derived hiPSCs, offering deepened understanding to disease mechanisms and an advanced model system for drug discovery. To our knowledge, this is the first GD brain organoid model derived from patient cells. These GD MLOs harbor prevalent $GBA1^{\text{L444P/P415R}}$ and $GBA1^{\text{L444P/RecNcil}}$ mutations, replicated key nGD phenotypes, including reduced GCase activity, glycosphingolipids accumulation, lysosomal abnormalities, and impaired dopaminergic neuron differentiation. Emerging interventions, CRISPR/Cas9-mediated gene correction, SapC-DOPS-fGCase nanoparticle, and AAV9-GBA1 gene therapy, significantly restored GCase function and mitigate disease hallmarks in GD MLOs, supporting their use in preclinical studies. Additionally, dysregulated Wnt signaling and impaired dopaminergic neuron differentiation in GD MLOs provides novel insights into nGD pathogenesis and potential therapeutic targets. This research underscores the utility of patient-derived MLOs as a human-relevant platform for modeling disease complexity and testing drug responses to bridg preclinical and clinical research.

Our findings revealed nGD-specific phenotypes in MLOs, including reduced GCase activity, GluCer and GluSph accumulation, and altered mTOR and autophagosome-lysosomal pathways, closely reflecting patient neuropathology [4,47]. Unlike *Gba1* knockout animal models, which lack patient-specific genetic and epigenetic diversity, patient-derived MLOs retain heterozygous or compound heterozygous background, enabling study of disease mechanisms and modifier genes. Partial rescue in isogenic iso-GD2-1260 MLOs, where mutation correction alleviated but did not fully resolve abnormalities, highlights the model's value in capturing nGD pathology and guiding personalized precision therapeutic strategies. This model offers a more advanced and

physiologically relevant platform than traditional gene knockout or transgenic models, better

capturing the complexity of nGD pathology.

Transcriptomic analysis of GD MLOs uncovered dysregulated Wnt signaling, a critical pathway

in early brain development, highlighting a mechanistic insight into nGD pathogenesis. Aligning

with Awad et al., who reported that downregulated Wnt/β-catenin signaling impairs DA neuron

differentiation in nGD hiPSC-derived neuronal progenitors [25], our MLOs exhibited reduced

FOXA2+ DA progenitors and TH+ mature DA neurons, along with decreased dopamine release

in culture medium. Transcriptome analysis of the GD MLOs identified 1,429 differentially

expressed genes enriched in Wnt signaling, axon guidance and neuron differentiation pathways,

indicating that GCase deficiency disrupts midbrain patterning and neuronal specification. This

finding is further supported by reduced expression of midbrain markers (e.g., FOXA2, TH),

reduced dopamine release from DA neurons, and skewed forebrain/midbrain patterning genes

(e.g., FOXP1, FOXG1) in GD MLOs. Compared to *GBA1* KO iPSC-derived midbrain-like

organoids, which showed only a marginal reduction in dopaminergic neuron differentiation and

no significant dysregulation of FOXG1 expression [23], our MLO model exhibits a distinct and

more disease-relevant phenotype. These results suggest that aberrant Wnt signaling contributes

to impaired dopaminergic neuron development and may drive progressive neurodegeneration in

nGD. Thus, targeting Wnt-related pathways could offer a therapeutic strategy to address early

developmental defects in nGD. Midbrain organoids at 35 days of development closely

correspond to the human embryonic midbrain at approximately 9 weeks, while 70-day-old

organoids align more closely with the 10-week stage [56]. These correlations highlight the

importance of initiating treatment during the fetal stage, which may be critical for preventing the

disease.

Isogenic MLOs generated by correcting the *GBA1* L444P mutation in GD2-1260 hiPSCs using

CRISPR/Cas9 revealed the key genetic mechanisms in nGD. The resulting WT/P415R genotype

in iso-GD2-1260 partially rescued nGD phenotypes. GCase activity restored to nearly 50% of

WT-75.1 levels, with normalized GluSph levels and improved TH expression. However, partial

recovery of FOXA2 (53% of WT-75.1), along with persistent LC3 and LAMP1 abnormalities,

may contribute to remaining pathology. These findings imply that the retained P415R mutation

in this isogenic line may contribute to residual lysosomal dysfunction, potentially exerting a

negative effect, as it has been reported to be associated with altered GCase stability and activity

in previous studies [57,58]. The partial rescue in iso-GD2-1260 MLOs suggests that the P415R

mutation may affect disease severity and therapeutic response, deserving further investigation

into the specific role of P415R mutation in GD and assessing whether dual mutation correction is

required for full phenotypic normalization.

Evaluation of GZ452, a CNS-accessible SRT drug [59], in GD2-1260 MLOs showed significant

reduction in GluCer and GluSph levels at a well-tolerated dose, alongside partial restoration of

lysosomal and autophagy markers (LAMP1, LC3-II). Unlike higher doses that affected organoid

growth, 0.3 μM preserved MLO size and expression of midbrain markers (ASCL1, TH, PLZF)

involved in organoid differentiation and growth. These findings underscore the value of the

MLOs for assessing drug safety (e.g., organoid size as toxicity indicator) and efficacy in a

human-specific, midbrain-relevant model before clinical studies. Our study demonstrated

positive   therapeutic   outcomes   using   AAV9-GBA1   gene   therapy   and   SapC-DOPS

nanoparticle-mediated GCase delivery in GD2-1260 and GD2-10-257 MLOs.AAV9-GBA1

restored GCase activity, while SapC-DOPS-fGCase elevated GCase activity to WT levels, both

approaches effectively reduced GluSph and improved lysosomal LAMP1 levels. Based on prior

studies showing no region-specific effects in GD patients and mice, these therapies may also be

effective in other brain organoid types, such as COs [4,33]. The novelty of employing MLOs lies in

their ability to concurrently test these diverse therapeutic modalities: SRT, gene therapy, and

nanoparticle-based enzyme delivery within a patient-derived system that replicates nGD

pathology, offering the unique patient-MLO system to assess complex cellular responses that

directly reflect patient specific conditions.

Unlike animal models, MLOs provide a controlled environment to evaluate dose-dependent

effects and cellular responses, enabling the identification of optimal therapeutic windows while

minimizing the risk of adverse outcomes in patients with nGD. Moreover, FDA has increasingly

encouraged the use of organoid systems in drug development, recognizing their potential to

improve preclinical testing and reduce reliance on animal models. Therefore, the adaptability of

MLOs to test advanced interventions, coupled with their alignment with regulatory trends

favoring human-relevant models, positions them as a great tool for accelerating the translation of

innovative treatments from preclinical research to clinical application.

There are limitations in our current MLOs, such as lacking vascular system and microglia.

Without vascularization, cells in the core may experience hypoxic and nutrient deprivation,

leading cell death or necrosis in deeper layers in long-term culture. This limitation may partially

explain the incomplete restoration of dopaminergic markers (TH, FOXA2) and partial recovery

of LAMP1 following SapC-DOPS-fGCase and AAV9-GBA1 treatments. Additionally, the lack

of microglia limits molding of neuroinflammatory, which contributes to the clearance of toxic

lipid substrates and mediating neuronal damage in GD. Elevated GluSph and impaired

dopaminergic differentiation observed in GD MLOs may not fully reflect inflammatory process.

Dysregulated Wnt signaling and lysosomal an autophagic markers (LAMP1, LC3-II/I) could be

amplified by microglial activation *in vivo*. To improve model fidelity, future studies would

incorporate vascularization (e.g., endothelial co-culture or microfluidics) to enhance oxygen and

nutrient distribution and integrate microglia via directed differentiation [60] or co-culture with

hiPSC-derived microglia [61] to enhance the relevance of MLOs for long-term nGD research and

drug testing.

In conclusion, patient-derived MLOs offer an advanced, human-relevant platform for elucidating

nGD pathogenesis and evaluating therapies, overcoming limitations of traditional models.

Dysregulated Wnt signaling and impaired dopaminergic neuron differentiation, along with partial

rescue of phenotypes through gene correction, highlight the complex genetic and developmental

underpinnings of nGD. The efficacy assessment of SapC-DOPS-fGCase, AAV9-GBA1 and

SRT-GZ452 therapies underscores the targeted interventions and safety evaluation of these

approaches in patient-derived brain organoid models. While limitations such as lack of

vascularization and microglia constrain modeling of neuroinflammation, these challenges can be

addressed using advanced culturing techniques. Collectively, our findings lay the groundwork for

developing patient-specific preclinical models to support personalized therapeutic strategies,

while also advocating for the continued refinement of MLO systems to accelerate the discovery

of effective treatments for nGD.

Acknowledgments

This work was supported by the Cincinnati Children's Pluripotent Stem Cell Facility (RRID: SCR_022634) and Transgenic Animal and Genome Editing Facility (RRID: SCR_022642). Authors acknowledge Freeline Therapeutics (now Spur Therapeutics) for providing the fGCase under a Material Transfer Agreement.

Additional information

Funding

This work is supported by awards from the National Institute of Health (grant number R21HD1027881, R21OD033660 and R01NS138309 to Y.S.; R01NS103931 partly to Y.S.; R21HD1027881 partly to C.N.M.; R01NS138309 partly to X.Q.); the National Center for Advancing Translational Sciences of the National Institutes of Health (Award Number 2UL1TR001425-05A1, CHMC-CTSA 00003827 to Y.S.); Cincinnati Children's Hospital Medical Center (CCHMC) Center of Pediatric Genomics Award to Y.S.; CCHMC Research Innovation Pilot Funding Program Award to Y.S., Y.L. and J.E.H. The funding source did not have any role in study design, data collection, data analyses, interpretation, or writing of report.

Author Contributions

Y.L. designed experiments, cultured hiPSCs and MLOs, performed experiments, analyzed data, and wrote the manuscript. B.L. measured GCase enzyme activity in hiPSCs and MLOs. V.F. carried out lipid extract from hiPSCs and MLOs, W.Z., X.Z., R.L.B. and K.DR.S performed quantification of GluCer/GluSph by LC-MS/MS and data analysis. S.A. assisted with immunofluorescence staining and immunoblot assays. C.N.M and YC.H made isogenic hiPSC

cells and conceived experimental designs and edited the manuscript. J.T. conceived experimental

designs and edited the manuscript. J.E.H. generated cerebral organoid and edited manuscript.

A.K. and X.Q. provided SapC-DOPS nanoparticles and guidance for the enzyme uptake assay.

R.A.F provided an nGD hiPSC cell line and edited the manuscript. Y.S. conceived experimental

designs, supervised the project, wrote the manuscript, and provided funding to the study.

Ethics

This work has been reviewed and approved by the Cincinnati Children's Hospital Medical

Center (CCHMC) and University of Cincinnati (UC) Embryonic Stem Cell Research Oversight

(ESCRO) committee under ESCRO protocol # EIPHG3114_R3. Primary fibroblast (GM01260)

for generating GD2-1260 iPSC was obtained from the Coriell Institute for Medical Research

under Material Transfer Agreement. WT-75.1 iPSC was generated using fibroblasts cultured

from discarded circumcision tissue obtained from a healthy donor (provided by Dr. Susanne

Wells, Division of Hematology/Oncology, CCHMC; institutional review board no. 02-9-29X) at

CCHMC. The procedures used to generate the GD2-1260 iPSC and WT-75.1 iPSC lines at

CCHMC have been reviewed and approved by CCHMC/UC ESCRO committee. Skin biopsies

to generate GD2-10-257 iPSCs were obtained from NIH with informed consent and were

approved by the Institutional Review Board (IRB). The procedures used to generate the

GD-2-10-257 nGD iPSC line at the University of Maryland Baltimore (UMB) have been

reviewed and approved by the UMB IRB and ESCRO committees.

Conflict of Interests

K.D.R.S. discloses outside this work equity in Asklepion Pharmaceuticals, Baltimore, and

Aliveris s.r.l. Italy and is a consultant to Mirum Pharmaceuticals. Other authors declare no

competing interests.

Fig. 1

Figures

Figure 1. Generation and characterization of midbrain-like organoids (MLOs) from

healthy human (h) iPSCs (WT-75.1 hiPSCs).

(A) A schematic overview of the procedures for generating MLOs from healthy WT-75.1

hiPSCs.

(B) Representative confocal images showing architectural structure of week (Wk) 8 MLOs

containing new-born neurons (Tuj1/NeuN), astrocytes (GFAP), dopaminergic neurons

(FOXA2/TH) and neural progenitor cells (SOX2/Ki67). Merge images show the distribution of

those cell markers.

(C) FOXG1 expression in MLO and cerebral organoid (CO). Transcription factor FOXG1

(forebrain marker) was enriched in CO at Wk8 of differentiation but absent in MLO.

Pan-neurons (NeuN) were both present in MLO and CO, as shown by NeuN immunostaining

(neuronal marker).

(D) Quantitative analysis of cell type specific genes expression for midbrain/dopaminergic

neuron (*FOXA2*/*ASCL1*/*LXM1A*/*PLZF*/*TH*), glial cells (*GLAST*/*S100B*) and multipotent stem

cells (*SOX2*/*NANOG*/*OCT4*) in Wk3 and Wk8 MLOs by qRT-PCR.

(E) Immunoblot of Sox2, Tuj1, MAP2, TH, GFAP and S100B in WT-75.1 hiPSCs and its derived

MLO (Wk8) lysate. β-Actin was used as a loading control.

Figure 2. GCase deficiency drives glycosphingolipids accumulation and transcriptomic alteration in GD MLOs.

The GD MLOs were generated as in Figure 1A.

(A) Reduced GCase protein in GD MLO (GD2-1260). Wk 8 MLOs (n=3) were pooled as a

biological sample. β-Actin was used as loading control.

(B) GCase activity in hiPSCs and MLOs. Data was normalized to WT-75.1 control.

(C) Representative images of WT-75.1 and GD2-1260 MLOs at Wks 4, 8 and 15 of

differentiation.

(D) MLO size was measured based on area of MLO spheres and normalized to WT-75.1 control

at each indicated time points. N ≥ 10 MLOs were quantified per group.

(E, F) Measurement of total glucosylceramide (GluCer) and GluCer species in Wk 15 MLO.

(G) Glucosylsphingosine (GluSph) levels in Wk15 and Wk28 MLOs. GluCer and GluSph levels

in the organoids were measured by LC-MS/MS and normalized by corresponding total protein of

MLO tissue lysate.

(H) 3D Principal Component Analysis (PCA) of bulk RNA sequencing (RNA-seq) data. The

Euclidean distance of the normalized gene expression among healthy control (WT-75.1) and GD

(GD2-1260) MLOs was used for sample clustering. Ellipsoids around each group indicate the

distribution and spread of the samples within the sample group. Wk 8 MLOs (n=3) were pooled

as one biological sample, and three samples were profiled in each group.

(I) MA plot showing the distinct genes differentially expressed in GD MLOs. Statistically

significant differentially expressed genes (DEGs; |fold change|≥ 1, p-adj ≤ 0.05 and base mean ≥

50) were highlighted in red. Number of DEGs downregulated and upregulated in GD2-1260

MLO against WT-75.1 MLO were shown. FC, fold-change.

(J, K) Dysregulated pathway in GD MLOs analyzed by GO (gene ontology) (J) and Kyoto

Encyclopedia of Genes and Genomes (KEGG) (K) enrichment of DEGs. Both gene counts and

level of significance (-log10 of p value) were shown as stacked columns for each category.

(L-O) Heat maps of dysregulated pathways or biological functions in GD MLO. Specifically,

aberrant expressions of genes involved in WNT signaling (L), anterior-posterior brain

specification (M), neuronal function (N) and lysosome-phagosome (O) were shown.

Figure 3. Skewed specification of midbrain patterning and dopaminergic neuron differentiation in GD MLOs.

(A) Gene expression of *FOXP1*, *FOXG1*, and *PAX6* in week 8 WT-75.1 and GD2-1260 MLOs.

Data were plotted using RNA sequencing counts. ***p < 0.001, unpaired Student's t test.

(B, C) Aberrant expression of FOXP1/FOXG1 transcription machinery for forebrain/midbrain

patterning in GD MLOs. Representative confocal images (B) and quantification of Wk8 WT-75.1

and GD2-1260 MLOs, immunostained for FOXP1 (red) and FOXG1 (green), with DAPI (blue)

labeling nuclei. Yellow arrows indicate FOXP1+FOXG1+ cells. Scale bar, 50 μm. *p < 0.05, **p

< 0.01, unpaired Student's t test.

(D) Confocal images of Wk6 MLOs, immunostained for midbrain patterning markers FOXA2

(green) or TH (red), with DAPI (blue) labeling nuclei. Scale bar, 100 μm.

(E) Representative images of differentiating DA neurons in MLOs derived from WT-75.1 and

GD2-1260 hiPSCs. TH (red), FOXA2 (green) were co-stained, with DAPI (blue) labeling nuclei.

Yellow arrows indicate TH+FOXA2+ cells. Scale bar, 50 μm.

(F) Quantification of midbrain progenitor markers *ASCL1*, *TH*, *LMX1A*, and *PLZF* expression in

WT-75.1 and GD2-1260 MLOs at Wk3 and Wk8, measured by qRT-PCR and normalized to

WT-75.1 hiPSC cells. Data are presented as mean ± SEM (n = 3-4 per group). *p < 0.05, **p <

0.01.

(G) Immunoblot analysis of midbrain/dopaminergic neuron markers TH, FOXA2, and MAP2 in

Wk16 MLOs. β-Actin was used as a loading control.

(H) Relative protein levels of TH, FOXA2, and MAP2 in Wk8 GD2-1260 MLOs compared to

WT-75.1. *p < 0.05, **p < 0.01, unpaired Student's t test.

(I) Dopamine levels in MLO culture medium assay by ELISA. Culture medium from 4

GD2-1260 MLOs or WT-75.1 MLOs at Wk12 cultured in 3 mL BGM medium for 72 hours was

Fig. 4

Figure 4. Mutation correction significantly rescued disease phenotypes in GD MLOs.

(A) Schematic overview of CRISPR/Cas9-mediated mutation correction of the *GBA1* L444P

mutation in GD2-1260 hiPSCs, converting the L444P (L444P/P415R) mutation (Proline, P to

Leucine, L) to the wild-type sequence (WT-P415R), generating isogenic iso-GD2-1260 hiPSCs.

The mutated base C in amino acid code 'CCG' for proline (P), was corrected to T to decode

leucine (L, CTG), which was confirmed by genome sequencing of *GBA1* locus.

(B, C) Immunoblot analysis of GCase protein and quantification in week 8 MLOs derived from

WT-75.1, GD2-1260, and iso-GD2-1260 hiPSCs. β-Actin was used as a loading control. Data are

presented as mean ± SEM (n = 3 biological replicates per group). ***$p < 0.001$.

(D) Relative GCase activity in GD2-1260 and iso-GD2-1260 hiPSCs and Wk8 MLOs,

normalized to WT-75.1 controls. Data are presented as mean ± SEM (2 MLOs pooled, n = 3 per

group). ***$p < 0.001$.

(E) Measurement of GluSph levels in WT-75.1, GD2-1260, and iso-GD2-1260 MLOs at Wk15

and Wk28 and their culture medium at Wk 15, quantified by LC-MS/MS and normalized to total

protein of tissue lysate. Data are presented as mean ± SEM (3 MLOs pooled, n = 3 per group for

MLO and 4 MLOs pooled, n = 3 per group for medium). **$p < 0.01$; ns, not significant.

(F) Representative bright-field images of GD2-1260 and iso-GD2-1260 MLOs at Wk15 of

differentiation. Scale bar, 1 mm.

(G) MLO size quantification for GD2-1260 and iso-GD2-1260 MLOs at Wks 4, 8, and 15,

measured as relative area of MLO spheres and normalized to WT-75.1 controls at indicated time

point. N ≥ 10 MLOs were quantified per group. Data are presented as mean ± SEM. *$p < 0.05$;

ns, not significant.

(H, I) Immunoblot analysis of midbrain/dopaminergic neuron markers TH and FOXA2 (H) and

their relative quantification (I) in Wk8 MLOs. GAPDH was used as a loading control.

(J) Dopamine levels in the culture medium of Wk12 MLOs derived from GD2-1260 and

iso-GD2-1260 hiPSCs, measured after 72 hours in BGM medium (n = 4 MLOs). Data are

presented as mean ± SEM (n = 5~7 per group). **$p < 0.01$.

(K, L) Immunoblot analysis of autophagy-lysosomal pathway markers LAMP1 and Cathepsin D

(K) and quantification (L) in Wk16 MLOs. GAPDH was used as a loading control. Data are

presented as mean ± SEM.

(M, N) Immunoblot analysis of LC3-I and LC3-II (M) and quantification (N) in Wk16 MLOs.

GAPDH was used as a loading control.

(O) Immunoblot analysis of mTOR signaling pathway components [4E-BP1,

P-4E-BP1(THR37/46), S6, and P-S6 (Ser235/236)] in Wk16 MLOs. β-Actin was used as a

loading control.

(P) Quantification of protein levels of mTOR signaling pathway components. Data are

normalized to WT-75.1 and presented as mean ± SEM.

Immunoblot analysis for panel H and I and K-P was performed using the lysate from 3 MLOs

pooled per group, 3 repeated experiments. ***$p < 0.001$; ns, not significant. One-way ANOVA

test with Tukey's test.

Fig. 5

Figure 5. Delivery of GCase to MLOs via SapC-DOPS nanoparticles corrects GD phenotypes.

(A) Schematic illustration of SapC-DOPS nanoparticle-mediated delivery of recombinant GCase

(fGCase) to MLOs. SapC-DOPS nanoparticles carrying fGCase or fluorescent label CVM were

cocultured with MLOs, followed by short-term (48 hours) or 2-week treatment period before

analysis.

(B) Confocal images of untreated and SapC-DOPS-CVM-treated MLOs, showing uptake of

CVM (magenta) with DAPI (blue) labeling nuclei. Scale bars: 200 μm (left panel), 50 μm (right

panels, magnified regions a and b).

(C) GCase activity in WT-75.1 and GD2-1260 MLOs following a 48-hour treatment with

SapC-DOPS-fGCase. Data are presented as mean ± SEM (3 MLOs pooled, n = 3 per group).

***$p < 0.001$.

(D) Confocal images of WT-75.1, GD2-1260, and GD2-10-257 MLOs treated with SapC-DOPS

or SapC-DOPS-fGCase for 2 weeks, immunostained for GCase (green) with DAPI (blue)

labeling nuclei. Scale bar, 200 μm.

(E-G) GCase activity and protein in WT-75.1 and GD (GD2-1260, GD2-10-257) MLOs treated

with SapC-DOPS or SapC-DOPS-fGCase for 2 weeks, measured by enzymatic assay and

immunoblot. Data are presented as mean ± SEM (3 MLOs pooled, n = 3~4 per group). ***$p <$

0.001; ns, not significant.

(H, I) GluSph levels in WT-75.1 and GD (GD2-1260, GD2-10-257) MLOs treated with

SapC-DOPS or SapC-DOPS-fGCase for 2 weeks, quantified by LC-MS/MS and normalized to

total protein. Data are presented as mean ± SEM (3 MLOs pooled, n = 3-4 per group). ***$p <$

0.001; **$p < 0.01$; ns, not significant.

(J) Immunoblot analysis of autophagy-lysosomal and mTOR pathway proteins in SapC-DOPS or

SapC-DOPS-fGCase treated GD2-1260 MLOs. GAPDH was used as a loading control.

Fig. 6

Figure 6. AAV9-GBA1 gene therapy mitigates disease phenotypes in GD MLOs.

(A) Schematic illustration of AAV9-*GBA1* gene therapy delivery to MLOs using a nanoliter

injector. AAV9 vectors carrying the GBA1 gene (AAV9-GBA1) are administered to Wk13

MLOs. The samples were analyzed after 3 weeks of treatment.

(B) GCase activity in WT-75.1, GD2-1260, and GD2-10-257 MLOs and AAV9-GBA1 treated

MLOs were measured by enzymatic assay. Data are presented as mean ± SEM (3 MLOs pooled,

n = 3 to 6 per group). ***p < 0.001.

(C) GluSph levels in AAV9-GBA1 treated GD and control MLOs were quantified by

LC-MS/MS and normalized to total protein. Data are presented as mean ± SEM (3 MLOs pooled,

n ≥ 3 per group). ***p < 0.001; ns, not significant.

(D) Immunoblot analysis of LAMP1 and TH in WT-75.1 and in GD2-1260 MLOs untreated or

treated with AAV9-GBA1. Protein levels are normalized to WT-75.1 untreated controls (set to

1.0).

(E) Transgene expressions (yellow arrows and enlarged insert) in neurons (NeuN), DA neurons

(TH) and astrocytes (GFAP) of AAV9-GBA1 treated GD2-1260 MLOs. Scale bar = 50 μm.

Fig. 7

Figure 7. Substrate Reduction Therapy with GZ452 reduces lipid accumulation and improves autophagic and lysosomal abnormalities in GD MLOs.

(A) Assessment of GZ452 tolerated dose in healthy MLO and the effect of GZ452 on organoid growth in WT-75.1 MLOs over 6 weeks.

(B, C) Total GluCer levels (B) and distribution of GluCer species (C) in WT-75.1 MLOs with various doses of GZ452 at Wk6.

(D) Schematic of the experimental timeline for short-term (2 weeks) GZ452 treatment of GD MLOs.

(E, F) GluCer (E) and GluSph (F) levels in WT-75.1 and GD2-1260 MLOs at Wk15 under

short-term GZ452 treatment. Data were normalized to protein mass.

(G) Schematic of the experimental timeline for long-term (28 weeks) GZ452 treatment in GD

MLOs.

(H) GluSph levels in MLOs at Wk15 under long-term GZ452 treatment. Data were normalized to

protein mass.

(I) Immunoblot analysis of LAMP1 and LC3-I/II in SRT treated GD2-1260 MLOs for 28 weeks,

with β-actin as loading control.

(J) Quantification of LAMP1 and LC3-II/I in MLOs. Protein levels are normalized to WT-75.1

untreated controls (set to 1.0).

All data presented as mean ± SEM (n = 3~4). $*p < 0.05$; $**p < 0.01$; ns, not significant. One-way

ANOVA with Tukey's test.

Additional files

Supplementary files

Supplementary Fig. 1. Correct L444P mutation in nGD iPSC by CRISPR-Cas9. (A) Single

stranded oligonucleotide design. Silent mutations in upper case. Phosphorothioate modified

bases (*). (B) Genomic sequence of mutant, wild type and corrected GBA1. Mutated codon

highlighted in red. T insertion for point mutation correction highlighted in yellow. PAM

highlighted in grey. sgRNA target sequence underlined. (C) DNA electrophoresis gels showing

genome editing and clone

screening for iso-GD2-1260 (clone #9). (D) Karyotyping of GD2-1260 and CRISPR/Cas9

corrected iso-GD2-1260 hiPSCs. Normal karyotype was observed in both hiPSC lines. (E)

Relative mRNA expression of genes (OCT4, NANOG and SOX2) required for generating and

maintaining hiPSCs pluripotency by quantitative RT-PCR. ns, not significant by One-way

ANOVA

analysis.

Supplementary Fig. 2. Neural rosette formation during MLO maturation were not affected by

GBA1 mutation. (A) Cryosections of Wk 6 MLOs derived from WT-75.1, GD2-1260 and

isogenic control iso-GD2-1260 hiPSC cells. Sections were stained with antibodies against Ki67

and Sox2 and nuclei were costained with DAPI. Scale bar, 100 μm. (B) Quantification of the

Sox2+, Ki67+ and Sox2+/Ki67+ cells (mean ± SEM; n = 3). ns, not significant by One-way

ANOVA analysis.

Supplementary Fig. 3. Restoration of GCase Expression in dopaminergic neuron and astrocytes

in SapC-DOPS-fGCase treated nGD MLOs. (A, B) Representative confocal images of

GD2-1260 (A) and GD2-10-257 MLOs (B) treated with SapC-DOPS-fGCase for 2 weeks,

immunostained for GCase (green) and dopaminergic neuron (TH, red) with DAPI (blue) labeling

nuclei. (C, D) Representative immunostaining images for GCase (green) and astrocytes (GFAP,

red) in GD2-1260 (C) and GD2-10-257 MLOs (D) treated with SapC-DOPS-fGCase. Scale bar,

50 µm. Yellow arrows indicate colocalized GCase in TH+ cells.

Supplementary Fig. 4. Restoration of GCase expression in lysosomal and autophagosomal

compartments in SapC-DOPS-fGCase treated nGD MLOs. (A, B) Representative confocal

images of GD2-1260 (A) and GD2-10-257 MLOs (B) treated with SapC-DOPS-fGCase for 2

weeks, immunostained for GCase (green) and lysosomal marker LAMP1 (red) with DAPI (blue)

labeling nuclei. Yellow arrows indicate colocalized GCase in LAMP1+ compartments. Scale bar,

50 µm. (C, D) Representative confocal images of GD2-1260 (C) and GD2-10-257 MLOs (D)

treated with SapC-DOPS-fGCase for 2 weeks, immunostained for GCase (green) and

autophagosomal marker LC3B (red) with DAPI (blue) labeling nuclei. Yellow arrows indicate

colocalized GCase in LC3B+ compartments. Scale bar, 50 µm.

Supplementary Fig. 5. Influence of SRT drug GZ452 on DA neuron differentiation in WT-75.1

MLOs. Relative mRNA expression of midbrain markers ASCL1, TH, and PLZF in WT-75.1 at

Wk6 in untreated or treated MLOs with indicated concentrations of GZ452. Relative gene

expression is normalized to untreated control WT-75.1 MLOs (set to 1.0).

Supplementary Table 1. Key Resources

Data availability

RNA-seq data generated in this study were deposited in Gene Expression Omnibus (GEO) at

National Institutes of Health (NIH) with accession number GSE303993.

References

1. Grabowski, G.A., Petsko, G.A., and Kolodny, E.H. (2010). Gaucher disease. In The
Online Metabolic and Molecular Bases of Inherited Diseases, D. Valle, A. Beaudet, B.
Vogelstein, K.W. Kinzler, S.E. Antonarakis, A. Ballabio, C.R. Scriver, W.S. Sly, B. Childs,
F. Bunz, et al., eds. (New York: The McGraw-Hill Companies, Inc.).
2. Mignot, C., Gelot, A., Bessieres, B., Daffos, F., Voyer, M., Menez, F., Fallet Bianco, C.,
Odent, S., Le Duff, D., Loget, P., Fargier, P., et al. (2003). Perinatal-lethal Gaucher
disease. American journal of medical genetics. Part A *120A*, 338-344.
10.1002/ajmg.a.20117.
3. Reissner, K., Tayebi, N., Stubblefield, B.K., Koprivica, V., Blitzer, M., Holleran, W.,
Cowan, T., Almashanu, S., Maddalena, A., Karson, E.M., and Sidransky, E. (1998). Type
2 Gaucher disease with hydrops fetalis in an Ashkenazi Jewish family resulting from a
novel recombinant allele and a rare splice junction mutation in the glucocerebrosidase
locus. Molecular genetics and metabolism *63*, 281-288. 10.1006/mgme.1998.2675.
4. Burrow, T.A., Sun, Y., Prada, C.E., Bailey, L., Zhang, W., Brewer, A., Wu, S.W., Setchell,
K.D.R., Witte, D., Cohen, M.B., and Grabowski, G.A. (2015). CNS, lung, and lymph
node involvement in Gaucher disease type 3 after 11 years of therapy: clinical,
histopathologic, and biochemical findings. Molecular genetics and metabolism *114*,
233-241. 10.1016/j.ymgme.2014.08.011.
5. Wong, K., Sidransky, E., Verma, A., Mixon, T., Sandberg, G.D., Wakefield, L.K.,
Morrison, A., Lwin, A., Colegial, C., Allman, J.M., and Schiffmann, R. (2004).
Neuropathology provides clues to the pathophysiology of Gaucher disease. Molecular
genetics and metabolism *82*, 192-207. 10.1016/j.ymgme.2004.04.011.
6. Conradi, N.G., Sourander, P., Nilsson, O., Svennerholm, L., and Erikson, A. (1984).
Neuropathology of the Norrbottnian type of Gaucher disease. Morphological and
biochemical studies. Acta neuropathologica *65*, 99-109.
7. Enquist, I.B., Lo Bianco, C., Ooka, A., Nilsson, E., Mansson, J.E., Ehinger, M., Richter,
J., Brady, R.O., Kirik, D., and Karlsson, S. (2007). Murine models of acute neuronopathic
Gaucher disease. Proceedings of the National Academy of Sciences of the United States
of America *104*, 17483-17488. 10.1073/pnas.0708086104.
8. Vardi, A., Zigdon, H., Meshcheriakova, A., Klein, A.D., Yaacobi, C., Eilam, R., Kenwood,
B.M., Rahim, A.A., Massaro, G., Merrill, A.H., Jr., Vitner, E.B., et al. (2016). Delineating
pathological pathways in a chemically induced mouse model of Gaucher disease. J Pathol
239, 496-509. 10.1002/path.4751.
9. Xu, Y.H., Reboulet, R., Quinn, B., Huelsken, J., Witte, D., and Grabowski, G.A. (2008).
Dependence of reversibility and progression of mouse neuronopathic Gaucher disease on
acid beta-glucosidase residual activity levels. Molecular genetics and metabolism *94*,
190-203. 10.1016/j.ymgme.2008.01.013.
10. Xu, Y.H., Quinn, B., Witte, D., and Grabowski, G.A. (2003). Viable mouse models of
acid beta-glucosidase deficiency: the defect in Gaucher disease. Am J Pathol *163*,
2093-2101. 10.1016/s0002-9440(10)63566-3.
11. Sun, Y., Liou, B., Ran, H., Skelton, M.R., Williams, M.T., Vorhees, C.V., Kitatani, K.,
Hannun, Y.A., Witte, D.P., Xu, Y.H., and Grabowski, G.A. (2010). Neuronopathic
Gaucher disease in the mouse: viable combined selective saposin C deficiency and
mutant glucocerebrosidase (V394L) mice with glucosylsphingosine and

glucosylceramide accumulation and progressive neurological deficits. Human molecular
genetics *19*, 1088-1097. 10.1093/hmg/ddp580.

12. Dasgupta, N., Xu, Y.H., Li, R., Peng, Y., Pandey, M.K., Tinch, S.L., Liou, B., Inskeep, V.,
Zhang, W., Setchell, K.D., Keddache, M., et al. (2015). Neuronopathic Gaucher disease:
dysregulated mRNAs and miRNAs in brain pathogenesis and effects of pharmacologic
chaperone treatment in a mouse model. Human molecular genetics *24*, 7031-7048.
10.1093/hmg/ddv404.

13. Liu, Y., Suzuki, K., Reed, J.D., Grinberg, A., Westphal, H., Hoffmann, A., Doring, T.,
Sandhoff, K., and Proia, R.L. (1998). Mice with type 2 and 3 Gaucher disease point
mutations generated by a single insertion mutagenesis procedure. Proceedings of the
National Academy of Sciences of the United States of America *95*, 2503-2508.
10.1073/pnas.95.5.2503.

14. Pasmanik-Chor, M., Laadan, S., Elroy-Stein, O., Zimran, A., Abrahamov, A., Gatt, S.,
and Horowitz, M. (1996). The glucocerebrosidase D409H mutation in Gaucher disease.
Biochemical and molecular medicine *59*, 125-133. 10.1006/bmme.1996.0077.

15. Grabowski, G.A., Zimran, A., and Ida, H. (2015). Gaucher disease types 1 and 3:
Phenotypic characterization of large populations from the ICGG Gaucher Registry.
American journal of hematology *90 Suppl 1*, S12-18. 10.1002/ajh.24063.

16. Birtele, M., Lancaster, M., and Quadrato, G. (2025). Modelling human brain
development and disease with organoids. Nat Rev Mol Cell Biol *26*, 389-412.
10.1038/s41580-024-00804-1.

17. Di Lullo, E., and Kriegstein, A.R. (2017). The use of brain organoids to investigate neural
development and disease. Nature reviews. Neuroscience *18*, 573-584.
10.1038/nrn.2017.107.

18. Raja, W.K., Mungenast, A.E., Lin, Y.T., Ko, T., Abdurrob, F., Seo, J., and Tsai, L.H.
(2016). Self-Organizing 3D Human Neural Tissue Derived from Induced Pluripotent
Stem Cells Recapitulate Alzheimer's Disease Phenotypes. PLoS One *11*, e0161969.
10.1371/journal.pone.0161969.

19. Lancaster, M.A., Renner, M., Martin, C.A., Wenzel, D., Bicknell, L.S., Hurles, M.E.,
Homfray, T., Penninger, J.M., Jackson, A.P., and Knoblich, J.A. (2013). Cerebral
organoids model human brain development and microcephaly. Nature *501*, 373-379.
10.1038/nature12517.

20. Smits, L.M., Reinhardt, L., Reinhardt, P., Glatza, M., Monzel, A.S., Stanslowsky, N.,
Rosato-Siri, M.D., Zanon, A., Antony, P.M., Bellmann, J., Nicklas, S.M., et al. (2019).
Modeling Parkinson's disease in midbrain-like organoids. NPJ Parkinson's disease *5*, 5.
10.1038/s41531-019-0078-4.

21. Allende, M.L., Cook, E.K., Larman, B.C., Nugent, A., Brady, J.M., Golebiowski, D.,
Sena-Esteves, M., Tifft, C.J., and Proia, R.L. (2018). Cerebral organoids derived from
Sandhoff disease-induced pluripotent stem cells exhibit impaired neurodifferentiation. J
Lipid Res *59*, 550-563. 10.1194/jlr.M081323.

22. Latour, Y.L., Yoon, R., Thomas, S.E., Grant, C., Li, C., Sena-Esteves, M., Allende, M.L.,
Proia, R.L., and Tifft, C.J. (2019). Human GLB1 knockout cerebral organoids: A model
system for testing AAV9-mediated GLB1 gene therapy for reducing GM1 ganglioside
storage in GM1 gangliosidosis. Mol Genet Metab Rep *21*, 100513.
10.1016/j.ymgmr.2019.100513.

23. Jo, J., Yang, L., Tran, H.D., Yu, W., Sun, A.X., Chang, Y.Y., Jung, B.C., Lee, S.J., Saw,

T.Y., Xiao, B., Khoo, A.T.T., et al. (2021). Lewy Body-like Inclusions in Human Midbrain Organoids Carrying Glucocerebrosidase and alpha-Synuclein Mutations. Ann Neurol *90*, 490-505. 10.1002/ana.26166.

24. Sun, Y., Florer, J., Mayhew, C.N., Jia, Z., Zhao, Z., Xu, K., Ran, H., Liou, B., Zhang, W., Setchell, K.D., Gu, J., et al. (2015). Properties of neurons derived from induced pluripotent stem cells of Gaucher disease type 2 patient fibroblasts: potential role in neuropathology. PLoS One *10*, e0118771. 10.1371/journal.pone.0118771.

25. Awad, O., Panicker, L.M., Deranieh, R.M., Srikanth, M.P., Brown, R.A., Voit, A., Peesay, T., Park, T.S., Zambidis, E.T., and Feldman, R.A. (2017). Altered Differentiation Potential of Gaucher's Disease iPSC Neuronal Progenitors due to Wnt/beta-Catenin Downregulation. Stem Cell Reports *9*, 1853-1867. 10.1016/j.stemcr.2017.10.029.

26. Kwak, T.H., Kang, J.H., Hali, S., Kim, J., Kim, K.P., Park, C., Lee, J.H., Ryu, H.K., Na, J.E., Jo, J., Je, H.S., et al. (2020). Generation of homogeneous midbrain organoids with in vivo-like cellular composition facilitates neurotoxin-based Parkinson's disease modeling. Stem cells *38*, 727-740. 10.1002/stem.3163.

27. Jo, J., Xiao, Y., Sun, A.X., Cukuroglu, E., Tran, H.D., Goke, J., Tan, Z.Y., Saw, T.Y., Tan, C.P., Lokman, H., Lee, Y., et al. (2016). Midbrain-like Organoids from Human Pluripotent Stem Cells Contain Functional Dopaminergic and Neuromelanin-Producing Neurons. Cell stem cell *19*, 248-257. 10.1016/j.stem.2016.07.005.

28. Ran, F.A., Hsu, P.D., Wright, J., Agarwala, V., Scott, D.A., and Zhang, F. (2013). Genome engineering using the CRISPR-Cas9 system. Nat Protoc *8*, 2281-2308. 10.1038/nprot.2013.143.

29. Chen, B., Gilbert, L.A., Cimini, B.A., Schnitzbauer, J., Zhang, W., Li, G.W., Park, J., Blackburn, E.H., Weissman, J.S., Qi, L.S., and Huang, B. (2013). Dynamic imaging of genomic loci in living human cells by an optimized CRISPR/Cas system. Cell *155*, 1479-1491. 10.1016/j.cell.2013.12.001.

30. Sun, Y., Liou, B., Chu, Z., Fannin, V., Blackwood, R., Peng, Y., Grabowski, G.A., Davis, H.W., and Qi, X. (2020). Systemic enzyme delivery by blood-brain barrier-penetrating SapC-DOPS nanovesicles for treatment of neuronopathic Gaucher disease. EBioMedicine *55*, 102735. 10.1016/j.ebiom.2020.102735.

31. Zhao, S., Chu, Z., Blanco, V.M., Nie, Y., Hou, Y., and Qi, X. (2015). SapC-DOPS nanovesicles as targeted therapy for lung cancer. Mol Cancer Ther *14*, 491-498. 10.1158/1535-7163.MCT-14-0661.

32. Marshall, J., Sun, Y., Bangari, D.S., Budman, E., Park, H., Nietupski, J.B., Allaire, A., Cromwell, M.A., Wang, B., Grabowski, G.A., Leonard, J.P., et al. (2016). CNS-accessible Inhibitor of Glucosylceramide Synthase for Substrate Reduction Therapy of Neuronopathic Gaucher Disease. Mol Ther *24*, 1019-1029. 10.1038/mt.2016.53.

33. Sun, Y., Zhang, W., Xu, Y.H., Quinn, B., Dasgupta, N., Liou, B., Setchell, K.D., and Grabowski, G.A. (2013). Substrate compositional variation with tissue/region and Gba1 mutations in mouse models--implications for Gaucher disease. PLoS One *8*, e57560. 10.1371/journal.pone.0057560.

34. Zhang, W., Oehrle, M., Prada, C.E., Schwartz, I.V.D., Chutipongtanate, S., Wattanasirichaigoon, D., Inskeep, V., Dai, M., Pan, D., Sun, Y., and Setchell, K.D.R. (2017). A convenient approach to facilitate monitoring Gaucher disease progression and therapeutic response. The Analyst *142*, 3380-3387. 10.1039/c7an00938k.

35. Zhao, X., Lin, Y., Liou, B., Fu, W., Jian, J., Fannie, V., Zhang, W., Setchell, K.D.R.,

Grabowski, G.A., Sun, Y., and Liu, C.J. (2023). PGRN deficiency exacerbates, whereas a brain penetrant PGRN derivative protects, GBA1 mutation-associated pathologies and diseases. Proceedings of the National Academy of Sciences of the United States of America *120*, e2210442120. 10.1073/pnas.2210442120.

36. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550. 10.1186/s13059-014-0550-8.

37. Sherman, B.T., Hao, M., Qiu, J., Jiao, X., Baseler, M.W., Lane, H.C., Imamichi, T., and Chang, W. (2022). DAVID: a web server for functional enrichment analysis and functional annotation of gene lists (2021 update). Nucleic Acids Res *50*, W216-W221. 10.1093/nar/gkac194.

38. Dai, M., Lin, Y., El-Amouri, S.S., Kohls, M., and Pan, D. (2018). Comprehensive evaluation of blood-brain barrier-forming micro-vasculatures: Reference and marker genes with cellular composition. PLoS One *13*, e0197379. 10.1371/journal.pone.0197379.

39. Prakash, N., and Wurst, W. (2006). Genetic networks controlling the development of midbrain dopaminergic neurons. J Physiol *575*, 403-410. 10.1113/jphysiol.2006.113464.

40. Green, D., Whitener, A.E., Mohanty, S., and Lekven, A.C. (2015). Vertebrate nervous system posteriorization: Grading the function of Wnt signaling. Dev Dyn *244*, 507-512. 10.1002/dvdy.24230.

41. Ryckman, A.E., Brockhausen, I., and Walia, J.S. (2020). Metabolism of Glycosphingolipids and Their Role in the Pathophysiology of Lysosomal Storage Disorders. Int J Mol Sci *21*. 10.3390/ijms21186881.

42. Srikanth, M.P., Jones, J.W., Kane, M., Awad, O., Park, T.S., Zambidis, E.T., and Feldman, R.A. (2021). Elevated glucosylsphingosine in Gaucher disease induced pluripotent stem cell neurons deregulates lysosomal compartment through mammalian target of rapamycin complex 1. Stem Cells Transl Med *10*, 1081-1094. 10.1002/sctm.20-0386.

43. Mato-Blanco, X., Kim, S.K., Jourdon, A., Ma, S., Choi, S.H., Giani, A.M., Paredes, M.I., Tebbenkamp, A.T.N., Liu, F., Duque, A., Vaccarino, F.M., et al. (2025). Early developmental origins of cortical disorders modeled in human neural stem cells. Nat Commun *16*, 6347. 10.1038/s41467-025-61316-w.

44. Ba, R., Yang, L., Zhang, B., Jiang, P., Ding, Z., Zhou, X., Yang, Z., and Zhao, C. (2023). FOXG1 drives transcriptomic networks to specify principal neuron subtypes during the development of the medial pallium. Sci Adv *9*, eade2441. 10.1126/sciadv.ade2441.

45. Ortiz, A., Ayhan, F., Khandelwal, N., Outland, E., Jankovic, M., Harper, M., and Konopka, G. (2025). Cell-type-specific roles of FOXP1 in the excitatory neuronal lineage during early neocortical murine development. Cell Rep *44*, 115384. 10.1016/j.celrep.2025.115384.

46. Peng, Y., Liou, B., Lin, Y., Mayhew, C.N., Fleming, S.M., and Sun, Y. (2023). iPSC-derived neural precursor cells engineering GBA1 recovers acid beta-glucosidase deficiency and diminishes alpha-synuclein and neuropathology. Mol Ther Methods Clin Dev *29*, 185-201. 10.1016/j.omtm.2023.03.007.

47. Brown, R.A., Voit, A., Srikanth, M.P., Thayer, J.A., Kingsbury, T.J., Jacobson, M.A., Lipinski, M.M., Feldman, R.A., and Awad, O. (2019). mTOR hyperactivity mediates lysosomal dysfunction in Gaucher's disease iPSC-neuronal cells. Dis Model Mech *12*. 10.1242/dmm.038596.

1166   48.   Panicker, L.M., Miller, D., Park, T.S., Patel, B., Azevedo, J.L., Awad, O., Masood, M.A.,
1167   Veenstra, T.D., Goldin, E., Stubblefield, B.K., Tayebi, N., et al. (2012). Induced
1168   pluripotent stem cell model recapitulates pathologic hallmarks of Gaucher disease.
1169   Proceedings of the National Academy of Sciences of the United States of America *109*,
1170   18054-18059. 10.1073/pnas.1207889109.
1171   49.   Comper, F., Miranda, C.J., Liou, B., Dodev, T., Jeyakumar, J.M., Canavese, M., Cocita,
1172   C., Khoshrou, K., Tiscornia, G., Chisari, E., Stotter, E., et al. (2025). FLT201, a novel
1173   liver-directed AAV gene therapy candidate for Gaucher disease type 1. Mol Ther *33*,
1174   3789-3807. 10.1016/j.ymthe.2025.05.003.
1175   50.   Hughes, D.A., and Ferrante, F. (2023). GALILEO-1: A Phase I/II safety and efficacy
1176   study of FLT201 gene therapy for Gaucher disease type 1. Future Rare Diseases *3*,
1177   FRD35.
1178   51.   Al-Zaidy, S., Pickard, A.S., Kotha, K., Alfano, L.N., Lowes, L., Paul, G., Church, K.,
1179   Lehman, K., Sproule, D.M., Dabbous, O., Maru, B., et al. (2019). Health outcomes in
1180   spinal muscular atrophy type 1 following AVXS-101 gene replacement therapy. Pediatr
1181   Pulmonol *54*, 179-185. 10.1002/ppul.24203.
1182   52.   Drag, S., Dotiwala, F., and Upadhyay, A.K. (2023). Gene Therapy for Retinal
1183   Degenerative Diseases: Progress, Challenges, and Future Directions. Invest Ophthalmol
1184   Vis Sci *64*, 39. 10.1167/iovs.64.7.39.
1185   53.   Aerts, J.M., Hollak, C.E., Boot, R.G., Groener, J.E., and Maas, M. (2006). Substrate
1186   reduction therapy of glycosphingolipid storage disorders. J Inherit Metab Dis *29*, 449-456.
1187   10.1007/s10545-006-0272-5.
1188   54.   Scott, L.J. (2015). Eliglustat: A Review in Gaucher Disease Type 1. Drugs *75*, 1669-1678.
1189   10.1007/s40265-015-0468-9.
1190   55.   Schiffmann, R., Mengel, E., Wallace, M., Rochmann, C., Turnbull, J., Krupnick, R.,
1191   Gwaltney, C., Pulikottil-Jacob, R., Batsu, I., Zheng, R., and Hamed, A. (2024).
1192   Qualitative Study of the Patient Experience with Venglustat for Gaucher Disease Type 3
1193   in a Phase 2 Open-Label, Multicenter, Multinational Study (LEAP). Adv Ther *41*,
1194   2907-2923. 10.1007/s12325-024-02881-2.
1195   56.   Zagare, A., Barmpa, K., Smajic, S., Smits, L.M., Grzyb, K., Grunewald, A., Skupin, A.,
1196   Nickels, S.L., and Schwamborn, J.C. (2022). Midbrain organoids mimic early embryonic
1197   neurodevelopment and recapitulate LRRK2-p.Gly2019Ser-associated gene expression.
1198   Am J Hum Genet *109*, 311-327. 10.1016/j.ajhg.2021.12.009.
1199   57.   Grace, M.E., Berg, A., He, G.S., Goldberg, L., Horowitz, M., and Grabowski, G.A.
1200   (1991). Gaucher disease: heterologous expression of two alleles associated with
1201   neuronopathic phenotypes. Am J Hum Genet *49*, 646-655.
1202   58.   Ron, I., Dagan, A., Gatt, S., Pasmanik-Chor, M., and Horowitz, M. (2005). Use of
1203   fluorescent substrates for characterization of Gaucher disease mutations. Blood Cells Mol
1204   Dis *35*, 57-65. 10.1016/j.bcmd.2005.03.006.
1205   59.   Peng, Y., Liou, B., Lin, Y., Fannin, V., Zhang, W., Feldman, R.A., Setchell, K.D.R.,
1206   Grabowski, G.A., and Sun, Y. (2021). Substrate Reduction Therapy Reverses
1207   Mitochondrial, mTOR, and Autophagy Alterations in a Cell Model of Gaucher Disease.
1208   Cells *10*. 10.3390/cells10092286.
1209   60.   Cakir, B., Tanaka, Y., Kiral, F.R., Xiang, Y., Dagliyan, O., Wang, J., Lee, M., Greaney,
1210   A.M., Yang, W.S., duBoulay, C., Kural, M.H., et al. (2022). Expression of the
1211   transcription factor PU.1 induces the generation of microglia-like cells in human cortical

1212   organoids. Nat Commun *13*, 430. 10.1038/s41467-022-28043-y.

1213   61. Sabate-Soler, S., Nickels, S.L., Saraiva, C., Berger, E., Dubonyte, U., Barmpa, K., Lan,
1214   Y.J., Kouno, T., Jarazo, J., Robertson, G., Sharif, J., et al. (2022). Microglia integration
1215   into human midbrain organoids leads to increased neuronal maturation and functionality.
1216   Glia *70*, 1267-1288. 10.1002/glia.24167.

1217