Experimental evidence of male-male interaction in laboratory swarms of *Anopheles gambiae* mosquitoes

Gloria Iacomelli[1†], Max Lombardi[1,2†], Leonardo Parisi[2,3†], Matteo Fiorini[3], Francesca Grucci[3], Alessio Lavorgna[1], Melania Ligato[1], Gaetano Zarcone[1], Matthew J. Peirce[1], Stefania Melillo[2,3*], Roberta Spaccapelo[1]

[1]Department of Medicine and Surgery, University of Perugia, Perugia, Italy.
[2]Institute for Complex Systems, National Research Council, Rome, Italy.
[3]Physics Department, Sapienza University, Rome, Italy.

*Corresponding author. E-mail: stefania.melillo@cnr.it;
†These authors contributed equally to this work.

Abstract

Mosquitoes mostly mate in the context of swarms: to facilitate encounters with females, males form disordered aggregations over a visual marker, which serves as a positional reference. Due to the scarcity of high-resolution data, it remains unclear whether swarms are the result of individuals interacting independently with the marker, or whether they are the expression of a genuine collective behavior produced by spontaneous interaction between individuals. Here, with a unique dataset comprising three–dimensional trajectories of 27 laboratory swarms of different size (ranging from 80 to 400 mosquitoes), we investigate swarming behavior of *Anopheles gambiae* mosquitoes. With a statistical physics approach, we provide experimental evidence of an effective male-male interaction. We find that individual speed fluctuations are strongly correlated in space, meaning that mosquitoes in close proximity tend to display similar deviations from the group average, effectively flying at a similar speed. We also prove that this correlation is not compatible with a random arrangement of individuals, indicating that an interaction process is at play. Our findings suggest that collective behavior in lab-based mosquito swarms indeed emerges from an interaction mechanism that, unexpectedly, does not involve the direction of flight, as observed in other biological systems, but rather their speed.

Keywords: collective behavior, male-male interaction, swarming behavior, 3D data

1 Introduction

Collective behavior is a widespread phenomenon, observed across many biological systems, from cells in a colony to vertebrates in a herd. It emerges when a large number of individuals self-organize and, through mutual interaction, exchange information that can be transferred to the entire group. In its most fascinating manifestations, interaction leads to global order. This is the case, among others, with the synchronized flashing of fireflies [1, 2], or the coordinated motion of birds and fish, [3–6].

While there is general consensus [7, 8] that order arises from the interaction between individuals, the debate on the nature of the behavior in disordered systems is still active, with insect swarms

serving as primary examples [9–11]. Swarms are indeed the paradigm of disordered systems: hundreds of insects - exclusively males - congregate about a common marker [12], to facilitate encounters with females, who will join the group only briefly to mate [13]. As a whole, the swarm maintains a stable position over the marker, which is used as a positional reference [14]. But, at an individual level, insects in the swarm follow apparently erratic flight patterns.

The lack of global order, combined with the presence of a marker, supports two competing interpretations. In one, the visual attraction towards the marker dominates - if not completely determines - the insects' dynamics [9, 15]. In the other, the marker plays a more marginal role, and the interaction between individuals gives rise to collective behavior [11].

In swarms of non-biting midges (family Chironomidae), the emergence of an authentic collective behavior has been proved through field experimental data [11, 16, 17], and confirmed by theory and simulations [18]. In mosquitoes, instead, there are still no conclusive studies. Research is mainly focused on the characterization of individual dynamics and overall swarm structure [19–21], as well as on the mechanisms of courtship and mating [22–24], with only a very few studies specifically addressing male-male interaction, [25–27]. Three dimensional field data on small swarms (less than 100 individuals) of *Anopheles gambiae* and *Anopheles coluzzii* [25, 26] suggest the presence of a short-range interaction, which leads to alignment in the direction of flight. A different form of interaction, mediated by the mosquito auditory system, has been proposed in *Aedes aegypti* [27], where the emergence of acoustic order is observed in short arrays (up to seven individuals) of tethered mosquitoes.

Here, with a unique dataset of 27 swarms, ranging from 80 to 400 individuals, we investigate swarming behavior in laboratory *Anopheles gambiae* mosquitoes. Using a statistical physics approach, we provide experimental evidence of a male-male interaction that, unlike previous studies [11, 25, 26], does not act on mosquitoes' direction of flight, but rather produces a long-range correlation of their speed.

2 Results

2.1 Experiments and data

We perform laboratory experiments on mosquitoes, *Anopheles gambiae* (strain G3), swarming in a large cage (a room $5.0m \times 3.5m \times 2.6m$), where we artificially reproduce visual stimuli needed for mosquitoes to swarm, see Methods and [20].

Data are collected with a $3D$ camera system: three synchronized high-speed cameras (IDT-M5) recording the same swarm from different positions, so that in post-processing, with the computer vision software GReTA [28], we are able to reconstruct individual trajectories of each individual in the swarm.

Data presented in the main section of this work comprise 27 swarms, ranging from 80 to 400 individuals. A distinct dataset (41 smaller swarms ranging from 10 to 80 individuals) is included in SI, to highlight the relevance of swarm size for behavioral analysis. Data belong to two different experimental campaigns: the first conducted in 2020/2021 with cameras shooting at 170fps and acquisitions lasting $\sim 15s$, the second conducted in 2024 at 80fps with acquisitions lasting $\sim 30s$.

2.2 Localization within the swarm

The fundamental assumption underpinning most advanced models for swarming dynamics [29], and more generally of statistical physics, is that individuals within a group are indistinguishable. At a given instant in time they may be located in different positions and display different kinematic attributes, but these differences are solely due to statistical fluctuations, and essentially vanish when observing the system over time. In contrast, we find that, despite being part of a swarm, mosquitoes maintain their individuality and exhibit specific characteristics that are not compatible with purely statistical fluctuations.

The analysis of individual trajectories shows that each mosquito performs consecutive loops, [20], around a specific position, slightly different from that of the other members of the group. In performing these loops, it explores a small portion of the volume occupied by the entire swarm, see Fig.1. Similarly, we find that each mosquito flies at its own characteristic speed, and it samples only

Fig. 1 Spatial localization. Swarm 20200928_ACQ3. Number of mosquitoes ~ 200, duration 2700 frames at 170fps, corresponding to 15.88s. **a.** Top view (xz-plane). In dark gray the $3D$ reconstruction of the swarm. The red circle represents the swarm centroid. The yellow circles indicate individual centroids, one for each mosquito in the swarm. **b.** Top view, xz–plane. In dark gray the $3D$ reconstruction of the swarm. In turquoise, yellow and purple three different trajectories highlight the spatial individual localization. Mosquitoes do not explore the entire volume occupied by the swarm, but only a small portion of it. The three black circles represent the individual centroids of the three trajectories. **c.** On the horizontal plane, the group spatial distribution (dark gray) is much broader than the individual spatial distribution (purple). Group radius is equal to $R = 0.24 \pm 0.14$m, the individual radius $r = 0.13 \pm 0.07$m. The vertical lines and vertical shaded areas represent the mean and standard deviation, gray for the group and purple for individuals. On the right of the two distributions, a schematic description of the two variables r (indicated by white within purple circle) and R (indicated by black within gray circle), where the large dark gray circle represents the swarm and the three small circles the volumes occupied by the three trajectories highlighted in yellow, purple and turquoise in panel b. **d.** On the horizontal plane, individual radii show low variability within the trajectories of the swarm, and are consistently smaller than the group radius (the black dashed line.) **e.** On the horizontal plane, the ratio between individual and group radii, r/R, is approximately equal to 0.5 (the black dashed line) in all the 27 swarms of the dataset, regardless of the number of mosquitoes in the swarm, N. **f.** Side view (xy-plane). In dark gray the $3D$ reconstruction of the swarm. The red circle represents the swarm centroid. The yellow circles indicate individual centroids, one for each mosquito in the swarm. **g.** Side view. In dark gray the $3D$ reconstruction of the swarm. In turquoise, yellow and purple three different trajectories, the same ones shown in panel b, highlight the spatial individual localization. **h.** Along the vertical axis, the group spatial distribution (dark gray) is much broader than the individual spatial distribution (purple). Group altitude is equal to $H = 0.16 \pm 0.10$m, the individual altitude $h = 0.05 \pm 0.03$m. The vertical lines and vertical areas represent the mean and standard deviation, gray for the group and purple for individuals. On the right of the two distributions, a schematic description of the two variables h (indicated by white within purple circle) and H (indicated by black within gray circle), where the large dark gray rectangle represents the swarm and the three small rectangles the volumes occupied by the three trajectories highlighted in yellow, purple and turquoise in panel g. **i.** Along the vertical axis, individual altitudes show low variability within the trajectories of the swarm, and they are consistently smaller than the group altitude (the black dashed line.) **j.** Along the vertical axis, the ratio between individual and group altitudes, h/H, is approximately equal to 0.5 (the black dashed line) in the 27 swarms of the dataset, regardless of the number of mosquitoes in the swarm, N.

a narrow range of the speeds observed at a group level, see Fig.2.

This leads to both spatial and speed *localization*.

Spatial localization. We give a quantitative description of the spatial localization, by estimating and comparing the extent of the swarm as a whole and that of individual trajectories.

To this aim, we compute the swarm centroid (the red circle in Fig.1a,f), averaging over the positions of all mosquitoes at all instants of time. We then measure the distance of each mosquito from

the swarm centroid, and we define the *group* spatial distribution as the probability distribution of these distances, see Methods. This spatial distribution describes how frequently an individual is found at a given distance from the swarm center. Its width provides a rough estimate of the spread of the swarm, while its mean is a measure of the swarm size.

In a similar way, we compute the centroid of each trajectory (the yellow circles in Fig.1a,f - one for each mosquito - and in the black circles in Fig.1b,g). We then measure the distance of each mosquito from its *own* centroid, and we define the *individual* spatial distribution as the probability distribution of these distances, aggregating all mosquitoes in the swarm, see Methods. The width of the individual spatial distribution gives a rough estimate of the typical spread of single trajectories, while its mean is a measure of the size of the region explored by each individual.

Mosquitoes move predominantly on the horizontal plane (xz-plane), with limited changes in altitude (y-axis), [20]. Because of this anisotropy, we measure group and individual spatial extents on the horizontal plane (radius) separately from the elongation along the vertical axis (altitude).

On the horizontal plane, the group spatial distribution is significantly broader than the individual one, see Fig.1c. For the swarm shown in Fig.1, the group radius is $R = 0.24 \pm 0.13$m, while the individual one is $r = 0.13 \pm 0.07$m, with very low variability between mosquitoes, see Fig.1d. This results in a ratio $r/R \sim 0.5$. Similarly, along the vertical axis, the swarm is much more elongated than individual trajectories. The typical vertical extent for the swarm in Fig.1h is equal to $H = 0.16 \pm 0.10$m, while the individual elongation is $h = 0.05 \pm 0.03$m, again with low variability between different mosquitoes and with a ratio $h/H \sim 0.5$, Fig.1i.

By performing this same analysis over all the swarms of our dataset, we find that r/R and h/H are consistently close to 0.5, see Fig.1e,j. This confirms that mosquitoes do not explore the entire volume occupied by the swarm, but instead remain localized in a small region, approximately one eighth of the total volume.

Speed localization. Through the analysis of individual speed, we find a phenomenon that, by

Fig. 2 Speed localization. a. In dark gray a typical group probability distribution. In purple, individual probability distributions of three different mosquitoes, whose mean speeds are highlighted with the purple vertical lines. **b.** The probability distribution of the individual mean speeds for the 200 mosquitoes belonging to the swarm 20200928_ACQ3. Mean speeds are in the range between 0.37m/s and 0.75m/s. This distribution displays large fluctuations, with standard deviation equal to 0.06m/s, highlighted with the light purple stripe. **c.** The fluctuations of individual mean speeds measured from the data (σ_S, in purple) and estimated from the group probability distribution (Σ, in gray) for all the swarms of our dataset, as a function of the size of the system. Measured standard deviations are consistently greater, up to ten times larger, than their expected values in the absence of localization. **d.** For the same swarm shown in panels a and b, the probability, P_0, that individual speed fluctuations are positive as a function of δs_i. P_0 spans the entire range from 0.1 to 0.9, confirming speed localization in mosquitoes. In the absence of localization, P_0 is equal to 0.5, the dashed horizontal line in the plot. As expected, P_0 increases with increasing δs_i. In the inset: the group probability distribution of the fluctuations.

analogy to what we have already observed for mosquito positions, we refer to as *speed localization*: individual mosquitoes do not explore the full range of speed covered by the entire swarm, but only a portion of it.

A first hint of speed localization emerges from the comparison between individual and group probability distributions, see Fig.2a, where we show in dark gray the group distribution and,

in purple, individual distributions for three different mosquitoes. In the absence of localization, we would expect the group and individuals to share the same statistical properties. All individuals would display a similar speed over time, so that their probability distributions would be identical and coincide with that of the group. Under this assumption, variations in individual mean speeds whose distribution is presented in Fig.2b, would reflect only statistical fluctuations and could be estimated by the standard error of the group distribution, Σ, see Methods. Having access to full trajectories, we are also able to measure these variations directly from the data, in the form of the standard deviation of individual mean speeds, σ_S, see Methods.

The comparison of these two quantities indicate that in our swarms the variation of the mean speed across individuals cannot be explained by statistical fluctuations alone. The values of σ_S, in the interval between 0.06m/s and 0.13m/s, are indeed not compatible with those of Σ, which we find in the interval between 0.01m/s and 0.02m/s, see Fig.2c.

The effect of speed localization becomes clearer when shifting our focus from absolute speeds to deviations from the group average. In particular, the speed fluctuation of mosquito i at time t is defined as:

$$\delta s_i(t) = s_i(t) - S(t) \tag{1}$$

where $S(t)$ represents the group average speed at time t, $s_i(t)$ the speed of mosquito i at time t, and $\delta s_i(t)$ its fluctuation. This transformation moves us from speed — always a positive quantity — to its fluctuation, which can be positive or negative. If a mosquito is flying faster than the group average, the fluctuation is positive; if slower, it is negative.

At a group level, the probability distribution of the fluctuations is centered around 0m/s, and it is nearly symmetric, see the inset of Fig.2d: At any one time, there is approximately a 50% chance of finding a positive fluctuation (a mosquito flying faster than the group average), and a 50% chance of a negative fluctuation (a mosquito flying slower than the group).

But when we examine speed fluctuations at an individual level, this symmetry breaks down. Individual distributions are not centered around zero, but they are unbalanced toward positive or negative fluctuations. This individual asymmetry is the hallmark of speed localization.

We quantify the asymmetry by computing, for each mosquito, the probability, P_0, that its speed fluctuations are positive. We essentially evaluate the proportion of time (relative to each trajectory's duration) that each mosquito spends flying faster than the group average. This corresponds to the area, highlighted in dark purple in the histograms in Fig.2d, under the distribution to the right of the y-axis, see Methods. In the absence of localization, P_0 should be approximately 0.5. We find instead that P_0 spans the full range from 0.1 - mosquitoes that fly slower than the group average for most of the time - to 0.9 - mosquitoes that fly faster than the group average for most of the time. This wide variation confirms the presence of speed localization in our swarms.

2.3 Experimental evidence of male-male interaction

In biological systems, interaction is typically driven by mechanosensory processes, involving vision, hearing, and other senses [30–33]. Individuals perceive the other members of the group, and adjust their own behavior, synchronizing their state. The physical limit of sensory perception defines the *interaction range*, which may be quite short (in mosquitoes hearing range is of the order of 10cm [34]). Despite the limited interaction range, information can travel much greater distances, thanks to a sort of *word of mouth* mechanism, see Fig.3a: A interacts with B, and B also interacts with C, who lies outside A's interaction range. By interacting, A and B adjust their behavior, as do B and C. As a result, the states of the three individuals are synchronized. Thanks to the indirect exchange of information between A and C, mediated by B, the system exhibits correlation over a distance (that between A and C) that exceeds the interaction range. In physics, this indirect synchronization, produced by a chain of interactions, is referred to as correlation.

Interaction and correlation are distinct, yet closely related, concepts. They both refer to an exchange of information that is direct in the case of interaction, but indirect in the case of correlation. Measuring how mosquitoes react to each other is relatively simple with tethered

mosquitoes [27, 35]. However, this kind of measurement is much more complicated in free–flying swarms. In these specific situations, where a direct measure of interactions is not feasible, correlation functions provide a powerful tool to reveal the presence of interaction within the system.

Fig. 3 Speed correlation. a. Interaction vs correlation. Interaction and correlation are both related to a transfer of information. Despite the short range of interaction, information can travel greater distances: A interacts with B. B interacts also with C, which exceeds A's interaction range. Due to the interaction, A and B synchronize their speed, so do B and C. As a result, the three particles are synchronized, hence correlated. The correlation range of the system is defined by the distance between A and C, that is larger than the interaction range. D exceeds the interaction ranges of the other particles. **b.** The group distribution of the speed fluctuations, with light and dark orange differentiating between negative and positive fluctuations. Two particles having same sign of fluctuations produce a positive correlation coefficient c_{ij}. **c.** The correlation function has a non–trivial trend. It is positive for short distances. It decreases at increasing distances, and crosses the x–axis at r_0, which defines the correlation range. r_0 is much larger than the nearest neighbor distance, d_{NN}. The correlation length, ξ is computed following [4]. Inset: correlation length, ξ, scales linearly with the correlation range ($\rho = 0.99$, p-value $< 10^{-6}$). **d.** Across all swarms of the dataset, the correlation length, ξ, scales linearly with the size of the system, L ($\rho = 0.85$, p-value $< 10^{-6}$). In the plot, bars are standard deviations. This linear trend represents the finite size expression of the scale–free property of the correlation function: there is no typical correlation length for our swarms; but the only relevant scale for the system is its size.

Spatial correlation functions of the speed fluctuations.

We measure whether and to what extent (in space), mosquitoes tend to synchronize deviations in their speed from the group average, by computing the spatial correlation function of the speed, in its connected declination, [11, 36]. It is worth noting that, unlike previous studies on insect swarms [11, 16], we investigate the correlation function of the speed, which does not refer to the direction of flight, as the velocity does. As shown in SI, our swarms display no significant correlation of the velocity.

We quantify the synchronization between two mosquitoes, by computing their correlation coefficient:

$$c_{ij}(t) = \delta s_i(t) \cdot \delta s_j(t) \qquad (2)$$

where $\delta s_i(t)$, $\delta s_j(t)$, defined as in eq.(1), denote the speed fluctuations of individuals i and j at time t. This product is positive if fluctuations are both positive or both negative (both mosquitoes flying faster or both fly slower than the group average), see Fig.3b. Conversely, it is negative if one is positive and the other is negative (one mosquito fly faster than the group average, the other flying slower). Therefore, a positive c_{ij} indicates a tendency towards synchronization.

We can now group pairs of mosquitoes based on their mutual distance, and compute the average of the correlation coefficient over each bin. With this procedure we associate a correlation value with each distance, thereby defining the spatial correlation function, $C(r)$ (see Methods for a formal definition). The $C(r)$ essentially defines what is the typical correlation coefficient of two mosquitoes at a mutual distance r. Across all swarms of our dataset, it exhibits a characteristic trend: it is positive for small values of r, meaning that mosquitoes in close proximity display similar speed fluctuations. It then decays with increasing distance, crossing the x–axis at a distance r_0, see Fig.3c, which defines the correlation range and gives a rough estimate of the size of the correlated domains.

We also give an estimate of the size of the correlated domains as the standard correlation length, ξ, that we compute using the integral form in [4]. This definition of ξ is general and robust, since it takes into account the entire shape of the correlation function up to r_0. In our case, across all swarms, ξ grows linearly with r_0, see

inset in Fig.3c, varies in the range between 0.04m and 0.16m (see Table S1) and it scales linearly (Pearson coefficient $\rho = 0.85$, $p-Value < 10^{-6}$) with the size of the swarm, L, see Fig.3d. This linear scaling is the finite-size expression of what is known in physics as *scale-free* correlation, a property also observed in other biological systems [4, 11]. It implies that there is no characteristic distance beyond which information transfer becomes negligible. Rather, the only relevant scale is the system size itself — meaning that information can propagate across the entire system, regardless of its size.

Evidence of interaction. In the absence of spatial and speed localization, non–trivial and scale–free spatial correlation would suggest that an *effective* interaction, acting on the speed domain, is at work in swarms of mosquitoes. But this correlation inherently involves both space and speed. In a system such as ours, that exhibits localization, it is then natural to wonder whether correlation is the genuine expression of interaction, or it is merely an epiphenomenon of the system localization properties.

To discriminate between these two scenarios, we need to test our data against a null case: an artificial swarm of particles that do not interact with each other, but do exhibit spatial and speed localization. We generate artificial swarms directly from the data. Specifically, we disrupt potential interaction links through a shuffling procedure applied to the mosquitoes' trajectory centroids, see Fig.4a for an example and Methods for formal details. With this procedure, artificial swarms present the same localization parameters, and the same speed probability distributions as real swarms, from which they differ only in the mosquitoes' positional arrangement.

For each real swarm, we generate 10^6 shuffled instances, and we compute their correlation ranges, r_{0S}. The probability distribution of r_{0S}, in Fig.4b, shows that in most of the cases r_{0S} is smaller than the distance to the nearest neighbor, d_{NN}, indicating that the correlation is negligible. From this same distribution we measure the probability $P(r_{0S} \geq r_0)$ that the correlation range is larger in shuffled swarms that in real ones, the purple area in Fig.4b. In doing so, we essentially compute the p–value for the correlation range.

Fig. 4 Evidence of interaction. a. We generate artificial swarms of non–interacting particles, that present the same localization properties and speed distributions of real swarms. We shift mosquito positions following a shuffling procedure on mosquito centroids. In the example, the centroid of the turquoise trajectory is assigned to the yellow mosquito, which is then shifted from the center of the swarm to the upper right corner; the centroids of the yellow and purple trajectories are assigned to the purple and turquoise mosquitoes respectively. The yellow and purple mosquitoes, originally close and potentially interacting, are located in two distant regions of the shuffled swarm, and thus no longer interact. **b.** The probability distribution of the correlation range for the 10^6 correlation functions of the shuffled swarms. In 80% of the cases, the system presents a negligible correlation, with a correlation range smaller than the nearest neighbor distance, d_{NN}. The area (highlighted in purple) below the distribution, on the right of the correlation range of the original real swarm, represents the probability of finding in the artificial swarm a correlation range larger than that of the original swarm. **c.** The logarithm of the probability $P(r_{0S} > r_0)$ for all the swarms of the dataset. Probability values are between $7 \cdot 10^{-6}$ and $1 \cdot 10^{-2}$, always smaller than the p–value threshold for significance of 0.05 (highlighted with the horizontal line in the plot). This suggests that the correlation range observed in real swarms is not compatible with a random arrangement of mosquitoes, providing evidence that the observed correlations are not merely a consequence of localization, and thus strongly indicate male–male interaction in our swarms.

In all the swarms of our dataset, we find values between $7 \cdot 10^{-6}$ and $1 \cdot 10^{-2}$, as shown in Fig.4c. Being below the standard threshold for p–value significance of 0.05, these results suggest that the correlation range observed in real swarms

is wholly incompatible with a random arrangement of mosquitoes, providing strong evidence that the observed correlations are not merely a consequence of localization. Rather, they reflect genuine interactions between mosquitoes within the swarm.

In concluding this Section, let us emphasize that the key to our results lies in the type of data we are working with: a variety of large swarms. As we show in SI, working only on smaller swarms (from 10 to 80 mosquitoes) the effect of interaction would not have been detectable, nor would it have been possible to assess the scale-free property of the correlation.

3 Discussion

With our work we take the first step towards a more complete understanding of swarming behavior in mosquitoes *Anopheles gambiae* (strain G3). While work remains to be done, our results pave the way to the design of new experimental investigations, and the development of new models of mosquito dynamics.

The two key features we identify - localization and speed correlation - both contribute to placing mosquito laboratory swarms within a framework distinct from that of previously studied systems, such as bird flocks, fish schools and, most notably, midge (*Diptera Chironomidae*) swarms, [4, 5, 11]. In these systems, correlation is primarily observed in the velocity domain, involving the direction of motion, and it is accompanied by speed correlation (see SI for an example comparing velocity and speed correlation in midge swarms). In contrast, our data exhibit strong correlation in speed, but no significant correlation in velocity (see SI). This difference may stem from biological factors, specifically from interaction mechanisms that depend on the system's taxonomic class and species. However, especially in comparison with midge swarms, it is also plausible that environmental conditions — laboratory versus field — influence dominant behavioral traits, such that some aspects of interaction are inhibited under laboratory conditions, which calls for benchmarking our dataset against field and semi–field experiments.

Localization and correlation of the speed require the design of novel models for swarm dynamics. Standard models assume an interaction acting on the velocity and homogeneity among individuals [29], such that differences, at a fixed instant in time, are solely due to statistical fluctuations. Localization instead implies heterogeneity and temporal persistence: individuals exhibit characteristic positions and speeds, which are slightly different from those of the other members of the group, and are maintained over time. From a biological perspective, localization may be seen as a natural property of the system. Environmental and genetic factors lead to differentiation of individual traits, which results in specific kinematic properties. As with humans - who may walk at different speeds depending on factors such as age, weight, or overall physical condition — mosquitoes could also exhibit preferences for specific locations and speeds.

However, the strong correlation we find and, even more, the negligible correlation in the shuffled swarms, indicate that mosquito spatial arrangement is not random, suggesting that localization is enhanced by interaction between individuals. Mosquitoes are likely to have individual specific speed determined by biological factors, but when they are in a swarm they are influenced by the presence of other individuals and tend to synchronize their speed, much as people do when walking together. Through a mechanism, which may be modeled as an imitation process, they modulate their speed to reduce differences with their neighbors, similarly to what is observed in bird flocks and fish schools [4, 5].

This imitation mechanism may appear to contradict speed localization: if mosquitoes synchronize their speed, we expect that they all tend to display the same speed probability distribution, hence they should not exhibit localization. This apparent contradiction may be explained with an interaction mechanism with low resolution. Due to mechanosensory constraints, mosquitoes may not be able to perceive the difference in speed from that of their neighbors, when it falls below a certain perception threshold. As a result, only when the difference exceeds this threshold, mosquitoes perceive themselves as out of sync and adjust their speed — such that the differences fall below the perception threshold. The system can therefore show synchronization and, at the same time, allow mosquitoes to keep their individuality within the swarm.

An interaction based on the speed, combined with a perception threshold, could serve as a effective mechanism when designing a model. But from a biological perspective, there is an alternative interpretation of our data: the observed correlation in speed may not emerge from an interaction directly acting on speed itself, but rather may be an epiphenomenon of a different kind of interaction, which indirectly forces individual speeds to be synchronized.

The parallel with the acoustic mechanism is compelling, but still not robustly supported by experimental data. The mosquito auditory system naturally presents the perception threshold we introduced above to reconcile localization and correlation [23]. More interestingly, mosquitoes play an active role in shaping the sound they perceive [37], in such a way that, by modulating their own wing beat frequencies, they can effectively modulate the environmental sounds to which they can respond. Through an acoustic interaction, males could then transform the noisy environment of the swarm, generated by the sounds of hundreds of individuals, into one where only biologically relevant signals (i.e., those from females) stand out, thereby enhancing mating efficiency and offering a great biological advantage to the group.

The link between acoustic and speed interaction lies in mosquito wing beat frequency, which defines the sound the mosquito emits and also affects their speed. Although a relation between wing beat frequency and speed appears reasonable, empirical data of simultaneous measurements in the context of swarms are still lacking. This gap highlights the need for new experiments aimed at determining whether wing beat frequency and speed are positively correlated when mosquitoes participate in a swarm—an essential step to confirm our biological interpretation.

4 Methods

In this section, we aim to provide all the formal definitions and computational details related to the results presented in Section 2. To facilitate the reader, the outline of this section reproduces the structure of Section 2, with a first subsection on the experiments, a second on the swarm localization properties, and a third on the experimental evidence of interaction.

4.1 Experiments

We performed $3D$ swarming experiments on mosquitoes *Anopheles gambiae* (*An. gambiae*) G3 strain (MRA-112). Mosquitoes were contained in a containment level 2 facility at the Department of Medicine and Surgery, Perugia, Italy, authorization N. PG/IC/Imp2/13/001-Rev2 from Ministry of Health.

As in [20], mosquitoes are reared using the MR4 protocol [38], with temperature at 28 ± 1°C and $75 \pm 10\%$ relative humidity under a $12/12 - h$ light/dark cycle. Details on feeding procedure and on the husbandry techniques may be found in [20].

Males and females are manually separated as pupae by microscopic examination of the terminalia and raised in separate cages, ensuring all males are virgins. For each experiment, between 800 and 1800 $2 - 3$-day-old males were released from 'Bug-Dorm' type cages approximately 1 h before the onset of the artificial sunset, and left in the large swarming cage containing multiple resting sites and sources of sugar/honey solution as described previously in [20], for the next few days for data collection.

Swarming arena setup. As described elsewhere [20], data collection took place in the Insectary Field 1 (measuring $8.5m \times 3.8m \times 3m$, $L \times W \times H$) at the Department of Medicine and Surgery of the University of Perugia, which contains one large cage of $5.0m \times 3.5m \times 2.6m$). Since swarms naturally occur at dusk, the swarming arena is illuminated using six 850nm IR LED lamps (RAYMAX 300). We reproduced the visual stimuli needed for mosquitoes to swarm; specifically, see [20]: i. a contrasting ground marker ($55cm \times 55cm$), at approximately 1.1m from the back wall; ii. two series of lights: three $2700K$ 8W lights located on the floor outside the cage; iii. three 2700K 9W LED lights placed on the floor at the back of the cage to simulate twilight and an artificial horizon, see [20] for a detailed scheme of the setup.

Data collection. We collect $3D$ data with a system of high–speed synchronized cameras (IDT-MotionScope M5) equipped with Schneider Xenoplan 28mm f/2.0 optics. Each camera is placed

on a plexiglass shelf, supported by a metal tripod (Manfrotto Pro Digital Tripod 475B), allowing the correct orientation of the cameras towards the swarming area. The three cameras are located outside of the large cage at a distance of $3.5 - 3.9$m from the ground marker. Cameras record the same swarm from different positions, so that, using the $3D$ tracking software GReTA [28], we are able to reconstruct three dimensional trajectories of each individual in the swarm. We calibrate the mutual position and orientation of the cameras after each experiment, following the post-calibration procedure defined in [39]; camera internal parameters (focal length, distortion coefficient and image center) are instead calibrated using a standard method [40]. As indicated in the text and in Supp. Table 1, two modes of data acquisition were employed: 1. using a frame rate of 170fps of duration approx. 15.9s 2. using a frame rate of 80fps and duration approx. 33 s.

Notations

We summarize here the notations we will use in the following two subsections:

- N represents the number of mosquitoes in the swarm;
- $i = 1, \ldots, N$ refers to a specific individual in the swarm;
- T_i, with $i = 1, \ldots, N$, refers to the duration of the trajectories of mosquito i, which is not always as long as the entire acquisition ($T_i \leq T$);
- $\vec{x}_i(t)$ is the $3D$ position vector of mosquito i at time t, expressed in m. In terms of the $3D$ coordinates, $\vec{x}_i(t) = (x_i(t), y_i(t), z_i(t))$;
- $\vec{v}_i(t)$ is the $3D$ velocity vector of mosquito i at time t, expressed in m/s, and computed from the spatial displacement between two consecutive frames with the following expression:

$$\vec{v}_i(t) = \frac{\vec{x}_i(t+1) - \vec{x}_i(t)}{\Delta t} \qquad (3)$$

with $\Delta t = 1/80$s or $\Delta t = 1/170$s, depending on the data;

- $s_i(t)$ is the speed of mosquito i at time t, namely the magnitude velocity vector $s_i(t) = |v_i(t)|$.

4.2 Localization within the swarm

Spatial localization. We quantify the spatial localization in our swarms, by computing group and individual spatial distributions on the horizontal plane and along the vertical axis.

Group spatial distributions - Swarm extent. We compute the swarm centroid, $\vec{X}$, as the average over the position of all mosquitoes at all instants of time:

$$\vec{X} = \frac{1}{N} \sum_{i=1}^{N} \frac{1}{T_i} \sum_{t=1}^{T_i} \vec{x}_i(t), \qquad (4)$$

and $\vec{X} = (X, Y, Z)$.

We measure the distance of each mosquito from $\vec{X}$ on the horizontal plane:

$$R_i(t) = \sqrt{(x_i(t) - X)^2 + (z_i(t) - Z)^2}. \qquad (5)$$

On the set of all these distances, we define the group probability distribution, $P_{\mathcal{R}}(R)$, the one highlighted in dark gray in Fig.1c.

We measure the radius of the swarm as the mean value, R, of the distribution $P_{\mathcal{R}}(R)$:

$$R = \frac{1}{N \sum_{i=1}^{N} T_i} \sum_{i=1}^{N} \sum_{t=1}^{T_i} R_i(t). \qquad (6)$$

R varies from swarm to swarm, ranging from 0.19m to 0.45m, see Table S1.

We measure the distance of each mosquito from $\vec{X}$ along the y-axis:

$$H_i(t) = \sqrt{(y_i(t) - Y)^2}. \qquad (7)$$

On the set of all these distances, we define the group probability distribution, $P_{\mathcal{H}}(H)$, the one highlighted in dark gray in Fig.1h.

We measure the elongation of the swarm as the mean value, H, of the distribution $P_{\mathcal{H}}(H)$:

$$H = \frac{1}{N \sum_{i=1}^{N} T_i} \sum_{i=1}^{N} \sum_{t=1}^{T_i} H_i(t) \qquad (8)$$

H varies from swarm to swarm, ranging from 0.10m to 0.19m, see Table S1.

Individual spatial distributions - Trajectory extent. We compute the centroid of each

trajectory, $\vec{x}_i$, by averaging the position of each individual over time:

$$\vec{x}_i = \frac{1}{T_i} \sum_{t=1}^{T} \vec{x}_i(t), \qquad (9)$$

and $\vec{x}_i = (x_i, y_i, z_i)$.

We then compute the distances of each mosquito from its own centroid, which we denote by $r_i(t)$:

$$r_i(t) = \sqrt{(x_i(t) - x_i)^2 + (z_i(t) - z_i)^2}, \qquad (10)$$

when the distance is computed on the horizontal plane, and by $h_i(t)$:

$$h_i(t) = \sqrt{(y_i(t) - y_i)^2}, \qquad (11)$$

when the distance is computed on the vertical axis.

We define the radius of trajectory i as:

$$r_i = \frac{1}{T_i} \sum_{t=1}^{T_i} r_i(t), \qquad (12)$$

and the elongation of trajectory i as:

$$h_i = \frac{1}{T_i} \sum_{t=1}^{T_i} h_i(t) \qquad (13)$$

In Fig.1d,i we show, with dark purple circles, the set of r_i and h_i for all the mosquitoes in one of our swarms. With the light purple shading we are instead showing their standard deviation:

$$\delta r_i = \sqrt{\frac{1}{T_i - 1} \sum_{t=1}^{T_i} (r_i(t) - r_i)^2}, \qquad (14)$$

and

$$\delta h_i = \sqrt{\frac{1}{T_i - 1} \sum_{t=1}^{T_i} (h_i(t) - h_i)^2}. \qquad (15)$$

Finally, we aggregate all individual distances $r_i(t)$ and $h_i(t)$ obtaining the two individual distributions, $P_r(r)$ and $P_h(h)$, shown in purple in Fig.1c,h. For each swarm, we then define the typical trajectory radius, r, and the typical elongation, h, as:

$$r = \frac{1}{N \sum_{i=1}^{N} T_i} \sum_{i=1}^{N} \sum_{t=1}^{T_i} r_i(t) \qquad (16)$$

and

$$h = \frac{1}{N \sum_{i=1}^{N} T_i} \sum_{i=1}^{N} \sum_{t=1}^{T_i} h_i(t). \qquad (17)$$

We compare the spatial extent of individual trajectories with that of the swarm, by computing the two ratios r/R (from eq.(16) and eq.(6)) and h/H (from eq.(17) and eq.(8)) that we show in Fig.1e,j as a function of the number of mosquitoes in the swarm.

Speed localization. We consider the set $\mathcal{S}_S = \{s_i(t), i = 1, \ldots, N \text{ and } t = 1, \ldots, T\}$, of the speeds of all mosquitoes at all instants of time. From this set, we define the group probability distribution, $P_S(s)$, (highlighted in dark gray in Fig.2a).

With a similar procedure we can also define individual probability distributions. We consider the sets $\mathcal{S}_i = \{s_i(t), t = 1, \ldots, T\}$, of the speed of mosquito i at all instants of time. From these sets, we define individual probability distributions, $P_{S_i}(s)$, one for each individual. In Fig.2a, we show the distributions of three different mosquitoes in purple.

For each mosquito i, we compute the mean speed as:

$$s_i = \frac{1}{T_i} \sum_{t=1}^{T_i} s_i(t). \qquad (18)$$

We define the distribution of these mean values, which we show in Fig.2b, where in light purple we highlight its standard deviation σ_S.

We also estimate the standard deviation of the mean from the group distribution, as the standard error, Σ:

$$\Sigma = \frac{\sigma}{\sqrt{N}} \qquad (19)$$

In the absence of speed localization, the two values (σ_S and Σ) should be compatible, and variations in individual mean speeds would reflect only statistical fluctuations.

Probability of positive speed fluctuations. To better quantify the localization, we move from absolute speed to their fluctuations with respect to the swarm average. To this aim, at each instant in time, we define the swarm average as:

$$S(t) = \frac{1}{N} \sum_{i=1}^{N} s_i(t), \qquad (20)$$

and the individual speed fluctuations for mosquito i as:

$$\delta s_i(t) = s_i(t) - S(t). \qquad (21)$$

From the set of all the fluctuations, we compute the probability distribution, see inset of Fig.2d. By definition the mean value of this distribution is equal to 0m/s.

For each individual, we compute the mean of its fluctuation, δs_i, as:

$$\delta s_i = \frac{1}{T_i} \sum_{t=1}^{T_i} \delta s_i(t). \qquad (22)$$

Surprisingly, these individual mean values are generally different from 0m/s, and are not symmetric with respect to the zero.

We give a measure of the individual asymmetry by computing, for each mosquito, the proportion of time that it spends having positive fluctuations. These quantity, denoted by P_{0i}, is computed as follows:

$$P_{0i} = \frac{1}{T_i} \sum_{t=1}^{T_i} \theta(\delta s_i(t)) \qquad (23)$$

where $\theta(\delta s_i)$ is the Heaviside step function defined as:

$$\theta(\delta s_i) = \begin{cases} 1 & \text{if } \delta s_i(t) \geq 0 \\ 0 & \text{otherwise .} \end{cases} \qquad (24)$$

In the absence of localization, P_{0i} should be equal to 0.5: a mosquito should spend half of the time flying faster than the group average (positive fluctuations) and half of the time flying slower (negative fluctuations). We find instead that in our swarms, it spans the entire range between 0.1 and 0.9, as shown in Fig.2d, where we plot the values of P_{0i} as a function of δs_i.

4.3 Experimental evidence of male–male interaction

Spatial correlation functions of the speed fluctuations. We start by computing the speed fluctuation of each individual at each instant in time, as in eq.(21). Then, for each pair of mosquitoes (i,j), we compute:

• the correlation coefficient defined as in eq.(2), $c_{ij}(t) = \delta s_i(t) \cdot \delta s_j(t)$;

• the mutual distance $r_{ij}(t) = |\vec{x}_i(t) - \vec{x}_j(t)|$ where $|\cdot|$ indicates the vector magnitude.

We then define the correlation at time t as:

$$\hat{C}(r,t) = \frac{\sum_{i,j=1}^{N} c_{ij}(t)\delta(r, r_{ij}(t))}{\sum_{i,j=1}^{N} \delta(r, r_{ij}(t))} \qquad (25)$$

where $\delta(r, r_{ij})$ is the Kronecker delta function:

$$\delta(r, r_{ij}(t)) = \begin{cases} 1 & \text{if } r_{ij}(t) = r \\ 0 & \text{otherwise .} \end{cases} \qquad (26)$$

With the definition in eq(25), we are averaging the correlation coefficients over pairs at mutual distance equal to r. Specifically, the factor $\delta(r, r_{ij}(t))$ selects only those pairs for which $r_{ij}(t) = r$, neglecting the contribution of all other pairs. In the numerator, we are then summing the correlation coefficient of all those pairs, while in the denominator we are counting how many they are.

The $\hat{C}(0,t)$ is a special case: at a fixed instant in time, the only pairs that participate in $\hat{C}(0,t)$ are the ones at a mutual distance $r = 0$, hence only the pairs where $i = j$:

$$\hat{C}(0,t) = \frac{1}{N} \sum_{i=1}^{N} c_{ii}(t) = \frac{1}{N} \sum_{i=1}^{N} \delta s_i^2(t) \qquad (27)$$

which is the average square fluctuation of the speed at time t.

We use $\hat{C}(0,t)$ to define the normalized correlation function at time t, $C(r,t)$, as follows:

$$C(r,t) = \frac{\hat{C}(r,t)}{\hat{C}(0,t)}. \qquad (28)$$

This normalized correlation function is such that $C(0,t) = 1$.

Finally, we average over time and obtain the spatial correlation function we show in Fig.3:

$$C(r) = \frac{1}{T} \sum_{t=1}^{T} C(r,t). \qquad (29)$$

When working on finite systems, the definition of the correlation function in eqs.(25-28) has two limitations. The first is that we cannot compute the $C(r)$ on continuous values: the mutual distances r_{ij} define a finite set of values (precisely

$N \cdot (N-1)$). The second is that if we strictly use the definition in eq.(25), we will not have statistically relevant samples, because for each value of r we will have essentially only one pair: it is very unlikely that two (or more) pairs are located exactly at the same mutual distance.

To overcome these two issues, we perform a binning in r of the $\hat{C}(r,t)$, and consequently of the $C(r)$. More specifically, we fix the size of the bin, Δr, and we define the discrete correlation function as:

$$\hat{C}(r_k, t) = \frac{\sum_{i,j=1}^{N} c_{ij}(t)\hat{\delta}(r_k, r_{ij}(t))}{\sum_{i,j=1}^{N} \hat{\delta}(r_k, r_{ij}(t))} \qquad (30)$$

where:

- $k = 1, \ldots, k_M$;
- k_M is defined as the smallest integer such that $k_M \Delta r$ is greater or equal to the size of the swarm, L, hence $k_{M-1}\Delta r < L \le k_M \Delta r$;
- $\hat{\delta}(r, r_{ij}(t))$ is a Kronecker delta function modified as follows:

$$\hat{\delta}(r_k, r_{ij}(t)) = \begin{cases} 1 & \text{if } r_{ij}(t) \in (r_k - \Delta r, r_k] \\ 0 & \text{otherwise .} \end{cases} \qquad (31)$$

For $r = 0$, we define the discrete correlation function as the continuous one defined in eq.(27).

We finally average over time, and obtain:

$$C(r_k) = \begin{cases} 1 & r_k = 0 \\ \dfrac{1}{T} \sum_{t=1}^{T} C(r_k, t) & r_k = k\Delta r, k = 1, \ldots, k_M \end{cases} \qquad (32)$$

We are essentially keeping the correlation for $r = 0$ separated from the other bins, since it has a special meaning. In the other bins we are instead averaging the correlation coefficient over intervals of size Δr. The choice of Δr is relevant here: if it is too small, the correlation function would be dominated by noise (as occurs in the continuous case); if it is too large, we would not be able to detect a correlation signal. Our choice is to define Δr as the typical nearest neighbor distance in the system.

Nearest neighbor distance, d$_{NN}$. We compute the distance of each mosquito from the other $N-1$, and we select the smallest. We do this, for all the mosquitoes in the swarm and for all the instants of times. We define the typical distance to the nearest neighbor, d_{NN}, as the median value of this set.

Correlation range, r$_0$. We compute the correlation range as the first point where the correlation function crosses the x-axis. To this aim, we find the smallest k such that $C(r_k) > 0$ and $C(r_{k+1}) < 0$, and we estimate r_0 with a linear interpolation between the two points $(r_k, C(r_k))$ and $(r_{k+1}, C(r_{k+1}))$.

Correlation length, ξ. Following [4] we computed the correlation length using the following integral definition:

$$\xi = \frac{\int_0^{r_0} r \cdot C(r) dr}{\int_0^{r_0} C(r) dr}. \qquad (33)$$

where r_0 is the correlation range. This definition has two main benefits: the first is that it uses not only the point where the correlation function crosses the x-axis, but the entire shape of the correlation function (up to r_0); the second, which is also the most relevant, is that this definition is general and it may be applied to correlation functions in different regimes. We compute the two integrals in eq.(33) with the trapezoid rule.

Swarm size, L. For consistency with [4, 11] we compute the size of the swarm L as the maximum distance between two mosquitoes. Specifically, at each instant in time we measure the distances between each pair of mosquitoes and we define the size at that frame, $L(t)$, as the maximum distance. We finally define L as the median value of the set of all these $L(t)$.

Standard deviation for ξ and L. We measure the standard deviation on the correlation length ξ and on the size of the swarm L, with a resampling method, see [41], already used in [18]. We randomly select half of the T different frames of the acquisition and compute, on this sub-sample, the correlation function, the correlation length and the size of the swarm. We repeat this procedure 10^5 times, obtaining a set of values for ξ and L. We extracted the standard deviations of the two sets, which are the estimates of the standard deviation for ξ and L shown in Fig.3d.

Evidence of male–male interaction. We designed and developed a shuffling procedure with which we were able to generate non–interacting

and realistic swarms, without the need to define a model. These artificial swarms present the same localization properties as real swarms, from which they differ only in the spatial arrangement of mosquitoes.

In eq.(9), we defined individual centroids as the average over time of the position of single mosquitoes. We can now rewrite the position at time t of a mosquito i, as a function of its centroid:

$$\vec{x}_i(t) = \vec{x}_i + \delta\vec{x}_i(t) \qquad (34)$$

where $\vec{x}_i$ is the trajectory centroid, hence not–dependent on time, and $\delta\vec{x}_i(t)$ represents the location of mosquito i at time t with respect to its own centroid.

We have then a set of centroids, $C = \{\vec{x}_i, i = 1, \dots, N\}$, each associated with a specific mosquito. In the shuffling procedure, we scramble the associations between centroids and mosquitoes. For instance, we associate the centroid of mosquito j to mosquito i, and we define a new trajectory, $\vec{x}_i^*$ such that:

$$\vec{x}_i^*(t) = \vec{x}_j + \delta\vec{x}_i(t). \qquad (35)$$

In doing so, the position of the new trajectory is shifted in space, in such a way that its new centroid is $\vec{x}_j$ instead of $\vec{x}_i$. But, its velocity and its speed in time are not affected by this shift, as well as its individual spatial distribution.

We repeat this procedure for all the mosquitoes in the swarm, associating each centroid with a different trajectory. In the example of Fig.4, the yellow trajectory is shifted on the turquoise centroid, the purple trajectory on the yellow centroid, and the turquoise trajectory on the yellow centroid.

At the end of the shuffling process, we have an artificial swarm where, by mixing up mosquito positions, we cut potential interaction links due to spatial proximity. But at the same time, we are not modifying individual spatial extent nor speed, which will both present the same localization properties as the original real swarm.

We also compute the probability of finding r_{0S} greater or equal to r_0. To this aim, we measured the ratio between the occurrences of the shuffled swarms with $r_{0S} \geq r_0$ and the total number of times we shuffled the swarm (10^6). This probability is essentially the p–value for r_0.

Acknowledgments.

We thank Andrea Cavagna and Irene Giardina for discussions and for the financial support, both in terms of resources and equipment. We thank CoBBS group, to which MF, FG, SM and LP belong, for the data on the midge swarm presented in SI. SM thanks Federica Ferretti, who unwittingly triggered part of this work with a question at the IntCha2025 conference. This work was supported by ERC Grant RG.BIO (Contract n. 785932), and MIUR Grant PRIN2022 funded by the European Union – Next Generation EU, under Italy's National Recovery and Resilience Plan (PNRR), Mission 4, Component 1 (CUP B53D23011780006).

References

[1] Sarfati, R., Hayes, J.C., Peleg, O.: Self-organization in natural swarms of *Photinus carolinus* synchronous fireflies. Science Advances **7**(28) (2021)

[2] Strogatz, S.H., Stewart, I.: Coupled oscillators and biological synchronization. Scientific american **269**(6), 102–109 (1993)

[3] Attanasi, A., Cavagna, A., Del Castello, L., Giardina, I., Grigera, T.S., Melillo, S., Parisi, L., Pohl, O., Shen, E., Viale, M.: Information transfer and behavioural inertia in starling flocks. Nature Physics **10** (2014)

[4] Cavagna, A., Culla, A., Feng, X., Giardina, I., Grigera, T.S., Kion-Crosby, W., Melillo, S., Pisegna, G., Postiglione, L., Villegas, P.: Marginal speed confinement resolves the conflict between correlation and control in collective behaviour. Nature Communications **13** (2022)

[5] Katz, Y., Tunstrøm, K., Ioannou, C.C., Huepe, C., Couzin, I.D.: Inferring the structure and dynamics of interactions in schooling fish. Proceedings of the National Academy of Sciences **108**(46), 18720–18725 (2011)

[6] Herbert-Read, J.E., Perna, A., Mann, R.P., Schaerf, T.M., Sumpter, D.J.T., Ward, A.J.W.: Inferring the rules of interaction of shoaling fish. Proceedings of the National

Academy of Sciences **108**(46), 18726–18731 (2011)

[7] Sumpter, D.J.T.: Collective Animal Behavior. Princeton University Press, Princeton (2011)

[8] Strogatz, S.: Sync: The Emerging Science of Spontaneous Order. Hyperion, New York (2003)

[9] Kelley, D., Ouellette, N.: Emergent dynamics of laboratory insect swarms. Scientific reports **3**, 1073 (2013)

[10] Gorbonos, D., Puckett, J., Vaart, K., Sinhuber, M., Ouellette, N., Gov, N.: Pair formation in insect swarms driven by adaptive long-range interactions. Journal of the Royal Society, Interface **17** (2020)

[11] Attanasi, A., Cavagna, A., Del Castello, L., Giardina, I., Melillo, S., Parisi, L., Pohl, O., Rossaro, B., Shen, E., Silvestri, E., Viale, M.: Collective behaviour without collective order in wild swarms of midges. PLoS Comput Biol **10(7)** (2014)

[12] Downes, J.A.: The swarming and mating flight of diptera. Annual Review of Entomology **14**, 271–298 (1969)

[13] Charlwood, J.D., Jones, M.D.R.: Mating behaviour in the mosquito, anopheles gambiae s.1.save. Physiological Entomology **4**(2), 111–120 (1979)

[14] Baeshen, R.: Swarming behavior in anopheles gambiae (sensu lato): Current knowledge and future outlook. Journal of Medical Entomology **59**(1), 56–66 (2021)

[15] Cribellier, A., Poda, B.S., Diabaté, A., Roux, O., Muijres, F.T.: The complex swarming dynamics of malaria mosquitoes emerges from simple minimally-interactive behavioral rules. bioRxiv (2024)

[16] Attanasi, A., Cavagna, A., Del Castello, L., Giardina, I., Melillo, S., Parisi, L., Pohl, O., Rossaro, B., Shen, E., Silvestri, E., Viale, M.: Finite-size scaling as a way to probe near-criticality in natural swarms. Phys. Rev. Lett. **113** (2014)

[17] Cavagna, A., Conti, D., Creato, C., Del Castello, L., Giardina, I., Grigera, T.S., Melillo, S., Parisi, L., Viale, M.: Dynamic scaling in natural swarms. Nature Physics **13** (2017)

[18] Cavagna, A., Di Carlo, L., Giardina, I., Grigera, T.S., Melillo, S., Parisi, L., Pisegna, G., Scandolo, M.: Natural swarms in 3.99 dimensions. Nature Physics **19** (2023)

[19] Manoukis, N., Diabate, A., Abdoulaye, A., Diallo, M., Dao, A., Yaro, A., Ribeiro, J., Lehmann, T.: Structure and dynamics of male swarms of anopheles gambiae. Journal of medical entomology **46**, 227–35 (2009)

[20] Cavagna, A., Giardina, I., Gucciardino, M.A., Iacomelli, G., Lombardi, M., Melillo, S., Monacchia, G., Parisi, L., Peirce, M.J., Spaccapelo, R.: Characterization of lab-based swarms of anopheles gambiae mosquitoes using 3d-video tracking. Scientific Reports **13** (2023)

[21] Poda, B.S., Cribellier, A., Feugère, L., Fatou, M., Nignan, C., Sales Hien, D.F., Müller, P., Gnankiné, O., Dabiré, R.K., Diabaté, A., Muijres, F.T., Roux, O.: Spatial and temporal characteristics of laboratory-induced anopheles coluzzii swarms: Shape, structure, and flight kinematics. iScience **27**(11) (2024)

[22] Gibson, G., Russell, I.: Flying in tune: Sexual recognition in mosquitoes. Current Biology **16**(13), 1311–1316 (2006)

[23] Somers, J., Georgiades, M., Su, M.P., Bagi, J., Andrés, M., Alampounti, A., Mills, G., Ntabaliba, W., Moore, S.J., Spaccapelo, R., Albert, J.T.: Hitting the right note at the right time: Circadian control of audibility in anopheles mosquito mating swarms is mediated by flight tones. Science Advances **8**(2) (2022)

[24] Poda, B., Nignan, C., Gnankine, O., Dabiré, R., Diabate, A., Roux, O.: Sex aggregation

and species segregation cues in swarming mosquitoes: Role of ground visual markers. Parasites & Vectors **12** (2019)

[25] Butail, S., Manoukis, N., Diallo, M., Ribeiro, J., Paley, D.: The dance of male *Anopheles gambiae* in wild mating swarms. In: Journal of Medical Entomology (2013)

[26] Shishika, D., Manoukis, N., Butail, S., Paley, D.: Male motion coordination in anopheline mating swarms. Scientific reports **4** (2014)

[27] Aldersley, A., Champneys, A., Homer, M., Bode, N.W.F., Robert, D.: Emergent acoustic order in arrays of mosquitoes. Current Biology **27**(22), 1208–1210 (2017)

[28] Attanasi, A., Cavagna, A., Del Castello, L., Giardina, I., Jelic, A., Melillo, S., Parisi, L., Pellacini, F., Shen, E., Silvestri, E., et al.: Greta - a novel global and recursive tracking algorithm in three dimensions. Pattern Analysis and Machine Intelligence, IEEE Transactions on **99** (2015)

[29] Shaebani, M.R., Wysocki, A., Winkler, R.G., Gompper, G., Rieger, H.: Computational models for active matter. Nature Reviews Physics **4** (2020)

[30] Clements, A.N.: The Biology of Mosquitoes: Volume 2: Sensory Reception and Behaviour. CABI Publishing, Wallingford (1999)

[31] Yuval, B.: Mating systems of blood-feeding flies. Annual Review of Entomology **51**(1), 413–440 (2006)

[32] Fawaz, E.Y., Allan, S.A., Bernier, U.R., Obenauer, P.J., Diclaro, J.W.: Swarming mechanisms in the yellow fever mosquito: aggregation pheromones are involved in the mating behavior of *aedes aegypti*. Journal of Vector Ecology **39**(2), 347–354 (2014)

[33] South, A., Catteruccia, F.: Sexual selection and the evolution of mating systems in mosquitoes. In: Advances in Insect Physiology vol. 51, pp. 67–92. Elsevier, London (2016)

[34] MC, G., D., R.: Nanometre-range acoustic sensitivity in male and female mosquitoes. Proc Biol Sci. **267** (2000)

[35] Gibson, G., Warren, B., Russell, I.: Humming in tune: Sex and species recognition by mosquitoes on the wing. Journal of the Association for Research in Otolaryngology **11** (2010)

[36] Cavagna, A., Giardina, I., Grigera, T.S.: The physics of flocking: Correlation as a compass from experiments to theory. Physics Reports **728**, 1–62 (2018). The physics of flocking: Correlation as a compass from experiments to theory

[37] Alampounti, A.C., Georgiades, M., Faber, J., Bagi, J., Andrés, M., Bozovic, D., Albert, J.T.: Distortion rules: audibility creation in the mosquito ear. bioRxiv (2025)

[38] Benedict, M.: Methods in Anopheles Research. MR4 (Malaria Research & Reference Reagent Resource Center), Atlanta, GA, USA (2010)

[39] Cavagna, A., Creato, C., Del Castello, L., Giardina, I., Melillo, S., Parisi, L., Viale, M.: Error control in the set-up of stereo camera systems for 3d animal tracking. The European Physical Journal Special Topics **224**(17-18), 3211–3232 (2015)

[40] Zhang, Z.: A flexible new technique for camera calibration. IEEE Transactions on Pattern Analysis and Machine Intelligence **22**(11), 1330–1334 (2000)

[41] Leuzzi, L., Marinari, E., Parisi, G.: Probability Theory for Quantitative Scientists. Cambridge University Press, Cambridge (2025)