A *De Novo* Algorithm for Allele Reconstruction from Oxford Nanopore Amplicon Reads, with Application to *CYP2D6*

Scott D. Brown[1], Lisa Dreolini[1], Agata Minor[2,3], Michelle Mozel[4], Nancy Wong[2], Sharon Mar[2], Amanda Lieu[2], Maimun Khan[2], Amanda Carlson[2], Monica Hrynchak[2,3], *Robert A. Holt[1,5,6], *Perseus I. Missirlis[4,7]

[1] Department of Basic and Translational Research; BC Cancer Research Institute; 675 W 10th Ave, Vancouver, BC, V5Z 1L3, Canada

[2] Department of Pathology and Laboratory Medicine; Molecular Cytogenetics Laboratory; Royal Columbian Hospital; 330 E Columbia St, New Westminster, BC, V3L 3W7, Canada

[3] Department of Pathology and Laboratory Medicine; University of British Columbia; G227-2211 Wesbrook Mall, Vancouver, BC, V6T 2B5, Canada

[4] Department of Anesthesiology and Perioperative Medicine; Royal Columbian Hospital; 330 E Columbia St, New Westminster, BC, V3L 0A2, Canada

[5] Department of Molecular Biology and Biochemistry; Simon Fraser University; SSB8166 – 8888 University Drive, Burnaby, BC, V5A 1S6, Canada

[6] Department of Medical Genetics; University of British Columbia; C201 – 4500 Oak Street, Vancouver, BC, V6H 3N1, Canada

[7] Department of Anesthesiology, Pharmacology & Therapeutics; University of British Columbia; 2775 Laurel Street, Vancouver, BC, V5Z 1M9, Canada

* Co-senior Authors - Robert A. Holt and Perseus I. Missirlis

Correspondence: sbrown@bccrc.ca, rholt@bccrc.ca or perseus.missirlis@fraserhealth.ca

Abstract

The Oxford Nanopore Technologies' sequencing platform offers a path towards bedside genomics, producing long reads that can completely cover a gene of interest, and thus detect any known or novel variant the gene contains. However, the analysis of these long reads to identify actionable genotypes remains challenging and typically requires customization depending on the target gene. Here, we describe a generic algorithm to accurately reconstruct allele sequences derived from long-reads of genomic-amplicon origin. Rather than calling variants directly from these long-reads, our method takes a "sequence-first" approach, performing an

unbiased reconstruction of the underlying amplicon sequences to generate high-confidence reconstructed allele sequences. This is

done without user input of the expected target gene, allowing for any source amplicon to be reconstructed. These high-confidence

reconstructed allele sequences are then compared to the genomic reference sequence of the gene to infer the specific diplotype

present in the sample. This approach is agnostic towards the number of genes and alleles present and readily detects novel variants.

We demonstrate our approach using three independent data sets for *CYP2D6*, a diverse and complex gene with over 175 known

alleles of clinical significance affecting drug dosing. We show how our approach can accurately recover validated *CYP2D6*

diplotypes from 20 Coriell samples sequenced using different primer sets, on different Oxford Nanopore Technologies flow cell

versions, and to different depths. This includes inferring occurrences of copy number variation from relative abundances of each

allele, a critical factor for ascribing functional effects to a diplotype. Further, we demonstrate our approach's utility for other

genomic regions, including *HLA*.

Keywords: MinION, pharmacogenomics, CYP2D6, long-read sequencing, allele reconstruction, haplotyping

---

Background

Gene diversity can significantly impact the therapeutic effect of drugs in patients. One such example is the cytochrome P450 (CYP)

superfamily of enzymes that play a key role in drug metabolism and can result in either supratherapeutic or subtherapeutic serum

drug concentrations (McDonnell and Dang, 2013). Genotyping of genes prior to treatment can inform drug selection, dosing

decisions, and avoidance of adverse side effects (Chang *et al.*, 2015). *CYP2D6* is one of the most clinically significant genes and

CYP2D6 genotype-guided dosing exists for a number of indications, including anti-fungal treatment in hematopoietic stem cell

transplantation, breast cancer chemotherapeutics, tricyclic antidepressants, and opioids for pain management (Teusink *et al.*, 2016;

Kiyotani *et al.*, 2008; Hicks *et al.*, 2017; Somogyi *et al.*, 2015). As such, a method for obtaining a fast and accurate readout of a

patient's *CYP2D6* genotype is of great importance. *CYP2D6* is highly polymorphic; currently there are over 175 known functional

alleles (each designated as a "star-allele", with *CYP2D6*1* being the reference), many of which differ from another allele by only

a single base substitution (Whirl-Carrillo *et al.*, 2021, 2012), and new alleles are constantly being added as they are discovered.

Traditional methods to interrogate genotypes are generally slow and expensive; typically relying on custom sets of PCR

reactions to interrogate specific sites along the length of each gene of interest. More contemporaneous approaches have leveraged

high-throughput sequencing data, either whole genome (Chen *et al.*, 2021; Deserranno *et al.*, 2023), or amplicon-based (Hari *et al.*,

2023; Liau, Cree, *et al.*, 2019; Liau, Maggo, *et al.*, 2019; Qiao *et al.*, 2016; Ammar *et al.*, 2015; Charnaud *et al.*, 2022). These

methods often make use of Pacific Biosciences or Oxford Nanopore Technologies (ONT) long reads, allowing the entire gene to

be sequenced in a single read, and thus, facilitating accurate DNA sequencing and haplotyping with a single assay. Regardless of

sequencing strategy, these methods take a "variant-first" analysis approach, relying on variant calling from the sequence reads first,

followed by phasing of the variants into haplotypes.

We sought to develop a method of diplotyping genes that uses a "sequence-first" rather than "variant-first" approach. We

aimed to perform allele sequence reconstruction from long read sequences by generating high-confidence consensus sequences of

each allele present in the original sample, then genotyping each resulting high-confidence consensus sequence to assign a haplotype.

Further, we ensured our method was unbiased and generalizable, not being restricted to a specific gene. As such, our only input

requirements are that the dataset is generated from genomic amplicon sequencing using ONT long reads and the dataset contains

reads which completely cover the genomic regions of interest. No information is needed *a priori* regarding the specific genomic

target, nor the number of targets. We demonstrate the utility of our approach using ONT sequencing data from *CYP2D6* amplicons

and scripts that automate *CYP2D6* haplotyping of reconstructed allele sequences to call the most likely star-allele and identify any

discrepant, potentially novel, variants.

Results

Selection of **CYP2D6** *amplicon.*

The *CYP2D6* gene locus is 4.3 kb long and is located on the negative strand of chromosome 22 (chr22:42,126,499-42,130,810;

hg38). Variants affecting CYP2D6 enzyme function may fall anywhere within this region, with the range of currently known

variants (insertions, deletions, and single nucleotide variants) spanning chr22:42,126,578-42,130,783 (Whirl-Carrillo *et al.*, 2021,

2012). We used primers previously developed by Ammar et al. (Ammar *et al.*, 2015) to selectively amplify a 5.1 kb region of

CYP2D6 (chr22:42,126,072-42,131,142), and not the nearly identical *CYP2D7* pseudogene (**Figure 1**).

Figure 1. Graphical summary of the genomic region encompassing *CYP2D6*. The *CYP2D6* gene is shown, with larger boxes for the exons. The
CYP2D7 pseudogene is shown upstream of the *CYP2D6* transcriptional start site. All known functional variants used by ClinPGx (Whirl-Carrillo
et al., 2021, 2012) to define specific *CYP2D6* alleles are shown as ticks. The three amplicon regions assessed in our work are shown as individual
tracks.
83

Oxford Nanopore Technologies sequence generation.

Genomic DNA from three Coriell samples with known *CYP2D6* haplotypes (NA02016; *2xN/*17*, NA10005; *17/*29*, NA17300;

*1/*6*) were used. Sequencing libraries were constructed using the ONT Ligation Sequencing Kit V14. We loaded each amplicon

library onto a separate R10.4.1 Flongle flow cell, each being run for 24 h on a MinION Mk1b USB sequencer.

Base calling of the raw .fast5 files was performed using the GPU-enabled version of Guppy (a base caller provided by

ONT). No quality filter was used at this stage, as quality filtering was performed at the per-base level later in our analysis pipeline.

Omitting the quality filter allows for high-quality bases in lower-quality reads to still contribute information, maximizing the depth

of coverage at all positions. Porechop was used to remove any adapter sequences and split chimeric reads. We obtained 87,906,

56,948, and 50,389 total reads from NA17300, NA10005, and NA02016 respectively. The average read length was 1,736 bases (±

2,078.35) (**Figure 2**). We observed a median 8.5 % error rate in these reads (range 7.0 – 9.0 %).

Figure 2. Distribution of read lengths for all samples. Samples ordered by decreasing numbers of total reads. In all cases, a population of reads
at the expected 5.1 – 6.6 kb range is seen. In-house generated samples (blue) show higher than expected numbers of short reads, suggestive of
degraded DNA amplicons during library preparation. The x-axis was truncated at 10,000 bp, removing an average of 0.37 % of reads per sample
from display.
100

Inclusion of existing **CYP2D6** *sequence data sets to test different flow cells, sequence qualities, and amplicons.*

To validate our approach on different ONT flow cell versions, *CYP2D6* amplicons, sequencing depths, and haplotypes, we obtained

data from Liau et al. (Liau, Maggo, *et al.*, 2019) for seven additional Coriell samples. Liau et al. generated a 6.6 kb amplicon for

CYP2D6 (**Figure 1**), which covered more of the upstream and downstream regions of *CYP2D6,* and sequenced these on older

MinION R9.4 flow cells, yielding an average of 2,292 reads per sample, and having a median error rate of 12.1 % (range 11.9 –

12.5 %). This larger 6.6 kb amplicon should have a greater ability to detect gene duplication events, being more likely to span the

junction site of a duplicated copy, though the exact location of this junction is unknown. Of note, Liau et al. also sequenced an

even larger 8.6 kb amplicon for samples with confirmed duplications, covering the repetitive region downstream of *CYP2D6*

(Gaedigk *et al.*, 2019), but this sequence data was unavailable.

Data for twelve additional Coriell samples were obtained from Wongsurawat et al. (Wongsurawat *et al.*, 2025), with an

average of 55,068 reads per sample, and a median error rate of 5.1 % (range 4.7 – 5.7 %). Wongsurawat et al. generated a 5.1 kb

amplicon using primers from Puaprasert et al. (Puaprasert *et al.*, 2018), covering a region similar to Ammar et al., and were

multiplexed on MinION R9.4.1 flow cells. Wongsurawat et al. included two additional primer sets, one designed to identify gene

deletions by producing a 3.5 kb amplicon only upon gene deletion (otherwise priming at sites 15 kb apart; too large to be efficiently

amplified), and another designed to amplify a 3.5 kb region between duplicated copies of *CYP2D6* (with priming sites on either

side of *CYP2D6* facing "outwards"). This data was generated using two different library construction kits; the Rapid Barcoding Kit

and the Ligation Sequencing Kit. We only used Ligation Sequencing Kit data for our analysis. Combined with our in-house

generated data, we have compiled 22 ONT sequence datasets covering 20 distinct Coriell samples.

Assigning reads to inferred amplicons based on whole genome alignment.

We designed our method to perform an unbiased reconstruction of allele sequences. We base this on an assumption that every

amplicon present in the sequence library will have a subset of reads supporting that amplicon. As such, rather than supplying the

algorithm with the anticipated amplicon, we used the sequence reads to infer the genomic loci that were amplified. To achieve this,

we aligned all reads to the human reference genome (hg38) using minimap2, allowing for only the single best mapping position of

each read. Supplementary alignments, or split reads, were included in order to detect large deletions or structural variants. The

depth of coverage across the whole genome was assessed, and the regions with high coverage were taken to be representative of

amplicons present in the source sequencing library (**Figure 3A**). Importantly, this approach does not assume any predefined number

of amplicons, allowing this approach to work, theoretically, for multiplexed samples of any number of amplicons. All of our

samples yielded an inferred amplicon region at the expected *CYP2D6* locus, and six samples yielded at least one second, distinct

inferred amplicon region (**Supplementary Table S1**). Two of these non-*CYP2D6* inferred amplicons (NA02016_ligate,

NA07439_ligate) were from the region between *CYP2D6* and *CYP2D7*, corresponding to the duplication-detecting amplicon from

Wongsurawat et al. One (NA17203_ligate) was from the *CYP2D7* locus, possibly derived from a *CYP2D6-CYP2D7* hybrid allele,

where the majority of the amplicon sequence was derived from *CYP2D7*. Three (NA02016, NA10005, NA17052_ligate) were from

elsewhere in the genome. NA02016 and NA10005 produced short (mean 609 bp) off-target amplicon regions from chromosomes

4, 9, 12, and 20. NA17052_ligate had an off-target amplicon fromfrom the *SORCS2* locus on chromosome 4. Careful inspection of

the identified *SORCS2* locus sequence shows it contains a string of 15 contiguous bases at the start (GCCCTGGGAGGTAGG),

and 13 contiguous bases at the end (GAAGCCTGCAAAG), which exactly match sequence within the *CYP2D6* locus, and are

present in the primers used by Wongsurawat et al. All inferred amplicons are carried forward in the analysis as possible loci of

interest.

Figure 3. Schematic overview of algorithm. (A) Sequence reads are aligned to the genome, and regions of high depth are extracted and inferred
as the source amplicon (blue). (B) These reads are clustered by length using the DBSCAN algorithm, and the most abundant cluster(s) are selected
for allele reconstruction (solid outlined box). (C) Selected reads are aligned to the inferred amplicon reference sequence (extracted from the genomic
reference; blue), and each position of the alignment is assessed sequentially, building a consensus sequence base by base (top underlined sequence).
Variants as compared to the reference are coloured orange. If a position is encountered as appearing heterozygous (having roughly equal support
for each of two or more bases), resulting in an unknown consensus base, the reads are partitioned into subsets based on the base they support at that
position (D). The reads in each subset are used, independently, to re-attempt allele reconstruction. In the example shown, this yields two consensus
sequences (top underlined sequence), each with homozygous support for each base. Small indels are treated the same way as bases, with partitioning
occurring similarly for reads supporting and not supporting the indel.

Reads clustered into groups based on length.

In our in-house generated data, we observed a higher than expected frequency of short reads (**Figure 2).** On average, 60.8 % of

reads were under 1 kb, compared to 0 % for data from Liau et al. (only aligned reads were provided) and 0.08 % from Wongsurawat

et al. We aligned our short reads to the *CYP2D6* reference sequence, demonstrating that they mapped to random positions spread

across the amplicon (**Supplementary Figure S1**). This suggested the short reads were derived from fragmented amplicons during

library construction, and were not representative of *bona fide* short amplicons within the sample. Liau et al. describe seeing many

short reads in their first run, which prompted them to use modified beads to remove any DNA fragments shorter than 3-4 kb (Liau,

Maggo, *et al.*, 2019), which could be an approach we take in the future. We assert that full-length (non-fragmented) reads which

were derived from the same amplicon would all have the same length (or highly similar lengths when allowing for sequence errors).

To select the subset of reads for each inferred amplicon most likely to be informative, we performed length-clustering on the reads

from each inferred amplicon using the DBSCAN algorithm (Schubert *et al.*, 2017) (**Figure 3B**). We performed a parameter space

exploration of the two DBSCAN parameters, EPS and Minimum Neighbours (see Materials and Methods), which revealed an

inverse relationship between the variability of read lengths and size of the resulting clusters. Low EPS and high minimum

neighbours result in a more strictly defined cluster, ensuring all reads have a similar length, at the sacrifice of number of reads,

whereas high EPS and low minimum neighbours allow larger clusters with more variable lengths. Since the purpose of length-

clustering here is to identify subsets of reads from the same amplicon, we chose the stricter parameter values to avoid using shorter,

degraded reads. After clustering, we first selected the cluster that had reads with a mean length that was closest to the source

inferred amplicon, regardless of abundance, representing reads that fully covered the amplicon. Secondly, we also assessed any

highly abundant clusters, based on the premise that there may be other smaller or larger amplicons that would otherwise go

undetected (**Supplementary Table S1**). All selected clusters were used for subsequent allele sequence reconstruction.

Allele sequence reconstruction performed through unbiased, recursive consensus sequence generation.

At this stage, we had obtained subsets of length-matched reads which were likely to have been derived from the same locus. These

subsets, however, contained a mix of reads supporting each of the underlying haplotypes present in the original sample. If all

sequence reads were error-free, this would have been a trivial exercise to partition reads based on their exact sequence. Due to

sequencing errors, reads derived from the same haplotype did not share the exact same sequence. In order to reconstruct the

underlying allele sequences, we took an empirical approach for determining which allele sequences the reads supported. This was

achieved by finding partitions of the reads such that all reads in a partition supported the same sequence (while allowing for some

noise due to sequence errors). First, we attempted to build a consensus sequence from all of the reads in the set, using similar logic

to our previous work (Brown *et al.*, 2023). Briefly, the reads were aligned to the inferred amplicon reference sequence, and the

pileup of base support at each position of the alignment was assessed, sequentially, to call a consensus base at that position. If a

position was reached that did not have sufficient evidence of a homozygous base (homozygosity defined as the top two bases

having a ratio of at least 2:1; see Materials and Methods) (**Figure 3C**), the reads were partitioned into subsets defined by the

observed bases at this heterozygous position (**Figure 3D**). The consensus sequence generation was restarted from the beginning,

recursively, on each of these resulting subsets. This process continued until a homozygous consensus sequence was successfully

generated from a subset of reads, or the size of a subset fell below the lower limit threshold for reporting (5% of the total starting

read set of length-clustered reads, selected empirically).

Our approach considered base quality in two ways. In order for a base to contribute to the consensus sequence, the base

must have had a minimum quality of 10 (selected empirically). If this criterion was not met, a secondary criterion was tested,

requiring a quality score above the 33[rd] percentile of observed qualities at that position (selected empirically). This ensured that

positions with lower-quality bases across all reads were not completely lost from the consensus due to lack of coverage, while still

ensuring that the consensus sequence was built using the best quality bases available at each position. Strand-specific errors are

also common in ONT data (Krishnakumar *et al.*, 2018; Branton *et al.*, 2008), resulting in a specific sequencing error occurring

primarily on one strand. When assessing all reads without considering the strand, such errors could appear as a heterozygous

position. To avoid erroneously calling a position heterozygous, we assess, separately, the positive and negative strand reads at each

position, requiring support of heterozygosity from both strands independently in order to classify a position as truly heterozygous.

All reconstructed consensus sequences are available in **Supplementary File S1**.

Focusing on the results for read sets derived from the correct genomic locus (*CYP2D6*) and of the correct mean length

(5.1 - 6.6 kb) for the 22 samples, all yielded at least two distinct consensus sequences (range 2-27). While having two sequences is

expected for a diploid genome, having greater than two consensus sequences could support gene duplication events where the

copies have sequence differences. This could also, however, be from a common sequencing error that is abundant enough to result

in a consensus sequence. For samples with a large number of consensus sequences, the distribution of read depths was highly

skewed, typically with only 2 consensus sequences having the majority of the reads. This is consistent with these lower-abundance

consensus sequences arising from common sequencing errors and not *bona fide* amplicons. Relative read depths may help

distinguish these two cases, as described later.

All three samples sequenced by our group had length-clusters that were shorter than the expected amplicon size, but were

abundant enough to have allele sequence reconstruction attempted (**Supplementary Table S1**). This is likely an artefact from our

high rate of short reads. One yielded two consensus sequences mapping to exon 9 and downstream of *CYP2D6* (NA02016), one

yielded two consensus sequences mapping to exons 8-9 of *CYP2D6* (NA10005), and one yielded a consensus sequence mapping

to exons 4-6 of *CYP2D6* (NA17300). This supports the notion that these short reads were derived from degraded DNA template

occurring during library construction. Indeed, when inspecting these short consensus sequences, they match the respective regions

of the long consensus sequences from the same sample (**Supplementary Figure S2**). Two samples had allele sequence

reconstruction attempted for reads that did not originate from the *CYP2D6* locus or supplementary amplicons. As described above,

these are most likely derived from off-target amplification from the genome during PCR, but could also be the result of *CYP2D6-*

CYP2D7 hybrid alleles (in the case of the *CYP2D7*-derived consensus sequence in NA17203_ligate). The *SORCS2*-derived

consensus sequences were run through BLASTn (Sayers *et al.*, 2025), and both mapped back to the *SORCS2* locus, supporting that

these reads were from off-target amplification.

Confirmation of deletion detection through simulated data.

To confirm our allele sequence reconstruction would generate a consensus sequence supporting a large deletion relative to the

genomic reference, should one exist, we simulated long reads representing our amplicon sequence with the entire *CYP2D6* sequence

deleted (see Materials and Methods). Encouragingly, our pipeline output a consensus sequence that perfectly matched the source

template sequence used to simulate the reads, despite the reads containing a 4.3 kb deletion relative to the human reference genome.

This demonstrates that our algorithm will accurately produce a consensus sequence representative of the sequence the reads support.

Automated reporting of the most likely* CYP2D6 *star alleles and putative novel variants.

To identify the specific *CYP2D6* alleles that each resulting consensus sequence supported, we first aligned each consensus sequence

to the *CYP2D6*1* reference sequence. All variants (indels and mismatches) were enumerated, and those occurring within coding

regions or splice sites were used for *CYP2D6* allele determination. We automated this process by using the *CYP2D6* Allele

Definition Table from ClinPGx (Whirl-Carrillo *et al.*, 2021, 2012). Briefly, for each known star allele, all canonical variants were

parsed from the ClinPGx definition table. For alleles with variants that have multiple possible bases allowed, some of which may

include the reference wildtype base, all possible allowed bases are stored. Then, the set of identified variants for the consensus

sequence was compared to the set of all required or possible variants from all known alleles, and each known star allele was given

a score by summing the number of disagreements in the identified variants from the consensus sequence. Thus, a consensus

sequence that had exactly the same variants as a defined star-allele would receive a score of zero for that allele. Silent mutations

found in the consensus would negatively impact the score across all alleles equally. The allele with the lowest score was returned

as the best candidate allele assignment for the consensus sequence. In the event of an imperfect match, missing and/or extra variants

in the consensus sequence compared to the candidate star allele reference were also returned, allowing for manual review of the

candidate allele assignment (**Table 1**). In this way, novel variants and alleles can also be readily identified, rather than simply the

best match to known alleles.

Inferring copy number from reconstructed alleles.

Copy number of genes can be an important factor for determining functional effects. In our results, the relative copy number of the

reconstructed alleles could be inferred based on the number of reads that contributed to each allele. We pooled the read counts for

all sequences of the same allele. The presence of multiple distinct consensus sequences for the same allele could be used as evidence

of duplication events; however, the sequences should be manually inspected to determine if the differences are believed to be from

true underlying sequence differences or from regions of poor sequence quality such as those near the ends of the amplicon.

Importantly, the lack of multiple distinct sequences does not preclude the presence of a duplication, as each duplicated copy may

indeed have an identical underlying sequence. For these reasons, we propose the read depths as a more reliable estimate of relative

copy number. To interpret the allele read depth ratios, we tested for ratios > 2:1. In our results, the exact number of duplication

events could not be reliably determined, likely due to biases in different allele sequences and sequencing errors, especially in

samples with lower read counts. Therefore, the alleles with a greater than 2:1 ratio were assigned an *xN* duplication.

Known diplotypes are recapitulated from Coriell samples.

Across the 22 distinct samples analyzed, our algorithm successfully reconstructed *CYP2D6* diplotypes that matched the reported

GeT-RM genotype in 15 cases (68.2 %; **Table 1**) (Pratt *et al.*, 2010, 2016; Gaedigk *et al.*, 2019). Three cases (13.6 %) did not

match GeT-RM, but were consistent with other reports from the literature (described below), suggesting they are also correct.

Two cases (9.1 %) only differed by GeT-RM due to copy number differences, and the final two cases (9.1 %) missed the hybrid

*68 allele which was not detectable due to limitations with the amplicon on those datasets. Overall, this results in an effective

90.9 % accuracy rate when taking into account the current literature and limitations of the amplicons tested. This high level of

concordance included 13 samples (59.1 %) containing either synonymous substitutions (C6015G, C6057T) or known

homopolymer-induced false-positive deletions (7598delC, 7599delC; **Figure 4A**), which are disregarded during interpretation.

This also included seven samples (31.8 %) where allele copy number carried inherent uncertainty in our calls or the GeT-RM

reference (NA02016, NA02016_ligate, NA07439, NA07439_ligate, NA18518, NA23297, NA23348). These cases with copy

number variation included the two Coriell samples from Wongsurawat et al., where the consensus sequence we generated from

their duplication-detecting amplicon supports the presence of a duplicated allele (**Supplementary Figure S3**). In one sample

(NA17300) our reconstruction identified an additional uncharacterized variant (G6996A) in a *6* allele **(Figure 4B)**. This variant

occurs downstream of the premature stop codon caused by the 6727delT frameshift mutation, and thus would not have further

functional effect. Four uncharacterized deletion variants were detected; 6033delG (NA12878), 5903delT (NA23297, NA23405),

6667delA (NA23405), and 6751delC (NA23405). These all preferentially occurred in consensus sequences with lower numbers

of reads, suggesting they may be artefacts due to insufficient coverage rather than true deletions. Inspecting the base support

around these regions supports this notion (**Figure 4C-F**), with low coverage in one or both strands.

 Table 1. Summary of *CYP2D6* allele reconstruction and diplotyping.

Sample ID	Flow cell	Total Reads	Reads aligning to *CYP2D6*	Star Allele Candidate	Identified Exon and Splice Variants*	Allele Depth	Copy Number Ratio	Genotype Call	GeT-RM Genotype
NA02016	Flongle R10.4.1	50,389	32,844	*2	G6681C, C7870T, G9200C	7,575	> 2:1	*2xN/*17	*2xN/*17 (Pratt *et al.*, 2010)
				*17	C6041T, G6681C, C7870T, G9200C	2,064			
NA02016_ligate	MinION R9.4.1	66,015	38,232	*2	G6681C, C7870T, G9200C	11,139	> 2:1	*2xN/*17	*2xN/*17 (Pratt *et al.*, 2010)
				*17	C6041T, G6681C, C7870T, G9200C	2,717			
				Duplication	3.5 kb duplication-detecting amplicon	20,356			
NA07439	MinION R9.4	1,467	1,413	*4	C5119T, C6057T‡, G6681C, 7599delC†, G9200C	574	> 2:1	*4xN/*41	*4x2/*41 (Gaedigk *et al.*, 2019)
				*41	G6681C, 7598delC†, 7599delC†, C7870T, G8008A, 8305delC†, G9200C	111			
NA07439_ligate	MinION R9.4.1	102,224	51,543	*4	C5119T, C6057T‡, G6681C, G6866A, G9200C	17,641	> 2:1	*4xN/*41	*4x2/*41 (Gaedigk *et al.*, 2019)
				*41	G6681C, C7870T, G8008A, G9200C	4,329			
				Duplication	3.5 kb duplication-detecting amplicon	36,853			
NA10005	Flongle R10.4.1	56,948	39,313	*29	G6679A, G6681C, C7870T, G8203A, G9200C	291	~ 1:1	*17/*29	*17/*29 (Pratt *et al.*, 2010)
				*17	C6041T, G6681C, C7870T, G9200C	215			
NA12244_ligate	MinION R9.4.1	49,461	41,956	*35	G5050A, G6681C, C7870T, G9200C	12,495	~ 1:1	*35/*41	*35/*41 (Pratt *et al.*, 2010)
				*41	G6681C, C7870T, G8008A, G9200C	8,956			
NA12813_ligate	MinION R9.4.1	47,070	42,013	*2	G6681C, C7870T, G9200C	6,006	~ 1:1	*2/*4	*2/*4 (Pratt *et al.*, 2016; Gaedigk *et al.*, 2019)
				*4	C5119T, C5992A, A6002G, C6015G‡, G6681C, G6866A, G9200C	5,800			
NA12878	MinION R9.4	486	486	*3	7569delA, 7598delC†, 7599delC†	153	~ 1:1	*3/*4	*3/(*68)+*4 (Gaedigk *et al.*, 2019)
				*4	C5119T, C5992A, A6002G, C6015G‡, **6033delG**, G6681C, G6866A, 7599delC†, G9200C	104			
NA17039_ligate	MinION R9.4.1	38,634	34,533	*2	G6681C, C7870T, G9200C	6,515	~ 1:1	*2/*17	*2/*17 (Pratt *et al.*, 2010)
				*17	C6041T, G6681C, C7870T, G9200C	6,312			
NA17052_ligate	MinION R9.4.1	70,348	40,480	*1	-	23,200	-	*1/*1	*1/*1 (Pratt *et al.*, 2010)
NA17058_ligate	MinION R9.4.1	66,012	60,635	*36	C5119T, C6057T‡, G6681C, G9144C, C9148G, A9151G, T9153C, C9175T, A9176C, G9178C, T9184G, T9186C, G9187A, C9188G, T9189C, C9192T, G9200C	16,107	~ 1:1	*10/**36**	*10/*10 (Pratt *et al.*, 2010)
				*10	C5119T, C6057T‡, G6681C, G9200C	15,263			
NA17119_ligate	MinION R9.4.1	55,696	48,829	*1	-	6,115	~ 1:1	*1/*2	*1/*2 (Pratt *et al.*, 2010)
				*2	G6681C, C7870T, G9200C	5,784			
NA17203_ligate	MinION R9.4.1	45,220	32,038	*4	C5119T, C5992A, A6002G, C6015G‡, G6681C, G6866A, G9200C	4,061	~ 1:1:1	*4/*35/**68**	*4/*35 (Pratt *et al.*, 2010)
				*35	G5050A, G6681C, C7870T, G9200C	3,328			
				*68	CYP2D6-CYP2D7 hybrid	2,440			

Brown et al. – preprint copy

Sample	Platform			Allele	Variants	Count	Ratio	Call	Reference
NA17240_ligate	MinION R9.4.1	44,930	37,732	*1	-	4,376	~ 1:1	*1/*10	*1/*10 (Pratt *et al.*, 2010)
				*10	C5119T, C6057T‡, G6681C, G9200C	3,897			
NA17280_ligate	MinION R9.4.1	51,996	44,679	*3	7569delA	5,724	~ 1:1	*3/**59**	*2/*3 (Pratt *et al.*, 2010)
				*59	G6681C, C7870T, G7959A, G9200C	4,401			
NA17300	Flongle R10.4.1	87,906	60,470	*1	-	4,308	~ 1:1	*1/*6	*1/*6 (Pratt *et al.*, 2010)
				*6	6727delT, **G6996A**	4,132			
NA18518	MinION R9.4	620	620	*17	C6041T, G6681C, 7599delC†, C7870T, G9200C	148	> 2:1	*17**xN**/*29	*17/*29 (Pratt *et al.*, 2016; Gaedigk *et al.*, 2019)
				*29	G6679A, G6681C, 7598delC†, 7599delC†, C7870T, G8203A, G9200C	68			
NA21781	MinION R9.4	3,018	3,018	*2	G6681C, 7598delC†, 7599delC†, C7870T, G9200C	619	~ 1:1	*2/*4	*2**x2**/(***68**)+*4 (Gaedigk *et al.*, 2019)
				*4	C5119T, C5992A, A6002G, C6015G‡, G6681C, G6866A, 7599delC†, G9200C	337			
NA23093_ligate	MinION R9.4.1	23,207	19,815	*10	C5119T, C6057T‡, G6681C, G9200C	3,485	~ 3:2:1**	*1/*10/*36	*1/*36+*10 (Gaedigk *et al.*, 2019)
				*36	C5119T, C6057T‡, G6681C, G9144C, C9148G, A9151G, T9153C, C9175T, A9176C, G9178C, T9184G, T9186C, G9187A, C9188G, T9189C, C9192T, G9200C	2,078			
				*1	-	1,020			
NA23297	MinION R9.4	6,833	6,833	*10	C5119T, C6057T‡, G6681C, 7599delC†, G9200C	2,644	> 2:1	*10x2/*17	*10x2/*17 (Gaedigk *et al.*, 2019)
				*17	**5903delT**, C6041T, G6681C 7599delC†, C7870T, G9200C	762			
NA23348	MinION R9.4	3,384	3,384	*7	7598delC†, 7599delC†, A7955C	1,203	> 2:1	*7**xN**/*35	*7/*35 (Pratt *et al.*, 2016; Gaedigk *et al.*, 2019)
				*35	G5050A, G6681C, 7599delC†, C7870T, G9200C	367			
NA23405	MinION R9.4	235	233	*1	**5903delT**, 7599delC†	78	~ 1:1	*1/*7	*1/*7 (Pratt *et al.*, 2016; Gaedigk *et al.*, 2019)
				*7	**6667delA**, **6751delC**, 7599delC†, A7955C	41			

269 Bolded and underlined text denotes discrepancy between expected variants or alleles and reported results. †Deletion occurring in a homopolymer (5+) base stretch, a known common
270 sequencing error, which is ignored. ‡Synonymous variant. *Variants identified compared to *CYP2D6*1*, from the most abundant consensus sequence supporting each allele. **Copy
271 number uncertain due to greater than 3 alleles.

Figure 4. Example signal-to-noise plots for different identified variants. In each panel, the top-observed base at each position is displayed as a solid point, and all other detected bases are hollow points. The size of the point denotes the average quality of the base from all reads used to generate the consensus. The top graph shows results from forward reads, and the bottom graph from reverse reads. (A) An example of the apparent decrease in coverage in homopolymer regions, shown here with a deletion 7599delC being called in NA23348. (B) The novel G6996A variant from NA17300 CYP2D6*6 allele is shown, with no evidence of this being an erroneous call due to poor signal-to-noise or poor quality. This variant occurs downstream of the premature stop codon due to the 6727delT mutation in CYP2D6*6, and thus will not have any effect on the resulting protein. (C-F) Four uncharacterized, non-homopolymer deletions from NA12878, NA23297, and NA23405, all due to low coverage from one or both strands, and are likely technical artefacts rather than true deletions in the sequence.

Of the five samples classified as inconsistent with GeT-RM diplotypes, two (NA17058_ligate, NA17280_ligate) yielded

reconstructions strongly suggestive of an error in the reported GeT-RM calls, which were originally based on targeted genotyping

panels rather than full-length sequencing, and are known to be imperfect (van der Maas *et al.*, 2025) (**Figure 5**). These inconsistent

calls are also supported by recent reports (Charnaud *et al.*, 2022; Wongsurawat *et al.*, 2025). One case (NA17203_ligate) yielded

a consensus from *CYP2D7*-derived reads. Recently, NA17203 has been reported as harboring a *CYP2D6-CYP2D7* hybrid allele

(**68*) (Wongsurawat *et al.*, 2025; Charnaud *et al.*, 2022), though this has not yet been reported by GeT-RM. Manually assessing

the alignment of the NA17203_ligate *CYP2D7*-derived consensus sequence to the *CYP2D6* and *CYP2D7* reference sequences

supports that this sequence may be representative of the **68 CYP2D6-CYP2D7* hybrid allele, with a higher number of variants

detected from exon 2 – 9 of *CYP2D6* compared to *CYP2D7* (**Supplementary Figure S4**). Indeed, comparing our consensus

sequence to the **68* sequence obtained by Rubben et al. (Rubben *et al.*, 2022) from NA12878, we see only a single indel difference,

occurring between exons 3 and 4 (**Supplementary Figure S5**), and thus our consensus sequence supports the same functional

hybrid **68* allele. The remaining two cases (NA12878 and NA21781) involved the **68 CYP2D6-CYP2D7* hybrid allele, which

was not detected by the Liau et al. primer design. While no hybrid allele reads were captured in this dataset, the underlying

constituent haplotypes were reconstructed correctly, with only the **68* classification absent. Taken together, these results

demonstrate that the algorithm is robust and largely concordant with reference diplotypes, with nearly all discrepancies attributable

to synonymous changes, sequencing artefacts, or structure variants beyond the scope of the amplicon design.

Comparison to existing CYP2D6 *genotyping tools.*

To put our method in context with existing *CYP2D6* genotyping tools, we ran all of the samples through Aldy4 (Hari *et al.*, 2023),

SpecImmune (S. Wang *et al.*, 2025), and CoLoRGen (Rubben *et al.*, 2022), three "variant-first" approaches. SpecImmune and

CoLoRGen are compatible with ONT amplicon data. Aldy4 (Hari *et al.*, 2023) does not officially support ONT amplicon data,

however can be run without phasing or copy number detection to allow genotyping. We did not use any PacBio-specific approaches,

as in our experience, ONT and PacBio data and algorithms are generally not cross-compatible due to different sequencing error

profiles. For completeness, we also compared our results to the reported genotypes from Liau et al. and Wongsurawat et al. source

dataset publications (**Table 2**).

Ignoring copy number inference, Aldy4 generated *CYP2D6* genotypes that matched the GeT-RM genotype in 18/22

(81.8 %) cases (**Table 2**), though we note that two of these are likely incorrect based on more current genotypes available in the

literature. Of the remaining four cases, two involved missing a **68* allele, and one matched the alternate genotype we and the source

publication obtained, further suggesting an error in the GeT-RM genotype. This left one sample with an incorrect genotype call by

Aldy4. We obtained *CYP2D6* genotypes for all 22 samples using SpecImmune, though only 5 (22.7 %) matched the reported GeT-

RM genotype. Seventeen (77.3 %) had at least one of the two alleles correct. For CoLoRGen, the analysis only completed for 13

319   samples. Three (13.6 %) matched the GeT-RM genotype, while 8 (36.4 %) had at least one matching allele. Overall, our pipeline

320   demonstrated a greater effective accuracy (90.9 % when considering limitations of the amplicons and current literature; green and

321   blue shading) compared to SpecImmune (31.8 % by the same metric), CoLoRGen (13.6 %), and Aldy4 (86.4 %), while also being

322   able to infer some copy number variations that Aldy4 was unable to do with this data.

323   **Table 2. Results from existing *CYP2D6* genotyping tools.**

Sample ID	GeT-RM	Brown *et al.*	Liau *et al.*	Wongsurawat *et al.*	Aldy4	SpecImmune	CoLoRGen
NA02016	*2xN/*17	*2xN/*17			*2/*17	*83+*1/*2	-
NA02016_ligate	*2xN/*17	*2xN/*17		*2x2/*17	*2/*17	*1/*2	-
NA07439	*4x2/*41	*4xN/*41	*4xN/*41		*4/*41	*4/*10	*4(x3?)/*41
NA07439_ligate	*4x2/*41	*4xN/*41		*4x2/*69+*41	*4/*41	*4/*41	-
NA10005	*17/*29	*17/*29			*17/*29	*29/*29	-
NA12244_ligate	*35/*41	*35/*41		*35/*41	*35/*41	*1/*68+*2	*83/*83
NA12813_ligate	*2/*4	*2/*4		*2+*65/*4	*2/*4	*1/*39	*83/*4
NA12878	*3/(*68)+*4	*3/*4	*3/*4		*3/*4	*3/*4	*3/*4
NA17039_ligate	*2/*17	*2/*17		*2/*17	*2/*17	*1/*68+*2	*83/*83
NA17052_ligate	*1/*1	*1/*1		*1/*1	*1/*1	*1/*1	-
NA17058_ligate	*10/*10	*10/*36		*10/*10+*36	*10/*10	*1/*36+*10	-
NA17119_ligate	*1/*2	*1/*2		*1/*2	*1/*2	*1/*1	*1/?
NA17203_ligate	*4/*35	*4/*35/*68		*4+*36/*35+*65	*4/*35	*1/*39	*83/*4
NA17240_ligate	*1/*10	*1/*10		*1/*10	*1/*10	*1/*1	?/?
NA17280_ligate	*2/*3	*3/*59		*3/*59	*3/*59	*1/*1	*83/*59
NA17300	*1/*6	*1/*6			*1/*6	*1/*1	-
NA18518	*17/*29	*17xN/*29	*17/*29		*17/*29	*17/*2	*17/*29
NA21781	*2x2/(*68)+*4	*2/*4	*2xN/*4		*2/*4	*2/*4	*4/*73(?)
NA23093_ligate	*1/*36+*10	*1/*10/*36		*1/*10+*36	*1/*10	*1/*36+*10	*83
NA23297	*10x2/*17	*10xN/*17	*17/*10xN		*10/*17	*39/*10	*17/?
NA23348	*7/*35	*7xN/*35	*7/*35		*7/*35	*7/*35	-
NA23405	*1/*7	*1/*7	*1/*7		*1/*7	*9/*7	-

324   Green cells are genotypes that are consistent with the reported GeT-RM genotype. Purple cells are consistent when ignoring copy
325   number changes. Blue cells are inconsistent with GeT-RM, but consistent with other genotypes reported in the literature with this
326   data.
327

***Assessment of another complex region;* HLA.**

We extended our analysis to explore three more complex loci; those of the *HLA* class I genes. We identified existing ONT sequence

data from 11 subjects who had pooled *HLA-A*, *HLA-B*, and *HLA-C* amplicon data (Anukul *et al.*, 2023). Conveniently, this data

containing three distinct amplicons also allows us to validate that our pipeline can handle multiple independent loci simultaneously.

These datasets had $43,865 \pm 6,587$ (mean $\pm$ standard deviation) reads each. We noted that the coverage of the three loci were

unequal; *HLA-A* being covered by 61 ± 12.6 % of the reads, *HLA-B* by 27 ± 11.6 % of the reads, and *HLA-C* by 10 ± 2.8 % of the

reads. We ran these 11 samples through our pipeline to reconstruct the allele sequences. To assign the *HLA* allele, we performed a

BLAST alignment of each consensus to a database of all *HLA* class I allele sequences obtained from IMGT (Barker *et al.*, 2026).

For *HLA-A*, we reported the correct two alleles for all samples (**Table 3**). *HLA-B* genotypes had one incorrect allele call in two

cases. A single case did not product any *HLA-C* consensus sequence or genotype, likely due to the lower coverage of this locus.

Taken together, we show that our allele reconstruction performs well for a second complex locus with sufficient read coverage,

with an accuracy of 100 % (*HLA-A*), 90.9 % (*HLA-B*), and 90.9 % (*HLA-C*).

Table 3. Summary of HLA genotyping results using data from Anukul et al.

SRA Run ID	Source Sample ID	HLA Genotype	Our Result
ERR11404059	NA02016	A*30:02:01/A*02:01:01 B*53:01:01/B*58:01:01 C*04:01:01/C*07:18	**A*30:02:01/A*02:01:01** **B*53:01:01/B*58:01:01** **C*04:01:01/C*07:18:01**
ERR11404060	NA07439	A*29:02:01/A*68:02:01 B*42:02:01/B*53:01:01 C*04:01:01/C*17:01:01	**A*29:02:01/A*68:02:01** **B*42:02:01/B*53:01:01** **C*04:01:01/C*17:01:01**
ERR11404061	NA12244	A*02:01:01/A*23:01:01 B*15:01:01/B*41:01:01 C*03:03:01/C*17:01:01	**A*02:01:01/A*23:01:01/** **B*15:01:01/B*41:01:01** **C*03:03:01/C*17:01:01**
ERR11404062	NA17039	A*30:01:01/A*31:01:02 B*35:08:01/B*42:02:01 C*04:01:01/C*17:01:01	**A*30:01:01/A*31:01:02** **B*35:08:01/B*42:02:01** **C*04:01:01/C*17:01:01**
ERR11404063	NA17052	A*02:01:01/A*33:03:01 B*39:04/B*44:03:01 C*07:02:01/C*14:03	**A*02:01:01/A*33:03:01/** **B*44:03:01/-** No consensus.
ERR11404064	NA17058	A*24:02:01/A*26:01:01 B*13:01:01/B*40:01:02 C*03:04:01/C*08:03:01	**A*24:02:01/A*26:01:01** **B*13:01:01/B*40:01:02** **C*03:04:01/C*08:03:01**
ERR11404065	NA17119	A*03:01:01/A*74:01:01 B*35:01:01/B*42:02:01 C*07:18/C*17:01:01	**A*03:01:01/A*74:01:01** **B*35:01:01/B*42:02:01** **C*07:18:01/C*17:01:01**
ERR11404066	NA17203	A*24:02:01/A*26:01:01 B*14:01:01/B*35:01:01 C*04:01:01/C*08:02:01	**A*24:02:01/A*26:01:01** **B*14:01:01/B*35:01:01** **C*04:01:01/C*08:02:01**
ERR11404067	NA17240	A*03:01:01/A*30:02:01 B*07:02:01/B*57:01:01 C*06:02:01/C*07:02:01	**A*03:01:01/A*30:02:01** **B*07:02:01/B*57:01:01** **C*06:02:01/C*07:02:01**
ERR11404068	NA17280	A*01:01:01/A*11:01:01 B*08:01:01/B*18:01:01 C*07:01:01/C*07:01:01	**A*01:01:01/A*11:01:01** **B*08:01:01/B*??** **C*07:01:01**
ERR11404069	NA23093	A*11:01:01/A*11:01:01 B*15:02:01/B*15:02:01 C*08:01:01/C*08:01:01	**A*11:01:01** **B*15:02:01** **C*08:01:01**

341   The HLA Genotype column provides the known genotypes for these samples, with orange text for those that were missed in Anukul et al. (Anukul
342   *et al.*, 2023) (by either their in-house pipeline or Type Stream). Our Result shows the result of our allele reconstruction approach, after using
343   BLAST to identify the best-matching HLA allele. Green text denotes a correct genotype call, while orange text denotes an incorrect call. Our
344   accuracy (percent of 22 alleles correctly identified): HLA-A – 100% (22/22 correct), HLA-B – 90.9% (20/22 correct), HLA-C – 90.9% (20/22
345   correct).
346

Allele reconstruction from other loci.

To further test the generalizability of our algorithm to reconstruct allele sequences from ONT amplicon sequence data of any loci,

we downloaded representative datasets from SRA. These included two sequence datasets from the *HLA-B* locus (Ton *et al.*, 2018),

one from *HTT* (Zobel *et al.*, 2025), three from *TRAC* (Nitulescu *et al.*, 2025), one from *EMX1* (B. Wang *et al.*, 2025), one from

CCR5 (Hodge *et al.*, 2025), and three from *ACP4* (Liu *et al.*, 2026). These datasets had variable sequencing depth (median 22,656.5

reads, range 3,503 – 439,236) and amplicon size (median 1.45 kb, range 289 bp – 9.65 kb). We processed all 11 of these datasets

through our pipeline without modification, and compared the resulting consensus sequences with the expected result based on each

source publication (**Table 4**). In all but one case, our resulting consensus sequence was consistent with the reported sequence. For

data from the *HTT* locus, our resulting consensuses sequence underreported the number of CAG repeats by 7.7 %. Overall, these

results demonstrate the utility of our approach for other, unrelated loci.

Table 4. Summary of miscellaneous amplicon reconstruction.

Source Sample ID	Source	Amplicon Description	Flow cell	Source Publication Result	Our Result
PI_B6	PRJNA449245 (Ton *et al.*, 2018)	A 943 bp region of *HLA-B* spanning exons 2 and 3, amplified from Maori and Pacific Island individuals.	MinION R9.4	B*40:01:01G, B*56:02:01	Consensus sequences (931-935 bp) match these two alleles.
PI_F6				B*35:01:01G, B*48:01:01G	Consensus sequences (926-933 bp) match these two alleles.
81Q No_Mod, Clone 8, Day 60	PRJNA1074134 (Zobel *et al.*, 2025)	A 1.95 kb region of the *HTT* locus, amplified from modified hES H9 cells.	MinION R9.4.1	$(CAG)_{78}$-CAACAG	Consensus sequence (1,929 bp) contains 72 CAG repeats.
Mock electroporated r3	PRJNA1257358 (Nitulescu *et al.*, 2025)	A 4.3 kb region of the *TRAC* locus, amplified from unedited CD3+ T cells.	MinION R10.4.1	Unedited TRAC locus	Consensus sequence (4,351 bp) consistent.
Mock electroporated r2				Unedited TRAC locus	Consensus sequence (4,349 bp) consistent.
Mock electroporated r1				Unedited TRAC locus	Consensus sequence (4,351 bp) consistent.
Ec78_inhib_R2	PRJNA1330876 (B. Wang *et al.*, 2025)	A 289 bp region of the *EMX1* locus, amplified from HEK293T cells modified by the Ec78 retron.	Plasmidsaurus (MinION R10.4.1)	GAAGGG converted to AAAGTT at 50% efficiency.	Two consensus sequences (288 bp) built, one with the edit, the other without. Approximate equal read depths.
CasX2_sgRNA_5 (20)_49	PRJNA1334413 (Hodge *et al.*, 2025)	An 814 bp region of the *CCR5* locus, amplified from HEK293-FT cells transfected with plasmids encoding CasX2 and sg5.	MinION R10.4.1	1.2% of loci edited.	Consensus sequence (808 bp) generated showing unedited locus.
Family 3 proband	PRJNA1403257 (Liu *et al.*, 2026)	A 9.65 kb region of the *ACP4* locus, amplified from patients with Amelogenesis imperfecta.	Flongle	c.254T>C, p.(Leu85Pro)	Consensus sequence (9,665 bp) contains this variant.
Family 2 proband				c.254T>C, p.(Leu85Pro)	Consensus sequence (9,665 bp) contains this variant.
Family 1 proband				c.254T>C, p.(Leu85Pro)	Consensus sequence (9,664 bp) contains this variant.

To further validate the generalizability of our approach for allele reconstruction, we used our existing sequence data from

three plasmids (Brown *et al.*, 2023); BCRxV.TF.1 (8.7 kb), BCRxV.VSVG.1 (6.4 kb), and BCRxV.GagPolRev.1 (10.8 kb). Since

these plasmids do not occur in the human genome, we built a synthetic genome reference comprised of a single chromosome

containing each of the plasmid reference sequences. Upon running these three datasets through our pipeline, we obtained a single

consensus sequence for each. All three plasmids resulted in a 99.9 % match to their expected reference. Errors were entirely due to

disagreements between strands, resulting in either an N base call or no base being called due to insufficient coverage. We note that

this is not unexpected, as our previous work using this data required an alternative pseudo-paired base-calling strategy to resolve

strand-specific errors and obtain perfect consensus sequences. Taken together, we have demonstrated the broad applicability of our

algorithm to reconstruct the underlying amplicon sequences from amplicon- or ligation-based ONT sequence datasets.

Discussion

We have presented our solution to reconstruct allele sequences from ONT-based amplicon sequencing, demonstrating its

effectiveness on 22 distinct *CYP2D6* sequence datasets encompassing different amplicons and flow cells, 11 pooled *HLA* class I

datasets, 11 existing datasets from other genomic loci, and three datasets from plasmid sequencing. Our approach is unbiased, only

requiring that a subset of reads exist which fully cover the amplicon of interest, but not requiring *a priori* knowledge of what the

specific amplicon is. The appeal of this approach in diverse genes such as *CYP2D6* is that potential duplications and structural

variants may be reconstructed without pre-existing knowledge of the sequence. For our application of this approach to the *CYP2D6*

locus, we provide post-hoc analysis scripts to automate the variant detection and star-allele assignment, facilitating rapid

interpretation of results. Our approach makes no assumptions regarding the number of gene amplicons (or alleles thereof) present

in the data, only requiring sufficient coverage of each amplicon in order to be reconstructed and reported. Since the primary output

of our approach is the reconstructed allele sequence, novel variants and alleles are readily detectable. In addition to the reconstructed

allele sequence and assigned star-allele, our pipeline also outputs a file with per-position base frequencies and average base

qualities, allowing interrogation of specific positions to assess the confidence of the resulting base. This gives the user more

information, allowing manual interpretation of putative novel variants (**Figure 4**), much like is possible with traditional Sanger

sequencing. Within the context of tamoxifen metabolism by CYP2D6, a recent study demonstrated that a neural-network model

given full-length gene sequences was better able to predict metabolism rates compared to rates predicted from the assigned star-

alleles alone (Van der Lee *et al.*, 2021). This demonstrates the increased power of allele sequence reconstruction, and suggests that

such reconstructions may be even more important in the future as more machine learning models are trained across a wider range

of genes and pathways.

The concept of building a consensus from clustered reads is not unique, however, our approach differs from existing

methods in the following ways. Buermans *et al.* describe the application of the Pacific Biosciences long-amplicon analysis tool

(pblaa) to *CYP2D6* (Buermans *et al.*, 2017)*,* an approach that is based upon the concept that multiple unique consensus sequences

are representative of multiple alleles. This approach is specific to PacBio sequence data, and is closed-source. Athlon (Liu *et al.*,

2018) is a tool developed to *HLA* type ONT data, performing an alignment to *HLA* reference alleles to partition reads, and variant

calling to build the consensus sequences. This is limited to two alleles per *HLA* gene, requires upfront segmentation of reads into

specific loci, and only uses a subset of total reads. Hui *et al.* (Hui *et al.*, 2025) describe an approach to generate consensus 16-23S

rRNA sequences for microbiome analysis. This approach orients sequence reads based on known primer sequences, clusters reads

by similarity, and generates a consensus using Racon or Medaka. While this approach performs well for 16S genes, it is unable to

distinguish separate alleles of human genes differing by a small number of variants, such as SNVs, since reads from both alleles

would be clustered together, and variants would be "error-corrected" away. Similarly, Sahlin *et al*. (Sahlin *et al.*, 2021) describe a

cluster-consensus approach for use in the microbiome field, which suffers from the same limitations when applied to highly similar

alleles of human genes, lacking the sensitivity to output separate consensus sequences that may legitimately differ by only a single

base.

We observed a higher than expected number of short reads in our data. In assessing these reads after mapping to the

genome, they appear to be spread across the whole expected *CYP2D6* amplicon region, and are not coming from any single

contaminant (**Supplementary Figure S1**). Such a signal is consistent with the concept of DNA amplicon fragmentation during

sequence library preparation, resulting in these DNA fragments having adapters ligated and subsequently sequenced. We note that

Liau et al. (Liau, Maggo, *et al.*, 2019) also observed a large fraction of short reads in their data (> 90 % of their reads were shorter

than the expected ~6 kb amplicon in their first sequencing run, ~45 % in their second run after changing the beads used for size

selection to remove DNA fragments shorter than 3-4 kb), so this may be an expected artefact of sequence library preparation which

would require handling in the downstream analysis pipeline. We opted to only consider short reads if they were evidenced to have

been derived from a single conserved, short amplicon.

Existing methods for diplotyping *CYP2D6* from amplicon or whole genome sequence data take a "variant-first" approach

(Hari *et al.*, 2023; Chen *et al.*, 2021; Deserranno *et al.*, 2023; Liau, Maggo, *et al.*, 2019; Ammar *et al.*, 2015), using algorithms

such as Clair3 (Zheng *et al.*, 2022) to call variants, and then these variants are subsequently phased into haplotypes using tools like

WhatsHap (Patterson *et al.*, 2015). Our method differs from existing methods in that it is not reliant on variant calling directly from

sequence reads. Rather, we distil the set of sequence reads into high-confidence, reconstructed allele sequences, which can be

directly compared to representative reference allele sequences for haplotype determination. This is especially helpful when

analyzing ONT reads, which have higher rates of sequencing errors than NGS approaches. Our allele sequence reconstruction

approach is not reliant on a specific number or identity of amplicons within the data, theoretically handling even extreme cases of

copy number variation or gene duplication. Our method uses commonly available Linux tools and open-source software, defined

in Anaconda environments and should be readily deployable on any Linux system.

We ran three existing *CYP2D6* diplotyping tools on our datasets; Aldy4, SpecImmune, and CoLoRGen. Aldy4, despite

not officially supporting ONT data, had similar accuracy as our method. Unfortunately, copy number detection was unable to be

performed by Aldy4 for amplicon data, and so results from Aldy4 do not capture the gene duplication events that our method was

able to identify. SpecImmune generally had lower accuracy compared to our method, with the reported genotype matching the

GeT-RM genotype in only five samples. CoLoRGen similarly had lower accuracy, with only three samples matching the GeT-RM

genotype. Further, CoLoRGen failed to generate a genotype for nine samples.

Our approach requires amplicon data, and specifically requires reads to be uniform in their coverage of the amplicon. As

such, the ONT Rapid (transposase-based) library construction kits are incompatible, as are "shotgun" based approaches such as

ONT adaptive sampling, as they result in read sets with variable start and end positions. Further, with data from a single amplicon,

there is no way to unambiguously assess copy number. Based on the number of reads contributing to each reconstructed allele, we

can approximate the copy number ratio if a sample has two distinct allele sequences. We note that this is a relative abundance

measure rather than an absolute measure of copy number. Despite this, relative abundance is likely sufficient to ascertain clinical

effect, as all *CYP2D6* alleles capable of ultrarapid metabolism do so by duplicating at least once (Hicks *et al.*, 2017). Thus,

determining the relative abundance of normal functioning alleles may be sufficient to designate an ultrarapid metabolizer. Further

copy number increases will elevate the "Activity Score" of the diplotype, but will not change the metabolizer status, nor drug

prescription recommendations. On the other hand, since there is no genomic reference to accurately estimate relative copy number,

if certain diplotypes consist of duplicated homozygous normal function alleles (such as *1/*1x2, *2/*2x2, *33/*33x2, etc.), it is

impossible to determine the copy number and thus the metabolizer phenotype. Multiplexing of multiple genes may help, assuming

that one locus is copy number invariant. Alternatively, a control region of the genome could be included that is known to be copy

number invariant, as is done for data used by Aldy 4 (Hari *et al.*, 2023). The data from Wongsurawat et al. (Wongsurawat *et al.*,

2025) demonstrated that our analysis of *CYP2D6* is not hindered by the presence of multiple amplicons, supporting this as a

reasonable approach in the future.

For the 22 *CYP2D6* datasets examined in this manuscript, none contained any detectable population of reads that

contained full-length gene duplication events, likely due to the standard *CYP2D6* primers resulting in too large of an amplicon to

be efficiently amplified in the event of a gene duplication. Wongsurawat et al. included specific primers targeting the region

between duplicated gene copies, which produced a readily identifiable 3.5 kb amplicon in their two samples with known gene

duplications. Two samples from Liau et al. with known *CYP2D6*68* alleles (a gene hybrid with *CYP2D7*) were both called

incorrectly, with no reads showing the expected fusion event. Alternative primer sets would be worth exploring in the future to

increase the chances of capturing these structural changes and other complex tandem rearrangements, which have not yet been

tested using our analysis approach. Gene deletions, such as *CYP2D6*5*, are also detectable by our pipeline as long as a sufficient

population of reads exist which span the deleted region. We validated this using simulated reads, demonstrating that reads spanning

a large deletion in the genomic reference sequence are effectively reconstructed into the true, deletion containing consensus

sequence. Overall, while we have not comprehensively tested the entire landscape of all known *CYP2D6* variants, we show

consistently positive performance reconstructing the underlying allele sequences for 14 distinct alleles from data sets for which

there are reads supporting these alleles.

Our method is extensible to other genomic loci. We ran our allele reconstruction on 11 samples containing pooled data

from *HLA-A*, *-B*, and *-C*, and show above 90 % accuracy in the resulting HLA genotypes. This supports the use of this method for

other complex loci in the human genome. More generally, we identified 11 existing ONT sequence datasets from six different

genomic loci, demonstrating that our approach faithfully reconstructs the underlying amplicon sequence in each case. Finally, we

show that this approach can be applied to any amplicon-like sequence where an expected reference sequence is known, running it

on plasmid sequence reads for 3 different plasmids. The reconstructed plasmid sequences showed 99.9 % accuracy.

We have demonstrated that *CYP2D6* allele sequence reconstruction and diplotyping can be achieved from modest long-

read sequencing (ONT Flongles), requiring limited specialized equipment, and having low cost. It is appealing that a singular

sequencing reaction coupled with this algorithm may be able to ascertain *CYP2D6* diplotypes with accurate phasing of known or

novel variants, as well as detect gene duplications and structural variants without the need for multiple experiments. This

information can then be used to designate metabolizer status in order to facilitate rational drug prescription as per CPIC guidelines

(Hicks *et al.*, 2017). Currently, to achieve this clinically, a laborious series of assays are conducted, such as Taqman real-time PCR

reactions for common SNPs and indels, copy number verification by qPCR and finally long-range PCR is used for confirmation of

duplications or deletions (Pratt *et al.*, 2021). These assays must have regulatory approval, being performed in a qualified lab using

approved medical devices. Regulatory approval of our approach is possible, particularly by transitioning to the GMP-ready Q-Line

of ONT devices. This would necessitate, at the expense of more complexity, barcoding and multiplexing of samples on larger

MinION flow cells to increase throughput.

<u>Materials and Methods</u>

CYP2D6 *gene amplification.* Genomic DNA was quantified using the Nanodrop, and diluted to 200 ng of DNA in 33.5 uL of nuclease-free

water. Primer sequences were obtained from Ammar et al. (Ammar *et al.*, 2015), using CYP2D6-2F (TAGCTCCCTGACGCCATGATTTGTCTT)

and CYP2D6-2R (CCTGGTTATCCCAGAAGGCTTTGCAG). We used the standard protocol for Phusion High-Fidelity DNA Polymerase (NEB

M0530): 5× Phusion HF Buffer, 10 mM dNTPs, 10 uM forward primer, 10 uM reverse primer, 200 ng template genomic DNA, and 0.5 uL Phusion

DNA polymerase in a 50 uL reaction. PCR cycling conditions were: 98°C for 30 sec., followed by 35 cycles of 98°C for 10 sec. and 72°C for 120

sec., followed by a final step of 72°C for 10 min. and hold at 4°C. PCR products were cleaned up using the Qiagen QIAquick PCR Purification Kit

following manufacturer's instructions.

Sequence library construction. Libraries were prepared using the ONT Ligation Sequencing Kit V14 (SQK-LSK114) following the provided

protocol (LSK114_ACDE_9163_v114_revN_29Jun2022). Samples were quantified with Qubit dsDNA HS Assay kit (Invitrogen Q32851). One

hundred fmols of amplicon DNA was repaired using NEBNext Ultra II End Prep Enzyme Mix provided in the NEBNext Companion Module for

Oxford Nanopore Technologies Ligation Sequencing kit (NEB E7180S). Samples were purified with 30 uL of Ampure XP beads (Beckman,

A63880), and eluted with 31 µL of nuclease-free water. Adapter ligation was performed with the NEBNext Quick T4 DNA Ligase provided in the

NEBNext Companion Module, ONT Ligation Buffer (LNB) and ONT adapter mix (LA). Samples were purified with 20 uL Ampure XP beads,

washed with ONT Long Fragment Buffer, and eluted with 7 µL ONT Elution Buffer (included in SQK-LSK114 kit).

Sequencing using ONT MinION. The Flongle R10.4.1 (FLO-FLG114) flow cell was placed into the Flongle Adapter provided with the Flongle

Starter Pack (FLGIntSP). Prior to sequencing samples, flow cell check was performed following the 'flow-cellpcheck-PQE_1004-V1' protocol to

ensure a minimum of 50 active pores were present. The flow cell was prepared and run according to ONT instructions: the recommended 5-10 fmol

of prepared library was loaded per flow cell, and the run time was set to 24 hours. The Flongle Adapter was USB connected to an M1 Macbook

Pro with 16 GB of RAM. On-device base calling was disabled, instead performing base calling in a later step on our higher-powered computational

resources.

Base calling raw sequence data with Guppy. We assembled all steps of our pipeline into a Snakemake v6.3.0 workflow. We performed initial

base calling of the raw .fast5 data using the GPU version of Guppy v6.5.7, selecting the "dna_r10.4.1_e8.2_400bps_sup.cfg" base calling

configuration file. We ran base calling on a GPU cluster comprised of eight NVIDIA GeForce RTX 3090 GPUs, 256 GB RAM, and two Intel Xeon

Silver 4212 12 core CPUs. All .fast5 files generated by MinKNOW for each sample were submitted to the cluster in its own job, requesting 20 GB

RAM, 1 GPU, and 4 CPUs. To obtain the most useable data possible, we retain the failed reads, as reads with poor quality at critical positions will

be filtered out later in our pipeline. We ran Porechop v0.2.4 (https://github.com/rrwick/Porechop) to remove any remaining adapter sequences and

separate chimeric reads.

Independent control data. We obtained existing base called sequence data of *CYP2D6* from other Coriell samples from Liau et al. (Liau, Maggo,

et al., 2019) and Wongsurawat et al. (Wongsurawat *et al.*, 2025). These datasets were generated in different labs, using different primers, and

sequenced on different versions of the ONT flow cells. These datasets are used to help us validate that our process is not dependent on a specific

amplicon or data source. Class I *HLA* data from Anukul et al. (Anukul *et al.*, 2023) was downloaded to test the performance on a different complex

locus, and to test the performance on datasets with pooled amplicons from multiple independent loci. Data from other human genomic loci were

obtained for the *HLA-B* locus (Ton *et al.*, 2018), the *HTT* locus (Zobel *et al.*, 2025), the *TRAC* locus (Nitulescu *et al.*, 2025), the *EMX1* locus (B.

Wang *et al.*, 2025), the *CCR5* locus (Hodge *et al.*, 2025), and the *ACP4* locus (Liu *et al.*, 2026). Data from plasmid sequencing of three plasmids

was obtained from Brown et al. (Brown *et al.*, 2023), using the subset of reads the authors made available on Zenodo.

Inference of genomic amplicon. All of these raw reads are aligned to the human genome (hg38) using minimap2 v2.17-r941 (Li, 2018) with

the following parameters: "-x map-ont --secondary=no –t 4". To infer the reference amplicon sequence in an unbiased manner, we use samtools

v1.15 (Danecek *et al.*, 2021) to get the depth of read coverage at every position of the genome, and extract out the regions that have depth above

10 % of the max observed depth. This parameter is tunable, potentially needing to be lowered for datasets comprised of amplicons from multiple

distinct loci with uneven coverage. A helper script is made available on the GitHub code repository to visualize the distribution of depths to help

select an appropriate threshold if needed. Any adjacent regions are merged using bedtools v2.26.0 (Quinlan and Hall, 2010) with the following

parameters: "merge -d 100". To aid in the detection of amplicons supporting large deletions, we check all pairs of high-coverage regions for reads

which align to both regions. If 50 % of the reads aligning to each region are shared, we create "chimeric regions" made by concatenating these

regions, which are used in addition to each region alone in subsequent steps.

Candidate read extraction. We extract all the reads that had a primary or supplemental alignment to the identified genomic region(s). We

observed a high number of short reads in our in-house-generated data. To further group reads into those completely covering the same amplicon

sequence, we perform length clustering on the candidate reads. Clustering is done using the DBSCAN algorithm (Schubert *et al.*, 2017) from the

scikit-learn library (v1.8.0), run on the read lengths, with the allowed neighbour length variability (EPS) set to 10 and the minimum core point

neighbours set to 0.1% of the total number of candidate reads, or 5, whichever is greater. This ensures that data sets with fewer reads still have a

reasonable value used. We performed a brief parameter space exploration on three samples, testing all pairwise combinations of EPS values 10, 20,

30, 40, and 50, and minimum neighbor values 5, 10, 15, 25, and 50. The maximum number of reads in the productive length cluster in each case

was obtained with EPS 50 and minimum neighbours 5, however, these large clusters had highly variable read lengths. Decreasing EPS to 10 and

increasing minimum neighbours to 50 dropped the number of reads by an average of 41.3 %, but resulted in clusters of reads with highly similar

lengths; optimal for consensus generation. However, we noted in small data sets, a minimum neighbour value of 50 was too stringent, thus we

implemented the value to scale with the input number of reads. Clusters are selected for consideration for downstream allele reconstruction if the

number of reads in the cluster is at least 25 % of the size of the largest cluster, selected empirically. Additionally, to ensure we do not miss the

cluster with reads the full-length of our inferred amplicon, we select the cluster with the length closest to the inferred amplicon region if it is not

already included.

Allele sequence reconstruction. Our allele reconstruction method is built as a Python (v3.11) script, and takes the alignments of each group of

candidate reads to their inferred amplicon reference sequence obtained using minimap2 v2.17-r941. These alignments are parsed, along with the

raw fastq files to extract per-base quality scores and unaligned sequence flanking each alignment. The unaligned flanking sequences are treated as

insertions, and are only used if at least 25% of the reads support sequence at that position. To facilitate efficient assessment of the pile-up of base

support at every reference position and possible insertion sites, we use byte matrices to store a representation of the multiple sequence alignment

of all reads. To generate this, we step through every alignment to track the maximum observed insertion at every position. We then create empty

byte matrices of reads × positions, where positions are offset relative to the reference sequence to allow for all observed insertions. We then step

through all alignments again to load the observed bases and base quality scores into the byte matrices. The element i,j of the matrix are integer

representations of the base found in the ith read at position j. Separate matrices are maintained for positive and negative strand alignments,

facilitating strand-specific assessment of coverage. The matrices are evaluated, position by position, to assess the number of reads supporting each

base at each position. If a position has evidence of heterozygosity (here, defined as the top two most-observed bases having less than a 2:1 ratio in

abundance, threshold determined empirically), the reads are partitioned into separate groups based on support of the heterozygous bases (**Figure**

3C-D), and consensus sequence construction tries again on each of the new read groups independently. This repeats, recursively, until pure (no

evidence of heterozygosity) consensus sequence(s) are derived, or the read groups are too small to continue. Due to the phenomenon of strand-

specific errors in ONT data (Krishnakumar *et al.*, 2018; Branton *et al.*, 2008), this test of purity is also performed on the forward and reverse reads

independently. Based on a 5kb reference sequence, our algorithm requires 1 CPU core and approximately 3 GB of RAM per 10,000 reads.

, The allele reconstruction script has a number of parameters that can be adjusted to fine-tune the behavior of the allele reconstruction.

Based on all of the control data, we used the following settings, determined empirically, and explained subsequently: --min_depth_factor 0.25, --

absolute_min_depth_thresh 10, --min_base_qual_global 10, --min_base_qual_percentile 33, --sig_to_noise_thresh 2. The "--min_depth_factor"

controls the minimum number of reads needed to support a base at any position, and represents the fraction of the maximum observed depth (here,

25 %). The "--absolute_min_depth_thresh" controls the minimum number of reads supporting a reconstructed allele sequence for it to be output,

here set to 10 reads. The "--min_base_qual_global" is the minimum base quality score needed to be used for allele sequence reconstruction. If this

threshold is not met, then the "--min_base_qual_percentile" is used, being a percentile threshold, per position, that is the lower base quality limit.

This allows for data to be obtained from positions with lower quality overall. Finally, the "--signal_to_noise_thresh" is the value of the minimum

ratio of the most frequent base to the second most frequent base at any position needed to be considered homozygous. This needs to be met when

considering all reads, and is multiplied by a factor of 1.5 when also testing reads from each strand separately. In the case of a position appearing

heterozygous when taking all reads together, but not when assessing each strand individually, the top observed base from the pooled strands is

tested against a softened signal to noise threshold (divided by a factor of 1.5) to call a consensus base, or an N base if a consensus base cannot be

determined. Based on quality thresholds, a read may be "ambiguous" for a given position (supporting a base, but due to low quality the base is not

used). Ambiguous reads do not contribute to depth calculations at that position. If the position is heterozygous, ambiguous reads for that position

are discarded. The average quality score for each consensus base across all reads used for the consensus is recorded and output in the detailed

results file, along with base frequencies, allowing manual inspection of the reconstructed allele sequences.

Automatic genotyping of consensus sequence. Each resulting reconstructed allele sequence is aligned to the *CYP2D6* reference sequence

using Biopython v1.79 pairwise2.align.localxs, and the alignment is parsed to enumerate all variants. All known *CYP2D6* variants are catalogued

from the *CYP2D6* Allele Definition Table from ClinPGx (Whirl-Carrillo *et al.*, 2021, 2012), downloaded from

https://www.pharmgkb.org/page/cyp2d6RefMaterials on January 21, 2025 (now available at https://www.clinpgx.org/page/cyp2d6RefMaterials).

This file was manually curated to ensure consistency in how each variant type was reported, allowing for straightforward parsing. For variants that

can have multiple possible bases allowed for a given allele, all possible allowed values are stored. The set of identified variants for each

reconstructed allele sequence is compared to those from each known allele, and a score is calculated for each known allele based on the number of

disagreements in identified variants within the exon and splice regions of the gene. The allele with the lowest score is returned as the best candidate

allele assignment for the reconstructed allele sequence, along with any missing or extra variants present in the consensus sequence compared to the

star allele reference should they exist. In this way, manual review of the candidate alleles and identified variants allows a user to confirm the allele

assignment, or characterize a novel allele. A helper script is made available on the GitHub code repository to report out the most likely genotype

for each sample based on all identified consensus sequences and abundances. This combines counts for all consensus sequences supporting the

same *CYP2D6* allele, and outputs any allele that represents at least 5 % of the reads of that length-cluster. Alleles with a greater than 2:1 ratio to

each other are reported as possibly supporting a duplication or copy number variation event.

Benchmarking to Aldy4, SpecImmune, and CoLoRGen. We directly compared our results for the 22 *CYP2D6* datasets to results we

obtained from running Aldy4 (Hari *et al.*, 2023), SpecImmune (S. Wang *et al.*, 2025), and CoLoRGen (Rubben *et al.*, 2022), three "variant-first"

approaches. Aldy4 was installed as described on the GitHub repository (https://github.com/0xTCG/aldy) using pip to install the required packages

into a conda environment. To allow Aldy4 to run on ONT amplicon data, copy number detection and variant phasing needed to be disabled. Aldy

genotype was run with "--gene cyp2d6 --profile illumina -o [samplename]-cyp2d6.aldy --cn 1,1 input.bam --param sam_long_reads=true

phase=False". We installed SpecImmune as described on the GitHub repository (https://github.com/deepomicslab/SpecImmune) using conda, after

simplifying the provided conda environment snapshot file to be compatible with our system. SpecImmune was run with "-r porechopped-reads.fastq

-j 16 -i CYP -n samplename -o output_directory -y nanopore --db CYP_ref_db --hg38 hg38_ref.fa". We used the

[samplename].CYP.merge.type.result.txt result file for each sample, taking the reported "CYP2D6 Diplotype". CoLoRGen was installed from the

GitHub repository (https://github.com/laurentijntilleman/CoLoRGen), creating a conda environment with python 3.8, medaka 1.4.3, pandas 1.1.5,

gtfparse 1.2.1, samtools 1.11, minimap2 2.18, tabix 1.11, py-bgzip, bowtie2, and pip to install bio 1.5.9. After manual adjustment of the provided

parameter file, removing unnecessary fields, we ran CoLoRGen with the provided parameters. CoLoRGen failed on nine samples for unknown

reasons. We used the haplotypes_haplotypes.txt file for the final genotype calls.

Simulated deletion detection. To test our algorithm's ability to reconstruct alleles that represent a large deletion relative to the reference genome,

we simulated ONT reads using PBSIM3 (v3.0.5; https://github.com/yukiteruono/pbsim3). We provided a reference template sequence derived from

our amplicon sequence with the entire 4.3 kb *CYP2D6* gene deleted, resulting in an 836 bp template. To simulate 1,000 reads, we created a template

reference fasta file containing this same sequence as 1,000 separate entries. We ran PBSIM3 using qshmm QSHMM-ONT-HQ.model, downloaded

from the GitHub repository. Upon running our pipeline on these simulated reads, we obtained a consensus sequence built from 910 reads which

exactly matched the provided template sequence.

Testing performance on* HLA *class I sequence data. We ran our pipeline with the following modifications to test our ability to generate

HLA consensus sequences from the data from Anukul et al. First, after inspecting the relative depths of the *HLA-A*, *HLA-B*, and *HLA-C* loci in these

samples, we lowered the LOCI_MIN_COVERAGE parameter from 0.10 to 0.04 to ensure that all *HLA* loci were retained. In this data, the coverage

from *HLA-C* is as low as ~4 % of the coverage of *HLA-A* in some samples. Second, to make *HLA* genotype calls on the resulting consensus

sequence, we replaced our automated *CYP2D6* genotyping steps with a BLAST alignment (blastn v2.5.0) (Camacho *et al.*, 2009) to a database of

HLA class I allele sequences downloaded from IMGT (Release 3.63; 2026-01) (Barker *et al.*, 2026), using the following parameters: "-word_size

11 –reward 2 –penalty -3 –gapopen 5 –gapextend 2 –dust no –soft-masking false –evalue 1e10 –max_target_seqs 50". The top scoring *HLA* allele(s)

were output as the best match for each consensus sequence, with genotypes collapsed at the 6-digit level.

Testing performance on other genomic loci. When testing our pipeline on sequence datasets from other, non-complex loci, we ran our

pipeline without modification, aside from skipping the final *CYP2D6* variant calling and allele assignment steps. Resulting consensus sequences

were manually assessed based on the reported results from each source publication, checking if our consensus sequences were consistent with the

results already reported.

Testing performance on plasmid sequence. To test our algorithm's ability to reconstruct sequence not derived from the human genome, we

used our existing plasmid sequence data. To perform this analysis, we created an appropriate reference sequence comprised of a single chromosome

containing the reference sequences of our plasmids. All other aspects of the pipeline were left unchanged, except for skipping of the final *CYP2D6*

variant calling and allele assignment. Resulting consensus sequences were compared directly to the known plasmid sequence for characterization

of any sequence errors. Importantly here, we did not perform the pseudo-paired basecalling as described in our previous work (Brown *et al.*, 2023),

which was required to obtain perfect sequence verification in these plasmids. We chose to analyze these sequences in a more standard manner here,

to better represent typical workflows.

List of Abbreviations

CPU   Central Processing Unit

CYP   Cytochrome P450

DNA   Deoxyribonucleic Acid

GPU   Graphics Processing Unit

HLA   Human Leukocyte Antigen

NGS   Next Generation Sequencing

ONT   Oxford Nanopore Technologies

PCR   Polymerase Chain Reaction

Declarations

Ethics approval and consent to participate

This study was approved by the Fraser Health Research Ethics Board and BC Cancer Agency Research Ethics Board (H21-01422).

Consent for publication

Not applicable.

Availability of data and materials

Custom code is available at the following GitHub repository, along with instructions for use and test data:
https://github.com/scottdbrown/allele-reconstruction-long-read-amplicon-data. A snapshot of the code at the time of publication is
available on Zenodo.org; doi 10.5281/zenodo.19716004. Raw .fastq sequence data for our three sequencing runs is available at the
SRA under Bioproject PRJNA1357883 (https://www.ncbi.nlm.nih.gov/bioproject/1357883).

Competing interests

The authors declare that they have no competing interests.

Funding

This project was supported by a Fraser Health Authority Strategic Priorities Grant and a directed donation funded by Royal
Columbian Hospital Foundation.

Authors' Contributions

RAH, PIM, and SDB designed the study. LD performed library preparation and sequencing optimization. SM, AL, MK, AC, and

MM performed library preparation and sequencing. AM, NW, and MH coordinated and oversaw the technical work conducted in

the Molecular Cytogenetics Laboratory at Royal Columbian Hospital. SDB performed the sequence analysis and algorithm

development. SDB and PIM wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgments

We would like to thank Dr. Martin Kennedy for sharing additional data from Liau et al. (Liau, Maggo, *et al.*, 2019).

References

Ammar,R. *et al.* (2015) Long read nanopore sequencing for detection of HLA and CYP2D6 variants and haplotypes. *F1000Research*, **4**, 1–17.

Anukul,N. *et al.* (2023) Ultrarapid and high-resolution HLA class I typing using transposase-based nanopore sequencing applied in
pharmacogenetic testing. *Front. Genet.*, **14**, 1–10.

Barker,D.J. *et al.* (2026) The IPD-IMGT/HLA database: recent developments in sequence submission. *Nucleic Acids Res.*, **54**, D1152–D1158.

Branton,D. *et al.* (2008) The potential and challenges of nanopore sequencing. *Nat. Biotechnol. 2008 2610*, **26**, 1146–1153.

Brown,S.D. *et al.* (2023) Complete sequence verification of plasmid DNA using the Oxford Nanopore Technologies' MinION device. *BMC*
Bioinformatics, **24**, 1–16.

Buermans,H.P.J. *et al.* (2017) Flexible and Scalable Full-Length CYP2D6 Long Amplicon PacBio Sequencing.

Camacho,C. *et al.* (2009) BLAST+: Architecture and applications. *BMC Bioinformatics*, **10**, 1–9.

Chang,K.L. *et al.* (2015) Pharmacogenetics: Using genetic information to guide drug therapy. *Am. Fam. Physician*, **92**, 588–594.

Charnaud,S. *et al.* (2022) PacBio long-read amplicon sequencing enables scalable high-resolution population allele typing of the complex CYP2D6
locus. *Commun. Biol.*, **5**, 1–10.

Chen,X. *et al.* (2021) Cyrius: accurate CYP2D6 genotyping using whole-genome sequencing data. *Pharmacogenomics J.*, **21**, 251–261.

Danecek,P. *et al.* (2021) Twelve years of SAMtools and BCFtools. *Gigascience*, **10**, 1–4.

Deserranno,K. *et al.* (2023) Targeted haplotyping in pharmacogenomics using Oxford Nanopore Technologies' adaptive sampling. *Front.*
Pharmacol., **14**, 1–11.

Gaedigk,A. *et al.* (2019) Characterization of Reference Materials for Genetic Testing of CYP2D6 Alleles: A GeT-RM Collaborative Project. *J.*
Mol. Diagnostics, **21**, 1034–1052.

Hari,A. *et al.* (2023) An efficient genotyper and star-allele caller for pharmacogenomics. *Genome Res.*, **33**, 61–70.

Hicks,J.K. *et al.* (2017) Clinical pharmacogenetics implementation consortium guideline (CPIC) for CYP2D6 and CYP2C19 genotypes and dosing
of tricyclic antidepressants: 2016 update. *Clin. Pharmacol. Ther.*, **102**, 37–44.

Hodge,C.A. *et al.* (2025) Enhanced cleavage of genomic CCR5 using CASX2Max. *RNA Biol.*, **22**, 1–18.

Hui,Y. *et al.* (2025) De novo clustering of long-read amplicons improves phylogenetic insight into microbiome data. *Gut Microbes*, **17**.

Kiyotani,K. *et al.* (2008) Impact of CYP2D6*10 on recurrence-free survival in breast cancer patients receiving adjuvant tamoxifen therapy. *Cancer*

Sci., **99**, 995–999.

Krishnakumar,R. *et al.* (2018) Systematic and stochastic influences on the performance of the MinION nanopore sequencer across a range of

nucleotide bias. *Sci. Reports 2018 81*, **8**, 1–13.

Van der Lee,M. *et al.* (2021) Toward predicting CYP2D6-mediated variable drug response from CYP2D6 gene sequencing data. *Sci. Transl. Med.*,

13.

Li,H. (2018) Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics*, **34**, 3094–3100.

Liau,Y., Cree,S.L., *et al.* (2019) A multiplex pharmacogenetics assay using the MinION nanopore sequencing device. *Pharmacogenet. Genomics*,

29, 207–215.

Liau,Y., Maggo,S., *et al.* (2019) Nanopore sequencing of the pharmacogene CYP2D6 allows simultaneous haplotyping and detection of

duplications. *Pharmacogenomics*, **20**, 1033–1047.

Liu,C. *et al.* (2018) Accurate Typing of Human Leukocyte Antigen Class I Genes by Oxford Nanopore Sequencing. *J. Mol. Diagnostics*, **20**, 428–

435.

Liu,L. *et al.* (2026) ACP4 Variants In Hypoplastic Amelogenesis Imperfecta Preprint.

van der Maas,S. *et al.* (2025) Dynamic star allele definitions in Pharmacogenomics: impact on diplotype calls, Phenotype predictions and statin

therapy recommendations. *Front. Pharmacol.*, **16**, 1–8.

McDonnell,A.M. and Dang,C.H. (2013) Basic Review of the Cytochrome P450 System. *J. Adv. Pract. Oncol.*, **4**, 263–268.

Nitulescu,A.M. *et al.* (2025) Single-stranded HDR templates with truncated Cas12a-binding sequences improve knock-in efficiencies in primary

human T cells. *Mol. Ther. Nucleic Acids*, **36**, 102568.

Patterson,M.D. *et al.* (2015) WhatsHap: Weighted Haplotype Assembly for Future-Generation Sequencing Reads. *J. Comput. Biol.*, **22**, 498–509.

Pratt,V.M. *et al.* (2010) Characterization of 107 genomic DNA reference materials for CYP2D6, CYP2C19, CYP2C9, VKORC1, and UGT1A1:

A GeT-RM and association for molecular pathology collaborative project. *J. Mol. Diagnostics*, **12**, 835–846.

Pratt,V.M. *et al.* (2016) Characterization of 137 Genomic DNA Reference Materials for 28 Pharmacogenetic Genes: A GeT-RM Collaborative

Project. *J. Mol. Diagnostics*, **18**, 109–123.

Pratt,V.M. *et al.* (2021) Recommendations for Clinical CYP2D6 Genotyping Allele Selection: A Joint Consensus Recommendation of the

Association for Molecular Pathology, College of American Pathologists, Dutch Pharmacogenetics Working Group of the Royal Dutch

Pharmacists Association. *J. Mol. Diagnostics*, **23**, 1047–1064.

Puaprasert,K. *et al.* (2018) Real time PCR detection of common CYP2D6 genetic variants and its application in a Karen population study. *Malar.*

J., **17**, 8–11.

Qiao,W. *et al.* (2016) Long-Read Single Molecule Real-Time Full Gene Sequencing of Cytochrome P450-2D6. *Hum. Mutat.*, **37**, 315–323.

Quinlan,A.R. and Hall,I.M. (2010) BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics*, **26**, 841–842.

Rubben,K. *et al.* (2022) Cas9 targeted nanopore sequencing with enhanced variant calling improves CYP2D6-CYP2D7 hybrid allele genotyping.
PLoS Genet., **18**, 1–21.

Sahlin,K. *et al.* (2021) NGSpeciesID: DNA barcode and amplicon consensus generation from long-read sequencing data. *Ecol. Evol.*, **11**, 1392–
1398.

Sayers,E.W. *et al.* (2025) Database resources of the National Center for Biotechnology Information in 2025. *Nucleic Acids Res.*, **53**, D20–D29.

Schubert,E. *et al.* (2017) DBSCAN Revisited, Revisited. *ACM Trans. Database Syst.*, **42**.

Somogyi,A.A. *et al.* (2015) Pharmacogenetics of opioid response. *Clin. Pharmacol. Ther.*, **97**, 125–127.

Teusink,A. *et al.* (2016) Genotype-directed dosing leads to optimized voriconazole levels in pediatric patients receiving hematopoietic stem cell
transplantation. *Biol. Blood Marrow Transplant.*, **22**, 482–486.

Ton,K.N.T. *et al.* (2018) Multiplexed Nanopore Sequencing of HLA-B Locus in Māori and Pacific Island Samples. *Front. Genet.*, **9**, 1–12.

Wang,B. *et al.* (2025) Structural basis for retron co-option of anti-phage ATPase-nuclease. *Nat. Struct. Mol. Biol.*, **33**.

Wang,S. *et al.* (2025) SpecImmune accurately genotypes diverse immune-related gene families using long-read data.

Whirl-Carrillo,M. *et al.* (2021) An Evidence-Based Framework for Evaluating Pharmacogenomics Knowledge for Personalized Medicine. *Clin.*
Pharmacol. Ther., **110**, 563–572.

Whirl-Carrillo,M. *et al.* (2012) Pharmacogenomics knowledge for personalized medicine. *Clin. Pharmacol. Ther.*, **92**, 414–417.

Wongsurawat,T. *et al.* (2025) Enhancing CYP2D6 genotyping with nanopore sequencing to address allele diversity P . vivax malaria elimination.
Sci. Rep., **15**, 33842.

Zheng,Z. *et al.* (2022) Symphonizing pileup and full-alignment for deep learning-based long-read variant calling. *Nat. Comput. Sci.*, **2**, 797–803.

Zobel,M. *et al.* (2025) A human CAGinSTEM platform for decoding HTT repeats ' somatic instability links CAG interruption to HD pathology in
neurons ll A human CAGinSTEM platform for decoding HTT repeats ' somatic instability links CAG interruption to HD pathology in
neurons. *CellReports*, **44**, 116685.

Supplemental

Supplementary Table S1. Summary table of all inferred amplicons, length-clusters, and allele reconstruction attempts with read

counts at each stage.

Supplementary File S1. All successfully constructed consensus sequences for all *CYP2D6* samples.

Supplementary Figure S1. *CYP2D6* **read coverage from short reads.** Screenshot from Geneious Prime. The length-cluster of

short reads from NA17300 (40,717 reads with mean length 445 bases) was aligned to the *CYP2D6* reference sequence. Coverage

is shown in blue, spanning the entire expected amplicon. Exons are marked with grey segments under the green *CYP2D6* gene.

Coverage is variable, and the region of highest coverage (exon 4-6) yielded a single reconstructed consensus sequence.

Supplementary Figure S2. Alignment of long and short consensus sequences to *CYP2D6*. Screenshots from Geneious Prime.

For NA10005 (top) and NA17300 (bottom), short consensus sequences were obtained from the respective length-clusters of short

reads. The sequences of these short consensus sequences match at least one of the long consensus sequences in each sample, with

the exception of mismatches at the beginning and end of the sequences.

Supplementary Figure S3. Dot plot of duplication-detecting consensus sequence and *CYP2D6*. Screenshot from Geneious

Prime. The consensus sequence from NA02016_ligate from the 3.5 kb duplication-detecting amplicon (y-axis) is compared to the

CYP2D6 reference sequence (x-axis), showing the consensus sequence supporting the region upstream and downstream of *CYP2D6*

as would be expected in the event of an allele duplication.

Supplementary Figure S4. Suspected *CYP2D6-CYP2D7* hybrid allele sequence aligned to *CYP2D6* and *CYP2D7*. Screenshots

from Geneious Prime. The *CYP2D7*-locus-derived consensus sequence from NA17203_ligate is aligned to the *CYP2D6* reference

(top) and *CYP2D7* reference (bottom), with the sequence identity as the green histogram above the sequences. Regions of lower

sequence identity to either reference are boxed in light blue. This consensus sequence appears to best support a *CYP2D6-CYP2D7*

hybrid allele, with exon 1 from *CYP2D6*, and the remainder of the sequence from *CYP2D7*.

758   **Supplementary Figure S5. Alignment of NA17203_ligate *CYP2D6-CYP2D7* hybrid allele sequence to *68 hybrid allele**

759   **sequence.** Screenshots from Geneious Prime. (A) Our NA17203_ligate *68 allele sequence shows a perfect match to the *68

760   sequence identified from NA12878 identified by Rubben et al, (B) with the exception of a C deletion observed in our sequence

761   within a homopolymer stretch. (C) When our *68 hybrid allele sequence is aligned to the *CYP2D6* reference sequence, this deletion

762   falls between exons 3 and 4, and thus would not be expected to change the function of the resulting protein.