Occurrence of *Nigrospora* spp. as the predominant causal agents of leaf spot disease in

Cavendish banana in banana plantations in Mindanao Island, Philippines

Short title: *Nigrospora* spp.: Major pathogens on Cavendish banana in Mindanao

Shunsuke Nozawa[1*], Yui Harada[1], Yoshiki Takata[1], Keiko Uchida[1], Mike Andre Malonzo[2],

Reynaldo Valle[2], Sherman M. Chavez[3], Aniway F. Penalosa[3], Kyoko Watanabe[1*]

[1] School of Agriculture, Tamagawa University, Machida, Tokyo, Japan

[2] BaCaDM project of College of Agriculture, Tamagawa University, 6-1-1 Tamagawa-

gakuen, Machida, Tokyo 194-8610, Japan

[3] Unifrutti Tropical Philippines, Inc., Km. 15, Panacan, Davao City 8000, Philippines

*Corresponding author:

E-mail: nozao802@gmail.com (N.S.) and wkyoko@agr.tamagawa.ac.jp (W.K.)

Abstract

Banana leaf diseases are a significant threat to Cavendish banana production. In the

Philippines, the main disease has been diagnosed as Black sigatoka disease caused by

Pseudocercospora fijiensis based on symptoms. However, our study showed that the main

pathogen in Mindanao island, the largest banana-producing region in the Philippines, belongs to

the genus *Nigrospora*, contradicting previous assumptions. We clarified the phylogenetic

positions of 160 *Nigrospora* isolates based on molecular phylogenetic analyses using ITS, *β-*

tubulin, and *tef1α* sequences, and compared their morphology with known species. Molecular

phylogenetic and morphological analysis revealed that *Nigrospora* isolates comprised *N.*

chinensis, *N. lacticolonia*, *N.* cf. *singularis*, *N. sphaerica*, *N. vesicularifera*, and a novel species,

N. nigrocolonia. Pathogenicity tests on banana leaves confirmed that these species are

pathogenic. Species other than *N. sphaerica* were for the first time reported as pathogens of

banana leaf. The results of the fungicide sensitivity test using 14 fungicides, including

pyrimethanil, spiroxamine, and tebuconazole, for the Sigatoka disease showed 100% inhibition

of all isolates at 100 ppm of active ingredients. However, low-sensitivity isolates were observed

for the remaining 11 fungicides. Our findings indicated the need for a comprehensive review of

banana leaf disease prevention strategies.

Introduction

Banana is a cash crop in Southeast Asia, Africa, and Latin America, with an annual production of 1.51 million tons in 2023, making it the most produced fruit globally (FAOSTAT 2023). The Cavendish banana is the most widely traded and globally distributed variety among all banana cultivars. Cavendish is exported in significant quantities among the banana cultivars traded internationally. More than 20 million tons of Cavendish banana are exported. The leading exporters of Cavendish banana include Colombia, Costa Rica, Ecuador, Guatemala, and the Philippines, and its major importers are the European Union and United States.

Banana production is affected by diseases such as Fusarium wilt caused by *Fusarium* spp. [1,2], leaf blight by *F. saccari* and *Alternaria* jacinthicola [3,4]; leaf spots by *Colletotrichum* spp., *Pseudocercospora eumusae*, and *Nigrospora* shaerica [5–7]; and Sigatoka disease by *Pseudocercospora* spp. [8]. Among these diseases, Sigatoka disease is one of the most serious, causing an estimated 3% annual reduction in banana yield and economic losses of approximately 1.6 billion dollars annually [9]. The Sigatoka disease is caused by several *Pseudocercospora* spp.; black Sigatoka by *P. fijiensis*, and yellow Sigatoka by *P.* musae [10]. Among these diseases, *P. fijiensis* (black Sigatoka) is the most virulent and causes serious damage. Since its discovery in Fiji in 1964 [11], the disease has spread globally owing to its virulence and infectiousness, becoming one of the most serious threats to banana production.

The Philippines is the world's fifth-largest producer of bananas, with an annual output

of approximately 5.9 million tons in 2023 (FAOSTAT 2023). Bananas are pivotal cash crops for

export, with an estimated annual export volume of approximately 2 million tons. However, the

country is grappling with a significant challenge: black Sigatoka disease, the management of

which has necessitated substantial financial investments [10]. To address this issue, a leaf

removal strategy was implemented to eliminate leaves exhibiting disease symptoms and

maintain a minimum of five leaves per plant. This approach involves fungicide applications,

sprayed approximately 50 to 70 times per year, to prevent disease transmission. According to

the Banana Working Group of the Fungicide Resistance Action Committee (FRAC), the

following fungicide groups are recommended for the control of Sigatoka disease: amines,

anilinopyrimidines (APs), benzimidazoles (BCMs), demethylase inhibitors (DMIs), guanidines,

multi-site inhibitors, N-Phenylcarbamates, Qi Inhibitors (QiIs), Qo Inhibitors (QoIs), Succinate

Dehydrogenase Inhibitors (SDHIs), and Toluamides. Fungicide application incurs high annual

costs and is the most expensive activity in banana production. However, there is a paucity of

surveys conducted in the Philippines to investigate the emergence of new pathogens despite the

presence of multiple diseases that can present spot symptoms similar to Sigatoka disease in the

world [5,7,12].

Our study shows that *Nigrospora* spp., rather than *Pseudocercospora* spp., is the

dominant pathogen on the island of Mindanao in the Philippines. This symptom closely

resembles that of Sigatoka disease, as it causes the formation of brown to black spots with a

yellow halo and the lesions expand along the leaf veins (Fig. 1). Consequently, locally, any

disease exhibiting such symptoms is diagnosed as Sigatoka disease.

To provide basic information for control measures against this disease, which is

mistakenly diagnosed as Sigatoka disease, we identified the pathogens belonging to the genus

Nigrospora at the species level and conducted haplotype analysis. In addition, in vitro

sensitivity tests were conducted to identify effective fungicides currently used to control black

Sigatoka disease and hypothesize the potential emergence of resistant *Nigrospora* spp. isolates.

Results

Frequency of the genus *Nigrospora*

We obtained 217 isolates from diseased banana leaves. As a preliminary identification to

determine the genus, a BLAST search of the ITS region in the NCBI database was conducted. A

Basic Local Alignment (BLAST) search showed that our isolates consisted of 18 genera (Table

1). Out of 217 isolates, 160 belonged to the genus *Nigrospora*, which was the most frequently

detected fungal genus. In addition, the genus *Nigrospora* was dominant at all the collection

locations. The proportions of *Nigrospora* were 96.6%, 70.2 %, 53.5 %, 92.9 %, 69 %, and 75%

in Dabong-dabong, Little Panay, Panambon, Patag-Alanib, Tugbok, and Ula, respectively. We

did not perform further analyses to identify the species of isolates that did not belong to the

genus *Nigrospora*.

Phylogeny of isolates belonging to genus *Nigrospora*

To reveal the phylogenetic position of our isolates, we constructed molecular phylogenetic

trees based on the combined data set of ITS, *β-tubulin*, and *tef1a* nucleotide sequences. The

alignment length, number of variable sites, and number of parsimony-informative sites in the

DNA data were 1,156, 545, and 461 bp, respectively (see Supplementary Data Set). The

phylogenetic trees (Fig. 2; see Supplementary Data Set) showed that our isolates belonged to a

clade comprised of *N. camelliae-sinensis* M. Wang & L. Cai and *N. singularis* M. Raza & L. Cai

(30 isolates: 13.4% of all isolates; BPP/ML = 0.99/92, *N. chinensis* clade (1 isolate: 0.5% of all

isolates; BPP/ML = 1.0/99), *N. lacticolonia* (118 isolates: 54.6% of all isolates; BPP/ML =

1.0/99), *N. nigrocolonia* Nozawa & Kyoko Watan. clade (9 isolates: 11.1% of all isolates; BPP =

1.0), *N. sphaerica* clade (1 isolate: 0.5% of all isolates; BPP/ML = 1.0/99), and *N.*

vesicularifera clade (1 isolate: 0.5% of all isolates; BPP/ML = 1.0/99). Regarding the isolates

belonging to a clade comprising of *N. camelliae-sinensis* and *N. singularis*, the evolutionary

distance between the isolates and the ex-type strain of *N. singularis* (CGMCC 3.19334) is closer

than that to *N. camelliae-sinensis*. However, identifying the isolates collectively as *N. singularis*

is not possible because *N. camelliae-sinensis* belongs within the same clade. Therefore, in this

study, these isolates were treated as strains closely related to *N. singularis* and designated as *N.*

cf. *singularis*. Although the clade comprising *N. camelliae-sinensis* and *N. singularis* warrants

further taxonomic examination, in the present study, we focused on identification and therefore

did not discuss this limitation.

The phylogenetic positions of the clades to which the isolates belong are as follows. *N.*

chinensis clade neighbored *N. humicola* Qin Yang & Ning Jiang (BPP/ML = 1.0/99), *N.*

lacticolonia clade neighbored *N. osmanthi* Mei Wang & L. Cai (BPP = 0.99/86), *N.*

nigrocolonia clade, a new lineage, neighbored a clade comprising *N. cooperae* Y.P. Tan, Bishop-

Hurley, Bransgr. & R.G. Shivas, *N. guilinensis* Mei Wang & L. Cai, *N. lacticolonia* Mei Wang

& L. Cai clade, *N. marylouisemclawsiae* Y.P. Tan, Ryley & Bishop-Hurley, and *N. osmanthi*, *N.*

sphaerica clade neighbored *N. brasiliensis* A.C.Q. Brito, C. Conforto & A.R. Machado

(BPP/ML = 1.0/99), and *N. vesicularifera* clade neighbored *N. saccharicola* M. Raza & L. Cai

(BPP/ML = 0.99/99).

Phylogenetic analyses were conducted based on three DNA regions (ITS, β-tubulin, and

tef1a; Supplementary Figure 1; see Supplementary Data Set) independently. Among the isolates,

thirty strains identified as *N.* cf. *sphaerica* formed a monophyletic clade with the reference

strains of *N. camelliae-sinensis*, *N. falsivesicularis* M. Raza & L. Cai, and *N. pyriformis* Mei

Wang & L. Cai in the phylogenetic tree based on ITS region (ITS tree; BPP < 0.95). In contrast,

these strains formed a strongly supported monophyletic clade (BPP = 1.0) with *N. camelliae-*

sinensis, *N. pyriformis*, and *N. sphaerica* (Sacc.) E.W. Mason in the phylogenetic tree based on

the *β-tubulin* region *(β-tubulin* tree; Supplementary Figure 2). In the phylogenetic tree based on

tef1a (*tef1a* tree; Supplementary Figure 3), although posterior probabilities were not indicated,

the isolates formed a monophyletic group with *N. singularis*.

The isolate identified as *N. chinensis* Mei Wang & L. Cai consistently formed a well-

supported monophyletic clade (BPP = 1.0 for all loci) with the reference strains of *N. chinensis*,

including the ex-type strain, across all ITS, β-tubulin, and *tef1a* trees (Supplementary Figure 1,

2, 3). The isolates identified as *N. lacticolonia* formed a monophyletic clade with *N. osmanthi*

and *N. vesicularifera* M. Raza & L. Cai in the ITS tree; however, they clustered with the

reference strains of *N. lacticolonia* (including the ex-type strain) in both the β-tubulin and *tef1α*

trees (BPP = 1.0/0.83: β-tubulin/*tef1a*). Similarly, the isolate identified as *N. sphaerica* formed a

monophyletic clade with the reference *N. sphaerica* (including the ex-type strain) across all loci

(BPP = 0.93/1.0/1.0 for ITS/*β-tubulin*/*tef1a*). The isolate identified as *N. vesicularifera* clustered

with the reference strains of *N. vesicularifera* (including the ex-type strain) in both the β-tubulin

and *tef1a* trees (BPP = 1.0/1.0), while in the ITS tree, it formed a monophyletic group with both

N. vesicularifera and *N. saccharicola* (BPP = 1.0). Although species-level resolution was not

always clearly delineated in phylogenies based on single locus, all isolates were positioned

within clades closely related to the corresponding identified species. Furthermore, the newly

established species *N. nigrocolonia* was independent from any known species in all

phylogenetic trees (BPP = 1.0/1.0/1.0: ITS/*β-tubulin* /*tef1a*).

Focusing on isolation frequency at each collection site, *N. lacticolonia* was the most

frequently isolated species in Little panay, Panambon, Tugbok, and Ula. Meanwhile, *N.*

singuralis M. Raza & L. Cai was the most frequently isolated species from Dabong-dabong and

Patag-Alanib (Fig. 3a, b).

Taxonomy

Nigrospora nigrocolonia Nozawa & Kyoko Watan., sp. nov. (Fig. 4.; Supplementary Figure 4)

MycoBank no: MB 858325.

Culture characteristics: Colonies on PDA reached 90 mm in diameter. After 5 days of incubation

at 25 ± 0.5 °C, the colonies had smooth margins, initially white to whitish cream with abundant

cottony aerial mycelia; the reverse was initially whitish cream (Fig. 4a, b). With age, the

colonies on the surface became grayish, and dark olive-to-black on the reverse (Fig. 4c, d).

Asexual morph on SNA: Hyphae are hyaline to brown, highly-branched, septate, smooth,

and 2.2–4.8 (mean ± SD = 3.1±0.1, n = 30) μm wide. Conidiophores are micronematous or

semi-macronematous, flexuous, olivaceous or brown, smooth (Fig. 4e), measuring 5.1–

10.6×2.5–4.3 μm (mean ± SD = 8.5±1.2×3.3±0.4, n = 4), frequently reduced to conidiogenous

cells (Fig. 4e-g). Conidiogenous cells are aggregated in clusters or dispersed on hyphae,

monoblastic, determinate, variable in shape, mostly subspherical to cylindrical, and olivaceous

or brown, measuring 5.4–15.2×5.9–10.8 μm (mean ± SD = 8.7±0.3×7.9±0.2, n = 30). Conidia

are solitary, aseptate, globose 13.2–18.7×14–18 (mean ± SD = 16.3±0.2×16.5±0.2, n = 30), and

subglobose to ellipsoidal 14.4–19.2×11.7–15 (mean ± SD = 16.2±0.2×13.1±0.2, n = 30) (Fig.

4e-g). No chlamydospores were observed in this study. Sexual morphs were not observed.

Etymology: Specific epithet refers to the black coloration of the colony on the back.

Type: Dried culture specimen grown on SNA (TAP H-24N202, holotype); ex-type strain

TAP24N202 isolated from diseased leaf of *Musa acuminata*, in Patag-Alanib, Lantapan,

Bukidnon, Philippines, on December 11, 2024, isolated by Shunsuke Nozawa. The holotype

specimen and ex-type strain were deposited in the culture collection of Tamagawa University,

Japan. The GenBank accession numbers for the type strain of Nigrospora nigrocolonia are as

follows: ITS: LC868094, *β-tubulin*: LC868147, and *tef1a*: LC868200.

Note: *N. nigrocolonia* forms a well-supported clade as a neighbor of a clade comprising *N.*

aurantiaca, *N. cooperae*, *N. falsivesicularis*, *N. guilinensis*, *N. lacticolonia*, *N.*

marylouisemclawsiae, *N. osmanthi*, *N. stoneae*, and *N. vesicularis* in the phylogenetic tree based

on cancate. The holotype of *N. nigrocolonia* exhibited sequence differences with other

Nigrospora species as follows: *N. aurantiaca* CGMCC 3.18130 showed ITS 1.1%, *β-tubulin*

12.4%, and *tef1a* 20.1%; *N. cooperae* BRIP 72440a showed ITS 1.9%, β-*tubulin* 13.0%, and

tef1a 21.1%; *N. falsivesicularis* CGMCC3.19678 showed ITS 2.4%, *β-tubulin* 12.3%, and *tef1a*

21.4%; *N. guilinensis* CGMCC 3.18124 showed ITS 1.6%, β-tubulin 11.7%, and *tef1a* 20.5%; *N.*

lacticolonia CGMCC 3.18123 showed ITS 1.6%, *β-tubulin* 9.0%, and *tef1a* 20.7%; *N.*

marylouisemclawsiae BRIP 74865b showed ITS 1.3%, *β-tubulin* 9.0%, and *tef1a* 20.4%; *N.*

osmanthi CGMCC 3.18126 showed ITS 1.6%, *β-tubulin* 10.0%, and *tef1a* 19.0%; *N. stoneae*

BRIP 75022a showed ITS 1.1%, *β-tubulin* 15.7%, and *tef1a* 20.3%; and *N. vesicularis* CGMCC

3.18128 showed ITS 1.3%, *β-tubulin* 12.1%, and *tef1a* 20.6%.

N. nigrocolona differed from related species in exhibiting dark olivaceous conidiogenous

cells upon maturation. The species is further distinguished by its conidial size. The conidial size

of *N. nigrocolonia* was notably bigger—measuring 13.2–18.7×14–18 (mean ± SD =

16.3±0.2×16.5±0.2)—than those of *N. aurantiaca* (mean ± SD = 14.82 ± 0.79×11.78 ± 1.07

µm), *N. cooperae* (10–12.5×8.5–10.5 µm), *N. falsivesicularis* (mean ± SD = 13.62 ± 1.8 µm), *N.*

guilinensis (mean ± SD = 12.46 ± 0.62 × 9.69 ± 0.71 µm), *N. lacticolonia* (mean ± SD = 14.36

± 1.04 µm), *N. osmanthi* (mean ± SD = 14.87 ± 0.63 µm), and *N. vesicularis* (mean ± SD = 14.8

± 0.76 µm). Morphological data for *N. marylouisemclawsiae* and *N. stoneae* are currently

unavailable and were therefore excluded from conidial comparisons.

In addition to the type strain identified as *N. nigrocolonia*, the following isolates—

TAP24N222, TAP24N224, TAP24N229, TAP24N232, TAP24N237, TAP24N246, TAP24N250,

and TAP24N252—have been deposited in the Tamagawa University Mycological Bio-bank of

Plant Diseases.

Morphological descriptions of isolates

In this section, species other than *N. nigrocolonia* isolated in this study are described. For

species from which multiple strains were obtained, one strain was randomly selected for

morphological characterization.

Nigrospora cf. *singularis*: Isolate no. TAP24N068: Culture characteristics: Colonies on

PDA reached 90 mm in diameter. After 4 days of incubation at 25 ± 0.5 °C, the colonies had

smooth margins and were white to whitish gray with abundant cottony aerial mycelia; the

reverse was initially whitish cream (Fig.5a, b). Asexual morph on SNA: Hyphae are hyaline to

pale brown, branched, septate, smooth, and 1.2–5.1 (mean±SD = 2.4±0.13, n = 30) μm wide.

Conidiophores are micronematous or semi-macronematous, flexuous, hyaline, and smooth.

Conidiogenous cells are dispersed on hyphae, monoblastic, determinate, variable in shape,

mostly subspherical to oblate, hyaline, measuring 6.4–16.1×6.4–17.3 (mean±SD =

9.4±0.41×8.1±0.35, n = 30) in size (Fig. 5c, d). Conidia are solitary, aseptate, and subglobose to

ellipsoidal 9.8–13×9–13 (mean±SD = 11.6±0.18×11±0.22, n = 30; Fig.4c, d). No

chlamydospores were observed in this study. Sexual morphs were not observed. The size of the

conidia of this strain closely matched the original description of *N. singularis* (CGMCC

3.19334) [13]: 9.5–13 × 11–15 (mean ± SD = 11.2 ± 1.11 × 13.4 ± 1.12) μm.

N. chinensis: Isolate no. TAP24N249: Culture characteristics: Colonies on PDA reached 90

mm in diameter. After 7 days of incubation at 25 ± 0.5 °C, the colonies had smooth margins and

were white to whitish cream, with abundant cottony aerial mycelia. the reverse was whitish gray

to pale brown (Fig. 6a, b). Asexual morph on SNA: Hyphae are hyaline to pale brown, highly-

branched, septate, smooth, and 1.5–4.6 (mean±SD = 2.7±0.14, n = 30) μm wide. Conidiophores

are micronematous or semi-macronematous, flexuous, hyaline, smooth, frequently reduced to

conidiogenous cells (Fig. 6c, d). Conidiogenous cells are dispersed on hyphae, monoblastic,

determinate, variable in shape, mostly subspherical to oblate, hyaline, and 4.8–12.6×6.9–10.5

(mean±SD = 8.8±0.31×8±0.16, n = 30) in size. Conidia are solitary, aseptate, and subglobose to

ellipsoidal 10.5–15.6×7.8–13 (mean±SD = 12.8±0.23×10.2±0.2, n = 30; Fig. 6c, d). No

chlamydospores were observed in this study. Sexual morphs were not observed. The size of

conidia of this strain closely matched the original description of *N. chinensis* (CGMCC

3.18127)[14]: 10–14.5 × 7.5–11 (mean±SD = 11.78 ± 0.75 × 9.18 ± 0.61) μm.

N. lacticolonia: Isolate no. TAP24N011: Culture characteristics: Colonies on PDA reached

90 mm in diameter. After 3 days of incubation at 25 ± 0.5 °C, the colonies had smooth margins,

and were white to whitish gray with abundant cottony aerial mycelia; the reverse was whitish

cream to pale brown (Fig. 7a, b). Asexual morph on SNA: Hyphae are hyaline to pale brown,

highly-branched, septate, smooth or verrucose, and 2–5.8 (mean±SD = 3.7±0.16, n = 30) μm

wide. Conidiophores are micronematous or semi-macronematous, flexuous, hyaline, smooth or

verrucose, frequently reduced to conidiogenous cells. Conidiogenous cells are dispersed on

hyphae, monoblastic, determinate, variable in shape, mostly subspherical to oblate, hyaline, and

4.6–14.4×5.7–10.9 (mean±SD = 8.2±0.35×7.8±0.2, n = 30) in size (Fig. 7c, d). Conidia are

solitary, aseptate, subglobose to ellipsoidal 13.2–16.5×12.2–16.1 (mean±SD =

15.3±0.17×14.9±0.15, n = 30; Fig 7c, d). No chlamydospores were observed in this study.

Sexual morphs were not observed. The size of conidia of this strain closely matched the original

description of *N. lacticolonia* (CGMCC 3.18123) [14]: 11.5–16.5 (mean±SD. = 14.36 ± 1.04)

μm diameter.

Nigrospora sphaerica: Isolate no. TAP24N107: Culture characteristics: Colonies on PDA

reached 90 mm in diameter. After 3 days of incubation at 25 ± 0.5 °C, the colonies had smooth

margins, and were white to whitish gray with abundant cottony aerial mycelia; the reverse was

whitish cream (Fig. 8a, b). Asexual morph on SNA: Hyphae are hyaline to pale brown, branched,

septate, smooth, and 1.9–4.2 (mean±SD = 3.1±0.12, n = 30) μm size. Conidiophores are

micronematous or semi-macronematous, flexuous, hyaline, smooth, and frequently reduced to

conidiogenous cells. Conidiogenous cells are dispersed on hyphae, monoblastic, determinate,

variable in shape, mostly subspherical to oblate, hyaline to pale brown, and 6.6–11.7×5.8–17.2

(mean±SD = 9.3±0.23×8.7±0.38, n = 30) in size (Fig. 8c, d). Conidia are solitary, aseptate,

subglobose to ellipsoidal 15.6–22.6×15.2–22.9 (mean±SD = 19.6±0.33×19.5±0.36, n = 30; Fig.

8c, d). No chlamydospores were observed in this study. Sexual morphs were not observed. The

size of conidia of this strain closely matched the description of *N. sphaerica* (LC2840)

measured by Wang et al. [14]: 16–21 (mean±SD = 18.22 ± 1.0) μm diameter.

N. vesicularifera: Isolate no. TAP24N023: Culture characteristics: Colonies on PDA

reached 90 mm in diameter. After 3 days of incubation at 25 ± 0.5 °C, the colonies had smooth

margins, and were white to whitish gray with abundant cottony aerial mycelia; the reverse was

whitish cream to brown (Fig. 9a, b). The isolate did not produce conidia. Sexual morphs were

not observed.

Haplotype analyses

Haplotype analysis based on the *tef1α* gene sequences was conducted for *N. lacticolonia*, *N. nigrocolonia*, and *N. singularis*, which were obtained as multiple isolates (Fig. 3a, b). *N. lacticolonia* included 118 isolates, and the alignment consisted of 616 sites. This species was divided into four haplotypes, characterized by six variable sites (S) and a nucleotide diversity (Pi) of 0.001586. Among all the isolates, 78 belonged to Hap_1, 38 to Hap_2, and 1 each to Hap_3 and Hap_4. Hap_1 was found at all six collection sites, whereas Hap_2 was found at five collection sites, excluding Patag-Alanib. Hap_3 was isolated solely from Little Panay, and Hap_4 was isolated solely from Panombon.

N. nigrocolonia included 9 isolates, and the alignment consisted of 714 sites. This species was divided into two haplotypes, with three variable sites (S) and a nucleotide diversity (Pi) of 0.001634. Two isolates belonged to Hap_1, and seven to Hap_2. Both haplotypes were isolated from Dabong-dabong and Patag-Alanib cultivars.

N. singularis included 30 isolates, and the alignment consisted of 604 sites. This species was divided into three haplotypes, with three variable sites (S) and a nucleotide diversity (Pi) of 0.001254. Among all isolates, 21 belonged to Hap_1, 4 to Hap_2, and 5 to Hap_3. Hap_1

was found at five of the collection sites, excluding Panombom, whereas Hap_2 was found at

Dabong-dabong and Patag-Alanib. Hap_3 was isolated from the panomboms and patag-alanib.

Pathogenicity to leaves

All 160 isolates were tested on wounded and non-wounded banana leaves. The

aggressiveness of the tested isolates on wounded leaves differed significantly, with lesion

diameters ranging from 2.33 mm to 22.61 mm (Fig. 10; Table 2; Supplementary Figure 5).

Based on diameter sizes of the disease spots, aggressiveness was classified as strongly virulent

($\geqq$6.6 mm), moderately virulent ($\geqq$3 to <6.6 mm), weakly virulent (> 0 to <3 mm), or very

weakly virulent (0 mm with brown or yellow discoloration at or around the wound site). Among

the 160 isolates investigated, 54 (33.75%) were strongly virulent, 52 (32.5%) moderately

virulent, 43 (26.9%) weakly virulent, and 8 (5%) very weakly virulent; 2 (1.25%) isolates were

non-pathogenic. No symptoms appeared on non-wounded banana leaves.

These 160 isolates included six *Nigrospora* species: *N. chinensiss*, *N. lacticolonia*, *N.* cf.

singularis, *N. sphaeric*a, *N. vesicularifera*, and *N. nigrocolonia*. Among them, *N. chinensis*, *N.*

sphaerica, and *N. vesicularifera* were each represented by a single isolate, while multiple

isolates were available for *N. lacticolonia*, *N.* cf. *singularis*, and *N. nigrocolonia*. The single

isolates of *N. chinensis* exhibited weak virulence. *N. sphaerica*, and *N. vesicularifera* exhibited

strong virulence. In contrast, isolates of *N. lacticolonia*, *N.* cf. *singularis*, and *N. nigrocolonia*

showed variable levels of aggressiveness. Among the *N. lacticolonia* isolates, 39.8% (47/118

isolates) were classified as strongly virulent, 37.3% (44/118 isolates) as moderately virulent,

18.6 (22/118 isolates) as weakly virulent, 2.5% (3/118 isolates) as very weakly virulent, and

1.7% (2/118 isolates) as non-pathogenic. For *N.* cf. *singularis*, 16.7% (5/30 isolates) were

classified as strongly virulent, 20% (6/30 isolates) as moderately virulent, 50% (15/30 isolates)

as weakly virulent, and 13.3% (4/30 isolates) as very weakly virulent. For the isolates of *N.*

nigrocolonia, 11.1% (1/9 isolates) were classified as strongly virulent, 22.2% (2/9 isolates) as

moderately virulent, 55.5% (5/9 isolates) as weakly virulent, and 11.1% (1/9 isolates) as very

weakly virulent.

In vitro fungicide sensitivity test

The heat map in Fig. 11 presents the sensitivity of each isolate of *Nigrospora*

lacticolonia (118 isolates), *N. singularis* (30 isolates), *N. nigrocolonia* (9 isolates), *N. chinensis*,

N. sphaerica, and *N. vesicularifera* (1 isolate each) to each fungicide (active ingredient), using

the rate of inhibition of mycelial growth as an indicator (Supplemental Table 1).

All the isolates exhibited high sensitivity (100% inhibition rate) to pyrimethanil (AP) and

tebuconazole (DMI). Against pyraclostrobin (QoI), one isolate of *N. lacticolonia* and one isolate

of *N. nigrospora* showed less than 100% sensitivity (72.7% and 92.3%, respectively). Twenty-

one isolates were less than 100% (78.7-96.7%) sensitive to chlorothalonil, and 30 isolates less

than 100% (74.4-88.5%) sensitive to chlorothalonil. Both fungicides resulted in 100% inhibition

of the remaining isolates. Seventy-seven isolates of *N. lacticolonia* exhibited less than 100%

(70.1-86.6%) sensitivity to boscalid (SDHI). One isolate (76.9%) of *N. singularis* and one

isolate (84.7%) of *N. nigrocolonia* exhibited less than 100% sensitivity. Sixty-nine isolates of *N.*

lacticolonia exhibited a range of 79.9–98% sensitivity to fluxapyroxad. One isolate (80.6%) of

N. singularis and two (83.2-85.8%) of *N. nigrocolonia* had less than 100% sensitivity. Twenty-

two isolates of *N. lacticolonia* displayed sensitivities ranging from 88.8–98.1% to fluopyram.

One isolate (67.9%) of *N. nigrocolonia* exhibited less than 100% sensitivity. All isolates were

highly sensitive to the GMI agents bitertanol, difenoconazole, and epoxiconazole, except for

one to two isolates of *N. nigrocolonia* (72-94.8% inhibition rate). With the exception of nine

isolates of *N. singularis* and one isolate each of *N. sphaerica* and *N. vesicularifera*, the

inhibition rates of fenpropimorph (SBI class 2) were less than 100% (16.9-90.7%). In contrast,

all fungi were highly sensitive (100%) to spiroxamine, which belongs to the same class as

fenpropimorph.

The dominant species, *N. lacticolonia*, exhibited low sensitivity to 8 of the 14 fungicides.

The second most dominant species, *N. singularis*, presented low sensitivity to the four species.

N. nigrocolonia exhibited low sensitivity to nine fungicides, while *N. chinensis* and *N.*

vesicularifera exhibited low sensitivity to one.

Principal component analysis (PCA) for fungicide sensitivety

PCA was performed based on inhibition rates (Supplemental Table 1). In the final PCA

results, data for TAP24N093, TAP24N094, TAP24N163, and TAP24N252, which appeared as

outliers, were excluded (Supplementary Figure 6). These outliers were determined based on

principal component scores obtained from PCA, using a threshold of three standard deviations

from the mean for PC1 and PC2. This criterion is grounded in the statistical property of the

normal distribution, where approximately 99.7% of data points fall within ±3 standard

deviations. PC1 and PC2 together explained 37.99% of the total variance in the dataset

(Supplementary Figure 7; Supplementary Table 2). For PC1, the top three contributors were

Chlorothalonil (22.7%), Mancozeb (21.6%), and Fluopyram (16.5%). For PC2, the top three

contributors included Flutriafol (53.8%), Fluxapyroxad (18.8%), and Difenoconazole (14.8%).

Cluster analysis, based on the distances between the coordinates in the two-dimensional scatter

plot of PC1 and PC2, revealed that no distinct clusters were formed within the specific regions

or fungal lineages (Fig. 12). In cases where PCA was performed including the four strains

identified as outliers, no clustering by geographic region or haplotype was observed, consistent

with the results obtained after excluding the outliers (Supplementary Figure 8).

Discussion

This study aimed to provide essential information for managing leaf spot diseases in the

Philippines. To this end, we identified pathogens responsible for banana leaf diseases in

Mindanao, the main banana-producing region in the Philippines, and evaluated the efficacy of

14 fungicides currently in use against these pathogens.

On Mindanao Island, the disease "black Sigatoka," caused by *Pseudocercospora fijiensis*, was

suspected but not detected in any of the sampled locations. Eighteen genera of fungi from

symptomatic leaves were collected from six farms in Mindanao, and the most prevalent genus

was *Nigrospora* spp., accounting for 72% of the total fungal isolates.

The isolated isolates were identified as six species: *N. chinensis*, *N. lacticolonia*, *N.*

nigrocolonia, *N. singularis*, *N. sphaerica*, and *N. vesicularifera*, all of which are pathogens that

infect banana leaves. To the best of our knowledge, this is the first report of *N. chinensis*, *N.*

lacticolonia, *N. nigrocolonia*, *N. singularis*, and *N. vesicularifera* as causative agents of banana

leaf spot worldwide. *N. sphaerica* has previously been reported as the causal agent of banana

"leaf spot" in Bangladesh[7], but this is the first report of *N. sphaerica* in the Philippines.

In the Philippines, chemical spraying is a common agricultural practice employed 50–70

times per year to manage the local incidence of Sigatoka disease. Frequent use of fungicides

may have altered the leaf biota community and created a selective pressure that allowed for the

persistence of resistant species or strains. However, this cross-sectional study cannot establish

causality. *Pseudocercosporaceae* fungi are inherently slow-growing; therefore, during this study,

they may have been overgrown by other co-infecting fungi and remained undetected. Therefore,

the presence of *P. fijiensis* at the survey sites is not discussed in this study. Nevertheless, it

should be hypothesized that *P. fijiensis* may have been subjected to selection pressure from

fungicide application, potentially contributing to the emergence of the genus *Nigrospora* as the

dominant pathogen at the sampling sites. Future verification through pathogen-specific PCR

detection on infected leaves is considered necessary.

An example of consequence fungicide application contributing to changes in the fungal

composition of pathogenic communities has been reported in wheat [15]. This raises concerns

regarding the selection of fungicides, as they may lead to a shift in the pathogenic fungi present

on banana leaves. To ensure the continued efficacy of fungicide applications in banana

cultivation, conducting regular surveys on disease incidence and fungicide sensitivity,

specifically targeting *Nigrospora* and *Pseudocercospora* species, is essential.

Among the 160 isolates of the six species of the genus *Nigrospora*, *N. lacticolonia* accounted

for 74% of the total (118 isolates). This species was identified as the dominant species at four

sites: Little Panay, Panambong, Tugbok, and Ula. Conversely, *N. singularis* was dominant in

Patag-Alanib and Dabong-dabong. The isolation frequencies of *N. nigrocolonia*, *N. sphaerica*,

and *N. vesicularifera* were comparatively lower than those of the two species, and they are not

currently considered primary pathogens of banana leaf spot.

Haplotype analysis revealed that *N. lacticolonia* Hap_1 and Hap_2 were detected in all six

regions and five regions, respectively, while *N. singularis* Hap_1 was detected in five regions.

The observation that the same haplotype was detected in multiple regions suggests that

genetically similar populations are widely distributed across Mindanao Island, indicating past

population migration and diffusion.

Furthermore, a comprehensive analysis of the sensitivity to 14 chemical substances using

principal component analysis (PCA) revealed that the distribution of PCA on the graph did not

reflect the classification of fungal species or haplotypes. Isolates exhibiting different

sensitivities to 11 chemicals (excluding tebuconazole, spiroxamine, and pyrimethanil) were

found within the species and haplotypes. Notably, the distribution of the isolates did not align

with the classification of species or haplotypes. This indicates that fungicide sensitivity among

the isolates may have evolved independently of species or haplotype differentiation. This

hypothesis is further supported by the findings of Hartmann et al. [16], who demonstrated that

mutations in CYP51, linked to azole fungicide resistance, and Cytb, associated with strobilurine

(QoI) resistance, emerged independently in distinct geographical regions. Therefore, this

observation suggests that the fungicide sensitivity in the isolates arose independently of species

and haplotype differentiation. However, because this study did not include a genetic analysis of

fungicide resistance, the ensuing discussion focuses on the potential mechanisms underlying the

acquisition of fungicide resistance. Hartmann et al. [16] noted that gene flow resulting from

human migration and spore dispersal contributes to the ectopic occurrence of resistant isolates.

The dispersal of diseases caused by *Nigropsora* spp. on Mindanao Island is not only attributable

to human migration resulting from the movement of soil and seedlings but also to soil runoff

caused by flooding and atmospheric migration. The potential contribution of soil runoff is

supported by the detection of this genus in both soil [17] and atmospheric environments [15].

Consequently, it can be posited that fungi with acquired resistance have a propensity for

dissemination.

Therefore, elucidating the mechanisms underlying the dissemination of pathogenic fungi is

imperative for effective disease control. This includes controlling resistant isolates and

determining the optimal timing for disinfectant application. Meredith [18] demonstrated that the

spores of *Nigrospora* spp. are released into the atmosphere in a diurnal cycle. Specifically, in

Jamaica, spore release commenced at approximately 7:00 a.m., reaching a peak between 8:00

and 10:00 a.m. This diurnal pattern suggests that *Nigrospora* spp. employ a release mechanism

that functions under decreasing vapor pressure [19], which is crucial information for disease

control. However, further research is necessary to investigate the environmental factors

associated with disease outbreaks in banana plantations in the Philippines.

The findings of this study regarding pathogen sensitivity to fungicides are fundamental data.

However, as this is an urgent issue, it is important to note that all 118 isolates of *N. lacticolonia*,

the most frequently isolated fungus species, exhibited moderate or low sensitivity to

fenpropimorph (inhibition rate 16.9–90.7%). The fungicide is unlikely to be effective because

the low in vitro sensitivity suggests that its inhibitory capabilities in the field will likely remain

low. In contrast, all fungi showed high sensitivity to spiroxamine, a fungicide of the same class

(SBI classification II group), with 100% inhibition of mycelial growth. Notably, the use of these

fungicides raises concerns regarding potential cross-resistance, and it is possible that fungi

resistant to spiroxamine may emerge in the future. Based on these results, we propose that the

emergence of cross-resistant fungi should be monitored.

In this study, we showed that banana leaf diseases occurring in Mindanao are not black

Sigatoka caused by *P. fijiensis* but are mainly leaf blight caused by six species belonging to the

genus *Nigrospora*. As important information for disease control, it was considered that fungal

senescence may occur due to fungicide treatments, as some of these are sensitive to the 14

fungicides used for black Sigatoka disease, while others are not. It is recommended that further

surveys of fungal diseases throughout Mindanao Island be conducted and that fungicide

selection for disease control be based on a comprehensive understanding of banana leaf diseases.

Materials and methods

Isolation of fungi

In July 2024, 175 of banana leaves (27 to 30 leaves from each banana plantation) with

Sigatoka-like symptoms were collected at 6 Cavendish banana plantations located in "Barangay

Dabong-dabong, Valencia City, Bukidnon (7°58′28″N 125°4′55″E; altitude: 313 m)" (29 leaves),

"Barangay Little Panay, Panabo, Davao del Norte (7°17′44″N 125°38′46″E; altitude: 17 m)" (30

leaves), "Barangay Panombon, Mati, Davao Oriental (7°0′32″N 126°16′28″E; altitude: 20 m)"

(30 leaves), "Barangay Patag-Alanib, Lantapan, Bukidnon (8°1′5″N 125°1′57″E; altitude: 851

m)" (27 leaves), "Barangay Tugbok, Davao City, Davao del Sur (7°11′45″N 125°27′23″E;

altitude: 217 m)" (29 leaves), and "Barangay Ula, Davao City, Davao del Sur (7°8′16″N

125°29′37″E; altitude: 291 m) (30 leaves)" in Mindanao Island, Philippines (Fig. 13). Banana

plants selected for sampling were estimated to be between five and eight months of age.

Symptoms included small spots with gray centers and yellow halos, as well as spots that turned

brown or dark brown as they expanded. The leaves also showed signs of coalescing lesions as

the spots expanded. Fungi were isolated from the symptoms, including black and brown spots

and yellow hallows (Fig. 7). Plant tissues with apparent symptoms were cut into 3-mm square

segments, sterilized with 0.6% (v/v) sodium hypochlorite for 1 min, rinsed twice with sterilized

water, and placed on water agar plates. To obtain monocultures of fungal isolates, a single hypha

was picked directly from the mycelium that emerged from the tissues on water agar. Colonies

with different appearances in a single sample were individually isolated, and then transplanted

onto potato dextrose agar (PDA) plates. The monoculture isolates were deposited in the culture

collection at Tamagawa University and maintained on PDA plates.

Polymerase chain reaction (PCR)

Template DNA for PCR was obtained from the mycelia of each isolate grown on potato

dextrose agar for 7 d using a CTAB method [20]. PCR was performed using a 10 μL PCR

mixture containing 7 μL distilled water, 1 μL 10× Ex Taq buffer with $MgCl_2$, 0.8 μL dNTP (10

mM each), 0.1 μL of each primer (50 μM), 0.05 μL Ex Taq DNA polymerase (5 U $μL^{-1}$, Takara,

Tokyo, Japan), and 1.0 μL of DNA template. For DNA sequencing of all 160 isolates, the partial

18S rRNA gene, ITS1, 5.8S rRNA gene, ITS2, and partial 28S rRNA gene internal transcribed

spacer (ITS) regions were amplified using primer pairs ITS4 and ITS5 [21]. For 107 isolates of

Nigrospora fungi, in addition to the ITS region, two DNA regions, partial sequences of the *β-*

tubulin and the *transcribed elongation factor 1 alfa* (*tef1*) genes, were amplified by using primer

pairs Bt2a and Bt2b [22] and Nigro_EF1 (5'-AAC ATC GTC GTT ATC GGC CA-3') and

Nigro_EF2 (5'-TGA TCT CCT TGT AAC GGG CC-3') (designed in this study), respectively.

As the commonly used primers EF1 and EF2 failed to amplify the tef1 region from the isolates

obtained in this study, Nigro_EF1 and Nigro_EF2 was newly designed in this study

(Supplementary Figure 9).

The PCR was performed under the conditions listed in Table 3. PCR reaction primers were

used for sequencing. PCR products were sequenced by the FASMAC Co. Ltd. in Japan

(https://fasmac.co.jp/). Obtained forward and reverse sequence reads were assembled using

BioEdit v7.7.1.0 software [23] (https://thalljiscience.github.io/page2.html).

Molecular phylogenetic analysis

To identify the 217 isolates at the genus level, we used ITS region sequences to conduct a

BLAST search in the NCBI database. Based on the results of the BLAST search, we selected

160 isolates belonging to the genus *Nigrospora* for further analyses to determine the dominant

species and diversity of pathogens in the genus *Nigrospora*. To construct phylogenetic trees, 160

isolates and 36 strains of *Nigrospora* spp., including ex-type strains, were used as ingroup taxa

(Table 4). As an out group, *Apiospora malaysiana* and *A. pseudoparenchymatica* were selected

from the genus *Apiospora*, which is a sister group to the genus *Nigrospora*. All of the ITS, *β-*

tubulin, and *tef1* sequences were used for multiple alignments by ClustalW with BioEdit [23]

and concatenated into one dataset (Supplementary data 1). Bayesian Inferences (BI) with

MrBayes 3.1.2 [24] (https://nbisweden.github.io/MrBayes/index.html) implementing the

Markov chain Monto Carlo technique were performed based on the combined dataset of ITS, *β-*

tubulin, and *tef1α*. The best-fitted model for each locus for BI was selected with

MrMODELTEST2.3 [25]. Each locus was treated as a separate partition. The parameters in

MrBayes were set as follows: lset nst = 6, 2, and 6 for ITS, β-tubulin, and *tef1α*, respectively;

rates = invgamma, prset statefreqpr = dirichlet (1,1,1,1), nchains = 4, temp = 0.2, swapfreq = 1,

and nswaps = 1. The Markov chains were continued until average standard deviation of split

frequencies fell below 0.0095 (7,305,000 generations). Only trees sampled after the average

standard deviation of split frequencies dropped below 0.01 were used for consensus tree

construction. Accordingly, the first 7,244,000 generations were discarded as burn-in. The

remaining trees were used to construct a 50% majority-rule consensus tree. Branches that

received Bayesian posterior probabilities (BPP) greater than or equal to 0.95 were considered as

significantly supported. Bayesian analysis was also performed for each locus with the model

described above, and using same parameters. For BI based on ITS and *β-tubulin* sequences, the

Markov chains were continued until average standard deviation of split frequencies fell below

0.0095 (27,815,000 and 1,835,000 generations, respectively). Only trees sampled after the

average standard deviation of split frequencies dropped below 0.01 were used for consensus tree

construction. Accordingly, the first 28,761,000 for ITS and 1,760,000 for *β-tubulin* generations

were discarded as burn-in. For BI based on *tef1α*, despite running the analysis for 40,000,000

generations, the average standard deviation of split frequencies did not fall below 0.01.

Therefore, we constructed the phylogenetic tree using 1,000 generations of data starting from

generation 28,761,000, which yielded the lowest average standard deviation of split frequencies

value (0.016862).

Phylogenetic analyses with maximum likelihood method were performed using Molecular

Evolutionary Genetics Analysis (MEGA10) software (Kumar 2018) [26]. Tamura-Nei model

was selected as the best substitution model based on the Akaike information criterion using

MEGA 10. All sites with gaps were treated as missing data. The reliability of each internal

branch in the phylogenetic trees was assessed using bootstrap analysis with 1,000 random-

addition replicates [27].

Morphological observation

Our isolates were cultured on synthetic low-nutrient agar (SNA) [28] at 28□ for 10 days to

induce sporulation in the dark. The shape and size of the microscopic structures were observed

using light microscopy (B51, Olympus, Tokyo, Japan), and the colonies were assessed. At least

30 conidiogenous cells and conidia were measured to determine mean colony size. Three SNA

plates were examined, and from each plate, ten or more conidiophores and conidia were

observed for measurement. These measurements were conducted using ImageJ 1.52a software

(Rasband, WS, ImageJ; US National Institutes of Health: https://imagej.net/ij/index.html).

Pathogenicity test

Each of our 160 isolates was tested for pathogenicity using mycelial discs (5 mm diameter)

obtained from the edges of colonies grown on PDA media for three days. These discs were then

applied to four sites on non-wounded and wounded segments of healthy banana leaves (approx.

5×5 cm in size; cv. Dwarf Cavendish). For each fungal strain, two non-wounded and two

wounded leaf segments were used. Three puncture sites were created on the wounded leaves

using a heated sewing needle. Each puncture site was spaced approximately 2 mm apart. The

inoculated leaves were covered with plastic boxes to maintain high humidity (85–90%) with wet

paper towel and maintained at approximately 25 °C under natural light condition for three days.

PDA was used as a control. Pathogenicity was evaluated based on the size and coloration of

lesions observed at 4 days after inoculation (dpi). Isolates that did not produce elliptical necrotic

lesions but caused brown or yellow discoloration at or around the wound site were considered

extremely weak in pathogenicity and were classified as "very weakly virulent". For isolates that

formed brown, spindle-shaped necrotic lesions, the lesion length at each of the eight inoculation

points was measured, and the mean lesion length was used as an index of pathogenicity. To

further categorize the degree of pathogenicity, the 33rd and 66th percentiles of the mean lesion

lengths were calculated. Isolates with mean values below the 33rd percentile were classified as

"weakly virulent", those between the 33rd and 66th percentiles as "moderate virulent", and

those above the 66th percentile as "strongly virulent". This classification was performed using R

(version 4.2.2), with data processing conducted via the dplyr package. Values of zero were

excluded prior to quantile calculation. Based on these criteria, all isolates were ultimately

categorized into five levels of pathogenicity: "non-pathogenic", "very weakly virulent",

"weakly virulent", "moderate virulent", and "strongly virulent". At 7 dpi, the symptoms

appeared, the fungi were re-isolated from the diseased tissue using the same method as that of

the description in "isolation of fungi" and re-identified based on morphologies of colony on

PDA.

Haplotype analyses

The haplotypes of *N. lacticolonia*, *N. singuralis*, and *N. nigrocolonia* were analyzed to

reveal the spread of pathogens on Mindanao Island. The sequence of the *tef1α* gene was used for

the analyses because mutations in each species were observed in the gene. The *tef1α* gene

sequences for each species were aligned using MEGA 10, and sites containing gaps at both ends

were trimmed. The total number of variable sites (S) and the nucleotide diversity (Pi) were

calculated using MEGA 10. The trimmed alignment data were exported to PopArt v1.7 [29] to

construct a haplotype network. The geographic distribution of the haplotypes was visualized

using the median-joining option in the PopArt program.

In vitro Fungicide Sensitivity test

In vitro tests were conducted to evaluate the inhibitory effects of the active ingredients in

fungicides, currently used in the Philippines, on the mycelial growth of *Nigrospora* spp.

Mycelial growth inhibition was used as an indicator of sensitivity. The active ingredients of 14

fungicides belonging to 6 classes of the Fungicide Resistance Action Committee (FRAC) were

dissolved in acetone or sterile distilled water and adjusted to a concentration of 1/5 in PDA

medium. Aryl hydroxamic acid (SHAM; Fluorochem Ltd.) was used in conjunction with the

QoI fungicide pyraclostrobin to inhibit alternative oxidase (AOX), which does not react with

cyano groups [30,31]. The final concentrations of the fungicides in the medium were as follows:

100 ppm for bitertanol, difenoconazole, epoxiconazole, flutriafol, tebuconazole, fenpropimorph,

spiroxamine, boscalid, and flusapirox; and 200 ppm for mancozeb and chlorothalonil. Media

without fungicides (0 ppm) were used as control to formulate inhibition rate. These fungicide

concentrations were determined based on the following reasons. In FRAC monitoring assays,

the upper concentration range typically falls between 1 and 30 ppm; however, concentrations as

high as 100 ppm are occasionally employed. By intentionally setting the concentration higher

than the standard FRAC monitoring range, we aimed to detect isolates with clearly low

sensitivity. Isolates were classified as sensitive or low-sensitive based on their growth response

at 100 ppm or 200 ppm. Those exhibiting visible growth at these concentrations were

designated as having low sensitivity.

Mycelial disks (6 mm in diameter) were obtained from the periphery of pre-cultured

colonies, which were cultured at 25°C for 3 days in 1/5 PDA. These disks were transferred to

1/5 PDA containing the chemicals at avobe mentioned concentrations. Following a 24-h

incubation period at 25°C, the diameters of the colonies were measured, and 6 mm was

subtracted from the diameter of the mycelial disc. The inhibition rate was defined as the rate of

growth when the control was 100%, and was calculated from the values of the four colonies.

Principal component analysis (PCA)

The PCA of standardized variables was performed using R (v4.2.2) to determine whether

there was a difference in the inhibition rate among species/haplotypes and sampling sites. We

conducted PCA using the inhibition rate data obtained from the 14 fungicides as variants. The

final principal components (PCs) were determined by excluding the data of individuals that fell

outside the ±3σ range as outliers. Euclidean distances were calculated based on the obtained

coordinates of PC1 and PC2, and a dendrogram was constructed using Ward's method. Outliers

were defined based on principal component scores obtained from PCA, using a threshold of

three standard deviations from the mean for PC1 and PC2. This criterion is grounded in the

statistical property of the normal distribution, where approximately 99.7% of data points fall

within ±3 standard deviations. Values exceeding this range are considered statistically rare and

thus indicative of potential outliers. A sample was classified as an outlier if either its PC1 or

PC2 score exceeded the threshold.

Acknowledgments

We are grateful to Ms Sayuri Sakurai (Tamagawa University BacaDM Project) for her

assistance in preparing the culture medium for this research.

Data Availability

The datasets generated during the current study are available in the GenBank provided by

NCBI/DDBJ. The GenBank accession numbers are listed in Table 4.

Author Contribution

S.N. conceived the study, designed the methodology, collected field samples, conducted

fungicide sensitivity tests, conducted all analyses, visualization of the data, and led the

manuscript preparation. Y.H. and Y.T. conducted fungicide sensitivity tests. K.U. designed the

methodology for fungicide sensitivity tests. M.A.M., R.V., S.M.C., and A.F.P. collected field

samples. K.W. supervised this study, acquired funding, conducted fungicide sensitivity tests, and

assisted in manuscript editing and project coordination. All authors reviewed and approved the

final manuscript

Competing Interests Statement

The authors declare no competing interests.

References

1. Nozawa, S. et al. *Fusarium mindanaoense* sp. nov., a new fusarium wilt pathogen of

cavendish banana from the Philippines belonging to the F. fujikuroi species complex. *J.*

Fungi (Basel) **9**, 443; 10.3390/jof9040443, PubMed: 37108898 (2023).

2. Ploetz, R. C. Fusarium wilt of banana. *Phytopathology* **105**, 1512–1521; 10.1094/PHYTO-

04-15-0101-RVW, PubMed: 26057187 (2015).

3. Wang, B. et al. A new leaf blight disease caused by *Alternaria jacinthicola* on banana in

China. *Horticulturae* **8**, 12 (2021).

4. Fu, B. Z. et al. First report of leaf spot caused by *Alternaria alternata* on Chinese dwarf

banana in China. *Plant Dis.* **98**, 691-691; 10.1094/PDIS-08-13-0831-PDN, PubMed:

30708541 (2014).

5. Huang, R. et al. Identification and characterization of Colletotrichum species associated

with anthracnose disease of banana. *Plant Pathol.* **70**, 1827-1837; 10.1111/ppa.13426

(2021).

6. Crous, P. W. & Mourichon, X. Mycosphaerella eumusae and its anamorph

Pseudocercospora eumusae spp. nov: causal agent of eumusae leaf spot disease of banana.

Sydowia **54**, 23-34 (2002).

7. Sadid, K. N. E., Hossain, M. A., Munshi, A. R. & Karim, M. R. First detection, isolation

and molecular characterization of banana leaf spot diseases caused by *Nigrospora sphaerica*

and *Neocordana musae* in Bangladesh. *Arch. Phytopathol. Plant Prot.* **56**, 1112-1125 (2023).

8. Fortune, M. P. et al. First report of black Sigatoka disease (causal agent *Mycosphaerella*

fijiensis) from Trinidad. *Plant Pathol.* **54**, 246; 10.1111/j.1365-3059.2005.01123.x (2005).

9. Strobl, E. & Mohan, P. Climate and the global spread and impact of bananas' black leaf

sigatoka disease. *Atmosphere* **11**, 947; 10.3390/atmos11090947 (2020).

10. Esguera, J. G., Balendres, M. A. & Paguntalan, D. P. Overview of the Sigatoka leaf spot

complex in banana and its current management. *Trop. Plant* **3**, e002; 10.48130/tp-0024-

0001 (2024).

11. Rhodes, P. L. A new banana disease in Fiji. *CPN* **10**, 38–41 (1964).

12. Pordel, A., Dehghani, K., Amirmijani, A. & Miri, K. Identification and pathogenicity of

Fusarium sensu lato species causeing wilting and leaf spot of Banana in South of Iran. *Iran.*

J. Plant Pathol. **58**, 96-108 (2022).

13. Raza, M. et al. Culturable plant pathogenic fungi associated with sugarcane in southern

China. *Fungal Divers.* **99**, 1–104; 10.1007/s13225-019-00434-5 (2019).

14. Wang, M et al. Phylogenetic reassessment of *Nigrospora*: Ubiquitous endophytes, plant and

human pathogens. *Persoonia* **39**, 118–142; 10.3767/persoonia.2017.39.06 (2017).

15. Karlsson, I., Friberg, H., Steinberg, C. & Persson, P. Fungicide effects on fungal community

composition in the wheat phyllosphere. *PLOS One* **9**, e111786;

10.1371/journal.pone.0111786, PubMed: 25369054 (2014).

16. Hartmann, F. E. et al. The complex genomic basis of rapid convergent adaptation to

pesticides across continents in a fungal plant pathogen. *Mol. Ecol.* **30**, 5390-5405;

10.1111/mec.15737, PubMed: 33211369 (2021).

17. Zhang, Y. Y. et al. *Nigrospora humicola* (Apiosporaceae, Amphisphaeriales), a new fungus

from soil in China. *Diversity* **16**, 118 (2024).

18. Meredith, D. S. Atmospheric content of nigrospora spores in Jamaican banana plantations. *J.*

Gen. Microbiol. **26**, 343-349; 10.1099/00221287-26-2-343, PubMed: 14472772 (1961).

19. Hirst, J. M. Changes in atmospheric spore content: diurnal periodicity and the effects of

weather. *Trans Br Mycol Soc.* **36**, 375–IN8; 10.1016/S0007-1536(53)80034-3 (1953).

20. Doyle, J. J. & Doyle, J. L. A rapid DNA isolation procedure for small quantities of fresh leaf

tissue. *Phytochem Bull.* 19A1 **A27**, 11–15 (1987).

21. White, T. J., Bruns, T., Lee, S. & Taylor, J. W. Amplification and direct sequencing of fungal

ribosomal RNA genes for phylogenetics. in 15 (ed. Gelfand, D. H., Shinsky, J. J. & White, T.

J.) PCR Protocols: A Guide to Methods and Applications Ch. 315–322 (M.A., Innis) (Acad.

Press (San Diego, 1990)).

22. Glass, N. L. & Donaldson, G. C. Development of primer sets designed for use with the PCR

to amplify conserved genes from filamentous Ascomycetes. *Appl Environ Microbiol.* **61**,

1323-1330; 10.1128/aem.61.4.1323-1330.1995, PubMed: 7747954 (1995).

23. Hall, T. A. BioEdit: a user-friendly biological sequence alignment editor and analysis

program. for Windows 95/98/NT. *Nucleic Acids Symp Ser.* 41 95–98, (1999).

24. Ronquist, F. & Huelsenbeck, J. P. MrBayes 3: Bayesian phylogenetic inference under mixed

models. *Bioinformatics* **19**, 1572–1574 (2003).

25. Nylander, J.A. Modeltest v2. Program distributed by the author. Evolutionary Biology

Centre, Uppsala University (2004).

26. Kumar, S., Stecher, G., Li, M., Knyaz, C., Tamura, K. et al. MEGA X: Molecular

evolutionary genetics analysis across computing platforms. *Mol. Biol. Evol.* **35**, 1547–1549;

10.1093/molbev/msy096, PubMed: 29722887 (2018).

27. Felsenstein, J. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*

39, 783–791; 10.1111/j.1558-5646.1985.tb00420.x, PubMed: 28561359 (1985).

28. Nirenberg, H. I. A simplified method for identifying Fusarium spp. occurring on wheat. *Can.*

J. Bot. **59**, 1599–1609 (1981).

29. Leigh, J. W. & Bryant, D. POPART: full-feature software for haplotype networkconstruction.

Methods Ecol. Evol. **6**, 1110-1116; 10.1111/2041-210X.12410 (2015).

30. Ishii, H. et al. Characterisation of QoI□resistant field isolates of Botrytis cinerea from citrus

and strawberry. *Pest Manag. Sci.* **65**, 916-922; 10.1002/ps.1773, PubMed: 19444805 (2009).

31. Hollomon, D. W., Wood, P. M., Reeve, C. & Miguez, M. Alternative oxidase and its impact

on the activity of Qo and Qi site inhibitors. in *Modern Fungicides and Antifungal*

Compounds IV (ed. Dehne, H. W., Gisi, U., Kuck, K. H., Russell, P. E. & Lyr, H.) 31–34

(BCPC, Alton, Hants, UK, 2005).

Fig. 1   Symptoms of leaf spot on banana leaves in fields. Naturally infected leaves in
banana plantations (a, b).

Fig. 2 Phylogenetic trees constructed based on the concatenated data of the DNA sequences
of ITS, *β-tubulin*, and *tef1α* genes inferred using MrBayes. HKY, GTR, and HKY was used as
models in MrBayes for ITS, *β-tubulin*, and *tef1α*, respectively. The numbers on the left and right
of each node are posterior probabilities estimated using the software MrBayes and bootstrap
values from ML method performed with MEGA 10 with Tamura-Nei model, respectively.
Nodes with a PPB of 1.0 are highlighted with green circles, while those with PPB ≥ 0.95 are
indicated by orange circles. Strain number with asterisk is ex-type strain. Species name and
strains with boldface is our isolate. Branch lengths are proportional to the number of
substitutions per site; the scale bar indicates expected substitutions per site.

Fig. 2   Continued.

Fig. 4 *Nigrospora nigrocolonia* (**TAP24N237**). Surface and reverse of colony on PDA after
3 days (a, b) and 7 days (c, d). Conidiogenous cells and conidia on SNA plate (e, f, g). Scale
bars: = 20 μm

Fig. 5 *Nigrospora chinensis* **(TAP24N249).** Surface and reverse of colony on PDA after 3
days (a, b). Conidiogenous cells and conidia on SNA plate (c, d). Scale bars: = 20 μm

738 **Fig. 6**   *Nigrospora lacticolonia* **(TAP24N011).** Surface and reverse of colony on PDA after 3
739 days (a, b). Conidiogenous cells and conidia on SNA plate (c, d). Scale bars: = 20 μm
740

743 **Fig. 7** *Nigrospora singuralis* **(TAP24N068).** Surface and reverse of colony on PDA after 3
744 days (a, b). Conidiogenous cells and conidia on SNA plate (c, d). Scale bars: = 20 μm
745

747 **Fig. 8** *Nigrospora sphaerica* **(TAP24N107).** Surface and reverse of colony on PDA after 3
748 days (a, b). Conidiogenous cells and conidia on SNA plate (c, d). Scale bars: = 20 μm
749

751   **Fig. 9**   *Nigrospora vesicularifera* **(TAP24N023).** Surface and reverse of colony on PDA after
752   3 days (a, b).

Fig. 10 Samples of banana leaves 4 days after inoculation with mycelial plugs. Inoculated leaves with *N. chinensis* (isolate TAP24N011) with wounds (a1, a2) and without wound (a3, a4), *N. lacticolonia* (isolate TAP24N078) with wounds (b1, b2) and without wound (b3, b4), *N. nigrocolonia* (isolate TAP24N237) with wounds (c1, c2) and without wound (c3, c4), *N.* cf. *singularis* (isolate TAP24N257) with wounds (d1, d2) and without wound (d3, d4), *N. sphaerica* (isolate TAP24N107) with wounds (e1, e2) and without wound (e3, e4), *N.vesicularifera* (isolate TAP24N023) with wounds (f1, f2) and without wound (f3, f4). Controls inoculated with potato dextrose agar discs with wounds (g1, g2) and without wound (g3, g4). Bars: 1 cm.

764   **Fig. 11   Heat map of the inhibition percentages for the isolates.**

767 **Fig. 12 Dendrogram based on coordinates on scatter plot of PCA derived from values of**
768 **inhibition rates of fungicide.** Label on the left and right of hyphen are species_haplotype and
769 sampling site, respectively.
770

Fig. 12   Continued.

773   **Fig. 13   Sampling sites in the Mindanao island, Philippines.**

Table 1 Preliminary identification of 217 fungal isolates from banana leaf samples from six cities of Mindanao Island.

Genus[a]	Isolate numbers from different cities of Mindanao Island					
	Dabong-dabong	Little Panay	Panambon	Patag-Alanib	Tugbok	Ula
Alternaria	0	0	0	0	1	0
Annulohypoxylon	0	0	0	1	0	0
Colletotrichum	0	0	0	0	2	3
Curvularia	0	0	1	0	1	0
Dimorphiseta	0	0	1	0	0	0
Exserohilum	0	1	0	0	0	0
Fusarium	0	3	6	0	0	4
Lasiodiplodia	1	0	0	0	1	0
Microdochium	0	0	3	0	0	0
Musicillium	0	0	0	0	0	1
Mycosphaerella	0	1	0	0	0	0
Neosetophoma	0	2	3	0	0	0
Nigrospora	28	33	21	25	20	33
Ochrocladosporium	0	1	0	0	0	0
Phaeosphaeria	0	5	6	0	3	2
Preussia	0	0	0	1	0	0
Pseudopestalotiopsis	0	1	0	0	1	0
Zasmidium	0	0	0	0	0	1

[a]The fungal isolates were identified based on ITS sequences.

Table 2 Lesion length and virulence of isolates based on pathogenicity tests.

Species	Isolate	Lesion length (mm)	Virulence
N. chinensis	TAP24N249	0-3.39 (mean±sd = 0.42±0.42)	Weak
N. lacticolonia	TAP24N001	6.62-12.27 (mean±sd = 9.52±0.81)	Strong
	TAP24N003	0-12.7 (mean±sd = 8.25±1.33)	Strong
	TAP24N005	6.58-11.04 (mean±sd = 8.6±0.55)	Strong
	TAP24N006	5.17-10.99 (mean±sd = 8.67±0.67)	Strong
	TAP24N008	8.88-11.37 (mean±sd = 10.28±0.29)	Strong
	TAP24N010	5.93-9.59 (mean±sd = 8.09±0.48)	Strong
	TAP24N011	8.87-10.35 (mean±sd = 9.38±0.17)	Strong
	TAP24N012	7.94-11.95 (mean±sd = 10.12±0.49)	Strong
	TAP24N013	0-16.05 (mean±sd = 10.5±2.37)	Strong
	TAP24N014	0-6.81 (mean±sd = 2.94±0.94)	Weak
	TAP24N016	4.44-8.56 (mean±sd = 6.8±0.58)	Strong
	TAP24N017	2.87-9.1 (mean±sd = 5.49±0.75)	Moderate
	TAP24N018	8.18-14.24 (mean±sd = 11.14±0.74)	Strong
	TAP24N019	0-6.72 (mean±sd = 2.96±1)	Weak
	TAP24N021	2.33-9.37 (mean±sd = 5.8±0.82)	Moderate
	TAP24N022	0-10.48 (mean±sd = 5.97±1.2)	Moderate
	TAP24N026	4.12-11.74 (mean±sd = 7.44±0.75)	Strong
	TAP24N027	3.61-6.65 (mean±sd = 5.03±0.41)	Moderate
	TAP24N028	5.64-12.31 (mean±sd = 8.11±0.79)	Strong
	TAP24N029	0-5.88 (mean±sd = 4.43±0.69)	Moderate
	TAP24N030	5.1-9.27 (mean±sd = 6.66±0.52)	Strong
	TAP24N031	4.91-9.32 (mean±sd = 7.61±0.52)	Strong
	TAP24N033	3.36-13.2 (mean±sd = 7.9±1)	Strong
	TAP24N035	4.3-11.41 (mean±sd = 9.17±0.84)	Strong
	TAP24N036	9.39-13.93 (mean±sd = 11.7±0.5)	Strong
	TAP24N038	2.9-5.39 (mean±sd = 3.83±0.28)	Moderate
	TAP24N039	0-8.2 (mean±sd = 2.98±1.26)	Weak
	TAP24N041	0-4.51 (mean±sd = 0.94±0.63)	Weak
	TAP24N043	0-8.45 (mean±sd = 2.97±1.19)	Weak
	TAP24N045	0.11-7.3 (mean±sd = 4.28±0.8)	Moderate
	TAP24N047	0-9.55 (mean±sd = 4.89±1.06)	Moderate

TAP24N048	0-7.82 (mean±sd = 4.08±1.01)	Moderate
TAP24N050	0-4.35 (mean±sd = 1.38±0.68)	Weak
TAP24N053	3.31-9.21 (mean±sd = 5.81±0.64)	Moderate
TAP24N054	0-9.22 (mean±sd = 6.04±1.36)	Moderate
TAP24N055	0-7.07 (mean±sd = 2.23±1.12)	Weak
TAP24N057	0-6.09 (mean±sd = 2.67±1.03)	Weak
TAP24N060	0-10.81 (mean±sd = 7.43±1.24)	Strong
TAP24N062	4.89-8.56 (mean±sd = 6.74±0.48)	Strong
TAP24N063	0-10.13 (mean±sd = 4.09±1)	Moderate
TAP24N065	3.1-11.57 (mean±sd = 6.78±1.13)	Strong
TAP24N067	4.56-16.65 (mean±sd = 11.04±1.35)	Strong
TAP24N069	0-8.13 (mean±sd = 5.4±1.19)	Moderate
TAP24N071	3.13-6.88 (mean±sd = 4.92±0.41)	Moderate
TAP24N073	0-10.55 (mean±sd = 3.83±1.54)	Moderate
TAP24N074	0-5.44 (mean±sd = 2.26±0.9)	Weak
TAP24N077	4.17-10.38 (mean±sd = 7.12±0.61)	Strong
TAP24N078	5.19-20.07 (mean±sd = 12.31±1.54)	Strong
TAP24N079	0	Very weak
TAP24N082	6.81-11.43 (mean±sd = 8.89±0.62)	Strong
TAP24N085	2.69-11.25 (mean±sd = 5.93±1.13)	Moderate
TAP24N086	8.2-14.77 (mean±sd = 10.64±0.98)	Strong
TAP24N088	3.95-9.66 (mean±sd = 7.13±0.58)	Strong
TAP24N090	3.56-4.72 (mean±sd = 4.17±0.17)	Moderate
TAP24N091	0-5.5 (mean±sd = 1.51±0.79)	Weak
TAP24N092	0-9.04 (mean±sd = 5.44±1.03)	Moderate
TAP24N093	2.87-8.09 (mean±sd = 5.62±0.63)	Moderate
TAP24N094	5.19-14.99 (mean±sd = 8.22±1.09)	Strong
TAP24N095	5.88-11.22 (mean±sd = 7.54±0.66)	Strong
TAP24N096	0-7.13 (mean±sd = 2.99±0.97)	Weak
TAP24N098	5.64-15.34 (mean±sd = 9.94±1.42)	Strong
TAP24N099	4.44-9.75 (mean±sd = 7.05±0.75)	Strong
TAP24N100	4.7-9.72 (mean±sd = 6.76±0.63)	Strong
TAP24N101	4.47-11.81 (mean±sd = 7.85±0.83)	Strong
TAP24N102	3.56-11.43 (mean±sd = 7.38±1.25)	Strong
TAP24N103	5.3-11.07 (mean±sd = 7.87±0.71)	Strong

TAP24N104	0-10.43 (mean±sd = 7.55±1.17)	Strong
TAP24N105	0	non-pathogenic
TAP24N108	0-7.4 (mean±sd = 1.97±1.03)	Weak
TAP24N112	0-6.78 (mean±sd = 3.24±1.01)	Moderate
TAP24N113	0-5.89 (mean±sd = 3.48±0.84)	Moderate
TAP24N115	0-8.26 (mean±sd = 3.83±1)	Moderate
TAP24N118	0-7.31 (mean±sd = 3.08±1.07)	Moderate
TAP24N123	6.6-14.57 (mean±sd = 9.18±0.94)	Strong
TAP24N125	3.15-7.5 (mean±sd = 5.34±0.59)	Moderate
TAP24N127	2.95-8.03 (mean±sd = 6.15±0.69)	Moderate
TAP24N128	6.44-10.25 (mean±sd = 8.31±0.42)	Strong
TAP24N129	0-2.73 (mean±sd = 0.34±0.34)	Weak
TAP24N131	0-6.06 (mean±sd = 3.33±0.8)	Moderate
TAP24N132	0-5.39 (mean±sd = 2.44±0.67)	Weak
TAP24N134	0-5.47 (mean±sd = 2.75±0.66)	Weak
TAP24N136	0-7.98 (mean±sd = 4.5±0.95)	Moderate
TAP24N139	0-2.43 (mean±sd = 0.3±0.3)	Weak
TAP24N141	0-4.42 (mean±sd = 2.08±0.63)	Weak
TAP24N145	4.05-8.24 (mean±sd = 6.85±0.54)	Strong
TAP24N146	2.75-4.57 (mean±sd = 3.92±0.21)	Moderate
TAP24N149	3.39-6.56 (mean±sd = 4.82±0.36)	Moderate
TAP24N154	5.07-8.36 (mean±sd = 6.29±0.38)	Moderate
TAP24N156	3.72-9.3 (mean±sd = 6.19±0.68)	Moderate
TAP24N157	4.96-9 (mean±sd = 6.36±0.49)	Moderate
TAP24N163	6.19-9.6 (mean±sd = 7.54±0.35)	Strong
TAP24N164	2.72-9.06 (mean±sd = 5.02±0.75)	Moderate
TAP24N165	5.21-12.48 (mean±sd = 8.13±0.84)	Strong
TAP24N166	2.54-6.84 (mean±sd = 4.31±0.44)	Moderate
TAP24N167	2.57-5.46 (mean±sd = 4.12±0.42)	Moderate
TAP24N168	0-5.11 (mean±sd = 1.96±0.79)	Weak
TAP24N174	2.68-7.63 (mean±sd = 4.98±0.59)	Moderate
TAP24N176	4.12-12.73 (mean±sd = 7.87±1.25)	Strong
TAP24N177	5.11-9 (mean±sd = 7.14±0.54)	Strong
TAP24N204	0-5.1 (mean±sd = 3.11±0.71)	Moderate

	TAP24N208	3.11-7.33 (mean±sd = 4.13±0.51)	Moderate
	TAP24N209	0-4.82 (mean±sd = 2.81±0.68)	Weak
	TAP24N226	0-2.56 (mean±sd = 0.32±0.32)	Weak
	TAP24N230	6.43-22.2 (mean±sd = 12.78±1.94)	Strong
	TAP24N231	0	non-pathogenic
	TAP24N233	4.62-10.03 (mean±sd = 8.12±0.73)	Strong
	TAP24N234	0-3.27 (mean±sd = 0.81±0.53)	Weak
	TAP24N235	0	Very weak
	TAP24N236	0-3.26 (mean±sd = 0.41±0.41)	Weak
	TAP24N238	7.77-11.65 (mean±sd = 9.39±0.46)	Strong
	TAP24N239	0-6.58 (mean±sd = 4.23±0.76)	Moderate
	TAP24N240	2.38-7.19 (mean±sd = 4.91±0.57)	Moderate
	TAP24N242	6.09-9.67 (mean±sd = 8.41±0.48)	Strong
	TAP24N243	3.94-9.79 (mean±sd = 6.37±0.69)	Moderate
	TAP24N245	0-10.44 (mean±sd = 4.47±1.21)	Moderate
	TAP24N251	2.88-4.48 (mean±sd = 3.45±0.17)	Moderate
	TAP24N254	0	Very weak
	TAP24N255	4.07-8.51 (mean±sd = 5.82±0.58)	Moderate
N. nigrocolonia	TAP24N202	0-3.2 (mean±sd = 0.79±0.52)	Weak
	TAP24N222	2.84-5.92 (mean±sd = 3.87±0.4)	Moderate
	TAP24N224	0-6.34 (mean±sd = 3.18±0.72)	Moderate
	TAP24N229	0-5.95 (mean±sd = 3.02±0.8)	Weak
	TAP24N232	0	Very weak
	TAP24N237	6.8-10.9 (mean±sd = 8.69±0.61)	Strong
	TAP24N246	0-4.45 (mean±sd = 0.99±0.65)	Weak
	TAP24N250	0-5.67 (mean±sd = 0.71±0.71)	Weak
	TAP24N252	0-0.11 (mean±sd = 0.01±0.01)	Weak
N. cf. *singularis*	TAP24N034	0-5.34 (mean±sd = 0.67±0.67)	Weak
	TAP24N049	0-3.65 (mean±sd = 1.84±0.57)	Weak
	TAP24N068	4.05-12.2 (mean±sd = 6.5±0.9)	Moderate
	TAP24N081	4.84-13.4 (mean±sd = 8.14±1.08)	Strong
	TAP24N120	0-3.64 (mean±sd = 0.46±0.46)	Weak
	TAP24N137	0-2.75 (mean±sd = 0.86±0.43)	Weak
	TAP24N201	0-4.62 (mean±sd = 2.33±0.71)	Weak

	TAP24N203	0-4.33 (mean±sd = 1.86±0.73)	Weak
	TAP24N205	0-4.09 (mean±sd = 1.96±0.59)	Weak
	TAP24N206	0-4.04 (mean±sd = 3.07±0.46)	Weak
	TAP24N207	2.77-14.02 (mean±sd = 5.78±1.36)	Moderate
	TAP24N210	5.98-15.06 (mean±sd = 9.96±1.19)	Strong
	TAP24N211	0	Very weak
	TAP24N212	5.84-9.79 (mean±sd = 7.62±0.43)	Strong
	TAP24N213	0-4.54 (mean±sd = 2.86±0.47)	Weak
	TAP24N214	0-2.11 (mean±sd = 0.26±0.26)	Weak
	TAP24N215	0-4.08 (mean±sd = 2.42±0.53)	Weak
	TAP24N217	0	Very weak
	TAP24N218	3.73-8.67 (mean±sd = 6.11±0.63)	Moderate
	TAP24N219	3.24-6.9 (mean±sd = 5.57±0.45)	Moderate
	TAP24N220	0-3.43 (mean±sd = 1.9±0.53)	Weak
	TAP24N221	0-5.39 (mean±sd = 1.2±0.8)	Weak
	TAP24N223	0-6.24 (mean±sd = 2.41±1)	Weak
	TAP24N227	0	Very weak
	TAP24N241	0-16.05 (mean±sd = 8.12±1.89)	Strong
	TAP24N244	3.36-10.8 (mean±sd = 6.38±0.97)	Moderate
	TAP24N247	0-3.87 (mean±sd = 2.31±0.54)	Weak
	TAP24N248	0-5.58 (mean±sd = 3.1±0.75)	Moderate
	TAP24N256	0	Very weak
	TAP24N257	7.34-24.96 (mean±sd = 16.44±2.11)	Strong
N. sphaerica	TAP24N107	11.61-22.61 (mean±sd = 14.33±1.25)	Strong
N. vesicularifera	TAP24N023	0-10.51 (mean±sd = 7.8±1.25)	Strong

Aggressiveness was classified based on lesions as follows: strongly virulent ($\geq$6.6 mm in mean length), moderately virulent ($\geq$3 to <6.6 mm), weakly virulent (>0 to <3 mm), and very weakly virulent (0 mm with brown or yellow discoloration at or around the wound site).

Table 3 PCR conditions for amplifying ITS, *β-tubulin*, and *tef1α* regions.

Loci	Primer name	Reaction condition	Reference
ITS	ITS4 ITS5	94□, 3m; 94□, 60s, 55□, 60s, 72□, 60s, 35 cycles; 72□, 5m	White et al. 1990

β-tubulin	Bt2a Bt2b	94□, 3m; 94□, 1m, 50□, 1m, 72□, 60s, 35 cycles; 72□, 5m	Glass and Donaldson (1995)
Tef1α	Nigro_EF1 Nigro_EF2	94□, 3m; 94□, 30s, 60□, 30s, 72□, 60s; 30 cycles; 72□, 5m	This study

Table 4 GenBank accession numbers of the DNA sequences used in phylogenetic analysis and pathogenicity of our isolates.

Species	Isolates	Locations	GenBank accession no.		
			ITS	β-tubulin	tef1a
Apiospora malaysiana	CBS 102053	-	KF144896	KF144988	KF145030
A. pseudoparenchymatica	LC7234*	-	KY494743	KY705211	KY705139
Nigrospora aurantiaca	CGMCC 3.18130*	-	KX986064	KY019465	KY019295
N. aurantiaca	LC7034	-	KX986093	KY019598	KY019394
N. bambusae	CGMCC 3.18327*	-	KY385307	KY385319	KY385313
	LC7245	-	KY385305	KY385321	KY385315
N. brasiliensis	CMM 1214*	-	KY569629	MK720816	MK753271
	CMM 1217	-	KY569630	MK720817	MK753272
N. camelliae-sinensis	CGMCC 3.18125*	-	KX985986	KY019460	KY019293
N. chinensis	LC6851	-	KX986049	KY019579	KY019450
	CGMCC 3.18127*	-	KX986023	KY019462	KY019422
	TAP24N249	Dabong-dabong	**LC868051**	**LC868104**	**LC868157**
N. covidalis	CGMCC 3.20538*	-	OK335209	OK431479	OK431485
	LC158337	-	OK335210	OK431480	OK431486
	LC2725	-	KX985960	KY019487	KY019313
	LC4566	-	KX986022	KY019545	KY019354
N. endophytica	A.R.M. 687	-	OM265226	OP572418	OP572415
	A.R.M. 973*	-	OM265233	OP572420	OP572416
N. falsivesicularis	CGMCC 3.19678*	-	MN215778	MN329942	MN264017
	LC13553	-	MN215779	MN329943	MN264018
N. globospora	CGMCC 3.20539*	-	OK335211	OK431481	OK431487
	LC15839	-	OK335212	OK431482	OK431488
N. gorlenkoana	CBS 480.73*	-	KX986048	KY019456	KY019420
N. guangdongensis	CFCC:53917*	-	MT017509	MT024495	MT024493

N. guilinensis	LC7301	-	KX986063	KY019608	KY019404
	CGMCC 3.18124*	-	KX985983	KY019459	KY019292
N. hainanensis	CGMCC 3.18129*	-	KX986091	KY019464	KY019415
	A.R.M.967	-	OM265228	OM793057	OM642834
	A.R.M.968	-	OM265229	OM793058	OM642835
	A.R.M.972	-	OM265232	OP572419	OM642837
	A.R.M.976	-	OM265236	OM793060	OP572417
N. lacticolonia	CGMCC 3.18123*	-	KX985978	KY019458	KY019291
	A.R.M. 921	-	OM265227	OM642838	OM642833
	TAP24N001	Ula	**LC846677**	**LC846784**	**LC846891**
	TAP24N003	Ula	**LC846678**	**LC846785**	**LC846892**
	TAP24N005	Ula	**LC846679**	**LC846786**	**LC846893**
	TAP24N006	Ula	**LC846680**	**LC846787**	**LC846894**
	TAP24N008	Ula	**LC846681**	**LC846788**	**LC846895**
	TAP24N010	Ula	**LC846682**	**LC846789**	**LC846896**
	TAP24N011	Ula	**LC846683**	**LC846790**	**LC846897**
	TAP24N012	Ula	**LC846684**	**LC846791**	**LC846898**
	TAP24N013	Ula	**LC846685**	**LC846792**	**LC846899**
	TAP24N014	Ula	**LC846686**	**LC846793**	**LC846900**
	TAP24N016	Ula	**LC846687**	**LC846794**	**LC846901**
	TAP24N017	Ula	**LC846688**	**LC846795**	**LC846902**
	TAP24N018	Ula	**LC846689**	**LC846796**	**LC846903**
	TAP24N019	Ula	**LC846690**	**LC846797**	**LC846904**
	TAP24N021	Ula	**LC846691**	**LC846798**	**LC846905**
	TAP24N022	Ula	**LC846692**	**LC846799**	**LC846906**
	TAP24N026	Ula	**LC846694**	**LC846801**	**LC846908**
	TAP24N027	Ula	**LC846695**	**LC846802**	**LC846909**
	TAP24N028	Ula	**LC846696**	**LC846803**	**LC846910**
	TAP24N029	Ula	**LC846697**	**LC846804**	**LC846911**
	TAP24N030	Ula	**LC846698**	**LC846805**	**LC846912**
	TAP24N031	Ula	**LC846699**	**LC846806**	**LC846913**
	TAP24N033	Ula	**LC846700**	**LC846807**	**LC846914**
	TAP24N035	Ula	**LC846702**	**LC846809**	**LC846916**
	TAP24N036	Ula	**LC846703**	**LC846810**	**LC846917**
	TAP24N038	Ula	**LC846704**	**LC846811**	**LC846918**
	TAP24N039	Ula	**LC846705**	**LC846812**	**LC846919**
	TAP24N041	Ula	**LC846706**	**LC846813**	**LC846920**
	TAP24N043	Ula	**LC846707**	**LC846814**	**LC846921**
	TAP24N045	Ula	**LC846708**	**LC846815**	**LC846922**
	TAP24N047	Ula	**LC846709**	**LC846816**	**LC846923**
	TAP24N048	Tugbok	**LC846710**	**LC846817**	**LC846924**
	TAP24N050	Tugbok	**LC846712**	**LC846819**	**LC846926**

TAP24N053	Tugbok	LC846713	LC846820	LC846927
TAP24N054	Tugbok	LC846714	LC846821	LC846928
TAP24N055	Tugbok	LC846715	LC846822	LC846929
TAP24N057	Tugbok	LC846716	LC846823	LC846930
TAP24N060	Tugbok	LC846717	LC846824	LC846931
TAP24N062	Tugbok	LC846718	LC846825	LC846932
TAP24N063	Tugbok	LC846719	LC846826	LC846933
TAP24N065	Tugbok	LC846720	LC846827	LC846934
TAP24N067	Tugbok	LC846721	LC846828	LC846935
TAP24N069	Tugbok	LC846723	LC846830	LC846937
TAP24N071	Tugbok	LC846724	LC846831	LC846938
TAP24N073	Tugbok	LC846725	LC846832	LC846939
TAP24N074	Tugbok	LC846726	LC846833	LC846940
TAP24N077	Tugbok	LC846727	LC846834	LC846941
TAP24N078	Tugbok	LC846728	LC846835	LC846942
TAP24N079	Tugbok	LC846676	LC846783	LC846890
TAP24N082	Little Panay	LC846730	LC846837	LC846944
TAP24N085	Little Panay	LC846731	LC846838	LC846945
TAP24N086	Little Panay	LC846732	LC846839	LC846946
TAP24N088	Little Panay	LC846733	LC846840	LC846947
TAP24N090	Little Panay	LC846734	LC846841	LC846948
TAP24N091	Little Panay	LC846735	LC846842	LC846949
TAP24N092	Little Panay	LC846736	LC846843	LC846950
TAP24N093	Little Panay	LC846737	LC846844	LC846951
TAP24N094	Little Panay	LC846738	LC846845	LC846952
TAP24N095	Little Panay	LC846739	LC846846	LC846953
TAP24N096	Little Panay	LC846740	LC846847	LC846954
TAP24N098	Little Panay	LC846741	LC846848	LC846955
TAP24N099	Little Panay	LC846742	LC846849	LC846956
TAP24N100	Little Panay	LC846743	LC846850	LC846957
TAP24N101	Little Panay	LC846744	LC846851	LC846958
TAP24N102	Little Panay	LC846745	LC846852	LC846959

TAP24N103	Little Panay	**LC846746**	**LC846853**	**LC846960**
TAP24N104	Little Panay	**LC846747**	**LC846854**	**LC846961**
TAP24N105	Little Panay	**LC846748**	**LC846855**	**LC846962**
TAP24N108	Little Panay	**LC846750**	**LC846857**	**LC846964**
TAP24N112	Little Panay	**LC846751**	**LC846858**	**LC846965**
TAP24N113	Little Panay	**LC846752**	**LC846859**	**LC846966**
TAP24N115	Little Panay	**LC846753**	**LC846860**	**LC846967**
TAP24N118	Little Panay	**LC846754**	**LC846861**	**LC846968**
TAP24N123	Little Panay	**LC846756**	**LC846863**	**LC846970**
TAP24N125	Little Panay	**LC846757**	**LC846864**	**LC846971**
TAP24N127	Little Panay	**LC846758**	**LC846865**	**LC846972**
TAP24N128	Little Panay	**LC846759**	**LC846866**	**LC846973**
TAP24N129	Little Panay	**LC846760**	**LC846867**	**LC846974**
TAP24N131	Little Panay	**LC846761**	**LC846868**	**LC846975**
TAP24N132	Panombon	**LC846762**	**LC846869**	**LC846976**
TAP24N134	Panombon	**LC846763**	**LC846870**	**LC846977**
TAP24N136	Panombon	**LC846764**	**LC846871**	**LC846978**
TAP24N139	Panombon	**LC846766**	**LC846873**	**LC846980**
TAP24N141	Panombon	**LC846767**	**LC846874**	**LC846981**
TAP24N145	Panombon	**LC846768**	**LC846875**	**LC846982**
TAP24N146	Panombon	**LC846769**	**LC846876**	**LC846983**
TAP24N149	Panombon	**LC846770**	**LC846877**	**LC846984**
TAP24N154	Panombon	**LC846771**	**LC846878**	**LC846985**
TAP24N156	Panombon	**LC846772**	**LC846879**	**LC846986**
TAP24N157	Panombon	**LC846773**	**LC846880**	**LC846987**
TAP24N163	Panombon	**LC846774**	**LC846881**	**LC846988**
TAP24N164	Panombon	**LC846775**	**LC846882**	**LC846989**
TAP24N165	Panombon	**LC846776**	**LC846883**	**LC846990**
TAP24N166	Panombon	**LC846777**	**LC846884**	**LC846991**
TAP24N167	Panombon	**LC846778**	**LC846885**	**LC846992**
TAP24N168	Panombon	**LC846779**	**LC846886**	**LC846993**
TAP24N174	Panombon	**LC846780**	**LC846887**	**LC846994**
TAP24N176	Panombon	**LC846781**	**LC846888**	**LC846995**
TAP24N177	Panombon	**LC846782**	**LC846889**	**LC846996**

	TAP24N204	Patag-Alanib	**LC868092**	**LC868145**	**LC868198**
	TAP24N208	Patag-Alanib	**LC868088**	**LC868141**	**LC868194**
	TAP24N209	Patag-Alanib	**LC868087**	**LC868140**	**LC868193**
	TAP24N226	Patag-Alanib	**LC868097**	**LC868150**	**LC868203**
	TAP24N230	Dabong-dabong	**LC868070**	**LC868123**	**LC868176**
	TAP24N231	Dabong-dabong	**LC868069**	**LC868122**	**LC868175**
	TAP24N233	Dabong-dabong	**LC868067**	**LC868120**	**LC868173**
	TAP24N234	Dabong-dabong	**LC868066**	**LC868119**	**LC868172**
	TAP24N235	Dabong-dabong	**LC868065**	**LC868118**	**LC868171**
	TAP24N236	Dabong-dabong	**LC868064**	**LC868117**	**LC868170**
	TAP24N238	Dabong-dabong	**LC868062**	**LC868115**	**LC868168**
	TAP24N239	Dabong-dabong	**LC868061**	**LC868114**	**LC868167**
	TAP24N240	Dabong-dabong	**LC868060**	**LC868113**	**LC868166**
	TAP24N242	Dabong-dabong,	**LC868058**	**LC868111**	**LC868164**
	TAP24N243	Dabong-dabong	**LC868057**	**LC868110**	**LC868163**
	TAP24N245	Dabong-dabong	**LC868055**	**LC868108**	**LC868161**
	TAP24N251	Dabong-dabong	**LC868049**	**LC868102**	**LC868155**
	TAP24N254	Dabong-dabong	**LC868047**	**LC868100**	**LC868153**
	TAP24N255	Dabong-dabong	**LC868046**	**LC868099**	**LC868152**
N. manihoticola	A.R.M. 645*	-	OM265224	OM869479	OM914791
N. musae	CBS 319.34*	-	KX986076	KY019455	KY019419
	LC6385	-	KX986042	KY019567	KY019371
N. nigrocolonia	**TAP24N202**	Patag-Alanib	**LC868094**	**LC868147**	**LC868200**
	TAP24N222	Patag-Alanib	**LC868075**	**LC868128**	**LC868181**
	TAP24N224	Patag-Alanib	**LC868073**	**LC868126**	**LC868179**
	TAP24N229	Dabong-dabong	**LC868071**	**LC868124**	**LC868177**
	TAP24N232	Dabong-dabong	**LC868068**	**LC868121**	**LC868174**

	TAP24N237	Dabong-dabong	**LC868063**	**LC868116**	**LC868169**
	TAP24N246	Dabong-dabong	**LC868054**	**LC868107**	**LC868160**
	TAP24N250	Dabong-dabong	**LC868050**	**LC868103**	**LC868156**
	TAP24N252	Dabong-dabong	**LC868048**	**LC868101**	**LC868154**
N. oryzae	LC2724	-	KX985959	KY019486	KY019312
	LC4265	-	KX985994	KY019518	KY019335
N. osmanthi	CGMCC 3.18126*	-	KX986010	KY019461	KY019421
	LC4487	-	KX986017	KY019540	KY019438
N. pernambucoensis	A.R.M.651	-	OM265225	OM869480	OM914792
	A.R.M. 974*	-	OM265234	OM869481	OM914793
N.philosophiae-doctoris	CGMCC 3.20540*	-	OK335214	OK431484	OK431490
N. pyriformis	CGMCC 3.18122*	-	KX985940	KY019457	KY019290
	A.R.M.970	-	OM265231	OM642839	OM513904
N. rubi	LC2698*	-	KX985948	KY019475	KY019302
N. saccharicola	LC12057	-	MN215789	MN329952	MN264028
	CGMCC 3.19362*	-	MN215788	MN329951	MN264027
N. sacchari-officinarum	CGMCC 3.19335*	-	MN215791	MN329954	MN264030
	LC13531	-	MN215792	MN329955	MN264031
N. singularis	CGMCC 3.19334*	-	MN215793	MN329956	MN264032
	LC12068	-	MN215794	MN329957	MN264033
	TAP24N034	Ula	**LC846701**	**LC846808**	**LC846915**
	TAP24N049	Tugbok	**LC846711**	**LC846818**	**LC846925**
	TAP24N068	Tugbok	**LC846722**	**LC846829**	**LC846936**
	TAP24N081	Little Panay	**LC846729**	**LC846836**	**LC846943**
	TAP24N120	Little Panay	**LC846755**	**LC846862**	**LC846969**
	TAP24N137	Panombon	**LC846765**	**LC846872**	**LC846979**
	TAP24N201	Patag-Alanib	**LC868095**	**LC868148**	**LC868178**
	TAP24N203	Patag-Alanib	**LC868093**	**LC868146**	**LC868199**
	TAP24N205	Patag-Alanib	**LC868091**	**LC868144**	**LC868197**
	TAP24N206	Patag-Alanib	**LC868090**	**LC868143**	**LC868196**
	TAP24N207	Patag-Alanib	**LC868089**	**LC868142**	**LC868195**
	TAP24N210	Patag-	**LC868086**	**LC868139**	**LC868192**

		Alanib			
	TAP24N211	Patag-Alanib	**LC868085**	**LC868138**	**LC868191**
	TAP24N212	Patag-Alanib	**LC868084**	**LC868137**	**LC868190**
	TAP24N213	Patag-Alanib	**LC868083**	**LC868136**	**LC868189**
	TAP24N214	Patag-Alanib	**LC868082**	**LC868135**	**LC868188**
	TAP24N215	Patag-Alanib	**LC868081**	**LC868134**	**LC868187**
	TAP24N217	Patag-Alanib	**LC868080**	**LC868133**	**LC868186**
	TAP24N218	Patag-Alanib	**LC868079**	**LC868132**	**LC868185**
	TAP24N219	Patag-Alanib	**LC868078**	**LC868131**	**LC868184**
	TAP24N220	Patag-Alanib	**LC868077**	**LC868130**	**LC868183**
	TAP24N221	Patag-Alanib	**LC868076**	**LC868129**	**LC868182**
	TAP24N223	Patag-Alanib	**LC868074**	**LC868127**	**LC868180**
	TAP24N227	Patag-Alanib	**LC868063**	**LC868125**	**LC868178**
	TAP24N241	Dabong-dabong	**LC868059**	**LC868112**	**LC868165**
	TAP24N244	Dabong-dabong	**LC868056**	**LC868109**	**LC868162**
	TAP24N247	Dabong-dabong	**LC868053**	**LC868106**	**LC868159**
	TAP24N248	Dabong-dabong	**LC868052**	**LC868105**	**LC868158**
	TAP24N256	Dabong-dabong	**LC868096**	**LC868149**	**LC868202**
	TAP24N257	Dabong-dabong	**LC868045**	**LC868098**	**LC868151**
N. sphaerica	LC2839	-	KX985964	KY019491	KY019317
	LC2840	-	KX985965	KY019492	KY019318
	TAP24N107	Little Panay	**LC846749**	**LC846856**	**LC846963**
Nigrospora sp.	LC6704	-	KX986047	KY019571	KY019373
N. vesicularis	LC0322	-	KX985939	KY019467	KY019296
	CGMCC 3.18128*	-	KX986088	KY019463	KY019294
N. vesicularifera	CGMCC 3.19333*	-	MN215812	MN329975	MN264051
	A.R.M.975	-	OM265235	OM642840	OM513905
N. vesicularifera	**TAP24N023**	Ula	**LC846693**	**LC846800**	**LC846907**
N. zimmermanii	CBS 290.62*	-	KY385309	KY385317	KY385311

CBS 984.69   -   KY385310   KY385322   KY385316