BIOLOGICAL SCIENCES, Neuroscience

Computational mechanisms for temporal integration in the anterior claustrum

Kuenbae Sohn, Donghyeon Yoon, Junghwa Lee[*] & Sukwoo Choi[*]

School of Biological Sciences, College of Natural Sciences, Seoul National University, 1 Gwanak-ro, Seoul 151-747, Korea,

K.S. and D.Y. contributed equally to this manuscript.

[*]*Correspondence should be addressed to Dr. Junghwa Lee and Dr. Sukwoo Choi, E-mail: antares0715@gmail.com or sukwoo12@snu.ac.kr*

Author contributions: K.S., J.L., and S.C. designed research; and K.S., and D.Y. performed research; K.S., D.Y., J.L., and S.C. analyzed data; and K.S., D.Y., J.L., and S.C. wrote the paper.

Key words: Claustrum, Recurrent Neural Networks, Integration, Trajectory, Dynamic coding

Abstract

The claustrum, with its extensive reciprocal connections to nearly all cortical regions, has long been hypothesized as a key hub for integrating diverse cognitive, sensory and motor information. However, despite its anatomical connectivity, whether and how it functionally integrates different inputs to generate coherent representations has remained unclear. Here, we developed a recurrent neural network (RNN) trained via supervised learning on behavioral metrics of delayed escape—a behavioral paradigm that requires integration of temporally separated task-relevant signals. A subset of RNN neurons exhibited dynamics similar to those of anterior claustral neurons during this behavior. These neurons formed a recurrent cluster, a structure supported by in vitro stimulation experiments in claustral brain slices. We analyzed the computational properties of this claustrum-like cluster via dimensionality reduction of population activity. The network showed nonlinear integration of temporally distributed inputs and increased synergistic information. Rather than settling into attractors, integrated information was dynamically encoded along continuously evolving neural trajectories. Notably, similar trajectory patterns associated with dynamic integration were observed in claustral recordings, suggesting the model's biological plausibility. We propose that the anterior claustrum dynamically integrates task-relevant input signals over time and broadcasts the evolving representation to downstream brain regions capable of reading and interpreting it in a context-dependent manner.

Introduction

The claustrum is a thin, elongated sheet of gray matter that maintains extensive reciprocal connections with nearly all cortical areas and several subcortical regions. This dense interconnectivity has led to the longstanding hypothesis that the claustrum serves as a hub for integrating internal and external signals into a unified percept, potentially supporting conscious awareness and higher-order cognition (1). However, direct experimental evidence for this hypothesis has remained scarce. One recent study reported that individual claustral neurons can receive convergent input from multiple cortical areas, which is critical for a behavioral task requiring multimodal sensory inputs (2). In contrast, numerous studies, including those from the Mathur and Citri groups, have implicated the claustrum in diverse cognitive processes such as attention, salience detection, and cognitive control (3-14). This apparent discrepancy may reflect

differences in the specific claustral subregions examined. Although the claustrum is less

differentiated in rodents than in primates, important distinctions exist even among rodent

species (15). In mice, the anterior boundary of the claustrum typically does not extend beyond

the rostral end of the striatum (16). In contrast, in rats, the claustrum extends further rostrally

beneath the forceps minor of the corpus callosum (fmi), where claustrum-specific gene

expression has been reported (17, 18).

Only a few previous studies have directly investigated this rostral segment of the rat claustrum

(rostral-to-striatum, rsCla) (18-22). However, recent findings suggest that neuronal activity in

this region plays a critical role in behavioral tasks in which two different kinds of information

must be integrated (17, 23), indicating a potentially distinct computational role for the rsCla.

Among these tasks, the newly developed delayed escape task provides a behavioral paradigm

in which rats must rapidly escape to a neutral zone following a fear-inducing conditioned

stimulus (CS), after a delay that separates the CS from the opening of an outlet. Crucially, the

task requires no prior training and instead depends on the flexible integration of two temporally

separated events that become jointly relevant for guiding escape behavior. Neural recordings

from this task revealed that CS-related information was maintained through persistent activity in

rsCla neurons, suggesting that these neurons may integrate the sustained internal threat signal

with the subsequent outlet-opening signal to guide future escape behavior.

To investigate how the anterior claustrum integrates information at the population level, we

employed a delayed escape task in Long-Evans rats. Because the task could only be performed

once per animal, and only a small number of single units were recorded per subject, direct

population-level analysis was severely limited. To overcome these constraints, we trained a

recurrent neural network (RNN) model solely on behavioral outputs—a strategy previously

shown to yield biologically plausible dynamics (24-30). We then compared the network's

emergent population activity patterns with in vivo claustral recordings, and used the model to

explore circuit-level mechanisms of integrating temporally separated inputs.

Results

RNNs trained on the Delayed Escape Task show claustrum-like dynamical patterns

The delayed escape task allows only a single test trial per animal, which means that

opportunities to obtain neurophysiological data are extremely limited. To overcome this

limitation, we used recurrent neural network (RNN) simulations, a strategy that has been widely

adopted in recent neuroscience research (26, 27, 31). In the delayed escape task, when a

conditioned stimulus (CS) previously associated with electric shock is presented for the first time

in a novel environment, animals infer that a value-neutral alternative space is likely to be safer

and therefore exhibit faster escape behavior (Fig. 1A, This figure is adapted from Fig. 1A of Han

et al. (17)). Importantly, the passage to this neutral space opens after a delay following CS

offset, making it essential for CS-related information to be maintained beyond the actual

presence of the CS. As a control group, animals were placed in the same environment during

the CS period but did not receive the CS. In previous experiments, animals that received the CS

escaped to the neutral space significantly faster than controls (17). To model this behavior, we

built upon the continuous-time RNN framework developed by Ehrlich et al. (2021) (24) and

Ehrlich & Murray (2022) (25) (Fig. 1B).

The trained RNN reproduced the escape latency pattern reported by Han et al. (2024) (17): the

CS + door-opening condition escaped faster than the control door-opening only group (Fig. 1C),

reflecting the in vivo behavioral results. Furthermore, it showed similarities to the claustral

network dynamics observed in that study (Fig. 1D–L, Fig. 1D, E, F and L(left) are adapted from

Figs. 4B, 4D, 4E and S4E of Han et al. (17)). In the previous biological study, the responses of

individual claustral units representing each behavioral epoch were subjected to dimensionality

reduction and clustering (Fig. 1D). One of these clusters showed persistent firing during and

after CS presentation, and the magnitude of this increase was inversely correlated with escape

latency (Fig. 1E–G). From this, we concluded that this cluster maintained CS signals until the

door to the escape path opened, thereby contributing to delayed escape.

To determine whether this cluster was reproduced in the trained RNN, unit responses in the

RNN representing each behavioral epoch were also subjected to dimensionality reduction and

clustering. One of the clusters displayed the same persistent activation during and after CS

presentation as observed in the in vivo claustrum (Fig. 1H–J). As in the biological in vivo data,

stronger activation in this model cluster predicted shorter escape latency (Fig. 1K). Single-unit

responses were also similar between real and simulated data: the persistence after the CS did

not arise from a single unit firing continuously but rather from many units intermittently

increasing their firing, which overlapped to form a persistent signal at the population level (Fig.

1L).

Even when no CS was presented and only the door opening cue was given (the door-opening

condition), the population activity pattern generated by the RNN showed a pattern of gradually

increasing after the door opened and then decreasing as the crossing approached (Fig. S1A).

The other two model clusters exhibited different patterns: Cluster 2 was generally suppressed

regardless of CS presence, and Cluster 3 gradually ramped up from the door opening cue to

just before the crossing, also regardless of CS presence (Fig. S1).

Next, we quantified the contributions of the three clusters to the RNN output when the CS was

presented. The claustrum-like Cluster 1 contributed very little to the output, whereas Cluster 3

was dominant in influencing cross behavior. This suggests that Cluster 1 may not directly drive

the output but instead processes the CS and door opening signals and broadcasts this

information to other clusters (Fig. S2). In our previous study (17), optogenetic inhibition of the

anterior claustrum during the delay period between CS offset and door opening significantly

increased escape latency (Fig. 1M, Fig. S3A). Introducing inhibitory input to suppress RNN

Cluster 1 during the same period produced a similar increase in latency (Fig. 1M), whereas

inhibiting the other clusters had no effect (Fig. S3B). Furthermore, when the interval between

CS and door opening was extended from 5 s to 180 s in vivo (17), the latency difference

between the CS and neutral groups disappeared. Indeed, the RNN also reproduced this result

under the same extended interval (Fig. S3C).

Taken together, the claustrum-like Cluster 1 in the RNN reproduced the firing pattern and

behavioral correlation of the persistent activity cluster observed in vivo and resembled several

key characteristics of actual claustral neurons in almost every respect, including latency

increases following inhibition. This trained RNN model therefore enables deeper analyses under

experimental conditions that are otherwise difficult to implement.

Claustral neurons exhibit recurrent connectivity similar to that of the RNN

Because a recurrent neural network (RNN), by definition, contains recurrent connections among

its units, finding a claustrum-like cluster whose properties mirror those of the real claustrum

suggests that biological claustral neurons may likewise be interconnected through recurrent

circuitry. As expected, the RNN exhibited strong recurrent weights within Cluster 1 (Fig. 2A). If a

similar architecture exists in vivo, the first prediction is that excitatory connectivity should be

detectable between claustral neurons.

To test this, we sparsely expressed ChrimsonR in a subset of claustral cells and prepared

sagittal brain slices for patch-clamp recordings to measure synaptic currents (Fig. 2B–C). We

activated ChrimsonR-expressing presynaptic neurons with 617 nm light, while patching

ChrimsonR-negative neurons to isolate synaptic responses. Patching ChrimsonR-positive cells
would have risked contamination by direct photocurrents in addition to synaptic responses.
Wide-field illumination induced excitatory postsynaptic currents (EPSCs) that were abolished by
NBQX + D-AP5, confirming that they were mediated by glutamatergic transmission (Fig. 2D–E).
One remaining concern was that these EPSCs might have originated from ChrimsonR-
expressing axons projecting into the claustrum from outside regions. To rule this out, we
employed high-resolution optical mapping. The wide field was divided into a 30 × 30 grid (900
pixels), and each pixel was stimulated for 2ms (Fig. 2F). EPSCs could be elicited not only near
the recorded soma but also from distant pixels, indicating the presence of local excitatory-to-
excitatory connectivity within the claustrum itself (Fig. 2G–H). These findings are consistent with
the previous study by Orman et al. (2015) (32) and Shelton et al. (2025) (2) but contrast with
reports that the posterior claustrum exhibits little excitatory-to-excitatory coupling (33). In
particular, Orman et al. (2015) reported that persistent increases in claustral activity were
observed only when brain slices were prepared at a specific slicing angle.

A more stringent prediction of recurrent wiring is that a brief, strong stimulus should induce
sustained neuronal activity. To test this, we expressed the calcium sensor GCaMP6f in the
anterior claustrum and delivered a 1 s train of electrical stimulation (20 Hz, 200 µA) in horizontal
slices (Fig. 2I–K, Fig. S4C). This stimulation evoked calcium signals that remained elevated for
over 10 s. Consistently, patch-clamp recordings from GCaMP6f-positive cells confirmed a
prolonged increase in firing rate over a similar period (Fig. S4B). Notably, coronal slices failed to
show such persistence, suggesting that horizontal slicing better preserves recurrent excitatory
circuits. To determine whether this persistent activity depends on synaptic reverberation, we
locally pressure-injected NBQX + D-AP5 at the stimulation site during the sustained phase of
the calcium signal. As predicted, the persistent response collapsed rapidly (Fig. 2L–M, Fig.
S4D–F), indicating that excitatory synaptic transmission within the claustrum is essential for
generating the sustained activity.

The RNN reproduced a similar phenomenon (Fig. 2N, Fig. S4F). Isolating Cluster 1 from the
trained network, we reduced all inhibitory weights by 60%. When a 1 s CS-strength input was
delivered, the cluster exhibited >10 s of persistent activity, matching the slice data. Reducing
inhibitory weights facilitates the persistent activity, which may reflect the reduced inhibitory
connectivity often present in horizontal slice preparations. Next, reducing 60% of excitatory
weights for only 10% of Cluster 1 units—mimicking the spatially restricted application of NBQX
and D-AP5—largely abolished the persistence (Fig. 2N, Fig. S4D–F). Together, these slice and

RNN experiments suggest that the anterior claustrum contains recurrent excitatory connections
capable of supporting prolonged activity.

PCA-based trajectory analysis characterizes RNN dynamics

Having established that claustral neurons exhibit recurrent connectivity capable of sustaining
activity, we next examined how such dynamics manifest at the population level in the RNN.
Based on the results so far, the claustrum-like cluster in the RNN resembles the cluster showing
persistent activity after CS presentation in the anterior claustrum. To visualize the population
activity dynamics of the RNN clusters—particularly the claustrum-like cluster— we applied
principal component analysis (PCA) to z-scored firing rates from three conditions: CS + door-
opening, door-opening only, and CS only (Fig. 3A–E, Fig. S5–6). At each time point, the
ensemble firing rates across all neurons in each cluster were treated as a population activity
vector and projected into a three-dimensional space defined by the top 3 principal components.
Because cross-latency (the time from door opening to actual crossing) varied across trials, the
interval from door opening to crossing was time-normalized within each trial. For the CS only
condition, in which cross-latency was undefined, we used the mean cross-latency from the CS +
door-opening condition to define a notional crossing time.

In the CS + door-opening condition, the mean trajectory moved away from the start region after
CS presentation and progressed along a curved path. Notably, a short-loop pattern appeared
immediately after the door opened—5 seconds after CS offset (Fig. 3A). In contrast, in the door-
opening only group, the trajectory remained near the start region until the door opened and then
followed a long-curved path toward the crossing point (Fig. 3B). As expected, the CS only group
exhibited a trajectory similar to that of the CS + door-opening condition until 5 seconds after CS
offset, after which it returned toward the start region (Fig. 3C). Interestingly, in two
manipulations known to delay crossing latency—(1) inhibition of neural activity during the 5-
second interval between CS offset and door opening, and (2) extension of this interval to 180
seconds—the short loop observed in the CS + door-opening group was not observed (Fig. 3D,
E).

Sudden changes in trajectory shape, such as the short loop observed in the CS + door-opening
condition, have been reported in previous studies of other brain regions to be associated with
specific brain functions (34). Therefore, based on the assumption that the persisting CS signal
and the door-opening signal combine to drive integration, we examined their interaction during

this period. To test this, we assumed that the post-CS segment of the CS only trajectory

(starting 5 seconds after CS offset) and the post-door-opening segment of the door-opening

only trajectory each represent the respective inputs to the integration process. We then tested

whether their combination could explain the post-door-opening trajectory in the CS + door-

opening condition (Fig. 3F-G, Fig S6). A linear combination model failed to reconstruct the early

time bins immediately after door opening, likely due to the distinctive short-loop shape. In

contrast, a nonlinear model (multilayer perceptron, MLP) achieved a lower residual sum of

squares (RSS), outperforming the linear model (Fig. 3G, Fig S6A). The difference in model fit

between the nonlinear and linear models was significantly larger in the CS + door-opening

condition than in either the inhibition or 180-second interval conditions (Fig. 3G). Notably, the

difference in the CS + door-opening condition was specific to the claustrum-like cluster, as the

other two clusters showed no comparable difference (Fig. S6B–D). Taken together, these

results suggest that nonlinear integration occurs in the claustrum-like cluster specifically during

the short-looping period immediately following door opening in the CS + door-opening condition.

Trajectory features predicted by the RNN are present in biological recordings
204
We next asked whether these trajectory features predicted by the RNN are also present in

biological claustral recordings. As mentioned earlier, our previous study (17) was limited by the

use of single-trial testing, a small number of recorded units per rat, and a limited total number of

animals, which restricted both the overall population size and the feasibility of in-depth analyses.

To address these limitations, we constructed a trained RNN model that produced reproducible

and testable trajectories, particularly those exhibiting short-loop dynamics suggestive of

information integration. To compare the model-derived trajectories with empirical neural data,

we applied Gaussian Process Factor Analysis (GPFA) (35, 36), a dimensionality reduction

method well-suited for single-trial time-series data. Although GPFA is typically optimized with

multiple trials, it can still provide a useful low-dimensional representation of neural activity. By

applying GPFA to single-trial unit data collected from multiple animals and embedding it into a

three-dimensional latent space, we were able to visualize population dynamics in a manner

comparable to the trajectories generated by the RNN, facilitating comparison between model

dynamics and experimental observations (Fig. 4A).

Interestingly, the trajectory derived from units recorded in CS rats—the experimental group

corresponding to the CS + door-opening condition in the RNN model—resembled the short-loop

structure observed in the RNN (Fig. 4A, left). The onset of the short loop was markedly delayed

compared to that in the claustrum-like cluster of the RNN. In contrast, trajectories from neutral

rats, which received a neutral CS (a light cue not associated with an electrical shock) during

testing, exhibited a gentler curvature (Fig. 4A, right). To confirm the reliability of this structure,

we partitioned the neurons into two subgroups and independently repeated the GPFA analysis.

Neurons with comparable basal firing rates were paired and subsequently classified into two groups.

To compare trajectories before and after the split, the behavioral reference points on the post-

split trajectories were aligned to those on the pre-split trajectories. The resulting latent

trajectories remained qualitatively similar in shape and evolution (Fig. 4B), suggesting that the

short-loop structure reflects coordinated activity across the neuronal population rather than

being driven by a small subset of neurons.

To further examine this similarity quantitatively, we performed PCA on the trajectories and

quantified their local geometry using an eigenvalue-based metric ($\lambda_3 / \Sigma\lambda$) (Fig. 4C−E). This

metric reflects the proportion of variance not explained by the plane defined by PC1 and PC2,

thereby capturing deviations from planarity in the trajectory. Importantly, this geometric metric

does not rely on recovering neuron-to-neuron noise correlations. Nevertheless, the estimation of

the latent space still depends on the covariance structure among claustral neurons, suggesting

that the inferred trajectories remain tied to biologically meaningful population dynamics. When

applied to the PCA trajectories of the RNN, the metric was significantly higher in the CS + Open

condition than in the Open-only condition (Fig. 4C). Similarly, when the GPFA trajectories from

real claustral neurons were reanalyzed using PCA and the same metric was applied, the CS

group showed significantly higher values than the Neutral group (Fig. 4D). Furthermore, a leave-

one-neuron-out analysis revealed that claustral neurons in the CS group contributed more

uniformly to this metric compared to those in the Neutral group (Fig. 4E). These findings support

the interpretation that the observed short-loop structure did not arise from a small subset of

neurons, outliers, or chance fluctuations.

Partial information decomposition uncovers neurons carrying synergy between CS and
door-opening signals

Having established that nonlinear integration emerges at the population level, we next asked

how this integration is implemented at the level of individual neurons. Because this analysis

requires full population observability and repeated trials, which are not feasible in the biological

dataset, we performed the following analyses using the trained RNN model. To examine the

contribution of individual neurons to this nonlinear integration, we measured the synergistic

information for each neuron—i.e., information encoded only when both stimuli, CS and door

opening, were present. To this end, we performed Partial Information Decomposition (PID)

analysis (37, 38). PID quantifies how two input variables (CS and door opening) contribute to

the information contained in the output neural activity by decomposing the total mutual

information into three components: (1) unique information provided independently by each input,

(2) redundant information shared by both inputs, and (3) synergistic information generated only

when both inputs are present simultaneously (Fig. 5C).

In the trained RNN model (as well as in real claustral neurons), the effect of an input stimulus

persists in neural activity even after the physical stimulus has ended. Therefore, to define

stimulus strength for PID analysis, we used the output of trained decoders—which infer the

presence or absence of each stimulus from neural activity—as the effective input variable in the

PID computation. We trained two decoders using the z-scored activity of neurons in the RNN

claustrum-like cluster: one to distinguish between CS-present and CS-absent trials, and another

to classify door-open versus no-door-open trials (Fig. 5A–B, Fig. S7; see Methods). The CS

decoder performed at chance level during the baseline period but sharply increased in accuracy

immediately after CS onset, with performance remaining high well beyond the end of CS

presentation (Fig. 5A). The door-opening decoder showed high accuracy during the period

when the door signal was present (Fig. 5B).

We used the CS decoding accuracy score and door-opening decoding accuracy score as

inputs, and the z-scored activity of claustrum-like cluster neurons in each trial as the output. PID

was computed for each neuron over time (Fig. 3D–H, see Methods). As expected, CS-unique

information rose following CS onset and gradually decayed, whereas door-opening-unique

information increased after door opening (Fig. S7C–D). Importantly, synergistic information—

representing new information generated only when CS and door-opening representations were

jointly present—increased immediately after door opening and then gradually decreased (Fig.

5E). This suggests that neurons in the claustrum-like cluster integrate CS and door-opening

signals to produce new, combined information. Moreover, synergistic signals were reduced in

two conditions where nonlinear integration was disrupted: when inhibition was applied to the

network, and when the CS–door interval was extended to 180 seconds (Fig. 3F–G).

We further divided claustrum-like cluster neurons into the top 25% with the highest synergy

("high-synergy" neurons) and the bottom 25% ("low-synergy" neurons) and analyzed their

trajectory features (i.e., the short loop). In high-synergy neurons, the curvature of the neural

trajectory immediately after door opening exhibited a more pronounced turning-angle change

compared to low-synergy neurons (Fig. S8A–B). Furthermore, when CS + door-opening trials

were predicted using trajectories from CS-only and door-opening-only conditions, nonlinear

prediction errors were greater for high-synergy neurons than for low-synergy neurons (Fig. 5H).

Together, these findings suggest that synergistic information at the single-neuron level is related

to the short loop and that, within the claustrum-like cluster, certain neurons play a more

dominant role in the nonlinear integration process.

294 Trajectory-based dynamic coding underlies integration

We next used the RNN model to examine how this integrated representation is dynamically

encoded over time. To assess the temporal evolution and stability of the integrated

representation, we applied cross-temporal decoding. This approach allowed us to examine

whether the combined information from CS and door-opening is encoded in a static manner or

in a time-varying fashion.

Using cross-temporal decoding, we examined how the neuronal population encodes information

unique to the CS + door-opening condition over time. For each corresponding time bin across

the three conditions (CS+Door, CS-only, and Door-only), we trained a decoder to distinguish the

CS+Door condition from the other two, and evaluated its classification accuracy across all time

bins. In the entire population of claustrum-like cluster, cross-temporal decoding revealed a

mixed coding regime: shortly after door opening, decoders trained at specific time bins

maintained discrimination power across relatively broad temporal windows (Fig. 6A). However,

as time progressed, the discriminative power of decoders became temporally confined,

performing well only near the training bin.

When neurons in the claustrum-like cluster were divided into high- and low-synergy groups and

decoders were trained and tested separately on each, high-synergy neurons exhibited a

dynamic coding pattern. Decoders trained on specific time bins in this group showed strong

performance at that time point, with limited generalization to neighboring bins. In contrast,

decoders trained on low-synergy neurons maintained more temporally stable performance.

These results suggest that high synergy neurons, which contribute critically to nonlinear

integration, encode CS and door-opening information using a dynamic coding scheme, whereas

low-synergy neurons may support a more stable representation (Fig. 6B–C).

This distinction was further supported by raw firing-rate heatmaps: both high and low synergy

neurons showed increased activity following CS onset, but after the door opened, only high-

synergy neurons displayed variable firing patterns (Fig. 6D, left). Notably, a similar pattern of

transient, variable spiking was observed in our in vivo recordings of claustral single units after
door opening in the CS + door-opening group (Fig. 6D, right).

Temporal evolution of the population state can be framed by two hypotheses (Fig. 6E). The
attractor hypothesis posits that neural dynamics consist of discrete or continuous attractors, with
the integrated state represented by transitions among them in response to varying inputs. The
trajectory hypothesis proposes that integration is embodied in the evolving trajectory itself,
independent of fixed attractors. To distinguish these views, we performed fixed-point analysis on
the RNN, locating states where the velocity vector is nearly zero (39). Fixed points were found
to lie far from the empirical trajectories, indicating that the network's dynamics are trajectory-
based rather than attractor-based when integrating CS and door-opening cues (Fig. 6F).
330
331

Discussion

In this study, we trained a vanilla recurrent neural network (RNN) solely on behavioral metrics
from the delayed escape task, which had previously been shown to require rsCla (17). The
trained network developed a claustrum-like cluster whose dynamics showed similarities to the
experimentally observed population activity patterns and reflected the relationship between
neural activity and behavior. Low-dimensional trajectory analysis revealed a nonlinear
integration of the conditioned stimulus (CS) and the temporally delayed door-opening signal,
accompanied by an increase in synergistic information. Rather than converging to stable
attractor states, the network encoded these signals through continuously evolving trajectories.
In the post-integration phase, the trajectories formed a sharply curved "short loop" immediately
after door opening—a trajectory pattern also observed in real claustrum recordings, thereby
supporting the biological plausibility of the model. Such temporal integration through dynamic
coding enables flexible responses to inputs that arrive in the claustrum with temporal gaps and
offers the advantage of generating richer representations compared with stable coding (27-29,
31, 34, 40). Moreover, contrary to the intuitive expectation that dynamically coded information
might be difficult to decode, it can, in fact, be easily read out using simple linear decoders (41),
which provides an additional advantage.

It is important to emphasize that behavior-trained recurrent neural networks (RNNs) can admit
multiple internal solutions capable of producing the same behavioral output. Accordingly, the

network analyzed in this study represents only one possible computational realization.
Nevertheless, the dynamical regime observed in the present model converged with several
independent lines of evidence from claustral recordings. These include persistent neural activity
during the delay period, the correlation between neural activity and escape latency, recurrent
connectivity, and the consistent geometric changes in population trajectories revealed through
population-level analyses. Taken together, these findings suggest that the present
computational model should not be interpreted as a direct implementation of the claustrum's
computational mechanism, but rather as a candidate model that captures a plausible
computational principle that may operate in the claustrum. In particular, conclusions derived
from the RNN analysis, such as increased synergy and dynamic coding, remain hypotheses that
could not be directly validated in the biological claustrum due to experimental constraints, and
therefore require future experimental testing. At the same time, the present model was
specifically designed to capture the computations required for the delayed escape task and
does not aim to account for the full range of claustral functions. In particular, it is not intended to
replace or encompass the diverse roles of the claustrum proposed in previous studies, including
those from the Mathur and Citri research groups (3-14). Future studies will be needed to
determine whether dynamical principles similar to those identified in the present model also
contribute to other proposed functions of the claustrum, including attention, salience detection,
cognitive control, sleep, and premotor control.

Previous studies have shown that the anterior claustrum is largely unresponsive to simple
sensory stimuli, becoming active only when such stimuli acquire behavioral relevance possibly
through convergent inputs from regions encoding memory, value, context, and motivational
states (7, 17, 23, 42-45). It also exhibits stronger responses to cross-modal stimuli when they
are semantically congruent (42, 46). These findings suggest that the claustrum preferentially
integrates behaviorally meaningful inputs via nonlinear operations to generate unified
representations. Within this framework, both the conditioned stimulus (CS) and the door-
opening cue in the delayed escape task may carry escape-relevant information. However, since
the behavioral meaning of the door-opening cue may require additional processing in the
biological circuit, the integration process may require additional time, which could explain the
delayed emergence of the short-loop trajectory in real neural data (see Fig. 4G, left). This
interpretation may be viewed as broadly compatible with prior theoretical proposals suggesting
that the claustrum contributes to the integration of semantically related information (1).
Furthermore, such integrated information is likely broadcast to higher-order cognitive regions

such as the PFC, ACC, and OFC, where it can be read out and interpreted (47), a possibility

that resonates with broader theoretical accounts of large-scale integrative brain function (1, 48).

Although our conclusions are mainly derived from modeling, the correspondence between

model predictions and empirical observations provides a rationale for future experimental

validation. Multi-trial behavioral paradigms and large-scale single-neuron recordings will be

essential to rigorously test these predictions. Furthermore, elucidating the nature of the

preprocessing that shapes inputs to the claustrum, as well as the downstream pathways

through which its integrated representations are accessed and utilized, will be critical for

understanding its role in brain-wide computation.

Materials and Methods

Animals

The following behavioral procedures were previously reported in Han et al. (2024) and are
summarized here for clarity (17). Male Long-Evans rats (Japan SLC Inc.) were dual-housed for
5–11 days before the experimental procedures began, under 12-hours inverted light/dark cycle
(light off at 9:00 a.m.) with ad libitum access to food and water. All behavioral experiments were
conducted during the dark portion of the cycle. The Institutional Animal Care and Use
Committee (IACUC) of the Seoul National University approved all experimental procedures, and
all experiments were performed following the guidelines for care and use of laboratory animals
of the Seoul National University. The in vivo single-unit recording experiments for each of the
CS and Neutral groups were performed separately.

Delayed escape task with in vivo single-unit recording

The followingrecording procedures were previously reported in Han et al. (2024) and are
summarized here for clarity (17). Rats (7 weeks old) were anesthetized with an intraperitoneal
(i.p.) injection of sodium pentobarbital (50 mg/kg) and maintained with isoflurane (1–1.5%) in
O2. Rats were mounted on a stereotaxic apparatus (Stoelting Co.). Fixed-wire electrodes were
bilaterally implanted into the rsCla (AP +3.35/ML ± 2.10/DV-4.50). A ground wire was implanted
in the cerebellum. The electrodes consisted of eight individually insulated nichrome microwires
(50 μm outer diameter, impedance 1-3 MΩ; California Fine Wire) contained in a stainless steel
guide cannula. The electrodes were affixed to the skull with screws using Poly-F zinc
polycarboxylate cement (Konstanz), vertex self-curing (vertex-dental, Zeist), and bond (Loctite
411, Henkel). Rats were allowed to recover for a week before they underwent experimental
procedures.
The rats in both the CS and Neutral groups were presented with the green light cue (10 s × 5
times) in the fear conditioning context with modification (a foamex floor instead of a metal grid
floor), which allowed for adaptation to the recording cable attachment. After 24 h, the rats were
presented with the green light cue (10 s × 5 times, average inter-trial interval: 90 s) in the same
context, to optimize recording parameters for each channel. Approximately 1 h later, the rats in
the CS group were fear-conditioned, and the rats in the Neutral group underwent pseudo-
conditioning in which the green light cues were presented but electrical foot shocks were
omitted.

On day 2, rats underwent the two-compartment chamber (54 × 23 cm area in total) exposure
session. The two compartments had identical floor sizes, but they were differently constructed
(compartment A vs. B: foamex floor vs. mesh grill floor; no visual cued walls vs. visual cued
walls with circles; 50 cm vs. 30 cm- height; a green LED bulb in the middle of the side wall and a
speaker right above the LED vs. no accessories installed on the wall, respectively). The two
compartments were connected by opposing square-shaped (7 × 7 cm) outlets, located 12 cm
above the floor. The outlet could be opened or closed by sliding. To ensure that the subject
localized a source of light within compartment A, the walls of the compartment were made of
non-reflective material (black foamex). Consequently, compartments A and B were illuminated
to a lesser intensity than in the fear conditioning performed on day 1, in which the walls were
made of reflective materials.
The rats were placed in compartment A to acclimate for 2 min on average on day 2. The outlet
in the wall (without any curtains) was then opened simultaneously with a tone pip sound (2.8
kHz, 200 ms, 85 dB). The rats were allowed to cross over to compartment B for a duration of 5
min. Once they crossed, the outlet was closed and the rats were allowed to stay in compartment
B for 2 min. The rats were then returned to their home cages. If a rat did not cross in 5 min, it
was removed from compartment A and temporarily moved to a small box near the two-
compartment chamber. Around 30 s later, the rat was re-introduced to compartment A and the
same procedure was performed. The rats that did not cross after 10 trials were excluded from
further experiments. We measured crossing latency and the number of trials in which the rat
succeeded in crossing. Crossing latency was defined as the duration from outlet opening to the
moment when all four paws of the rat touched the floor of compartment B. The chamber was
cleaned with 70% ethanol and then with distilled water prior to the beginning of each trial.
For the delayed escape test session (day 3), the two-compartment chamber was modified to
further prohibit any potential association between the outlet opening and crossing behavior. The
opened outlet was visually blocked by a black curtain installed at the outlet side of compartment
B. The rat head could go through the curtain to see compartment B since the curtain consisted
of two pieces of fabric adjoined at the centreline of the outlet. The rats repetitively put their
heads into compartment B through the curtain during the test session, but this behavior did not
appear to be associated with outlet crossing. In the modified chamber, a 20 s CS was presented
220 s after the placement of the rat in compartment A. The CS presentation was delayed when
the rat showed immobility at the time of CS onset. The outlet was opened with a tone pip sound
(2.8 kHz, 200 ms, 85 dB) 5 s after the CS offset, and the rat was allowed to cross the outlet for a
duration of 5 min. The test session was performed only once. The rats that did not cross during
the test session were considered to omit the task, and they were excluded from data analysis.
463
464   **RNN modeling**

465   The RNN model was implemented in PsychRNN (24, 25) as a continuous-time network with a
466   1s integration step. Each network consisted of a single hidden layer of 100 ReLU units, of which
467   80% were excitatory and 20% inhibitory.

468   Hidden-state $\boldsymbol{h}(t)$ dynamics followed

469
$$\tau\,\boldsymbol{\dot{h}}(t) = -\boldsymbol{h}(t) + f\left[\boldsymbol{W}_{\text{rec}}\boldsymbol{h}(t) + \boldsymbol{W}_{in}x(t) + \boldsymbol{b}_{\text{rec}}\right] + \boldsymbol{\eta}(t)$$
470
471
$$f(x) = \max(x, 0)$$
472
473
$$\boldsymbol{y}(t) = W_{\text{out}}\boldsymbol{h}(t) + b_{\text{out}}$$
474

An internal time constant of τ = 1000ms was adopted, in accordance with electrophysiological
measurements (Fig. S4), and ReLU activation f(x)=max(0,x) was used throughout the network.
Here, W_{rec}, W_{in}, and W_{out} denote the recurrent, input, and output weight matrices, respectively,
while b_{rec} and b_{out} are constant biases applied to the recurrent and output units. Gaussian noise
$\boldsymbol{\eta}(t)$ with variance $\sigma^2 = 0.05^2$ was injected into the recurrent layer. Simulations were conducted
using a continuous-time recurrent neural network with a discretized time step of 1000ms. To
ensure stable signal propagation and gradient flow during early training, the recurrent weight
matrix ($\boldsymbol{W}_{rec}$) and output weight matrix (W_{out})were initialized using a Glorot normal distribution.
Specifically, initial weights were drawn from a normal distribution centered on 0 with a standard
deviation of $\sigma = \sqrt{2/(N_{in} + N_{out})}$ where N_{in} and N_{out} denote the number of input and output
units for the given connectivity matrix. Input weights (W_{in}) were initially drawn uniformly from the
interval [0, 0.15], after which all rows corresponding to inhibitory units were zeroed out to
enforce Dale's principle. To strictly enforce Dale's principle throughout training, rows in (W_{out})
corresponding to inhibitory units were zeroed out, and recurrent weights were constrained such
that excitatory and inhibitory neurons maintained their respective positive and negative synaptic
projections.
We trained the RNN in a supervised manner using the behavioral data from rat experiments
(49). We provided the RNN with sequential inputs that matched the experimental timeline: a 30-
second baseline period with no input, followed by a 20-second CS input period, then a 5-second
interval with no input, and finally a sustained door-open signal until the end of the trial. In the
control group (neutral group), no input was provided during the period corresponding to CS
presentation, and trials from the CS and neutral groups were presented in a randomized order.
The network's output was treated as the probability of crossing; when this probability exceeded
0.5, the model was considered to have initiated a cross, allowing us to compute latency for each
trial.
The network received four input channels, denoted as $\mathbf{x}(t) = [x_{CS}, x_{door}, x_{inh}, x_{ctx}]^\mathsf{T}$. To reflect
the intense and salient nature of the fear-conditioning experience, the magnitude of the CS
channel ($x_{CS}(t)$) was set to 2 during the presentation of the CS, specifically from t = 70 s to t =
90 s (20 s duration), serving as a relatively stronger sensory drive compared to the simple
mechanical cue of the door opening. The door signal ($x_{door}(t)$) was set to 1 starting at t = 90 +
d s, where d was selected from [0, 1, 2.5, 5] seconds depending on curriculum learning, and
remained 0 beforehand. The inhibition signal ($x_{inh}(t)$) was set to -1 for the 5 seconds
immediately following the CS offset (t = 90-95 s) on inhibition trials. Finally, the context input
($x_{ctx}(t)$) was maintained at 1 throughout the entire trial.

Input weights were first drawn uniformly from the interval [0, 0.15], after which all rows
corresponding to inhibitory units were zeroed out to enforce Dale's principle.

A cross was defined as the first time point t^* at which the network output $y(t^*) \geq 0.5$. The
desired output trajectory $\tilde{y}(t)$ was specified as a step function transitioning at the target escape

$$\tilde{y}(t) = \begin{cases} 0, & t < t_{tar,} \\ 1, & t \geq t_{tar.} \end{cases}$$

513   time t_{tar}, which combines the door-opening time and the experimentally measured behavioral
514   latency measured in rats (latency for CS present: 48.7s, latency for CS absent: 111.3s):

$$t_{tar} = \begin{cases} t_{door} + \ell_{CS} = 90 + d + 48.7 \text{ s}, & \text{CS present,} \\ t_{door} + \ell_{noCS} = 90 + d + 111.3 \text{ s}, & \text{CS absent,} \end{cases}$$

516 Using fixed empirical latencies anchored the network to the average rat behavior, such that
517 remaining variability in output timing arose solely from the network's internal dynamics.
518 The per-trial loss function minimized during training was:

519
$$\mathcal{L} = -\sum_t [\tilde{y} \ln y + (1 - \tilde{y}) \ln(1 - y)] + \lambda(\|\mathbf{W}_{\text{in}}\| + \|\mathbf{W}_{rec}\|) + \lambda_{FR} \langle \|\text{h}(t)\|_2^2 \rangle_t$$

The L2 weight regularization coefficient $\lambda = 0.01$ and the firing rate regularization coefficient
$\lambda_{FR} = 0.95$. were systematically optimized via a grid search procedure to maintain biologically
plausible firing rates while preventing overfitting. The network was trained using TensorFlow
2.1.0 and the Adam optimizer (learning rate $\alpha = 5 \times 10^{-4}$) for 1.2 million iterations with a batch
size of 256. Each training batch comprised trials assigned randomly to CS-present or CS-absent
conditions with equal probability (50% each). The delay d between CS offset and door-opening
was incrementally increased (0,1, 2.5, and 5 s) once the batch-wise accuracy exceeded
thresholds of 0.5, 0.5, 0.5 and 1, respectively. All simulations were executed on a single RTX-
2080 Ti GPU using XLA just-in-time compilation.
After training was completed, the network was saved for downstream testing under various
experiment conditions, including inhibition, extended (180-s) interval, CS-only, no-CS/no-Door-
opening, and stimulation of the slice experiments. The resulting cross latencies t^*, neuronal
activity $\mathbf{h}(t)$, and weight matrices were exported to MATLAB for further analyses such as
clustering, principal component analysis (PCA), and Partial Information Decomposition (PID).
534
RNN Population Analysis
The recurrent neural network (RNN) simulation was processed with the same population-level
pipeline used for the experimental data (49).The first 10 s of each simulation were discarded to
eliminate variability caused by initial states. Activity from the next 30 s served as the baseline for
Z-score normalization.
For every CS trial (n = 147), Z-scores were averaged neuron-wise and then summarized across
five task epochs—cue, interval, early-open, late-open, and the 5 s after the cross. These epoch-
wise vectors were embedded into a three-dimensional t-SNE space (perplexity = 24,
exaggeration = 48, RNG = mt19937ar). The optimal number of clusters was determined with the
gap statistic and applied in a k-means clustering step. Through this procedure, we obtained
three clusters consisting of 24, 38, and 38 units (with 24 units in the claustrum-like cluster).
Here, the term claustrum-like refers to the fact that the mean response of the units remained
persistent not only during the CS presentation but also throughout the subsequent period until
crossing. The main difference between the empirical experiment and the RNN results is that, in
the actual experiment, each rat was tested with only one latency value, so correlations had to be
calculated across animals, whereas in the simulation, multiple trials were generated from a
single RNN, allowing correlations to be calculated across trials. Specifically, for each trial, the
mean z-scored activity across the 24 units during the cue-to-cross window was calculated, and
this value was paired with the crossing latency from that trial., and these z-scores were paired
with the crossing latency from that trial. With 100 trials, this yielded 100 data points. The
variability of model latency reflects both the injected noise and differences in initial states.
Notably, only a subset of the supervised RNNs we trained produced a claustrum-like cluster;
among 100 independent runs, only 5 met this criterion, and all results reported here are based
on one representative RNN network from those 5 cases.
As shown in Fig S2, the output contribution of each cluster for each time points was computed
as the product of the hidden-state vector $\mathbf{h}(t)$ and the output-weight matrix W_{out}.
561
Slice electrophysiology
Male Long-Evans rats (3–4 weeks old) were anesthetized (sodium pentobarbital, 50 mg/kg, i.p.)
and mounted in a stereotaxic frame. For optogenetic circuit mapping, AAV encoding ChrimsonR

(pAAV-CamKIIa-ChrimsonR-mScarlet-KV2.1; Addgene; 1×10^{13}–1×10^{14} vg/mL) was diluted
1:4 in sterile saline for sparse labeling and injected unilaterally (200 nL) into the rsCla (AP
+2.95, ML ±1.95, DV −3.85 mm). For calcium imaging, AAV encoding GCaMP6f
(pENN.AAV.CamKII.GCaMP6f.WPRE.SV40; Addgene; $\geq1\times10^{13}$ molecules/mL) was injected
bilaterally (400 nL) into the same coordinates. Viruses were delivered at 20 nL/min via a pulled
glass capillary using a pressure injection system (Nanoject II, Drummond; or Nanoliter 2010,
WPI). The pipette was left in place for 10 min post-infusion to ensure diffusion. Animals were
dual-housed for 3–4 weeks for viral expression. Subsequently, rats were anesthetized with
isoflurane and decapitated for acute slice preparation. The isolated whole brains were placed in
the slicing chamber filled with a warm (34~36 °C) artificial cerebrospinal fluid (aCSF) solution
containing 120 mM NaCl, 3.5 mM KCl, 1.25 mM NaH_2PO_4, 26 mM $NaHCO_3$, 1.3 mM $MgCl_2$, 2
mM $CaCl_2$, and 11 mM D-(+)-glucose, and continuously bubbled at room temperature with 95%
O_2/5% $CO_2.$ Acute sagittal or horizontal slices containing the rsCla (300 μm thickness) were
prepared using a vibratome (VT1200, Leica). Slices were then stored under submerged
conditions at 34 °C for 1.5 hour before recordings. For recording, slices were transferred to the
recording chamber in which aCSF was continuously perfused. The perfused aCSF was
continuously aerated with 95% O_2/5% $CO_,$ and maintained at 32 °C.
Whole-cell patch clamp recordings were performed using micropipettes (4~6 MΩ) which were
pulled from borosilicate glass capillaries (1.2 mm OD, 0.94 mm ID, Warner Instruments;
Massachusette, USA) with a micropipette puller (Pc-10, Narishige). The pipettes were filled with
the following solution: 120 mM potassium D-gluconate, 0.2 mM EGTA, 10 mM HEPES, 5 mM
NaCl, 2 mM Mg-ATP, 0.3 mM Na-GTP, and 1 mM $MgCl_2$, with the pH adjusted to 7.2 with KOH
and osmolality adjusted to approximately 297 mmol/kg with sucrose. Recordings were made
under infrared differential interference contrast (IR-DIC)–enhanced visual guidance from
neurons located in the rsCla slices. Neurons were first current-clamped neurons and then used
with membrane potentials lower than -50 mV. The neurons were voltage-clamped at -70 mV,
and solutions were delivered to slices via superfusion driven by gravity at a flow rate of 1.3
ml/min. The pipette series resistance was monitored throughout the experiments, and if it
changed by >20%, the data were discarded. To avoid direct light stimulation, whole cell
recordings were conducted using neurons in which the expression of ChrimsonR was absent.
Wide-field photostimulation was executed via a microscope's objective lens (40×/0.80NA) using
a power of 6.4 mW (measured at the lens level) and a wavelength of 617 nm. This stimulation
was achieved through a digital mirror device (Polygon 400, mightex) for a duration of 2 ms. To
stimulate each of the divided parts of the entire optical field, the same digital mirror device
(DMD) was used. The entire optical field was partitioned into a 30 by 30 grid, yielding a total of
900 equivalent sections. Each section received 2 ms of stimulation, and the heatmap was
generated using the Matlab interpolation algorithm with the EPSC data. The resultant EPSC
amplitudes were used to generate an EPSC heatmap encompassing the entire field. Whole-cell
currents were filtered at 2 kHz, digitized at up to 10 kHz, and stored on a microcomputer
(Clampex 10.7 software, Molecular Devices). We used 10 μM of NBQX (Tocris), and 50 μM of
D-AP5 (Tocris). One or two neurons were recorded per animal (a single neuron per slice). All
recordings were completed within 5 hours after slice preparation.
In brain slices where GCaMP6f was expressed, electrical stimulation was applied using a
Concentric Bipolar Electrode (FHC) via an isolator (WPI, A360). Stimulation was delivered at
150- 200 μA, 20 Hz, 1sec duration. The puffer pipette was positioned 75 μm below the slice
surface and 50 μm away from the electrode. At the 10-second mark post electrical stimulation,
pressure injection was carried out under a pressure of 20 psi. We used 100 μM of NBQX
(Sigma) and 500 μM of D-AP5 (Tocris) for this experiment. The analysis of fluorescence
changes in the recorded images follows these steps: Initially, cell boundaries were manually
marked for the 35 recorded GCaMP6f slice samples. Subsequently, a U-Net deep learning

model was trained on these images to develop a cell segmentation model. Using this model, the
attached cells were segmented and separated utilizing the watershed algorithm. Finally, all cells
were manually reviewed once more through video inspection.
618
RNN simulation adjusted for slice-stimulation conditions
To mimic the effect of electrical stimulation and applied NBQX+D-AP5 in the acute slice, we
constructed a reduced circuit composed exclusively of Cluster 1 (C1) units. All recurrent weights
outside C1 were zeroed, leaving only C1 → C1 connections in W_{rec}. Because inhibitory
neurons may lose efficacy in acute slices, every inhibitory weight in W_{rec} was scaled to 40 % of
its original value.
Stimulation was modeled by injecting an external drive $x_{stim}(t) = 2$ for 1 s. The corresponding
input weights were drawn uniformly from 0 to 0.15 for excitatory neurons only. Input weights to
the inhibitory neurons were set to zero.
In our model, we simulated the effects of applying NBQX + D-AP5 by introducing recurrent
inhibition, which began 10 seconds after the initial stimulation. The inhibition was targeted,
affecting only 10% of the neurons in the network. For these selected neurons, the strength of
the excitatory inputs they received from other neurons (W_{rec}) was reduced by 60%. This
simulation was designed to replicate key findings from our slice experiments (Fig. S4): First, the
effect of NBQX + D-AP5 is localized and does not spread widely; and second, drug application
resulted in an approximately 60% reduction in the amplitude of EPSCs.
635
Visualization of PCA trajectories
To visualize neural dynamics, trial-wise Z-scored firing rates from a specific neuronal cluster
were extracted for each simulation group (CS + Open, Open-only, and CS-only). Trials were
concatenated across groups and reshaped into a 2D matrix of size (trial x time) × neuron for
principal component analysis (PCA). PCA was performed on the combined matrix, and the first
three PCs were used for visualization.
Since each trial had a different latency to the cross event, trajectories were time-normalized.
Each trial was divided into two segments: a fixed-length pre-open segment (Segment A, from
the start to the open event; 55s) and a variable-length post-open segment (Segment B, from the
open event to the cross event). For trials without a defined Cross event (e.g., CS-only group),
the average cross latency from the CS + Open group was used. Segment B was interpolated to
a fixed length of 30 bins using shape-preserving piecewise cubic interpolation. The two
segments were concatenated and smoothed using a moving average. The smoothing window
size was determined heuristically to attenuate approximately 25% of the energy of the PCA
scores for each trial.
Time-normalized trajectories were averaged within each group to produce group-level mean
trajectories and corresponding standard errors of the mean (SEM). These trajectories, uniform
in the number of time bins, were visualized either separately for each group or overlaid for direct
comparison.
For both the inhibition simulation and the 180s interval simulation, corresponding trial data were
combined with CS + Open trials. These combined datasets were reshaped and processed using
the same PCA and time-normalization procedures described above.
658

Prediction of CS+Open Trajectories from Component Conditions

To evaluate how well CS+Open trajectories could be reconstructed from their component
conditions, we extracted time-normalized PCA trajectories from the CS+Open, Open-only, and
CS-only groups. For the CS+Open condition, trajectories were aligned to the door-opening

event, which occurred 5 seconds after CS offset; for the CS-only condition, trajectories were
taken beginning at the corresponding time point—5 s after CS offset. All trajectories were
preprocessed using the same procedures described in the trajectory visualization section. To
standardize across conditions, trajectories were trimmed to include only the period from the
open event to the cross event (30 bins), which served as the input for modeling.

We trained two models to predict individual CS+Open trial trajectories from the group-averaged
Open-only and CS-only trajectories:

Linear regression model: At each time bin, the trial trajectory was regressed onto the
corresponding bins of the mean Open-only and CS-only trajectories, yielding a bin-wise linear
estimate.

Multilayer perceptron (MLP): To test for nonlinear integration, we implemented a feedforward
neural network with two hidden layers ([6, 3] units) and L2 regularization ($\lambda \in [0, 0.25, 0.5, 0.75,$
$1]$), with the optimal λ chosen to minimize the RSS for each condition. For each condition
(CS+Open, inhibition, 180 s interval), the λ yielding the lowest residual sum of squares (RSS)
was selected. The MLP received concatenated mean trajectories from the Open-only and CS-
only conditions as inputs and was trained to predict the trial trajectory of the CS+Open
condition. Training was performed independently for each trial using a 70/15/15% split of bins
for training, validation, and test sets, with min–max normalization applied to the inputs.

Prediction performance was quantified as the RSS at each time bin. For the MLP, training was
repeated 10 times per trial, and RSS values were averaged across repetitions to yield a trial-
wise mean RSS time series. To assess the benefit of nonlinear integration, improvement by the
MLP over the linear model was calculated as the bin-wise difference between their RSS values,
normalized by the linear model RSS. These normalized scores were averaged across bins to
obtain a single summary value for each trial.

687
Decoding analysis (for Fig. 5A-B)
CS Decoding:
We first gathered the time-normalized Z-scores of all cluster neurons from two trial types: CS +
Open (cue present) and Open-only (cue absent). Time normalization from open to crossing was
identical to the procedure described above.
For each time bin we formed a trial-by-neuron matrix and trained a linear discriminant model
with fitcdiscr('linear') in MATLAB R2022b. Trials in which a cue was delivered were labelled 1
from cue onset onward, whereas cue-absent trials were labelled 0 throughout. Model
performance was assessed by five-fold cross-validation, and accuracy was reported as 1 –
kfoldLoss. Repeating this procedure over all bins produced a temporal profile of CS-decoding
accuracy.
Door Open Decoding:
A Door-open decoder was generated in the same way, but the training data comprised CS +
Open versus CS-only trials. Here, bins occurring after the outlet opened were labelled 1, and all
earlier bins (as well as every bin in CS-only trials) were labelled 0. Five-fold cross-validated
linear discriminant analysis again yielded a time-resolved accuracy curve that reflected when
RNN activity best predicted door opening.
705
Partial Information Decomposition (PID) Analysis

To quantify how CS and door-open information were distributed across RNN neurons, we
applied Partial Information Decomposition (PID) analysis to decoding accuracies as X variables
and neural activity as a Y in cluster 1–3. For each cluster, z-scored activity was extracted and
time-normalized using a fixed window of 85 bins, which included a 55s pre-open segment and a
30-bin post-open segment interpolated to a fixed length. Trials from four conditions were
included: CS + Open, Open-only, CS-only, and None.
For each neuron and each time bin, we computed PID terms (redundancy, CS-unique
information, Open-unique information, and synergy) using MATLAB code(the Neuroscience
Information Theory Toolbox (37)).
At each time bin, we used the neuron's z-scored activity rounded to one decimal place across
all trials as the Y (time × trial), and the CS and door-open decoding scores as X_1 and X_2,
respectively (time × trial) (see Fig. 5D).
Specifically, the construction of X_1 (cue-related events) was as follows.
For the CS+Open and CS-only conditions, decoding accuracy were computed as described
above. Trials in which a cue was delivered were labeled 1 from cue onset onward, whereas cue-
absent trials were labeled 0 throughout. A decoder was trained to classify the z-scored firing
rates in each time bin of CS+Open and Open-only trials according to these labels. To express
decoding accuracy as a score ranging from 0 to 1, we subtracted 0.5 from the decoding
accuracy, multiplied the result by 2, and set any negative values to 0. The resulting scores were
smoothed with a Gaussian kernel (window size = 15). To compute the probability of coincident
events, smoothed scores were rounded to the first decimal place. For Open-only and None
conditions, all time bins were assigned a value of 0.
The construction of X_2 (door-open-related events) was as follows.
For CS+Open trials, as described above, time bins after the outlet opening were labeled 1, and
all earlier bins as well as bins in CS-only trials were labeled 0. A decoder was trained to classify
each time bin's z-scored activity in CS+Open and CS-only trials based on these labels.
Decoding scores were then computed using the same method as described for X_1.
For Open-only trials, time bins following outlet opening were labeled 1, and all earlier bins as
well as bins in None trials were labeled 0. A decoder was trained to classify z-score in Open-
only and None trials, and decoding scores were computed identically.
For CS-only and None conditions, all bins were assigned a value of 0.
For each discrete state j of Y and each combination of states k from a source A (either X_1 alone,
X_2 alone, or their joint distribution [X_1, X_2]), we computed the corresponding conditional
probabilities required for PID analysis.
741
742    $P(A = k|Y = j) = P(Y = j, A = k)/P(Y = j), P(Y = j|A = k) = P(Y = j, A = k)/P(A = k)$
743
744   The information that source A conveys about Y when Y is in state j is then
745

746
$$I(Y = j; A) = \sum_k P(Y = j, A = k) \log_2 \frac{P(Y = j|A = k)}{P(Y = j)}$$

747
We defined its redundancy by, for each j, taking the minimum of $I(Y = j; A)$ over the X1 and X2,
and then computing the dot product of that vector of minima with the state-probabilities $P(Y =$
$j)$. X_1-unique information is the mutual information between Y and X1 minus the redundancy.
X2-unique information is the mutual information between Y and X2 minus the redundancy.
Synergy is obtained by taking the joint mutual information, subtracting both unique-information
terms, and then adding back the redundancy.

PID terms were averaged across neurons within each cluster. For visualization, time-resolved
PID terms were shown as mean ± SEM across neurons.
756
Cross-temporal Decoding Analysis
Cross-temporal decoding of CS+Open information was performed as follows: For each time bin,
a decoder was trained to classify z-scored neural activity from CS+Open trials (all time bins
labelled as 1) versus CS-only and Open-only trials (all time bins labelled as 0). The trained
decoder was then tested across all time bins to predict trial type. Decoding accuracy was
computed as (correct trials / total trials × 100) and visualized as a two-dimensional heatmap
indexed by training and testing time bins.
764
765 **Fixed Point Analysis**
766 We implemented the FixedPointFinder toolbox to identify fixed points in the trained RNN
767 model(39).
768 Fixed point analysis was performed separately for the following input epochs, defined by
769 specific timestep intervals and corresponding input vectors:
770
771 Baseline (40–69 s): [0, 0, 0, 1, 0, 0, 0, 0]
772 CS (70–89 s): [2, 0, 0, 1, 0, 0, 0, 0]
773 Interval (90–94 s): [0, 0, 0, 1, 0, 0, 0, 0]
774 Open (95–239 s): [0, 1, 0, 1, 0, 0, 0, 0]
775
The recurrent and input weights, bias vector, and the alpha parameter ($\alpha = \Delta t / \tau$) were used to
represent the trained dynamics throughout the epochs.
For each epoch, the neural state trajectories x and corresponding input vector u were
constructed. Initial conditions for fixed point searches were formed by combining all neural
states within the epoch with a randomly sampled subset of global network states (n_trial=
(number of epoch states)+ 512 randomly sampled states; x: n_trial × n_neurons matrix,u:n_trial
x 8 matrix). The input vector u was tiled across trials to match the dimensionality.
Fixed points were then identified jointly via optimization To find fixed point candidates, x and
their one-step forward states F were computed, and the difference between x and F was
minimized in an iterative gradient descent loop, where x was updated using the Adam optimizer.
If change in loss became smaller than a convergence tolerance($1x10^{-4}$) times the learning rate
or if the loss is smaller than tolerance, then the iteration was stopped. The maximum number of
iterations was set to 10,000. We discarded the candidate as an outlier if the distance between a
fixed point candidate x* and the centroid of the initial states was larger than 10.0 times the
average distance between the centroid and each initial state. Redundant candidates were
further merged if the Euclidean distance between any pair of x* was less than $1x10^{-2}$.The
remaining fixed points x* and their Jacobian eigenvalues were collected. Fixed points were
classified based on their dynamical stability: points were labeled as stable (attractors) if all
eigenvalues had absolute magnitudes less than 1.0, and as saddle points otherwise.
795
Trajectory Analysis of Biological Data Using GPFA
For the biological data, z-scored activity pooled from in vivo recordings of nonexploratory cluster
1 neurons was reduced to a three-dimensional space using Gaussian-Process Factor Analysis
(GPFA), with Gaussian smoothing kernel size of 20 bins (implemented using MATLAB code
from Lakshmanan et al. 2015 (50)). Each pseudopopulation trajectory from CS or Neutral rats
was subsequently smoothed with a Gaussian kernel (10-bin window) and visualized. We
employed GPFA because it estimates latent states shared across neurons while explicitly
modeling the noise characteristics of individual neurons, making it well-suited for single-trial
trajectory visualization. Unlike PCA, which identifies orthogonal axes that explain maximal

variance without incorporating temporal structure, GPFA accounts for temporal dynamics,
providing a key advantage in analyzing population activity over time.
Validation was performed by independently applying GPFA to two neuronal subsets derived
from in vivo CS group recordings. Neurons were ranked in descending order based on their
mean z-scored firing rates during the baseline to 5-second post-crossing window. The ranked
list was then divided into two subsets comprising odd- and even-indexed neurons, respectively.
GPFA was applied to each subset using identical parameter settings as described above. For
visualization, neural trajectories were smoothed with a 10-bin Gaussian window. The trajectory
of one subset (subset 1) was chosen as the reference, and the others were aligned to it using
the Procrustes method (MATLAB function). Specifically, anchor points from each trajectory were
mapped to the corresponding anchors in the reference trajectory by minimizing the sum of
squared errors under a linear transformation. The procedure included centering and
normalization, followed by singular value decomposition (SVD) to estimate the optimal rotation,
and finally applying translation and scaling. Anchor points consisted of the trial start, CS onset,
CS offset, door opening, and crossing time.
820
821
822
823

Acknowledgements

This work was supported by the NRF (https://www.nrf.re.kr) of Korea grant funded by the Korean

Government Ministry of Education (https://english.moe.go.kr), Science and Technology (NRF-

2021R1A2B5B03002345), awarded to SC.

Reference

1.   F. C. Crick, C. Koch, What is the function of the claustrum? *Philos Trans R Soc Lond B*
Biol Sci **360**, 1271–1279 (2005).
2.   A. M. Shelton *et al.*, Single neurons and networks in the mouse claustrum integrate input
from widespread cortical sources. *Elife* **13** (2025).
3.   M. G. White *et al.*, Anterior Cingulate Cortex Input to the Claustrum Is Required for Top-
Down Action Control. *Cell Rep* **22**, 84–95 (2018).
4.   G. Atlan *et al.*, Claustrum neurons projecting to the anterior cingulate restrict engagement
during sleep and behavior. *Nat Commun* **15**, 5415 (2024).
5.   C. A. Faig *et al.*, Claustrum projections to the anterior cingulate modulate nociceptive and
pain-associated behavior. *Curr Biol* **34**, 1987–1995 e1984 (2024).
6.   M. Niu *et al.*, Claustrum mediates bidirectional and reversible control of stress-induced
anxiety responses. *Sci Adv* **8**, eabi6375 (2022).
7.   M. Chevee, E. A. Finkel, S. J. Kim, D. H. O'Connor, S. P. Brown, Neural activity in the
mouse claustrum in a cross-modal sensory selection task. *Neuron* **110**, 486–501 e487
(2022).
8.   K. Narikiyo *et al.*, The claustrum coordinates cortical slow-wave activity. *Nat Neurosci* **23**,
741–753 (2020).
9.   M. G. White *et al.*, The Mouse Claustrum Is Required for Optimal Behavioral Performance
Under High Cognitive Demand. *Biol Psychiatry* **88**, 719–726 (2020).

10.   H. Qadir *et al.*, The mouse claustrum synaptically connects cortical network motifs. *Cell Rep* **41**, 111860 (2022).

11.   Y. Goll, G. Atlan, A. Citri, Attention: the claustrum. *Trends Neurosci* **38**, 486–495 (2015).

12.   G. Atlan *et al.*, The Claustrum Supports Resilience to Distraction. *Curr Biol* **28**, 2752–2762 e2757 (2018).

13.   A. Terem *et al.*, Claustral Neurons Projecting to Frontal Cortex Mediate Contextual Association of Reward. *Curr Biol* **30**, 3522–3532 e3526 (2020).

14.   M. B. Madden *et al.*, Serotonin and psilocybin activate 5-HT(1B) receptors to suppress cortical signaling through the claustrum. *Nat Commun* **16**, 7733 (2025).

15.   M. H. Chong, R. Gămănuţ, Anatomical and physiological characteristics of claustrum neurons in primates and rodents. *Frontiers in Mammal Science* **3**, 1309665 (2024).

16.   Q. Wang *et al.*, Organization of the connections between claustrum and cortex in the mouse. *J Comp Neurol* **525**, 1317–1346 (2017).

17.   Y. Han *et al.*, Delayed escape behavior requires claustral activity. *Cell Rep* **43**, 113748 (2024).

18.   C. M. Dillingham *et al.*, The Anatomical Boundary of the Rat Claustrum. *Front Neuroanat* **13**, 53 (2019).

19.   X. Zhang *et al.*, Susceptibility to kindling and neuronal connections of the anterior claustrum. *J Neurosci* **21**, 3674–3687 (2001).

20.   M. M. Jankowski, S. M. O'Mara, Dynamics of place, boundary and object encoding in rat anterior claustrum. *Front Behav Neurosci* **9**, 250 (2015).

21.   M. M. Jankowski, M. N. Islam, S. M. O'Mara, Dynamics of spontaneous local field potentials in the anterior claustrum of freely moving rats. *Brain Res* **1677**, 101–117 (2017).

22.   K. Grasby, A. Talk, The anterior claustrum and spatial reversal learning in rats. *Brain Res* **1499**, 43–52 (2013).

23.   S. Park, K. Sohn, D. Yoon, J. Lee, S. Choi, Single-unit activity in the anterior claustrum during memory retrieval after trace fear conditioning. *PLoS One* **20**, e0318307 (2025).

24.   D. B. Ehrlich, J. T. Stone, D. Brandfonbrener, A. Atanasov, J. D. Murray, PsychRNN: An Accessible and Flexible Python Package for Training Recurrent Neural Network Models on Cognitive Tasks. *eNeuro* **8** (2021).

25.   D. B. Ehrlich, J. D. Murray, Geometry of neural computation unifies working memory and planning. *Proc Natl Acad Sci U S A* **119**, e2115610119 (2022).

26.   R. Kim, T. J. Sejnowski, Strong inhibitory signaling underlies stable temporal dynamics and working memory in spiking neural networks. *Nat Neurosci* **24**, 129–139 (2021).

27.   J. P. Stroud, K. Watanabe, T. Suzuki, M. G. Stokes, M. Lengyel, Optimal information loading into working memory explains dynamic coding in the prefrontal cortex. *Proceedings of the National Academy of Sciences* **120**, e2307991120 (2023).

28.   C. D. Brody, R. Romo, A. Kepecs, Basic mechanisms for graded persistent activity: discrete attractors, continuous attractors, and dynamic representations. *Current opinion in neurobiology* **13**, 204–211 (2003).

29.   W. Chaisangmongkon, S. K. Swaminathan, D. J. Freedman, X.-J. Wang, Computing by robust transience: how the fronto-parietal network performs sequential, category-based decisions. *Neuron* **93**, 1504–1517. e1504 (2017).

30.   K. Mohan, O. Zhu, D. J. Freedman, Interaction between neuronal encoding and population dynamics during categorization task switching in parietal cortex. *Neuron* **109**, 700–712. e704 (2021).

31.   J. P. Stroud, J. Duncan, M. Lengyel, The computational foundations of dynamic coding in
working memory. *Trends Cogn Sci* **28**, 614–627 (2024).
32.   R. Orman, Claustrum: a case for directional, excitatory, intrinsic connectivity in the rat. *J*
Physiol Sci **65**, 533–544 (2015).
33.   J. Kim, C. J. Matney, R. H. Roth, S. P. Brown, Synaptic Organization of the Neuronal
Circuits of the Claustrum. *J Neurosci* **36**, 773–784 (2016).
34.   J. Voigts *et al.*, Spatial reasoning via recurrent neural dynamics in mouse retrosplenial
cortex. *Nat Neurosci* **28**, 1293–1299 (2025).
35.   B. M. Yu *et al.*, Gaussian-process factor analysis for low-dimensional single-trial analysis
of neural population activity. *J Neurophysiol* **102**, 614–635 (2009).
36.   K. C. Lakshmanan, P. T. Sadtler, E. C. Tyler-Kabara, A. P. Batista, B. M. Yu, Extracting
Low-Dimensional Latent Structure from Time Series in the Presence of Delays. *Neural*
Comput **27**, 1825–1856 (2015).
37.   N. M. Timme, C. Lapish, A Tutorial for Information Theory in Neuroscience. *eNeuro* **5**
(2018).
38.   N. Timme, W. Alford, B. Flecker, J. M. Beggs, Synergy, redundancy, and multivariate
information measures: an experimentalist's perspective. *J Comput Neurosci* **36**, 119–140
(2014).
39.   M. D. Golub, D. Sussillo, FixedPointFinder: A Tensorflow toolbox for identifying and
characterizing fixed points in recurrent neural networks. *Journal of open source software*
3, 1003 (2018).
40.   C. D. Brody, A. Hernández, A. Zainos, R. Romo, Timing and neural encoding of
somatosensory parametric working memory in macaque prefrontal cortex. *Cerebral cortex*
13, 1196–1207 (2003).
41.   E. M. Meyers, Dynamic population coding and its relationship to working memory. *J*
Neurophysiol **120**, 2260–2268 (2018).
42.   H. R. Naghavi, J. Eriksson, A. Larsson, L. Nyberg, The claustrum/insula region integrates
conceptually related sounds and pictures. *Neurosci Lett* **422**, 77–80 (2007).
43.   M. M. Reus-Garcia *et al.*, The Claustrum is Involved in Cognitive Processes Related to the
Classical Conditioning of Eyelid Responses in Behaving Rabbits. *Cereb Cortex* **31**, 281–
300 (2021).
44.   D. R. Ollerenshaw *et al.*, Anterior claustrum cells are responsive during behavior but not
passive sensory stimulation. *BioRxiv*, 2021.2003. 2023.436687 (2021).
45.   B. W. Stewart *et al.*, Pathological claustrum activity drives aberrant cognitive network
processing in human chronic pain. *Curr Biol* **34**, 1953–1966 e1956 (2024).
46.   M. Froesel *et al.*, Macaque claustrum, pulvinar and putative dorsolateral amygdala support
the cross-modal association of social audio-visual stimuli based on meaning. *Eur J*
Neurosci **59**, 3203–3223 (2024).
47.   C. Seguin, O. Sporns, A. Zalesky, Brain network communication: concepts, models and
applications. *Nat Rev Neurosci* **24**, 557–574 (2023).
48.   S. Dehaene, M. Kerszberg, J. P. Changeux, A neuronal model of a global workspace in
effortful cognitive tasks. *Proc Natl Acad Sci U S A* **95**, 14529–14534 (1998).
49.   Y. Han *et al.*, Delayed escape behavior requires claustral activity. *Cell Reports* **43** (2024).
50.   K. C. Lakshmanan, P. T. Sadtler, E. C. Tyler-Kabara, A. P. Batista, B. M. Yu, Extracting
low-dimensional latent structure from time series in the presence of delays. *Neural*
computation **27**, 1825–1856 (2015).

Fig. 1

Figure 1 | RNN simulation capable of performing the Delayed Escape Test

(A) Schematic diagram of the Delayed Escape Task. On Day 1, animals were conditioned with a light cue paired with an electric footshock. On Day 2, they were adapted to the test context with only a single crossing permitted. On Day 3, animals were placed in one compartment of the test context and presented with a 20-s CS, followed by a 5-s delay, after which the outlet door was opened to allow escape. The test was conducted only once.

(B) Schematic diagram of the task structure that the RNN model was designed to simulate.

(C) Graphs of crossing latency, measured as the interval from door opening to crossing in rats and as derived from RNN simulations. (*p = 0.0335, unpaired *t*-test; RNN, ****p < 0.0001, Mann–Whitney).

(D) Clustering of all recorded claustral neurons in rats (n = 203).

(E) Mean Z-scored firing rates of non-exploratory Cluster 1 neurons in the CS group (n = 25, rat data). Non-exploratory Cluster 1 denotes the cluster showing increased persistent activity both during CS presentation and thereafter. The CS group corresponds to the CS + door-opening condition in the RNN model.

(F) Average Z-score for each behavioral period shown in E. Bars: Friedman test, ****p < 0.0001; Dunn's multiple comparisons—Base vs Cue, ****p < 0.0001; Base vs Interval, **p = 0.0085; Base vs Early Open, **p = 0.0075; Base vs Late Open, *p = 0.0455; Base vs Post-cross, p > 0.9999.

(G) Correlation between crossing latency and Cluster 1 activity in the CS group (n = 12; Spearman r = 0.6503, *p = 0.0257).

(H) Clustering for simulated neurons (n = 100 neurons).

(I) Mean Z-scored firing rates of Cluster 1 neurons in the CS + door-opening condition (n = 24 neurons, RNN) across all simulated trials.

(J) Average Z-score for each behavioral period shown in I. Bars: Friedman ****p < 0.0001; Dunn's—Base vs Cue, ****p < 0.0001; Base vs Interval, ***p = 0.0001; Base vs Early Open, ****p < 0.0001; Base vs Late Open, **p = 0.0034; Base vs Post-cross, p = 0.7134.

(K) Correlation between crossing latency and Cluster 1 activity in the CS + door-opening condition (147 trials; Spearman r = –0.1628, *p = 0.0488).

(L) Heatmaps of single-neuron activity corresponding to panels E and I. Left: rat data (firing rate, Hz); Right: RNN data (Z-score). Neurons are ordered by overall activity.   (M) Crossing latency in rats with anterior claustrum inhibition during the 5-s period between CS offset and door opening, compared with control virus–expressing animals (left, **p = 0.0079, Mann–Whitney). Corresponding results in the RNN are shown for selective inhibition of Cluster 1 neurons during the same 5-s window simulation (right, ****p < 0.0001, Mann–Whitney).

Panels A, D, E, F, G, and L(left) are adapted from Figs. 1A, 4B, 4D, 4E, and S4E of Han et al. (12), *Cell Reports*, © Elsevier.

A postsynaptic cluster / Presynaptic cluster / Synaptic Weight (A.U)

B 617nm LED / ChrimsonR-expressing cell

C Sagittal slice / fmi / rsCla / ChrimsonR-expressing cell / 1000μm

D aCSF / NBQX+D-AP5 / wash / 50pA / 0.05s

E EPSC Amplitude (pA) / * / aCSF / NBQX + D-AP5

F 617nm LED / ChrimsonR-expressing cell

G Top / Bottom

H EPSC (pA) / Latency (ms) / EPSC / Latency / Top / Bottom

I GCaMP6f-expressing cell / Stimulation 200mA 20Hz 1sec / Stimulation / 2s after stimulation

J Stimulation / Neuron# / Time(s) / ΔF/F

K stimulation / ΔF/F / Time (s)

L NBQX + D-AP5 / GCaMP6f-expressing cell / Stimulation 150mA 20Hz 1sec

M Stimulation aCSF / Neuron# / Time(s) / ΔF/F — stimulation aCSF / ΔF/F / Time (s) — Stimulation NBQX + D-AP5 / Neuron# / Time(s) / ΔF/F — Stimulation NBQX + D-AP5 / ΔF/F / Time (s)

N Stimulation / Neuron# / Time(s) / Z-score — Stimulation / z-score / Time (s) — Stimulation Recurrent inhibition / Neuron# / Time(s) / Z-score — Stimulation Recurrent Inhibition / z-score / Time (s)

Slice   **RNN**

Figure 2 | Comparison of recurrent connectivity in the claustrum and in the RNN simulation
(A) Mean absolute inter-cluster synaptic weight in the RNN.
(B) Schematic representation of whole-cell recordings of ChrimsonR-non-expressing neurons during optogenetic stimulation using LEDs to stimulate ChrimsonR-expressing neurons across the entire optical field.
(C) Confocal image showing the expression pattern of the ChrimsonR virus in sagittal claustrum slices.
(D) Representative EPSCs evoked by 2ms LED pulses before and after pharmacological treatments (red bar indicates stimulation).
(E) Pooled data of EPSC amplitudes (n=7) with statistical analysis using the Wilcoxon matched-pairs signed rank test (*p=0.0156).
(F) Schematic depiction of local stimulation using a digital mirror device (DMD).
(G) Optical stimulation of each divided part produces variable amplitudes of EPSCs in whole-cell patched neurons expressing no ChrimsonR. Heatmap showing EPSC amplitudes upon stimulation of the designated part of optical field. Please note that when stimulating closer to the recorded neuron, larger amplitude EPSCs were induced.
(H) The graph displays EPSC amplitudes and latencies in the region marked with dashed lines on the heatmap shown in G.
(I) Left: schematic representation showing an experimental configuration in which brief electrical stimulation produces a persistently enhanced activity in rsCla slices. Right: representative images of GCaMP6f fluorescence changes immediately before and 2 seconds after stimulation.
(J) Heatmap of fluorescence changes for individual puncta in the slice shown in I.
(K) Population-averaged fluorescence trace of all puncta in J (mean ± SEM).
(L) Schematic representation showing an experimental configuration in which effects of the blockers for AMPA/NMDA receptors on the persistent response were examined.
(M) Calcium imaging before and after an aCSF puff (top panels) and an NBQX + D-AP5 puff (bottom panels): heatmaps (left) and population-averaged traces (right).
(N) RNN analogue: heatmaps (left) and averaged Z-scores (right) for Cluster 1 neurons following brief excitation (top panels) and after the addition of recurrent inhibition (bottom panels).

Fig. 3

Figure 3 | RNN PCA Trajectories.
(A-C) Time-normalized, trial-averaged PCA trajectories for each simulation condition. All trajectories begin at the Start event (black circle), proceed through CS onset and offset (CSon & CSoff; green circle) in the CS-containing and inhibition simulations, then through the Open event (orange circle) in the Open-containing and inhibition simulations, and terminate either at the Cross event (red circle) for crossing conditions or at the End event (black circle) for the CS-only condition.
(A) PCA trajectory for the CS+door-opening condition. The mean latency from door opening to crossing was 44.16 ± 0.9583 s.
(B) PCA trajectory for the door-opening only condition.
(C) PCA trajectory for the CS only condition.
(D) Simulated inhibition applied during a 5-s interval between CSoff and door-opening. Top: schematic of the simulated task. Bottom: time-normalized, trial-averaged PCA trajectory.
(E) Simulation with a 180-s interval between CSoff and door-opening (no inhibition applied). Top: schematic of the simulated task. Bottom: time-normalized, trial-averaged PCA trajectory.
(F) Schematic of the trajectory-combination model: the predicted CS + door-opening trajectory (purple line)—obtained by model of which input are the CS only and door-opening only trajectories—is plotted against the actual CS + door-opening trajectory (orange line).
(G) Model-fit comparison for cluster 1: difference in residual sum of squares (ΔRSS) between the linear regression and MLP models, normalized to the mean RSS of the linear model. Bar colors denote condition (CS + door-opening = green, inhibition = red, 180-s interval = gray). One-way ANOVA with Holm–Sidak's multiple comparisons test: CS + door-opening vs. inhibition, **p = 0.0063; CS + Open vs. 180-s interval, ****p < 0.0001. Bars show mean ± SEM. Solid lines represent mean PCA trajectories; shaded areas denote SEM.

Fig. 4

Figure 4 | Biological Neural GPFA Trajectories and Quantification of Local Geometry.

(A) GPFA trajectories of in vivo single-unit recording data: (left) CS rats, (right) neutral rats (n = 25 neurons, 16 rats). The mean latency from door opening to crossing was 56.363 ± 8.22 s.

(B) Validation of GPFA trajectories in CS group recordings. Neurons were ranked in descending order of mean z-scored firing rates (baseline to 5-s post-crossing) and split into odd- and even-indexed subsets. The trajectory from subset 1 (left; n = 13 neurons, 6 rats) served as the reference, while the trajectory from subset 2 (right; n = 12 neurons, 10 rats) was aligned accordingly. For clarity, only the segment from door opening to crossing is displayed. The mean latency from door opening to crossing for subset 1 was 53.306 ± 14.857 s. The mean latency from door opening to crossing for subset 2 was 50.969 ± 7.417 s.

(C) Third eigenvalue divided by the sum of all eigenvalues (i.e., the proportion of total three-dimensional variance explained by PC3), computed using sliding windows from the post-open RNN trajectory, comparing the CS+Open (green; n = 20 windows) and Open-only (gray; n = 20 windows) conditions. **** Unpaired t test, p < 0.0001.

(D) $\lambda_3/\Sigma\lambda$ of the biological neural trajectory within the sliding window, comparing the CS (green) and Neutral (gray) group pseudopopulations. $\lambda_3/\Sigma\lambda$ (the third eigenvalue divided by the sum of all eigenvalues) was computed within a sliding window of 10 time bins across the 20 post-open time bins of the averaged PCA trajectory (RNN) or the GPFA trajectory (biological pseudopopulation). **** Mann–Whitney test, p < 0.0001.

(E) Neuron-wise contribution to the mean $\lambda_3/\Sigma\lambda$, defined using a leave-one-neuron-out analysis of the biological neural trajectory, comparing CS (green; n = 25 neurons) and Neutral (gray; n = 17 neurons). For the leave-one-neuron-out analysis, the GPFA trajectory was recomputed separately for each biological group after excluding one neuron at a time. For each neuron, its contribution was defined as the difference between the full-population mean $\lambda_3/\Sigma\lambda$ and the leave-one-out mean $\lambda_3/\Sigma\lambda$. The mean $\lambda_3/\Sigma\lambda$ was calculated by averaging $\lambda_3/\Sigma\lambda$ across sliding windows. ** Mann–Whitney test, p = 0.0012.

Raw data from Fig. 4 of Han et al. (12), *Cell Reports*, were reanalyzed using GPFA. © Elsevier.

Fig. 5

Figure 5 | Decoder Accuracy, and Partial Information Decomposition of Cluster 1 RNN Neurons under different simulation conditions.
(A) Left: Schematic of the CS decoder. A classifier was trained to distinguish CS + door-opening trials from door-opening only trials using cluster 1 neuron activity. Right: CS decoding accuracy (%) across time bins.
(B) Left: Schematic of the door-opening decoder. A classifier was trained to distinguish CS + door-opening trials from CS only trials using cluster 1 neuron activity. Right: Door-opening decoding accuracy (%) across time bins.
(C) Conceptual diagram of Partial Information Decomposition. Mutual information about CS and door-opening is decomposed into CS-unique information, door-opening-unique information, redundancy, and synergy.
(D) Schematic of the PID analysis. CS decoding accuracy and door-opening decoding accuracy were computed for each trial type (CS + door-opening, CS only, door-opening only, and None), and used as input variables for neuron-wise PID. PID terms were then averaged across neurons.
(E) Synergy information of cluster 1 neurons.
(F) Comparison of synergy between inhibition and no-inhibition simulations.
(G) Comparison of synergy between the 180-s interval and the 5-s interval simulations. Green line: 5-s interval; green shaded box indicates CS presentation window. Gray line: 180-s interval. Solid lines represent means; shaded areas indicate SEM.
(H) Model fit comparison between cluster 1 RNN neurons with high synergy and those with low synergy. The difference in residual sum of squares (ΔRSS) between linear regression and MLP models is shown, normalized to the mean RSS of the linear model. Bar colors indicate condition (high synergy = green, low synergy = gray). (High synergy: n = 147 trials; Low synergy: n = 147.) Mann-Whitney test: high vs. low synergy, ****$p < 0.0001$. Bars show mean $\pm$ SEM.
In this figure, Open denotes door-opening.

Fig. 6

Figure 6 | Trajectory Coding Hypothesis and Biological Neural Trajectories.
(A) Cross-temporal decoding of integration-specific information from cluster 1 RNN neurons.
(B) Cross-temporal decoding of integration-specific information from cluster 1 RNN neurons with high synergy.
(C) Cross-temporal decoding of integration-specific information from cluster 1 RNN neurons with low synergy.
(D) Left: Heatmaps of raw firing rates for RNN neurons from a representative CS + door-opening trial. Right: Heatmaps for biological cluster 1 non-exploratory neurons in the CS group during the delayed escape task (right). Neurons in the right heatmap are ordered by overall activity.
(E) Hypotheses on dynamic coding. Left: Attractor hypothesis: Multiple or continuous stable converging points exist, and changes in input cause neural dynamics to move from one point to another, thereby encoding varying states. Right: Trajectory hypothesis: Neural dynamics are not, or are only minimally, related to attractors (which may not exist); instead, changes in input cause changes in trajectory, which encode varying states.
(F) Fixed-point analysis of a representative trial in an RNN. Green dots represent stable fixed points (CS period), while black and yellow X-marks denote unstable fixed points (interval and open periods).
Panel D (Right) are adapted from Fig. S4E of Han et al. (12), Cell Reports, © Elsevier.

Figure S1 | RNN activity of Clusters 1–3 under CS and door-opening only conditions

(A) **Cluster 1, door-opening only.** Left, Z-scored firing rate of Cluster 1 neurons (n = 24) in a representative trial. Middle, average Z-score for each behavioral period across all trials; Friedman test, ****$p < 0.0001$; Dunn's post hoc—Base vs Cue, ****$p < 0.0001$; Base vs Interval, ***$p = 0.0001$; Base vs Early Open, ****$p < 0.0001$; Base vs Late Open, **$p = 0.0034$; Base vs Post-escape, $p = 0.7134$. Right, Spearman correlation between crossing latency and Cluster 1 activity (n = 147; r = 0.1143, p = 0.2388).

(B) **Cluster 2, CS.** Left, Z-scored firing rate of Cluster 2 neurons (n = 38) in a representative trial. Middle, average Z-score for each behavioral period across all trials; Friedman ****$p < 0.0001$; Dunn's—Base vs Cue, ***$p < 0.003$; Base vs Interval, ****$p < 0.0001$; Base vs Early Open, **$p = 0.0072$; Base vs Late Open, ***$p = 0.0002$; Base vs Post-escape, ****$p < 0.0001$. Right, correlation between crossing latency and Cluster 2 activity (n = 147; r = 0.0618, p = 0.4573).

(C) **Cluster 2, door-opening only.** Left, Z-scored firing rate of Cluster 2 neurons (n = 38) in a representative trial. Middle, average Z-score for each behavioral period across all trials; Friedman ****$p < 0.0001$; Dunn's—Base vs Cue, $p > 0.9999$; Base vs Interval, $p > 0.9999$; Base vs Early Open, ***$p = 0.0009$; Base vs Late Open, ****$p < 0.0001$; Base vs Post-escape, ****$p < 0.0001$. Right, correlation between crossing latency and Cluster 2 activity (n = 147; r = –0.0717, p = 0.4607).

(D) **Cluster 3, CS.** Left, Z-scored firing rate of Cluster 3 neurons (n = 38) in a representative trial. Middle, average Z-score for each behavioral period across all trials; Friedman ****$p < 0.0001$; Dunn's—Base vs Cue, $p > 0.9999$; Base vs Interval, $p > 0.9999$; Base vs Early Open, ****$p < 0.0001$; Base vs Late Open, ****$p < 0.0001$; Base vs Post-escape, ****$p < 0.0001$. Right, correlation between crossing latency and Cluster 3 activity (n = 147; r = 0.3335, ****$p < 0.0001$).

(E) **Cluster 3, door-opening only.** Left, Z-scored firing rate of Cluster 3 neurons (n = 38) in a representative trial. Middle, average Z-score for each behavioral period across all trials; Friedman ****$p < 0.0001$; Dunn's—Base vs Cue, $p = 0.2867$; Base vs Interval, $p > 0.9999$; Base vs Early Open, $p > 0.9999$; Base vs Late Open, ****$p < 0.0001$; Base vs Post-escape, ****$p < 0.0001$. Right, , correlation between crossing latency and Cluster 3 activity (n = 147; r = 0.1861, p = 0.0539).

Fig. S2

Figure S2 | Contribution of each cluster to the output during the delayed escape test
Relative contributions of Clusters 1–3 to the network output across behavioral periods in the delayed escape test.

Fig. S3

Figure S3 | RNN activity of Clusters 2 and 3 under inhibition and escape latency under 180-s delay condition

(A) Schematic of the inhibition-simulation protocol. In the RNN model, inhibition was applied for 5 s during the delay interval after CS offset and before the door-opening signal.

(B) Bar graphs showing crossing latency under inhibition of Cluster 2 or Cluster 3 neurons.: Cluster 2, **p = 0.0023; Cluster 3, p = 0.1511 (both Mann–Whitney).

(C) Cross-latency under a prolonged 180-s CS–door-opening interval. Bar graphs show crossing latencies in rats (left) and in the RNN model (right): rats, p = 0.9720 (unpaired t-test); RNN, p = 0.3574 (unpaired t-test).

Fig. S4

Figure S4 | Slice physiology and pharmacological manipulation of claustral persistent activity

(A) Membrane-potential decay τ during whole-cell recordings.

(B) Top: voltage trace showing persistent spiking after 20 Hz, 200 μA, 1-s stimulation; Bottom: corresponding GCaMP6f fluorescence changes recorded from the same cell.

(C) Relationship between the distance from the stimulating electrode to the recorded cell and the persistence ratio (peak fluorescence / fluorescence at 10 s), calculated from the dataset shown in Fig. 2J.

(D) Schematic of pressure injection of drugs onto whole-cell patch-clamped neurons in horizontal claustral slices.

(E) Representative EPSCs recorded under control (aCSF, no pressure injection) and after NBQX + D-AP5 pressure injection with the drug pipette positioned 30 μm or 70 μm from the recorded cell. EPSCs were evoked by electrical stimulation at 0.5 s and 3.5 s after injection. (F) Decay kinetics measured from the graphs in Fig. 2M and N under NBQX + D-AP5 or aCSF treatment. Left, decay τ after application of NBQX + D-AP5 (n = 16) or aCSF (n = 12) 10 s after electrical stimulation; **p = 0.0012, Mann–Whitney. Right, decay τ with versus without recurrent inhibition in the RNN (n = 24 each); ****p < 0.0001, Mann–Whitney.

Fig. S5

Figure S5 | PCA trajectories of Clusters 2 and 3 under various simulation conditions

(A–F) Cluster 2: time-normalized, trial-averaged PCA trajectories for (A) CS + door-opening, (B) door opening only, (C) CS only, (D) overlay of A–C, (E) inhibition simulation, and (F) 180 s delay between CS offset and door-opening.
(G–L) Cluster 3 trajectories arranged as in panels A–F.

Fig. S6

Figure S6 | Model-fit comparison across clusters

(A) Residual sum of squares (RSS) from a linear regression model and a multilayer perceptron (MLP) across normalized time bins. Data are taken from the Open event to the Cross event (for the CS only case, from the open event to the mean cross latency of the CS + door-opening condition). Green line: RSS of the linear regression model; blue line: RSS of the MLP.

(B–D) Difference in residual sum of squares (ΔRSS) between linear regression and MLP fits, normalized by the mean RSS of the linear model, across conditions.; bars, mean ± SEM.

(B) Cluster 1: CS + door-opening vs inhibition, **p = 0.0063; CS + door-opening vs 180 s delay, ****p < 0.0001.

(C) Cluster 2: CS + door-opening vs inhibition, p = 0.6735; CS + door-opening vs 180 s delay, **p = 0.0021.

(D) Cluster 3: CS + door-opening vs inhibition, p = 0.4067; CS + door-opening vs 180 s delay, ****p < 0.0001 (one-way ANOVA with Holm–Šidák).

Fig. S7

Figure S7 | Decoding performance and PID analysis across clusters

(A) Decoding accuracy for classifying CS-present versus CS-absent trials using cluster-specific decoders.
(B) Decoding accuracy for classifying Open-present versus Open-absent trials using cluster-specific decoders.
(C–E) PID metrics (CS-unique, Open-unique, synergy) for neurons in each cluster.
(F–J) Same analyses as in panels A–E under the inhibition condition.
(K–O) Same analyses as in panels A–E under a 180 s CS–Open interval.

Fig. S8

Figure S8 | PCA trajectories of high- and low-synergy neurons

(A–B) Time-normalized, trial-averaged PCA trajectories for (A) high-synergy and (B) low-synergy Cluster 1 RNN neurons.