Lower engagement of cognitive control, attention, modulation networks and lower creativity in children while using ChatGPT: an fMRI study

Tzipi Horowitz-Kraus[1,2,3,4,5,6], Yoed N. Kenett[3,7], Dafna Link[2,3], Emeel Ashqar[2,3], Rola Farah[1]

[1] Educational Neuroimaging Group, Faculty of Education in Science and Technology, Technion – Israel Institute of Technology, Haifa, Israel;

[2] Faculty of Biomedical Engineering, Technion – Israel Institute of Technology, Haifa, Israel

[3] May-Blum-Dahl Technion Human MRI Research Center (TecHMRC);

[4] Kennedy Krieger Institute, Baltimore, MD, USA;

[5] Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, MD, USA

[6] Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, MD, USA

[7] Faculty of Data and Decision Sciences, Technion – Israel Institute of Technology, Haifa, Israel

Corresponding author:

Tzipi Horowitz-Kraus, PhD
Educational Neuroimaging Group, Faculty of Education in Science and Technology, Technion – Israel Institute of Technology, Haifa, Israel; Faculty of Biomedical Engineering, Technion – Israel Institute of Technology, Haifa, Israel
Email: Tzipi.kraus@technion.ac.il. Tel: +972-522-989298

Highlights

- Children demonstrate a lower within-network connectivity in the frontoparietal, salience, and dorsal attention systems while using ChatGPT compared to adults.

- Reduced between-network integration in cognitive control and modulation networks (frontoparietal–salience and frontoparietal–dorsal) in children vs adults while using ChatGPT.

- In adults, but not in children, higher creativity was associated with stronger connectivity within cognitive control networks.

Abstract

Generative AI can scaffold idea generation, but how it engages control, attention, and memory systems in the developing versus mature brain, and how this relates to creativity, remains unclear. We compared 6-7-year-old (N = 15) children and adults (N = 16) during a naturalistic 5-minute co-creative dialogue with ChatGPT using the same prompt, in-scanner conversation, undergoing functional MRI scanning and their dialogues were recorded. Functional connectivity (Fisher's z) was computed within and between seven large-scale networks: cingulo-opercular, default mode (DMN), memory retrieval, frontoparietal (FP), salience, ventral attention (VAN), and dorsal attention (DAN). Creativity scores, assessed outside of the scanner, did not differ between groups. Adults showed stronger within-network connectivity in control/attention systems, most robustly in FP, with additional effects in salience and DAN. Adults also exhibited greater cross-network integration, particularly FP-salience and FP-DAN couplings, alongside broader adult-greater effects among memory- and attention-related pairs. During a co-creative AI-assisted conversation, children exhibit a less coherent and integrated control–attention–memory architecture than adults. The combination of similar creativity scores with divergent neural profiles suggests that developmental differences in the implementation of creative cognition may be more pronounced than baseline capacity. Our findings highlight FP-centric pathways and their salience-gated interactions as candidate substrates by which creativity supports goal-directed co-creative dialogue with AI.

Keywords: Generative AI, Children, ChatGPT, Conversation, Development, Functional connectivity, Creativity, fMRI, Cognitive control

Introduction

Digital tools have become deeply embedded in children's daily lives worldwide. Research shows that these technologies, ranging from tablets and laptops to smartphones and emerging generative AI, have a multifaceted impact on children's learning, well-being, and development[1-7]. While digital tools may offer substantial opportunities to enhance educational experiences, they also introduce challenges such as distraction, cognitive overload, social-emotional strain, and negative impact on literacy development [1].

In many countries, fostering digital competence and inclusion is a strategic priority, with a vision that this implementation will promote a resilient, innovative, knowledge-based society and will spark creativity. Technology can support learning, but it cannot lead it. Digital tools are not a substitute for educators; they must be used in partnership with them. Children learn best when accompanied by someone who guides, encourages, challenges, and believes in them, a feeling often evoked by generative AI tools[8]. However, how much children are actually engaged cognitively and creatively when interacting with Chat-GPT is yet to be known, which is the topic of the current study.

Generative AI use by children

Children's interactions with generative AI (GenAI) are emerging as both opportunities for creativity and challenges for critical engagement. Experimental studies have shown that tools like ChatGPT can support young children's storytelling by providing interactive, adaptive dialogue [8]. For example, a study of 5–6-year-olds found

that children enjoyed storytelling with ChatGPT, which provided more frequent positive feedback and open-ended prompts than parents, suggesting potential for supporting children's creative expression when parental time is limited [8]. Similarly, interventions using games such as *Bot VoyAlge* engaged 10–12-year-olds in critically examining ChatGPT's responses, revealing that children could identify gender, cultural, and age biases embedded in its outputs [9]. These findings highlight that while children can find GenAI engaging and even preferable in certain contexts, structured support is necessary to help them navigate its biases and limitations.

Beyond creative engagement, researchers emphasize the developmental, societal, and ethical implications of children's growing exposure to GenAI. Cross-national work applying a child rights framework has underscored the need to prioritize children's voices and experiences in shaping AI governance, pointing to rights such as privacy, safety, participation, and development that are often overlooked in current policy debates [10]. Broader reviews caution that AI may reshape learning, attention, and social interactions, particularly during sensitive developmental windows, requiring careful regulation and proactive safeguards [11]. Educational interventions that build AI literacy, such as teaching middle schoolers to recognize deepfakes and misinformation, show promise in fostering critical awareness of GenAI's socio-technical dimensions [12]. Collectively, these studies demonstrate that children can be both active creators and critical evaluators of GenAI, but ensuring safe, equitable, and developmentally appropriate use will require interdisciplinary collaboration and child-centered policy.

Many fields, including the field of creativity research, are now experiencing and contending with the rise of AI models such as ChatGPT [13] which are becoming increasingly popular tools in many linguistic Human-Computer Interactions. The emergence of publicly accessible AI models such as ChatGPT has given rise to global conversations on the implications of AI capabilities and the potential for co-creative cooperation between human and machine agents [14,15]. Co-creativity is generally defined as multiple parties contributing to a creative process in a blended manner [16]. Co-creativity is becoming especially relevant in the age of generative AI models [14], where highly advanced generative AI can now begin to match humans in creativity tests [15]. However, increasing evidence highlights not only the benefits [17-20], but also the potential costs of AI on human creativity [21-25]. Thus, elucidating potential differences between children and adult co-creative interactions is greatly needed, and especially elucidating the neural mechanisms that relate to such potential differences in co-creative acts. This, as children's development continues to shape large-scale neural wiring and may be affected by the rapid AI technological immersion.

Brain development

A recent study utilizing Electroencephalogram (EEG) examined the cognitive engagement in adult individuals who were instructed to write an essay under several conditions: while using ChatGPT, search engines, or without any assistance. Another condition allowed those who used ChatGPT to write without any assistance, and those who wrote without any assistance were allowed to use ChatGPT for writing [26]. EEG coherence data demonstrated a cognitive underengagement in ChatGPT users: For

writing assignments, individuals were found to engage fewer cognitive resources than individuals who were using other tools to gain knowledge (such as Google) [26]. Adults, however, have well-developed cognitive, attention, and modulation systems [27-29]. Neurobiologicaly, brain networks supporting these cognitive control abilities include the frontoparietal and congulo-opercular (Executive Functions; EF system), Dorsal and Ventral Attention Networks (DAN, VAN, bottom-up attention networks), and the modulation system (Default Mode Network; DMN, and the Salience network) [30],reach their full developmental maturation at the age of 26 years [31-33]. The long-range connections and the efficiency of communication between these networks increase markedly through development, reaching maturity in adulthood [34]. While generating new ideas, as happens when individuals communicate or when they speak to generative AI, there is a need to plan, inhibit, and remember what was already discussed (i.e., working memory)- all parts of EF, which are not yet mature among young 6-year-old children[35]. However, creativity is also important for generating new ideas [36]. What are the neural mechanisms that support the involvement of EF during a co-creative task such as an interaction with ChatGPT, how do they differ in children versus adults, and how might they be impacted with relation to interaction with ChatGPT?

Creativity in children

Creativity is broadly defined as the ability to generate ideas that are both novel and useful within a given social context [37-39]. This construct relies on cognitive processes, including divergent thinking (DT), which encompasses fluency, originality, and flexibility [40,41], memory [42,43], cognitive control [44], attention [45], and associative

thinking [46]. Developmental perspectives emphasize that creativity is not a fixed trait but an emergent process, with executive functions (EF) providing regulatory mechanisms to support idea generation and sustain attention [47].

However, empirical findings suggest that young children's DT may rely heavily on bottom-up associative processes, with EF sometimes playing an inverse or constraining role [41]. Critically, current research focusing on the creative process emphasize dull-process theories, mainly those including creative idea generation and idea evaluation stages [42,48-50]. A growing scientific focus regards creative idea evaluation, and how it is related to metacognitive capacities, which develop later in life [51]. Longitudinal studies show substantial gains in fluency and originality between ages 4 and 6, supported by both memory retrieval and mental operations, but also highlight a potential "slump" in creativity when children enter first grade and encounter more formalized, convergent instructional practices [40].

The assessment of creativity has often relied on DT tasks such as the Alternative Uses Task (AUT), which capture both product-level measures (e.g., fluency and originality) and process-level mechanisms underlying idea generation [52]. These tasks are advantageous in their sensitivity to developmental change and in revealing not only the quantity but also the quality of children's ideas. Complementary measures of evaluative ability are also necessary, as they assess whether children can judge the originality or appropriateness of their own ideas [52]. Such current assessment tools are based on computational methods such as large language models to provide automatic and quantitative neasures of originality and appropriateness of responses in open-ended creativity tasks such as the AUT [53-56].

Contemporary accounts highlight that creativity is shaped by social interaction and cultural context, positioning it as a developmental process embedded in children's everyday environments [37,57]. While studies using AI with children demonstrate that generative models can scaffold storytelling and imaginative play, the extent to which such tools enhance or constrain DT remains insufficiently understood, and whether cognitive processes are engaged equally as in adults further requires empirical work.

The neuroscience of creativity

Over the past few decades, extensive neuroscientific research has been conducted to elucidate the neural mechanisms that realize creative thinking [58,59]. Such research has been conducted via multiple neuroscientific methods [60], and extensively via functional MRI research to elucidate cuntional brain connectivity that supports creativity [61-65].

Functional magnetic resonance imaging (fMRI) has significantly advanced the understanding of creativity by highlighting how distributed brain networks support creative cognition. Recent studies have emphasized functional interactions among large-scale brain systems composed of multiple regions. This approach has identified several key networks—specifically, the EF system, and the modulation one (DMN, and salience network)- whose coordinated activity underlies various facets of creative thinking [63].

The DMN, as part of the modulation system, comprising regions such as the medial prefrontal cortex, posterior cingulate cortex, angular gyrus, and hippocampus— has emerged as central to creative cognition due to its role in memory retrieval and

associative thinking [66]. The DMN supports memory processes essential to creative ideation, especially remote associative thinking—the ability to link conceptually distant ideas [46,67,68]. Beyond idea generation, the DMN also contributes to evaluating ideas by comparing them to stored knowledge to determine novelty and by mentally simulating their feasibility to assess appropriateness [42]. The salience network is comprised of the anterior insula and dorsal anterior cingulate cortex—orchestrates interactions between the DMN and the EF system, dynamically transitioning between generative and evaluative modes [61,63,64]. Supporting this gating function, multilayer network analyses demonstrate extensive reconfiguration among the EF system, DMN, and SN during creative idea generation, while evaluation engages a more integrated yet stable network architecture [69].

Alongside the modulation system, the EF networks —comprising primarily lateral prefrontal and posterior parietal cortices—provides top-down regulation to steer creative thinking toward task goals, such as response inhibition and idea revision that are critical for refining creative ideas. The EF system is particularly involved in suppressing dominant but less original responses, enabling individuals to consider novel, more creative alternatives [68]. Such cognitive control engagement tends to vary with creative task demands [70], with flexible down-regulation often facilitating originality when dominant responses must be overcome [44]. Creative thinking is characterized by heightened cooperation among these networks, with stronger coupling between the DMN and EF consistently predicting higher creative performance [61,63]. Connectome-based predictive modeling reliably links enhanced functional connectivity across DMN, EF, and SN regions to individual differences in creative ability [62,65,71].

Highly creative individuals exhibit greater network flexibility, showing more frequent

transitions between the DMN and EF system [64] and increased temporal variability,

especially within the DMN [72]. How critical is the involvement of these cognitive control

networks for the realization of creativity potentially impacted by interaction with

ChatGPT, and even more differentiated in children and adults in the question of the

current study.

The current study

The current study aimed to determine the differences in cognitive engagement

when children and adults are engaged while communicating with ChatGPT during a co-

creative task. The study also examined whether the level of creativity was related to the

level of cognitive control engagement. We hypothesized that children would show a

lower engagement of cognitive control networks and that a connection between

creativity and cognitive control functional connectivity would be observed in adults, but

not in children.

Methods

Participants

Fifteen-6-7 year-old children, all first graders (mean age: 6.44 years, SD=4

months, 11 maels) and 16 adults (mean age: 38.12 years, SD=7.25 years, 4 maels)

from an average-above-average socioeconomic status participated in the current study.

For both age groups, the following inclusion criteria were applied: (1) intact vision and

hearing, (2) no history of psychiatric or neurological delays (3) intact non-verbal and

verbal IQ (>85), (4) born term (>37 weeks gestational age), (5) no contradiction to MR

scanning: claustrophobia, any metal object in the body, including cardiac pacemaker,

dental braces, and cochlear implants. Participants underwent a comprehensive

behavioral assessment of their EF and creativity abilities, as well as an fMRI scan.

Children and their parents signed informed assent and consent forms, respectively, and

adults signed written consents before participating. Data was collected at the Technion

Human MRI research center (TechMRC) at the Technion. The protocol was approved

by the Ministry of Health Helsinki committee (for children) and by the institutions' ethical

committee at the Technion. Participants were compensated for their participation.

Study procedure

Once children and adults were recruited and consented to participate in the

study, they were invited to the Educational Neuroimaging Group (ENIG) facility at the

Technion, where their cognitive and creative abilities were assessed. Then, upon

completion, families were invited to the MRI facilities at the Technion, where ChatGPT

was set up on the MRI console, and the child's voice and all communication were

recorded. All conversations with ChatGPT were recorded and analyzed for semantic

complexity, a factor previously linked to creativity (following [73]).

Behavioral measures

To verify intact nonverbal IQ, children completed the Test of Nonverbal

Intelligence (TONI [74]) which includes a completion of missing parts in given matrices. To

evaluate language abilities, the vocabulary test from the Peabody Picture Vocabulary

Test (PPVT [75]) was used. For adults, equivalent tests were administered: the Matrix

task and the vocabulary task from the Wechsler Adult Intelligence Scale (WAIS) [76].

Creativity was assessed using the divergent association task (DAT; [77],

https://www.datcreativity.com/task), conducted outside of the scanner. In this task,

participants were required to retrieve 10 words that are as unrelated as possible,

without any time constraints. The semantic relations between the words were calculated

with a higher value representing higher creativity abilities (more divergent thinking).

Neuroimaging measures

Talking to ChatGPT paradigm

The fMRI data were collected using a Prisma Magneton 3 Tesla scanner

(Siemens Healthineers, Erlangen, Germany). A unique infrastructure set-up at the

TechMRC allowed the data collection during the communication wth ChatGPT: an

internet connection to the audio-visual system in the scanner, a real-life speech

recording while fMRI data is collected using Optoacoustics system (Optoacoustics Ltd.,

Mazor, Israel), an in-house noise reduction system developed at the Technion allowing

a clear sound transmission of the ChatGPT audio stimuli, and another in-house

developed algorithm which allows a noise elimination from the audio recording allowing

a direct transcript of the verbal output of the child. After desensitization and acclimation

to the scanner environment (following our previous experience in this age group [78]),

anatomical data (T1) was collected for coregistration purposes. Upon connecting to

ChatGPT, children were informed that they were about to speak to the "smartest

computer in the world," also known as ChatGPT, whereas adults were simply told that

they would be speaking to ChatGPT (see prompts in the supplemental materials). Both groups were told that their goal is to open a store; It can be any store they would like, it can be anywhere, for anyone. They need to consult with ChatGPT regarding the type of store for 5 minutes while the child speaks to ChatGPT. Note, the instruction was given only when they were in the scanner, immediately before the start of the scan. An audio recording of the entire conversation was conducted via Audacity software (https://www.audacityteam.org/) and the transcript of the conversation was generated using ChatGPT.

Neuroimaging data

Each fMRI session lasted 5 minutes exactly and included a Time Repetition (TR) of 1 sec and the echo time (TE) was 28 milliseconds. The imaging setup included a field of view (FOV) of 20 × 20 × 13.6 cm, with a 100 × 100 matrix and slices of 2 mm thickness. High-resolution T1 images were collected for each participant to align the functional images accurately. The T1 scan parameters were a TR of 1750 milliseconds, a TE of 2.8 milliseconds, an inversion time of 900 milliseconds, and a flip angle of 9°, with a field of view (FOV) of 22.4 × 22.4 × 16 cm, a 224 × 224 × 160 matrix, and a slice thickness of 1 mm. All scans included a total of 300 volumes per session.

Neuroimaging data processing

The fMRI data were corrected for geometrical distortion and signal intensity variation due to static B0 field variations in T2* weighted fMRI data using corresponding B0 field map images acquired in every participant. Rigid boundary-based registration [79]

to a high-resolution 3D, T1-weighted anatomical volume acquired within the same

session, was performed. Nonlinear warping of this anatomical volume into standard

space using a pediatric template was performed using ANTs [80] [81]. Functional MRI data

were subjected to the same preprocessing strategy outlined in [82-88]. Using a model-

based approach, we constructed a model for the functional connections among brain

regions involved in the reading network [89-95]. This included the export of functional

connectivity values for the selected networks, as well as EF, attention and modulation

networks (CO, FP, memory, DAN, VAN, DMN, and Salience [96]) using the CONN toolbox

implemented in SPM 12 ([97]; http://www.nitrc.org/projects/conn/). Functional connectivity

between brain regions was quantified by the temporal correlation between their

spontaneous BOLD signal fluctuations [98] based on Power's atlas [99] containing 264

putative functional parcels, covering cognitive, sensory, and language/auditory

functional networks (see Figure 1 for the networks spatial masks). To control for motion

artifacts, functional connectivity analysis included a rigorous quality control (QC)

process, which involved calculating the mean framewise displacement (FD) by summing

the absolute values of the derivatives of the translational and rotational realignment

estimates (after converting the rotational estimates to displacement at a 50mm radius).

The QC vector was correlated across participants and conditions with the pairwise

functional connectivity correlations between all parcels to ensure that motion artifacts do

not drive the connectivity (see [100]). None of the volumes reached FD > 0.5mm, and

hence segments were not removed from the data (per the acceptable definition of data

lasting fewer than five contiguous volumes were flagged and removed; [100,101]). In

addition, motion parameters were included as covariates in the first-level analyses. Due

to significant difference in FD in children (X = .306, SD = .06) and adults (X = .226, SD = .05), t(30) = 3.65, *p* < .001), FD was controlled for in all analyses involving neuroimaging data.

Figure 1. Executive functions, attention and modulation networks' masks

Divergent Sematic Integration (DSI) generated from the conversations with ChatGPT

DSI calculates the average semantic distance between all word pairs in a response using embeddings from a pre-trained model and cosine similarity. DSI has been shown to significantly predict originality on other creative tasks involving long-form text responses [102,103]. Semantic-distance models work by placing words, sentences, or questions into a shared "semantic space," where the distance between them reflects how similar or different their meanings are [56]. For example, cat and dog will appear

closer together in this space because they share many contextual similarities, while cat

and democracy will be much farther apart. This is done in an unsupervised manner

meaning they learn patterns of word co-occurrence and meaning directly from large text

corpora without relying on predefined human ratings or labeled training data [46]. DSI

scores were computed for each participant for their entire discussion with ChatGPT

based on their narrative transcripts. This was done to ensure enough text in each

narrative to reliably compute DSI score. Thus, each DSI score represents the general

diversity, or richness, of the conversation with ChatGPT. A higher score denotes a

richer, more original narrative interaction with ChatGPT.

Statistical analyses

Behavioral data analysis

Several *t*-test analyses were conducted to determine the differences in creativity

levels assessed inside (DSI) and outside (DAT) of the scanner, among children and

adults.

Neuroimaging data analysis

To determine differences in within- and between-network functional connectivity

between children and adults, a multivariate analysis of variance (MANOVA) was

conducted with group (children, adults) as a between-subjects factor and framewise

displacement (FD**)** as a covariate to control for head motion. Dependent variables were

mean Fisher-z-transformed connectivity values within seven canonical large-scale brain

networks: cingulo-opercular (CO), default mode (DMN), memory retrieval, frontoparietal

(FP), salience, ventral attention (VAN), and dorsal attention (DAN) networks, as well as their combinations. Univariate tests followed significant group effects and post hoc pairwise comparisons. Partial eta-squared (η^2_p) values were reported as effect sizes.

Moderation analysis for the relationship between neuroimaging and behavioral data (creativity percentile scores)

To examine whether creativity moderates group differences in functional connectivity, we extended the above models by adding the creativity percentile score as a covariate and including the group-by-creativity interaction term.

Separate multivariate GLM models were estimated for both within- and between-network connectivity: 1) Model predictors**:** group (children vs. adults), creativity (centered), FD, and the group × creativity interaction, and 2) Dependent variables: within- or between-network connectivity measures as described above. For significant interactions, simple slope analyses and predicted values were examined to probe the direction and nature of the moderation effects.

Results

Behavioral results

Both children and adults demonstrated average verbal and nonverbal general abilities, with no significant differences between adults and children in these measures, as determined by independent-samples *t*-tests. Group comparisons revealed no significant difference in creativity performance (DAT) outside of the scanner between

children and adults. However, creative interaction with ChatGPT (DSI) were significantly

higher in adults compared to children (Table 1).

Table 1. Comparisons between children and adults in general abilities and creativity

scores.

Domain	Measure	Groups		t-test
		Children	Adults	
		Mean(SD)	Mean(SD)	
Nonverbal abilities	Matrices	10.41 (1.99)	12.93(2.2)	-3.354, $p<.01$
Verbal abilities	Vocabulary	11.2(3.09)	13.92(3.25)	-2.72, $p<.05$
Creativity	DAT	28.61 (19.59)	26.16 (21.53)	0.33, NS
	DSI (while speaking to ChatGPT)	.81 (.01)	.82 (.005)	-3.67, $p < .001$

Note. SS = Standard Score (M = 100, SD = 15), ScS = Scaled Score (M = 10, SD = 3) *

$p < .05$, ** $p < .01$, *** $p < .001$, n.s. = non significant

Neuroimaging measures

<u>Within networks analysis</u>: After statistically controlling for head motion (FD), significant

group differences in functional connectivity were observed in the FP network, $F(1,27) =$

8.77, $p = .006$, partial $\eta^2 = .245$, the salience network, $F(1,27) = 5.65$, $p = .025$, partial η^2

$= .173$, and the DAN, $F(1,27) = 5.49$, $p = .027$, partial $\eta^2 = .169$, with adults showing

higher FC within these networks compared to children. These effects correspond to

medium-to-large effect sizes, indicating that a substantial proportion of variance in

connectivity was explained by group differences even after adjusting for motion.

Table 2. Within networks functional connectivity of EF, attention, and sensory networks

Network	Children (Mean ± SD)	Adults (Mean ± SD)	ANCOVA Results (F(1,27), p, partial η²)	Contrast
Cingulo-Opercular	0.307 ± 0.128	0.306 ± 0.106	F = 0.006, p = .940, partial η² = .000	C ≈ A
Frontoparietal (FP)	0.113 ± 0.036	0.201 ± 0.085	**F = 8.773, p = .006, partial η² = .245**	**A > C**
Memory	0.265 ± 0.168	0.342 ± 0.124	F = 1.938, p = .175, partial η² = .067	A > C
Default Mode (DMN)	0.181 ± 0.054	0.187 ± 0.057	F = 0.497, p = .487, partial η² = .018	A > C
Salience	0.198 ± 0.073	0.289 ± 0.078	**F = 5.653, p = .025, partial η² = .173**	**A > C**
Dorsal Attention (DAN)	0.162 ± 0.098	0.211 ± 0.086	**F = 5.487, p = .027, partial η² = .169**	**A > C**
Ventral Attention (VAN)	0.369 ± 0.107	0.344 ± 0.120	F = 1.562, p = .222, partial η² = .055	C > A

VAN: Ventral attention network; DAN: Dorsal attention network; FP: Fronto-parietal; CO: Cingulo-opercular; DMN: default mode network; C: children; A: adults

<u>Between Networks analysis:</u> After statistically controlling for head motion (FD), several significant group differences emerged in between-network functional connectivity (Table 3). Compared with children, adults exhibited significantly stronger connectivity between the CO-FP networks, $F(1,27)$ = 5.628, p = .025, $η^2_p$ = .173) and between the CO-Salience networks ($F(1,27)$ = 8.182, p = .008, $η^2_p$ = .233. Similarly, adults showed greater coupling between the DMN-Memory, $F(1,27)$ = 6.491, p = .017, $η^2_p$ = .194, and DMN-FP, $F(1,27)$ = 5.570, p = .026, $η^2_p$ = .171. After correcting for multiple comparisons (using FDR correction [104], Benjamini–Hochberg, q = .05), the results for the FP findings remained significant.

Enhanced between networks connectivity in adults was also observed between the Memory Retrieval network and multiple systems, including the FP, $F(1,27) = 6.449$, $p = .017$, $\eta^2_p = .193$, VAN, $F(1,27) = 5.514$, $p = .026$, $\eta^2_p = .170$, and DAN $F(1,27) = 5.944$, $p = .022$, $\eta^2_p = .180$. Finally, significant group effects were found for connectivity between the FP-VAN, $F(1,27) = 5.820$, $p = .023$, $\eta^2_p = .177$, and FP-DAN, $F(1,27) = 13.252$, $p = .001$, $\eta^2_p = .329$, with adults again showing higher connectivity.

Overall, these results suggest a broader and more integrated pattern of between-networks connectivity in adults relative to children, particularly involving the FP, DAN,VAN, memory, and salience systems. After correcting for multiple comparisons (FDR correction [104], Benjamini–Hochberg, q = .05), the results for the FP-salience and FP-DAN findings remained significant (Table 3). See also Figure 2.

Table 3. Between networks, functional connectivity of EF, attention, and sensory after controlling for FD

Network Pair	Children (Mean ± SD)	Adults (Mean ± SD)	ANCOVA Results ($F(1,27)$, p, η^2_p)	Contrast
CO–DMN	0.133 ± 0.070	0.119 ± 0.123	$F = 0.006$, $p = .937$, $\eta^2_p = .000$	C > A
CO-Memory	0.015 ± 0.096	0.085 ± 0.110	$F = 3.453$, $p = .074$, $\eta^2_p = .113$	A > C
CO–FP	0.043 ± 0.074	0.105 ± 0.052	**$F = 5.628$, $p = .025$, $\eta^2_p = .173$**	**A > C**
CO-Salience	0.109 ± 0.076	0.207 ± 0.079	**$F = 8.182$, $p = .008$, $\eta^2_p = .233$**	**A > C**
CO-VAN	0.165 ± 0.123	0.182 ± 0.103	$F = 0.311$, $p = .582$, $\eta^2_p = .011$	A > C
CO-DAN	0.121 ± 0.077	0.116 ± 0.086	$F = 0.494$, $p = .488$, $\eta^2_p = .018$	C > A

Network Pair	Children (Mean ± SD)	Adults (Mean ± SD)	ANCOVA Results ($F(1,27)$, p, η^2_p)	Contrast
DMN–Memory	0.111 ± 0.045	0.150 ± 0.091	**$F = 6.491$, $p = .017$, $\eta^2_p = .194$**	**A > C**
DMN – FP	0.090 ± 0.039	0.115 ± 0.065	**$F = 5.570$, $p = .026$, $\eta^2_p = .171$**	**A > C**
DMN–Salience	0.111 ± 0.060	0.115 ± 0.081	$F = 0.480$, $p = .494$, $\eta^2_p = .017$	A > C
DMN–VAN	0.196 ± 0.072	0.188 ± 0.091	$F = 0.015$, $p = .902$, $\eta^2_p = .001$	C > A
DMN–DAN	0.116 ± 0.073	0.118 ± 0.086	$F = 0.896$, $p = .352$, $\eta^2_p = .032$	A > C
Memory–FP	0.079 ± 0.064	0.104 ± 0.096	**$F = 6.449$, $p = .017$, $\eta^2_p = .193$**	**A > C**
Memory-Salience	0.049 ± 0.059	0.079 ± 0.103	$F = 3.990$, $p = .056$, $\eta^2_p = .129$	A > C
Memory-VAN	0.091 ± 0.073	0.133 ± 0.152	**$F = 5.514$, $p = .026$, $\eta^2_p = .170$**	**A > C**
Memory-DAN	0.086 ± 0.076	0.144 ± 0.100	**$F = 5.944$, $p = .022$, $\eta^2_p = .180$**	**A > C**
FP–Salience	0.138 ± 0.040	0.180 ± 0.060	$F = 2.434$, $p = .130$, $\eta^2_p = .083$	A > C
FP–VAN	0.095 ± 0.059	0.167 ± 0.088	**$F = 5.820$, $p = .023$, $\eta^2_p = .177$**	**A > C**
FP–DAN	0.039 ± 0.059	0.126 ± 0.043	**$F = 13.252$, $p < .001$, $\eta^2_p = .329$**	**A > C**
Salience–VAN	0.012 ± 0.000	0.189 ± 0.007	$F = 1.687$, $p = .205$, $\eta^2_p = .059$	A > C
Salience–DAN	0.010 ± 0.005	0.145 ± 0.005	$F = 1.824$, $p = .188$, $\eta^2_p = .063$	A > C
VAN–DAN	0.002 ± 0.007	0.186 ± 0.007	$F = 0.338$, $p = .566$, $\eta^2_p = .012$	A > C

VAN: Ventral attention network; DAN: Dorsal attention network; FP: Fronto-parietal; CO: Cingulo-opercular; DMN: default mode network; C: children; A: adults

Figure 2. Averages for within and between networks' functional connectivity (Z scores)

Figure 2. The Z-score averages for within and between networks functional connectivity for children (A), adults (B) and the difference between the groups (C). The hot color represents a higher Z values whrease the cool color represents lower Z values.

Moderation results

In order to determine whether creativity scores moderated the group effect on within-network connectivity (controlling for FD), we conducted moderation analyses, which revealed no significant group × creativity interaction effects across any network (p > .05).

Moderation analyses revealed that creativity outside the scanner (DAT) significantly interacted with group in predicting between-network functional connectivity. Specifically, significant moderation effects emerged for the FP–Salience, $F(1,26)$ = 5.869, p = .023, partial η^2_p = .184, and Salience–VAN, $F(1,26)$ = 4.575, p = .042, partial

η^2_p = .150, pairs, indicating that the relationship between creativity and connectivity strength differed between children and adults for these networks.

To determine which group (children vs adults) shows a greater association between creativity skills (DAT) and functional connectivity between FP-Salience networks, an indepth slope analysis for the significant interactions was conducted, and revealed that adults' show a positive significant slope (slope B = +0.028 per 1 SD creativity, p = .022, 95% CI [0.004, 0.052]) whrease children demonstrate a negative non significant slope (B = −0.014, p = .293, 95% CI [−0.042, 0.013] (n.s.)). Similar directions were found for Salience–VAN networks as well (Adults: B = +0.037, p = .062, 95% CI [−0.002, 0.077]; Children: B = −0.025, p = .273, 95% CI [−0.070, 0.021] (n.s.)). Overall, higher creativity is associated with stronger FP–Salience coupling in adults, but not in children (Figure 3).

Figure 3. Association between creativity, functional connectivity and group differences

Figure 3. A. Correlation analysis between the finctional networks and creativity score. B. Slope differences analysis, comparing children to adults.

Power analysis

Power calculations were anchored to published between-group differences in large-scale network connectivity. In younger versus older adults, Avelar-Pereira et al. reported a robust task-state difference in left FP-DAN coupling during the Multi-Source Interference Task, with groups of 29 and 30 and a large effect (Cohen's d ≈ 1.3)[105]. In a separate high-versus low-creative adult comparison (12 per group), Beaty et al. reported ROI-to-ROI group differences spanning t values of 2.38–3.75 for IFG–DMN links, corresponding to large effects (d ≈ 1.0–1.5)[106]. To adopt a conservative planning assumption, we powered a priori for a moderate effect (Cohen's d = 0.60–0.70), which requires approximately 44–33 participants per group for 80% power at α = .05 (two-sided). Our achieved sample (children, n = 15; adults, n = 16) is therefore well-powered

for large effects, consistent with prior literature, but underpowered for moderate effects, guiding the interpretation of null or small effects accordingly.

Discussion

How old do you need to be to use generative AI? With the invasion of these tools into our lives and even into the educational system [107], the current study aimed to explore the differences in cognitive control engagement when children vs. adults were orally communicating with ChatGPT during a creative task. In line with our hypothesis, children showed a lower engagement of cognitive control networks while communicating with ChatGPT compared to adults, despite an equal creativity level (age normalized) across groups. Moreover, and in line with our results, positive relations between creativity levels and cognitive control functional connectivity strength were observed in adults, but not in children.

Reduced cognitive control, attention, and modulation networks during conversations with GenAI tools in children

Talking to ChatGPT or any other generative AI tools provides a unique opportunity to quickly gather information and build upon it to generate more higher-level, creative ideas. Whereas knowledge was previously gathered using search engines or other printed paper sources, today, gathering basic information is faster, thanks to the efficiency prompts provided by the user[108]. This process demands planning the utterance, monitoring your response and ChatGPT's response, reorienting one's attention to unpredictable responses, and maybe inhibiting responses when the conversation takes a different direction - these abilities all fall under the EF system

definition [109]. Whereas these stages may not dramatically differ in children and adults during human-to-human interaction, the speed of information flow and responses from ChatGPT and the lack of gestures usually received from humans but equipped with other verbal gestures[110], might make this task very different than communicating with humans.

It was previously suggested that using generative AI tools results in lower alpha and beta waves generated from EEG data, with the claim that these tools demand less cognitive effort than an actual active search (engines vs self-search with no external tools) [111]. The current study suggests that where adults do show an engagement of cognitive control (FP), attention (DAN), and modulation (salience) systems while speaking to ChatGPT, children show reduced functional connections within these networks. These findings are in line with developmental theories previously suggested [112].

Children also demonstrated reduced functional connectivity between the selected cognitive networks (FP-salience, FP-DAN, DMN-memory). In line with the reported increased between-networks connectivity during development [34], it appears that, when communicating with ChatGPT, children are less capable of retrieving information and internal mentation (DMN-memory), or adapting the behavior/thoughts in response to changes in a given stimulus (FP-DAN). These results, combined with studies demonstrating a lack of maturation of the cognitive control system at the age of 6 years [35] (first graders, the age of the participants in the current study), suggest that speaking to ChatGPT, which seems not to be so complicated, actually demands cognitive control, which is lower in children who participated in our study. This is especially true as the

children in the current study are in their first grade, and there is a debate regarding the most appropriate age for using ChatGPT in school. Additional studies are needed to determine the exact time window in development when children gain the most from interacting with ChatGPT, also in terms of cognitive demands.

Differential engagement of cognitive control networks in creative thinking:
children versus adults

Creativity, measured in this study as DAT [77], is related to cognitive control, with evidence demonstrating increased creative abilities with increased frontal activation (specifically the Dorsolateral prefrontal cortex) from adolescence to young adulthood [112]. Interestingly, although children and adults in the current study exhibited comparable levels of creativity measured outside of the scanner (age-normalized DAT), only adults showed significant associations between functional connectivity among cognitive control, attention, and modulation networks (FP–salience, salience–VAN) networks involved in the switching of executive functions and attention [113] critcal for creative thinking [62-64,69,114]. In contrast, no such relations were observed in children.

Interestingly, previous research has linked creative thinking to the involvement of the salience network, which identifies and filters important or novel ideas before relaying them to the executive control system for goal-directed processing [62,63,114]. Notably, the DMN, proposed as a key component in models of creativity [66,115] that support spontaneous thought and flexible memory retrieval, did not differ between adults and children, either overall or in relation to their DAT scores. This lack of difference may reflect the novelty of interacting with ChatGPT for both groups.

An additional possible explanation for these findings between-networks

connectivity results and the absence of relations to creativity measures in children might

be that children still do not have the neurobiological infrastructure to benefit from the

conversation with ChatGPT, which also aligns with the lower DSI scores from their

generated ChatGPT script. This is critical given the extended attention given to co-

creative human-AI interaction [14], and their potential costs [21-23,25] and benefits [17-20] on

human creativity. Future studies should examine the interaction between familiarity with

ChatGPT and DMN involvement to determine whether the absence of engagement of

DMN in relation to creativity is due to the equal novelty shared among children and

adults using ChatGPT.

Limitation

The results of the current study should be considered in light of the following

limitations. First, the instructions for both age groups were slightly different: children

were told they'd speak to "the smartest computer in the world," whereas adults were

simply told "ChatGPT." In addition, the speech behavior, i.e., the conversation features

such as turn count, words/min, response latency, were not accounted for and might also

drive the results of the study (i.e., higher engagement and back and forth, might explain

the greater cognitive control engagement). Moreover, there are apparent differences in

vocabulary knowledge that change across the lifespan [116,117]. Such a difference, which

was not controlled here, might impact the co-creative interaction within the scanner and

provide an alternative explanation for our results. Future studies are needed to replicate

and extend our results, for example, by controlling for vocabulary knowledge in a similar

study design. Finally, Post-hoc power analysis indicated 80% power to detect medium-to-large effects (Cohen's d ≥ 0.73) for primary between-group comparisons. The moderation analyses were exploratory and likely underpowered for interaction effects, which typically require larger samples. Future studies with a sample size of ~ 50 children per group are warranted to adequately test creativity × group interactions.

Conclusions

This novel study examined the level of cognitive engagement of children while speaking to ChatGPT compared to adults, as well as how this cognitive engagement is related to creativity. The results point to the overengagement of the self-regulated networks in 6-year-old children, which warrants additional longitudinal studies to determine the turning point where cognitive control is more engaged and that creative thinking is related to cognitive control networks, similarly to adults. Given the rapid emergence of AI in our daily lives, alongside increased discussion on the costs and benefits of AI on creativity, our results highlight the significance of between-network functional coupling during human-AI creative acts. Decision makers, especially in the education system, as well as those related to children's health (pediatricians), should pay particular attention to the official age when children can use ChatGPT, similarly to the attention given to screen exposure.

Acknowledgement

The authors would like to thank the children, the parents, and the students participating in the current project. The authors would also like to thank the research team and students at the Educational Neueoimahging Group (of 2025) for helping with data collection and tasks' administration. The authors would also like to thank Tuval Raz for his help in analyzing the transcripts of participants' conversations with ChatGPT in the scanner.

Conflict of interest

The authors declare no conflict of interest.

Author contribution

THK: Conceptualization, data collection, funding, analysis, wrote the first version of the paper, reviewed the final version of the paper

RF: Analysis, wrote the first version of the paper, reviewed the final version of the paper

YK: Analysis, wrote the first version of the paper, reviewed the final version of the paper

DL: Data collection, reviewed the final version of the paper

EA: Data collection, reviewed the final version of the paper

Funding sources

No funding to declare.

References

1. Zivan M, Vaknin S, Peleg N, Ackerman R, Horowitz-Kraus T. Higher theta-beta ratio during screen-based vs. printed paper is related to lower attention in children: An EEG study. *Plos one.* 2023;18(5):e0283863.
2. Hahnefeld A, Fink M, Le Beherec S, Baur MA, Bernhardt K, Mall V. Correlation of screen exposure to stress, learning, cognitive and language performance in children. *European child & adolescent psychiatry.* 2024:1-10.
3. Horowitz-Kraus T, Fotang J, Niv L, Apter A, Hutton J, Farah R. Executive functions abilities in preschool-age children are negatively related to parental EF, screen-time and positively related to home literacy environment: an EEG study. *Child Neuropsychology.* 2024;30(5):738-759.
4. Nagata JM, Al-Shoaibi AA, Leong AW, et al. Screen time and mental health: a prospective analysis of the Adolescent Brain Cognitive Development (ABCD) Study. *BMC Public Health.* 2024;24(1):2686.
5. Sundqvist A, Barr R, Heimann M, Birberg-Thornberg U, Koch FS. A longitudinal study of the relationship between children's exposure to screen media and vocabulary development. *Acta Paediatrica.* 2024;113(3):517-522.
6. Jomaa F, Ebraheem F, Horowitz-Kraus T. Greater Parent–Child Brain Synchronisation During Printed Book Versus Screen Reading Using Hyperscanning Electroencephalograph Data. *Acta Paediatrica.* 2025.
7. Wan X, Kang X, Chen S, Du J, Yan F, Bai Y. Analysis of the impact of parents' electronic screen time habits, young children's screen exposure and parent-child interaction on language development delay in young children. *Frontiers in Pediatrics.* 2025;13:1667048.
8. Chin JH, Lee S, Ashraf M, et al. Young children's creative storytelling with ChatGPT vs. Parent: comparing interactive styles. Paper presented at: Extended Abstracts of the CHI Conference on Human Factors in Computing Systems2024.
9. Shrivastava V, Sharma S, Chakraborty D, Kinnula M. Is a sunny day bright and cheerful or hot and uncomfortable? young children's exploration of chatGPT. Paper presented at: Proceedings of the 13th Nordic Conference on Human-Computer Interaction2024.
10. Stoilova M, Livingstone S, Ringmar Sylwander K. Researching children's experiences of generative artificial intelligence: a child rights approach. 2025.
11. Neugnot-Cerioli M, Laurenty OM. The future of child development in the AI era. Cross-disciplinary perspectives between AI and child development experts. *arXiv preprint arXiv:240519275.* 2024.
12. Ali S, DiPaola D, Lee I, et al. Children as creators, thinkers and citizens in an AI-driven future. *Computers and Education: Artificial Intelligence.* 2021;2:100040.
13. Achiam OJ, Adler S, Agarwal S, et al. GPT-4 Technical Report. *arXiv preprint arXiv:230308774.* 2023.
14. Rafner J, Beaty RE, Kaufman JC, Lubart T, Sherson J. Creativity in the age of generative AI. *Nature Human Behaviour.* 2023;7(11):1836-1838.
15. Hubert KF, Awa KN, Zabelina DL. The current state of artificial intelligence generative language models is more creative than humans on divergent thinking tasks. *Sci Rep.* 2024;14(1):3440.

16.   Wan Q, Hu S, Zhang Y, Wang P, Wen B, Lu Z. *"It Felt Like Having a Second Mind": Investigating Human-AI Co-creativity in Prewriting with Large Language Models.* 2023.
17.   Kreisberg-Nitzav A, Kenett YN. Creativeable: Leveraging AI for Personalized Creativity Enhancement. *AI.* 2025;6(10):247.
18.   de Chantal P-L, Beaty RE, Laverghetta A, et al. Artificial intelligence enhances human creativity through real-time evaluative feedback. *PsyArxiv.* 2025.
19.   de Chantal P-L, Houde-Labrecque C, Leblanc M-C, Organisciak P. Investigating Lasting Effects of Real-Time Feedback on Originality and Evaluation Accuracy. *Creativity Research Journal.* 2024:1-18.
20.   de Chantal P-L, Organisciak P. Automated feedback and creativity: On the role of metacognitive monitoring in divergent thinking. *Psychology of Aesthetics, Creativity, and the Arts.* 2023.
21.   Wenger E, Kenett YN. We're Different, We're the Same: Creative Homogeneity Across LLMs. *arXiv.* 2025.
22.   Moon K, Kushlev K, Bank A, Green A. Impersonal Statements: LLM-Era College Admissions Essays Exhibit Deep Homogenization Despite Lexical Diversity. *PsyArxiv.* 2025.
23.   Doshi AR, Hauser OP. Generative AI enhances individual creativity but reduces the collective diversity of novel content. *Science Advances.* 2024;10(28):eadn5290.
24.   Cropley D. Is artificial intelligence more creative than humans? ChatGPT and the Divergent Association Task. *Learning Letters.* 2023;2:13-13.
25.   Moon K, Green AE, Kushlev K. Homogenizing effect of large language models (LLMs) on creative diversity: An empirical comparison of human and ChatGPT writing. *Computers in Human Behavior: Artificial Humans.* 2025;6:100207.
26.   Kosmyna N, Hauptmann, E., Yuan, Y. T., Situ, J., Liao, X., Beresnitzky, A. V., Braunstein, I., & Maes, P. . Your Brain on ChatGPT: Accumulation of Cognitive Debt when Using an AI Assistant for Essay Writing Task. *ArXiv https://arxivorg/abs/250608872.* 2025.
27.   Posner MI, & Petersen, S. E., . The attention systems of the human brain *Annu Rev Neurosci* 1990;13:25– 42.
28.   Posner MI. *Cognitive Neuroscience of Attention, Second Edition.* Guilford Publications; 2011.
29.   Petersen SE, & Posner, M. I. The Attention System of the Human Brain: 20 Years After. *Annual Review of Neuroscience.* 2012;35:73-89.
30.   Dosenbach NU, Fair DA, Cohen AL, Schlaggar BL, Petersen SE. A dual-networks architecture of top-down control. *Trends Cogn Sci.* 2008;12(3):99-105.
31.   Casey BJ, Giedd JN, Thomas KM. Structural and functional brain development and its relation to cognitive development. *Biol Psychol.* 2000;54(1-3):241-257.
32.   Giedd JN. Structural magnetic resonance imaging of the adolescent brain. *Ann N Y Acad Sci.* 2004;1021:77-85.
33.   Lenroot RK, Giedd JN. Brain development in children and adolescents: Insights from anatomical magnetic resonance imaging. *Neuroscience & Biobehavioral Reviews.* 2006;30(6):718-729.

34. Fair DM, Dosenbach NU, Petersen SE, Schlaggar BL. Resting state studies on the development of control systems. In: Posner MI, ed. *Cognitive Neuroscience of Attention.* 2nd ed. New York: Guilford Press; 2012:291-311.

35. Anderson P. Assessment and development of executive function (EF) during childhood. *Child Neuropsychol.* 2002;8(2):71-82.

36. Benedek M, Jauk E, Fink A, et al. To create or to recall? Neural mechanisms underlying the generation of creative new ideas. *NeuroImage.* 2014;88:125-133.

37. Leggett N. Early childhood creativity: Challenging educators in their role to intentionally develop creative thinking in children. *Early Childhood Education Journal.* 2017;45(6):845-853.

38. Runco MA, Jaeger GJ. The standard definition of creativity. *Creativity Research Journal.* 2012;24(1):92-96.

39. Green AE, Beaty RE, Kenett YN, Kaufman JC. The process definition of creativity. *Creativity Research Journal.* 2024;36(3):544-572.

40. Bai H, Mulder H, Moerbeek M, Leseman PP, Kroesbergen EH. The development of divergent thinking in 4-to 6-year-old children. *Creativity Research Journal.* 2024;36(4):656-674.

41. Vaisarova J, Carlson SM. When a spoon is not a spoon: Examining the role of executive function in young children's divergent thinking. *Trends in Neuroscience and Education.* 2021;25:100161.

42. Benedek M, Beaty RE, Schacter DL, Kenett YN. The role of memory in creative ideation. *Nature Reviews Psychology.* 2023;2(4):246-257.

43. Kenett YN. The role of knowledge in creative thinking. *Creativity Research Journal.* 2025;37(2):242-249.

44. Chrysikou EG. Creativity in and out of (cognitive) control. *Current Opinion in Behavioral Sciences.* 2019;27:94-99.

45. Benedek M, Fink A. Toward a neurocognitive framework of creative cognition: the role of memory, attention, and cognitive control. *Current Opinion in Behavioral Sciences.* 2019;27:116-122.

46. Beaty RE, Kenett YN. Associative thinking at the core of creativity. *Trends in Cognitive Sciences.* 2023;27(7):671-683.

47. Tzachrista M, Gkintoni E, Halkiopoulos C. Neurocognitive profile of creativity in improving academic performance—A scoping review. *Education Sciences.* 2023;13(11):1127.

48. Kleinmintz OM, Ivancovsky T, Shamay-Tsoory SG. The two-fold model of creativity: the neural underpinnings of the generation and evaluation of creative ideas. *Current Opinion in Behavioral Sciences.* 2019;27:131-138.

49. Sowden PT, Pringle A, Gabora L. The shifting sands of creative thinking: Connections to dual-process theory. *Thinking & Reasoning.* 2014;21(1):40-60.

50. Boot N, Baas M, van Gaal S, Cools R, De Dreu CK. Creative cognition and dopaminergic modulation of fronto-striatal networks: Integrative review and research agenda. *Neuroscience & Biobehavioral Reviews.* 2017;78:13-23.

51. Lebuda I, Benedek M. A systematic framework of creative metacognition. *Physics of Life Reviews.* 2023;46:161-181.

52. Runco M. The evaluative, valuative, and divergent thinking of children. *The Journal of Creative Behavior.* 1991.

53. Beaty RE, Johnson DR. Automating creativity assessment with SemDis: An open platform for computing semantic distance. *Behavior Research Methods.* 2021;53(2):757-780.

54. Patterson JD, Pronchick J, Panchanadikar R, et al. CAP: The creativity assessment platform for online testing and automated scoring. *Behavior Research Methods.* 2025;57(9):264.

55. Organisciak P, Acar S, Dumas D, Berthiaume K. Beyond semantic distance: Automated scoring of divergent thinking greatly improves with large language models. *Thinking Skills and Creativity.* 2023;49:101356.

56. Kenett YN. What can quantitative measures of semantic distance tell us about creativity? *Current Opinion in Behavioral Sciences.* 2019;27:11-16.

57. Sawyer RK, John-Steiner V, Moran S, et al. *Creativity and development.* Oxford University Press; 2003.

58. Benedek M, Beaty RE, Chrysikou EG, et al. Neuroscience of creativity. In: Reiter-Palmon R, Katz-Buonincontro J, eds. *APA Handbook of Creativity* in press.

59. Abraham A. The promises and perils of the neuroscience of creativity. *Frontiers in Human Neuroscience.* 2013;7:246.

60. Benedek M, Christensen AP, Fink A, Beaty RE. Creativity assessment in neuroscience research. *Psychology of Aesthetics, Creativity, and the Arts.* 2019;13(2):218.

61. Beaty RE, Benedek M, Silvia PJ, Schacter DL. Creative Cognition and Brain Network Dynamics. *Trends in Cognitive Sciences.* 2016;20(2):87-95.

62. Beaty RE, Kenett YN, Christensen AP, et al. Robust prediction of individual creative ability from brain functional connectivity. *Proceedings of the National Academy of Sciences.* 2018;115(5):1087-1092.

63. Beaty RE, Seli P, Schacter DL. Network neuroscience of creative cognition: mapping cognitive mechanisms and individual differences in the creative brain. *Current Opinion in Behavioral Sciences.* 2019;27:22-30.

64. Chen Q, Kenett YN, Cui Z, et al. Dynamic switching between brain networks predicts creative ability. *Communications Biology.* 2025;8(1):54.

65. Frith E, Elbich DB, Christensen AP, et al. Intelligence and creativity share a common cognitive and neural basis. *Journal of Experimental Psychology: General.* 2021;150(4):609-632.

66. Luchini SA, Volle E, Beaty RE. The role of the default mode network in creativity. *Current Opinion in Behavioral Sciences.* 2025;65:101551.

67. Marron TR, Berant E, Axelrod V, Faust M. Spontaneous cognition and its relationship to human creativity: A functional connectivity study involving a chain free association task. *NeuroImage.* 2020;220:117064.

68. Beaty RE, Christensen AP, Benedek M, Silvia PJ, Schacter DL. Creative constraints: Brain activity and network dynamics underlying semantic interference during idea production. *NeuroImage.* 2017;148:189-196.

69. Kenett YN, Chrysikou EG, Bassett DS, Thompson-Schill SL. Neural dynamics during the generation and evaluation of creative and non-creative ideas. *bioRxiv.* 2024:2024.2004.2015.589621.

70. Chrysikou EG, Weber MJ, Thompson-Schill SL. A matched filter hypothesis for cognitive control. *Neuropsychologia.* 2014;62:341-355.

71. Ovando-Tellez M, Kenett YN, Benedek M, et al. Brain connectivity–based prediction of real-life creativity is mediated by semantic memory structure. *Science Advances.* 2022;8(5):eabl4294.

72. Sun J, Liu Z, Rolls ET, et al. Verbal creativity correlates with the temporal variability of brain networks during the resting state. *Cerebral Cortex.* 2019;29(3):1047-1058.

73. Raz T, Luchini S, Beaty R, Kenett Y. Automated scoring of open-ended question complexity: A large language Model approach. 2024.

74. Brown L, Sherbenou R, Johnsen S. Test of Nonverbal Intelligence–Fourth Edition (TONI-4). *Austin, TX: Pro-Ed.* 2010.

75. Dunn LM, Dunn DM. *PPVT-4: Peabody picture vocabulary test.* Pearson Assessments; 2007.

76. Wechsler D. *The Wechsler Adult Intelligence Scale (WAIS) -IV.* New York: The Psychological Corporation.; 2008.

77. Olson JA, Nahas J, Chmoulevitch D, Cropper SJ, Webb ME. Naming unrelated words predicts creativity. *Proceedings of the National Academy of Sciences.* 2021;118(25):e2022340118.

78. Kraus D, Horowitz-Kraus T. Functional MRI research involving healthy children: Ethics, safety, and recommended procedures. *Acta Paediatrica.* 2022.

79. Greve DN, Fischl B. Accurate and robust brain image alignment using boundary-based registration. *Neuroimage.* 2009;48(1):63-72.

80. Avants BB, Epstein CL, Grossman M, Gee JC. Symmetric diffeomorphic image registration with cross-correlation: evaluating automated labeling of elderly and neurodegenerative brain. *Med Image Anal.* 2008;12(1):26-41.

81. Klein A, Andersson J, Ardekani BA, et al. Evaluation of 14 nonlinear deformation algorithms applied to human brain MRI registration. *Neuroimage.* 2009;46(3):786-802.

82. Taran N, Farah, R., DiFrancesco, M., Altaye, M., Vannest, J., Holland, S., Rosch, K. Schlaggar, B. L., & Horowitz-Kraus, T. . The role of visual attention in dyslexia: Functional connectivity analysis. *Human Brain Mapping.* Accepted.

83. Meri R. F, R., & Horowitz-Kraus, T. Children with dyslexia utilize both top-down and bottom-up networks equally in contextual and isolated word reading. *Neuropsychologia* 2020;147:107574.

84. Farah R, Horowitz-Kraus T. Increased functional connectivity within and between cognitive-control networks from early infancy to nine years during story listening. *Brain connectivity.* 2019;9(3):285-295.

85. Farah R, Coalson RS, Petersen SE, Schlaggar BL, Horowitz-Kraus T. Children Use Regions in the Visual Processing and Executive Function Networks during a Subsequent Memory Reading Task. *Cerebral Cortex.* 2019;29(12):5180-5189.

86. Twait T, Farah, R., & Horowitz-Kraus, T. Decreased functional connectivity of the salience network during narrative comprehension in children with reading difficulties: An fMRI study. *Clinical Neuroimage.* 2018;20:987-992.

87. Levinson O, Hershey A, Farah R, Horowitz-Kraus T. Altered functional connectivity of the executive functions network during a stroop task in children with reading difficulties. *Brain connectivity.* 2018;8(8):516-525.

88. Horowitz-Kraus T, & Holland, S.K. . Greater Functional Connectivity between Reading and Error-Detection Regions Following Training with the Reading Acceleration Program in Children with Reading Difficulties. *Annals of Dyslexia.* 2015;10.1007/s11881-015-0096-9

89. Price CJ. Current themes in neuroimaging studies of reading. *Brain Lang.* 2013;125(2):131-133.

90. Twomey T, Kawabata Duncan KJ, Price CJ, Devlin JT. Top-down modulation of ventral occipito-temporal responses during visual word recognition. *Neuroimage.* 2011;55(3):1242-1251.

91. Turkeltaub PE, Gareau L, Flowers DL, Zeffiro TA, Eden GF. Development of neural mechanisms for reading. *Nat Neurosci.* 2003;6(7):767-773.

92. Richlan F. Developmental dyslexia: dysfunction of a left hemisphere reading network. *Frontiers in human neuroscience.* 2012;6:120.

93. Vogel AC, Church JA, Power JD, Miezin FM, Petersen SE, Schlaggar BL. Functional network architecture of reading-related regions across development. *Brain Lang.* 2013;125(2):231-243.

94. Karunanayaka P, Schmithorst VJ, Vannest J, Szaflarski JP, Plante E, Holland SK. A linear structural equation model for covert verb generation based on independent component analysis of FMRI data from children and adolescents. *Front Syst Neurosci.* 2011;5:29.

95. Freedman L, Zivan M, Farah R, Horowitz-Kraus T. Greater functional connectivity within the cingulo-opercular and ventral attention networks is related to better fluent reading: A resting-state functional connectivity study. *NeuroImage: Clinical.* 2020;26:102214.

96. Vogel AC, Petersen SE, Schlaggar BL. The VWFA: it's not just for words anymore. *Front Hum Neurosci.* 2014;8:88.

97. Whitfield-Gabrieli S, Nieto-Castanon A. Conn: a functional connectivity toolbox for correlated and anticorrelated brain networks. *Brain connectivity.* 2012;2(3):125-141.

98. Power JD, Schlaggar BL, Petersen SE. Studying Brain Organization via Spontaneous fMRI Signal. *Neuron.* 2014;84(4):681-696.

99. Power AD, Fair, D.A., Schlaggar, B. L., Petersen, S. E. The Development of Human Functional Brain Networks. *Neuron.* 2010;67:735-748.

100. Power JD, Schlaggar BL, Petersen SE. Recent progress and outstanding issues in motion correction in resting state fMRI. *Neuroimage.* 2015;105C:536-551.

101. Gordon EM, Laumann TO, Adeyemo B, Huckins JF, Kelley WM, Petersen SE. Generation and Evaluation of a Cortical Area Parcellation from Resting-State Correlations. *Cereb Cortex.* 2014.

102. Johnson DR, Kaufman JC, Baker BS, et al. Divergent semantic integration (DSI): Extracting creativity from narratives with distributional semantic modeling. *Behavior Research Methods.* 2023;55(7):3726-3759.

103. Luchini SA, Maliakkal NT, DiStefano PV, et al. Automated scoring of creative problem solving with large language models: A comparison of originality and quality ratings. *Psychology of Aesthetics, Creativity, and the Arts.* 2025.

104. Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B.* 1995;57(1):289-300.

105. Avelar-Pereira B, Bäckman L, Wåhlin A, Nyberg L, Salami A. Age-Related Differences in Dynamic Interactions Among Default Mode, Frontoparietal Control, and Dorsal Attention Networks during Resting-State and Interference Resolution. *Front Aging Neurosci.* 2017;9:152.

106. Beaty RE, Benedek M, Wilkins RW, et al. Creativity and the default network: A functional connectivity analysis of the creative brain at rest. *Neuropsychologia.* 2014;64:92-98.

107. Sahu MN. The genai revolution: Unleashing The role of information technology in education. *Sudarshan Research Journal.* 2024;2(5):55-75.

108. Huang Y, Huang JX. Exploring ChatGPT for next-generation information retrieval: Opportunities and challenges. Paper presented at: Web Intelligence2024.

109. Cirino PT. Framing executive function as a construct and its relation to academic achievement. *Mind, Brain, and Education.* 2023;17(4):226-237.

110. Welivita A, Pu P. Is ChatGPT more empathetic than humans? *arXiv preprint arXiv:240305572.* 2024.

111. Kosmyna N, Hauptmann E, Yuan YT, et al. Your brain on chatgpt: Accumulation of cognitive debt when using an ai assistant for essay writing task. *arXiv preprint arXiv:250608872.* 2025;4.

112. Jolles DD, van Buchem MA, Crone EA, Rombouts SA. A comprehensive study of whole-brain functional connectivity in children and young adults. *Cerebral cortex.* 2011;21(2):385-391.

113. Schimmelpfennig J, Topczewski J, Zajkowski W, Jankowiak-Siuda K. The role of the salience network in cognitive and affective deficits. *Frontiers in human neuroscience.* 2023;17:1133367.

114. Beaty RE, Benedek M, Kaufman SB, Silvia PJ. Default and Executive Network Coupling Supports Creative Idea Production. *Scientific Reports.* 2015;5:10964.

115. Shofty B, Gonen T, Bergmann E, et al. The default network is causally linked to creative thinking. *Molecular Psychiatry.* 2022;27(3):1848-1854.

116. Cosgrove AL, Beaty RE, Diaz MT, Kenett YN. Age differences in semantic network structure: Acquiring knowledge shapes semantic memory. *Psychology and Aging.* 2023;38(287-102).

117. Li Y, Beaty RE, Luchini S, Hu W, Kenett YN. The role of semantic memory networks in crystallized intelligence and creative thinking ability. *Learning and Individual Differences.* 2024;111:102426.

Supplemental materials

Prompt for children

You are having a conversation with a six-year-old Hebrew-speaking child.
The child will ask you questions related to opening a store, kindergarten, or school that they wish to start.
You should always reply in **only one sentence**, and **end your sentence with a question** to encourage the conversation.
The conversation should last **five minutes**.
Do **not** end the conversation on your own initiative.
Do **not** start until I tell you to begin — first, the child will say "hello / hi" for a test, and you should just reply "hello" in return.
Wait for my instruction to start.

Prompt for adults

You are having a conversation with a Hebrew-speaking adult.
The adult will ask you questions related to opening a store, kindergarten, or school that they wish to start.
You should always reply in **only one sentence**, and **end your sentence with a question** to encourage the conversation.
The conversation should last **five minutes**.
Do **not** end the conversation on your own initiative.
Do **not** start until I tell you to begin — first, the adult will say "hello / hi" for a test, and you should just reply "hello" in return.
Wait for my instruction to start.