Two-Year Field Testing of Genetically Engineered American Chestnut Reveals Fungal Blight Tolerance

Thomas Klak[1], Steven E Travis[2], Virginia Grace May[3], Ek Han Tan[4], Mathew WH Chatfield[5], Miranda Wheeler[6]

Abstract

The American chestnut (*Castanea dentata* [Marsh.] Borkh.) was a foundational forest canopy species in eastern

North America until an accidentally imported fungal blight (caused by *Cryphonectria parasitica* [Murr.] Barr.)

rendered it functionally extinct across its native range. Biotechnological approaches, such as the bioengineered

Darling 54 line, have potential for future restoration of American chestnut, but field-based evaluations of blight

tolerance have been limited. Progress has been slowed by the many years it takes for seedlings to grow to saplings,

then to full-fledged trees. Current regulatory restrictions also constrain the testing of transgenic chestnuts to within

permitted orchards. This research reports on a two-year field trial of Darling 54, their non-transgenic wild-type

siblings and Chinese chestnut (*Castanea mollissima* Blume) controls deployed a randomized block design to test for

blight tolerance. In the two years, three replicates each of 261 trees were branch-inoculated with EP-155, a highly

virulent isolate of the fungal blight. Based on canker length, Darling 54 trees consistently outperformed their non-

transgenic wild-type siblings and Chinese chestnut. To our knowledge, this is the first report of a multi-year field

trial of fungal blight inoculations comparing advanced generation Darling 54 families. This field-based evaluation

suggests that reintroduction programs using Darling 54 American chestnuts, which can commence after federal

approval, may offer a promising path to success.

Key words: genetic engineering · field trials · fungal inoculation · restoration · eastern US forests

Corresponding author: Thomas Klak tklak@une.edu

[1] School of Marine and Environmental Programs, University of New England, Biddeford, Maine USA 0009-0008-4639-9551

[2] School of Biological Sciences, University of New England, Biddeford, Maine USA 0000-0001-9338-8953

[3] School of Marine and Environmental Programs, University of New England, Biddeford, Maine USA 0009-7 0000-7500-0695

[4] School of Biology and Ecology, University of Maine-Orono, Orono, Maine USA 0000-0001-8872-7404

[5] School of Biology and Ecology, University of Maine-Orono, Orono, Maine USA 0000-0001-5808-5332

[6] School of Marine and Environmental Programs, University of New England, Biddeford, Maine USA

Author Contributions Statement: T.K., V.M., E.H.T., M.C. and M.W. conducted the field trial. T.K., S.T. and V.M. prepared the data. S.T. and V.M. analyzed the data. T.K. and S.T. prepared the tables and figures. T.K., E.H.T, and V.M. wrote the manuscript. All authors reviewed and edited the manuscript.

Funding Declaration: To support this work, the first author received funding from the PW Sprague Memorial Foundation, the Quimby Family Foundation, and individual donors.

Acknowledgments: We thank colleagues at SUNY-ESF (State University of New York – Environmental Science and Forestry) for providing Darling 54 seeds and seedlings, and at West Virginia University's Forest Pathology and Environmental Microbiology lab for providing the chestnut blight fungus. For helpful suggestions on earlier drafts, the authors thank Jack Kennell, Jay Wason III, and three anonymous reviewers.

Competing Interests Statement: The authors declare that they have no competing interests as defined by Springer, or other interests that might be perceived to influence the results and/or discussion reported in this paper. The first author is a member of the board of directors of American Chestnut Restoration, Inc. and the Maine Chapter of the American Chestnut Foundation.

Introduction

For millennia, the American chestnut (*Castanea dentata* [Marsh.] Borkh.*, Cd*) was a foundational species of eastern US forests stretching from Maine to Mississippi and west to Indiana (Russell 1987; Wang et al. 2013). In the late 1800s, a fungal blight that American chestnut is completely susceptiple to (caused by *Cryphonectria parasitica* [Murr.] Barr.) was accidentally introduced (Rigling and Prospero 2017). By the 1950s, the fungal blight had spread throughout the native range to kill the vast majority of the estimated four billion American chestnut trees (Powell et al. 2019). The species survives today mainly as stump sprouts, as the fungal blight does not kill the roots. However, trees rarely grow to maturity before succumbing to the blight (Dalgleish et al. 2016).

Genetic engineering is one strategy being developed to restore the American chestnut. Aiming to give American chestnuts enhanced fungal blight tolerance, researchers at SUNY-ESF[7] inserted the Oxalate Oxidase (OxO) gene from wheat driven by various promoters via callus transformation into Ellis 1, a pure American chestnut line (Polin et al. 2006; Zhang et al. 2013; Newhouse et al. 2014). The expression of OxO appears to detoxify the

---

[7]The research team at State University of New York Environmental Science and Forestry College that produced the genetically engineered American chestnut was headed by the late Dr. William Powell and now by Dr. Andrew Newhouse.

oxalic acid that is excreted by *Cryphonectria parasitica* which is a necrotroph, thus conferring blight tolerance

(Steiner et al. 2017). The transformation of Ellis 1, using a construct containing OxO driven by the strong,

constitutive CaMV 35S promoter (Guilley et al. 1982) generated several transgenic events, including Darling 54

which is currently under consideration for federal deregulation (Jacobs et al. 2023; Newhouse et al. 2024; Klak et al.

2025). For this field trial, we utilized diversified trees derived from Darling 54 that were three or four breeding

generations from the transformed T0 generation (Westbrook et al. 2020). We also included wild type full-siblings

(null segregants) that did not inherit the Darling 54 transgene.

To date, no definitive fungal blight tolerance assays have been developed, but there are methods to

approximate the impacts of natural fungal blight. Prior empirical work has determined that the optimal fungus

inoculation methodologies vary significantly based on the age and stem diameter of the chestnut trees under study.

Three of these inoculation assays, in order of the age and size of the tree for which they are best suited, are small

stem, branch, and main stem. For young trees, a greenhouse based small stem assay has often been used, although

this method has drawbacks. It has been found to have a large "no take" rate (i.e., inoculations that failed to transmit

the fungus to the tree). It has also outright killed many tested seedlings and has tended to have a low correlation with

later tests of larger trees (Conn et al. 2023; Newhouse et al. 2024). In fact, Steiner et al. (2017) showed that only T0

Darling 54 plants did not exhibit wilting or death after performing small stem assays, while the 50% of resistant

'Qing' variety of *Castanea mollisima* exhibited wilting or death. Second, for smaller trees, a branch inoculation

assay known as the alternative small stem assay (AltSSA) can be utilized to measure the extent of fungal cankers

and tissue injury as the distance it spreads down stems from the inoculation sites (Cipollini et al. 2021). It was

developed and previously tested in Georgia, near the southern edge of the American chestnut's range (Conn et al.

2023). The AltSSA yielded canker production at the site of inoculation of over 90% (i.e. few "no take" inoculations

that are excluded from the data). This rate of successful inoculation is higher in many cases than other blight

inoculation methods (Conn et al. 2023). Third, main stem inoculation has been used extensively for trees grown in

the field with a 38mm (1.5in.) or larger diameter at breast height and is not recommended for smaller trees (Georgi

and Fitzsimmons 2015; Penn State 2025).

In the present study, we utilized the AltSSA method because 69% of inoculated trees had stem diameters

that are too small (<38mm) for main stem inoculation. For trees under 38mm diameter at breast height, which is the

case for all the trees in this study, even highly resistant genotypes may be lethally girdled when inoculated at too

small a diameter (Conn et al. 2023). Blight can aggressively penetrate through a small tree's living layers rather than

surrounding them as is typical fungal blight behavior with larger trees (Hebard 2012). The present study is also the

first to deploy the AltSSA outside of the south, near the northern edge of the chestnut's native range. Besides the

AltSSA's appropriateness for plants of sapling size, it has another advantage discovered in prior studies. Previous

publications on the AltSSA describe it as "intended primarily for forward selection" which means that young

chestnuts can be screened with the AltSSA and only those with small cankers can be outplanted, thus saving orchard

space and effort. At the same time, the AltSSA deployed in previous work correlated with other measures of blight

tolerance, including a multidimensional method gauging the blight tolerance of the young chestnut's mother tree; the

AltSSA also differentiated susceptibility between trees ranging from American chestnut to Chinese chestnut

(*Castanea mollissima* Blume) and hybrids in between (Cipollini et al. 2021; Conn et al. 2023). To our knowledge,

this is the first application of the AltSSA to compare genetically engineered chestnuts to full-sibling wild types.

To summarize, the AltSSA is deployed in the present research as the best nondestructive tool available to

assess trees beyond seedling-size but not yet large enough for main stem inoculation in a field setting. The two main

objectives of this study are: (1) to compare the fungal blight tolerance between Darling 54, non-transgenic wild-type

American chestnut, and Chinese chestnut trees grown in a field plot in Maine, and (2) to evaluate the use of the

AltSSA, a branch inoculation assay for quantifying the blight tolerance of American chestnut towards the ongoing

efforts to breed blight tolerant trees for restoration.

Materials and Methods

Transgenic American Chestnut and Chinese Chestnut Trees Tested

The two-year field trial was conducted on a one-acre, APHIS[8]-permitted orchard (permit #124-833JJQQ) in

long-term partnership with a private landowner in Cape Elizabeth, Maine (Klak 2018; TACF 2025; **Supp. Figs. 1**

and 2 for general location). Chestnuts in this orchard came from wild type mother trees that had been bred the

previous summer with Darling 54 pollen produced in labs at the University of New England and SUNY-ESF (Klak

et al. 2025). The resulting seedlings were grown from nuts assembled from crosses with wild type trees in Maine

and elsewhere provided by collaborators in 2021. A full-sibling family is defined here as a group of transgenic and

non-trangenic (wild-type) progeny from a cross between a single pistillate tree and a single staminate hemizygous

---

[8]Animal and Plant Health Inspection Service, US Department of Agriculture

transgenic tree in the same season. Not all seedlings in the trial orchard survived the first or even the second year of

outplanting and required replacement plantings of the following summers.

Tree height was measured to the nearest half centimeter from the top of the soil at the base of the tree to the

woody tip of the tallest branch, which was usually the main stem. Stem diameter was measured with an electronic

caliper to the nearest tenth millimeter approximately 16cm from the top of the soil. Because chestnut stems are often

not exactly round, two measures were taken perpendicular to each other and averaged together.

One methodological improvement for Year 2 of the trial was the inclusion of additional surviving 4- to 5-

year-old Darling 54 and wild-type orchard trees from the same families. This allowed for more intra-family

comparisons of Darling 54 and wild-type siblings. At the same time, some trees that were challenged in Year 1 but

did not have a full-sibling counterpart were excluded in Year 2. Lastly, all surviving trees possessing some

proportion of Chinese chestnut alleles in the orchard were inoculated in Year 2. This increased the number of plants

with Chinese chestnut alleles in the study from 19 in Year 1 to 30 in Year 2.

Branch Inoculation Method based on the Alternative Small Stem Assay (AltSSA)

For this study, we deployed the EP-155 isolate of *Cryphonectria parasitica* (Crouch et al. 2020). It is the

most virulent documented isolate of the chestnut blight and in recent years has been widely used in American

chestnut resistance testing assays including the AltSSA (Hebard 2012; Conn et al. 2023). We performed branch

inoculation using the AltSSA method that Cipollini et al. (2021) described, with minor changes noted here. EP-155

spores were obtained from West Virginia University's Forest Pathology and Environmental Microbiology lab and

were reconstituted in potato dextrose agar in petri dishes, following standard protocol (Kheyrodin 2024). The oven-

plating procedure from Midwest Grow Kits (940 Dieckman St, Woodstock, IL) was used to plate spores onto agar in

an aseptic environment. In the field, 4mm inoculum discs were cut to fit within the diameter of 5 or 6mm crimped

straws and positioned so that the fungus was in direct contact with each newly cut branch (**Supp. Fig. 3**). In both

years of the trial, we selected branches that were, whenever possible, south-oriented and mid-level on the tree.

Branch diameters needed to be in the 4.5–5.5mm range to firmly secure the crimped straw containing the fungus

plug against the fresh wound. Due to limited availability of branches, as well as internal checks of known resistant

and susceptible trees (Chinese chestnut, wild-type siblings within Darling 54 families), we only conducted the

AltSSA with the presence of EP-155.

Canker Scoring Methods

In both years, trees were inoculated on June 5, and the crimped straws containing the inoculum were removed one week later as in previous studies (**Supp. Fig. 3**). After 90 days (September 5), inoculated stems were removed, and cankers and injured tissue were measured. The sites from where the stems were cut were disinfected to reduce the likelihood of further infection to the tree. Following protocol, two measurements were taken. One was the discolored canker zone where the fungus had advanced from the inoculation site down the branch. A second measure called full canker length was taken of a longer canker if there was evidence of additional, patchy discoloration up the branch as described by Cipollini et al. (2021) and Conn et al. (2023). The presence of sporulation on cankers was also recorded when orange clumps of spores, forming over the branches' normally white lenticels, were visible (**Supp. Fig. 4**). If spores were present on one of the AltSSA inoculations, the entire individual was recorded for the presence of sporulation.

Statistical Methods

Data were analyzed using R and followed a standard analysis of covariance (ANCOVA) approach, which treated chestnut type (i.e., Darling 54, wild-type siblings, or trees with Chinese chestnut alleles) as the principal factor of interest. Tree height was treated as a covariate of both of the canker length responses to fungal blight infection. In order to fully meet model assumptions for normality, Full Canker length measured in Year 1 and Height (both years) were square-root transformed, whereas Full Canker length measured in Year 2, Darkened Canker length and Orange Canker length were log-transformed.

ANCOVAs were first run using just the Darling 54 and wild-type families from both trial years to rule out family and year effects, and to thereby justify treating all trees used in our experiments as independent indicators of fungal blight tolerance. Although overall levels of replication were similar between our Year 1 and 2 datasets, the greater amount of replication *within* families conducted in Year 2 allowed for a comparison of chestnut types nested within their full-sibling families and year of planting. The lesser amount of within-family replication in Year 1 did not provide sufficient full-sibling families to allow for the same approach, but still allowed for separate comparisons of combined full- and half-sibling families by mother tree and by pollen source. The latter comparisons were made for Year 2 as well to assure consistency of results. These analyses indicated that neither year nor family was a

significant variable in either the single Year 2 full-sibling model (**Supp. Table 1**) nor any of the Year 1 or 2

combined full- and half-sibling models (for mother tree results see **Supp. Table 2**; for pollen source results, see

Supp. Table 3). These two factors, family and year, were therefore omitted from subsequent analyses, which

allowed us to broaden our comparisons to include a third chestnut type, trees with Chinese chestnut alleles.

Significant overall effects of chestnut type on discolored canker or full canker length were followed by multiple

pairwise comparisons of type, which were evaluated by applying a Bonferroni correction to test-wise alpha levels.

We sought to strike a balance between minimizing the probability of committing Type I and Type II errors and thus

set our test-wise α-levels at 0.033 (α=0.10/3).

167   Prior to this study, genomic analysis (unpublished data obtained from genomics labs at the University of

Notre Dame and Virginia Tech University) revealed that three of the wild type mothers of the orchard trees in the

study had hybridity with European chestnut (*Castanea sativa* Mill.).[9] Thirty seven of the sixty five (57%) trees

tested in the Darling 54 families analysis in Year 2 of the trial had some European chestnut alleles, with the

percentage ranging from approximately 10.5 to 20%. To determine if the European-American hybrids had a

differing response to the blight than the 100% American chestnuts, a series of ANCOVAs was run, once again

treating tree height as a covariate. Both chestnut type and percent European chestnut were run as uncrossed factors.

Each blight response, as well as the tree height covariate, were log- or square-root transformed prior to analysis in

order to fully meet model assumptions for normality.

Results

Tree height analysis

179   For both years, tree height and stem diameter were significant positive covariates with canker length on the

180   branches that has been infected via the alternative small stem assay (AltSSA). In addition, tree height and stem

181   diameter were highly correlated, as expected. We determined which of these two covariates had a more significant

182   relationship with each response variable using a simple linear regression (response ~ covariate). Tree height was

---

[9] In 2020, leaf samples from three mother trees in Maine that phenotypically appeared to be American chestnut were part of a larger group of samples sent to genomics labs at the University of Notre Dame and Virginia Tech University for chestnut species analysis. The independent results from the two labs closely matched, revealing that one of the mother trees was approximately 21% European chestnut (*Castanea sativa* Mill.) while two others were 40% European chestnut. When bred with 100% American chestnut pollen, offspring from these three trees that are included in this study are on average 10.5-20% European chestnut.

more highly correlated with canker length than stem diameter so height was included in the model. Within family

effects and the influence from the height covariate were factored out for all chestnut types, so the only difference

between the Darling 54 and wild-type siblings was their presence or absence of the OxO transgene. When height

was held constant and included in the model, Darling 54 trees had statistically shorter discolored cankers than wild-

type trees (**Table 1**). The wild-type trees produced orange/darkened cankers that were on average $\geq$10mm (2024)

and $\geq$6mm (2025) longer than their Darling 54 siblings (**Supp. Table 4**).

Table 1. Full model and pairwise ANCOVA results from comparisons of all chestnut types measured for
Orange/Darkened and Full Canker length in Year 1, and Darkened and Full Canker length in Year 2 based on type
and tree height. Chestnut types include Darling 54, wild-type siblings (WT), and trees with Chinese alleles (CH).

Comparison	Source	Year 1 Orange Canker			Year 1 Full Canker			Year 2 Darkened Canker			Year 2 Full Canker		
		F	DF	p	F	DF	p	F	DF	p	F	DF	p
All	Type	17.58	2,161	<0.0001***	22.10	2,161	<0.0001***	8.85	2,92	0.0003***	7.19	2,92	0.0013**
	Height	5.99	1,161	0.0155*	8.98	1,161	0.0032**	8.07	1,92	0.0056**	6.60	1,92	0.0118*
Darling 54/WT	Type	32.90	1,143	<0.0001***	39.55	1,143	<0.0001***	5.61	1,63	0.0209*	3.31	1,63	0.0737
	Height	8.39	1,143	0.0044**	11.69	1,143	0.0008***	10.86	1,63	0.0016**	6.46	1,63	0.0135*
Darling 54/CH	Type	2.12	1,110	0.1478	6.58	1,110	0.0117*	16.80	1,61	0.0001***	13.81	1,61	0.0004***
	Height	4.02	1,110	0.0473*	6.06	1,110	0.0154*	4.22	1,61	0.0443*	6.05	1,61	0.0167*
WT/CH	Type	8.34	1,68	0.0052**	5.13	1,68	0.0266*	3.84	1,59	0.0549	3.97	1,59	0.051
	Height	0.35	1,68	0.5575	0.73	1,68	0.3964	3.14	1,59	0.0814	1.67	1,59	0.2008

193
The tree height-stem diameter correlation among all Darling 54 and wild-type trees was 0.833 (n=70).

Chinese chestnuts tend to grow wider and shorter than American chestnuts. When trees with Chinese alleles were

included (n=96), and the correlation was lower (0.785) but still high. Between height and diameter, height is worth

examining more closely to see if the Darling 54 were shorter than their wild-type siblings, which has been reported

elsewhere (TACF n.d.), and was attributed to costs of defending against pathogens that can consume a plant's

energy and resources that would otherwise be put to growth (Monson et al. 2022). In our experimental orchard,

Darling 54 trees had a mean height of 195.3cm (sd=85.3cm) while their wild-type siblings had a similar mean height

of 190.4cm (sd=70.3cm). The standard deviations indicate considerable height variation in each group, but Darling

54 trees were not on average shorter than their wild-type siblings. As has been recently reported in a study of

chestnut lines variously planted in open field and shelterwood sites in three other US states, chestnut growth and

performance will vary per location and site conditions (Olichney et al. 2026).

Year 1 AltSSA Canker Measurement and Analysis

Canker lengths were measured from the inoculation site on the branch at two distances: orange canker and full

canker (**Supp.Fig 4**). The most significant contrast observed was between the Darling 54 and wild-type siblings.

They differed statistically for both measures of canker length (orange canker length: $F_{1,143}=32.90$, p<0.0001; full

canker length: $F_{1,143}=39.55$, p<0.0001; **Table 1** and **Fig. 1**). Darling 54 also had significantly shorter full canker

lengths than trees with Chinese alleles ($F_{1,110}=6.58$, p=0.0117; **Table 1** and **Fig. 1**).

.

Fig. 1 90-day mean canker lengths for the three chestnut types in Year 1. Fig1A: Orange canker. Fig1B: Full canker.

The number inside each bar graph is the sample size. Darling 54 are hemizygous lines that carry the 35S::OxO

transgene, WT refers to null segregants from the same families that are wild-type, and Chinese refers to plants with

Chinese chestnut alleles. Error bars represent standard errors. Shared letters above the bars indicate non-statistically-

significant differences in means.

The presence of sporulation on cankers was also recorded. If spores were present on one of a plant's

inoculations, the entire individual was recorded for the presence of sporulation. A $\chi 2$ test was run on the presence or

absence of sporulation to compare chestnut types, then a series of odds ratio tests was run as planned contrasts to

determine if Darling 54 trees differed in sporulation presence from their wild-type siblings or the chestnuts with

Chinese alleles. The only significant result of the three planned contrasts was between Darling 54 and wild-type

siblings. There was significantly less sporulation observed on Darling 54 trees compared to wild-type siblings (odds

ratio 0.403, confidence interval 95% [0.181, 0.901], p=0.013).

Year 2 AltSSA Canker Measurement and Analysis

We observed that canker discoloration from fungal blight infection manifested differently in the two years

of the study. In Year 1, the zone appeared orange and aligned with coloration reported in other studies (May 2025).

In Year 2, the discoloration was dark and the texture was more shriveled. We speculate that the more shriveled and

darker-colored branches were impacted by the 2025 drought. Consistent with this interpretation is the fact that

whereas fungal sporulation was observed on most (i.e., 69%) orange zone branches in Year 1, it was entirely absent

in the Year 2 drought year. We observed that in Year 2 the fungal inoculations appeared phenotypically drier on the

branches. To capture the inter-year differences, we analyzed the darkened canker for the Year 2 on branches

inoculated via AltSSA.

Based on this analysis, the darkened cankers on Darling 54 trees in Year 2 were significantly shorter than

those on wild-type siblings ($F_{1,63}$=5.61, p=0.0209) and also significantly shorter than those of trees with Chinese

chestnut alleles ($F_{1,61}$=16.80, p=0.0001; **Table 1** and **Fig. 2**). Whether the canker zone was orange as in Year 1 or

darkened as in Year 2, there is considerable consistency in the inter-year findings when comparing Darling 54 trees

to their wild type siblings (**Figs. 1** and **2**). As noted, fungal sporulation differed between Years 1 and 2. Sporulation

was observed on most (i.e., 69%) orange zone branches in 2024, it was entirely absent at 90 days in 2025. Like the

chestnut hosts, the fungal blight appeared phenotypically drier during the 2025 drought.

Fig. 2 90-day mean canker lengths for the 3 chestnut types in Year 2. Fig2A: Darkened canker. Fig2B: Full canker.

The number inside each bar graph is the sample size. Darling 54 are hemizygous lines that carry the 35S::OxO

transgene, WT refers to null segregants from the same families that are wild-type, and Chinese refers to plants with

Chinese chestnut alleles. Error bars represent standard errors. Shared letters above the bars indicate non-statistically-

significant differences in means.

Another question this study addressed was whether cankers differed between trees that were 4- versus 5-

years-old in Year 2. A well-replicated test of outplanting year and family via comparison of Darling 54 and wild-

type siblings showed no effect of either factor on blight canker metrics (e.g., for darkened canker length, Year:

$F_{1,10}$=0.06, p=0.8064; Family: $F_{10,12}$=0.31, p=0.9646; **Supp. Table 1**). After these alternative explanations were

discounted, the conclusion was that both canker measurements showed that Darling 54 trees outperformed their

wild-type siblings ($F_{12,40}$=2.69, p=0.0095; **Supp. Table 1**).

Influence from European chestnut on AltSSA canker length

As mentioned earlier, genomic analysis revealed that three of the wild type mothers of the orchard trees in

the study had hybridity with European chestnut (*Castanea sativa* Mill.). Thirty seven of the sixty five (57%) trees

tested in the Darling 54 families analysis in Year 2 of the trial had some European chestnut alleles, with the

percentage ranging from approximately 10.5 to 20%. European chestnut is also thought to be susceptible to *Cp* and

should therefore not contribute resistance alleles in these trees (Ježić et al. 2024). We therefore investigated the

question of whether the percentage European chestnut genes could explain fungal AltSSA canker length. A series of

ANCOVAs was run, once again treating tree height as a covariate. In both years, canker lengths from trees that

inherit European chestnut alleles was not statistically significant compared to canker lengths of 100% American

chestnut trees (with p-values ranging from 0.30–0.47; **Supp. Table 4**).

Discussion

In this study, we showed that Darling 54 outperformed wild-type American chestnut and Chinese chestnut

over two years of field trial based on the alternative small stem assay (AltSSA), using the most virulent

Cryphonectria parasitica isolate, EP-155. Over the two years, we measured canker lengths on 261 trees in triplicates

from AltSSA infections and showed that Darling 54 exhibited significantly smaller canker lengths compared to

checks. Previous field tests of Darling 54 on fungal blight tolerance are few and based on small-scale field tests

(e.g., only n=3 Darling 54 trees and n=3 wild-type siblings) that were conducted on first or second generation

Darling 54 trees in a single year (Powell et al. 2020; ESF 2023; Newhouse et al. 2024).

Methodologically, this study provides further support for the efficacy of Cipollini et al.'s (2021) branch

inoculation assay (AltSSA), specifically as a tool for assessing chestnut trees in an orchard setting. The successful

inoculate rate of the AltSSA is high. The two previous AltSSA studies had very low "no take" rates and ≥98%

inoculation success (Cipollini et al. 2021; Conn et al. 2023). In our study, the "no take" rates were slightly higher,

with a 91% inoculation success rate in Year 1 and 95% in Year 2, which is comparable to these studies. In

comparison, our AltSSA "no take" rates were much lower than the ~40% rate produced by traditional small stem

assays (Gentner 2018). This study demonstrated the ability of the AltSSA to yield meaningful results in a different

outdoor environment at the northern edge of the species' range. Furthermore, we were able to improve on previous

studies by inoculating each sapling three times, with each being treated as a subsample. The subsampled canker

measurements were averaged together to reduce variation and more inclusively detect blight response. Lastly, the

AltSSA has previously been used to compare blight resistance among American and Chinese chestnuts and their

hybrids. This study was the first to establish a baseline response for blight tolerance of advanced generation Darling

54 trees compared to full-sibling wild-type trees in the field.

292   Our results from both years showed that Darling 54 trees outperformed trees with Chinese alleles when

using the AltSSA method and EP-155, although this finding requires further investigation because of the small

number of trees with Chinese alleles used in this comparison. In three of the four comparisons in the two years,

Darling 54 significantly outperformed trees with Chinese chestnut alleles (**Table 1** and **Figs. 1-4**). This might be

viewed as an unexpected finding, considering that the Chinese chestnut coevolved with the fungal blight and

therefore should have natural blight tolerance. However, some previous studies using the small stem assay found a

similar pattern, including the SUNY-ESF submission to APHIS for nonregulated status for Darling 54 (Newhouse et

al. 2024). The American Chestnut Foundation also reported that Darling 54 (erroneously referred to then as Darling

58) had shorter cankers than full Chinese chestnuts, although the difference was not statistically significant in their

small study (TACF 2023). The comparison of Darling 54 and trees with Chinese chestnut alleles is a subject

especially in need of further testing.

303   A major complication for assessing fungal blight tolerance is the American chestnut's natural lifespan of

five hundred or more years in the absence of blight. Predictions about long-term survivability based on evidence

from young trees must be taken with caution. Nonetheless, it is valuable for making incremental progress toward the

goal of chestnut restoration to assess the fungal blight tolerance of younger trees. This is particularly true in the field

where, instead of the relatively stable, consistent, mild and idealized lab/greenhouse conditions, trees are exposed to

harsher weather, seasonality, herbivory, a wider range of environmental conditions, and vicissitudes such as late

freeze and summer drought. The young trees evaluated in this study have already experienced all of the above.

Unfavorable weather conditions significantly impacted younger chestnut trees across Maine in 2025 (Year

2). Whereas 2024 (Year 1) was a relatively normal summer for temperatures and rainfall by historical standards,

2025 was exceptionally difficult for younger chestnuts. Compared to 2024, 2025 had greater swings from a wet

early summer followed by prolonged drought. Commenting on the impacts on agriculture in 2025, which would also

apply to young chestnut trees, Maine's Drought Task Force explained that "Crops typically require about 1" of

water per week and this has not happened during June through August" (Maine 2025; see also **Supp Figs. 1** and **2**).

The timing of the drought corresponded with the field trial in Year 2 and these impacts on chestnut trees of the

extreme weather were visible during the 90-day period in Year 2. Wilted leaves, leaf drop, seedling death, and, for

mature trees outside the orchard, bur drop and the aborting of flowers were observed. Across five monitored WT

orchards that host 5- to 8-year-old trees in different Maine locations, the production of burs was severely diminished

and even zero in some places (data not shown). In prior, more normal years, there was an increasing crop of flowers

followed by burs and fertile nuts on trees as they matured, as expected with chestnut maturation. These observations,

along with what we observed in terms of disease development on these trees in 2025 compared to 2024, are

potentially associated with drought, resulting in an atypical growing season in 2025.

In conclusion, we suggest that experimentally challenging Darling 54 with highly virulent fungal isolates

such as EP-155 should extend from the baseline established in this two year trial. Additional trials that can provide

further documentation of how Darling 54 trees perform under varying site conditions is warranted. One

recommended next step is to continue to challenge Darling 54 trees as they grow older and larger. Another is to

challenge trees growing in different geographical locations across the American chestnut native range. A third is to

evaluate how differing site specific environmental conditions, such as soils and varying amounts of light and shade,

impact Darling 54 performance. Indeed, other recent empirical work focused on tree growth and survival has

documented that geographical and site conditions shape the relative performance of various chestnut lines including

Darling 54 (Olichney et al. 2026). As the American chestnut restoration project continues to move incrementally

forward, additional trials will provide further insights as to the extent to which the OxO gene offers sufficient fungal

blight protection to Darling 54 American chestnuts.

Reference List

Cipollini, ML, Moss, JP, Walker, W, Bailey N, Foster C, Reece H, Jennings C. (2021) Evaluation of an alternative

small stem assay for blight resistance in American, Chinese, and hybrid chestnuts (*Castanea* spp.). Plant Disease.

105:576-584. https://doi.org/10.1094/PDIS-06-20-1272-RE

Conn, CE, Howie, N, Lynch, M, M, Lee S, Young, E, Westbrook J, Holliday, J, Zhang Q, Cipollini ML. (2023)

Validation of an alternative small stem assay for blight resistance in chestnut seedlings and recommendations for

broader use. Plant Disease. 107:1576-1583. https://doi.org/10.1094/PDIS-06-22-1489-RE

Crouch, JA et al. (2020) Genome Sequence of the Chestnut Blight Fungus *Cryphonectria parasitica* EP155: A

Fundamental Resource for an Archetypical Invasive Plant Pathogen. Phytopathology. 110(6):1180-1188.

https://doi.org/10.1094/PHYTO-12-19-0478-A

Dalgleish HJ, Nelson CD, Scrivani JA, Jacobs DF (2016) Consequences of Shifts in Abundance and Distribution of

American Chestnut for Restoration of a Foundation Forest Tree. Forests. 7(1):4.

https://doi.org/10.3390/f7010004

ESF (State University of New York-Environmental Science and Forestry). (2023) ESF Darling chestnut science

summary. State University of New York College of Environmental Science and Forestry.

https://www.esf.edu/chestnut/science-update/index.php#garden Accessed 25 Mar 2025

Gentner K (2018) Evaluation of blight resistance in chestnut F2 half-sibling and full-sibling families via small stem

assay. Honor's Thesis, University of Tennessee-Chattanooga

https://scholar.utc.edu/honors-theses/153/

Georgi L, Fitzsimmons S. (2015) Inoculation best practices. Chestnut Magazine (Summer) https://tacf.org/wp-

content/uploads/2017/03/Chestnut_Summer2015_FINAL.pdf

365 Guilley H, Dudley RK, Jonard G, Balàzs E, Richards KE. (1982) Transcription of Cauliflower mosaic virus DNA:

366 detection of promoter sequences, and characterization of transcripts. Cell. 30(3):763–773 https://www-sciencedirect-

367 com.une.idm.oclc.org/science/article/pii/0092867482902811

369 Hebard FV. (2012) The American Chestnut Foundation Breeding Program. Proceedings of the 4th International

370 Workshop on Genetics of Host-Parasite Interactions in Forestry

371 https://www.fs.usda.gov/psw/publications/documents/psw_gtr240/psw_gtr240_221.pdf

373 Jacobs, DF et al. (2023) Reintroduction of at☐risk forest tree species using biotechnology depends on regulatory

374 policy, informed by science and with public support. New For. 54:587–604

375 https://doi.org/10.1007/s11056-023-09980-y

377 Ježić, M., Nuskern, L., Peranić, K., Popović, M., Ćurković-Perica, M., Mendaš, O., Škegro, I., Poljak, I., Vidaković,

378 A., & Idžojtić, M. (2024). Regional Variability of Chestnut (*Castanea sativa*) Tolerance Toward Blight

379 Disease. *Plants*, *13*(21), 3060. https://doi.org/10.3390/plants13213060

381 Kheyrodin H. (2024) How to make Potato Dextrose Agar from Potato. J. Biol. Chem. Research 41(2): 35-39

382 http://jbcr.co.in/Current_Issue/Volume%2041%20(2),%20July%20to%20December,%202024/4.%20Potato%20Dex

383 trose%2035-39.pdf

385 Klak T. (2018) TACF-Landowner Partnerships: An Unfolding Story From Maine. Chestnut: The Journal of the

386 American Chestnut Foundation, 32(3):10-12 https://tacf.org/wp-content/uploads/2020/04/WEB-

387 Chestnut_Fall2018.pdf

389 Klak T, Pilkey H, May VG, Matthews D, Oakes AD, Tan EH, Newhouse AE (2025) Speed breeding transgenic

390 American chestnut trees towards restoration. Plant Direct 9(12):e70129 https://doi.org/10.1002/pld3.70129

Maine, State of. (2025) Drought Task Force Report on Current Drought Conditions September 4, 2025

https://www.maine.gov/mema/sites/maine.gov.mema/files/inline-

files/Drought%20Task%20Force%20Report%2020250904%20%281%29.pdf Accessed 6 Jan 2026

May, VG. (2025) Speed Breeding and Fungal Blight Testing of The Darling 54 American Chestnut (*Castanea*

dentata). Master's Thesis, School of Marine & Environmental Programs, UNE https://dune.une.edu/theses/449

Monson, R.K., Trowbridge, A.M., Lindroth, R.L., Lerdau, M.T. (2022) Coordinated resource allocation to plant

growth–defense tradeoffs. N. Phytol. 233 (3), 1051–1066 https://nph-onlinelibrary-wiley-

com.une.idm.oclc.org/doi/full/10.1111%2Fnph.17773

Newhouse AE et al. (2014) Transgenic American chestnuts show enhanced blight resistance and transmit the trait to

T1 progeny. Plant Sci. 228:88–97 https://www-sciencedirect-

com.une.idm.oclc.org/science/article/pii/S016894521400079X

Newhouse, AE et al. (2024) Petition For Determination Of Nonregulated Status For Blight-Tolerant Darling 54

American Chestnut (*Castanea dentata*). downloadable at: https://www.regulations.gov/document/APHIS-2020-

0030-17585

Olichney JA, Newhouse AE, Franklin MZ, Holliday JA, Klopf SK, Powell WA, Stehman SV, John E. Drake, JE.

(2026) Evaluating restoration candidates: Performance of blight-tolerant American chestnut varieties in open field

and shelterwood silvicultural conditions. Forest Ecology and Management, 599:123302

https://doi.org/10.1016/j.foreco.2025.123302

Polin LD. et al. (2006) Agrobacterium-mediated transformation of American chestnut (*Castanea dentata* (Marsh.)

Borkh.) somatic embryos. Plant Cell Tissue Organ Cult. 84:69–79 https://link-springer-

com.une.idm.oclc.org/article/10.1007/s11240-005-9002-1

Powell WA, Newhouse AE, Coffey V. (2019) Developing blight-tolerant American chestnut trees. Cold Spring

Harb Perspect Biol. 11(7):a034587. https://pubmed.ncbi.nlm.nih.gov/31110131/

Powell WA. et al. (2020) Petition for determination of nonregulated status for blight-tolerant Darling 58 American

chestnut. https://www.aphis.usda.gov/sites/default/files/19-309-01p.pdf

Rigling D, Prospero S. (2017) *Cryphonectria parasitica*, the causal agent of chestnut blight: invasion history,

population biology and disease control. Mol Plant Pathol. 19(1):7-20. https://bsppjournals-onlinelibrary-wiley-

com.une.idm.oclc.org/doi/full/10.1111/mpp.12542

Russell EWB. (1987) Pre-Blight Distribution of Castanea dentata (Marsh.) Borkh Bulletin of the Torrey Botanical

Club. (114)2:183-190 https://doi.org/10.2307/2996129

Steiner KC, Westbrook JW, Hebard FV, Georgi LL, Powell WA, Fitzsimmons SF. (2017) Rescue of American

chestnut with extraspecific genes following its destruction by a naturalised pathogen. New For. 48(2):317-336.

https://link-springer-com.une.idm.oclc.org/article/10.1007/s11056-016-9561-5

TACF (The American Chestnut Foundation). (2023) 2022 Annual Report.

https://tacf.org/wp-content/uploads/2021/05/2022_TACF_AnnualReport.pdf

TACF(The American Chestnut Foundation). (2025) Plot: Cape Elizabeth - D58 Common Garden 2021.

*dentata*Base. https://dentatabase-staging.herokuapp.com/orchards/1101/plots/3360 Accessed 13 Mar 2025

TACF (The American Chestnut Foundation). n.d. Darling 58/54. https://tacf.org/darling-58/ Accessed 17 Dec 2025

Wang GG, Knapp BO, Clark SL, Mudder, BT. (2013) The Silvics of *Castanea dentata* (Marsh.) Borkh., American

Chestnut, *Fagaceae* (Beech Family). U.S. Department of Agriculture, Forest Service, Southern Research Station.

https://doi.org/10.2737/SRS-GTR-173.

Westbrook JW, Holliday JA, Newhouse AE, Powell WA. (2020) A plan to diversify a transgenic blight□tolerant

American chestnut population using citizen science. Plants, People, Planet 2:84–95.

https://doi.org/10.1002/ppp3.10061

Zhang B, Oakes AD, Newhouse AE, Baier KM, Maynard CA, Powell WA (2013) A threshold level of oxalate

oxidase transgene expression reduces *Cryphonectria parasitica*-induced necrosis in a transgenic American chestnut

(*Castanea dentata*) leaf bioassay. Transgenic Res 22:973–982. https://link-springer-

com.une.idm.oclc.org/article/10.1007/s11248-013-9708-5

Supplemental Material

Supplemental Fig. 1 Moisture Conditions in 2024 and 2025 by the end of the 90-day inoculation period. Arrow

points to the field trial orchard's location

Source: US Drought Monitor

Supplemental Fig. 2 Precipitation deficit during the 2025 90-day inoculation test. Arrow points to the field trial

orchard's location

Base Map Source: Maine, 2025

Supplemental Fig. 3 Flagged branch immediately after inoculation, with crimped, clear plastic straw sealing the

inoculum disc snugly to the cut end of branch

Supplemental Fig. 4 Example cankers from 2024 (top; A=orange canker length including normally-white lenticels

turned orange indicating sporulation in the right photo; B=full canker length); and 2025 (bottom) showing darkened,

more shriveled canker where in this case both canker lengths are nearly the same

Supplemental Tables

Supp.Table 1. Full-sib Nested Analysis of Covariance (ANCOVA) results evaluating the effects of Year,
Family within Year, and Chestnut Type within Family (including Darling 54 and Wild-Type) on
Darkened and Full Canker length measured in Year 2. Due to the nested nature of the factors, the
ANCOVA's were run in stepwise fashion, first evaluating the effect of the covariate, height, on each
branch inoculation assay via Simple Linear Regression analysis, and then running Nested ANOVAs on
the residuals from the regressions. In order to fully meet model assumptions, Height was square-root
transformed, whereas Darkened and Full Canker length were log-transformed.

	2025 Darkened Canker			2025 Full Canker		
Source	F	DF	p	F	DF	p
Height	6.91	1,62	0.0108*	4.02	1,62	0.0495*
Year	0.06	1,10	0.8064	0.15	1,10	0.7096
Family	0.31	10,12	0.9646	0.63	10,12	0.7672
Type	2.69	12,40	0.0095**	1.67	12,40	0.1116

Supp.Table 2. Combined full- and half-sib Nested Analysis of Covariance (ANCOVA) results evaluating
the effects of Year, Mother Tree within Year, and Chestnut Type within Mother Tree (including Darling
54 and Wild-Type) on Orange and Full Canker length (Year 1 dataset) and Darkened and Full Canker
length (Year 2 dataset). Due to the nested nature of the factors, ANCOVAs were run in stepwise fashion,
first evaluating the effect of the covariate, height, on each branch inoculation assay via Simple Linear
Regression analysis, and then running Nested ANOVAs on the residuals from the regressions. In order to
fully meet model assumptions, Full Canker length measured in Year 1 and Height (both years) were
square-root transformed, whereas Full Canker length measured in Year 2, Darkened Canker length and
Orange Canker length were log-transformed.

	Year 1 Orange Canker			Year 1 Full Canker			Year 2 Darkened Canker			Year 2 Full Canker		
Source	F	DF	p	F	DF	p	F	DF	p	F	DF	p
Height	16.68	1,144	<0.0001***	12.89	1,144	0.0005***	6.91	1,62	0.0108*	4.02	1,62	0.0495*
Year	3.23	1,7	0.1153	3.33	1,7	0.1110	0.18	1,8	0.6811	0.04	1,8	0.8506
Mother	0.30	7,9	0.9356	0.33	7,9	0.9192	0.41	8,10	0.8894	0.63	8,10	0.7371
Type	3.93	9,128	0.0002**	4.63	9,128	<0.0001***	3.04	10,44	0.0052**	2.17	10,44	0.038*

Supp.Table 3. Combined full- and half-sib Nested Analysis of Covariance (ANCOVA) results evaluating
the effects of Year, Pollen Source within Year, and Chestnut Type within Pollen Source (including
Darling 54 and Wild-Type) on Orange and Full Canker length (Year 1 dataset) and Darkened and Full
Canker length (Year 2 dataset). Due to the nested nature of the factors, ANCOVAs were run in stepwise
fashion, first evaluating the effect of the covariate, height, on each branch inoculation assay via Simple
Linear Regression analysis, and then running Nested ANOVAs on the residuals from the regressions. In
order to fully meet model assumptions, Full Canker length measured in Year 1 and Height (both years)
were square-root transformed, whereas Full Canker length measured in Year 2, Darkened Canker length
and Orange Canker length were log-transformed.

Source	Year 1 Orange Canker			Year 1 Full Canker			Year 2 Darkened Canker			Year 2 Full Canker		
	F	DF	p	F	DF	p	F	DF	p	F	DF	p
Height	16.22	1,134	<0.0001***	13.56	1,134	0.0003***	6.91	1,62	0.0108*	4.02	1,62	0.0495*
Year	2.52	1,6	0.1634	1.54	1,6	0.2611	0.39	1,6	0.5563	0.07	1,6	0.8059
Pollen	0.38	6,6	0.8653	0.69	6,6	0.6704	0.18	6,8	0.9752	0.30	6,8	0.9197
Type	5.31	6,122	0.0001***	6.64	6,122	<0.0001***	2.89	8,48	0.0103*	2.46	8,48	0.0256*

Supp.Table 4. Summary inoculation assay statistics for Chestnut Types measured in Years 1 and 2.

	Year 1 Orange Canker				Year 1 Full Canker				Year 2 Darkened Canker				Year 2 Full Canker			
	Mean	SD	SE(mean)	n	Mean	SD	SE(mean)	n	Mean	SD	SE(mean)	n	Mean	SD	SE(mean)	n
Chinese	15.28	6.35	1.46	19	29.59	9.45	2.17	19	28.44	26.35	4.81	30	33.53	27.56	5.03	30
Wild-Type	21.61	12.86	1.78	52	39.91	14.39	2.00	52	15.46	12.84	2.27	32	18.78	16.02	2.83	32
Darling 54	11.71	10.19	1.05	94	25.37	13.33	1.37	94	9.21	6.91	1.19	34	12.96	13.04	2.24	34