Impact of gestational antibiotics on maternal and offspring gut microbiota and growth in pigs

Md. Rayhan Mahmud[1], Toomas Orro[2], Camilla Munsterhjelm[1,3], Tiina Pessa-Morikawa[4], Kristina Ahlqvist[1,3], Sami Junnikkala[4], Emilia König[1,3], Meritxell Pujolassos[5], M. Luz Calle[5,6], Huyk Nam Kwon[7,8], Claudio Oliviero[1], Mari Heinonen[1,3], Mikael Niku[4]

1. Department of Production Animal Medicine, Faculty of Veterinary Medicine, University of Helsinki, Helsinki, Finland
2. Institute of Veterinary Medicine and Animal Sciences, Estonian University of Life Sciences, Tartu, Estonia
3. Research Centre for Animal Welfare, Department of Production Animal Medicine, Faculty of Veterinary Medicine, University of Helsinki, Helsinki, 00014, Finland.
4. Department of Veterinary Biosciences, Faculty of Veterinary Medicine, University of Helsinki, Helsinki, Finland
5. Bioscience Department, Faculty of Sciences, Technology and Engineering, University of Vic – Central University of Catalunya, Vic, 08500, Spain
6. Institut de Recerca i Innovació en Ciències de la Vida i de la Salut a la Catalunya Central (IRIS-CC), Vic, 08500, Spain
7. School of Biological Sciences, University of Ulsan, Ulsan 44610, Republic of Korea
8. Basic-Clinical Convergence Research Institute, University of Ulsan, Ulsan 44033, Republic of Korea

Abstract

Maternal microbiota modulates the development of the microbiota in the offspring. Effects of gestational antibiotics are not well understood, as most studies have focused on the perinatal period. We treated sows with penicillin, tetracycline or saline on days 78-80 of the 114-119-day gestation, a critical period in the fetal immune system development. Microbiotas were analyzed by 16S rRNA gene amplicon sequencing in sow feces and vagina at days 77 and 113, in colostrum, and in piglet feces at three days, three weeks and ten weeks of age. Sow fecal microbiota changed during pregnancy, but less in antibiotic groups. No significant effects on

sow microbiota remained on day 113. The piglets of the antibiotic-treated sows had lower Firmicutes+Actinobacteriota to Bacteroidota+Proteobacteria ratio, lower alpha diversity and higher relative abundance of *Escherichia* at three days. At ten weeks, they exhibited higher alpha diversity, had higher of *Prevotella* and *Clostridium sp. CAG-127,* and smaller increase of *Limosilactobacillus* than the control. Increased alpha diversity at 10 weeks was associated with lower weight gain during nursing. *Oliverpabstia* and *Mitsuokella* were positively associated with weight gain, while *CAG-127* and *Campylobacter B* were negatively associated. Sow antibiotic treatment decreased the positive effect of *Oliverpabstia* and *Mitsuokella* and increased the negative effects of *CAG-127* and *Campylobacter B*. Gestational antibiotics may have had adverse effects on microbiota and growth of the offspring, even if their effects on maternal microbiota were undetectable by parturition.

Keywords: pregnancy, antibiotics, penicillin, tetracycline, fetal development, intestinal microbiota, growth, pig

Introduction

Proper development and function of the immune system, digestive system, and metabolism require microbiota (Bosch, 2021; Esser et al., 2019). Early-life exposure to microbial signals program the immune system and energy metabolism (Kimura et al., 2020; Torow et al., 2023). The immune system then reciprocally controls the composition of the microbiota by mucosal immunoglobulin A (IgA) and other regulatory mechanisms (Pabst et al., 2016). Interactions

with microbes begin already before birth: maternal microbiota modulates fetal development by circulating metabolites (De Agüero, 2016; Husso et al., 2023) and possibly also by macromolecular components (Kaisanlahti et al., 2023). This prepares the immune system for the rapid microbial colonization at birth and for the transition towards adult diet and microbiota at weaning. Disturbance of the early host-microbe interactions can have long-lasting consequences on microbiota composition and host physiology (Schokker et al., 2015).

Maternally administered antibiotics can also perturb the microbiota in the offspring and increase the risk of immunological and metabolic disorders (Dierikx et al., 2020; Miyoshi and Hisamatsu, 2025). The dysbiotic microbiota of antibiotic-treated mothers at peripartum is transferred to the offspring (Miyoshi&Hisamatsu 2025). However, little is known about the effects of prenatal maternal microbiota perturbation on fetal development. Experimental evidence is available mostly from mice, in which the short gestation does not allow the recovery from antibiotics before parturition, and therefore prenatal and postnatal effects cannot be distinguished. Complete lack of maternal microbiota during gestation caused broad changes in fetal gene expression and in postnatal intestinal innate immunity (De Agüero, 2016; Husso et al., 2023). In pigs, maternal antibiotic administration affected intestinal mucosa in the offspring (de Greeff et al., 2020), but in this study, the exposure occurred shortly before birth. In a prospective cohort study, 2[nd] pregnancy trimester antibiotic exposure was associated with altered long-term microbiota composition and obesity in humans (Zhang et al., 2019).

Pigs serve as a valuable model for investigating human gut microbiota and immune system due to the physiological and microbial resemblance to humans (Heinritz et al., 2013; Pabst, 2020), while understanding the effects of maternal antibiotic treatment on piglet microbiota is important for animal husbandry. Use of antibiotics in pig farming has raised concerns about

their effects on gut microbiota, antimicrobial resistance, and the overall health of the animals (Holman and Chenier, 2015). Investigating how maternal antibiotic treatment affects the microbiota of piglets is crucial, as the early colonization of microbes plays a significant role in immune system development and long-term health (Mueller et al., 2015; Schokker et al., 2015). Prior research has examined the impact of antibiotic exposure during pregnancy and early life on pigs, pointing out potential disruptions in microbial diversity and function (Guevarra et al., 2019). Antibiotic administration to sows during late pregnancy affected the development of offspring's intestine and this change persisted for at least five weeks after birth (de Greeff et al., 2020). Weaning is a critical period for the development and growth of piglets and is accompanied by a large change in the microbiota, and alteration of the maternal microbiota can exacerbate dysbiosis at this stage (Guevarra et al., 2019; Madany et al., 2022b).

In this study, we exposed pregnant sows to narrow-spectrum (penicillin) or broad-spectrum (tetracycline) antibiotics at days 78-80 of the 114 to 119-day gestation. The aim was to perturb the maternal microbiota at a critical period in the fetal development of the pig immune system (Butler et al., 2017; Sinkora and Butler, 2009), allowing recovery of the material microbiota to pre-treatment status before parturition, and to investigate the effects of maternal antibiotic treatment on development of the intestinal microbiota and growth of the offspring.

Methods

Experimental design

This study was conducted on a commercial farm in Finland. The sows (Yorkshire × Landrace; n = 98) had an average parity of 4.5 ± 1.6 (first parity sows were excluded) and had not received any antibiotics during their ongoing pregnancy. They were housed in two pregnancy rooms,

both with partially slatted concrete floors. In room one, sows were kept in groups of 6 to 16, each with an individual feeding cage, while in room two, the group size ranged from 4 to 7, and they fed from long troughs. In both rooms, they had access to some roughage from a hay rack and water ad libitum from water nipples. At day 77 of gestation, they were weighed with a cage scale (DV203E digital, Danvaegt, Hinnerup, Denmark) and a veterinarian examined them clinically to include only healthy individuals not needing any medical treatments into the study. Body conditions were scored as in Norring et al. (2019): 1 = emaciated, 2 = thin, 3 = fit, 4 = fat and 5 = very fat. The sows were assigned systematically to three groups ensuring same parity distribution to each group. The following day, a three-day antibiotic treatment regimen was initiated, with sows receiving either an intramuscular injection of a saline solution (group CON, *Natriumchloride Braun Ecoflac* 9 mg/ml, 0.6 ml/10 kg) as a control, procaine penicillin (group PEN, *Ethacillin®* 300 mg/ml, 15 mg/kg), or tetracycline (group TET, *Engemycin LA®* 100 mg/ml, 5 mg/kg).

On day 113 of gestation, a second clinical examination was performed, and only healthy sows that had not received antimicrobial treatments after the first sampling were retained in the study. Out of 98 initially enrolled sows, 37 remained in the three study groups (CON n=14, PEN n=12, TET n=11). To minimize the workload in farrowing supervision, 30 sows with farrowing during the same calendar days, were selected for further monitoring (10 per group; average parities: CON 4.4 ± 1.8, PEN 4.6 ± 1.8, TET 4.8 ± 1.8). The farrowings of the sows were supervised, and the birth time of the first piglet was recorded.

A total of 82 piglets (CON n= 30, PEN n=24, TET n=28) from 19 sows (CON n=7, PEN n=6, TET n=6) were included in the laboratory analyses, ensuring a balanced sex distribution across the sow medication groups. We only included piglets which received no antimicrobial

treatments during the follow-up period and remained with their own dam. No piglets from other litters were moved into the study litters during the suckling period. After weaning, the study piglets were housed together in pens according to their dam's medication group.

Piglets were weighed at one and three days as well as at three weeks of age using a digital hook scale (Pesola PHS040, China) and a custom-made fabric bag. At the age of ten weeks, they were weighed with the same cage scale, which was used for the sows earlier. Using weighting results we calculated average daily gain (ADG) for two growth periods: from birth to weaning (ADGsuckling) and from weaning to end of the follow-up period (ADGnursing) and these were used as outcomes to study microbiota associations with piglets weight gain.

Metadata for sows and piglets is included as Table S1. The animal experiment permit was granted by the Regional State Administrative Agency for Southern Finland (ESAVI/22394/2022).

Sow sampling

Sows were sampled at two timepoints: before antibiotic administration at an average of 77 ± 0.2 days of gestation and again shortly before farrowing at 113 ± 0.2 days of gestation. Fecal and vaginal samples were collected at both stages, while colostrum samples were obtained within six hours of the birth of the first piglet. Vaginal samples were collected using sterile cotton swabs (Applimed SA, Châtel-St-Denis) and inserted into sterile milk sample tubes (Greiner Bio-One Gmbh, Germany). Fecal samples were obtained manually from rectum using factory clean gloves and placed in sterile 76 × 20 mm feces tubes (Sarstedt, Nümbrecht, Germany). Colostrum samples were collected following the procedure described in (Kaiser, 2019), after carefully wiping the nipple with disinfectant, into milk sample tubes (Greiner Bio-

One Gmbh, Germany) and placed immediately after sampling in refrigerator temperature. After that the samples were aliquoted to sterile microcentrifuge tubes, immediately frozen to −20 °C, and transferred to −80 °C within one week.

Piglet sampling

Piglets were sampled at three timepoints: at three days of age, just before weaning (three weeks = 23–27 days of age), and before the piglets were transferred to the finishing herd (ten weeks = 70–74 days of age). Fecal samples were collected using sterile cotton swabs (Applimed SA, Châtel-St-Denis) inserted into the rectum and placed into sterile cryotubes. All samples were placed immediately after sampling to refrigerator temperature, transferred to −20°C within one hour and transported in three days to laboratory storage in −80°C until analysis.

One of the day three samples in the CON group was excluded because the piglet was castrated before sampling. Four week ten samples in TET group and two in the PEN group were excluded due to antibiotic treatments before the last sampling. Thus, the final microbiota sample numbers were: day three, CON n= 29, PEN n=24, TET n=28; week three, CON n= 30, PEN n=24, TET n=28; week ten, CON n= 30, PEN n=22, TET n=24.

DNA extraction and 16S rRNA gene amplicon sequencing

DNA was extracted from 125 mg of piglet feces, sow sampling swabs, and 200 µl of colostrum using the ZymoBIOMICS DNA Miniprep Kit based on mechanical cell lysis (Zymo Research, Irvine, CA), with minor modifications to the manufacturer's protocol as described previously (Husso et al., 2020). Negative controls (DNA extracted from unused sampling instruments, and no-template controls), a ZymoBIOMICS Microbial Community Standard (Zymo Research), and an in-house complex microbiota standard (adult cow fecal DNA extract) were included.

DNA concentration and purity was measured using a Nanodrop spectrophotometer 2000 (Thermo Fisher Scientific, Waltham, MA, USA).

For sequencing, the hypervariable regions V3 and V4 of the bacterial 16S rRNA genes were preamplified as described previously (Piirainen et al., 2025). Fecal and rectal swab samples were preamplified with 14 cycles, vaginal swabs with 18 cycles, colostrum samples with 21 cycles, along with negative controls for each sample type, and ZymoBIOMICS standard with 12 cycles. The sequencing was performed at the DNA Sequencing and Genomics Laboratory of the University of Helsinki, Finland. Excess primers were removed, and the indexing PCR was performed with 18 cycles. The PCR products were pooled and purified using MagSI-NGS plus beads (Magtivio, HK Nuth, Netherlands). The library pool was sequenced on the AVITI instrument (Element Biosciences, San Diego, CA) with paired-end reads (326 bp read 1 and 278 bp read 2) at a final library concentration of 7.5 pM, using the AVITI 2x300 Cloudbreak FS High Output Sequencing Kit.

Amplicon sequencing data processing

Primers and spacers were trimmed using Cutadapt version 1.10 (Martin, 2011). The mapping file was validated using Keemei (Rideout et al., 2016). QIIME2 version 2024.2 (Bolyen et al., 2019) was used to process the sequencing data on the Puhti supercomputer of the CSC – IT Center for Science, Finland. Read quality was inspected and forward reads truncated to 243 bp and reverse reads to 204 bp to optimize data quality. The DADA2 (Callahan et al., 2016) plugin was used to merge and denoise the reads to amplicon sequence variants (ASVs). Taxonomic classification was performed using Greengenes2 version 2022.10 (McDonald et al., 2024) and a Naïve Bayes classifier trained on the target region. ASVs with less than 10 counts, mitochondrial and chloroplasts reads, and reads with undefined phylum were

removed. The colostrum sequencing data was decontaminated using an R script, accepting ASVs which had at least 2× prevalence or at least 10× relative abundance in samples versus the negative controls. This removed 6.5% ± 5% of reads (mean ± standard deviation) in colostrum samples. The other sample types were not affected by contaminants due to their high microbial DNA content. The expected microbiota composition was observed for the ZymoBIOMICS Microbial Community Standard. The ASV count table is available as Table S2 and the metadata as Table S1.

Statistics

The microbiota compositions were analyzed in R version 4.5.0, using phyloseq 1.52.0 (McMurdie and Holmes, 2013), vegan 2.7-1 (Martinez Arbizu, 2020) and microbiome 1.30.0 (Lahti, 2017). Compositions were visualized using microViz 0.12.7 (Barnett et al., 2021). Alpha diversities were analyzed by the Shannon index in phyloseq, using non-rarefied ASVs with prevalence ≥5% in a sample type. Beta diversities were analyzed as Bray-Curtis distances of the relative abundances of species which were detected >10 times in ≥5% of samples and visualized using the plot_ordination function of phyloseq. PERMANOVA was performed using adonis2 and beta_disper of vegan, including group, sow parity, sow body condition score (BCS) (as a 4-level categorical variable), pregnancy room, farrowing room, and piglet sex.

Differential abundances were analyzed at genus level, in each sample type and each timepoint separately, using MaAsLin3 version 1.0.0, with total sum scaling (TSS, relative abundances) log_2 transformation and Benjamini-Hochberg correction, without median comparisons (Nickols et al., 2024). Read depths and covariates which were significant in beta diversity analysis were included as fixed factors: group and sow parity (as three levels) in all analyses; for sow feces, also sow BCS (as four levels) and pregnancy room; for day three and week three

piglets, also sow BCS, pregnancy room and farrowing room; and for week ten piglets, also sow BCS and piglet sex.

Longitudinal changes in genus abundances were analyzed by linear mixed-effects models (LMER) using the lme4 R package (Bates, 2015) for log-transformed relative abundances of genera with ≥25% prevalence and ≥0.1% mean relative abundance: lmer(abundance ~ group * timepoint + parity + pregnancy_room + bcs + (1 | animal_ID), data = df, REML = TRUE, control = lmerControl (optimizer = "bobyqa", optCtrl = list (maxfun = 2e5))). Type III ANOVA on primary effects and interactions was performed using lmerTest (Kuznetsova et al., 2017). P values were adjusted using the Benjamini–Hochberg correction (Benjamini, 1995).

Longitudinal changes in overall genus-level compositions were assessed as Aitchison distances of clr transformed data, using the volatility R package (Bastiaanssen et al., 2021). Genera which were detected >10 times in ≥5% of samples were included. Kruskal-Wallis and Wilcoxon tests were used for determining the significances of the observed differences.

The effects of treatment on alpha diversities and phylum-level compositions [log-transformed ratios of (Firmicutes+Actinobacteriota) / (Bacteroidota+Proteobacteria)] were analyzed by Generalized Linear Mixed Models package (GLMM) in SPSS 29.0.2.0. Variances explained by dam and pregnancy room were counted for by testing them as random intercepts to find the strategy resulting in the best model fit. Timepoint was included as repeated factor. Timepoint and treatment group as well as their interaction were included as fixed factors. Piglet sex as well as dam parity were controlled for if showing at least a statistical trend. Models were approved when homoscedasticity and normality of residuals could be established. Post-hoc pair-wise contrasts were interpreted with LSD adjustment for multiple comparisons.

Models were approved when homoscedasticity and normality of residuals could be established. Post-hoc pair-wise contrasts were interpreted with LSD adjustment for multiple comparisons.

General linear multivariate models (GLM) were used to analyze the associations between piglets' fecal microbiota and ADG. Initially, to identify potential associations between sow- and piglet-related variables and piglet growth (analyzed separately for ADG during the suckling and nursing periods), a mixed linear regression model was applied, with sow included as a random intercept. After incorporating sow-related variables—such as parity, body condition score (BCS) group, and pregnancy room—as fixed factors, the random effect of the sow was no longer significant. Therefore, in subsequent models analyzing growth and ADG associations, these sow-related variables were included as fixed effects in all GLM models to control for potential confounding effects.

Four separate GLM models were constructed to investigate associations between fecal microbiota and piglet growth. For the suckling period, ADGsuckling was used as the outcome variable, with selected fecal genera at birth and at weaning as explanatory variables (two separate models). For the nursing period, ADGnursing was used as the outcome variable, with selected fecal genera at weaning and at the end of the study as explanatory variables (two additional models).

Potential confounders: sow parity (2–8), sow body condition score (BCS) group (2.5, 3, 3.5, and 4), and pregnancy room (a two-level categorical variable) were included in all models.

To select the most relevant genera for inclusion in the multivariable GLM models, stability selection (Meinshausen and Bühlmann, 2010) was applied using the "stabiliser" package in R. Only genera with at least 10% prevalence and a minimum relative abundance of 0.1% were

considered (at birth: n = 52; at weaning: n = 143; at the end of the study: n = 123). Sow parity, BCS, and pregnancy room were included as dummy variables.

A triangulation procedure (Lima et al., 2021), involving three variable selection models, was used to identify the most stable genera associated with ADG. The ten most stable genera were included in the final multivariate GLM models. Multicollinearity among explanatory variables was assessed. The linearity of the relationship between the outcome and continuous explanatory variables (genera relative abundances) was visualized using LOESS scatter plots. If a nonlinear relationship was observed, a $\log_2$ transformation of genera relative abundances was applied. A stepwise backward elimination procedure was used for model selection, with a significance threshold of $p \leq 0.05$. Model assumptions were evaluated by assessing residual normality and scatter plots.

To assess the potential confounding effect of sow antibiotic treatment on the associations between bacterial genera and ADG, the sow treatment group (CON, PEN, TET) was included in the final models. Changes in the coefficients of the genera were calculated, and a change of $\geq 10\%$ was considered indicative of a confounding effect, suggesting that some of the associations between genera and ADG could be attributed to the sow's antibiotic treatment.

The described GLM analyses were performed using preselected taxa based on their associations with ADG, as identified by the stability selection models. Taxa associated with the treatment group but not selected by the stability models were not included in these analyses.

To further explore associations between microbiota and growth, we conducted additional GLM analyses to assess the relationship between the relative abundance of bacterial genera and ADGsuckling or ADGnursing. These analyses focused on genera that exhibited significantly

different relative abundances or significant timepoint × treatment group interactions at either three or ten weeks of age but were not among the 10 most stable genera identified by the stability selection models and thus were not included in the initial GLM models. In these additional models, sow-related confounders (parity, BCS, and pregnancy room) were included, and each genus was analyzed individually. The sow treatment group was also included in the final models to assess its potential confounding effect on the associations between these genera and ADG.

Similarly, we evaluated the associations between the Shannon diversity index and the Firmicutes+Actinobacteriota to Bacteroidota+Proteobacteria ratio with ADGsuckling and ADGnursing using GLM models. These models also included the sow treatment group to assess its potential confounding influence on the observed associations.

All GLM models were constructed using Stata/IC 14.2 (StataCorp, College Station, TX, USA). Generative AI (Claude Sonnet 4.5, Anthropic; and ChatGPT R Code Helper 5, OpenAI; both San Francisco, USA) was used for R coding assistance. All code was reviewed and validated by authors.

Results

Microbiota of sow feces, vagina, and colostrum

In sow fecal microbiota, there were no significant phylum or genus-level differences between the treatment groups (CON, PEN, and TET) at either timepoint (Fig 1). No significant differences were detected in alpha diversities between the treatment groups, nor were they different by beta diversity analysis.

A substantial change in the sow fecal microbiota was however observed between the two phases of pregnancy. The microbiota compositions were significantly different as examined by beta diversity (ADONIS2 p<0.004; Fig. 1D). The relative abundances of 187 genera were significantly different between the sampling timepoints in LMER analysis, including lower abundances of the major genera *Clostridium_T*, *Romboutsia_B*, *Turicibacter*, and *Cryptobacteroides* and higher of *Sodaliphilus*, *Faecousia*, *Prevotella*, *Terrisporobacter,* and *Lactobacillus* at the end of pregnancy (Table S3).

Interestingly, in both antibiotic groups the fecal microbiota compositions changed significantly less between the sampling timepoints in comparison to CON, as measured by Aitchison distances; p = 0.043 for both comparisons.

Fig. 1. Fecal microbial profiles of pregnant sows treated with saline (CON = control group), penicillin (PEN) or tetracycline (TET) or during days 78-80 of pregnancy. Pregnancy day 77 (77 D) shows the results before treatments and 113 D shortly before farrowing. Microbiota compositions at (A) phylum and (B) genus level, numbers below the graph indicate parity. (C) alpha diversities of sow fecal microbiota (D) Beta diversity (PCoA plots of Bray-Curtis dissimilarity; 77D solid lines, 113D dashed lines).

In sow vaginal microbiota, the ratio of Firmicutes to Bacteroidota was significantly higher in TET vs CON at day 77 (Fig. 2). No other significant differences were observed between the experimental groups at either phase of pregnancy by Shannon alpha diversity index, beta diversity, or genus-level abundances (Fig 2). The vaginal microbiota changed significantly between the phases of pregnancy by beta diversity (ADONIS2 $p < 0.002$; Fig. 2D), but there was no significant difference in the magnitude of the change between the experimental groups. The relative abundances of *Fusobacterium_C* and *Veillonella_A* increased significantly from day 77 and both were among the most abundant at D 113, while the relative abundance of *Clostridium_AN*, one of the minor genera, decreased (Fig. 2B and Table S4). Other significant changes were not detected.

Fig. 2. Vaginal microbial profiles of pregnant sows treated with saline (CON = control group), penicillin (PEN) or tetracycline (TET) during days 78-80 of pregnancy. Pregnancy day 77 (77 D) shows the results before treatments and 113 D shortly before farrowing. Microbiota compositions at (A) phylum and (B) genus level, numbers below the graph indicate parity. (C) alpha diversities of sow fecal microbiota (D) Beta diversity (PCoA plots of Bray-Curtis dissimilarity; 77D solid lines, 113D dashed lines).

In sow colostrum microbiota, neither the alpha diversity by Shannon nor beta diversity, nor the phylum ratio differed significantly between CON and the antibiotic-treated groups (Fig 3). No significant differences were detected in the microbial composition between the treatment

groups at genus level. Three sows (two PEN and one TET sows) formed a separate cluster in the PCoA plot due to high abundances of *Streptococcus* as seen in Fig 6C (left). The ASVs matching the species *Streptococcus dysgalactiae* was very highly abundant in these three sows.

Fig. 3. Colostrum microbial profiles within 6 hours after delivery of the first piglet in sows treated with saline (CON = control group), penicillin (PEN) or tetracycline (TET) during days 78-80 of pregnancy. Microbiota compositions at (A) phylum and (B) genus level, numbers below the graph indicate parity. (C) alpha diversities of sow fecal microbiota (D) Beta diversity PCoA plots of Bray-Curtis dissimilarity.

Development of piglet fecal microbiota and impacts of maternal antibiotics

The microbiota of the 3-day-old piglets was characterized by high relative abundance of Proteobacteria (Fig 4A). *Clostridium_P, Escherichia_710834, Streptococcus,* and *Bacteroides_H* were the most abundant genera (Fig. 4B). Alpha diversity was significantly lower in PEN versus CON (p=0.007) (Fig. 4C). The difference between CON and TET was not significant. The treatment groups differed in beta diversity (ADONIS2 p=0.001). In PCoA of Bray-Curtis distances CON was slightly separated from PEN and TET (Fig 4D).

The phylum ratio of (Firmicutes+Actinobacteriota) / (Bacteroidota+Proteobacteria) was significantly higher in CON vs PEN (p=0.006) and CON vs TET (p=0.016). At genus level, PEN had significantly higher abundance of *Escherichia _710834* and *Phocaeicola _A_858004*, and lower abundances of *Peptostreptococcus*, *Streptococcus*, Unclassified Cellulosilyticaceae, *Streptococcus*, and *Veillonella_A* in MaAsLin3 (Fig. 5, Table S5). These genera also showed significant timepoint × treatment group interactions, i.e. differences in change between the treatment groups by time in LMER analysis (Fig. 5, Table S6). No significantly different genera were detected in TET vs CON.

In the end of the suckling phase, at three weeks of age, the most abundant phyla were Firmicutes_A, Firmicutes_D, and Bacteroidota, while Proteobacteria had much lower relative abundance (Fig 4A). *Clostridium_T, Fimimorpha, Romboutsia_B, Unclassified Lachnospiraceae, Limosilactobacillus* and *Faecousia* were the most abundant genera. (Fig 4B). Alpha diversity of the fecal microbiota of the piglets had increased during suckling (p=0.000). No significant differences in Shannon diversity were detected between the treatment

groups (Fig. 4C). The treatment group had significant effect by ADONIS2 (p=0.008), but the groups overlapped in PCoA (Fig. 4D).

The phylum ratio (Firmicutes+Actinobacteriota) / (Bacteroidota+Proteobacteria) tended to be lower in TET compared with CON (p=0.097). At genus level, *Egerieousia* and *CAG-873* were significantly less abundant in TET vs CON, and *Phocaeicola_A_858004* more abundant in PEN vs CON by MaAsLin3. Significant differences between the treatment groups in the change by time (timepoint × treatment group interactions) were detected by LMER analysis for *Egerieousia* and *CAG-873* and additionally for *Alloprevotella* and *VUNA01*, which were not significantly different at three weeks by MaAsLin3 (Fig. 5 & Table S5).

At ten weeks the most abundant phyla were Firmicutes_A, Firmicutes-D, and Bacteroidota. Phylum Proteobacteria was almost absent (Fig 4A). *Lactobacillus*, *Clostridium_T*, *Prevotella*, *Limosilactobacillus, Blautia_A_141781, and Terrisporobacter* were the most abundant genera (Fig 4B). Alpha diversity by Shannon diversity index had increased further from week three to week ten (p=0.017) and was higher in TET (p=0.036) compared to CON (Fig. 4C). No significant effects of treatment on beta diversity were detected by ADONIS2. CON separated partially from PEN in PCoA, TET overlapped with both the other groups (Fig. 4D).

At ten weeks, the ratio Firmicutes + Actinobacteriota / Bacteroidota + Proteobacteria was higher in CON (p=0.014) compared to TET (and tended to be higher in CON vs PEN (p=0.063)). There were pronounced differences in the relative abundances of several genera between CON and the antibiotic groups (Fig. 5 and Table S5). Both PEN and TET had significantly higher abundances of *Agathobacter_164117, Alloprevotella, Anaerovibrio, CAG-127,* and *Prevotella* than CON. Additionally, *Butyribacter*, *Roseburia*, and *Turicibacter* were

significantly more abundant and *Oliverpabstia* less abundant in PEN vs CON in MaAsLin3. The abundance changes by time for *Agathobacter_164117, Alloprevotella, CAG-127, Roseburia*, and *Turicibacter* were significally different between the treatment groups (significant timepoint × treatment group interactions in LMER, Fig. 5 and Table S6). In addition, the increase of the relative abundance of the major genus *Limosilactobacillus* from three to ten weeks was significantly stronger in CON than in the antibiotic groups (timepoint × treatment group interaction p=0.003 in LMER, Fig. 5 and Table S6).

The microbiota changed significantly more in TET compared to CON and PEN between three days and ten weeks (p.adj < 0.03 for Aitchison distances in both comparisons). In PEN, the change was not different from CON.

Fig. 4. Piglet fecal microbiota composition and diversity at 3 days (3 D), 3 weeks (3 W) and 10 weeks (10 W) of age of **sows treated with saline (CON = control group), penicillin (PEN) or tetracycline (TET)during days 78-80 of pregnancy.** Microbial compositions at (A) phylum and (B) genus level. (C) Alpha diversity, as Shannon diversity index, (D) Beta diversity, as PCoA plots of Bray-Curtis dissimilarity.

Fig 5. Trajectories of fecal microbial genera in piglets of sows treated with saline (CON = control group), penicillin (PEN) or tetracycline (TET) during days 78-80 of pregnancy. Horizontal axis age 3 days (3 D), 3 weeks (3 W) and 10 weeks (10 W), vertical axis relative abundance. Significance in MaAsLin3 at each timepoint and significance of timepoint × treatment group interaction in LMER are indicated in the plots: ** $q < 0.01$, *$q < 0.05$, † $q < 0.1$.

Piglet weight and growth

Piglets were born at an average weight of 1.48 ± 0.18 kg (SD), weaned at 8.51 ± 1.35 kg, and transferred to the fattening unit at 29.6 ± 5.1 kg. The ADGsuckling was 0.29 ± 0.05 kg/day (n = 80; birth weight was unavailable for two piglets), and ADGnursing 0.44 ± 0.10 kg/day (n = 82).

Associations of piglet fecal microbiota with growth

We evaluated the effects of sow antibiotic treatment on the associations of piglet microbiota and growth. Treatment was included in all models evaluating these associations. There was a consistent and significant effect of sow antibiotic treatment on piglet ADGnursing across all GLM models evaluating associations between microbiota at three and ten weeks of age and weight gain. PEN piglets grew less than CON piglets, with model coefficients ranging from –0.129 to –0.074 kg/day and p-values between <0.001 and 0.031. TET group piglets also showed reduced growth compared to CON piglets, with coefficients ranging from -0.077 to -0.017 kg/day; however, p-values varied from 0.017 to 0.643 (see Tables S9 – S12).

Sow antibiotic treatment was not associated with ADGsuckling in any of the models (see Tables S7 and S8).

We detected associations between microbial diversity and ADGnursing. An increase in Shannon diversity at three weeks by one unit was associated with an average increase of 0.037 kg/day in ADGnursing (SE = 0.017, p = 0.038), based on a GLM model that included sow-related variables (parity, BCS, pregnancy room) and piglet sex as confounders (n = 82). Adding treatment to this model did not substantially alter the coefficient (<10% change). Conversely, an increase in Shannon diversity at ten weeks (n = 76) by one unit was associated with an

average decrease of 0.064 kg/day in ADGnursing (SE = 0.029, p = 0.029). Including treatment in this model decreased the coefficient by 17%, as Shannon diversities were higher in TET group than in CON (Fig. 4) suggesting that the negative effect of Shannon diversity on weight gain was enhanced by sow antibiotic treatment (see Tables S13 and S14).

No associations were found between the phylum ratio of (*Firmicutes* + *Actinobacteriota*) / (*Bacteroidota* + *Proteobacteria*) at any of the three timepoints and ADGsuckling or ADGnursing (data not shown).

Associations of genus-level relative abundances at the three time-points to ADG were studied separately for the suckling and the nursery period using the GLM multivariable models (multiple genera were included as explanatory variables). Most notably, the relative abundances of *Oliverpabstia* and *Mitsuokella* at 10 weeks of age had significant positive associations with ADGnursing, and *CAG-127*, *UBA6382,* and *Campylobacter_B* had significant negative associations (Table S10, Fig 6). When treatment was included in the GLM model, the association coefficient for *Oliverpabstia* with ADG nursing decreased by 11%, while the negative association coefficient of CAG-127 with ADGnursing decreased by 33%, and the coefficient for *Campylobacter_B* decreased by 25% (Table S10). The relative abundance of *Oliverpabstia* was higher in the CON group compared to the PEN treatment group, while the relative abundance of *CAG-127* was higher in the PEN and TET groups compared to CON, and *Campylobacter_B* increased more in in the PEN and TET groups compared to CON (Fig. 5).

Treatment also modulated the association coefficients of UBA6382 and *Mitsuokella* with ADGnursing decreasing the coefficient of UBA6382 by 36% and increasing that of *Mitsuokella*

by 13%. However, their relative abundances were not significantly affected by the sow antibiotic treatment at 10 weeks by MaAsLin3 analysis (Fig. 5).

Fig 6. Loess regression curves of association of relative abundances of bacterial genera at 10 weeks with ADGnursing in piglets of sows treated with saline (CON = control group), penicillin (PEN) or tetracycline (TET) during days 78-80 of pregnancy.

The relative abundance of *Actinobacillus C 733574,* among a few other genera, at three days was negatively associated with ADGsuckling. Relative abundance of *Butyricimonas* at three weeks was negatively and *Romboutsia_B* positively associated with ADGsuckling. Relative abundances of *Paludicola* and *Sodaliphilus* at three weeks were positively associated with

ADGnursing. However, inclusion of treatment to the GLM model had only small effects on the association coefficients of any of these genera (Tables S7-S8). *Faecousia* at three weeks was positively associated with ADGnursing, and inclusion of sow treatment in the model increased the association coefficient by 27%. In contrast, *Limousia* was negatively associated with ADGnursing, and adding the treatment variable decreased the coefficient by 19%. (Fig S1, Table S9).

Additional GLM models were constructed to evaluate genera that showed significantly different relative abundances between treatment groups or significant timepoint × treatment group interactions but were not among the 10 most stable genera identified by the stability selection models. These were included individually in the models to assess their associations with ADGnursing. Several of these genera showed significant associations, and sow antibiotic treatment was identified as a significant confounder in multiple cases (see Tables S11 and S12). LOESS regression curves illustrating these associations are presented in Fig 6.

The relative abundance of *VUNA01* at three weeks was significantly and positively associated with ADGnursing in the CON and PEN groups (p = 0.007) (Fig S1, Table S12). When treatment was included in the model, the association coefficient was reduced by 40%. Although the relative abundance of this genus was lower in the PEN and TET groups compared to CON, the difference was not statistically significant in MaAsLin3. However, a significant timepoint × treatment interaction was detected using LMER (Fig. 5). *Alloprevotella* at three weeks showed a similar but weaker positive association with ADGnursing (p = 0.100). Including treatment in the model reduced the association coefficient by 30% (see Table S7).

Turicibacter had higher relative abundances in PEN than CON at ten weeks of age (q=0.07). The genus had negative association with ADGnursing (p = 0.056) and the coefficient was increased by 33% when treatment was included. Similar results were obtained for *Clostridium_T*, which was one of the most abundant genera at ten weeks with higher relative abundance in PEN than CON (LMER q=0.12 for the timepoint × treatment group interaction). This genus had significant negative association with ADGnursery (p = 0.018) and the coefficient increased by 15% when treatment was included.

Two additional genera, *Faecalibacterium* and *Megasphaera_A_38685*, which have been shown to be positively associated with growth in earlier studies (refs), were also included in the analysis. The relative abundance of *Faecalibacterium* at ten weeks had a positive association with ADGnursery (p = 0.041), which inclusion of treatment increased by 33%. However, no effect on the relative abundance of this genus by antibiotic treatment was observed. The relative abundance of *Megasphaera_A_38685* at ten weeks also had a positive association with ADGnursery (p = 0.008), which was reduced 12% by inclusion of treatment. At ten weeks it had lower abundances in PEN and TET vs CON, but the difference was not significant.

Discussion

In this study, we treated pregnant sows with saline, penicillin, or tetracycline five weeks before parturition and analyzed their fecal, vaginal, and colostrum microbiota and the fecal microbiota of their offspring using 16S rRNA gene amplicon sequencing. No significant differences were detected in sows between the treatment groups in fecal or vaginal microbiota compositions shortly before parturition at day 113 or colostrum microbiota, but antibiotic treatment interfered with the dynamic change in maternal fecal microbiota during pregnancy. Significant differences in microbial diversities and compositions were observed in the fecal microbiotas of the piglets. The differences persisted and even increased after weaning and separation of the piglets from their dams, suggesting a possible impact on early programming of the mechanisms regulating host-microbe interactions. Microbial composition especially at ten weeks of age was associated with growth: microbial diversity at ten weeks was negatively associated with ADGnursing. Furthermore, relative abundances of several genera such as *Oliverpabstia*, *CAG-127* and *Campylobacter_B* at ten weeks and *Faecousia* and *Limousia* at three weeks were associated with ADGnursing. Antibiotic treatment modified these associations.

Effect of antibiotic treatment on sow microbiota

Our results on the sow microbiota compositions show that the effects of the antibiotic treatment at day 77 of gestation on the fecal microbiota had subsided by day 113, as we expected. No major differences were observed between control and antibiotic-treated groups in fecal, vaginal or colostrum microbiota, making it unlikely that direct effects of antibiotics on sow microbiota compositions persisting until the time of delivery would have affected the microbiota of the piglets. While the microbial composition changed during pregnancy, we did not detect significant change in microbial richness, or in the relative abundances at phylum level between

days 78 and 113. Substantial changes in the composition of the fecal microbiota are known to take place during pregnancy both in human and in pigs. The most robust differences involve decrease of the phyla Proteobacteria (Pseudomonadota) and Actinobacteria (Actinomycetota) in human (Koren et al., 2012)and significant reduction of microbial richness both in human and pigs (Koren 2012, Kong 2016) accompanied by differential abundances of several individual genera (Koren 2012, Kong 2016, Liu 2025). The most significant differences are detected between early and late pregnancy. There was little overlap in the differentially abundant genera in our study compared to those reported in earlier studies (Kong et al., 2016; Liu et al., 2025). However, the fecal microbiota changed less towards parturition in antibiotics-treated sows. Such changes may affect the early priming of the piglet immune system and the subsequent microbial colonization.

Previous studies on the effects of antibiotics during pregnancy have mainly focused on treatments given to sows at late pregnancy and/or during lactation (de Greeff et al., 2020; Ma et al., 2020; Wang et al., 2021; Wang et al., 2020; Zhu, 2024) directly modifying the sow intestinal and colostrum microbiota compositions during delivery and lactation (de Greeff 2020). Further studies will be needed to determine how transient antibiotics-induced perturbations are mediated and affect the development of the piglet's microbiota and its future health.

Vaginal microbiota is involved in the initial colonization of the newborn (Chen et al., 2022; Li et al., 2022; Liu et al., 2019) and changes in this microbiota may have implications for the piglet health. However, we did not detect any significant differences between the treatment groups that could be associated with the differences detected in the piglet fecal microbial compositions. Colostrum is another important source of microbes for colonization of the

newborn piglet gut (Liu et al., 2019). We did not detect differences in the colostrum microbiota composition between treatment groups, indicating that antibiotic treatment about one month earlier had minimal impact on milk microbiota composition. A few sows belonging to the antibiotic groups showed high levels of *Streptococcus dysgalactiae*, a mastitis-associated bacterium, despite showing no clinical signs. These sows were not among those whose piglets were included in the study. Similar findings were made in sow colostrum in an earlier study not involving the use of antibiotics (Piirainen et al., 2025).

Early-life microbiota: Initial colonization

Earlier research has shown that antibiotic treatment of dams during pregnancy has impacts on the development of the commensal microbiota of their offspring (de Greeff et al., 2020; Dierikx et al., 2020; Zhang et al., 2019). In most experimental animal studies addressing this question, antibiotics were administered orally shortly before or at the time of parturition (Alhasan et al., 2020; Alhasan et al., 2023; Champagne-Jorgensen et al., 2020; Chen et al., 2021; de Greeff et al., 2020; Gonzalez-Perez et al., 2016; Madany et al., 2022a, b; Wang et al., 2021; Zou et al., 2018) and often also during lactation (Alhasan et al., 2020; Chen et al., 2021; Wang et al., 2020). Effects often included reduced microbial richness and lower relative abundance of Firmicutes and Actinobacteria in relation to Proteobacteria in the early gut microbiota of the offspring (Chen et al., 2021; Gonzalez-Perez et al., 2016; Zou et al., 2018) Analogous findings have been recorded in many studies on the effect of maternal therapeutic antibiotics on the microbiota development of human infants (Dierikx et al., 2020). Interestingly, we observed similar effects on the developing microbiota of the offspring when the dams were treated with antibiotics intramuscularly as early as five weeks before parturition. At three days of age the fecal microbiota of piglets of PEN-treated sows had

lower alpha diversity and lower relative abundance of the phyla Firmicutes +

Actinobacteriota in relation to Bacteroidota + Proteobacteria.

Significant differences in bacterial abundance were detected between the treatment groups.

The PEN group piglets had higher relative abundances of *Escherichia _710834 and*

Phocaeicola _A_858004. While there is considerable variation in taxa reported to be

differentially abundant in the offspring without or with maternal antibiotic treatment, higher

relative abundance of the genus *Escherichia* or family Escherichia-Shigella has been reported

in the newborn progeny of antibiotic-treated vs control dams in several species (Alhasan et

al., 2023; Chen et al., 2021; Zou et al., 2018). Species belonging to this taxon have been

linked to dysbiosis and opportunistic infections (Denamur et al., 2021; Luo Y, 2022; Pantazi

et al., 2025). The PEN group piglets also had lower abundances of *Peptostreptococcus*,

Streptococcus, Unclassified Cellulosilyticaceae, and *Veillonella_A* than controls.

Peptostreptococcus and *Veillonella* have been reported to belong to the most abundant genera

in weaned piglets with diarrhea (Kong et al., 2022). The genus *Streptococcus* contains several

potentially pathogenic species, such as *S. suis* and *S. dysgalactiae* (Ma et al., 2025) but the

ASVs matching these species were not significantly different between the treatment groups.

Pre-weaning transition: three weeks

At three weeks of age, the differences between the antibiotic groups and the controls had

subsided. The alpha diversity had increased substantially in all the groups and no effects of

antibiotic treatment on alpha diversity measures were detectable. There was only a tendency

to lower phylum ratio in TET vs CON. The genera *Phocaeicola*, *CAG-873*, *VUNA01* and

Alloprevotella were significantly differentially abundant between the groups.

Phocaeicola_A_858004, which had lower abundance in the antibiotic groups, is a gut

commensal identified as *P. vulgatus,* which is recognized as a potential carbohydrate utilizer in pigs (Holman DB, 2022). The species has been shown to utilize milk oligosaccharides *in vitro* (Rumeau et al., 2025). *Egerieousia,* with lower abundance in TET, is a relatively uncharacterized genus that was identified by metagenomic analysis in chicken gut (Gilroy et al., 2021). *Egerieousia* sp004561775 has been identified as a potential butyrate producer in pigs (Holman DB, 2022). Little information is available on *Prevotella CAG-873,* also less abundant in TET, except that a strong positive correlation of sow *Akkermansia* during lactation to piglet fecal *CAG-873* after weaning has been shown (Saladrigas-Garcia et al., 2022). *VUNA01* is a bacterial taxon identified in pigs (Holman DB, 2022) but its properties are not known.

Post-weaning: maturation

By ten weeks the alpha diversity had increased further towards more mature levels in all groups but was significantly higher in TET compared to CON. The Firmicutes + Actinobacteriota / Bacteroidota + Proteobacteria ratio was lower in PEN than CON. The relative abundance of *Oliverpabstia* was significantly higher in CON than in the antibiotic groups. In addition, the increase in the relative abundance of *Limosilactobacillus* by ten weeks was much more pronounced in CON than in the antibiotic groups. Several genera, including *Agathobacter_164117, Alloprevotella, Anaerovibrio, Butyribacter, CAG-127, Roseburia, Prevotella, and Turicibacter* were more abundant in one or both antibiotic groups.

Limosilactobacillus is a genus containing potential probiotic species (Ksiezarek et al., 2022). Species belonging to this genus have been shown to improve the intestinal barrier function in piglets (Li et al., 2023) and other species (Wu, 2022). *Oliverpabstia* (Wylensek et al., 2020) is

a genus of the family Lachnospiraceae that contains species enriched in piglet feces after weaning (Holman DB, 2022).

Agathobacter, Anaerovibrio, Alloprevotella, Prevotella, and *Turicibacter* are all relatively abundant commensal bacteria. *Anaerovibrio* contains species with lipolytic activity growing on sugars (Strömpl, 1999). *Alloprevotella* utilizes carbohydrates and produces succinic acid (Downes et al., 2013). *Agathobacter (*Rosero et al., 2016*)* and *Prevotella* (Ivarsson et al., 2014) are involved in fiber degradation and carbohydrate metabolism and production of short-chain fatty acids. The relative abundance of *Prevotella* has been reported to increase after weaning in piglets (Guevarra et al., 2018; Mach et al., 2015), with positive correlations of its relative abundance to body weight and soluble IgA concentration (Mach et al., 2015). Decreased abundance of *Prevotella* in the offspring after maternal antibiotic treatment has been reported in several studies with maternal antibiotic treatment (Alhasan et al., 2023; Champagne-Jorgensen et al., 2020; Zhu, 2024) . *Turicibacter* has the capability of decreasing the levels of lipids and modulating bile acid profiles of in the serum of mammalian hosts *(*Lynch et al., 2023*)*. *CAG-127* is a member of the family Lachnospiraceae known for production of short-chain fatty acids (Brito Rodrigues et al., 2025).

Associations with Growth Performance

The associations of microbiota and growth were strongest after weaning. *Oliverpabstia* and *Mitsuokella* were positively associated with growth, and *CAG-127* and *Campylobacter_B* negatively associated. Inclusion of sow antibiotic treatment to the statistical models had substantial effects on the association coefficients. The results suggests that antibiotic treatment reduced the positive effect of *Oliverpabstia* and enhanced the negative effect of *CAG-127* and *Campylobacter_B* on ADG-nursing. As the relative abundance of *Oliverpabstia* was higher in

the CON group compared to the PEN treatment group, the positive effect of this genus on growth in CON vs PEN may be due to its higher relative abundance in CON. The relative abundances of *CAG-127* and *Campylobacter_B* were higher in the PEN and TET groups compared to CON, and their negative effect on growth during the nursery period may be due to their higher abundances in the antibiotic groups. Although treatment also modulated the association coefficients of *UBA6382* and *Mitsuokella* with ADGnursing decreasing the coefficient of *UBA6382* and increasing that of *Mitsuokella,* their relative abundances in piglets were not significantly affected by the sow antibiotic treatment. Therefore, the effect of treatment on growth through these genera cannot be interpreted directly. However, the results suggest that sow antibiotic treatment may have enhanced the negative impact of *UBA6382* and increased the positive impact of *Mitsuokella* on growth during the nursery period.

Turicibacter and *Clostridium_T* at ten weeks had significant negative association to ADGnursery, and treatment increased the coefficient, suggesting that treatment enhanced their negative impact on growth. The relative abundances of these genera were higher in PEN than CON, suggesting a link between the abundance and the negative effect. In contrast, *VUNA01* at three weeks had a positive association with growth during the nursery period, which was reduced by antibiotic treatment. Similar observation was made for *Alloprevotella* though its association with growth was weaker.

The positive association of *Faecousia* at three weeks with ADGnursing was increased and the negative association of *Limousia* with ADGnursing was decreased by inclusion of sow treatment in the model. These results suggest that sow antibiotic treatment enhanced the positive effect of *Faecousia* and intensified the negative effect of *Limousia* on growth. However, their relative abundances did not differ significantly between the treatment groups.

To our knowledge, associations of *Oliverpabstia* or *CAG-127* with piglet growth, or maternal antimicrobial influence on their abundances in piglet gut have not been reported earlier. However, we have observed a significantly higher relative abundances of *Mitsuokella* in high ADG weaning piglets in an earlier study (Mahmud MR, 2023). Moreover, a positive association of *Mitsuokella* with body weight and ADG in piglets has been reported (Yang et al., 2021). *Campylobacter* is a genus containing intestinal pathogens and it has been associated with post-weaning diarrhea in piglets (Yang et al., 2020; Zheng et al., 2023) with accompanying metabolic dysfunction (Yang et al., 2020). *Oliverpabstia* and *Faecousia* have been implicated in maternal obesity in a rat model (Li et al., 2025). *Turicibacter* belongs to the genera reported to be related to post-weaning transition of piglets (Yi 2025). The major ASV corresponding to *Clostridium_T* in our data was identified as *C. disporicum*, which is a saccharolytic commensal species (Horn, 1987) with pathogenic potential (Woo et al., 2005).

Limitations of the study

The sows were given antibiotics at days 78-80 of gestation, and the subsequent sampling was done only on day 113. Hence, any short-term alterations in microbiota, that were likely to have taken place following the treatment, could not be traced. Secondly, we employed 16S rRNA gene amplicon sequencing to assign taxonomy to near-genus level, which is less discriminatory than shotgun metagenomic sequencing. Lastly, the experiment was carried out on a commercial farm, with many environmental and farm management factors inducing experimental variation. This may have prevented detection of some treatment effects and associations due to insufficient statistical power. Despite these limitations, this research provides valuable insights into the effects of maternal antibiotic exposure on neonatal gut microbiota and lays the groundwork for future studies.

Conclusion

Our study highlights that maternal antibiotic use during pregnancy has a long-term impact on the gut microbiota of piglets and impairs growth, even if the maternal microbiota largely recovers by parturition. Significant treatment-related differences in the fecal microbial compositions were evident in the piglets, especially at ten weeks of age. This suggests that maternal antibiotic exposure affects not only the establishment of the initial gut microbiota but also its progressive development in the offspring.

Maternal antibiotic treatment in the beginning of the last third of pregnancy did not significantly alter the microbiota of sows at farrowing but impaired the dynamic development of the fecal microbiota over gestation, which may affect the early priming of the piglet immune system and subsequent microbial colonization. This study stresses the importance of cautious antibiotic use during pregnancy in livestock to protect offspring health and productivity. Future research should explore interventions such as probiotics or modulation to balance the microbiome in piglets.

Acknowledgements
We thank Kirsi Lahti for expert technical assistance. We acknowledge DNA Sequencing and Genomics Laboratory (supported by HiLIFE and Biocenter Finland funding), Institute of Biotechnology, University of Helsinki for sequencing, and CSC – IT Center for Science, Finland, for generous computational resources.

Disclosure statement
No potential conflict of interest was reported by the authors.

Funding

This work was supported by the Finnish Ministry of Agriculture and Forestry (project number VN/7587/2021), Research Council of Finland (project 347925), A-Producers Ltd, and Vetcare Ltd.

Author contributions

Conceptualization and design the study: M.H., M.N., K.A., S.J., T.P.M.; sample collection: M.R.M., M.H., K.A., E.K., S.J., M.N., T.P.M., H.N.K.; DNA extraction and quantification: M.R.M.; metadata preparation: M.H., K.A., M.N., M.R.M.; bioinformatics analysis, M.R.M & M.N.; statistical analysis: M.R.M., M.N., T.O, C.M., statistical analysis supervised M.P, M.L.C, M.N., T.O.; manuscript writing: M.R.M. T.P.M, T.O. C.M.; editing and reviewing manuscript: M.N, T.P.M, M.H., C.M., T.O, K.A., S.J., C.O., E.K. C.M; supervision, M.N., M.H., C.O. Approval of the final manuscript: All the authors read and approved the final manuscript.

Data availability

The 16S rRNA gene amplicon sequencing data will be available at ENA upon publication of the manuscript.

References

Alhasan, M.M., Cait, A.M., Heimesaat, M.M., Blaut, M., Klopfleisch, R., Wedel, A., Conlon, T.M., Yildirim, A.O., Sodemann, E.B., Mohn, W.W., Bereswill, S., Conrad, M.L., 2020. Antibiotic use during pregnancy increases offspring asthma severity in a dose-dependent manner. Allergy 75, 1979-1990.

Alhasan, M.M., Holsken, O., Duerr, C., Helfrich, S., Branzk, N., Philipp, A., Leitz, D., Duerr, J., Almousa, Y., Barrientos, G., Mohn, W.W., Gamradt, S., Conrad, M.L., 2023. Antibiotic use during pregnancy is linked to offspring gut microbial dysbiosis, barrier disruption, and altered immunity along the gut-lung axis. Eur J Immunol 53, e2350394.

Barnett, D., Arts, I., Penders, J., 2021. microViz: an R package for microbiome data visualization and statistics. Journal of Open Source Software 6.

Bastiaanssen, T.F.S., Gururajan, A., van de Wouw, M., Moloney, G.M., Ritz, N.L., Long-Smith, C.M., Wiley, N.C., Murphy, A.B., Lyte, J.M., Fouhy, F., Stanton, C., Claesson, M.J., Dinan, T.G., Cryan, J.F., 2021. Volatility as a Concept to Understand the Impact of Stress on the Microbiome. Psychoneuroendocrinology 124, 105047.

Bates, D., Mächler, M., Bolker, B., & Walker, S. , 2015. Fitting Linear Mixed-Effects models usinglme4. Journal of Statistical Software 67, 1–48.

Benjamini, Y., Hochberg Y., 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B (Methodological) 57, 289–300.

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J.E., Bittinger, K., Brejnrod, A., Brislawn, C.J., Brown, C.T., Callahan, B.J., Caraballo-Rodriguez, A.M., Chase, J., Cope, E.K., Da Silva, R., Diener, C., Dorrestein, P.C., Douglas, G.M., Durall, D.M., Duvallet, C., Edwardson, C.F., Ernst, M., Estaki, M., Fouquier, J., Gauglitz, J.M., Gibbons, S.M., Gibson, D.L., Gonzalez, A., Gorlick, K., Guo, J., Hillmann, B., Holmes, S., Holste, H., Huttenhower, C., Huttley, G.A., Janssen, S., Jarmusch, A.K., Jiang, L., Kaehler, B.D., Kang, K.B., Keefe, C.R., Keim, P., Kelley, S.T., Knights, D., Koester, I., Kosciolek, T., Kreps, J., Langille, M.G.I., Lee, J., Ley, R., Liu, Y.X., Loftfield, E., Lozupone, C., Maher, M., Marotz, C., Martin, B.D., McDonald, D., McIver, L.J., Melnik, A.V., Metcalf, J.L., Morgan, S.C., Morton, J.T., Naimey, A.T., Navas-Molina, J.A., Nothias, L.F., Orchanian, S.B., Pearson, T., Peoples, S.L., Petras, D., Preuss, M.L., Pruesse, E., Rasmussen, L.B., Rivers, A., Robeson, M.S., 2nd, Rosenthal, P., Segata, N., Shaffer, M., Shiffer, A., Sinha, R., Song, S.J., Spear, J.R., Swafford, A.D., Thompson, L.R., Torres, P.J., Trinh, P., Tripathi, A., Turnbaugh, P.J., Ul-Hasan, S., van der Hooft, J.J.J., Vargas, F., Vazquez-Baeza, Y., Vogtmann, E., von Hippel, M., Walters, W., Wan, Y., Wang, M., Warren, J., Weber, K.C., Williamson, C.H.D., Willis, A.D., Xu, Z.Z., Zaneveld, J.R., Zhang, Y., Zhu, Q., Knight, R., Caporaso, J.G., 2019. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol 37, 852-857.

Bosch, T.C., & McFall-Ngai, M., 2021. Animal development in the microbial world: Re-thinking the conceptual framework. Current Topics in Developmental Biology 399–427.

Brito Rodrigues, P., de Rezende Rodovalho, V., Sencio, V., Benech, N., Creskey, M., Silva Angulo, F., Delval, L., Robil, C., Gosset, P., Machelart, A., Haas, J., Descat, A., Goosens, J.F., Beury, D., Maurier, F., Hot, D., Wolowczuk, I., Sokol, H., Zhang, X., Ramirez Vinolo, M.A., Trottein, F., 2025. Integrative metagenomics and metabolomics reveal age-associated gut microbiota and metabolite alterations in a hamster model of COVID-19. Gut Microbes 17, 2486511.

Butler, J.E., Wertz, N., Sinkora, M., 2017. Antibody Repertoire Development in Swine. Annu Rev Anim Biosci 5, 255-279.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J., Holmes, S.P., 2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods 13, 581-583.

Champagne-Jorgensen, K., Mian, M.F., Kay, S., Hanani, H., Ziv, O., McVey Neufeld, K.A., Koren, O., Bienenstock, J., 2020. Prenatal low-dose penicillin results in long-term sex-specific changes to murine behaviour, immune regulation, and gut microbiota. Brain Behav Immun 84, 154-163.

Chen, C.M., Chou, H.C., Yang, Y.S.H., 2021. Maternal Antibiotic Treatment Disrupts the Intestinal Microbiota and Intestinal Development in Neonatal Mice. Front Microbiol 12, 684233.

Chen, W., Ma, J., Jiang, Y., Deng, L., Lv, N., Gao, J., Cheng, J., Liang, J.B., Wang, Y., Lan, T., Liao, X., Mi, J., 2022. Selective Maternal Seeding and Rearing Environment From Birth to Weaning Shape the Developing Piglet Gut Microbiome. Front Microbiol 13, 795101.

De Agüero, M.G., Ganal-Vonarburg, S. C., Fuhrer, T., Rupp, S., Uchimura, Y., Li, H., Steinert, A., Heikenwalder, M., Hapfelmeier, S., Sauer, U., McCoy, K. D., & Macpherson, A. J. , 2016. The maternal microbiota drives early postnatal innate immune development. Science 351, 1296–1302.

de Greeff, A., Schokker, D., Roubos-van den Hil, P., Ramaekers, P., Vastenhouw, S.A., Harders, F., Bossers, A., Smits, M.A., Rebel, J.M.J., 2020. The effect of maternal antibiotic use in sows on intestinal development in offspring. J Anim Sci 98.

Denamur, E., Clermont, O., Bonacorsi, S., Gordon, D., 2021. The population genetics of pathogenic Escherichia coli. Nat Rev Microbiol 19, 37-54.

Dierikx, T.H., Visser, D.H., Benninga, M.A., van Kaam, A., de Boer, N.K.H., de Vries, R., van Limbergen, J., de Meij, T.G.J., 2020. The influence of prenatal and intrapartum antibiotics on intestinal microbiota colonisation in infants: A systematic review. J Infect 81, 190-204.

Downes, J., Dewhirst, F.E., Tanner, A.C.R., Wade, W.G., 2013. Description of Alloprevotella rava gen. nov., sp. nov., isolated from the human oral cavity, and reclassification of Prevotella tannerae Moore et al. 1994 as Alloprevotella tannerae gen. nov., comb. nov. Int J Syst Evol Microbiol 63, 1214-1218.

Esser, D., Lange, J., Marinos, G., Sieber, M., Best, L., Prasse, D., Bathia, J., Ruhlemann, M.C., Boersch, K., Jaspers, C., Sommer, F., 2019. Functions of the Microbiota for the Physiology of Animal Metaorganisms. J Innate Immun 11, 393-404.

Gilroy, R., Ravi, A., Getino, M., Pursley, I., Horton, D.L., Alikhan, N.F., Baker, D., Gharbi, K., Hall, N., Watson, M., Adriaenssens, E.M., Foster-Nyarko, E., Jarju, S., Secka, A., Antonio, M., Oren, A., Chaudhuri, R.R., La Ragione, R., Hildebrand, F., Pallen, M.J., 2021. Extensive microbial diversity within the chicken gut microbiome revealed by metagenomics and culture. PeerJ 9, e10941.

Gonzalez-Perez, G., Hicks, A.L., Tekieli, T.M., Radens, C.M., Williams, B.L., Lamouse-Smith, E.S., 2016. Maternal Antibiotic Treatment Impacts Development of the Neonatal Intestinal Microbiome and Antiviral Immunity. J Immunol 196, 3768-3779.

Guevarra, R.B., Hong, S.H., Cho, J.H., Kim, B.R., Shin, J., Lee, J.H., Kang, B.N., Kim, Y.H., Wattanaphansak, S., Isaacson, R.E., Song, M., Kim, H.B., 2018. The dynamics of the piglet gut microbiome during the weaning transition in association with health and nutrition. J Anim Sci Biotechnol 9, 54.

Guevarra, R.B., Lee, J.H., Lee, S.H., Seok, M.J., Kim, D.W., Kang, B.N., Johnson, T.J., Isaacson, R.E., Kim, H.B., 2019. Piglet gut microbial shifts early in life: causes and effects. J Anim Sci Biotechnol 10, 1.

Heinritz, S.N., Mosenthin, R., Weiss, E., 2013. Use of pigs as a potential model for research into dietary modulation of the human gut microbiota. Nutr Res Rev 26, 191-209.

Holman, D.B., Chenier, M.R., 2015. Antimicrobial use in swine production and its effect on the swine gut microbiota and antimicrobial resistance. Can J Microbiol 61, 785-798.

Holman DB, K.A., Tingley JP, Abbott DW. , 2022. Novel Insights into the Pig Gut Microbiome Using Metagenome-Assembled Genomes. Microbiol Spectr. 10.

Horn, N., 1987. Clostridium disporicum sp. nov. , a Saccharolytic Species Able to Form Two Spores per Cell, Isolated from a Rat Cecum. Int J Syst Bacteriol 37, 398–401.

Husso, A., Pessa-Morikawa, T., Koistinen, V.M., Karkkainen, O., Kwon, H.N., Lahti, L., Iivanainen, A., Hanhineva, K., Niku, M., 2023. Impacts of maternal microbiota and microbial metabolites on fetal intestine, brain, and placenta. BMC Biol 21, 207.

Ivarsson, E., Roos, S., Liu, H.Y., Lindberg, J.E., 2014. Fermentable non-starch polysaccharides increases the abundance of Bacteroides-Prevotella-Porphyromonas in ileal microbial community of growing pigs. Animal 8, 1777-1787.

Kaisanlahti, A., Turunen, J., Byts, N., Samoylenko, A., Bart, G., Virtanen, N., Tejesvi, M.V., Zhyvolozhnyi, A., Sarfraz, S., Kumpula, S., Hekkala, J., Salmi, S., Will, O., Korvala, J., Paalanne, N., Erawijantari, P.P., Suokas, M., Medina, T.P., Vainio, S., Medina, O.P., Lahti, L., Tapiainen, T., Reunanen, J., 2023. Maternal microbiota communicates with the fetus through microbiota-derived extracellular vesicles. Microbiome 11, 249.

Kaiser, M.Ø., 2019. Postpartum dysgalactia syndrome in sows - Pathogenesis and early diagnosis. PhD thesis. University of Copenhagen

Kimura, I., Miyamoto, J., Ohue-Kitano, R., Watanabe, K., Yamada, T., Onuki, M., Aoki, R., Isobe, Y., Kashihara, D., Inoue, D., Inaba, A., Takamura, Y., Taira, S., Kumaki, S., Watanabe, M., Ito, M., Nakagawa, F., Irie, J., Kakuta, H., Shinohara, M., Iwatsuki, K., Tsujimoto, G., Ohno, H., Arita, M., Itoh, H., Hase, K., 2020. Maternal gut microbiota in pregnancy influences offspring metabolic phenotype in mice. Science 367.

Kong, Q., Zhang, W., An, M., Kulyar, M.F., Shang, Z., Tan, Z., Xu, Y., Li, J., Liu, S., 2022. Characterization of Bacterial Microbiota Composition in Healthy and Diarrheal Early-Weaned Tibetan Piglets. Front Vet Sci 9, 799862.

Kong, X.F., Ji, Y.J., Li, H.W., Zhu, Q., Blachier, F., Geng, M.M., Chen, W., Yin, Y.L., 2016. Colonic luminal microbiota and bacterial metabolite composition in pregnant Huanjiang mini-pigs: effects of food composition at different times of pregnancy. Sci Rep 6, 37224.

Koren, O., Goodrich, J.K., Cullender, T.C., Spor, A., Laitinen, K., Backhed, H.K., Gonzalez, A., Werner, J.J., Angenent, L.T., Knight, R., Backhed, F., Isolauri, E., Salminen, S.,

Ley, R.E., 2012. Host remodeling of the gut microbiome and metabolic changes during pregnancy. Cell 150, 470-480.

Ksiezarek, M., Grosso, F., Ribeiro, T.G., Peixe, L., 2022. Genomic diversity of genus Limosilactobacillus. Microb Genom 8.

Kuznetsova, A., Brockhoff, P.B., Christensen, R.H.B., 2017. lmerTest Package: Tests in Linear Mixed Effects Models. Journal of Statistical Software 82.

Lahti, L., Shetty, S. 2017. Tools for microbiome analysis in R. Microbiome package version 1.23.1. (Bioconductor).

Li, J., Feng, S., Pi, Y., Jiang, X., Li, X., Zhou, Z., Liu, X., Wei, H., Tao, S., 2023. Limosilactobacillus johnsoni and Limosilactobacillus mucosae and Their Extracellular Vesicles Alleviate Gut Inflammatory Injury by Mediating Macrophage Polarization in a Lipopolysaccharide-Challenged Piglet Model. J Nutr 153, 2497-2511.

Li, X., Zhao, Y., Liu, M., Zhang, M., Zhu, Z., Liao, A., Pan, L., Lv, X., Liu, F., Huang, J., 2025. Fermented Wheat Germ Ameliorates High-Fat Diet-Induced Maternal Obesity in Rats: Insights from Microbiome and Metabolomics. J Agric Food Chem 73, 12666-12678.

Li, Y., Liu, Y., Ma, Y., Ge, X., Zhang, X., Cai, C., Yang, Y., Lu, C., Liang, G., Guo, X., Cao, G., Li, B., Gao, P., 2022. Effects of Maternal Factors and Postpartum Environment on Early Colonization of Intestinal Microbiota in Piglets. Front Vet Sci 9, 815944.

Lima, E., Hyde, R., Green, M., 2021. Model selection for inferential models with high dimensional data: synthesis and graphical representation of multiple techniques. Sci Rep 11, 412.

Liu, H., Zeng, X., Zhang, G., Hou, C., Li, N., Yu, H., Shang, L., Zhang, X., Trevisi, P., Yang, F., Liu, Z., Qiao, S., 2019. Maternal milk and fecal microbes guide the spatiotemporal development of mucosa-associated microbiota and barrier function in the porcine neonatal gut. BMC Biol 17, 106.

Liu, Z., Tsai, T., Zuo, B., Howe, S., Farrar, J.E., Randolph, C.E., Maxwell, C.V., Zhao, J., 2025. The sow vaginal and gut microbiota associated with longevity and reproductive performance. J Anim Sci Biotechnol 16, 6.

Luo Y, R.W., Smidt H, Wright AG, Yu B, Schyns G, McCormack UM, Cowieson AJ, Yu J, He J, Yan H, Wu J, Mackie RI, Chen D. , 2022. Dynamic Distribution of Gut Microbiota in Pigs at Different Growth Stages: Composition and Contribution. Microbiol Spectr. 10.

Lynch, J.B., Gonzalez, E.L., Choy, K., Faull, K.F., Jewell, T., Arellano, A., Liang, J., Yu, K.B., Paramo, J., Hsiao, E.Y., 2023. Gut microbiota Turicibacter strains differentially modify bile acids and host lipids. Nat Commun 14, 3669.

Ma, C., Gao, Q., Zhang, W., Zhu, Q., Tang, W., Blachier, F., Ding, H., Kong, X., 2020. Supplementing Synbiotic in Sows' Diets Modifies Beneficially Blood Parameters and Colonic Microbiota Composition and Metabolic Activity in Suckling Piglets. Front Vet Sci 7, 575685.

Ma, J., Wu, H., Ma, Z., Wu, Z., 2025. Bacterial and host factors involved in zoonotic Streptococcal meningitis. Microbes Infect 27, 105335.

Mach, N., Berri, M., Estelle, J., Levenez, F., Lemonnier, G., Denis, C., Leplat, J.J., Chevaleyre, C., Billon, Y., Dore, J., Rogel-Gaillard, C., Lepage, P., 2015. Early-life establishment of the swine gut microbiome and impact on host phenotypes. Environ Microbiol Rep 7, 554-569.

Madany, A.M., Hughes, H.K., Ashwood, P., 2022a. Antibiotic Treatment during Pregnancy Alters Offspring Gut Microbiota in a Sex-Dependent Manner. Biomedicines 10.

Madany, A.M., Hughes, H.K., Ashwood, P., 2022b. Prenatal Maternal Antibiotics Treatment Alters the Gut Microbiota and Immune Function of Post-Weaned Prepubescent Offspring. Int J Mol Sci 23.

Mahmud MR, J.C., Uddin MK, Huhtinen M, Salonen A, Peltoniemi O, Venhoranta H, Oliviero C. Impact of Intestinal Microbiota on Growth Performance of Suckling and Weaned Piglets., 2023. Impact of Intestinal Microbiota on Growth Performance of Suckling and Weaned Piglets. Microbiol Spectr 11e, 03744-03722.

Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal 17, 10-12.

Martinez Arbizu, P. 2020. pairwiseAdonis: Pairwise multilevel comparison using adonis. R package version 0.4.

McDonald, D., Jiang, Y., Balaban, M., Cantrell, K., Zhu, Q., Gonzalez, A., Morton, J.T., Nicolaou, G., Parks, D.H., Karst, S.M., Albertsen, M., Hugenholtz, P., DeSantis, T., Song, S.J., Bartko, A., Havulinna, A.S., Jousilahti, P., Cheng, S., Inouye, M., Niiranen, T., Jain, M., Salomaa, V., Lahti, L., Mirarab, S., Knight, R., 2024. Greengenes2 unifies microbial data in a single reference tree. Nat Biotechnol 42, 715-718.

McMurdie, P.J., Holmes, S., 2013. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. PLoS One 8, e61217.

Meinshausen, N., Bühlmann, P., 2010. Stability Selection. Journal of the Royal Statistical Society Series B: Statistical Methodology 72, 417-473.

Miyoshi, J., Hisamatsu, T., 2025. Effect of maternal exposure to antibiotics during pregnancy on the neonatal intestinal microbiome and health. Clin J Gastroenterol 18, 1-10.

Mueller, N.T., Bakacs, E., Combellick, J., Grigoryan, Z., Dominguez-Bello, M.G., 2015. The infant microbiome development: mom matters. Trends Mol Med 21, 109-117.

Nickols, W.A., Kuntz, T., Shen, J., Maharjan, S., Mallick, H., Franzosa, E.A., Thompson, K.N., Nearing, J.T., Huttenhower, C., 2024. MaAsLin 3: Refining and extending generalized multivariable linear models for meta-omic association discovery. bioRxiv.

Norring, M., Valros, A., Bergman, P., Marchant, J.N., Heinonen, M., 2019. Body condition, live weight and success in agonistic encounters in mixed parity groups of sows during gestation. Animal 13, 392-398.

Pabst, O., Cerovic, V., Hornef, M., 2016. Secretory IgA in the Coordination of Establishment and Maintenance of the Microbiota. Trends Immunol 37, 287-296.

Pabst, R., 2020. The pig as a model for immunology research. Cell Tissue Res 380, 287-304.

Pantazi, A.C., Mihai, C.M., Lupu, A., Balasa, A.L., Chisnoiu, T., Mihai, L., Frecus, C.E., Ungureanu, A., Chirila, S.I., Nori, W., Lupu, V.V., Stoicescu, R.M., Baciu, G., Cambrea, S.C., 2025. Gut Microbiota Profile and Functional Gastrointestinal Disorders in Infants: A Longitudinal Study. Nutrients 17.

Piirainen, V., Konig, E., Husso, A., Heinonen, M., Iivanainen, A., Pessa-Morikawa, T., Niku, M., 2025. Bacterial profiles of the oral, vaginal, and rectal mucosa and colostrum of periparturient sows. PLoS One 20, e0317513.

Rideout, J.R., Chase, J.H., Bolyen, E., Ackermann, G., Gonzalez, A., Knight, R., Caporaso, J.G., 2016. Keemei: cloud-based validation of tabular bioinformatics file formats in Google Sheets. Gigascience 5, 27.

Rosero, J.A., Killer, J., Sechovcova, H., Mrazek, J., Benada, O., Fliegerova, K., Havlik, J., Kopecny, J., 2016. Reclassification of Eubacterium rectale (Hauduroy et al. 1937)

Prevot 1938 in a new genus Agathobacter gen. nov. as Agathobacter rectalis comb. nov., and description of Agathobacter ruminis sp. nov., isolated from the rumen contents of sheep and cows. Int J Syst Evol Microbiol 66, 768-773.

Rumeau, M., Chadi, S., Pepke, F., Beaumont, M., Vicente, C.M., Juppeau, A., Vandecasteele, C., Cauquil, L., Pascal, G., Langella, P., Knudsen, C., Combes, S., Martin, R., 2025. Structure-dependent degradation of milk oligosaccharides by newly isolated intestinal commensal bacterial strains from suckling piglets and rabbits. BMC Microbiol 25, 513.

Saladrigas-Garcia, M., Sola-Oriol, D., Lopez-Verge, S., D'Angelo, M., Collado, M.C., Nielsen, B., Faldyna, M., Perez, J.F., Martin-Orue, S.M., 2022. Potential effect of two Bacillus probiotic strains on performance and fecal microbiota of breeding sows and their piglets. J Anim Sci 100.

Schokker, D., Zhang, J., Vastenhouw, S.A., Heilig, H.G., Smidt, H., Rebel, J.M., Smits, M.A., 2015. Long-lasting effects of early-life antibiotic treatment and routine animal handling on gut microbiota composition and immune system in pigs. PLoS One 10, e0116523.

Sinkora, M., Butler, J.E., 2009. The ontogeny of the porcine immune system. Dev Comp Immunol 33, 273-283.

Strömpl, C., Tindall, B. J., Jarvis, G. N., Lünsdorf, H., Moore, E. R., & Hippe, H. , 1999. A re-evaluation of the taxonomy of the genus Anaerovibrio, with the reclassification of Anaerovibrio glycerini as Anaerosinus glycerini gen. nov., comb. nov., and Anaerovibrio burkinabensis as Anaeroarcus burkinensis [corrig.] gen. nov., comb. nov. International journal of systematic bacteriology 49, 1861–1872.

Torow, N., Hand, T.W., Hornef, M.W., 2023. Programmed and environmental determinants driving neonatal mucosal immune development. Immunity 56, 485-499.

Wang, K., Hu, C., Tang, W., Azad, M.A.K., Zhu, Q., He, Q., Kong, X., 2021. The Enhancement of Intestinal Immunity in Offspring Piglets by Maternal Probiotic or Synbiotic Supplementation Is Associated With the Alteration of Gut Microbiota. Front Nutr 8, 686053.

Wang, K., Zhu, Q., Kong, X., Song, M., Azad, M.A.K., Xiong, L., Zheng, Y., He, Q., 2020. Dietary Probiotics or Synbiotics Supplementation During Gestation, Lactation, and Nursery Periods Modifies Colonic Microbiota, Antioxidant Capacity, and Immune Function in Weaned Piglets. Front Vet Sci 7, 597832.

Woo, P.C., Lau, S.K., Chan, K.M., Fung, A.M., Tang, B.S., Yuen, K.Y., 2005. Clostridium bacteraemia characterised by 16S ribosomal RNA gene sequencing. J Clin Pathol 58, 301-307.

Wu, J., Lin, Z., Wang, X., Zhao, Y., Zhao, J., Liu, H., Johnston, L. J., Lu, L., & Ma, X., 2022. Limosilactobacillus reuteri SLZX19-12 Protects the Colon from Infection by Enhancing Stability of the Gut Microbiota and Barrier Integrity and Reducing Inflammation. . Microbiol Spectr. 10.

Wylensek, D., Hitch, T.C.A., Riedel, T., Afrizal, A., Kumar, N., Wortmann, E., Liu, T., Devendran, S., Lesker, T.R., Hernandez, S.B., Heine, V., Buhl, E.M., P, M.D.A., Cumbo, F., Fischoder, T., Wyschkon, M., Looft, T., Parreira, V.R., Abt, B., Doden, H.L., Ly, L., Alves, J.M.P., Reichlin, M., Flisikowski, K., Suarez, L.N., Neumann, A.P., Suen, G., de Wouters, T., Rohn, S., Lagkouvardos, I., Allen-Vercoe, E., Sproer, C., Bunk, B., Taverne-Thiele, A.J., Giesbers, M., Wells, J.M., Neuhaus, K., Schnieke, A., Cava, F., Segata, N., Elling, L., Strowig, T., Ridlon, J.M., Gulder, T.A.M., Overmann, J., Clavel, T., 2020. A collection of bacterial isolates from the pig intestine reveals functional and taxonomic diversity. Nat Commun 11, 6389.

Yang, G., Yan, Y., Zhang, L., Ruan, Z., Hu, X., Zhang, S., Li, X., 2020. Porcine circovirus type 2 (PCV2) and Campylobacter infection induce diarrhea in piglets: Microbial dysbiosis and intestinal disorder. Anim Nutr 6, 362-371.

Yang, Y., Gao, H., Ran, J., Huang, Y., Yang, M., Yang, L., Zhao, S., Qiao, S., Pan, H., 2021. Comparison of early-life jejunal microbiota diversity in two pig breeds. Czech Journal of Animal Science 66, 459-469.

Zhang, M., Differding, M.K., Benjamin-Neelon, S.E., Ostbye, T., Hoyo, C., Mueller, N.T., 2019. Association of prenatal antibiotics with measures of infant adiposity and the gut microbiome. Ann Clin Microbiol Antimicrob 18, 18.

Zheng, X., Nie, K., Xu, Y., Zhang, H., Xie, F., Xu, L., Zhang, Z., Ding, Y., Yin, Z., Zhang, X., 2023. Fecal Microbial Structure and Metabolic Profile in Post-Weaning Diarrheic Piglets. Genes (Basel) 14.

Zhu, Q., Azad, M. A. K., Li, R., Li, C., Liu, Y., Yin, Y., & Kong, X. , 2024. Dietary probiotic and synbiotic supplementation starting from maternal gestation improves muscular lipid metabolism in offspring piglets by reshaping colonic microbiota and metabolites. . mSystems 9, e0004824.

Zou, Z.H., Liu, D., Li, H.D., Zhu, D.P., He, Y., Hou, T., Yu, J.L., 2018. Prenatal and postnatal antibiotic exposure influences the gut microbiota of preterm infants in neonatal intensive care units. Ann Clin Microbiol Antimicrob 17, 9.