1 **Breaking the "one nucleus, one whole genome" rule: *Neurospora crassa***
2 **separates its haploid chromosomes into different nuclei**
3
4
5

Jinyi Tan[1,2], Yan Xu[3], Lei Tian[1,2], Xuanye Chen[1,2], Monika Fischer[2], Yuelin Zhang[3] and Xin

Li[1,2]*

[1] Michael Smith Laboratories, University of British Columbia, Vancouver BC V6T 1Z4, Canada

[2] Department of Botany, University of British Columbia, Vancouver BC V6T 1Z4, Canada

[3] The College of Life Sciences, Sichuan University, Chengdu, 610064, Sichuan, China.

*For correspondence: xinli@msl.ubc.ca

Abstract

Neurospora crassa has long served as a key fungal genetics model, facilitating discoveries like

the "one gene, one enzyme" hypothesis. It produces many conidia, mostly with two or three

nuclei and each nucleus was once thought to contain a full haploid genome. Using flow

cytometry and FISH, we found that *N. crassa* haploid chromosomes are divided among nuclei,

each conidium is haploid – challenging the "one nucleus, one whole genome" rule.

Main

Nuclei define eukaryotes. They assist the separation of genetic materials and gene expressions. It has been established in molecular genetics and cell biology that each nucleus contains at least one complete set of chromosomes necessary for proper genomic function and cell development, the "one nucleus, one whole genome" rule. However, our recent study challenges this long-standing principle. In two plant pathogenic fungi belonging to the Sclerotiniaceae family, soilborne *Sclerotinia sclerotiorum* and airborne *Botrytis cinerea*, we uncovered that their haploid chromosomes are separated among different nuclei (1). Such finding prompted the next question on how widespread such phenomenon is. As both *S. sclerotiorum* and *B. cinerea* have multinuclear spores but also produce uninuclear spermatia that are not viable likely due to chromosome losses, we searched the literature to identify fungi with bi- or multinuclear spores and frequent nonviable uninuclear cells, which are potential candidates that may also be breaking the "one nucleus, one whole genome" rule. *Neurospora crassa* came out of our literature analysis.

N. crassa is a non-pathogenic ascomycete commonly known as bread mold. It has been serving as a classical model organism in fungal genetics and molecular biology (2,3). It was the first filamentous fungus fully sequenced, making it a preferred model for genome-wide experimental approaches (4,5). Its genome is approximately 40 Mb in size and comprises 7 chromosomes (4–7).

N. crassa can be easily recognized by their abundant orange conidial spores, which are typically multinucleate, majority of which have 2 to 3 nuclei (8,9) (Figure 1A). Although it does produce

uninuclear conidia as well, less than 15% of them are viable likely due to incomplete

chromosome sets (10,11). To quantify the DNA amount in each nucleus of *N. crassa*, we

employed flow cytometry through fluorescence-activated cell sorting (FACS) using purified

nuclei from mycelia. Haploid and diploid *Saccharomyces cerevisiae* (baker's yeast) strains

BY4741 and BY4743 were used as controls. Based on the relative fluorescence intensity (RFI)

of nuclei (Figure 1B and C), each *N. crassa* mycelial nucleus contained about a quarter of its

total genome, suggesting that each nucleus within *N. crassa* mycelia harbors only a subset of its

full genome (Figure 1C).

**Figure 1. *N. crassa* conidia harbor multiple nuclei and each nucleus only contain parts of the total haploid
genome.**

(A) Quantification of nuclei number in the conidium of *N. crassa* (left). The fluorescence microscopy images show
the nuclei in conidia of the *N. crassa* histone H1-eGFP strain (right). Scale bar, 10 μm.
(B) Propidium iodide (PI) fluorescence intensity of sorted purified nuclei from *Saccharomyces cerevisiae* BY4741
(haploid yeast), BY4743 (diploid yeast), mycelia of *N. crassa*. One hundred nuclei from each strain were sorted into

each well of a 96-well plate by a Sony MA900 cell sorter. Data from 15 wells were collected for each strain. The PI
fluorescence intensity values were calculated based on the mean fluorescence intensity of each individual well. Error
bars indicate the standard deviation of the values from 15 wells.
(C) Calculated DNA content per nucleus in *S. cerevisiae* BY4741 (haploid), BY4743 (diploid) and *N. crassa*. The
known genome sizes of these strains are 12, 24, and 40 Mb, respectively. The DNA content in each nucleus was
calculated by multiplying the genome size of the haploid yeast strain BY4741 by the fold-change values derived
from data in Fig B. Error bars indicate the standard deviation ($n = 15$).
67

To further test whether each nucleus contains a full set of the haploid chromosomes in *N. crassa*

conidia, we performed fluorescence in situ hybridization (FISH) using a chromosome 1 (Chr1)-

specific probe (Supplementary figure 1A). As shown in Figure 2, hybridization signals were

consistently detected in only one nucleus per conidium. Notably, Chr1-specific signals were

never detected in multiple nuclei within the same conidium (Figure 2A and C). As a control, a

non-specific probe that is present in all chromosomes (Supplementary figure 1B) yielded signals

in all nuclei with positive hybridizations, confirming that probe accessibility was not limiting

(Figure 2B and C). These FISH results support that the chromosome inside the nuclei of the bi-

or tri- nuclear conidium are different. Together with the flow cytometry results, each *N. crassa*

conidium seems to carry a complete set of haploid chromosomes distributed among one or

multiple nuclei. The conidia are thus haploid regardless of their nuclei number. This feature will

offer a major practical advantage. For gene replacement experiments to knock out or modify

target genes, protoplasts made from conidia can be used, skipping the step of uninucleate

microconidia purification in current methods. The same with mutagenesis for forward genetic

analyses. Just as with *S. sclerotiorum*, binucleate ascospores or multinuclear conidia can be used

with UV or chemical mutagens to generate pure haploid mutants (12).

Probe	No. of conidia without probe signal	No. of conidia with probe signal in one nucleus	No. of conidia with probe signals in two or more nuclei	Total numbers of conidia
Chr1-specific probe	20	122	0	144

Probe	No. of conidia without probe signal	No. of conidia with probe signals in over half of nucleus	No. of conidia with probe signals in all nuclei	Total numbers of conidia
Non-specific probe	15	29	79	123

Figure 2. A *N. crassa* conidium can separate its haploid chromosomes into one, two, or three nuclei.

(A) FISH signals of Chr1-specific (left) and non-specific (right) probes in representative conidia. Blue and red
signals are from DAPI-stained nuclei and hybridized probe, respectively. The hybridized Cy5 probe signals are
pointed out by arrows. Scale bars, 5 μm.
(B) Summary of FISH results for Chr1-specific (top) and non-specific (bottom) probes in one or multiple nuclei of
examined conidia.

Interestingly, although uninucleate *N. crassa* conidia (microconidia) are viable, they exhibit a

very low germination rate (10,11), whereas uninucleate spermatia cells in *S. sclerotiorum* and *B.*

cinerea are nonviable. This led to the improvement of *N. crassa* mutagenesis protocols where

uninucleate spores were firstly purified before mutagenesis, reinforcing the "one nucleus, one

whole genome" dogma (11,13). We believe the reason why a small fraction of the uninucleate

cells is viable in *N. crassa* is probably due to its low haploid chromosome number. Its 7

chromosomes may occasionally be packed into just one nucleus, allowing the spore to be viable.

As for *S. sclerotiorum* and *B. cinerea*, 16 or 18 chromosomes are seemingly almost impossible to

be packed simultaneously into a single nucleus during their nuclear division.

In summary, our findings in *N. crassa* suggest that dividing haploid chromosomes across

multiple nuclei may be more widespread than previously recognized. It is not restricted to

sclerotia-forming plant fungal pathogens of the Sclerotiniaceae family. Other Ascomycetes and

Basidiomycetes with known bi- or multi-nucleate ascospores and basidiospores, such as

Morchella, Poronia, Oudemansiella and *Cortinarius*, also warrant further investigation (14–18).

Studying these organisms could provide important insights into the spread of this phenomenon

and advance our understanding of chromosome biology and genome evolution. Moreover, the

precise molecular mechanisms underlying such chromosome splitting phenomenon remain

puzzling and represent an intriguing area for future research. For instance, how do different

nuclei housing the full complement of chromosomes communicate and coordinate with each

other? What are the key genes and proteins involved in nuclear coordination? Are there special

checkpoints to ensure proper chromosomal distribution? Advanced live-cell imaging and single-

nuclei sequencing could be particularly valuable for tracking chromosomal segregation and

assessing genetic variation or stability across multiple nuclei.

Methods

Relative fluorescence intensity (RFI) determination by flow cytometry

Flow cytometry was performed following the previous study using nuclei from *Saccharomyces*

cerevisiae and mycelia of *N. crassa* (1).

FISH analyses

Through BLAST searches of the *N. crassa* genome, a chromosome 1 (Chr1)-specific probe was

identified, which is located in a non-coding region with a single successive hybridization site

(Supplementary figure 1A). Additionally, a non-specific repetitive probe was identified that can

target all chromosomes (Supplementary figure 1B).

Conidia fixation and FISH were performed following the previous study with minor changes (1).

100 ng of probe was used. The hybridization was performed overnight at 37 to 42°C for non-

specific probe, and at 45°C for 22 hours for the Chr1-specific probe. Nuclei were stained with 1

µg/ml DAPI before imaging.

Microscopy observation

Images of the Histone-GFP strain and FISH samples were performed with the ZEISS AXIO

Imager M2 microscope at 200× or 1000× magnification. Scale bars are shown in each image and

specified in the figure legends.

138
139 **Supplementary figure 1. Chromosomal locations of the specific and non-specific FISH probes.**

140 **(A)-(B)** The chromosomal regions in *N. crass* where the Chr1-specific (A), and non-specific (B) probes can
141 hybridize to are indicated by red triangles. The Chr1-specific probe occurs once, while the non-specific probe is
142 repeated from seven to eleven times across different chromosomes. The probe sequences are shown at the bottom of
143 each panel. Scale bars showing chromosome lengths are shown on the right.

References

1. Xu, Y., Tian, L., Tan, J., Huang, W., Li, J., O'Neil, N., Hirst, M., Hieter, P., Zhang, Y., &
Li, X. Distribution of haploid chromosomes into separate nuclei in two pathogenic fungi.
Science **388**, 784–788 (2025). https://doi.org/10.1126/science.abn7811
2. Davis, R. H., & Perkins, D. D. Neurospora: A model of model microbes. *Nature Reviews*
Genetics **3**, 397–403 (2002). https://doi.org/10.1038/nrg797

3.   Seiler, S., & Plamann, M. The Genetic Basis of Cellular Morphogenesis in the Filamentous
Fungus Neurospora crassa. *Molecular Biology of the Cell* **14**, 4352–4364 (2003).
https://doi.org/10.1091/mbc.e02-07-0433
4.   Schulte, U., Becker, I., Mewes, H. W., & Mannhaupt, G. Large scale analysis of sequences
from Neurospora crassa. *Journal of Biotechnology* **94**, 3–13 (2002).
https://doi.org/10.1016/S0168-1656(01)00415-1
5.   Galagan, J. E., Calvo, S. E., Borkovich, K. A., Selker, E. U., Read, N. O., Jaffe, D.,
FitzHugh, W., Ma, L. J., Smirnov, S., Purcell, S., Rehman, B., Elkins, T., Engels, R., Wang,
S., Nielsen, C. B., Butler, J., Endrizzi, M., Qui, D., Ianakiev, P., … Birren, B. The genome
sequence of the filamentous fungus Neurospora crassa. *Nature* **422**, 859–868 (2003).
https://doi.org/10.1038/nature01554
6.   Selker, E. U. Neurospora crassa. *Reference Module in Life Sciences* , (2017).
https://doi.org/10.1016/b978-0-12-809633-8.06790-x
7.   Kelkar, H. S., Griffith, J., Case, M. E., Covert, S. F., Hall, R. D., Keith, C. H., Oliver, J. S.,
Orbach, M. J., Sachs, M. S., Wagner, J. R., Weise, M. J., Wunderlich, J. K., & Arnold, J.
The Neurospora crassa genome: Cosmid libraries sorted by chromosome. *Genetics* **157**,
979–990 (2001). https://doi.org/10.1093/genetics/157.3.979
9.   Roca, M. G., Kuo, H. C., Lichius, A., Freitag, M., & Read, N. D. Nuclear dynamics,
mitosis, and the cytoskeleton during the early stages of colony initiation in Neurospora
crassa. *Eukaryotic Cell* **9**, 1171–1183 (2010). https://doi.org/10.1128/EC.00329-09
10.   Schmit, J. C., & Brody, S. Biochemical genetics of Neurospora crassa conidial germination.
Bacteriological Reviews **40**, 1–41 (1976). https://doi.org/10.1128/mmbr.40.1.1-41.1976
11.   Maheshwari, R. Microconidia of Neurospora crassa. *Fungal Genetics and Biology* **26**, 1–18
(1999). https://doi.org/10.1006/fgbi.1998.1103
12.   Xu, Y., Ao, K., Tian, L., Qiu, Y., Huang, X., Liu, X., Hoy, R., Zhang, Y., Rashid, K. Y.,
Xia, S., & Li, X. A forward genetic screen in Sclerotinia sclerotiorum revealed the
transcriptional regulation of its sclerotial melanization pathway. *Molecular Plant-Microbe*
Interactions **35**, 244–256 (2022). https://doi.org/10.1094/MPMI-10-21-0254-R
13.   Ebbole, D., & Sachs, M. S. A rapid and simple method for isolation of Neurospora crassa
homokaryons using microconidia. *Fungal Genetics Reports* **37**. (1990).
https://doi.org/10.4148/1941-4765.1472
14.   Horton, T. R. The number of nuclei in basidiospores of 63 species of ectomycorrhizal
Homobasidiomycetes. *Mycologia* **98**, 233–238 (2006).
https://doi.org/10.1080/15572536.2006.11832695
15.   Rebecca, R., Gao, Q., Cui, Y., Rong, C., Liu, Y., Zhao, W., Kumara, W., & Wang, S.
Nuclear conditions of basidiospores and hyphal cells in the edible mushroom
Oudemansiella aparlosarca. *MicrobiologyOpen* **10**, 1–11 (2021).
https://doi.org/10.1002/mbo3.1233
16.   Rogers, J. D. Cytology of Poronia oedipus and P. punctata. *Canadian Journal of Botany* **48**,
1665–1668 (1970). https://doi.org/10.1139/b70-246
17.   Du, X. H., Wang, S. Y., Ryberg, M., Guo, Y. J., Wei, J. Y., Pfister, D. H., & Johannesson,
H. Cytological studies reveal high variation in ascospore number and shape and conidia
produced directly from ascospores in Morchella galilaea. *Frontiers in Microbiology* **14**, 1–
20 (2023). https://doi.org/10.3389/fmicb.2023.1286501
18.   Liu, W., He, P., Zhang, J., Wu, L., Er, L., Shi, X., Gu, Z., Yu, F., & Pérez-Moreno, J.
Ultrastructure and Physiological Characterization of Morchella Mitospores and Their

Relevance in the Understanding of the Morel Life Cycle. *Microorganisms* **11**. (2023).
https://doi.org/10.3390/microorganisms11020345
203

Acknowledgements

We thank Josh Li for assistance with the cultivation of *N. crassa* conidia. ChatGPT 5.0 (OpenAI,

San Francisco, CA, USA) was used for correcting grammar mistakes and language polishing.

This work was supported by funds to X.L. from the Canadian Natural Sciences and Engineering

Research Council (NSERC) Discovery program, NSERC-CREATE-PRoTECT, Canada

Research Chair (CRC), and the Canadian Foundation for Innovation (CFI) funds. J.T. is partly

supported by a University of British Columbia Four-year Fellowship.

Author Contributions

213 X.L. conceived the idea; J.T. and X.L. wrote the manuscript draft; J.T. and Y.X. conducted the

214 experiments. L.T. and Y.Z. revised the original manuscript. All authors edited and approved the

215 final manuscript.

Competing financial interests

217 The authors declare no competing interests.

Correspondence and requests for materials should be addressed to XL.

A Chr1-specific probe

Bright Field | DAPI | Cy5 | Merged

B Non-specific probe

Bright Field | DAPI | Cy5 | Merged

C

Probe	No. of conidia without probe signal	No. of conidia with probe signal in one nucleus	No. of conidia with probe signals in two or more nuclei	Total numbers of conidia
Chr1-specific probe	20	122	0	144

Probe	No. of conidia without probe signal	No. of conidia with probe signals in over half of nucleus	No. of conidia with probe signals in all nuclei	Total numbers of conidia
Non-specific probe	15	29	79	123