Structural basis and physiological significance of **non-canonical** Gs coupling to the prototypical Gi-coupled melatonin MT$_1$ receptor

Atsuro Oishi*[1,2]✉,, Hiroyuki H. Okamoto*[3], Keisuke Ikegami*[4,5], Ronan McHugh[6], Bernard Masri[1], Tsukasa Kusakizako[3], Kazuhiro Kobayashi[3], Angeliki Karamitri[1], Erika Cecon[1], Julie Dam[1], Miki Nagase[2], Irina G. Tikhonova[6], Osamu Nureki[3,7,✉], and Ralf Jockers[1,7,✉],

Affiliations
[1] Université de Paris, Institut Cochin, INSERM, CNRS, F-75014 PARIS, France
[2] Department of Anatomy, Kyorin University School of Medicine, 181-8611, Tokyo, Japan
[3] Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Bunkyo-ku, Tokyo, Japan.
[4] Department of Physiology, School of Medicine, Aichi Medical University, 1-1 Yazako-karimata, Nagakute, Aichi, 480-1195, Japan
[5] Laboratory of Regulation in Metabolism and Behavior, Faculty of Agriculture, Kyushu University, 744 Motooka, Nishi-ku, Fukuoka, 819-0395, Japan
[6] School of Pharmacy, Queen's University Belfast, Belfast BT9 7BL, Northern Ireland, U.K.
[7] These authors jointly supervised this work: Osamu Nureki, Ralf Jockers
* equal contributions
✉ e-mail: aoishi@ks.kyorin-u.ac.jp; nureki@bs.s.u-tokyo.ac.jp; ralf.jockers@inserm.fr

Abstract

G protein-coupled receptors (GPCRs) transduce extracellular stimuli into intracellular signals by coupling to various heterotrimeric G proteins. However, the rules governing G protein preference remain largely elusive. MT_1 and MT_2 are prototypical $G_{i/o}$-coupled GPCRs responding to melatonin, a hormone secreted in a circadian manner. We show here that MT_1, but not MT_2, couples also to G_s proteins *in vitro* and activates the G_s/cAMP pathway upon long-term melatonin exposure *in vivo*, mimicking physiological dawn conditions. We solved the cryo–electron microscopy structure of the melatonin-MT_1-G_s complex at 3.0Å resolution, which revealed a strikingly distinct binding mode compared to the MT_1–G_i complex. The third intracellular loop of MT_1 emerges as a key stabilizer for G_s coupling, a feature previously unrecognized. This is the first solved receptor-G_s complex of a primary G_i-coupled GPCRs, providing new structural and functional insights into G protein selectivity and circadian switch of G protein coupling.

(Introduction)

G protein-coupled receptors (GPCRs) represent the largest family of membrane proteins involved in cellular signal transduction. GPCRs activate their primary transducers, heterotrimeric G proteins, that are classified in four families: G_s, $G_{i/o}$, $G_{q/11}$, and $G_{12/13}$. Among those, G_s and $G_{i/o}$ proteins have opposing roles as they are stimulating or inhibiting adenylyl cyclases (AC) and cyclic AMP production, respectively. Large-scale common G protein coupling datasets show that a particular GPCR can couple to one or several G protein types[1]. G_s-coupled GPCRs show the lowest coupling promiscuity and rarely co-couple to $G_{i/o}$ proteins or with a large difference in their strength of activation[1]. The few exceptions of G_s/$G_{i/o}$ co-coupling include the primarily G_s-coupled 5-HT4, EP4 and ß1AR[2-4] , as well as the primarily $G_{i/o}$-coupled GPR139[5].

MT_1 and MT_2 are high-affinity receptors for melatonin (5-methoxy-N-acetyl-tryptamine) that primarily couple to $G_{i/o}$ proteins, leading to inhibition of cAMP production and the PKA/CREB pathway, activation of ERK1/2, and regulation of ion channels[6-8]. The molecular structures of MT_1 and MT_2 receptors have been solved, all in the presence of synthetic agonists, with some in complex with the G_i protein[8-13]. The stimulation profile of melatonin receptors is unique, governed

by the circadian secretion pattern of melatonin in the pineal gland with peak levels during the night[14]. Due to this rhythmic pattern of melatonin secretion, melatonin receptors regulate and synchronize a variety of physiological functions including sleep/awake rhythm, seasonal reproduction, and retina physiology, with currently marketed drugs indicated for sleep and circadian disorders and major depression[15,16]. Whereas most effects of melatonin receptors appear to be transmitted by $G_{i/o}$ proteins, coupling to G_s-dependent pathways has been suggested for MT_1[17-19]. However, whether this is due to direct G_s coupling or to downstream crosstalk is unknown. The physiological relevance of dual G_i/G_s coupling, with apparently opposing signaling outcomes, also remains elusive and intriguing. Here, we provide functional evidence for G_s coupling to MT_1 under physiologically relevant conditions in the mouse hypophysial *pars tuberalis* (PT), and report the cryo-electron microscopy (cryo-EM) structure of the MT_1–G_s complex in the presence of the natural ligand melatonin. The unique interaction mode of G_s depends on the third intracellular loop of MT_1 (ICL3) which is required to stabilize the interaction with MT_1.

Results

Melatonin MT_1 receptor activates both the inhibitory G_i/cAMP and the stimulatory G_s/cAMP pathway

Acute stimulation with melatonin of HEK293 cells expressing low levels of human MT_1 receptor inhibited forskolin (FSK)-stimulated cAMP production with an pEC_{50} of 9.96±0.34 (n=6), as expected for this prototypical $G_{i/o}$-coupled receptor (Fig. 1a). In contrast, cells expressing high MT_1 levels showed a biphasic behavior with inhibition of cAMP production at low melatonin concentrations (pEC_{50}=10.3±0.35; n=6) and stimulation of cAMP production at higher melatonin concentrations (pEC_{50}=8.6 ±0.66; n=6) (Fig. 1a). Similar experiments in cells expressing high levels of the closely related $G_{i/o}$-coupled human MT_2 melatonin receptor showed only the inhibitory component (pEC_{50}=9.93±0.23; n=6), indicating that the biphasic behavior is a unique feature of MT_1.

Pretreatment of HEK293 cells expressing high levels of MT_1 and MT_2 with pertussis toxin (PTX), a $G_{i/o}$ inhibitor, abolished the inhibitory effect of melatonin for both receptors, while the stimulatory effect was preserved for MT_1 with a

pEC_{50} of 8.51±0.38; n=6) (Fig. 1b), which was similar to the value observed in the absence of PTX. This indicates that MT_1 and MT_2 couple to PTX-sensitive $G_{i/o}$ proteins and MT_1 to an additional PTX-insensitive G protein, possibly the stimulatory G_s protein. To test this hypothesis, we measured G_i and G_s coupling directly to MT_1 and MT_2 with the nanoluciferase (Nluc) complementation assay[20,21]. MT_1 and MT_2 were fused at their C-terminus to the NP fragment of Nluc ($MT_{1(2)}$-NP) and coexpressed with $miniG\alpha_s$ or $miniG\alpha_i$ proteins fused to the LgBiT fragment of Nluc (LgBit-$miniG\alpha_{s/i}$) to monitor G protein recruitment through Nluc complementation[16,17]. Melatonin-promoted recruitment was observed between LgBit-$miniG\alpha_i$ and both receptors (Fig. 1c,e) and between LgBit-$miniG\alpha_s$ and MT_1-NP, but not MT_2-NP (Fig. 1d,f), indicating that MT_1 couples to G_i and G_s while MT_2 couples only to G_i. Consistently, bi- vs monophasic melatonin concentration-response curves of FSK-stimulated cAMP production were also recapitulated with NP-tagged human MT_1 and MT_2, respectively (Supplementary Fig. S1a,b). Similar results were obtained for mouse MT_1 and MT_2 indicating that mice could be a relevant model for in vivo validation (Supplementary Fig. S1c,d).

MT_1 is abundantly expressed in the rodent pituitary PT whereas MT_2 is absent[22,23]. Acute melatonin treatment has been shown to inhibit FSK-stimulated cAMP production in cultured PT cells[24]. To simulate physiologically relevant long-term stimulation conditions, we first established ex-vivo cultures from mouse PT and stimulated them for 16h with melatonin (Fig. 1g). Under these conditions, melatonin promoted cAMP production (Fig. 1h) and further downstream CREB phosphorylation (Fig. 1i,j). Intraperitoneal administration of melatonin increased CREB phosphorylation in the mouse PT 16h after treatment (Fig. 1k-m). Collectively, these data indicate that endogenously expressed MT_1 couples to the G_s/cAMP pathway in vivo when stimulated for 16h mimicking the end of the dark phase under short-day conditions.

Structural insights into the MT_1-$miniG_s$ signaling complex

To obtain structural evidence for G_s coupling to MT_1, we determined the structure of the MT_1-$miniG_s$ complex by cryo-EM at 3.0 Å global resolution, together with the endogenous ligand melatonin, $G\beta_1$, $G\gamma_2$ subunits and the Nb35 fragment (Fig. 2a, b). The overall structure of MT_1 is similar to the MT_1 structure in the previously reported MT_1-Gi complex[12]. In the MT_1-$miniG_s$ complex, MT_1 is in an active state, supported by the structural similarity at the

conserved activation motifs within the MT$_1$-G$_i$ complex (Extended Data Fig. 3a-e). At the G$_s$ interface, the MT$_1$-miniG$_s$ displayed a novel binding mode, where Arg125$^{3.50}$, in the DRY motif of MT$_1$ made no interaction with Tyr391$^{G.H.5.23}$ of G, while this interaction is observed in almost all of the reported structures of class A GPCR-G$_s$ complex (Fig. 2c). Instead of the typical interaction between Arg$^{3.50}$ and Tyr391$^{G.H.5.23}$, Tyr128$^{3.53}$ at TM3 of MT$_1$ formed a hydrogen bond with Gln390$^{G.H.5.22}$ of G$_s$, and Ser132$^{34.50}$ at intracellular loop (ICL)2 of MT$_1$ formed a hydrogen bond with His387$^{G.H.5.19}$ of G$_s$. In parallel, Leu133$^{34.51}$ at ICL2 of MT$_1$ formed a hydrophobic interaction with the pocket of miniG$_s$ formed by H41$^{G.S1.02}$, F219$^{G.S3.03}$, F376$^{G.H5.08}$, R380$^{G.H5.12}$ and I383$^{G.H5.15}$, which is observed in most of the G$_s$ complexes (Fig. 2c).

The importance of these interacting residues was tested by the real-time cAMP assay with the Y128$^{3.53}$A, S132$^{34.50}$A, L133$^{34.51}$A MT$_1$ mutants (Fig. 2d-h). All mutated receptor variants showed similar expression levels (Supplementary Fig. S2).. Whereas the inhibitory G$_i$-dependent phase was maintained for all mutants, the stimulatory G$_s$-dependent phase was completely lost for Y128$^{3.53}$A, not affected for S132$^{34.50}$A and substantially reduced for L133$^{34.51}$A. This suggests the importance of Tyr128$^{3.53}$ and to a lesser extend Ser132$^{34.50}$ in cAMP production via MT$_1$-G$_s$ coupling, while maintaining MT$_1$-G$_i$ coupling.

Despite this unique binding modes at the MT$_1$-miniG$_s$ interface, miniG$_s$ in the MT$_1$-miniG$_s$ showed a similar conformation as the previously reported β$_2$AR-G$_s$ complex (PDB 3SN6) and was distinct from the GDP-bound inactive G$_s$ protein structure (PDB 6EG8) (Fig. 2i-k). Phe376$^{G.H5.08}$ moved upward to the C-terminal of the α5 helix in a typical way, causing the extension of α5 helix together with the movement of the TCAT motif at the nucleotide binding site (Fig. 2i-k). From these observations, we concluded that the MT$_1$-miniG$_s$ complex is a canonical signaling complex rather than an artifact.

Structural comparison with the previous structure of MT$_1$ in complex with its primary signal transducer, the G$_i$ trimer (PDB 7DB6), revealed that MT$_1$ exhibits a distinct binding mode when bound to miniG$_s$ (Fig. 2l). While the overall structure of MT$_1$ remains similar between complexes, the G protein α5 helixes adopt dramatically different entry angles: G$_s$ α5 inserts from the ICL2 side, whereas G$_i$ α5 penetrates from the ICL3 side (Fig. 2l). This differential positioning enables unique interactions in each complex. In the MT$_1$-G$_s$ structure, Leu133$^{34.51}$ forms hydrophobic contacts with the Gα pocket - an interaction absent in the MT$_1$-G$_i$ complex (Fig. 2m,n). This observation aligns

with preserved cAMP inhibition in L133A and Y128A mutants. Conversely, G_i residues at equivalent positions do not form similar interactions between Cys351$^{G.H.5.23}$ of G_i and Arg125$^{3.50}$ of MT$_1$ (Fig. 2n). Critically, G_s α5 adopts shallow binding with its C-terminal tip oriented toward ICL2/TM3, while G_i α5 penetrates more deeply toward TM5/TM6 regions[12]. We conclude from these observations that the binding mode of the MT$_1$-G$_s$ complex is distinctive from that of the MT$_1$-G$_i$ complex.

MT$_1$ shares a similar G$_s$ entry angle with GPR61 but shows a different binding mode from GPR61

To identify other GPCRs coupling to G$_s$ in a similar way to MT$_1$, we performed an amino acid sequence alignment of TM3 and ICL2 of class A GPCRs using the GPCRdb server [25]. Surprisingly, only 5 class A GPCRs contain a Tyr at position 3.53, including MT$_1$ and MT$_2$ (Fig. 3a). Among the 5 GPCRs, GPR61 is the only receptor for which the structure of G$_s$ complex is available in PDB. Structural comparison with the recent cryo-EM structure of the GPR61-G$_s$ complex (PDB 8KGK) revealed that MT$_1$ and GPR61 share a similar G$_s$ entry angle (Fig. 3b, d). However, G$_s$ enters deeper into GPR61 than into MT$_1$ both at ICL2 and the cytoplasmic pocket of the receptor (Fig. 3c, e). At the ICL2 interface, residues around Tyr148$^{34.51}$ of GPR61 make closer contacts than MT$_1$, and Glu151$^{3.54}$ of GPR61 interacts with G$_s$ via Gln35$^{G.HN.52}$, which i not observed in the MT$_1$-minG$_s$ (structure Fig. 3c). At the cytoplasmic pocket, the backbone of Tyr143$^{3.53}$ of GPR61 forms a hydrogen bond with His387$^{G.H.5.19}$ of G$_s$, while Tyr128$^{3.53}$ of MT$_1$ forms a hydrogen bond with Gln390$^{G.H.5.22}$ of G$_s$ (Fig. 3e). Associated with this conformational difference, Pro147$^{34.50}$ of GPR61 does not interact with G$_s$, which is at the same position as Ser132$^{34.50}$ of MT$_1$ interacting with His387$^{G.H.5.19}$ of G$_s$. As a result, G$_s$ enters deeper into GPR61 than MT$_1$, associated with additional interactions; Gln347$^{8.49}$ to Gln390$^{G.H.5.22}$, and Arg346$^{8.48}$ and Arg348$^{8.50}$ to Glu392$^{G.H.5.24}$. These structural insights may reflect the constitutive G$_s$ signaling activity of GPR61 and the fact that G$_s$ corresponds only to the secondary coupling of MT$_1$, which is primarly coulpled to G$_i$.

Binding features among G_s- and G_i- signaling complexes

To understand how GPCRs couple to different G protein subtypes, we compared structures of five receptors with both G_s-bound and G_i-bound complexes: 5HT₄ [2] (G_i: PDB 7XTA, G_s: PDB 7XT9), FFAR4 (G_i: PDB 8ID9, G_s: PDB 8H4I), β₁AR [3] (G_i: PDB 7S0F, G_s: PDB 7S0G), CCKAR [26] (Gi: PDB 7EZH, G_s: PDB 7EZK) and EP4 [4] (G_i: PDB 8GCP, G_s: PDB 8GDA). 5HT4, β₁AR and EP4 primarily couple to G_s, while CCKAR and FFAR4 primarily couple to G_q. MT₁ is unique as the first primarily G_ii-coupled GPCR also complexed with G_s.

5HT4-G_s/-G_i and FFAR4-G_s/-G_i displayed significant differences in the N-terminal region of α5 helix while maintaining similar C-terminal conformations, a pattern closely resembling the structural variations observed in MT₁-G_s/-G_i complexes. In contrast, EP4-G_s/-Gi and CCKAR-G_s/-Gi complexes exhibited more pronounced conformational differences at the C-terminal "wavy hook" region of α5. In β₁AR complexes, G_s and G_i showed the closest overlay of α5, likely due to the engineered Gi protein containing N-terminal G_s substitutions.

These structural comparisons reveal receptor-dependent mechanisms for dual G protein coupling with no universal structural determinant emerging from current structures, highlighting the complexity of GPCR-G protein selectivity.

Structural basis of MT₁ subtype specific G_s-coupling is in the TM5-ICL3-TM6 region

Tyr128$^{3.53}$ is a key feature of MT₁ for G_s coupling, yet Tyr$^{3.53}$ exists also in MT₂, which does not couple to G_s, suggesting that features other than Tyr$^{3.53}$ explain the exclusive MT₁ coupling to G_s and not MT₂. Recent structural studies demonstrated that the distance from the latter half of TM5 to TM6 is the key for G_s and G_i selectivity of serotonin receptor subtypes, as swapping of the TM5-ICL3-TM6 domain between the G_s-coupled 5HT₄ and the G_i-coupled 5HT$_{1A}$ switched also their G protein coupling profiles[2]. Inspired by this study, we exchanged the TM5-ICL3-TM6 regions between MT₁ and MT₂ (Fig. 4a,b). The **MT₁**-MT₂(TM5-6) chimera lost the melatonin-induced cAMP elevation phase seen in wild-type MT₁, whereas the **MT₂**-MT₁(TM5-6) chimera gained the cAMP elevation phase of the wild-type MT₁ (Fig. 4c). The **MT₂**-MT₁(TM5-6) chimera coupled to both miniGα$_i$ and miniGα$_s$ proteins (Fig. 4d). These data indicate that the molecular determinants of G_s coupling are located in the TM5-ICL3-TM6 region of MT₁ and are absent in MT₂.

To better understand the binding mode of G_s to the **MT₂**-MT₁(TM5-6) chimera,

we determined its cryo-EM structure in complex with miniG$_s$^{star} [27] (Fig. 4e). The overall structure of the **MT$_2$**-MT$_1$(TM5-6)-G$_s$ complex was similar to the MT$_1$-G$_s$ complex (Fig. 4f), with the exception of the location of the α5 helix of the chimera complex which was slightly different from that of the MT$_1$-G$_s$ (Fig. 4f). Associated with this conformational change, Arg138$^{3.50}$ formed a stacking interaction with Tyr391$^{G.H.5.23}$ of G$_s$, though Tyr141$^{3.53}$ and Ser145$^{34.50}$ kept interactions with Gln390$^{G.H.5.22}$ and His387$^{G.H.5.19}$ of G$_s$, respectively (Fig. 4h,i). In parallel, Met146$^{34.51}$ at the same position of Leu133$^{34.51}$ maintained a hydrophobic interaction with the pocket of G$_s$ (Fig. 4j). To understand why the substitution of TM5-ICL3-TM6 enabled MT$_2$ to couple to G$_s$, we compared the **MT$_2$**-MT$_1$(TM5-6)-G$_s$ structure with the previously solved structure of MT$_2$ in complex with G$_i$ (Fig. 4g, k). Similar to MT$_1$, different entry angles were observed between G$_s$ and G$_i$. No other substantial changes were found. The position of TM5 and TM6 was similar between MT$_2$ and the **MT$_2$**-MT$_1$(TM5-6) chimera structures indicating that the positioning and the rotation of TM5 and TM6 is not responsible for G$_s$ coupling of the chimera (Fig. 4k). Collectively, these data suggest that TM5 and TM6 do not play an important role in G$_s$ coupling of MT$_1$ vs MT$_2$, and suggest that key molecular determinants are likely located within ICL3.

ICL3 of MT$_1$ supports coupling to G$_s$ but not to G$_i$

To provide experimental evidence for the importance of ICL3 over TM5 and TM6 of MT$_1$ for G$_s$ coupling, we generated MT$_2$-MT$_1$ chimera containing TM5, ICL3 or TM6 of MT$_1$ alone or a combination of ICL3 and TM6 (Fig. 5a-e, Supplementary Fig.S3). All chimeras were expressed at comparable levels as the MT$_2$-MT$_1$(TM5c-ICL3-TM6) reference chimera (Supplementary Fig. S4). All chimeras showed the melatonin-induced cAMP inhibition phase (Fig. 5b-e, Supplementary Fig.S3). The MT$_2$-MT$_1$(ICL3) chimera displayed, in addition, the cAMP elevation phase (Fig. 5d). Among the 20 amino acids of the ICL3 region, the MT$_2$-MT$_1$(coreICL3) chimera containing the N-terminal 12 amino acids (VRQRVKPDRKPK) of MT$_1$ maintained the G$_s$ phase (Fig. 5e), whereas the MT$_2$-MT$_1$(C-ICL3+TM6) chimera containing the 8 C-terminal amino acids did not contribute to the G$_s$ phase (Supplementary Fig. S3b). Further delimitation of the 12 amino acid stretch failed to induce the G$_s$ phase in the MT$_2$ context (Supplementary Fig. S3c,d), defining the 12 N-terminal amino acid of the MT$_1$-ICL3 as the minimal region for the G$_s$ phase of cAMP signaling. Coupling of the

minimal MT$_2$-MT$_1$(coreICL3) chimera to both G$_s$ and G$_i$ was confirmed directly in the miniG$_{s/i}$ coupling assays (Fig. 5f,g).

Since the flexible ICL3 could not be resolved in our MT$_1$-G$_s$ structure, we performed molecular dynamics (MD) simulations of MT$_1$-G$_s$, MT$_1$-G$_i$, MT$_2$-G$_i$, and MT$_2$-MT$_1$(TM5-ICL3-TM6)-G$_s$ complexes using AlphaFold3-modeled ICL3 structures. A MT$_2$-G$_s$ test model was generated to understand why wild-type MT$_2$ cannot stabilize G$_s$ interactions. Unlike many GPCRs with extensive ICL3 regions (>100 residues), the melatonin receptors possess short, 20-residue, loops with high AlphaFold3 confidence (>60%). MD simulations revealed that the ICL3 forms dynamic polar and hydrophobic interactions with G protein α5 helices, with notable hydrophobic contacts between MT$_1$ V217^{ICL3} and L388$^{G.H5.20}$ of G$_s$ (Fig. 6a,b, Extended Data Fig. 6a). Together with I129$^{3.54}$, V214$^{5.65}$, and L213$^{5.64}$, these residues create a hydrophobic patch that stabilizes the receptor-G$_s$ interface. While V217^{ICL3} also interacts with G$_i$, these contacts are less essential because Gi α5 penetrates deeper into TM5/TM6 regions, achieving stability through intrinsic hydrophobic interactions involving L348$^{G.H5.20}$, F354$^{GH5.26}$, and L353$^{GH5.25}$ with I210$^{5.61}$, whose alanine mutation substantially reduces G$_i$ coupling [12]. The predicted importance of V217^{ICL3} for G$_i$ and G$_s$ coupling was tested in the context of the MT$_2$-MT$_1$(coreICL3) chimera representing the minimal MT$_1$ sequence enabling notable MT$_2$-G$_s$ coupling, by the V217A^{ICL3} mutant replacing Val 217 by Ala present in MT$_2$ (Fig. 6c). While the inhibitory phase was preserved, the stimulatory phase was abolished consistent with the important role of V217^{ICL3} for G$_s$, but not G$_i$, coupling (Fig. 6c). Deletion of ICL3 (ΔQRVKPDRKPK) in MT$_1$ resulted in the loss of the melatonin-induced cAMP stimulation phase while maintaining the inhibition phase further confirming the essential role of ICL3 for G$_s$ coupling but not for G$_i$ coupling to MT$_1$ (Fig. 6d). This ICL3 deletion chimera also shows that V217^{ICL3} alone is not sufficient for G$_s$ coupling as this residue as well as the other residues involved in the hydrophobic patch (I129$^{3.54}$, L213$^{5.64}$, V214$^{5.65}$) are present in this chimera (Fig. 6b). This is consistent with the observation that chimeric replacement of TM5 alone does not confer G$_s$ coupling to MT$_2$ (see Fig. 5b). Our MT$_2$-G$_s$ test model MD simulations support this mechanism, showing that equivalent alanine residues in MT$_2$ ICL3 cannot provide adequate stabilization of G$_s$, resulting in high ICL3 and α5 mobility with substantially reduced interaction energies (Extended Data Fig. 6b-d). Since VRQRVKPDRKPK represents the minimal sequence enabling notable MT$_2$-G$_s$

coupling, specific ICL3 conformation appears also essential for establishing hydrophobic interactions necessary for G_s recognition. Our findings demonstrate that ICL3 is essential for G_s coupling but dispensable for G_i coupling, in a context where G_s coupling to MT_1 relies heavily on ICL3 interactions, which likely compensates for its shallow binding mode.

Discussion/conclusion

We describe here the molecular determinants enabling G_s coupling to MT_1, a prototypical G_i-coupled GPCR. The binding mode of the MT_1-G_s complex is structurally distinctive from the MT_1-G_i coupling: while G_i binding involves the C-terminal end of TM5 and TM6 of MT_1[12,13], G_s forms essential interactions with TM3 (Y128[3.53]) and ICL2 (L133[34.51]) of MT1, which were experimentally validated. Since the flexible ICL3 could not be resolved in our MT_1-G_s structure, we investigated its role through MT_2-MT_1 chimera studies and MD simulations. Chimera studies defined the N-terminal 12 amino acids (VRQRVKPDRKPK) of MT_1-ICL3 as minimal region required for G_s coupling with V217[ICL3] establishing hydrophobic contacts participating in G_s recognition, stabilizing the otherwise shallow binding mode of G_s in terms of depth of penetration into the TM5/TM6 region. Consistently with the absence of G_s coupling to MT_2, the stabilizing role of ICL3 is not seen for this receptor in MD simulations. While the ICL3 is essential for G_s coupling to MT_1, it is dispensable for G_i coupling as demonstrated with the MT_1-ICL3 deletion mutant.

The contribution of ICL3 in G protein coupling has been suggested previously by functional and bioinformatics studies[28-30]. On the other hand, for different serotonin receptor subtypes, the length of TM5 and TM6 has been suggested as important determinant of G_s-G_i selectivity[2]. For MT_1, the contribution of TM5 and TM6 could be excluded by solving the cryo-EM structure of the **MT_2**-**MT_1**(TM5-6) chimera which couples to G_s, but shows no significant differences in TM5 and TM6 length, position and rotation compared to the MT_2, which does not couple to G_s. The study of Sadler et al. [30] suggested a gating function of ICL3 controlling G access to receptors. Our data suggest different roles of ICL3 for G protein coupling to MT1. Whereas the MT_1-G_s complex suggests rather a supportive role of ICL3 for G_s coupling, ICL3 is dispensable for G_i coupling to

MT_1.

Coupling of MT_1 to G_i occurs upon acute melatonin stimulation. Coupling to G_s is also of physiological relevance as it occurs upon long-term melatonin stimulation, mimicking the conditions at dawn, when receptors are exposed to melatonin throughout the night. Stimulation of the G_s/cAMP/PKA pathway has been observed in our study *ex vivo* and *in vivo* in the mouse PT expressing endogenous MT_1. Previously, melatonin has been shown to have a dual impact on the rhythmic transcription of PT cells [31]. In the beginning of the night melatonin acutely suppresses cAMP levels which is consistent with the well-documented G_i-coupling of MT_1 receptor. At dawn, melatonin has a delayed heterologous sensitizing effect on the adenylyl cyclase system[32] which was shown by the sensitization of the signaling of adenosine A_{2b} receptor, the most prominently expressed G_s-coupled receptor in the PT[33]. Our study indicates that apart from this heterologous sensitization by amplifying the A_{2b} receptor response, the G_i-G_s switch of MT_1 coupling is likely to contribute to the increased activity of the cAMP/PKA pathway at the end of the night, thus enhancing the contrast of cAMP-dependent rhythmic transcription in PT cells at the night/day transition. These results provide new insights on the molecular mechanisms underlying the synchronization effects of melatonin, where the acute $G_{i/o}$ pathway upon melatonin rise would signal the beginning of the dark phase of the light/dark cycle, and the switch to G_s-coupling would signal the night/day transition.

Materials and Methods

Material

HEK293T cells were used as described previously[34]. The expression constructs of melatonin receptors used in this study were as follows. For Fig. 1a,b, C-terminal untagged Flag-MT_1 and HA-MT_2 constructs were used as described previously[34]. For other experiments, C-terminal–tagged constructs (Flag-MT_1-NP and Flag-MT_2-NP) were generated by PCR-based modification using plasmids obtained from Addgene (Plasmid #66443[35] and #66444[35]). Melatonin receptors (MTR) mutants were generated by PCR using PNK-phosphorylated primers carrying the intended mutation, followed by self-ligation with T4 DNA ligase (Nippon Gene), as described previously[36]. Chimeric MTRs were

generated by PCR using the In-Fusion HD cloning kit (Clontech) as described previously[37]. cDNA encoding LgBit-miniGs protein or –miniGs/i chimera protein was provided by Dr. Nevin Lambert (Augusta, Georgia, USA).

Cell Culture and Transfection

HEK293T cells were transiently transfected with Jet-PEI or PEI-Max according to the manufacturer's protocol[36,37]. Briefly, approximately 0.3×10^6 cells per well were seeded into 12-well plates. After 6 h, a total of 1 µg DNA (expression plasmid supplemented with empty vector) was mixed with 3 µL of transfection reagent, incubated for 15 min, and then added to the cells. On the following day, cells were transferred to white 96-well plates, and experiments were performed 48 h post-transfection.

miniG protein Nanoluc Complementation assay

Nanobit complementation assay was performed as following: HEK293T cells seeded in 12 well plate were co-transfected with LgBiT-fused miniGs (5ng/well) or miniGs/i (15ng/well) together with 5ng/well NP-fragment–tagged[21] human Flag-MT$_1$ or Flag-MT$_2$ receptors or their chimera proteins. Cells were reseeded into white 96-well plates on the following day. After 46 h, the culture medium was replaced with phenol red–free DMEM. Two hours later, 5 µM coelenterazine H was added, and baseline luminescence was recorded. Cells were then stimulated with vehicle or 1 µM melatonin, and luciferase activity was measured. Data represent luminescence values obtained 2 min after melatonin stimulation.

In Vitro cAMP Assay by HTRF

cAMP accumulation was assessed by HTRF (Fig.1a,b) under suspension conditions as described previously[34]. HEK293T cells were transfected with varying amounts of Flag-MT$_1$ without any C-terminal tag (1 µg, "high"; or 10 ng plus 990 ng empty vector, "low") or 1 µg HA-MT$_2$ without any C-terminal tag plasmids in 12-well plates. After 48 h, cells were dispensed into 384-well plates (4,000 cells per well) and stimulated with 2 µM forskolin in the presence of the indicated melatonin concentrations for 30 min at room temperature in PBS supplemented with 1 mM 3-isobutyl-1-methylxanthine (IBMX; Sigma-Aldrich, France). Cells were then lysed, and cAMP levels were quantified following the manufacturer's instructions (Cisbio Bioassays, France). Where indicated, cells

were pretreated with PTX (200 ng/mL, 4 h). Luminescence was recorded using an Infinite F500 microplate reader (Tecan).

cAMP monitoring by CAMYEL-BRET sensor

cAMP measurement by CAMYEL-BRET sensor was performed as previously described[38]. HEK293T cells were co-transfected with 500 ng CAMYEL sensor and 500 ng melatonin receptor (Flag-MT$_1$-NP or Flag-MT$_2$-NP and their mutants or chimera receptor) plasmids in 12-well plates. The following day, cells were reseeded into 96-well plates. After 46 h, the culture medium was replaced with phenol red–free DMEM. Two hours later, 5 µM coelenterazine H was added, and baseline BRET signals were recorded. Cells were then treated with vehicle or 5 µM forskolin in the presence of the indicated melatonin concentrations (without IBMX), and real-time BRET changes were monitored. Data shown correspond to measurements taken 10 min after melatonin stimulation. BRET was measured by TECAN SPARK.

Animals

Five-week-old male C57BL/6JJmsSlc mice (Japan SLC Inc., Shizuoka, Japan) were purchased and housed in plastic cages (170 W × 240 D × 125 H mm, Clea, Tokyo, Japan) under a 12-h light (200 lx of fluorescent light) / 12-h dark cycles (12L12D, 0800 light ON, 2000 light OFF), maintained at a constant temperature (23 ± 1 °C). Food (CE-2; CLEA) and water were provided *ad libitum*. All animal experiments were approved by the Committee of Animal Care and Use of the Aichi Medical University. All experimental procedures were conducted in accordance with the institutional guidelines for the use of experimental animals.

Ex- and *In-Vivo* Melatonin treatment

For analysis of melatonin effect on cultured PT/MBH, brain of 10–12-week-old mice were quickly removed under general anesthesia at 6 hours after dawn and immersed in ice-cold Hank's buffered saline (HBS; #09735-75, Nacalai tesque, Kyoto, Japan). The hypothalamus, containing the PT and MBH was sectioned at 1 mm thickness by brain matrix. Then, the PT/MBHs were placed on a culture membrane (Millicell-CM PICM0RG50, Merck Millipore, Darmstadt, Germany) in a 35-mm petri dish and cultured at 37°C with 1.2 mL of medium containing Dulbecco's modified Eagle medium (DMEM) (Gibco, Carlsbad, CA), 100 U/mL

penicillin, and 100 µg/mL streptomycin. After 6 hours preincubation, slices were then added with the culture medium containing melatonin at ZT12 (finally 10 µM, 0.1 % DMSO; Wako, Tokyo, Japan). After 16 hours, proteins of the PT/MBH were extracted for using cAMP measurement and western blotting.

For *in vivo* analysis of melatonin effect, mice were kept under 8-h light / 16-h dark cycles (8L16D) for 3 weeks, and melatonin (0.26 mM / 0.1mL) were intraperitoneally (i.p.) injected as previous report 1, once at ZT8. After 16 hours (ZT0), mice were perfused then fixed on slides with 4% paraformaldehyde (PFA) in 0.1 M phosphate buffer (pH 7.4), and brains were removed and put into the 4% PFA for 48 hours, and then transfer to 20 % sucrose / PBS for using frozen sectioning of brain.

Ex vivo cAMP measurement at PT

Protein extraction of cultured PT/MBH was performed using cell lysis buffer (#9803, Cell Signaling Technology, Tokyo, Japan) containing protease inhibitor cocktail (#P8340, Sigma) and phosphatase inhibitor cocktail 1 (#P2850, Sigma) according to the manufacturer's instructions.

cAMP measurements were performed with a homogeneous TR-FRET immunoassay using the LANCE cAMP Detection Kit (#AD0262, PerkinElmer, USA), according to the manufacturer's instructions (PerkinElmer). Cultured PT/MBH tissues homogenates (0.2 µg/µL) were diluted 10 times with stimulation buffer, and added to 10 µL of Alexa Fluor 647 anti-cAMP antibody diluted with stimulation buffer. After incubation for 45 min at room temperature, the reaction was stopped by the addition of 20 µL working solution (10 µL Eu-cAMP and 10 µL ULight-anti-cAMP), and incubated for 1 h at room temperature. The TR-FRET signal was read using a microplate reader SpectraMax M5 (Molecular Devices). cAMP concentrations were determined using GraphPad Prism 8 software (GraphPad Software Inc., San Diego, CA).

Western blot analysis

Western blot analysis was performed as follows: homogenized PT/MBH was loaded as 2 µg protein samples per lane onto 8 % SDS-polyacrylamide gels. Following separation at 100 mA, membranes were incubated with the following primary antibodies: rabbit monoclonal antibodies against phospho-CREB (Ser133) (1:1000; #9198, Cell Signaling Technology), and CREB (1:1000; #9192, Cell Signaling Technology). Membranes were washed and then

incubated with HRP-conjugated goat polyclonal antibody against rabbit IgG (1:10,000; #7074, Cell Signaling Technology). By using ImmunoStar LD (#292-69903, Fijifilm), chemiluminescent images were detected using an Amersham Imager 600 (Cytiva Lifescience).

Immunohistochemistry

Immunohistochemistry was performed by using the Vectastain Elite ABC rabbit IgG kit (Vector, Burlingame, CA) as our previous reports[39]. Frozen sections of paraformaldehyde-fixed trimmed mice brain (14 μm) including the PT/MBH were used. Briefly, after drying, the sections were immersed in HistoVT One (pH7.0, #06380, Nakarai) and heated in a microwave oven for 15 min at low voltage. After blocking with normal rabbit serum (Vectastain), sections were incubated with rabbit polyclonal antibody against phospho-CREB (Ser133) (1:500; #9198, Cell Signaling Technology) at 4°C for 48 h. After encapsulation with mounting medium, images were detected using a BZ-X800 (Keyence). ImageJ software (NIH) was used to analyze pCREB-immunoreactive (ir) cell numbers as previous report[40]. Data are shown as percent of maximum values (pCREB-ir cells /mm2).

Expression and purification of Gβ$_1$-Gγ$_2$ dimer

In the previous study, we expressed and purified the Gi1 heterotrimer including human Gαi1, bovine Gβ$_1$ and mouse Gγ$_2$ by Ni-NTA affinity chromatography and anion exchange chromatography. As a byproduct of the anion exchange chromatography step, the fraction of Gβ$_1$-Gγ$_2$ dimer was eluted before the fraction of the Gi$_1$ heterotrimer.

Expression and purification of Nb35

The plasmid encoding Nb35 was prepared as previously reported[41]. The protein was expressed in the periplasm of E. coli C41 (Rosetta) cells cultured in LB medium supplemented with 1 mM IPTG for 20 h at 25 °C. After 20 h, the cells were collected and disrupted by ultrasonication in hypotonic buffer (20 mM Tris-HCl, pH 7.5, 150 mM NaCl and 2 mM MgCl2), and the Nb35 protein was purified by Ni-NTA affinity chromatography and then subjected to size-exclusion chromatography on a HiLoad Superdex75 16/600 column. Peak fractions were pooled and concentrated to 3 mg ml^{-1}.

Expression and purification of MT$_1$-miniG$_s$-Gβ$_1$-Gγ$_2$ complex

The HA signal sequence, the FLAG epitope tag (DYKDDDDK), the ALFA tag (SRLEEELRRRLTE)[42] and GSGSG linker were fused to the N-terminus of MT$_1$, in this order. In addition, we fused the miniGs fragment to the C-terminus of MT$_1$.

In the MT$_2$-MT$_1$ chimera construction, we substituted TM5-ICL3-TM6 of MT$_2$ to the MT$_1$ sequence (from Met200$^{5.51}$ to Phe256$^{6.53}$) between Pro212$^{5.50}$ and Ile270$^{6.54}$. And we fused the same fragments as MT$_1$-G$_s$ to the N-terminus of MT$_1$. In contrast, we fused mini-Gs^{star} protein created in our previous study[41] to the C-terminus of MT$_1$. All the above constructions were cloned into the pEGBacmam vector.

Then we infected a one tenth volume of a solution containing the virus encoding the above construction to the HEK293F cells at a density of 3–4 × 10^6 cells ml^{-1}, and grown at 37 °C for 20 h in FreeStyle 293 Expression Medium (Gibco). After sodium butyrate (FujiFilm Wako Pure Chemical Corporation) was added to 10 mM, the cells were incubated further at 30 °C for 48 h.

The HEK293F cells were collected by centrifugation at 5,000g for 10 min, lysed in buffer containing 50 mM Tris (pH 8.0), 150 mM NaCl and 10% glycerol and fresh frozen by liquid nitrogen. The cells were thawed and directly solubilized at 4 °C for 2 h in the solubilization buffer, containing 50 mM Tris (pH 8.0), 150 mM NaCl, 10% glycerol,1.5% DDM, 0.15% CHS, 5.2 μg ml^{-1} aprotinin, 2.0 μg ml^{-1} leupeptin, 1.4 μg ml^{-1} pepstatin A, 100 μM PMSF and 10 μM melatonin (Wako). The soluble fraction containing MT$_1$-miniG$_s$ was separated by ultracentrifugation (186,000g for 30 min), and the supernatant was mixed with the purified Gβ$_1$-Gγ$_2$ dimer (0.8 mg l−1 HEK cells), the purified Nb35 (0.5 mg l−1 HEK cells) and final 50 mU/mL of Apyrase (New England Biolabs). Then they were stirred at 4 °C for overnight. The solution was supplemented with 2 mM CaCl$_2$ at the final concentration, mixed with the M1 anti-FLAG resin and rotated at 4 °C for 2 h.

The resin was collected by centrifugation (500g for 3 min) and washed with 20 column volumes of buffer, containing 50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 10% glycerol, 0.03% LMNG, 0.003% CHS, 2 mM CaCl$_2$ and 1 μM melatonin. The complex was eluted with buffer, containing 50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 10% glycerol, 0.03% LMNG, 0.003% CHS, 2 mM CaCl$_2$, 1 μM melatonin, 5 mM EDTA (pH 8.0) and 125 μg ml^{-1} FLAG peptide (Synthesized by GenScript), concentrated and purified by size-exclusion chromatography on a Superdex200 10/300 Increase (GE) column, using buffer containing 50 mM

HEPES-NaOH (pH 7.5), 150 mM NaCl, 100 μM TCEP, 0.01% LMNG, 0.001% CHS and 1 μM melatonin. Peak fractions were pooled and concentrated to around 7 mg ml^{-1} with a centrifugal filter device (Millipore 50 kDa molecular weight cutoff).

Grid preparation and cryo-EM data collection

The purified complex solution was applied to freshly glow-discharged Au 300 mesh R1.2/1.3 grids (Quantifoil), using a Vitrobot Mark IV (FEI) at 4 °C, with a blotting time of 4 s under 100% humidity conditions. The grids were then plunge-frozen in liquid ethane cooled to the temperature of liquid nitrogen.

Cryo-EM data were collected using a Titan Krios G3i microscope (Thermo Fisher Scientific), running at 300 kV and equipped with a Gatan Quantum-LS Energy Filter (GIF) and a Gatan K3 Summit direct electron detector in the electron counting mode (The University of Tokyo, Japan). Movies were recorded at a nominal magnification of 105,000×, corresponding to a calibrated pixel size of 0.83Å, with a total dose of approximately 50 electrons per Å^2 per 48 frames. The data were automatically acquired using the EPU software (Thermo Fisher Scientific), with a defocus range of −0.8 to −1.6 μm. For MT1-miniGs, 10,373 movies were obtained. For MT1-MT2-miniGs, 8,174 movies were obtained.

Image processing

All acquired movies were dose-fractionated and subjected to beam-induced motion correction implemented in RELION 3.1[43]. The contrast transfer function (CTF) parameters were estimated using patch CTF estimation in cryoSPARC v3.3[44].

For MT$_1$-miniG$_s$, particles were initially picked from a small fraction with the Blob picker and subjected to several rounds of two-dimensional (2D) classification in cryoSPARC using 897 movies. Selected particles were used for template picking on the full dataset, and 8,196,012 particles were picked and extracted with a pixel size of 3.32 Å, followed by 2D classification, ab initio reconstruction, homogeneous refinement and heterogeneous refinement, displayed in the Extended Data Fig. 1. A total of 624,555 particles were used for training of topaz model[45], and 3,524,939 particles were extracted. After 2D classification, heterogenous refinement and non uniform (NU) refinement, 449,204 particles were transferred into RELION 3.1 and were processed by 3D classification without alignment using TMD mask with several T parameters. After 3D

refinment with SIDESPLITTER[46], particle polishing, re-extract as 1.0 Å/pix and removal of the duplicated particles, 85,855 particles were transferred into cryoSPARC.

Finally, these 85,855 particles were reconstructed using NU refinement, resulting in a 3.03 Å overall resolution reconstruction, with the gold standard Fourier Shell Correlation (FSC = 0.143) criteria in cryoSPARC. Moreover, the 3D model was refined with a mask on the receptor by local refinement. As a result, the local resolution of the receptor portion was estimated as 3.24 Å by cryoSPARC. In parallel, the 3D model was refined with a mask on the G protein by local refinement, resulting in the local resolution at 2.9 Å estimated by cryoSPARC.

For MT_2-MT_1-$miniG_s$, particles were initially picked from a small fraction with the Blob picker and subjected to several rounds of 2D classification in cryoSPARC using 2,429 movies. Selected particles were used for template picking on the full dataset, and 54,229,298 particles were picked and extracted with a pixel size of 3.32 Å, followed by 2D classification, ab initio reconstruction, homogeneous refinement and heterogeneous refinement, displayed in the Extended Data Fig. 2. A total of 272,585 particles were selected as the best class and were transferred into RELION 4.0. After 3D classification without alignment using TMD mask with several T parameters, 74,839 particles were selected, processed by 3D refinement with SIDESPLITTER[46], particle polishing and re-extract as 1.0375 Å/pix and transferred into cryoSPARC.

Finally, these 74,839 particles were reconstructed using NU refinement, resulting in a 2.94 Å overall resolution reconstruction, with the gold standard Fourier Shell Correlation (FSC = 0.143) criteria in cryoSPARC. Moreover, the 3D model was refined with a mask on the receptor by local refinement. As a result, the local resolution of the receptor portion was estimated as 3.43 Å by cryoSPARC. In parallel, the 3D model was refined with a mask on the G protein by local refinement, resulting in the local resolution at 2.82 Å estimated by cryoSPARC.

Model building and validation

The cryo-EM structures of the MT_1-Gi complex (PDB 7DB6)[12], MT_2-Gi complex (PDB 7VH0)[13] and OR51E2-miniGs (PDB 8F76)[47] were used as starting templates for modeling the MT_1, MT_2-MT_1, miniGs, Gβγ and Nb35 components. Initially, these models were positioned into the density map using jiggle fit in

COOT[48]. Then these models were modified by COOT and refined with phenix.real_space_refine (v1.19)[49,50] and the secondary structure restraints from phenix.secondary_structure_restraints. Cif format definition (ML1) downloaded from PDB was used for modeling melatonin molecule at ligand binding site.

Computer Simulations

MD simulations were conducted on melatonin receptor systems derived from cryo-EM structures. All simulated systems and their stability parameters are listed in **(Supplementary Table 1)**. MD simulations were performed on five distinct melatonin receptor-G protein complexes: MT_1+Gi (PDB: 7DB6), MT_1+Gs, MT_2+Gi (PDB: 7VH0), MT_2_MT_1(TM5-ICL3-TM6)+Gs, and MT_2_WT+Gs. The cryo-EM structures were prepared for simulations using Schrödinger Maestro (2025-1)[51]. ICL3 regions were rebuilt by grafting the corresponding segments from GPCRdb active state AlphaFold2-Multistate models [52,53] (version 15-05-2024): MT_1 AlphaFold ICL3 (V217-P231) with an average pLDDT confidence score of 65.17, and MT_2 AlphaFold ICL3 (A230-P244) with an average pLDDT confidence score of 73.62. The grafting boundaries were determined by identifying structured TM5/TM6 regions flanking the missing ICL3 density in the cryo-EM structures. AlphaFold segments were integrated as follows: MT_1+Gi [V198-P244], MT_1+Gs [V198-N236], MT_2+Gi [L211-L251], MT_2_MT_1(TM5-ICL3-TM6)+Gs [V198-P254]. The experimentally determined chimeric structure was used as the template for both MT_2_MT_1(TM5-ICL3-TM6)+Gs and MT_2_WT+Gs systems, with the MT_2_WT+Gs variant generated by mutating the MT_1-derived TM5-ICL3-TM6 segment back to MT_2 wild-type sequence using Schrödinger Maestro 3D builder utility. In all systems, D55$^{2.50}$ was protonated as per PROPKA3 predictions[54,55].

CHARMM-GUI[56-64] server was used to prepare simulation systems. Membrane protein orientations were determined using the Positioning of Proteins in Membranes (PPM) 2.0[56] web server prior to embedding in a 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC) bilayer membrane. Membrane dimensions were optimized for each receptor-G protein complex, ranging from 130 Å × 130 Å to 140 Å × 140 Å, to maintain adequate lipid-protein boundaries and prevent periodic artifacts. Each system was solvated with TIP3P water extending 30-35 Å from the membrane surfaces, with ionic strength adjusted to physiological conditions using 0.15 M NaCl. The resulting simulation systems

comprised approximately 250,000-300,000 atoms, with total system size dependent on the specific receptor-G protein complex dimensions and corresponding membrane requirements.

All MD simulations were performed using AMBER20[65] under NPT ensemble conditions at 310K and 1 bar. The FF19SB[66] force field was employed for protein parameters, while lipid21[67] and GAFF[68] force fields described lipid and ligand interactions, respectively. The OPC[69] water model was used for solvation, with long-range electrostatic interactions calculated via particle mesh Ewald. A 9 Å van der Waals cutoff was applied throughout all simulations.

System preparation involved energy minimization followed by 47.5 ns of equilibration. Energy minimization was performed using 2,500 steepest descent steps followed by 2,500 conjugate gradient steps. Temperature equilibration employed the Langevin thermostat (γ = 1.0 ps^{-1}) with gradual heating to 310 K over two consecutive 2.5 ns NVT phases using 1 fs timesteps. Pressure equilibration utilized the Berendsen barostat (τ_p = 1.0 ps) through a series of NPT phases: an initial 2.5 ns phase at 1 fs timestep, followed by two consecutive 10 ns phases at 2 fs timestep, and a final 20 ns phase at 2 fs timestep with restraints applied only to protein and ligand atom. Harmonic restraints were systematically reduced across the equilibration stages. Lipid restraints decreased progressively to 2.5, 2.5, 1.0, 0.5, 0.1, and 0.0 kcal/mol/Å² across the six equilibration phases. Protein and ligand restraints followed a similar pattern: 10.0, 5.0, 2.5, 1.0, 0.5, and 0.1 kcal/mol/Å². Production simulations were conducted for 1,000 ns per replica with five independent replicates per system, using 2 fs timesteps and coordinates saved every 1,000 ps.

Structural dynamics analyses were performed on simulation trajectories using AmberTools23 CPPTRAJ[70] for trajectory processing, imaging, and alignment. The simulation trajectories were aligned to the post-minimization starting structure using Cα atoms of residues (MT1 systems: S28–R54, F65–V84, S103–Y128, L145–N162, I189–Q216, F234–G258 and V278–Y295; and MT2 systems: A42–V65, F78–I101, A115–Y139, P158–P174, Y200–A230, F257–L272 and F290–I306) of the 7-transmembrane bundle as the reference framework. Root-mean-square deviation (RMSD) and root-mean-square fluctuation (RMSF) calculations were conducted relative to these aligned post-minimization structures to assess structural stability and regional flexibility throughout the simulation period.

For interaction energy analysis, noncovalent interaction energies between individual receptor residues and the G protein α5 helix were calculated using the NAMDenergy utility[71,72] (v1.6) distributed with VMD 1.9.3 [73]. The energy calculations employed a 9 Å cutoff distance with a 7.5 Å van der Waals smoothing switch distance. Energy analyses were performed on every 10th frame of the trajectories (100 frames total from each 1000-frame simulation), corresponding to a 10 ns sampling frequency, across 3 independent replicas of 1 μs MD simulations. Results are reported as mean ± standard deviation values calculated across all replicas to ensure statistical robustness of the findings.

Statistical analysis

All data are presented as the mean ± standard error of the mean (SEM). Statistical significance was assessed with paired t-test, and two-tailed p-values are reported (Fig. 1c–f, 4d, 5c), or multiple t-tests with Holm–Šidák correction for multiple comparisons (Fig. 2h) and exact p-values are reported throughout. One-way ANOVA followed by Bonferroni's multiple comparison test (Fig. 5b, 6c) or two-way ANOVA followed by the Holm–Šidák test (Fig. 2d) was used as appropriate. For unpaired comparisons in ex vivo and in vivo experiments, Welch's correction was applied, and one-tailed p-values are reported (Figures 1h, j, m). P-values < 0.05 were considered statistically significant, and exact p-values are indicated in the figures. All analyses were performed using GraphPad Prism versions 6 and 8 (GraphPad Software Inc., San Diego, CA). Multiple t-tests with Holm–Šidák correction for multiple comparisons and exact p-values are reported throughout. A p-value < 0.05 was considered statistically significant.

Declarations

Acknowledgments

We are grateful to Dr. Nevin Lambert (Augusta, Georgia, USA) for sharing some reagents for the G protein recruitment assay, Akihide Yoshimi (National Cancer Center, Tokyo, Japan) for sharing some materials of melatonin receptors, and Takashi Yoshimura (Nagoya University, Aichi, Japan) for helpful suggestions. We are grateful to the Institute of Comprehensive Medical Research, Division of

Animal Research Promotion Division (Aichi Medical University) for maintaining the mice, providing advanced research promotion for the expression analysis of proteins, and performing measurements of cAMP in *pars tuberalis* samples.

Funding

This work was supported by the Agence Nationale de la Recherche (ANR-19-CE16-0025-01 « mitoGPCR », ANR-21-CE18-0023 «alloGLP1R», GPCR-metab, the Fondation pour la Recherche Médicale (Equipe FRM DEQ.20130326503), the "Projet de recherche international" (PRI) Inserm édition 2023, the Institut National de la Santé et de la Recherche Médicale (INSERM) and the Centre National de la Recherche Scientifique (CNRS) to R.J. This work was supported by funding from the by the JST Fusion-Oriented Research for Disruptive Science and Technology (FOREST) (grant number JPMJFR215U), the Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for Scientific Research (C) (grant number 22K06132, 25K10187), the Vehicle Racing Commemorative Foundation, Japan Diabetes Foundation(Novo Nordisk Research grant), Suzuken Memorial Foundation, Daiwa Securities Health Foundation, SENSHIN Medical Research Foundation, Takahashi Industrial and Economic Research Foundation, Precise Measurement Technology Promotion Foundation, Hokuto Foundation for Bioscience, The Cell Science Research Foundation, The Mitsubishi Foundation and the Asahi Glass Foundation to AO.

This work was supported by JSPS KAKENHI grants 21H05037 (O.N.), JP20K15754 (T.K.), JP22K15072 (T.K.), JP24K01961 (T.K.), and 21J20897 (H.H.O.) ; the Japan Science and Technology Agency (JST) grant numbers JPMJCR20E2 (O.N.) ; JST PRESTO grant number JPMJPR22E4 (T.K.) ; the Japan Agency for Medical Research and Development (AMED), grant numbers JP233fa627001 (O.N.) ; the Platform Project for Supporting Drug Discovery and Life Science Research (Basis for Supporting Innovative Drug Discovery and Life Science Research (BINDS)) from AMED grant numbers JP25am121002 (support no. 3272, O.N.), JP25ama121012 (O.N.)

This study was supported by the JSPS KAKENHI (Grant Number 19K09962), Takeda Science Foundation to KI.

This work was supported by the Biotechnology and Biological Sciences Research Council (U.K.), grant BB/R007101/1 (to I.G.T.), R.M.'s PhD study is supported by the strategic priority PhD studentship from Northern Ireland

Department for Employment and Learning. This project made use of computational time on Kelvin-2 supported by Engineering and Physical Sciences Research Council (U.K.), grants: EP/T022175/1 and EP/W03204X/1.

Author Contribution

A.O., H.H.O., K.I., B.M. J.D., M.N., I.G.T., O.N. and R.J. conceived and designed experiments or contributed to critical discussion. A.O., B.M. performed biochemical in vitro experiments including the design of receptor mutants and chimera. K.I. performed in vivo studies. H.H.O., T.K. and K.K. performed structural studies including receptor expression and purification, cryoEM data collection and image processing and model building.

T.K. and K.K. performed G protein and Nb35 expression and purification. R.M. and I.G.T. performed molecular simulation studies. A.K., E.C. performed initial in vitro cAMP studies. A.O., H.H.O., K.I., R.M., B.M. I.G.T. and R.J. analysed data.

A.O., K.I., J.D., I.G.T. and R.J. obtained the funding. A.O., H.H.O., and R.J. wrote the first draft of the manuscript which was edited by K.I., B.M., A.K., E.C., J.D. I.G.T., O.N. and approved by all authors.

Competing interests

O.N. is a cofounder and scientific advisor for Curreio. All other authors declare no competing interests.

Figure legends

Fig 1. Melatonin MT$_1$ receptor activates the stimulatory G$_s$/cAMP pathway *in vitro*, *ex vivo* and *in vivo*

(a-f) Melatonin MT$_1$ receptor activated G$_s$-cAMP pathway in HEK293T cells. (a) (b) Concentration–response effect of melatonin on forskolin (FSK)-induced cAMP production (30 min, FSK [2 µM], IBMX [1 mM]) in HEK293T cells expressing MT$_1$-WT or MT$_2$-WT, assessed by HTRF assay without (a) or with (b) PTX treatment (400 ng/mL, 4 h). Data represent means ± SEM of at least 4 independent experiments. (c-f) NanoBiT complementation assay showing recruitment of LgBiT–miniG$_i$ (c,e) or LgBiT–miniG$_s$ (d,f) to human MT$_1$-NP (c, d) or MT$_2$-NP (e, f) upon melatonin stimulation (1 µM, 2 min). Data are presented as means ± SEM of five independent experiments. Statistical significance was determined using paired two-tailed t-tests, exact p-values are reported.

(g-j) Melatonin (MLT) induces cAMP production in mouse pituitary *pars tuberalis* (PT) *ex vivo*. (g) MLT (10 µM) were exposed to sliced PT / mediobasal hypothalamus (MBH) tissue complex of B6 mice kept under 12L/12D conditions. (h) Sixteen-hour exposure of MLT significantly increased cAMP levels in the PT/MBH compared with control 0.1 % DMSO exposure group. Data are means ± SEM of 4 independent experiments. (i) Western blot analysis of phosphorylate CREB (pCREB [Ser133]) in cultured PT/MBH tissues. MLT significantly increased pCREB. Data are means ± SEM of 4 independent experiments. Unpaired t-tests with Welch's correction were used and one-tailed p-values are reported.

(k-m) MLT induces cAMP production in mouse pituitary PT *in vivo*. (k) MLT (0.26 mM / 0.1 mL) was injected intraperitoneally (i.p.) at ZT8 to B6 mice kept under 8L/16D conditions. After 16 h, brains were extracted and sectioned coronally for immunohistochemistry. ZT, Zeitgeber time; ZT1 indicates light onset time. (l) Representative images of pCREB immunoreactivity (ir) in the PT and MBH of MLT-injected mice. 3V, Third ventricle. (m) Relative changes of pCREB-ir cells normalized by area in the PT. MLT injection increased pCREB (**$p < 0.01$, t-test). Data are means ± SEM of 6-7 independent experiments. Unpaired t-tests with Welch's correction were used and one-tailed p-values are reported.

Fig. 2 Cryo-EM structure of the MT$_1$-miniGs complex

(a) CryoEM density map of MT$_1$-miniG$_s$ with Gβ$_1$, Gγ$_2$ and Nb35. MT$_1$, miniG$_s$, Gβ$_1$, Gγ$_2$ and Nb35 were shown as blue, yellow, red, pink and brown, respectively. Map including micelle density was shown in transparent style.

(b) Overall structure of MT$_1$-miniG$_s$ with Gβ$_1$, Gγ$_2$ and Nb35. Melatonin molecule is shown as light purple CPK model.

(c) The interface between MT$_1$ and miniG$_s$.

(d–g) Concentration–response effect of melatonin on FSK-induced cAMP production (5 min, FSK (5µM) in HEK293T cells expressing MT1-WT (d) or mutant receptors Y128A (e), S132A (f), and L133A (g), monitored with the BRET-based CAMYEL biosensor. Data represent means ± SEM from six independent experiments.

(h) Melatonin (1 µM, 10 min)–induced cAMP production in the absence of FSK in HEK293T cells pretreated with PTX (400 ng/mL, 4 h) and expressing MT1-WT or mutant receptors (Y128A, S132A, L133A) assessed with the BRET-based CAMYEL biosensor. Data represent means ± SEM from four independent experiments. Multiple t-tests with Holm–Šidák correction were used for statistical comparisons, exact p-values are indicated.

(i-k) Structural comparison of miniG$_s$ protein among MT$_1$-miniG$_s$ (yellow), β$_2$AR-G$_s$ (green; PDB 3SN6) and GDP bound G$_s$ (gray; PDB 6EG8), focusing on (i) overall structure, (j) α5 helix and (k) TCAT motif. (l-n) Structural comparison with MT$_1$-miniG$_s$ (light blue and purple; PDB 7DB6), displaying (l) overall structure, (m) ICL2 and (n) TM3.

Fig. 3 Structural comparison with GPR61-G$_s$ complex and other G$_s$/G$_i$ complexes with the same receptor

(a), Amino acid sequences alignment of human MT$_1$, MT$_2$, GPR50, GPR61 and GPR151.

(b-e), Structural comparison between MT$_1$-miniG$_s$ and GPR61-miniG$_s$ (PDB 8KGK) displaying (b) overall, (c) ICL2 interface, (d) overall structure from cytoplasmic side and (e) α5 helix interface.

(f, g), Structural comparison among reported GPCR-G complexes with the same receptor showing (f) primarily G$_s$ coupling receptors including 5HT$_4$ (PDB 7XT9 and 7XTA), β$_1$AR (PDB 7S0G and 7S0F), EP4 (PDB 8GCP and 8GDA), or (g) primarily G$_q$ coupling receptors including CCKAR (PDB 7EZM, 7EZH and 7EZK) and FFAR4 (PDB 8H4L, 8ID9 and 8H4I).

Fig. 4 Structural basis of MT$_2$-MT$_1$-chimera Gs coupling

(a) Amino acid sequences alignment of human MT$_1$ and MT$_2$ chimeric receptors in TM5-ICL3-TM6 region.

(b,c) Concentration–response effect of melatonin on FSK-induced cAMP production (5 min, FSK [5µM]) in HEK293T cells expressing (b) MT$_1$–MT$_2$(TM5–ICL3–TM6) or (c) MT$_2$–MT$_1$(TM5–ICL3–TM6) chimeric receptors, assessed with the BRET-based CAMYEL biosensor. Data represent means ± SEM of six independent experiments.

(d) NanoBiT complementation assay showing recruitment of LgBiT–miniG$_i$ (left) or LgBiT–miniG$_s$ (right) to the MT$_2$–MT$_1$(TM5–ICL3–TM6)–NP receptor upon melatonin stimulation (1 µM, 2 min). Data present means ± SEM of five independent experiments. Statistical significance was determined using paired two-tailed t-tests, exact p-values are reported.

(e) CryoEM density map (left) and overall structure (right) of MT$_2$-MT$_1$(TM5-TM6)-miniG$_s$ with Gβ$_1$, Gγ$_2$ and Nb35. MT$_2$, MT$_1$(TM5-TM6), miniG$_s$, Gβ$_1$, Gγ$_2$ and Nb35 were shown as light green, blue, orange, red, pink and brown, respectively. Map including micelle density was shown in transparent style.

(f, g) Structural comparison of MT$_2$-MT$_1$(TM5-TM6)-miniG$_s$ with (f) MT$_1$-miniG$_s$ or (g) MT$_2$-G$_i$ (PDB 7VH0), including lateral and cytoplasmic viewpoints. MT$_2$ in MT$_2$-G$_i$ is shown as light purple, and G$_i$ in MT$_2$-G$_i$ is shown as green.

(h) The interface between MT$_2$-MT$_1$(TM5-TM6) and miniG$_s$ around TM3 and ICL2.

(i, j) Structural comparison between MT$_2$-MT$_1$(TM5-TM6)-miniG$_s$ and MT$_1$-miniG$_s$ at (i) TM3 and α5 helix or (j) ICL2.

(k) Structural comparison between MT$_2$-MT$_1$(TM5-TM6)-miniG$_s$ and MT$_2$-G$_i$ at TM5, TM6 and α5 helix.

Fig. 5 Identification of key determinants of MT$_1$-G$_s$ interaction by MT$_2$-MT$_1$ chimera receptors

(a) Amino acid sequences alignment of human MT$_2$ chimeric receptors.

(b-e) (upper) Schematic representation of chimera. (middle) Concentration–response effect of melatonin on FSK-induced cAMP production (10 min, FSK [5µM]) in HEK293T cells expressing (b) MT2–MT1(TM5c), (c) MT2–MT1(TM6), (d) MT2–MT1(ICL3) or (e) MT2–MT1(core ICL3) chimeric receptors, assessed with the BRET-based CAMYEL biosensor. Data represent means ± SEM of

seven independent experiments. (lower) Statistical comparison of concentration–response data at 0, 1 nM and 1 µM of melatonin. Data represent means ± SEM of six independent experiments. Statistical analysis was performed by one-way ANOVA followed by Bonferroni's multiple comparison test.

(f, g) NanoBiT complementation assay showing recruitment of LgBiT–miniGi (f) or LgBiT–miniGs (g) to the MT$_2$–MT$_1$(coreICL3)–NP receptor upon melatonin stimulation (1 µM, 2 min). Data represent means ± SEM of five independent experiments. Statistical significance was determined using paired two-tailed t-tests, exact p-values are reported.

Fig. 6 Molecular dynamics simulations support differential G protein binding modes and ICL3-dependent Gs coupling of MT1

(a) Representative MD simulation snapshots of MT$_1$-G protein complexes. Carbon colours indicate electrostatic energies, stick thickness represents van der Waals interactions with α5 helix. Only residues >1 kcal/mol in TM5-ICL3-TM6 are shown. AlphaFold3-modeled ICL3 (pink) highlights non-conserved hydrophobic residues. TM5 and TM6 are highlighted in darker colour. Key contacts: I210$^{5.61}$ with L348$^{G.H5.20}$, F354$^{GH5.26}$, and L353$^{GH5.25}$ (Gi deep binding) versus V217^{ICL3} with L388$^{G.H5.20}$ (Gs shallow binding).

(b) Amino acid sequence alignment of human MT$_2$ chimera and human MT$_1$ deletion mutant in TM5-ICL3 region.

(c,d) (upper) Concentration–response effect of melatonin on FSK-induced cAMP production (10 min, FSK [5µM]) in HEK293T cells expressing (c) MT2-MT1(coreICL3-V217A) and (d) MT1-ICL3-deletion mutant chimeric receptors, assessed with the BRET-based CAMYEL biosensor. Data represent means ± SEM of seven (c) or four (d) independent experiments. (Lower) Statistical comparison of concentration–response data (Upper) at 0, 1 nM and 1 µM of melatonin. Data represent means ± SEM of six independent experiments. Statistical analysis was performed by one-way ANOVA followed by Bonferroni's multiple comparison test.

Extended Data Fig. 1 Cryo-EM analysis of the MT$_1$-miniG$_s$ complex

The workflow of Cryo-EM analysis of MT$_1$-miniG$_s$ complex using RELION 3.1 and cryoSPARC v3.3.

Extended Data Fig. 2 Cryo-EM analysis of the MT$_2$ chimera-miniG$_s$ complex

The workflow of Cryo-EM analysis of MT$_2$ chimera-miniG$_s$ complex using RELION 3.1, RELION 4.0 and cryoSPARC v3.3.

Extended Data Fig. 3 Structural comparison of the MT$_1$-miniG$_s$ and MT$_2$-MT$_1$-miniG$_s$ complexes

a-e, Structural comparison of MT$_1$-miniG$_s$, MT$_1$-G$_i$ (PDB 7DB6) and crystal structure of MT$_1$ (PDB 6ME2), on (a) overall receptor, (b) ligand binding site, (c) PIF motif, (d) N(D)RY motif and (e) Na$^+$ binding site and NPxxY motif.

f-j, Structural comparison of MT$_2$-MT$_1$-miniG$_s$, MT$_2$-G$_i$ (PDB 7VH0) and crystal structure of MT$_2$ (PDB 6ME9), on (f) overall receptor, (g) ligand binding site, (h) PIF motif, (i) N(D)RY motif and (j) Na$^+$ binding site and NPxxY motif.

k-m, Structural comparison of G$_s$ protein among MT$_2$-MT$_1$-miniG$_s$, β$_2$AR-G$_s$ (PDB 3SN6) and GDP-bound inactive state (PDB 6EG8), on (k) overall region, (l) α5 helix region and (m) GDP/GTP binding site.

Extended Data Fig. 4 Structural comparison with GPR174-G$_s$ complex and 5HT$_4$-G$_s$/-G$_i$ complex

a, Structural comparison between MT$_1$-miniG$_s$ and GPR174-G$_s$ (PDB 7VX3).

b, Structural comparison among all the structures of GPR174-G$_s$ (PDB 7VX3, 8KH5 and 8IZB).

c, Structural comparison between MT$_1$-miniG$_s$, GPR174-G$_s$ (PDB 7VX3), 5HT$_4$-G$_s$ (PDB 7XT9) and 5HT$_4$-G$_i$ (PDB 7XTA).

Extended Data Fig. 5 Structural comparison of GPCR-G$_s$/-G$_q$ and GPCR-G$_i$/-G$_q$ complex with the same receptor

a, Structural comparison between NK1R-G$_s$ (PDB 7P02) and NK1R-G$_q$ (PDB 7P00, primary G$_q$ coupling).

b, Structural comparison among GPCR-G$_i$/-G$_q$ complexes with the same receptor; ETBR-G$_i$ (PDB 8IY5, primary coupling)/-G$_q$ (PDB 8HCX), GASR-G$_q$ (PDB 7F8W, primary coupling)/-G$_i$ (PDB 7F8V), GHSR-G$_q$ (PDB 7W2Z, primary coupling)/-G$_i$ (PDB 7F9Y), MRGPRX2-G$_q$ (PDB 7S8N)/-G$_i$ (PDB 7S8M), NMUR2-G$_q$ (PDB 7XK8)/-G$_i$ (PDB 7W57), NTSR1-G$_i$ (PDB 6OS9, primary coupling)/-G$_q$ (PDB 8FMZ), OXYR-G$_q$ (PDB 7RYC)/-G$_i$ (PDB 7QVM), US28-G$_q$ (PDB 7RKF)/-G$_i$ (PDB 7RKM) and SSTR2-G$_i$ (PDB 7Y24, primary coupling)/-G$_q$

(PDB 7Y26).

Extended Data Fig. 6 Molecular dynamics simulations support differential G protein binding modes and ICL3-dependent Gs coupling selectivity in the melatonin receptors

a, Representative MD simulation snapshots of the melatonin receptor-G protein complexes. Carbon colours indicate electrostatic energies, stick thickness represents van der Waals interactions with α5 helix. Only residues >1 kcal/mol in TM5/TM6/ICL3 are shown. AlphaFold3-modeled ICL3 (pink) highlights non-conserved hydrophobic residues. TM5 and TM6 are highlighted in darker colour. Key contacts: $I210^{5.61}$ with $L348^{G.H5.20}$, $F354^{GH5.26}$, and $L353^{GH5.25}$ (Gi deep binding) versus $V217^{ICL3}$ with $L388^{G.H5.20}$ (G_s shallow binding). The MT2-Gs test model shows compromised interactions due to alanine substitutions.

b, Root mean square fluctuation (RMSF) values of the G protein α5 helix across different melatonin receptor-G protein complexes.

c, G protein α5 helix flexibility (RMSF).

d, ICL3-α5 helix interaction energies. Data in b-d: mean ± SEM, 5×1μs simulations.

References

1. Hauser, A.S. et al. Common coupling map advances GPCR-G protein selectivity. *Elife* **11** (2022).
2. Huang, S. et al. GPCRs steer G(i) and G(s) selectivity via TM5-TM6 switches as revealed by structures of serotonin receptors. *Mol Cell* **82**, 2681-2695 e6 (2022).
3. Alegre, K.O. et al. Structural basis and mechanism of activation of two different families of G proteins by the same GPCR. *Nat Struct Mol Biol* **28**, 936-944 (2021).
4. Huang, S.M. et al. Single hormone or synthetic agonist induces G(s)/G(i) coupling selectivity of EP receptors via distinct binding modes and propagating paths. *Proc Natl Acad Sci U S A* **120**, e2216329120 (2023).
5. Zhou, Y. et al. Molecular insights into ligand recognition and G protein coupling of the neuromodulatory orphan receptor GPR139. *Cell Res* **32**, 210-213 (2022).

6. Dubocovich, M.L. et al. International Union of Basic and Clinical Pharmacology. LXXV. Nomenclature, classification, and pharmacology of G protein-coupled melatonin receptors. *Pharmacol Rev* **62**, 343-380 (2010).

7. Hegron, A. et al. Identification of Key Regions Mediating Human Melatonin Type 1 Receptor Functional Selectivity Revealed by Natural Variants. *ACS Pharmacol Transl Sci* **4**, 1614-1627 (2021).

8. Okamoto, H.H., Cecon, E., Nureki, O., Rivara, S. & Jockers, R. Melatonin receptor structure and signaling. *J Pineal Res* **76**, e12952 (2024).

9. Johansson, L.C. et al. XFEL structures of the human MT2 melatonin receptor reveal the basis of subtype selectivity. *Nature* **569**, 289-292 (2019).

10. Stauch, B. et al. Structural basis of ligand recognition at the human MT1 melatonin receptor. *Nature* **569**, 284-288 (2019).

11. Ishchenko, A. et al. Toward G protein-coupled receptor structure-based drug design using X-ray lasers. *IUCrJ* **6**, 1106-1119 (2019).

12. Okamoto, H.H. et al. Cryo-EM structure of the human MT1-Gi signaling complex. *Nat Struct Mol Biol* **28**, 694-701 (2021).

13. Wang, Q. et al. Structural basis of the ligand binding and signaling mechanism of melatonin receptors. *Nat Commun* **13**, 454 (2022).

14. Zhao, D. et al. Melatonin Synthesis and Function: Evolutionary History in Animals and Plants. *Front Endocrinol (Lausanne)* **10**, 249 (2019).

15. Liu, J. et al. MT1 and MT2 Melatonin Receptors: A Therapeutic Perspective. *Annu Rev Pharmacol Toxicol* **56**, 361-83 (2016).

16. Cabello, N. et al. Metabotropic glutamate type 5, dopamine D2 and adenosine A2a receptors form higher-order oligomers in living cells. *J Neurochem* **109**, 1497-507 (2009).

17. Morgan, P.J. et al. Melatonin receptors couple through a cholera toxin-sensitive mechanism to inhibit cyclic AMP in the ovine pituitary. *J Neuroendocrinol* **7**, 361-9 (1995).

18. Chen, L. et al. Melatonin receptor type 1 signals to extracellular signal-regulated kinase 1 and 2 via Gi and Gs dually coupled pathways in HEK-293 cells. *Biochemistry* **53**, 2827-39 (2014).

19. Tse, L.H., Cheung, S.T., Lee, S. & Wong, Y.H. Real-Time Determination of Intracellular cAMP Reveals Functional Coupling of G(s) Protein to the Melatonin MT(1) Receptor. *Int J Mol Sci* **25**(2024).

20.	Dixon, A.S. et al. NanoLuc Complementation Reporter Optimized for Accurate Measurement of Protein Interactions in Cells. *ACS Chem Biol* **11**, 400-8 (2016).

21.	Laschet, C., Dupuis, N. & Hanson, J. A dynamic and screening-compatible nanoluciferase-based complementation assay enables profiling of individual GPCR-G protein interactions. *J Biol Chem* **294**, 4079-4090 (2019).

22.	Klosen, P. et al. MT1 and MT2 melatonin receptors are expressed in nonoverlapping neuronal populations. *J Pineal Res* **67**, e12575 (2019).

23.	Yasuo, S., Yoshimura, T., Ebihara, S. & Korf, H.W. Melatonin transmits photoperiodic signals through the MT1 melatonin receptor. *J Neurosci* **29**, 2885-9 (2009).

24.	Morgan, P., Lawson, W., Davidson, G. & Howell, H. Melatonin inhibits cyclic AMP in cultured ovine pars tuberalis cells. *J Mol Endocrinol* **3**, R5-R8 (1989).

25.	Herrera, L.P.T. et al. GPCRdb in 2025: adding odorant receptors, data mapper, structure similarity search and models of physiological ligand complexes. *Nucleic Acids Res* (2024).

26.	Liu, Q. et al. Ligand recognition and G-protein coupling selectivity of cholecystokinin A receptor. *Nat Chem Biol* **17**, 1238-1244 (2021).

27.	Kobayashi, K. et al. Endogenous ligand recognition and structural transition of a human PTH receptor. *Mol Cell* **82**, 3468-3483 e5 (2022).

28.	Inoue, A. et al. Illuminating G-Protein-Coupling Selectivity of GPCRs. *Cell* **177**, 1933-1947 e25 (2019).

29.	Matic, M., Miglionico, P., Tatsumi, M., Inoue, A. & Raimondi, F. GPCRome-wide analysis of G-protein-coupling diversity using a computational biology approach. *Nat Commun* **14**, 4361 (2023).

30.	Sadler, F. et al. Autoregulation of GPCR signalling through the third intracellular loop. *Nature* **615**, 734-741 (2023).

31.	von Gall, C. et al. Rhythmic gene expression in pituitary depends on heterologous sensitization by the neurohormone melatonin. *Nature Neuroscience* **5**, 234-238 (2002).

32.	Hazlerigg, D.G., Gonzalez, B.A., Lawson, W., Hastings, M.H. & Morgan, P.J. Prolonged exposure to melatonin leads to time-dependent sensitization of adenylate cyclase and down-regulates melatonin receptors in pars tuberalis cells from ovine pituitary. *Endocrinology* **132**,

285-92 (1993).

33. Stehle, J.H. et al. Molecular cloning and expression of the cDNA for a novel A2-adenosine receptor subtype. *Mol Endocrinol* **6**, 384-93 (1992).

34. Oishi, A. et al. Orphan GPR61, GPR62 and GPR135 receptors and the melatonin MT2 receptor reciprocally modulate their signaling functions. *Sci Rep* **7**, 8990 (2017).

35. Kroeze, W.K. et al. PRESTO-Tango as an open-source resource for interrogation of the druggable human GPCRome. *Nat Struct Mol Biol* **22**, 362-9 (2015).

36. Oishi, A., Dam, J. & Jockers, R. beta-Arrestin-2 BRET Biosensors Detect Different beta-Arrestin-2 Conformations in Interaction with GPCRs. *ACS Sens* **5**, 57-64 (2020).

37. Kusama, K. et al. Electrically Driven, Bioluminescent Compliant Devices for Soft Robotics. *ACS Appl Mater Interfaces* **17**, 11248-11258 (2025).

38. Somalo-Barranco, G. et al. Mitochondria-targeted melatonin photorelease supports the presence of melatonin MT1 receptors in mitochondria inhibiting respiration. *Cell Chem Biol* **30**, 920-932 e7 (2023).

39. Ikegami, K., Shigeyoshi, Y. & Masubuchi, S. Circadian Regulation of IOP Rhythm by Dual Pathways of Glucocorticoids and the Sympathetic Nervous System. *Invest Ophthalmol Vis Sci* **61**, 26 (2020).

40. Ikegami, K., Katou, Y., Higashi, K. & Yoshimura, T. Localization of circadian clock protein BMAL1 in the photoperiodic signal transduction machinery in Japanese quail. *J Comp Neurol* **517**, 397-404 (2009).

41. Kobayashi, K. et al. Endogenous ligand recognition and structural transition of a human PTH receptor. *Mol Cell* **82**, 3468-3483 e5 (2022).

42. Gotzke, H. et al. The ALFA-tag is a highly versatile tool for nanobody-based bioscience applications. *Nat Commun* **10**, 4403 (2019).

43. Zivanov, J. et al. New tools for automated high-resolution cryo-EM structure determination in RELION-3. *Elife* **7**(2018).

44. Punjani, A., Rubinstein, J.L., Fleet, D.J. & Brubaker, M.A. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296 (2017).

45. Bepler, T. et al. Positive-unlabeled convolutional neural networks for particle picking in cryo-electron micrographs. *Nat Methods* **16**, 1153-1160 (2019).

46. Ramlaul, K., Palmer, C.M., Nakane, T. & Aylett, C.H.S. Mitigating local over-fitting during single particle reconstruction with SIDESPLITTER. *J Struct Biol* **211**, 107545 (2020).

47. Billesbolle, C.B. et al. Structural basis of odorant recognition by a human odorant receptor. *Nature* **615**, 742-749 (2023).

48. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr* **60**, 2126-32 (2004).

49. Afonine, P.V. et al. Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta Crystallogr D Struct Biol* **74**, 531-544 (2018).

50. Adams, P.D. et al. PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-21 (2010).

51. Schrödinger, L. *Maestro*, (Schrödinger, LLC, New York, NY, 2025).

52. Jumper, J. et al. Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589 (2021).

53. Heo, L. & Feig, M. Multi-state modeling of G-protein coupled receptors at experimental accuracy. *Proteins* **90**, 1873-1885 (2022).

54. Olsson, M.H., Sondergaard, C.R., Rostkowski, M. & Jensen, J.H. PROPKA3: Consistent Treatment of Internal and Surface Residues in Empirical pKa Predictions. *J Chem Theory Comput* **7**, 525-37 (2011).

55. Sondergaard, C.R., Olsson, M.H., Rostkowski, M. & Jensen, J.H. Improved Treatment of Ligands and Coupling Effects in Empirical Calculation and Rationalization of pKa Values. *J Chem Theory Comput* **7**, 2284-95 (2011).

56. Lomize, M.A., Pogozheva, I.D., Joo, H., Mosberg, H.I. & Lomize, A.L. OPM database and PPM web server: resources for positioning of proteins in membranes. *Nucleic Acids Res* **40**, D370-6 (2012).

57. Jo, S., Kim, T., Iyer, V.G. & Im, W. CHARMM-GUI: a web-based graphical user interface for CHARMM. *J Comput Chem* **29**, 1859-65 (2008).

58. Wu, E.L. et al. CHARMM-GUI Membrane Builder toward realistic biological membrane simulations. *J Comput Chem* **35**, 1997-2004 (2014).

59. Jo, S., Lim, J.B., Klauda, J.B. & Im, W. CHARMM-GUI Membrane Builder for mixed bilayers and its application to yeast membranes. *Biophys J* **97**, 50-8 (2009).

60. Jo, S., Kim, T. & Im, W. Automated builder and database of

protein/membrane complexes for molecular dynamics simulations. *PLoS One* **2**, e880 (2007).

61. Lee, J. et al. CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. *J Chem Theory Comput* **12**, 405-13 (2016).

62. Kim, S. et al. CHARMM-GUI ligand reader and modeler for CHARMM force field generation of small molecules. *J Comput Chem* **38**, 1879-1886 (2017).

63. Lee, J. et al. CHARMM-GUI Membrane Builder for Complex Biological Membrane Simulations with Glycolipids and Lipoglycans. *J Chem Theory Comput* **15**, 775-786 (2019).

64. Lee, J. et al. CHARMM-GUI supports the Amber force fields. *J Chem Phys* **153**, 035103 (2020).

65. Case, D.A.B., K.; Ben-Shalom, I. Y.; Brozell, S. R.; Cerutti, D. S.;Cheatham, T. E., et al. *AMBER 2020*, (University of California, San Francisco, San Francisco, 2020).

66. Tian, C. et al. ff19SB: Amino-Acid-Specific Protein Backbone Parameters Trained against Quantum Mechanics Energy Surfaces in Solution. *J Chem Theory Comput* **16**, 528-552 (2020).

67. Dickson, C.J., Walker, R.C. & Gould, I.R. Lipid21: Complex Lipid Membrane Simulations with AMBER. *J Chem Theory Comput* **18**, 1726-1736 (2022).

68. He, X., Man, V.H., Yang, W., Lee, T.S. & Wang, J. A fast and high-quality charge model for the next generation general AMBER force field. *J Chem Phys* **153**, 114502 (2020).

69. Gowers, R. et al. MDAnalysis: A Python Package for the Rapid Analysis of Molecular Dynamics Simulations. 98-105 (2016).

70. Michaud-Agrawal, N., Denning, E.J., Woolf, T.B. & Beckstein, O. MDAnalysis: a toolkit for the analysis of molecular dynamics simulations. *J Comput Chem* **32**, 2319-27 (2011).

71. Phillips, J.C. et al. Scalable molecular dynamics on CPU and GPU architectures with NAMD. *J Chem Phys* **153**, 044130 (2020).

72. Phillips, J.C. et al. Scalable molecular dynamics with NAMD. *J Comput Chem* **26**, 1781-802 (2005).

73. Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular

dynamics. *J Mol Graph* **14**, 33-8, 27-8 (1996).

Figure 1 | Melatonin MT1 receptor activates both the inhibitory Gi/cAMP and the stimulatory Gs/cAMP in vitro and in the pituitary pars tuberalislex

Figure 2 | Cryo-EM structure of the MT$_1$-miniG$_s$ complex

Figure 3 | Structural comparison with GPR61-Gs complex and other Gs/Gi complexes with the same receptor

Fig. 4 | Structural basis of MT₂-MT₁-chimera-G_s coupling

Figure 5 | Identification of key determinants of MT$_1$-G$_s$ interaction by MT$_2$-MT$_1$ chimera receptors

Figure 6 | Molecular dynamics simulations support differential G protein binding modes and ICL3-dependent G$_s$ coupling of MT$_1$