Virus induced transgene- and tissue-culture free heritable genome editing in tomato

Ye Liu[1]*, Trevor Weiss[1], Jinhee Lee[1], Jessica Powell[1], Shi Ying Charlize Choo[1], Elnaz Roshannai[1], Maris Kamalu[1], Jasmine Amerasekera[1], Suhua Feng[1], Steven E. Jacobsen[1,2]*

[1] Department of Molecular, Cell and Developmental Biology, University of California Los Angeles, Los Angeles, CA, USA.

[2] Howard Hughes Medical Institute, University of California Los Angeles, Los Angeles, CA, USA.

* Correspondence: ly22@ucla.edu and jacobsen@ucla.edu

Abstract

Genome editing has emerged as a powerful tool for crop breeding. However, most commonly used approaches rely on tissue culture and the use of transgenic materials, which are time-consuming and labor-intensive, and often exhibit strong genotype dependency. Here we utilized tobacco rattle virus (TRV)-mediated ISYmu1 genome editing, coupled with *in planta* shoot regeneration, to achieve somatic and heritable genome editing in different tomato cultivars. By targeting the *SlPDS* gene, we successfully generated virus- and transgene-free homozygous mutant progeny in one generation. We further verified the method by generating somatic edits in *SlDA1*, a gene with potential agronomic relevance. Given the wide host range of TRV and its compatibility with *in planta* shoot regeneration, our system could be broadly applicable for rapid, non-transgenic and heritable genome editing, thereby advancing functional genomics and crop improvement.

Introduction

The emergence of genome editing technology presents tremendous potential for accelerating crop improvement[1, 2]. However, despite the rapid adoption of genome editing in many major crops, two significant bottlenecks remain: efficient delivery of gene-editing reagents and the efficient generation of transgene-free plants. Conventional plant genome editing often relies on the use of tissue culture and plant transformation, which typically involves introducing the editing construct into plants through *Agrobacterium tumefaciens*-mediated transformation, selecting transgenic plants carrying the desired mutation, and subsequently removing the transgenic components by crossing[3, 4]. This method is time- and labor- intensive and is limited to specific crop species and genotypes, which hinders the application of genome editing in crop plants.

To overcome these barriers, several plant viruses have been engineered to deliver single-guide RNAs (gRNAs) directly to transgenic plants expressing SpCas9 for virus induced genome editing (VIGE) in both dicotyledonous and monocotyledonous crop species[5-8]. For example, some RNA viral vectors, such as *Tobacco Rattle Virus* (TRV) and *Barley Stripe Mosaic Virus* (BMSV) have been shown to deliver the gRNA to meristematic and germline cells to achieve heritable editing, in part with the help of RNA mobility sequences[9-12]. However, most plant viruses are excluded from meristematic tissues, limiting the access to the germline and thus reducing the potential to generated gene-edited progeny plants[13, 14]. Moreover, due to the limited cargo capacity of most viruses, the application of these methods still requires the additional steps of using tissue-culture and transformation to obtain the overexpressing-Cas9 line as well as crosses to segregate away the transgenes after edits are obtained[15].

A recent strategy to circumvent these limitations was to engineer TRV to deliver the compact ISYmu1 TnpB nuclease and the gRNA[16], to enable transgene-free germline editing in *Arabidopsis*[17]. It remains unknown, however, whether this approach can be applied to additional species. Alternatively, large-capacity vectors[18], such as *Barley Yellow Striate Mosaic Virus* (BYSMV) have delivered both Cas9 and gRNAs to achieve

heritable editing in wheat[19]. However, this approach faces limitations due to its

complexity and dependence on insect transmission, making it difficult to adapt to other

plant species. Thus, the development of simple, rapid, and broadly applicable methods

for producing transgene-free, heritable genome-edited crops without the need for tissue

culture remains an urgent priority.

In this study, we established a strategy that utilized TRV-mediated delivery of the

compact ISYmu1 TnpB endonuclease and gRNA in tomato. Using this system, we first

targeted the *SlPDS* gene to successfully generate somatically edited shoots, and also

observed transmission of edited alleles to the next generation. We further extended the

approach to *SlDA1*, a gene of potential agronomic relevance, and achieved highly

efficient somatic editing. Together, these results demonstrate that our platform enabled

efficient somatic editing and stable transmission of targeted mutations in a crop, thereby

providing a streamlined one-step strategy for heritable and transgene-free genome

editing in tomato that bypass tissue-culture.

Results

Viral Delivery of ISYmu1for somatic genome editing in tomato.

Encouraged by our recent work demonstrating that TRV delivery of both ISYmu1 and

gRNA induced heritable transgene-free germline editing in *Arabidopsis*[17], we reasoned

that this strategy represents an ideal system for genome modification that can

completely bypass transgenics and tissue culture steps in crops like tomato. To this end,

we engineered a pTRV2-TnpB-gRNA vector fused with a truncated mutant FT

sequence of tomato (mSlFT) (Fig. 1A) , which has been shown to enhance mobility of

gRNA between cells[20]. We then generated a pTRV2-TnpB-gRNA^SlPDS-mSlFT vector

that expresses a gRNA targeting the tomato phytoene desaturase gene (*SlPDS*), which

was delivered to a red cherry cultivar via an *Agrobacterium tumefaciens* cotyledon

infiltration method (Fig. 1B, S1A). Three weeks post infiltration, DNA was isolated

from young leaves and used as a template for PCR, followed by NGS analysis of PCR

amplicons flanking the *SlPDS* target site. Of the 20 plants infiltrated we observed

3(15%) plants with somatic editing, ranging from 0.8%~6.93% indels in systemically

infected leaves (Fig. S1B). We collected seeds from #13 plants showing 6.93% somatic

editing to screen for heritability of edited alleles, however, these somatic mutations

failed to be transmitted to the progeny seedlings. These data indicated that TRV can

deliver ISYMu1 and gRNA and induce somatic genome editing in tomato.

Induction of Gene Edited *De Novo* Shoots via Viral Delivery of ISYMu1 and

gRNA[SlPDS]

Recent studies have demonstrated that heritable editing was achieved by inducing *de*

novo shoots from edited somatic cells through trimming TRV-infected plants, or

through TRV-mediated gRNA delivery to axillary meristems in tomato[21]. In addition,

emerging *in planta* regeneration approaches offer the potential for more efficient,

induction of *de novo* shoots[22, 23]. This inspired us to explore a heritable editing strategy

in which genome editing was first induced in somatic cells, followed by the

regeneration of *de novo* shoots from the edited cells. To do this we removed the upper

shoot and all the axillary shoots, and then injected *Agrobacterium* encoding TRV-TnpB-

gRNA[SlPDS]-mSlFT into all wounding sites of the main stem and axillary regions (Fig.

1C, S2 A-C). Five weeks after injection, both callus-like tissue or new shoots arose

from the wounding sites, suggesting these *de novo* shoots arose from either direct or

indirect regeneration or both (S2 D, 2 E). A total of 49 plants were injected, from which

72 *de novo* shoots were initiated (Table S1). One of the *de novo* shoots (#3) showed

fully or partial white sectors on the plant similar to the known *slpds* mutant phenotype

(Fig. 1D-G), suggesting biallelic mutations in the *SlPDS* gene. To confirm this, we first

performed Sanger sequencing on three randomly selected green leaves of this shoot,

which all revealed 5-bp deletion as indicated by the presence of two sets of peaks in the

sequencing traces at the target site (Fig. 1I). Further amplicon-sequencing (amp-seq)

analysis showed that this plant exhibited 56.66% somatic editing efficiency in green

leaves (Fig. 1J), consisting of 48.51% of the 5-bp deletion and another 8.15% with a 4-

bp deletion. Interestingly, as the plant grew, the petals of all of the flowers were white

(Fig. 1H), but the fruit did not display a photobleaching phenotype. We also isolated genomic DNA from different photobleached organs to measure the mutation frequency. Amp-seq analysis showed that different tissues displayed different mutation patterns and editing efficiencies (S3A, 3B), indicated that TRV-TnpB-gRNA[SlPDS]#3 was chimeric and that genome editing was continuing to occur late in the development of this shoot.

For other new developed shoots without photobleaching phenotype (Fig. 1K-M), we first performed Sanger sequencing as a preliminary screen. The results showed mixed peaks in TRV-TnpB-sgRNA[SlPDS]#7, #17, #43 and #70 lines out of 71 shoots (S4A-F), indicating a high level of editing. To quantify this editing, tissue samples from these lines were collected for amp-seq analysis, which revealed somatic mutation frequencies ranging from 25.21% to 49.69% (Fig. 1N-P, S5). Notably, #7 and #17 shoots appeared to carry a single predominant deletion with a frequency of roughly 50% (Fig. 1N, 1O), suggesting that these lines were heterozygous for a mutant allele that likely arose very early in the development of the *de novo* shoot. These data demonstrate TRV-TnpB-gRNA[SlPDS]-mSlFT can induce a high level of somatic edits in *de novo* shoots.

Heritable Genome Editing via Viral Delivery of ISYmu1 and gRNA[SlPDS]

We next investigated whether targeted mutations could be transmitted to the next generation. A total of 96 seeds from four lines (#3, #7, #17, #43) were sown on 1/2 MS plates. After one week, albino seedings were observed in the progeny of shoots 3# and #17, suggesting biallelic mutation in the *SlPDS* gene (Fig. 2A, 2B, 2E, 2G). To characterize the genotypes of the seedlings we performed Sanger sequencing. Analysis of seedlings from shoot #3 revealed 27 homozygous mutants (28%) with a 5-bp frame-shift deletion, 39 heterozygous mutants (41%) and 30 wild type (31%), consistent with a roughly 3:1 segregation ratio and confirming that the parent plant was likely heterozygous for the 5-bp deletion (Fig. 2I). Similarly, the seedlings of shoot #17 showed 23 homozygotes for a 4-bp deletion (24%), 54 heterozygotes (51%) and 19

wild type (20%) (Fig. 2I), and shoot #7 produced 27% homozygotes (9-bp deletion) and 51% heterozygotes. The progeny of shoot #47 on the other hand showed a lower transmission of mutations, with only 2% homozygotes and 24% heterozygotes for a 3-bp deletion. It is not surprising that we did not observe albino plants in the progeny of #7 and #43 plants (Fig. 2B, 2D, 2F, 2H), since these lines carried 9-bp or 3-bp deletions, respectively, which would be in frame deletions. Together these data indicate that TRV deliver of ISYMu1 and gRNA fused to mSlFT is capable of inducing heritable editing in tomato.

Previous studies have shown that TRV is not transmitted to the next generation in plants[17, 24]. To confirm that the TRV was absent in the progeny of a TRV-infected plants, reverse transcription PCR was performed on 5 albino progenies of both #3 and #17 parent shoots (Fig. 2J). As expected, TRV and TnpB mRNA signals were not present in any of the albino offspring seedings, confirming that the TRV-TnpB-gRNA-mSlFT system can be used to generate edited virus-free progeny.

Efficient Somatic Editing of *SlDA1* in *De Novo* Shoots Using TRV To Express ISYMu1 and gRNA

Many prior demonstrations of novel genome-editing approaches have relied on targeting genes with an easy phenotypic readout, such as *PDS* [25-27]. To demonstrate that our system can support editing of other genes, we targeted the *DA1* gene of tomato, whose homologous genes play an important role in the regulation of seed and organ size in other plant species[28-31], but which remains uncharacterized in tomato. To this end, we first conducted a Blast search using the *Arabidopsis* DA1 protein sequence as a query against the tomato genome database and, combined with phylogenetic analysis, identified that *Solyc04g079840* is the likely tomato *DA1* homolog (Fig. 3A), hereafter referred to as *SlDA1*. We then generated a TRV–TnpB-gRNA[SlDA1]-mSlFT construct and utilized the same method (Fig. S2) to infect a different tomato cultivar, M82, so that we could simultaneously test whether our system functions effectively in different tomato cultivars (Fig. 3B). Approximately 5 weeks after injection, 41 *de novo* shoots were initiated from 27 injected M82 plants (Table S2). Three random leaves from each of the

41 *de novo* shoots were collected for genomic DNA extraction and Sanger sequencing analysis. We observed mixed sequencing peaks near the *SlDA1* target site in four of the TRV–TnpB-gRNA[SlDA1]-mSlFT infected shoots, #3, #23, #32 and #38 (S6 A-F), suggesting the presence of mutations. Further amp-seq analysis of these four shoots detected indel frequencies from 63.36% to 99.7% in the *SlDA1* gene (Fig. 3C, S7). Notably, shoots #3 and #23 contained more than 99% edited reads consisting of roughly 50% of one type of indel and 50% of another type of indel, suggesting that these two shoots are biallelic mutants with two different mutations arose on the two homologous chromosomes very early during shoot development. These results show that the TRV– TnpB-gRNA-mSlFT system can efficiently induce somatic genome editing in *de novo* shoots targeting *SlDA1*.

Discussion

To advance modern crop genome editing it is important to develop rapid and convenient strategies to generate transgene-free and heritable genome edited crop plants without relying on tissue culture. In a number of recent studies, tissue-culture-free heritable gene editing was achieved by delivering the gRNA to Cas9-expressing tomato plants using TRV[20, 21, 32]. However, this approach still requires the generation of Cas9-expressing plants via tissue culture and plant transformation, as well as self- or back-crossing to remove the transgenic components after the edits are obtained. In this study, we delivered TRV vectors expressing TnpB and its gRNA into wounding sites, which efficiently induced targeted mutations at the *SlPDS* and *SlDA1* in *de novo* shoots. Further, we demonstrated that the *SlPDS* mutations were stably transmitted to the progeny. Thus, our method simultaneously overcomes two major obstacles for tomato genome editing, dependence on tissue culture and the requirement of transgenic components. We targeted both the *SlPDS* gene, a commonly used marker gene with an easily screenable phenotype, as well as the *SlDA1* gene not yet studied in tomato, and utilized two tomato ecotypes in the study. This suggests that our system will be useful for gene function analysis and crop breeding, and that it may be effective in a wide range of tomato genotypes.

The TRV vector in this study possesses a broad host range of more than 400 plant species[33], suggesting that the approach described here should work in many additional crops. In addition, ISYmu1 is compact in size and can likely therefore be incorporated into other viral vectors with cargo capacities comparable to TRV, such as *Potato virus X* (PVX) and *Pea early browning virus* (PEBV)[7, 34], further expanding the range of crops that could benefit from this approach. Given that *in planta* viral infection strategies have already been demonstrated with many different viruses and plant species, the potential number of systems that could utilize ISYmu1 is large. Our approach should be especially useful in recalcitrant species that are difficult to genetically transform. In the future, further improving the editing efficiency and target sequence range of ISYMu1 and related TnpB enzymes will be essential to fully realize the potential of this strategy.

In summary, by integrating the advantages of a compact RNA-guided nuclease, viral infection, and would-induced shoot regeneration, this work demonstrates a method that could advance functional genomics and precision breeding in tomato and other crops.

Methods and Materials

Plant material and growth conditions

In this study, two tomato genotypes were used for viral infection, a red cherry type tomato (Amazon.com) and M82 (USDA). Seeds were soaked in water at 55°C for 20 minutes, then planted in germination trays and placed in a long-day growth chamber (16-h light and 8-h dark; day 25°C/night 21°C; light intensity: 150 μmol/m^2/s; 50% humidity) for about 3 weeks.

A different procedure was followed for growing plants used for seed harvesting. Two weeks after germination, small seedlings were transferred from germination trays into large pots. They were grown in a greenhouse with long-day conditions (16-h light and 8-h dark; day 24°C/night 21°C; light intensity:150 μmol/m^2/s; 50% humidity) for about 3 months.

Plasmids construction

To generate the TRV2-TnpB-wRNA-mSlFT (pYL001) plasmid, the TnpB-wRNA-*ccdb*-HR-mSlFT sequence with 20 bp overlapping as a gene block was synthesized from Integrated DNA Technologies (IDT). The pMK435 vector was digested by SacI (NEB, R3156S) and PmlI (NEB, R0532S) for 2 hours and the largest fragment was purified using QIAQuick Gel Cleanup kit (Qiagen, 28704). Next, the gene block and digested purified pMK435 plasmid fragment were used to make the pYL001 plasmid using NEBuilder HiFi DNA Assembly (NEB, E2621) according to the manufacturer's protocol. Finally, the reaction was transformed into the ccdB Survival2 T1R Competent Cells (Invitrogen, A10460).

The pYL001 plasmid was used as a base vector to construct the TRV2 vector with *SlPDS* and *SlDA1* gRNA using the Anneal and Golden Gate strategy. First, two target site oligos were synthesized from Integrated DNA Technologies (IDT) and diluted to 100 μm, and then phosphorylated and annealed. Next, the annealed double-stranded DNA and the pYL001 plasmid were used in a Golden Gate reaction using PaqCI® (NEB, R0745) and T4 DNA Ligase (NEB, M0202), followed by the Golden Gate Assembly Protocol. Finally, the reaction was transformed into the 10-beta competent *E. coli* (NEB, C3019). All the plasmids were confirmed using Primordium whole-plasmid sequencing. The target site primers are listed in Table S2

Viral infection and *de novo* shoot induction

TRV1 and TRV2 related plasmids were transformed into *Agrobacterium* strain GV3101 electrocompetent cells (Golabio, CC-207) and grown on LB plates with Kan (50 μg/ml) and RIF (30 μg/ml) for 48 hours at 28°C. A single colony was selected and moved to 2 ml LB liquid media with antibiotics and shaken overnight at 28°C. After confirmation by bacterial PCR, 1 ml of Agrobacterium liquid was transferred into 100 ml of LB liquid media containing KAN (50 μg/ml), RIF (30 μg/ml), 10mM MES and 25uM acetosyringone and grown overnight at 28°C incubator. The next day, the Agrobacterium culture was centrifuged for 15 min at 4000 rpm. The supernatant was

discarded and the pellet was resuspended in infection buffer containing 10 mM MgCl2,

10 mM 2-(*N*-morpholino) ethanesulfonic acid and 250 µM acetosyringone to an OD_{600}

of 0.6. The *Agrobacterium* cells were then incubated at room temperature for 3 hours

in darkness. After approximately 3 hours, the TRV1 and TRV2 Agrobacterium were

mixed in a 1:1 ratio and prepared for injection into the tomato plants.

Three-week-old tomato seedlings with about 4 true leaves were pruned, removing

the main shoot and all the axillary shoots. This produced wounding sites on the stem

and axillary sites, only leaving two true leaves and two cotyledons. TRV1 and TRV2

Agrobacterium mix was injected near the wounding site using 1 ml syringes. Successful

injection was confirmed by observing the *Agrobacterium* buffer seeping from the

wounding site. Plants were then placed in dark and high-humidity (80%) growth

chamber at 25°C for 2 days, and then grown at 20°C for 2 days. Next, plants were grown

in a greenhouse at 16/8-hour day/night cycle at 24/21°C. We removed any new shoots

that appeared at the wounding sites in the first month to promote the *de novo* shoot

initiation. One month later, all the new shoots were genotyped.

Edited shoots screening and mutation frequency analysis

Approximately 35 DPI, 3 random green leaves from each shoot were harvested for

extracting genomic DNA by using Qiagen DNeasy Plant Mini Kit (Qiagen, 69106)

according to manufacturer instructions. Target sites were amplified using Q5 DNA

polymerase (NEB, M0491s) and PCR products were directly sent for Sanger

sequencing. If edited was observed, plants were transferred to larger pots. Otherwise,

the unedited shoots were cut at their base to promote new shoot initiation.

Deep next-generation amplicon sequencing was performed for analysis of the

mutation frequency of edited plants. DNA libraries were preparted by using a 2-step

PCR amplification method. For the first round of PCR, target regions were amplified

with primers close to the target site using Phusion™ High-Fidelity DNA Polymerases

(Thermo Fisher, F530S) and the reaction was performed under the following cycling

conditions: 98 °C for 30s, 98 °C for 30 s, 55°C for 20 s, 72 °C for 30s, 25 cycles (Step2);

72 °C for 5 min. Then, the PCR products were purified by using 1.0× Ampure XP bead purification (Beckman Coulter, A63881). Purified PCR products were then used as the template for the second round PCR by using Illumina indexing primers under the following cycling conditions: 98 °C for 30s, 98 °C for 30s, 60°C for 20s, 72 °C for 30s, 12 cycles (Step2); 72 °C for 5 min. Subsequently, the second-round PCR products were purified and quantified. Finally, equal amounts of the second-round PCR products were mixed and single-end next-generation sequencing on the Illumina NovaSeqX platform was performed.

Next-generation amplicon sequencing analysis

Amplicon sequencing analysis was conducted as previously described[35]. Single-end reads were processed by adapter trimming with Trim Galore using default parameters, and the resulting reads were aligned to the target genomic region with BWA (v0.7.17) employing the BWA-MEM algorithm. The resulting BAM files were sorted and indexed, and subsequently analyzed with the CrispRvariants R package (v1.14.0). For each sample, distinct mutation patterns and their associated read counts were extracted using CrispRvariants. Based on the assessment of control samples, a stringent classification criterion was applied; only reads containing indels of ≥1 bp (insertions or deletions of identical size starting at the same position) with a minimum of 10 supporting reads per sample were considered edited. Single-nucleotide variants were excluded from the analysis.

Progeny genotyping

Seeds were harvested from the editing parent plants and dried in a 37°C incubator for 3 days. Seeds were sterilized using a 55°C water incubator for 10mins, 75% ethanol for 1 min, and washed three times with sterile water, 3% NaClO for 7 mins. Seeds were then put in 1/2 MS plates and incubated in darkness at 23°C for 3 days, followed by growth with long-day conditions (16-h light and 8-h dark; day 25°C/night 20°C; light intensity:150 μmol/m^2/s; 50% humidity) for about 5 days. Once the cotyledons were

fully expanded, half of a cotyledon was collected for DNA extraction using the Invitrogen Platinum Direct PCR Universal Master Mix (A44647500) following the manufacturer's protocol. The extracted DNA was subsequently used as a template for PCR, and the PCR products were subjected to Sanger sequencing. Quantification of indel frequencies was performed with the ICE CRISPR Analysis Tool.

Genotypes were classified according to indel frequencies as follows: Shoots or seedlings with a total indel frequency of 0–10% were considered wild type (WT); samples with a single unique sequence modification at 35–65%, the nonmodified sequence frequency between 35% and 65%, and the total frequency of single unique sequence modification and nonmodified sequence between 85%~100% was designated as heterozygous (Het); samples with a single unique sequence modification at 85–100% were classified as homozygous (Ho); and samples with two distinct sequence modifications each at 35–65% and the total modification frequency at 85–100% were designated as biallelic for two different alleles.

RT-PCR

Total RNA from progeny seedlings was extracted using Zymo Research Direct-zol RNA MiniPrep kit (R2052). Total RNA was reverse transcribed to synthesize first-strand cDNA using the Invitrogen SuperScript IV VILO Master Mix (11766050). The synthesized cDNA was diluted and then used for PCR with specific primers to detect the TRV virus and TnpB expression by using New England Biolabs Q5 High-Fidelity 2× Master Mix (M0492L) according to manufacturer instructions. The *SlActin* gene was used as the internal control. PCR products were fractionated by 2% agarose gel electrophoresis.

Author contributions: Y.L. and S.E.J. designed research. Y.L., J.L., J.P., S.Y.C.C., E.R., J.A. and S.H.F. performed experiments; Y.L., T.W. and M.K. analyzed data; Y.L. and S.E.J. wrote the paper; S.E.J. and T.W. reviewed and edited the manuscript.

The authors declare no competing interest.

Acknowledgments: We thank all members of the Jacobsen Lab for their helpful insights and valuable suggestions. We also thank Mahnaz Akhavan for support with Next generation sequencing at the UCLA Broad Stem Cell Research Center BioSequencing Core Facility. We are grateful for helpful discussion from members of Jennifer A. Doudna lab (University of California, Berkeley). This work was supported by an NSF Plant Genome Research Program grant (2334027). Steven E. Jacobsen is an investigator of the Howard Hughes Medical Institute.

References

1.   Tuncel, A., Pan, C., Clem, J.S., Liu, D. & Qi, Y. CRISPR–Cas applications in agriculture and plant research. *Nature Reviews Molecular Cell Biology* **26**, 419–441 (2025).

2.   Li, B., Sun, C., Li, J. & Gao, C. Targeted genome-modification tools and their advanced applications in crop breeding. *Nature Reviews Genetics* **25**, 603–622 (2024).

3.   Altpeter, F. et al. Advancing Crop Transformation in the Era of Genome Editing. *Plant Cell* **28**, 1510–1520 (2016).

4.   Nasti, R.A. & Voytas, D.F. Attaining the promise of plant gene editing at scale. *Proc Natl Acad Sci U S A* **118** (2021).

5.   Ali, Z. et al. Efficient Virus-Mediated Genome Editing in Plants Using the CRISPR/Cas9 System. *Mol Plant* **8**, 1288–1291 (2015).

6.   Liu, Q., Zhao, C., Sun, K., Deng, Y. & Li, Z. Engineered biocontainable RNA virus vectors for non-transgenic genome editing across crop species and genotypes. *Mol Plant* **16**, 616–631 (2023).

7.   Uranga, M. et al. Efficient Cas9 multiplex editing using unspaced sgRNA arrays engineering in a Potato virus X vector. *Plant J* **106**, 555–565 (2021).

8.   Lee, S.Y. et al. Development of virus-induced genome editing methods in Solanaceous crops. *Hortic Res* **11**, uhad233 (2024).

9.   Nagalakshmi, U., Meier, N., Liu, J.Y., Voytas, D.F. & Dinesh-Kumar, S.P. High-efficiency multiplex biallelic heritable editing in Arabidopsis using an RNA virus. *Plant Physiol* **189**, 1241–1245 (2022).

10.   Li, T. et al. Highly efficient heritable genome editing in wheat using an RNA virus and bypassing tissue culture. *Mol Plant* **14**, 1787–1798 (2021).

11.   Tamilselvan-Nattar-Amutha, S. et al. Barley stripe mosaic virus-mediated somatic and heritable gene editing in barley (Hordeum vulgare L.). *Front Plant Sci* **14**, 1201446 (2023).

12.   Ellison, E.E. et al. Multiplexed heritable gene editing using RNA viruses and mobile single guide RNAs. *Nat Plants* **6**, 620–624 (2020).

13.   Bradamante, G., Mittelsten Scheid, O. & Incarbone, M. Under siege: virus control in plant

meristems and progeny. *Plant Cell* **33**, 2523–2537 (2021).

14.   Wu, H. et al. WUSCHEL triggers innate antiviral immunity in plant stem cells. *Science* **370**, 227–
231 (2020).

15.   Steinberger, A.R. & Voytas, D.F. Virus-induced gene editing free from tissue culture. *Nature*
Plants **11**, 1241–1251 (2025).

16.   Xiang, G. et al. Evolutionary mining and functional characterization of TnpB nucleases identify
efficient miniature genome editors. *Nature Biotechnology* **42**, 745–757 (2024).

17.   Weiss, T. et al. Viral delivery of an RNA-guided genome editor for transgene-free germline
editing in Arabidopsis. *Nature Plants* **11**, 967–976 (2025).

18.   Yoshida, T., Ishikawa, M., Toki, S. & Ishibashi, K. Heritable Tissue-Culture-Free Gene Editing in
Nicotiana benthamiana through Viral Delivery of SpCas9 and sgRNA. *Plant and Cell Physiology*
65, 1743–1750 (2024).

19.   Qiao, J.-H. et al. Transgene- and tissue culture-free heritable genome editing using RNA virus-
based delivery in wheat. *Nature Plants* **11**, 1252–1259 (2025).

20.   Lee, H., Baik, J.E. & Kim, K.-N. Development of an efficient and heritable virus-induced genome
editing system in Solanum lycopersicum. *Horticulture Research* **12** (2024).

21.   Liu, D. et al. Heritable gene editing in tomato through viral delivery of isopentenyl transferase
and single-guide RNAs to latent axillary meristematic cells. *Proceedings of the National*
Academy of Sciences **121**, e2406486121 (2024).

22.   Sebiani-Calvo, A., Hernández-Soto, A., Hensel, G. & Gatica-Arias, A. Crop genome editing
through tissue-culture-independent transformation methods. *Frontiers in Genome Editing*
Volume 6 - 2024 (2024).

23.   Bélanger, J.G., Copley, T.R., Hoyos-Villegas, V., Charron, J.-B. & O'Donoughue, L. A
comprehensive review of in planta stable transformation strategies. *Plant Methods* **20**, 79
(2024).

24.   Kang, G.H., Ko, Y. & Lee, J.M. Enhancing virus-mediated genome editing for cultivated tomato
through low temperature. *Plant Cell Rep* **44**, 22 (2025).

25.   Shan, Q. et al. Targeted genome modification of crop plants using a CRISPR-Cas system. *Nature*
Biotechnology **31**, 686–688 (2013).

26.   Kang, B.-C. et al. Precision genome engineering through adenine base editing in plants. *Nature*
Plants **4**, 427–431 (2018).

27.   Zhong, Z. et al. Efficient plant genome engineering using a probiotic sourced CRISPR-Cas9
system. *Nature Communications* **14**, 6102 (2023).

28.   Li, Y., Zheng, L., Corke, F., Smith, C. & Bevan, M.W. Control of final seed and organ size by the
DA1 gene family in Arabidopsis thaliana. *Genes Dev* **22**, 1331–1336 (2008).

29.   Liu, H. et al. TaDA1, a conserved negative regulator of kernel size, has an additive effect with
TaGW2 in common wheat (Triticum aestivum L.). *Plant Biotechnology Journal* **18**, 1330–1342
(2020).

30.   Wang, J.-L. et al. Down-regulation of BnDA1, whose gene locus is associated with the seeds
weight, improves the seeds weight and organ size in Brassica napus. *Plant Biotechnology*
Journal **15**, 1024–1033 (2017).

31.   Li, C. et al. OsDA1 positively regulates grain width in rice. *The Crop Journal* **12**, 92–101 (2024).

32.   Oh, Y. et al. Heritable virus-induced germline editing in tomato. *Plant J* **122**, e70115 (2025).

33.   Macfarlane, S.A. Tobraviruses--plant pathogens and tools for biotechnology. *Mol Plant Pathol*

11, 577–583 (2010).

34.   Ali, Z., Eid, A., Ali, S. & Mahfouz, M.M. Pea early-browning virus-mediated genome editing via

432   the CRISPR/Cas9 system in Nicotiana benthamiana and Arabidopsis. *Virus Res* **244**, 333–337

433   (2018).

434   35.   Li, Z. et al. Genome editing in plants using the compact editor CasΦ. *Proc Natl Acad Sci U S A*

435   **120**, e2216822120 (2023).

Fig. 1. Virus-induced genome editing in *de novo shoots* of tomato using TRV-TnpB-gRNA^SlPDS -mSlFT. (**A**) Schematic diagram of the TRV1 and TRV2 plasmids. The tomato mutated *FT* mobility element (mSlFT) was fused to the 3' end of the gRNA in TRV2. RdRP, RNA dependent RNA polymerase; MP, movement protein; CP, coat protein; sgP, *Pea Early Browning Virus* subgenomic promoter; HR, HDV ribozyme, mSlFT; mutated tomato *FT* mobility element. (**B**) Schematic representation of *SlPDS* genomic structure. The light blue box represents exons, light grey represents the UTR sequence, and dark grey represents introns. The TAM sequence and target sequence are

highlighted by the pink and blue lines, respectively. (**C**) The procedure for TRV-TnpB-gRNA-mSlFT mediated genome editing system in tomato plants. Edited *de novo* shoots were screened and validated by Sanger and NGS amplicon sequencing. (**D-H**) Different growth stages of TRV-TnpB-gRNA^SlPDS#3 plant displaying albino sectors in leaves and flowers. White leaflets in leaves are indicated by blue arrows, and the flowers exhibited white petals (**H**). **G** is a close-up view of **F**; DAI: Days After Injection. (Scale bars, 1 cm). (**I**) Sanger sequencing of the *SlPDS* gene target site in TRV-Control (top) and TRV-TnpB-gRNA^SlPDS#3 (bottom) plants. (**J**) Indel frequency analysis of the *SlPDS* gene in the TRV-TnpB-gRNA^SlPDS#3 shoot. Genomic DNA was extracted from green leaf tissue and subjected to amplicon sequencing using next-generation sequencing (NGS). The TAM sequence and target sequence are highlighted by the pink and blue lines, respectively. Deleted residues are marked by black dotted lines. The different indel types are listed on the left. The percentage of reads corresponding to each indel type is shown on the right. The indel efficiency was calculated by the total indel reads number divided by total reads number. (**K-M**) Phenotype of the TRV-TnpB-gRNA^SlPDS#7 (**K**), #17 (**L**) and #43 (**M**) shoots, respectively., showing no obvious albino sectors. Images include the whole plant (left), a leaf (top right) and flowers (bottom right). (Scale bars, 1 cm). (**N-P**) Indel frequency analysis of the *SlPDS* gene from the TRV-TnpB-gRNA^SlPDS#7 (**N**), #17 (**O**) and #43 (**P**) shoots. The reads of sequences with frequencies >5% are shown as well as the total indel frequency.

Fig. 2. Transgene-free and heritable genome editing in TRV-TnpB-gRNA^{SlPDS}-mSlFT infected tomato plants. (**A-D**) Phenotype of 1-week old progeny seedlings from TRV-TnpB-gRNA^{SlPDS}-mSlFT#3 (**A**), #7 (**B**), #17 (**C**) and #43 (**D**) parent plants. The albino seedlings are marked with yellow arrows. (Scale bars, 2 cm). (**E-H**) Phenotyping and genotyping analysis of progeny seedings from TRV-TnpB-gRNA^{SlPDS}-mSlFT#3 (**E**), #7 (**F**), #17 (**G**) and #43 (**H**) parent plants. Representative image of 1-week old progeny seedlings (top panel). Sanger sequencing trace file of PCR product from progeny seedlings (bottom panel). Ref: Reference sequence; WT: Wild Type; Het: Heterozygous; Ho: Homozygous. (Scale bars: 1 cm). (**I**) Table summarizing the percentage of progeny seedlings with heterozygous and homozygous mutations from the four parent plants shown in panels A-H. (**J**) RT-PCR performed using total RNA from 2 control plants, 2 TRV injected plants (tissue shortly after injection), and 10 albino progeny seedlings to detect *TnpB* (upper panel) or TRV mRNA (middle panel) expression. The *SlActin* gene served as an endogenous control (bottom panel). The experiment was repeated three times with similar results.

Fig.3. Somatic editing in tomato using TRV to deliver TnpB and guide RNA targeting *SlDA1*. (**A**) Phylogenetic analysis of tomato DA1 (SlDA1) and its homologs from other species. The SlDA1 is marked with cyan color. (**B**) Schematic diagram of genomic structure. The cyan box represents exons, light grey represents the UTR sequence and dark grey represents introns, purple box represents the gRNA site. The TAM and target sequences are highlighted by the pink and blue lines, respectively. (**C**) Indel frequency analysis of the *SlDA1* gene from the TRV-TnpB-gRNA[SlDA1] #3, #23, #32 and #38 shoots. The mutant sequences with frequencies >5% are shown as well as the total indel frequency.