Developing a Molecular Toolkit to ENABLE all to apply CRISPR/Cas9-based Gene Editing *in planta*

Birhan Addisie Abate[1,2#], Florian Hahn[3#], Daniele Chirivì[4,5], Camilla Betti[4], Fabio Fornara[4], Jennifer C Molloy[1], and Kate M Creasey Krainer[4]

[1]Biomanufacturing Group, International Centre for Genetic Engineering and Biotechnology, Trieste, Italy
[2]Plant Biotechnology Directorate, Bio and Emerging Technology Institute (BETin), Addis Ababa, Ethiopia
[3]Grow More Foundation, New York, NY, United States
[4]Department of Biosciences, University of Milan, Milan, Italy
[5]Present address: Centre for Research in Agricultural Genomics (CRAG), Barcelona, Spain

[#]These authors contributed equally as first author
Corresponding author: Kate M Creasey Krainer

ABSTRACT

Gene editing, especially the CRISPR/Cas9 (<u>C</u>lustered <u>R</u>egulatory <u>I</u>nterspaced <u>S</u>hort <u>P</u>alindromic <u>R</u>epeats/<u>C</u>RISPR <u>as</u>sociated protein <u>9</u>) system, has revolutionized trait development in crops. However, large parts of the world are missing out on applying CRISPR *in planta*. There is an obvious lack of gene editing applications in locally relevant crops in the Global South which tend to be neglected by mainstream agricultural research and development. Access barriers to these new breeding technologies need to be removed to allow the potential impact of these technologies on food security to happen. Here, we present the ENABLE® Gene Editing *in planta* toolkit, a minimal molecular toolbox allowing users to create a CRISPR knockout vector for transient or stable plant transformation in two simple cloning steps. We validate the toolkit in rice (*Oryza sativa*) protoplasts and in *Arabidopsis thaliana* plants. The ENABLE® kit is designed to be utilized specifically by users in the Global South who are new to CRISPR technology by providing a simple workflow, extensive accompanying protocols as well as options for low cost methods for cloning verification and gene editing verification *in planta*. We hope that our toolkit helps bridge the gap between the recent biotechnological advancements in plant breeding that high income countries can access and the lack of those technologies in low and middle income countries.

Keywords: CRISPR, ENABLE, gene editing, crops, plants

INTRODUCTION

Around 2.3 billion people, mostly in the Global South, faced moderate to severe food insecurity in 2024. While the situation in Asia and Latin America improved in recent times, the issue is still worsening in Africa (FAO, IFAD, UNICEF, WFP and WHO, 2025). On top of that, climate change is likely to increase food insecurity around the globe with countries in the Global South being disproportionately affected by it (Onyeaka et al., 2024). Investments in R&D are needed to bring resilience into the food producing sector (FAO, IFAD, UNICEF, WFP and WHO, 2025). Specifically, plant breeding can make agriculture more resilient to the consequences of erratic environmental conditions, abiotic and biotic stress factors, as well as improve the nutritional quality of crops (Huang et al., 2002; Qaim, 2020). The impact of improvements in plant varieties on food security has been shown in the Green Revolution starting in the 1960s, where newly developed cultivars (as well as fertilizer input regimes) helped boost crop production and calorie intake in many countries. However, the impact of the Green Revolution was diminished in some parts of the world, such as sub-Saharan Africa, due to its primary focus on rice and wheat and neglection of locally relevant, underutilized crops (Evenson and Gollin, 2003; Tadele, 2019). Hence, we need a second Green Revolution to ensure food security in low and middle income countries (LMIC), with a strong focus on local varieties, which will be essential for success. For this, we need to ensure that we make use of all available breeding technologies as no single silver bullet exists to solve those issues.

Unfortunately, many plant scientists from countries with the highest share of food insecurity have problems with accessing and making use of new technologies in the same way as it is done e.g. in the Global North or China. Africa for example has a lower percentage of scientists in the population and accordingly low numbers of publications and patents, partially due to lack of adequate STEM education (UN, 2024). Outdated or missing regulatory frameworks also delay application of modern breeding technologies (Qaim, 2020). On top of that, geopolitical issues, financial hurdles and VISA issues can exclude scientists from international collaborations and knowledge transfer (Marks et al., 2023). Practical issues like access to reagents and lack of cold chain delivery options additionally hinder uptake of new technologies. However, it is crucial that plant scientists from disadvantaged countries in the Global

South, particularly in Africa, are able to work on the traits and crops that they deem relevant as they have the best understanding of the plants, the issues that local farmers face, as well as consumers' needs.

One of the technologies that has become highly relevant in the last decade as a new player in the crop breeding toolbox is the CRISPR/Cas9 technology. The CRISPR/Cas9 technology is based on a bacterial defence system against phages and was developed into a gene editing tool in 2012 (Jinek et al., 2012). It is a simple two component system, consisting of the DNA cleaving enzyme Cas9 from *Streptococcus pyogenes* (Sp) as well as an artificial programmable single guide RNA (sgRNA) molecule that guides the Cas9 endonuclease to a 20 bp target site of choice in the genome through base pairing. By artificially introducing SpCas9 and a sgRNA into a crop of interest, DNA breaks are introduced into the crop genome at the target site. In plants and other eukaryotes, those breaks are then usually repaired via the cell's error-prone non homologous end joining pathway, which can randomly introduce insertions or deletions into the target site (Pacher and Puchta, 2017). Due to its simplicity and efficiency, the CRISPR/Cas9 technology quickly became the method of choice for trait development in many organisms. Over the last decade, it has been further adapted for many applications, such as base editing, gene insertion or gene activation/repression (Liu et al., 2022; Wang and Doudna, 2023). Despite the variety of CRISPR use cases and nuclease variants available these days, most users still employ the SpCas9 nuclease, primarily for introduction of gene knockouts.

CRISPR/Cas9 has proven to be highly impactful in plant sciences, partially also due to relaxed regulatory hurdles compared to transgenic modifications in many parts of the world (see Gene Editing Regulatory Tracker by Genetic Literacy Project: https://crispr-gene-editing-regs-tracker.geneticliteracyproject.org/#jet-tabs-control-1401 [retrieved: 11.08.2025]) as well as the higher availability of genome sequence resources these days. Traits are developed around resistance to biotic and abiotic stresses, yield improvement, biofortification or shelf life extension in a variety of crops and with some improved varieties having been cleared for market entrance already (Ahmar et al., 2024; Chen et al., 2024; Liu et al., 2021; Polidoros et al., 2024). Its potential impact on the Global South, e.g. in Africa has been recognised (Amoah et al., 2024; Missanga and Mmbando, 2025; Ndudzo et al., 2024). However, while first

use cases in locally relevant crops such as yam (Syombua et al., 2021), foxtail millet (Liang et al., 2022), sorghum (Hao et al., 2023), pigeonpea (Senthil et al., 2025) and cassava (Gomez et al., 2019; Mukami et al., 2024) have been described, the overall number of CRISPR studies in underutilized crops and particularly those performed from research groups in sub-Saharan Africa remains low. No CRISPR studies have been published to our knowledge in locally relevant crops such as African eggplant, Bambara groundnut, mungbean, hyacinth bean, and white fonio (Venezia and Creasey Krainer, 2021, EU SAGE database: https://www.eu-sage.eu/genome-search [retrieved: 11.08.2025]). Plant scientists in LMIC need improved access to the CRISPR/Cas9 technology to benefit from its potential impact on (underutilized) crop improvement.

The **Grow More Foundation** (www.growmorefoundation.org) is a non-profit NGO, dedicated to equitable access to plant science and technology to support food security and sustainability in LMIC. With its ENABLE® program, the foundation aims to translate plant science technologies to underutilized staple crops by providing the resources and community to enable plant scientists in the Global South to solve agricultural problems that are relevant in their countries. In this study, we developed the **ENABLE® gene editing *in planta* toolkit**, a simple molecular toolkit consisting of eight CRISPR related modules, as well as extensive corresponding documentation. Our toolkit allows for low cost methods of cloning verification as well as editing verification. With this toolkit, we want to enable users at every stage of gene editing proficiency to generate CRISPR vectors for *in planta* usage. We prove the efficiency of our toolkit in the model plants Arabidopsis and rice and make it accessible to the academic community via the public repository Addgene (www.addgene.org, Addgene ID 1000000270). Our vision is for ENABLE® kits to be incorporated in training and research programs to empower the next generation of plant scientists to develop new traits in relevant local crop varieties.

METHODS

Vectors of the *pro bono* ENABLE Kit

The following Golden Gate modules form part of the *pro bono* ENABLE kit: The vectors pGMF1-M (Addgene ID: 242203) and pGMF2-M (Addgene ID: 242204) for expression of two sgRNAs in monocots using the rice *U6-2* promoter, and pGMF6 (Addgene ID: 242208) expressing enhanced-nuclear localized GFP, were synthesized by Integrated DNA Technologies, Inc. (Coralville, USA). pGMF1-D (Addgene ID: 242205) and pGFM2-D (Addgene ID: 242206), for expression of two sgRNAs in dicots using the Arabidopsis *U6-26* promoter, were synthesized by Vectorbuilder (Chicago, USA). pGMF3 (Addgene ID: 242207) provides the T-DNA binary backbone vector and was adapted from the vector pICSL4723 (Castel et al., 2019, pICSL4723 was gifted by Jonathan Jones, the Sainsbury Laboratory). pGMF4 (Addgene ID: 128183; also designated CasC in this study and previously published as pFH54; gifted by Vladimir Nekrasov, Rothamsted Research; Hahn et al., 2020) functions as *CAS9* expression vector, and pGMF5 (previously published as pICSL11059; Hygromycin resistance cassette, Lawrenson et al., 2015; Addgene ID: 68263) by Nicola Patron (University of Cambridge). All those vectors can be ordered as a combined kit from Addgene (Addgene ID: 1000000270).

In addition to these vectors, the following Golden Gate modules were tested for their gene editing efficiency in this study: CasA, previously published as pFH52 (Hahn et al., 2020; Addgene ID: 128181, gifted by Vladimir Nekrasov), CasB, previously published as pFH66 (Hahn et al., 2020; Addgene ID: 131765, gifted by Vladimir Nekrasov), CasD, previously published as Level 1 P2 OsUbiP:Cas9:NosT (Smedley et al., 2021; Addgene ID: 165424, gifted by Wendy Harwood).

All sequences can be found in Suppl. File 1.

sgRNA sequences

The following target sites were used to create mutations in the Arabidopsis and rice genome:

Species	Gene name	Gene ID	sgRNA ID	5'-3' sequence
Oryza sativa (cv.	*Phytoene Desaturase 1*	LOC_Os03g08 570	*OsPDS1* sgRNA1	GTGGAGGTC TTGGAAAGTC

Nipponbare)				C
			OsPDS1 sgRNA2	CCAGCAATC ACGACCTGT AA
Arabidopsis thaliana (Col-0)	*Glabrous 1*	At3g27920	*AtGL1* sgRNA1	GGAAAAGTT GTAGACTGA GA
			AtGL1 sgRNA2	TTGAGCCCTA ATGTGAACAA

AtGL1 sgRNA1 target site was first described in Hahn et al. (2017). Other sgRNA target sites were designed with the help of CRISPOR online target design tool (Concordet and Haeussler, 2018) following guidelines summarized in Suppl. File 2.

DNA construct assembly

To test CRISPR editing efficiencies of the Golden Gate modules described above, they were combined to binary T-DNA constructs using MoClo Golden Gate DNA assembly (Engler et al., 2009; Weber et al., 2011). The Golden Gate cut-ligation reactions were performed according to the described protocol in Suppl. File 2 and 3. In summary, for each experiment, two annealed primer pairs (Suppl. File 3, Eurofins Genomics) encoding two sgRNA target sequences for each gene were subcloned into pGMF1-D and pGMF2-D, or pGMF1-M and pGFM2-M respectively by Golden Gate cut-ligation reactions using BsaI-HFv2® and T4 DNA Ligase (New England Biolabs). To generate constructs for stable Arabidopsis plant transformation, the resulting two vectors were combined with a vector mix of pGMF3, a *CAS9* expression vector, and pGMF5 by Golden Gate cut-ligation reactions using BbsI-HF® and T4 DNA Ligase (New England Biolabs) leading to the final binary vector. To generate constructs for rice protoplast transformation, the resulting two vectors were combined with a vector mix of pGMF3, a *CAS9* expression vector, and pGMF6 by GG cut-ligation reactions using BbsI-HF® and T4 DNA Ligase. All sequences can be found in Suppl. File 1.

All cut-ligation reactions were transformed into high efficiency 5-alpha competent *E. coli* cells (New England Biolabs), cells were plated on suitable selection plates (Suppl. File 2), single white colonies were inoculated and plasmids were isolated from

overnight cultures using the Monarch® plasmid miniprep kit (New England biolabs). All constructs were verified by full plasmid sequencing (Eurofins Genomics).

Rice protoplast isolation and transformation

Rice cv. Nipponbare seeds were dehusked, surface sterilized with ethanol and bleach, and grown in tin foil wrapped Magenta vessels on ½ MS media at 28 °C for 10 days. Rice protoplasts were isolated from 100 seedlings following the protocol by He et al. (2016) with the following change: During the protoplast isolation process, the stem strips in enzyme solution were placed in air tight food boxes (Zwilling Fresh & Save) and vacuum was applied using a manual vacuum pump (Zwilling Fresh & Save) instead of a classic laboratory vacuum pump setup. The isolated protoplasts were split into 9 batches and each batch but one control batch (ctrl) was transformed with 10 μg of one of the eight CRISPR constructs using PEG-mediated transformation. Successful transformation was verified by eGFP analysis 3 days post transformation using a Nikon Eclipse Ti inverted microscope fitted with an Andor Neo sCMOS camera and using a FITC filter for eGFP detection. Microscopy pictures were analysed using NIS-Elements software (Nikon). DNA was isolated from protoplast pellets by resuspending the pellet in 300 μL TPS buffer (100 mM Tris-HCl, pH 8, 10 mM EDTA, 1 M KCl) followed by incubation for 3 min at 50 °C. DNA was precipitated by addition of an equivolume of isopropanol for 20 min. The DNA pellet was washed with 70% ethanol before resuspension in 20 μL TE buffer (10 mM Tris-HCl, pH 7.5, 1 mM EDTA). DNA was used for PCR based methods of editing verification.

Agrobacterium transformation

Constructs designed for stable Arabidopsis transformation were transformed into competent Agrobacteria (strain GV3101) and selected on LB plates containing 50 μg/mL Rifampicin, Gentamicin, Tetracycline and Kanamycin. Presence of constructs in Agrobacteria was verified using colony PCR on single colonies using primers V_79 and V_80 (Suppl. File 3).

Arabidopsis transformation

Arabidopsis thaliana ecotype Col-0 plants were grown in pots on general potting compost in growth chambers (Jumo LPF200) under long day conditions (16/8 h cycles of light (70 μmol m^{-2} sec^{-1}) at 22°C in the day and at 18°C in the night). Plants were

transformed using floral dipping following standard procedures (Clough and Bent, 1998). T0 seeds were sown on ½ MS plates supplemented with 10 mg/L hygromycin, 0.1 % (w/v) sucrose, 0.05 % (w/v) MES, pH 5.8 adjusted with KOH and solidified with 0.8 % (w/v) of plant agar (no hygromycin was added for WT control plants). Plants were grown for 10 days on plates before transfer on soil. Pictures were taken of 3-4 week old plants using a Fujifilm XT200 camera and a Zeiss Discovery V8 stereomicroscope. DNA was isolated from one of the first four leaves of 3-4 week old plants by crushing the leaf in an Eppendorf tube with 300 μL TPS buffer (100 mM Tris-HCl, pH 8, 10 mM EDTA, 1 M KCl) and a metal bead using a mixer mill (Retsch, 30 Hz, 1 min) followed by incubation for 3 min at 50 °C. The solution was transferred into a new tube and DNA was precipitated by addition of an equivolume of isopropanol for 20 min. The DNA pellet was washed with 70% ethanol before resuspension in 50 μL TE buffer (10 mM Tris-HCl, pH 7.5, 1 mM EDTA). T-DNA presence in T0 lines was verified by PCR using DreamTaq Green PCR Mastermix (Thermo Fisher Scientific) and primers V_79/V_80 (Suppl. File 3). DNA of verified T0 transformants was used for PCR based methods of editing verification.

Mutation analysis

For Arabidopsis, the *GL1* target region was amplified by PCR with Q5® DNA Polymerase (New England Biolabs) or DreamTaq Green PCR Mastermix (Thermo Fisher Scientific) using primers GMF133/GMF134 (Suppl. File 3). PCR products were purified using ExoSAP-IT™ PCR Product Cleanup Reagent (Applied Biosystems) and mutation presence was verified on sequence level by Sanger sequencing (Eurofins Genomics, Germany). Sanger chromatograms were analysed for edits using the ICE Sanger Sequencing Analysis tool (https://ice.editco.bio/#/). For rice, the *PDS1* target region was amplified by PCR with Q5® DNA Polymerase (New England Biolabs) using primers GMF147/GMF148 (Suppl. File 3). PCR products were purified using Monarch® PCR & DNA Cleanup Kit and sequenced using Amplicon EZ Next Generation Sequencing (Genewiz, Germany). NGS data was analysed using CRISPResso2 (https://crispresso2.pinellolab.org, Clement et al., 2019).

RESULTS

Development of an accessible toolkit for *in planta* gene editing

We aimed at developing an easy to use molecular toolkit which provides users new to *in planta* CRISPR gene editing the necessary tools for simple and inexpensive gene editing experiments both in monocot and dicot species. While plant gene editing using ribonucleoproteins or RNA only based approaches has been shown to be feasible in some species (Zhang et al., 2021), in most cases, it is still performed using stable transformation of a vector containing three major components: a **CAS9** gene, one or more genes for **sgRNA** expression and a **selection marker** gene for successful transformation. Also, DNA is more stable than RNA or protein when considering shipping of CRISPR components around the world. Hence, we decided to use vector based provision of CRISPR components for our toolkit. To be able to test different components for our vector system and to allow the end user a degree of flexibility regarding the selection marker, or to adapt the toolkit further to their own needs, we opted for a modular vector system, where multigene plasmids can be assembled in one-pot reactions using Golden Gate cloning. Golden Gate cloning has become the method of choice for large multigene assembly and uses libraries of defined basic modules with unique restriction overhangs, which are then seamlessly assembled into a receptor vector in a defined linear order using type IIS restriction enzymes (Engler et al., 2009; Weber et al., 2011). The method is characterized by its high efficiency and simplicity compared to earlier cloning procedures. In our ENABLE® toolbox, we aimed to provide all basic CRISPR modules (sgRNA modules, *CAS9* module, selection marker module) on separate level 1 (L1) vectors, which can be easily combined into the level 2 (L2) backbone (pGMF3) in a Golden Gate one-pot restriction ligation reaction to create the final CRISPR vector for *in planta* gene editing (Fig. 1a, 1b, Suppl. File 2).

We aimed to provide the following parts in our toolkit: **pGMF1** and **pGFM2** encoding each an optimized sgRNA scaffold (Dang et al., 2015) under the control of a Polymerase III (Pol III) promoter, which allow users to subclone their 20 base pair target sequences seamlessly upstream of the sgRNA scaffold via a BsaI-guided restriction ligation reaction (Fig. 1b, Suppl. File 2). We opted to provide two sgRNA genes in our ENABLE® toolkit to increase chances of successful gene editing,

allowing targeting of two independent genes or creating targeted larger deletions between two sgRNA sites. Both vectors use a IPTG/X-Gal based blue/white screening system for low cost assessment of successful target sequence subcloning (Suppl. Fig. 1a, Suppl. File 2). **pGMF3** is the binary T-DNA backbone for the final CRISPR construct, as it was derived from a frequently used vector backbone for gene editing experiments in a variety of plants (Castel et al., 2019; Dai et al., 2020; Luginbuehl et al., 2020; Jethva et al., 2024) and due to its red/white selection system to screen for successfully assembled CRISPR constructs in *E. coli* (Suppl Fig. 1a, Suppl. File 2). This backbone can be used for Agrobacterium mediated plant transformation, as well as protoplast transfection or particle bombardment. **pGMF4** encodes a *CAS9* gene under a strong, constitutive promoter. **pGMF5** and **pGMF6** encode selection markers for stable (pGMF5: Hygromycin resistance, Lawrenson et al., 2015) and transient (pGMF6: nuclear localised eGFP) plant transformation, allowing the user to choose which of the two cassettes to include in their final CRISPR vector based on their downstream application.

Figure 1: Concept of the ENABLE® Gene Editing *in planta* toolkit. (a) The major components envisioned for the ENABLE gene editing toolkit to allow the user to clone a CRISPR vector for transient or stable plant transformation. (b) The 2-step Golden Gate cloning strategy involves subcloning of two CRISPR target sequences into the two vectors pGMF1 and pGMF2 via annealed oligos encoding the 20 bp target site, followed by assembly of the final T-DNA binary vector. (c) Four different *CAS9* cassettes with different promoters, nuclear localization signals (NLS), tags, codon optimizations (Ara = Arabidopsis, wheat or human codon optimized) and terminators were tested in this study. (d) sgRNA expression via an Arabidopsis or rice polymerase III promoter was tested in this study.

CAS9 variants and sgRNA expression cassettes

Where possible, we aimed at using publicly available basic modules for our ENABLE®
toolkit, to ensure that the parts have been validated independently. It has been shown
previously that there can be a high variability in gene editing efficiencies when using
different *CAS9* expression cassettes on the same target gene. These differences in
editing activity could be caused by the choice of promoters and terminators, but also
by *CAS9*-intrinsic differences, such as codon optimization, presence/absence of
protein tags, the choice of nuclear localisation signal (NLS) and intronization (Mikami
et al., 2015; Castel et al., 2019; Raitskin et al., 2019; Hahn et al., 2020; Grützner et
al., 2021). To provide a *CAS9* module which ensures high editing efficiency in both
monocots and dicots, we aimed at comparing four publicly available *CAS9* modules
which were all compatible with the Golden Gate cloning system, but differed in regards
to promoter/terminator choice, codon optimization, and choice of NLS (labelled CasA-
D in this study, Fig. 1C, Suppl. File 1, Hahn et al., 2020; Smedley et al., 2021).

For sgRNA expression, the main issue when designing a simple toolkit which should
cater to as many plant species as possible is the choice of a Pol III promoter which
works in monocots and dicots. While ideally one would use a Pol III promoter derived
from the plant species one is working on, this is not feasible for a "one solution fits all"
toolkit. Hence, we decided to design and test sgRNA modules where expression was
either driven by the rice *OsU6-2* promoter, or the Arabidopsis *AtU6-26* promoter (Fig.
1D), as both have been used extensively across monocot and dicot species,
respectively (Feng et al., 2013; Fauser et al., 2014; Mikami et al., 2015; Lawrenson
et al., 2015; Zhou et al., 2018; Li et al., 2019; Abe et al., 2019; Nie et al., 2022) and
have been shown to express across both groups of angiosperms (Connelly et al.,
1994; Lin et al., 2018).

Cloning of binary CRISPR constructs

To find the best working combination of parts and to verify the straightforward
assembly of our vector system, we decided to test all possible combinations of the four
CAS9 cassettes and two sgRNA setups on the *PHYTOENE DESATURASE 1* (*PDS1*)
gene in *Oryza sativa* cv. Nipponbare protoplasts (as a monocot model) and the
GLABROUS 1 (*GL1*) gene in stably transformed *Arabidopsis thaliana* (as a dicot
model), both of which have been frequently used in the past as CRISPR targets due

to their distinct phenotype in case of a gene knockout (Zhang et al., 2014; Hahn et al., 2017; Banakar et al., 2020; Kong et al., 2023). For this, we cloned 8 Level 1 vectors containing the various sgRNA-promoter combinations. Those were then used to clone 16 binary L2 CRISPR vectors (8 per target gene) containing two sgRNAs targeting either *AtGL1* or *OsPDS1* under the control of *OsU6-2* or *AtU6-26* promoters, one of the four *CAS9* cassettes and a selection marker (Hygromycin resistance for Arabidopsis experiments, eGFP for rice experiments, Fig. 2a, Fig. 3a). For selected constructs, we verified cloning efficiencies based on blue/white screening for L1 constructs and red/white screening for L2 constructs. Using the protocol described (Suppl. File 2, Suppl. File 3), we achieved in most cases white colony rates of > 90% (Suppl. Fig. 1b), which underlines the efficiencies of the MoClo Golden Gate syntax and the associated cloning protocol.

Toolkit testing in transiently transformed rice protoplasts

Rice cv. Nipponbare protoplasts were transformed with eight different CRISPR vectors containing different combinations of Pol III promoters for sgRNA expression and *CAS9* cassettes. In all cases, the vectors contained two sgRNAs targeting the second exon of the *PDS1* gene (Fig. 2b) and an eGFP selection marker. Fluorescence microscopy of protoplasts after 3 days showed nuclear localized eGFP expression (Fig. 2c) and proved successful transformation at efficiencies of ~ 80% across all samples (Suppl. Fig. 2). We amplified the targeted genomic region via PCR in all samples to check for editing activity. While all samples including the untransformed control sample produced a main amplicon with the expected length of 497 bp, we could detect a smaller, additional amplicon in all samples transformed with CRISPR vectors containing sgRNAs expressed by the *OsU6-2* promoter. The size of the smaller amplicon corresponds to alleles containing a deletion between both sgRNA targets (97 bp distance). The additional bands were not visible in the control sample or samples where sgRNAs were expressed using the *AtU6-26* promoter (Fig. 2d). We also submitted the PCR amplicons of the target region of all samples to Amplicon NGS (Fig. 2e, Suppl. File 4). Constructs where sgRNAs were expressed by the *AtU6-26p* showed very low frequencies of mutated reads between 1 and 4.8% depending on the *CAS9* cassette. In contrast, expressing the same sgRNAs with the *OsU6-2* promoter boosted the frequencies of mutated reads to at least 13.2 % (CasA) and up to 35.3 % (CasC).

Figure 2: Testing of the ENABLE® Gene Editing *in planta* toolkit in rice protoplasts. (a) Eight different CRISPR vectors were assembled to target the *OsPDS1* gene with each containing a unique combination of *CAS9* cassette (CasA – CasD) and sgRNA expression cassette (sgRNA expression via *OsU6-2* promoter or *AtU6-26* promoter). (b) The *OsPDS1* gene. Black boxes indicate exons. The position of the two sgRNA target sites in Exon 2 is annotated. The PAM sequence is underlined. (c) Exemplary close-up from untransformed rice protoplasts (top) and a protoplast transformed with one of the eight CRISPR vectors containing an *eGFP* gene (CasA_OsU6-2p::OsPDS1_sgRNAs). (d) *OsPDS1* PCR amplicon analysis of protoplast cultures transformed with different CRISPR constructs. Expected product size in non-edited rice cells: 498 bp, distance between sgRNA1 and sgRNA2 cut site: 97 bp. M = Marker. (e) NGS analysis reveals high proportions of modified reads in PCR amplicons derived from protoplast cultures transformed with CRISPR constructs expressing sgRNAs using the *OsU6-2* promoter.

Toolkit testing in stably transformed *Arabidopsis* plants

Arabidopsis plants were transformed with 8 different CRISPR vectors (Fig. 3a). In all cases, the vectors contained two sgRNAs targeting the second exon of the *GL1* gene (Fig. 3b) and a Hygromycin selection cassette but contained different combinations of Pol III promoters for sgRNA expression and *CAS9* cassettes. After selection on Hygromycin, independent T0 lines for each vector were phenotypically analysed for loss of trichomes. Amongst the analysed lines, we found plants exhibiting WT-like trichome covering, chimeric phenotypes of partial glabrousness as well as full glabrousness (Fig. 3c, Suppl. File 5). In summary, all plants transformed with constructs expressing sgRNAs via the *OsU6-2* promoter but one did show WT-like trichome patterning. In contrast, most plants expressing the sgRNAs via the *AtU6-26* promoter showed mutational phenotypes. CasC proved to be especially efficient in creating fully mutant plants (Fig. 3d). We also isolated leaf genomic DNA of each plant and amplified the target region via PCR. We submitted the PCR amplicon to Sanger sequencing. Analysis of the Sanger sequencing histograms revealed high insertion-deletion (InDel) rates in all plants expressing the *GL1* sgRNAs under the control of the *AtU6-26* promoter, corresponding to the phenotypic analysis (Fig. 3e, Suppl. File 5 & 6). In contrast to the phenotypic analysis, we also detected induced mutations in some of the plants expressing sgRNAs under the *OsU6-2* promoter, albeit mostly at low frequencies. Those might not be sufficient to induce a phenotypic change, e.g. because cells are not homozygously mutated.

Figure 3:Testing of the ENABLE® Gene Editing *in planta* toolkit in stably transformed Arabidopsis plants. (a) Eight different CRISPR vectors were assembled to target the *AtGL1* gene with each containing a unique combination of *CAS9* cassette (CasA – CasD) and sgRNA expression cassette (sgRNA expression via *OsU6-2* promoter or *AtU6-26* promoter). (b) The *AtGL1* gene. Black boxes indicate exons. The position of the two sgRNA target sites in Exon 2 is annotated. The PAM sequence is underlined. (c) Exemplary pictures from Arabidopsis plants with WT trichome patterning (left), full glabrousness (middle, transformed with CasB_AtU6-26p::AtGL1_sgRNAs construct) or chimeric phenotype (right, transformed with CasD_AtU6-26p::AtGL1_sgRNAs construct). (d) Results of visual inspection of trichome phenotype in independent primary Arabidopsis transformants. (e) Results of *AtGL1* genotyping in independent primary Arabidopsis transformants.

Choosing the final toolkit components

The experiments in rice and Arabidopsis gave crucial insights into the optimal configuration for the ENABLE® vector system. (1) The main influence on gene editing efficiencies in our experiments came from the choice of the right sgRNA promoter. Hence, dicot- and monocot-specific versions of the vectors pGMF1 and pGMF2 (pGMF1-M and pGMF2-M, containing the *OsU6-2p* for sgRNA expression in monocots; pGMF1-D and pGMF2-D, containing the *AtU6-26p* for sgRNA expression in dicots) will be included in the final toolkit. (2) All previously published *CAS9* cassettes showed editing activity when combined with a suitable sgRNA setup. However, differences were visible in editing efficiencies with CasC outperforming the other *CAS9* modules in both rice and Arabidopsis. Given that the CasC cassette expresses the *CAS9* gene using a dicot promoter (*AtUBQ10* promoter), the strong activity of this module compared to other *CAS9* modules with monocot derived promoters in rice protoplasts was unexpected. However, the *AtUBQ10* promoter has been used successfully in the past for transgene expression and gene editing experiments in monocots such as rice and maize (Behera et al., 2015; Feike et al., 2019; Li et al., 2023). Hence, only the CasC cassette will be included as CAS9 expression vector pGFM4 vector in our ENABLE® toolkit. (3) All other toolkit components (pGMF3 backbone, pGMF5/6 selection markers) could be verified for their functionality in transient and stable gene editing experiments. In summary, our final toolkit will provide eight vectors, offering flexibility regarding use in monocots and dicots and in terms of transient or stable transformation for the end user (Fig. 4).

Figure 4: The final ENABLE® Gene Editing *in planta* toolkit. Eight plasmids are provided to the user. The user chooses five of those plasmids to assemble a T-DNA binary vector, based on the crop species they are working in and if they aim for transient or stable plant transformation.

DISCUSSION

Benefits of the ENABLE® Gene Editing *in planta* toolkit

Ongoing inequities hinder the uptake of plant biotechnology methods in LMIC and lead to research biases against crops relevant for those markets (Marks et al., 2023; Auge et al., 2024). These inequities range from lack of funding in general, over practical issues in scientific exchange with other researchers (e.g. because of VISA issues) to access to reagents and resources. The ENABLE® program of the Grow More Foundation aims at enabling plant scientists in LMIC to solve problems of their choice in their crop of interest by ensuring they have access to relevant technologies. One key emerging technology in plant breeding is the CRISPR/Cas9 technology, as precision breeding via targeted nucleases can play a pivotal role in supporting food

security in the Global South by tackling key traits such as resistance to biotic stresses, drought tolerance or nutrient content of crops (Chen et al., 2024; Ogbu et al., 2024; Amombo et al., 2025). Hence, the goal of this study was to develop a straightforward easy to use molecular toolkit to improve access to gene editing technology following these key requirements: (1) Work across monocot and dicot species to cover as many relevant crop species as possible. (2) Suitable for scientists new to the CRISPR technology. (3) Simplified cloning procedure for all levels of experience. (4) Where possible, the toolkit should enable low cost ways of cloning and mutation detection compared to "gold-standard", but pricey sequencing technologies. (5) Should be accessible via open repositories to allow simple distribution. And finally and most critically, (6) provide real life translational applicability. With the current ENABLE® toolkit version, we believe we have achieved these goals. By providing a set of only eight vectors, we allow the user to clone a unique CRISPR vector tailored to their target sequence of interest, which can be used for knockout generation in monocot or dicot species, in both transient or stable transformation capacities, hence covering the most common use cases for CRISPR in plants. We tested its application in a monocot and dicot species and benchmarked the efficiency of the toolkit CAS9 module against other plant CAS9 modules. While many plant CRISPR vector collections have been described in the past years (see e.g. https://www.addgene.org/crispr/plants), we wanted to focus on the gap between sets of vectors limited to specific CRISPR use cases, e.g. in a particular plant/one of the two groups of angiosperms only (e.g. Jacobs et al., 2015; Ordon et al., 2017; Wu et al., 2018) and large collections of plasmids (Lowder et al., 2015; Čermák et al., 2017; Hahn et al., 2020; Cheng et al., 2025) offering dozens of different modules, giving experienced users a plethora of options for their experiment but being potentially overwhelming for CRISPR beginners. To address this, as well as provide a starting point for users new to the CRISPR technology, we developed a detailed guide for the construct design process, from sgRNA target site selection to construct assembly and editing verification (Suppl. File 2). Yet despite being beginner-friendly, our toolkit can be applied for actual gene editing experiments *in planta*, setting it apart from purely educational classroom CRISPR kits (Collins et al., 2024). Our ENABLE® toolkit makes use of the MoClo Golden Gate assembly technology, a modular cloning method that is proven to function efficiently (Suppl. Fig. 1b), uses just three different enzymes and a basic thermocycler, independently of methods such as PCR or gel extraction (Weber et al.,

2011; Engler et al., 2014; Patron et al., 2015), thereby eliminating the typical pain points in cloning workflows and reducing costs. Additionally, the selection of (likely) positive clones via blue/white and red/white screening during the cloning process (Suppl. Fig. 1a) is beneficial for labs with limited resources. Similarly, as our ENABLE toolkit allows generation of large deletions by targeting a gene with two sgRNAs at once, we also provide a low cost way for mutation verification via PCR based methods (Fig. 2d). Finally, our ENABLE® kit is accessible on Addgene (Addgene ID: 1000000270) to (almost) every country in the world.

Limitations of the ENABLE® Gene Editing toolkit

Our ENABLE® toolkit focuses mainly on the generation of functional knockouts, the most common CRISPR use case *in planta*. While the use of two sgRNAs allows for some additional specialised scenarios (multiplexing, targeted deletions of gene regions), other gene editing applications, such as base editing, gene activation/repression or gene knock-ins are outside the scope of the ENABLE® kit. Similarly, while the two transformation selection marker genes within the toolkit (Hygromycin resistance and eGFP fluorescence) allow for selection of stable or transient transformants in most plant species, other selection markers with specific use cases, such as FAST (Shimada et al., 2010) or RUBY (He et al., 2020) might be advantageous for some users. Due to the modular nature of the ENABLE® vectors and the use of the widely used MoClo syntax (Weber et al., 2011; Patron et al., 2015), different nuclease modules as well as resistance markers from existing resources (e.g. Engler et al., 2014; Hahn et al., 2020) could be easily ordered from platforms such as Addgene to replace the CAS9 cassette and/or selection marker cassettes in our toolkit if needed. Our study showed the dramatic impact of the choice of a suitable Pol III promoter for sgRNA expression on gene editing efficiencies. While we provide monocot- and dicot-specific modules for sgRNA expression, it has been shown in various studies that species-specific promoters tend to provide the best editing efficiencies when using Pol III promoters for sgRNA expression (Sun et al., 2015; Ren et al., 2021; Zhang et al., 2022; Riu et al., 2023). However, species-specific promoters are not suitable for a simplified toolkit with broad applicability. Expression of sgRNAs using Pol II promoters coupled with RNA processing systems, such as the tRNA processing system or self-cleaving ribozymes can provide an alternative way of achieving high sgRNA expression rates across species (Gao and Zhao, 2014; Xie et

al., 2015; Milner et al., 2024) and will be considered for future versions of the ENABLE® toolkit. Our current distribution of the ENABLE® toolkit via Addgene provides the possibility for (almost) worldwide delivery of the toolkit. Still, we also acknowledge that the pricing system of Addgene (currently around 440 USD for the toolkit) can be a barrier for some researchers in LMIC. While Addgene provides case-by-case discounts on its plasmids for laboratories experiencing financial difficulties, we also aim to explore further ways to distribute the plasmids, e.g. via local distribution hubs in the future. A final limitation is the accessibility of the enzymes and buffers needed for the cut-ligation reactions. While major enzyme providers have local distribution partners in many countries, there are still countries which have difficulties obtaining basic molecular biology restriction enzymes. Also sublimation of dry ice in shipments is a significant barrier for LMIC accessibility. Hence, future versions of the toolkit could contain lyophilized versions of the relevant enzymes that are shipped together with the eight vectors. Any updated versions of the ENABLE® toolkit will be published on https://www.growmorefoundation.org/enable-crispr-kits .

The ENABLE® program in the context of ongoing initiatives and recent developments to enable scientists in the Global South

We see that the ENABLE® toolkit complements existing initiatives, such as the Beneficial Bio network (https://beneficial.bio/) aiming to provide laboratory reagents and consumables to resource-limited laboratories worldwide or the ReClone network (https://reclone.org/), an open source plasmid sharing platform with a focus on the Global South, and that it provides an additional angle with its focus on plant sciences. Additionally, ongoing initiatives such as the African BioGenome Project (https://africanbiogenome.org/), which aims at sequencing huge collections of locally relevant plants (and other species) or the sequencing efforts by the African Orphan Crops Consortium (https://africanorphancrops.org/) will provide much needed sequence information for CRISPR technology usage specifically in local crop varieties. Various local initiatives, e.g. SynBio4ALL Africa (https://synbio4all.wixsite.com/synbio4all), focus on training delivery around synthetic and molecular biology methods for the future generation of scientists. Another noteworthy community-driven effort is the GetGenome initiative, which provides *pro bono* whole plasmid sequencing for LMIC (https://getgenome.net/callforprojects/ggplasmidseq), which could be specifically

useful in combination with CRISPR plasmid cloning projects using the ENABLE® toolkit. These initiatives, combined with the impact of new large language models on CRISPR experiments (Qu et al., 2025; Ruffolo et al., 2025), with recent improvements in plant transformation (Lowe et al., 2016; Debernardi et al., 2020; Che et al., 2022; Thorpe et al., 2025; Weiss et al., 2025) and with the proven track record of CRISPR-enhanced crops over the last decade (Akanmu et al., 2024), make us certain that now is the right time to improve the accessibility of this technology for everyone.

Food production will always face biological, environmental, and societal challenges, and we need to apply the most efficient strategies to face these challenges. CRISPR/Cas9 is an accessible and affordable way to do so, especially when made accessible to LMIC and being applied to underutilized crops. Our ENABLE® toolkit provides the first molecular step towards equitable access, bridging the gap between the biotechnological advancements and geographical location. We envision that the ENABLE® Gene Editing *in planta* kit will be widely utilized for teaching purposes for newcomers to this technology at first and then grow from basic to applied science.

ACKNOWLEDGEMENTS

We want to thank Laura Luoni (University of Milan) for the stable transformation of Arabidopsis plants with our CRISPR constructs and plant care. We would like to thank Vittorio Venturi (ICGEB Trieste) for providing us access to rice growth room facilities and Nezka Kavcic (ICGEB Trieste) for supporting fluorescence microscopy procedures. We want to thank Laura Dalle Carbonare, Vinay Shukla and Francesco Licausi, who tried out early versions of the ENABLE® toolkit at the University of Oxford. We want to acknowledge Daria Chrobok for illustration of some of our concepts. We want to thank Vladimir Nekrasov (Rothamsted Research), Nicola Patron (University of Cambridge), Jonathan Jones (The Sainsbury Laboratory) and Wendy Harwood (John Innes Centre) for provision of plasmids used in this study. We would like to thank Tom Speedy from Integrated DNA Technologies (IDT) as well as Penny Devoe and Carole Keating from New England Biolabs (NEB) for *pro bono* provision of gene synthesis and reagents.

CONTRIBUTIONS

Wet lab experiments were executed and analysed by Birhan Abate with support from Florian Hahn and Daniele Chirivì. Florian Hahn designed the Golden Gate constructs, CRISPR targets and experimental strategy. Kate Creasey Krainer conceptualised the idea of the ENABLE® Gene Editing *in planta* toolkit. Camilla Betti, Fabio Fornara, Jenny Molloy, Florian Hahn and Kate Creasey Krainer supervised the project progress and contributed to data interpretation. Florian Hahn wrote the manuscript with contributions from Kate Creasey Krainer and Birhan Abate.

FUNDING STATEMENT

This work was funded by the Grow More Foundation and ICGEB with additional funds awarded to Camilla Betti through PSR_Linea 3 829 FLORICE_My First SEED (MUR DM6 737/2021).

CONFLICT OF INTEREST STATEMENT

Kate Creasey Krainer and Florian Hahn are named inventors on a filed patent (US-20230348921-A1) and a filed provisional patent application (US63/913,876) relating to gene editing molecular toolkits. Jennifer Molloy is Executive Director of Beneficial Bio and Global Co-Lead of the ReClone Network.

REFERENCES

Abe, F., Haque, E., Hisano, H., Tanaka, T., Kamiya, Y., Mikami, M., Kawaura, K., Endo, M., Onishi, K., Hayashi, T., Sato, K., 2019. Genome-Edited Triple-Recessive Mutation Alters Seed Dormancy in Wheat. Cell Reports 28, 1362-1369.e4. https://doi.org/10.1016/j.celrep.2019.06.090

Ahmar, S., Usman, B., Hensel, G., Jung, K.-H., Gruszka, D., 2024. CRISPR enables sustainable cereal production for a greener future. Trends in Plant Science 29, 179–195. https://doi.org/10.1016/j.tplants.2023.10.016

Akanmu, A.O., Asemoloye, M.D., Marchisio, M.A., Babalola, O.O., 2024. Adoption of CRISPR-Cas for crop production: present status and future prospects. PeerJ 12, e17402. https://doi.org/10.7717/peerj.17402

Amoah, P., Oumarou Mahamane, A.-R., Byiringiro, M.H., Mahula, N.J., Manneh, N., Oluwasegun, Y.R., Assfaw, A.T., Mukiti, H.M., Garba, A.D., Chiemeke, F.K., Bernard Ojuederie, O., Olasanmi, B., 2024. Genome editing in Sub-Saharan Africa: a game-changing strategy for climate change mitigation and sustainable agriculture. GM Crops & Food 15, 279–302. https://doi.org/10.1080/21645698.2024.2411767

Amombo, E., Mumm, R.H., Nilahyane, A., Hirich, A., Hijri, M., Ntui, V.O., 2025. Enhancing Crop Nutrition in Arid and Semiarid Regions of Africa Through Genome Editing Using CRISPR /Cas. Food and Energy Security 14, e70068. https://doi.org/10.1002/fes3.70068

Auge, G., Sunil, R.S., Ingle, R.A., Rahul, P.V., Mutwil, M., Estevez, J.M., 2024. Current challenges for plant biology research in the Global South. New Phytologist 244, 1168–1174. https://doi.org/10.1111/nph.20083

Banakar, R., Schubert, M., Collingwood, M., Vakulskas, C., Eggenberger, A.L., Wang, K., 2020. Comparison of CRISPR-Cas9/Cas12a Ribonucleoprotein Complexes for Genome Editing Efficiency in the Rice Phytoene Desaturase (OsPDS) Gene. Rice 13, 4. https://doi.org/10.1186/s12284-019-0365-z

Behera, S., Wang, N., Zhang, C., Schmitz-Thom, I., Strohkamp, S., Schültke, S., Hashimoto, K., Xiong, L., Kudla, J., 2015. Analyses of Ca^{2+} dynamics using a ubiquitin-10 promoter-driven Yellow Cameleon 3.6 indicator reveal reliable transgene expression and differences in cytoplasmic Ca^{2+} responses in Arabidopsis and rice (*Oryza sativa*) roots. New Phytologist 206, 751–760. https://doi.org/10.1111/nph.13250

Castel, B., Tomlinson, L., Locci, F., Yang, Y., Jones, J.D.G., 2019. Optimization of T-DNA architecture for Cas9-mediated mutagenesis in Arabidopsis. PLOS ONE 14, e0204778. https://doi.org/10.1371/journal.pone.0204778

Čermák, T., Curtin, S.J., Gil-Humanes, J., Čegan, R., Kono, T.J.Y., Konečná, E., Belanto, J.J., Starker, C.G., Mathre, J.W., Greenstein, R.L., Voytas, D.F., 2017. A Multipurpose Toolkit to Enable Advanced Genome Engineering in Plants. Plant Cell 29, 1196–1217. https://doi.org/10.1105/tpc.16.00922

Che, P., Wu, E., Simon, M.K., Anand, A., Lowe, K., Gao, H., Sigmund, A.L., Yang, M., Albertsen, M.C., Gordon-Kamm, W., Jones, T.J., 2022. Wuschel2 enables highly efficient CRISPR/Cas-targeted genome editing during rapid de novo shoot regeneration in sorghum. Commun Biol 5, 344. https://doi.org/10.1038/s42003-022-03308-w

Chen, F., Chen, L., Yan, Z., Xu, J., Feng, L., He, N., Guo, M., Zhao, J., Chen, Z., Chen, H., Yao, G., Liu, C., 2024. Recent advances of CRISPR-based genome editing for enhancing staple crops. Front. Plant Sci. 15, 1478398. https://doi.org/10.3389/fpls.2024.1478398

Cheng, Y., Li, G., Qi, A., Mandlik, R., Pan, C., Wang, D., Ge, S., Qi, Y., 2025. A comprehensive all-in-one CRISPR toolbox for large-scale screens in plants. The Plant Cell 37, koaf081. https://doi.org/10.1093/plcell/koaf081

Clement, K., Rees, H., Canver, M.C., Gehrke, J.M., Farouni, R., Hsu, J.Y., Cole, M.A., Liu, D.R., Joung, J.K., Bauer, D.E., Pinello, L., 2019. CRISPResso2 provides accurate and rapid genome editing sequence analysis. Nat Biotechnol 37, 224–226. https://doi.org/10.1038/s41587-019-0032-3

Clough, S.J., Bent, A.F., 1998. Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. The Plant Journal 16, 735–743. https://doi.org/10.1046/j.1365-313x.1998.00343.x

Collins, M., Lau, M.B., Ma, W., Shen, A., Wang, B., Cai, S., La Russa, M., Jewett, M.C., Qi, L.S., 2024. A frugal CRISPR kit for equitable and accessible education in gene editing and synthetic biology. Nat Commun 15, 6563. https://doi.org/10.1038/s41467-024-50767-2

Concordet, J.-P., Haeussler, M., 2018. CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens. Nucleic Acids Research 46, W242–W245. https://doi.org/10.1093/nar/gky354

Connelly, S., Marshallsay, C., Leader, D., Brown, J.W.S., Filipowicz, W., 1994. Small Nuclear RNA Genes Transcribed by Either RNA Polymerase II or RNA Polymerase III in Monocot Plants Share Three Promoter Elements and Use a Strategy To Regulate Gene Expression Different from That Used by Their Dicot Plant Counterparts. Molecular and Cellular Biology 14, 5910–5919. https://doi.org/10.1128/mcb.14.9.5910-5919.1994

Dai, Y., Hu, G., Dupas, A., Medina, L., Blandels, N., San Clemente, H., Ladouce, N., Badawi, M., Hernandez-Raquet, G., Mounet, F., Grima-Pettenati, J., Cassan-Wang, H., 2020. Implementing the CRISPR/Cas9 Technology in Eucalyptus Hairy Roots Using Wood-Related Genes. IJMS 21, 3408. https://doi.org/10.3390/ijms21103408

Dang, Y., Jia, G., Choi, J., Ma, H., Anaya, E., Ye, C., Shankar, P., Wu, H., 2015. Optimizing sgRNA structure to improve CRISPR-Cas9 knockout efficiency. Genome Biol 16, 280. https://doi.org/10.1186/s13059-015-0846-3

Debernardi, J.M., Tricoli, D.M., Ercoli, M.F., Hayta, S., Ronald, P., Palatnik, J.F., Dubcovsky, J., 2020. A GRF–GIF chimeric protein improves the regeneration efficiency of transgenic plants. Nature Biotechnology 38, 1274–1279. https://doi.org/10.1038/s41587-020-0703-0

Engler, C., Gruetzner, R., Kandzia, R., Marillonnet, S., 2009. Golden Gate Shuffling: A One-Pot DNA Shuffling Method Based on Type IIs Restriction Enzymes. PLoS ONE 4, e5553. https://doi.org/10.1371/journal.pone.0005553

Engler, C., Youles, M., Gruetzner, R., Ehnert, T.-M., Werner, S., Jones, J.D.G., Patron, N.J., Marillonnet, S., 2014. A Golden Gate Modular Cloning Toolbox for Plants. ACS Synth. Biol. 3, 839–843. https://doi.org/10.1021/sb4001504

Evenson, R.E., Gollin, D., 2003. Assessing the Impact of the Green Revolution, 1960 to 2000. Science 300, 758–762. https://doi.org/10.1126/science.1078710

FAO, IFAD, UNICEF, WFP and WHO, 2025. In Brief to The State of Food Security and Nutrition in the World 2025. FAO; IFAD; UNICEF; WFP; WHO; https://doi.org/10.4060/cd6015en

Fauser, F., Schiml, S., Puchta, H., 2014. Both CRISPR/Cas-based nucleases and nickases can be used efficiently for genome engineering in Arabidopsis thaliana. Plant J. 79, 348–359. https://doi.org/10.1111/tpj.12554

Feike, D., Korolev, A.V., Soumpourou, E., Murakami, E., Reid, D., Breakspear, A., Rogers, C., Radutoiu, S., Stougaard, J., Harwood, W.A., Oldroyd, G.E.D., Miller, J.B., 2019. Characterizing standard genetic parts and establishing common principles for engineering legume and cereal roots. Plant Biotechnology Journal 17, 2234–2245. https://doi.org/10.1111/pbi.13135

Feng, Z., Zhang, B., Ding, W., Liu, X., Yang, D.-L., Wei, P., Cao, F., Zhu, S., Zhang, F., Mao, Y., Zhu, J.-K., 2013. Efficient genome editing in plants using a CRISPR/Cas system. Cell Res 23, 1229–1232. https://doi.org/10.1038/cr.2013.114

Gao, Y., Zhao, Y., 2014. Self-processing of ribozyme-flanked RNAs into guide RNAs in vitro and in vivo for CRISPR-mediated genome editing. J. Integr. Plant Biol. 56, 343–349. https://doi.org/10.1111/jipb.12152

Gomez, M.A., Lin, Z.D., Moll, T., Chauhan, R.D., Hayden, L., Renninger, K., Beyene, G., Taylor, N.J., Carrington, J.C., Staskawicz, B.J., Bart, R.S., 2019. Simultaneous CRISPR /Cas9-mediated editing of cassava *EIF 4E* isoforms *NCBP -1* and *NCBP -2* reduces cassava brown streak disease symptom severity and incidence. Plant Biotechnology Journal 17, 421–434. https://doi.org/10.1111/pbi.12987

Grützner, R., Martin, P., Horn, C., Mortensen, S., Cram, E.J., Lee-Parsons, C.W.T., Stuttmann, J., Marillonnet, S., 2021. High-efficiency genome editing in plants mediated by a Cas9 gene containing multiple introns. Plant Communications 2, 100135. https://doi.org/10.1016/j.xplc.2020.100135

Hahn, F., Korolev, A., Sanjurjo Loures, L., Nekrasov, V., 2020. A modular cloning toolkit for genome editing in plants. BMC Plant Biology 20, 179. https://doi.org/10.1186/s12870-020-02388-2

Hahn, F., Mantegazza, O., Greiner, A., Hegemann, P., Eisenhut, M., Weber, A.P.M., 2017. An Efficient Visual Screen for CRISPR/Cas9 Activity in Arabidopsis thaliana. Front. Plant Sci. 8. https://doi.org/10.3389/fpls.2017.00039

Hao, J., Yang, Y., Futrell, S., Kelly, E.A., Lorts, C.M., Nebie, B., Runo, S., Yang, J., Alvarez, S., Lasky, J.R., Schachtman, D.P., 2023. CRISPR/Cas9-Mediated Mutagenesis of *Carotenoid Cleavage Dioxygenase* (*CCD*) Genes in Sorghum Alters Strigolactone Biosynthesis and Plant Biotic Interactions. Phytobiomes Journal PBIOMES-08-22-0053-R. https://doi.org/10.1094/PBIOMES-08-22-0053-R

He, F., Chen, S., Ning, Y., Wang, G., 2016. Rice (*Oryza sativa*) Protoplast Isolation and Its Application for Transient Expression Analysis. CP Plant Biology 1, 373–383. https://doi.org/10.1002/cppb.20026

He, Y., Zhang, T., Sun, H., Zhan, H., Zhao, Y., 2020. A reporter for noninvasively monitoring gene expression and plant transformation. Hortic Res 7, 152. https://doi.org/10.1038/s41438-020-00390-1

Jacobs, T.B., LaFayette, P.R., Schmitz, R.J., Parrott, W.A., 2015. Targeted genome modifications in soybean with CRISPR/Cas9. BMC Biotechnol 15, 16. https://doi.org/10.1186/s12896-015-0131-2

Jethva, J., Hahn, F., Giuliani, R., Peeters, N., Cousins, A., Kelly, S., 2024. Realisation of a key step in the evolution of C_4 photosynthesis in rice by genome editing. https://doi.org/10.1101/2024.05.21.595093

Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J.A., Charpentier, E., 2012. A Programmable Dual-RNA–Guided DNA Endonuclease in Adaptive Bacterial Immunity. Science 337, 816–821. https://doi.org/10.1126/science.1225829

Kong, W., Wang, M., Huang, L., Wu, F., Tao, J., Mo, B., Yu, Y., 2023. A high-efficient and naked-eye visible CRISPR/Cas9 system in Arabidopsis. Planta 257, 30. https://doi.org/10.1007/s00425-022-04060-5

Lawrenson, T., Shorinola, O., Stacey, N., Li, C., Østergaard, L., Patron, N., Uauy, C., Harwood, W., 2015. Induction of targeted, heritable mutations in barley and Brassica oleracea using RNA-guided Cas9 nuclease. Genome Biol 16, 258. https://doi.org/10.1186/s13059-015-0826-7

Li, C., Nguyen, V., Liu, J., Fu, W., Chen, C., Yu, K., Cui, Y., 2019. Mutagenesis of seed storage protein genes in Soybean using CRISPR/Cas9. BMC Res Notes 12, 176. https://doi.org/10.1186/s13104-019-4207-2

Li, Z., Zhong, Z., Wu, Z., Pausch, P., Al-Shayeb, B., Amerasekera, J., Doudna, J.A., Jacobsen, S.E., 2023. Genome editing in plants using the compact editor CasΦ. Proc. Natl. Acad. Sci. U.S.A. 120, e2216822120. https://doi.org/10.1073/pnas.2216822120

Liang, Z., Wu, Y., Ma, L., Guo, Y., Ran, Y., 2022. Efficient Genome Editing in Setaria italica Using CRISPR/Cas9 and Base Editors. Front. Plant Sci. 12, 815946. https://doi.org/10.3389/fpls.2021.815946

Lin, C.-S., Hsu, C.-T., Yang, L.-H., Lee, L.-Y., Fu, J.-Y., Cheng, Q.-W., Wu, F.-H., Hsiao, H.C.-W., Zhang, Y., Zhang, R., Chang, W.-J., Yu, C.-T., Wang, W., Liao, L.-J., Gelvin, S.B., Shih, M.-C., 2018. Application of protoplast technology to CRISPR/Cas9 mutagenesis: from single-cell mutation detection to mutant plant regeneration. Plant Biotechnology Journal 16, 1295–1310. https://doi.org/10.1111/pbi.12870

Liu, G., Lin, Q., Jin, S., Gao, C., 2022. The CRISPR-Cas toolbox and gene editing technologies. Molecular Cell 82, 333–347. https://doi.org/10.1016/j.molcel.2021.12.002

Liu, Q., Yang, F., Zhang, J., Liu, H., Rahman, S., Islam, S., Ma, W., She, M., 2021. Application of CRISPR/Cas9 in Crop Quality Improvement. IJMS 22, 4206. https://doi.org/10.3390/ijms22084206

Lowder, L.G., Zhang, D., Baltes, N.J., Paul, J.W., Tang, X., Zheng, X., Voytas, D.F., Hsieh, T.-F., Zhang, Y., Qi, Y., 2015. A CRISPR/Cas9 Toolbox for Multiplexed Plant Genome Editing and Transcriptional Regulation. Plant Physiol 169, 971–985. https://doi.org/10.1104/pp.15.00636

Lowe, K., Wu, E., Wang, N., Hoerster, G., Hastings, C., Cho, M.-J., Scelonge, C., Lenderts, B., Chamberlin, M., Cushatt, J., Wang, L., Ryan, L., Khan, T., Chow-Yiu, J., Hua, W., Yu, M., Banh, J., Bao, Z., Brink, K., Igo, E., Rudrappa, B., Shamseer, P.M., Bruce, W., Newman, L., Shen, B., Zheng, P., Bidney, D., Falco, C., Register, J., Zhao, Z.-Y., Xu, D., Jones, T., Gordon-Kamm, W., 2016. Morphogenic Regulators Baby boom and Wuschel Improve Monocot Transformation. The Plant Cell 28, 1998–2015. https://doi.org/10.1105/tpc.16.00124

Luginbuehl, L.H., El-Sharnouby, S., Wang, N., Hibberd, J.M., 2020. Fluorescent reporters for functional analysis in rice leaves. Plant Direct 4, e00188. https://doi.org/10.1002/pld3.188

Marks, R.A., Amézquita, E.J., Percival, S., Rougon-Cardoso, A., Chibici-Revneanu, C., Tebele, S.M., Farrant, J.M., Chitwood, D.H., VanBuren, R., 2023. A critical analysis of plant science literature reveals ongoing inequities. Proc. Natl. Acad. Sci. U.S.A. 120, e2217564120. https://doi.org/10.1073/pnas.2217564120

Mikami, M., Toki, S., Endo, M., 2015. Comparison of CRISPR/Cas9 expression constructs for efficient targeted mutagenesis in rice. Plant Mol Biol 88, 561–572. https://doi.org/10.1007/s11103-015-0342-x

Milner, M.J., Sharma, M., Bates, R.E., Whiting, M., Craze, M.S., Miller, P., Brooks, J., Kouidri, A., Wallington, E.J., 2024. Differential editing efficiencies in cereal crops: a comparative analysis of tRNA and ribozyme multiplexed guide delivery. Front. Plant Sci. 15, 1426184. https://doi.org/10.3389/fpls.2024.1426184

Missanga, J.S., Mmbando, G.S., 2025. Challenges of food security in Africa in the face of climate change: potential of the clustered regularly interspaced short

palindromic repeats (CRISPR)-cas9 technology to address these challenges. Discov Sustain 6, 305. https://doi.org/10.1007/s43621-025-01092-1

Mukami, A., Juma, B.S., Mweu, C., Oduor, R., Mbinda, W., 2024. CRISPR-Cas9-induced targeted mutagenesis of feruloyl CoA 6′-hydroxylase gene reduces postharvest physiological deterioration in cassava roots. Postharvest Biology and Technology 208, 112649. https://doi.org/10.1016/j.postharvbio.2023.112649

Ndudzo, A., Sibanda Makuvise, A., Moyo, S., Bobo, E.D., 2024. CRISPR-Cas9 genome editing in crop breeding for climate change resilience: Implications for smallholder farmers in Africa. Journal of Agriculture and Food Research 16, 101132. https://doi.org/10.1016/j.jafr.2024.101132

Nie, H., Shi, Y., Geng, X., Xing, G., 2022. CRISRP/Cas9-Mediated Targeted Mutagenesis of Tomato Polygalacturonase Gene (SlPG) Delays Fruit Softening. Front. Plant Sci. 13, 729128. https://doi.org/10.3389/fpls.2022.729128

Ogbu, A.C., Ume, O.A., Nwogiji, O.C., Akeredolu, S.O., Eze, O.G., Onwe, C.V., Oduguwa, O.I., Chigozie, U.V., 2024. Advances in the use of Genome Editing Tools in Africa: A Review. Asian J. Biochem. Gen. Mol. Biol. 16, 48–67. https://doi.org/10.9734/ajbgmb/2024/v16i12422

Onyeaka, H., Nwauzoma, U.M., Akinsemolu, A.A., Tamasiga, P., Duan, K., Al-Sharify, Z.T., Siyanbola, K.F., 2024. The ripple effects of climate change on agricultural sustainability and food security in Africa. Food and Energy Security 13, e567. https://doi.org/10.1002/fes3.567

Ordon, J., Gantner, J., Kemna, J., Schwalgun, L., Reschke, M., Streubel, J., Boch, J., Stuttmann, J., 2017. Generation of chromosomal deletions in dicotyledonous plants employing a user-friendly genome editing toolkit. Plant J 89, 155–168. https://doi.org/10.1111/tpj.13319

Pacher, M., Puchta, H., 2017. From classical mutagenesis to nuclease-based breeding – directing natural DNA repair for a natural end-product. The Plant Journal 90, 819–833. https://doi.org/10.1111/tpj.13469

Patron, N.J., Orzaez, D., Marillonnet, S., Warzecha, H., Matthewman, C., Youles, M., Raitskin, O., Leveau, A., Farré, G., Rogers, C., Smith, A., Hibberd, J., Webb, A.A.R., Locke, J., Schornack, S., Ajioka, J., Baulcombe, D.C., Zipfel, C., Kamoun, S., Jones, J.D.G., Kuhn, H., Robatzek, S., Van Esse, H.P., Sanders, D., Oldroyd, G., Martin, C., Field, R., O'Connor, S., Fox, S., Wulff, B., Miller, B., Breakspear, A., Radhakrishnan, G., Delaux, P.-M., Loqué, D., Granell, A., Tissier, A., Shih, P., Brutnell, T.P., Quick, W.P., Rischer, H., Fraser, P.D., Aharoni, A., Raines, C., South, P.F., Ané, J.-M., Hamberger, B.R., Langdale, J., Stougaard, J., Bouwmeester, H., Udvardi, M., Murray, J.A.H., Ntoukakis, V., Schäfer, P., Denby, K., Edwards, K.J., Osbourn, A., Haseloff, J., 2015. Standards for plant synthetic biology: a common syntax for exchange of DNA parts. New Phytol 208, 13–19. https://doi.org/10.1111/nph.13532

Polidoros, A., Nianiou-Obeidat, I., Tsakirpaloglou, N., Petrou, N., Deligiannidou, E., Makri, N.-M., 2024. Genome-Editing Products Line up for the Market: Will Europe Harvest the Benefits from Science and Innovation? Genes 15, 1014. https://doi.org/10.3390/genes15081014

Qaim, M., 2020. Role of New Plant Breeding Technologies for Food Security and Sustainable Agricultural Development. Applied Eco Perspectives Pol 42, 129–150. https://doi.org/10.1002/aepp.13044

Qu, Y., Huang, K., Yin, M., Zhan, K., Liu, D., Yin, D., Cousins, H.C., Johnson, W.A., Wang, X., Shah, M., Altman, R.B., Zhou, D., Wang, M., Cong, L., 2025. CRISPR-GPT for agentic automation of gene-editing experiments. Nat. Biomed. Eng. https://doi.org/10.1038/s41551-025-01463-z

Raitskin, O., Schudoma, C., West, A., Patron, N.J., 2019. Comparison of efficiency and specificity of CRISPR-associated (Cas) nucleases in plants: An expanded toolkit for precision genome engineering. PLOS ONE 14, e0211598. https://doi.org/10.1371/journal.pone.0211598

Ren, C., Liu, Y., Guo, Y., Duan, W., Fan, P., Li, S., Liang, Z., 2021. Optimizing the CRISPR/Cas9 system for genome editing in grape by using grape promoters. Hortic Res 8, 52. https://doi.org/10.1038/s41438-021-00489-z

Riu, Y.-S., Kim, G.H., Chung, K.W., Kong, S.-G., 2023. Enhancement of the CRISPR/Cas9-Based Genome Editing System in Lettuce (Lactuca sativa L.) Using the Endogenous U6 Promoter. Plants 12, 878. https://doi.org/10.3390/plants12040878

Ruffolo, J.A., Nayfach, S., Gallagher, J., Bhatnagar, A., Beazer, J., Hussain, R., Russ, J., Yip, J., Hill, E., Pacesa, M., Meeske, A.J., Cameron, P., Madani, A., 2025. Design of highly functional genome editors by modelling CRISPR–Cas sequences. Nature 645, 518–525. https://doi.org/10.1038/s41586-025-09298-z

Senthil, K., Rathinam, M., Parashar, M., Dokka, N., Tyagi, S., Mathur, V., Sharma, S., Gaikwad, K., Bhattacharya, R., Sreevathsa, R., 2025. Establishing a CRISPR/Cas9 genome editing framework in pigeonpea (Cajanus cajan L.) by targeting phytoene desaturase (PDS) gene disruption. Journal of Genetic Engineering and Biotechnology 23, 100465. https://doi.org/10.1016/j.jgeb.2025.100465

Shimada, T.L., Shimada, T., Hara-Nishimura, I., 2010. A rapid and non-destructive screenable marker, FAST, for identifying transformed seeds of *Arabidopsis thaliana*. The Plant Journal 61, 519–528. https://doi.org/10.1111/j.1365-313X.2009.04060.x

Smedley, M.A., Hayta, S., Clarke, M., Harwood, W.A., 2021. CRISPR-Cas9 Based Genome Editing in Wheat. Current Protocols 1, e65. https://doi.org/10.1002/cpz1.65

Sun, X., Hu, Z., Chen, R., Jiang, Q., Song, G., Zhang, H., Xi, Y., 2015. Targeted mutagenesis in soybean using the CRISPR-Cas9 system. Sci Rep 5, 10342. https://doi.org/10.1038/srep10342

Syombua, E.D., Zhang, Z., Tripathi, J.N., Ntui, V.O., Kang, M., George, O.O., Edward, N.K., Wang, K., Yang, B., Tripathi, L., 2021. A CRISPR/Cas9-based genome-editing system for yam (*Dioscorea* spp.). Plant Biotechnology Journal 19, 645–647. https://doi.org/10.1111/pbi.13515

Tadele, Z., 2019. Orphan crops: their importance and the urgency of improvement. Planta 250, 677–694. https://doi.org/10.1007/s00425-019-03210-6

Thorpe, C., Luo, W., Ji, Q., Eggenberger, A.L., Chicowski, A.S., Xu, W., Sandhu, R., Lee, K., Whitham, S.A., Qi, Y., Wang, K., Jiang, S., 2025. Enhancing biolistic plant transformation and genome editing with a flow guiding barrel. Nat Commun 16, 5624. https://doi.org/10.1038/s41467-025-60761-x

UN, 2024. The Indispensability of STEM for Sustaining Economic Growth in the 21st Century - Africa Needs to Intensify Efforts. UN.

Venezia, M., Creasey Krainer, K.M., 2021. Current Advancements and Limitations of Gene Editing in Orphan Crops. Front. Plant Sci. 12, 742932. https://doi.org/10.3389/fpls.2021.742932

Wang, J.Y., Doudna, J.A., 2023. CRISPR technology: A decade of genome editing is only the beginning. Science 379, eadd8643. https://doi.org/10.1126/science.add8643

Weber, E., Engler, C., Gruetzner, R., Werner, S., Marillonnet, S., 2011. A Modular Cloning System for Standardized Assembly of Multigene Constructs. PLOS ONE 6, e16765. https://doi.org/10.1371/journal.pone.0016765

Weiss, T., Kamalu, M., Shi, H., Li, Z., Amerasekera, J., Zhong, Z., Adler, B.A., Song, M.M., Vohra, K., Wirnowski, G., Chitkara, S., Ambrose, C., Steinmetz, N., Sridharan, A., Sahagun, D., Banfield, J.F., Doudna, J.A., Jacobsen, S.E., 2025. Viral delivery of an RNA-guided genome editor for transgene-free germline editing in Arabidopsis. Nat. Plants 11, 967–976. https://doi.org/10.1038/s41477-025-01989-9

Wu, R., Lucke, M., Jang, Y., Zhu, W., Symeonidi, E., Wang, C., Fitz, J., Xi, W., Schwab, R., Weigel, D., 2018. An efficient CRISPR vector toolbox for engineering large deletions in Arabidopsis thaliana. Plant Methods 14, 65. https://doi.org/10.1186/s13007-018-0330-7

Xie, K., Minkenberg, B., Yang, Y., 2015. Boosting CRISPR/Cas9 multiplex editing capability with the endogenous tRNA-processing system. PNAS 112, 3570–3575. https://doi.org/10.1073/pnas.1420294112

Zhang, Hui, Zhang, J., Wei, P., Zhang, B., Gou, F., Feng, Z., Mao, Y., Yang, L., Zhang, Heng, Xu, N., Zhu, J., 2014. The CRISPR / C as9 system produces specific and homozygous targeted gene editing in rice in one generation. Plant Biotechnology Journal 12, 797–807. https://doi.org/10.1111/pbi.12200

Zhang, S., Wu, S., Hu, C., Yang, Q., Dong, T., Sheng, O., Deng, G., He, W., Dou, T., Li, C., Sun, C., Yi, G., Bi, F., 2022. Increased mutation efficiency of CRISPR/Cas9 genome editing in banana by optimized construct. PeerJ 10, e12664. https://doi.org/10.7717/peerj.12664

Zhang, Y., Iaffaldano, B., Qi, Y., 2021. CRISPR ribonucleoprotein-mediated genetic engineering in plants. Plant Communications 2, 100168. https://doi.org/10.1016/j.xplc.2021.100168

Zhou, J., Wang, G., Liu, Z., 2018. Efficient genome editing of wild strawberry genes, vector development and validation. Plant Biotechnology Journal 16, 1868–1877. https://doi.org/10.1111/pbi.12922