A Two-Component Regulatory System Mediates Quorum Sensing–Dependent Morphology and Motility Transitions in the Archaeon *Haloferax volcanii*

Jacob A. Cote[1], Priyanka Chatterjee[1], Marco Garcia[2], Ran Tao[3], Arnold Mathijssen[3], and Mechthild Pohlschroder[1]*

[1]Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania 19104
[2]Department of Biology, University of Pennsylvania, Philadelphia, Pennsylvania 19104
[3]Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, Pennsylvania 19104

Corresponding author: Mechthild Pohlschröder

Email: pohlschr@sas.upenn.edu

40
ABSTRACT

Quorum sensing (QS) enables microorganisms—including bacteria, eukaryotes, and

viruses—to coordinate collective behaviors in response to population density. Despite their

ecological and evolutionary significance, QS mechanisms in Archaea remain poorly

characterized. The halophilic archaeon *Haloferax volcanii* provides a model for archaeal QS,

transitioning from motile rods to non-motile disks in a density-dependent response to a

secreted disk-forming signal (DFS). To identify components of the DFS regulatory network,

we screened for spontaneous mutants that retained motility in DFS-containing soft-agar

medium. One candidate, *HVO_1357*, encodes a predicted response regulator located

adjacent to a histidine kinase (*HVO_1356*) and a second response regulator (*HVO_1358*),

consistent with an extended two-component regulatory system (TCS). Based on our results,

these genes encode quorum-sensing associated regulators (Qar), therefore, we propose

rename them *qarA* (*HVO_1357*), *qarB* (*HVO_1356*), and *qarC* (*HVO_1358*). Deletion of *qarA*

enabled cells to swim on DFS-containing soft-agar plates and conferred hypermotility on

standard soft-agar media; however, these phenotypes were not due to changes in motility-

related parameters, but a reduced sensitivity to DFS for induction of the non-motile, disk-

shaped state. In contrast, Δ*qarB* and Δ*qarC* strains were non-motile and exhibited

premature disk formation during normal growth. Suppressor mutations that restored

motility to Δ*qarB* and Δ*qarC* mapped exclusively to *qarA*, suggesting QarA is the central

regulator of this system. Transcriptomic analyses revealed that *qarA* deletion leads to

upregulation of genes involved in motility and rod-shape formation. Together, these

findings reveal *qarABC* as a DFS-responsive regulatory module and represent the first TCS

in archaea shown to control QS-dependent behavior.

64
IMPORTANCE

Archaea are ubiquitous and play key roles across diverse ecosystems—including

human microbiomes—yet little is known about how they communicate with one another

and with other organisms, or how these interactions shape their ecological impact. Such

intercellular communication, including quorum sensing (QS), allows microorganisms to

coordinate behaviors critical for survival, adaptation, and community organization. In this

study, we identify the first archaeal two-component regulatory system that is involved in

QS-dependent regulation, providing a foundation for understanding how organisms in this

domain sense and respond to population cues. By revealing a previously unknown aspect

of archaeal biology, this work represents an important step toward understanding how

archaeal communication shapes both their physiology and their interactions within

complex microbial communities.

77
78
79

INTRODUCTION

Microorganisms rely on intercellular communication to coordinate behaviors that promote survival, competition, and adaptation in fluctuating environments. One of the best-characterized mechanisms of microbial communication is quorum sensing (QS), a process that enables density-dependent regulation of gene expression through the production and detection of secreted signaling molecules. When signal concentrations reach a threshold, cells alter transcriptional programs to coordinate behaviors like bioluminescence, biofilm formation, competence, motility, and sporulation (1-4). In gram-negative bacteria, QS often involves acyl-homoserine lactones (AHLs), exemplified by the LuxI/LuxR system in *Vibrio* species, where LuxI synthesizes AHLs that are bound by LuxR, which, once a threshold level is exceeded, activates QS-regulated genes (5). In contrast, gram-positive bacteria typically use secreted peptides as signals, and two-component regulatory systems (TCS) as signal transduction and response pathways (6). QS-like mechanisms have also been identified in fungi and viruses (7, 8), but in archaea quorum sensing remains largely unexplored, representing a major gap in our understanding of microbial communication.

Archaea possess unique physiologies and enzymes with considerable biotechnological potential (9, 10) and, due to their close evolutionary relationship to eukaryotes, serve as powerful models for studying fundamental cellular processes and the origins of eukaryotic biology (11-13). While archaea are recognized as ubiquitous members of environmental microbiomes that play critical roles in global biogeochemical cycles (14-17), they are also increasingly detected in human and animal microbiomes, where their ecological functions and impacts on host health remain poorly understood (18, 19). Despite these remarkable features, archaea remain deeply understudied compared to their bacterial and eukaryotic counterparts. Processes such as quorum sensing and other forms of intercellular signaling may provide mechanisms for communication within archaeal communities in microbiomes, as well as for interdomain communication with bacteria and eukaryotes (20). Understanding the molecular signals and regulatory pathways that mediate these interactions is therefore essential for revealing the broader ecological and physiological roles of archaea across diverse environments.

To date, QS-like behaviors have been reported in only a few archaea. These include *Natrialba magadii*, which exhibits growth phase–dependent Nep protease secretion (21); *Halorubrum lacusprofundi*, where culture supernatants enhance biofilm formation (22, 23); and *Methanosaeta harundinacea*, in which carboxy-AHLs are proposed to mediate density-dependent morphological changes (24). The authors of the latter study also identified a TCS that might be involved in the quorum sensing regulatory pathway, but in vivo data have not yet supported this, potentially due to the challenges to genetically manipulate this archaeon (25). In contrast, the model archaeon *Haloferax volcanii* combines a recently

identified quorum sensing–like behavior (20) with genetic tractability and facile growth
conditions, making it an ideal platform to study archaeal intercellular signaling (20, 26).
In *Hfx. volcanii*, cell morphology changes in a density-dependent manner: low-
density cultures are composed of motile, rod-shaped cells, whereas high-density cultures
consist of non-motile, pleomorphic disk-shaped cells (27-30). This transition can be
artificially induced at low cell densities by adding cell-free conditioned medium (CM) (20),
a hallmark of quorum sensing. Work done by Chatterjee *et al.* found that these cell shape
transitions are mediated by a secreted, yet-to-be-defined small molecule "disk-forming
signal" (DFS) (20). This study also identified two *Hfx. volcanii* proteins, DdfA and CirA, as
potential components of the DFS regulatory network (20). DdfA (disk-determining factor
A) is predicted to function as a shape-associated regulatory protein, as Δ*ddfA* cells remain
persistently rod-shaped (31). CirA, a KaiC-like signal transduction protein (32), shows a
similar phenotype, with Δ*cirA* cells also remaining rod-shaped (33). CirA has additionally
been linked to archaella gene regulation (33) and is more abundant in cells exposed to CM
(20). Proteomic data also identified a set of putative TCS proteins that may participate in
DFS sensing, analogous to the TCS-based quorum sensing systems seen in gram-positive
bacteria (20). One of these proteins is HVO_1356, a sensor histidine kinase encoded near
two response regulators: HVO_1357 and HVO_1358. Notably, mutations in *HVO_1357* were
previously identified in hypermotile isolates by Collins *et al.* (34)*,* suggesting these proteins
may play a role in regulating motility-related responses to DFS.
TCS are among the most versatile and widespread signaling mechanisms in
prokaryotes, enabling cells to sense and respond to environmental and intercellular cues
through reversible phosphorylation cascades. Each system comprises a sensor histidine
kinase, which autophosphorylates at a conserved histidine residue, and a response
regulator that receives the phosphoryl group at a conserved aspartate in the receiver (REC)
domain to modulate its activity (35, 36). The histidine kinase detects various signals—
either directly or indirectly—and adjusts the phosphorylation state of the response
regulator by balancing its kinase and phosphatase activities (35-37). The response
regulator in turn orchestrates cellular functions depending on the type of output domain it
contains (37). In *E. coli*, as many as 30 TCS have been experimentally defined (38, 39). In
contrast, archaeal TCSs remain poorly understood, with only a few experimental studies
reported to date (25, 40, 41). Bioinformatic analyses, however, reveal striking differences
between bacterial and archaeal TCS components. In bacteria, sensor kinases are typically
membrane-bound, response regulators commonly possess DNA-binding domains, and
kinases and regulators generally occur in pairs (42). In archaea, by contrast, sensor kinases
are often cytosolic, response regulators frequently lack output domains, and kinases
commonly outnumber response regulators in the genome (42). These distinctions suggest
that archaeal TCSs may have evolved fundamentally different signaling architectures (42),
underscoring the need for further investigation into these systems.

In this study, we used phenotypic screens and reverse genetics approaches to
expand our understanding of the DFS regulatory network. Our results show that *HVO_1356*,
HVO_1357, and *HVO_1358* form a regulatory module that controls motility and cell-shape
responses to DFS. HVO_1357 functions as the central regulator, promoting rod-to-disk
shape transitions in response to DFS. In contrast, HVO_1356 and HVO_1358 are essential
for motility and appear to modulate the activity of HVO_1357. Together, these results
identify HVO_1356–HVO_1358 as quorum-sensing–associated regulators (Qar) that form a
DFS-responsive two-component system mediating quorum-sensing–dependent behaviors
in *Hfx. volcanii*—the first such system linked to quorum sensing and among the earliest
characterized in Archaea.
167

RESULTS

Identification of mutants less responsive to DFS-dependent motility inhibition

We previously showed that wild-type *Hfx. volcanii* cannot swim on soft-agar plates
containing conditioned media (CM), whereas knockout strains of *cirA* and *ddfA* can (20). In
an attempt to identify additional proteins involved in sensing or responding to the quorum-
sensing signal in CM, we adopted a screen which previously was used to identify
hypermotile mutants (34). Mid-log liquid cultures were inoculated as a line across soft-agar
plates containing 25% CM. Under these conditions, wild-type cells were non-motile;
however, spontaneous mutants with reduced response to DFS regained motility and moved
away from the inoculation line, forming flares (Fig. 1). Cells from the edge of such flares
were isolated and re-stabbed on CM-containing plates to confirm that they are able to
swim in the presence of DFS. Mutants isolated by this procedure were then subjected to
whole-genome sequencing. Two independent rounds of this screen yielded 23 isolates, and
sequencing identified mutations in 21 of these, including disruptions in *cirA* (five isolates)
and *ddfA* (one isolate), validating that our screen can identify genes of interest in the DFS
sensing and response pathway (Table 1; Table S1).

One mutant carried a large deletion spanning *HVO_2001–HVO_2069*, which includes
genes involved in cell-shape regulation such as *cetZ5*, *volA*, and components of the *agl15*-
dependent *N*-glycosylation pathway (Table 1, Table S1) (31). Additional mutations were
identified in *cdc48a* (*HVO_2380*, AAA-type ATPase; one isolate), *HVO_2649* (one isolate),
and *HVO_2248* (two isolates)—genes previously linked to hypermotile phenotypes by
Collins *et al.* (34)—as well as in *HVO_A0350* (two isolates), *pgsA1* (*HVO_1136*, CDP-alcohol
1-architidyltransferase; one isolate), and *gltP2* (*HVO_2885*, SDF family transporter; one
isolate) (Table 1). For two isolates, we were unable to identify any mutations despite their
ability to swim in the presence of CM, likely due to variant calling challenges associated
with polyploid genomes.

In addition to the mutations described above, a final set of mutants highlighted
HVO_1357, which appeared in four independent isolates (Table 1); notably, this gene was

one of two major loci previously reported by Collins *et al.* to cause hypermotility when
disrupted (34). *HVO_1357* is predicted to encode a DNA-binding response regulator and is
structurally similar to the Bat protein of *Halobacterium salinarum* (OE3101R), which
regulates expression of the bacteriorhodopsin apoprotein (34, 43-45). Unlike Bat, which
contains a non-functional REC domain (Fig. S1) and has yet to be associated with other TCS
components, *HVO_1357* appears to function as part of a TCS as it is adjacent to two
additional TCS-associated genes—a sensor histidine kinase (*HVO_1356*) and a standalone
REC-domain response regulator (*HVO_1358*) (Fig. 2). As demonstrated below, because all
three genes in this cluster are related to quorum sensing, we propose naming them
quorum-sensing associated regulators (Qar), designating HVO_1357—which has the
strongest impact—as *qarA*, the histidine kinase HVO_1356 as *qarB*, and the response
regulator HVO_1358 as *qarC*.
Multiple sequence alignments with experimentally characterized archaeal and
bacterial regulators show QarA contains all key features of a response regulator (37),
including a conserved phosphorylatable aspartate (D67) and other characteristic motifs of
functional REC domains (Fig. S1a). In addition to the REC domain (IPR001789), QarA
contains single copies of HalX (IPR013971), GAF (IPR029016), BAT (IPR031803), and
HTH_10 (IPR007050) domains, making it a complex multi-domain response regulator (Fig.
2)(34). HalX is haloarchaea-specific and often fused to REC domains (42), while GAF
domains typically sense stimuli–such as cyclic nucleotides, small molecules, and light–and
appear to be common features of archaeal TCS proteins (42, 46). The BAT domain, of
unknown function, is usually paired with GAF and HTH_10, a DNA-binding helix-turn-helix
specific to Bat-type regulators (42). When gene disruptions in *qarA* from this study and
from Collins *et al.* (34) are mapped to these domains, most of the disruptions occur in the
BAT domain followed by the GAF and HTH domains (Fig. 2). Interestingly, no mutations
were observed in the REC or HalX domains.

QarA negatively regulates swimming and restricts motility in the presence of CM
To confirm that disruption of QarA induces hypermotility and enables swimming
motility in the presence of CM, we constructed deletion strain Δ*qarA*. Consistent with the
motility phenotypes observed by Collins *et al.* (34) and in our CM motility screens, Δ*qarA*
exhibits hypermotility on soft-agar plates (Fig. 3a) and is motile on soft-agar plates
supplemented with CM (Fig. 3b). Complementation of the deletion strain with *qarA*
expressed from pTA963, the commonly used overexpression plasmid in *Hfx. volcanii*, could
not be evaluated as the plasmid itself induces hypermotility for unknown reasons (47, 48).
Interestingly, we found pTA963 also enabled wild-type cells to swim in the presence of CM,
consistent with its ability to induce hypermotility (Fig. S3a). However, at least partial
complementation of Δ*qarA in trans* was observed on CM-containing soft-agar plates, where
overexpression of *qarA* reduced motility (Fig. S3a). Furthermore, the observed
hypermotility of the ΔqarA strain was not attributable to growth differences; the deletion

strain was similar to the wild-type upon OD_{600} measurements and upon determination of
viable cell counts (Fig. S2a,b).
To assess whether loss of *qarA* affects microscopic swimming behaviors that could
thereby influence motility on soft-agar plates (49, 50), we used live-cell tracking
microscopy to quantify various motility parameters in liquid culture (Movie S1). Consistent
with previous studies, we found both wild-type and Δ*qarA* displayed two distinct
swimming velocities around 2.2μm/s and 4.4μm/s, corresponding to forward and reverse
swimming modes respectively (Fig. 4d)(29). However, we could find no biologically
meaningful differences between wild-type and Δ*qarA* cells when comparing other
parameters such as average swimming speed (2.197μm/s vs 1.975μm/s) or motile
fractions (48.5% vs 46%) (Fig. 4b,c,d). We also attempted to quantify the average time
between directional reversals (e.g. "turns or tumbles") for both strains; although the
standard deviation was high—likely due to background noise—the mean values were
similar between wild-type and Δ*qarA* cells and consistent with previously reported values
for wild-type *Hfx. volcanii* (Fig. S4) (29). These results suggest that the increased motility of
the Δ*qarA* strain is not due to changes in the motility characteristics of any given cell while
it is in the motile phase, but rather it may have a defect in regulating when to adopt the
motile-rod versus non-motile-disk state.
253
Δ*qarA* raises the DFS threshold for motility arrest
We hypothesized that the hypermotility of Δ*qarA* in regular soft-agar medium and
its ability to swim on CM-containing plates result from altered sensing of the DFS signal,
such that mutant cells require higher DFS concentrations to slow or stop motility. This
delay in sensing allows Δ*qarA* cells to remain motile longer, with the rod-to-disk transition
occurring only later in growth or upon addition of higher DFS levels. To test this, we
monitored cell shape at different growth stages and with exogenous CM. In early-log phase,
both wild-type and Δ*qarA* cells were rod-shaped, though Δ*qarA* cells were noticeably
longer (Fig. 5a, b). By mid-log, many wild-type cells had converted to disks, whereas Δ*qarA*
cells remained largely rod-shaped (Fig. 5b). In stationary phase or with 1% (10μL/mL) CM,
Δ*qarA* cultures fully transitioned to disks (Fig 5b, c, d), thus precluding a defect in shape
transitions. This makes it likely that QarA may modulate DFS sensitivity. We quantified this
by exposing cells to low CM concentrations (0.2–4 μL/mL) and measuring early-log disk
formation. Wild-type cells responded even at 0.2–0.6 μL/mL CM, whereas Δ*qarA* required
~1 μL/mL CM to initiate disk formation, reaching wild-type levels by 2 μL/mL (Fig. 5e).
The shifted response curve demonstrates that deletion of *qarA* raises the DFS threshold
needed for shape conversion, explaining why mutants are hypermotile in standard medium
and retain motility on CM-containing plates, yet show no change in single-cell swimming
behavior.
273
QarB and QarC are required for motility and maintenance of the motile rod shape

Previous bioinformatic analyses revealed that *qarB*, *qarA*, and *qarC* are located in
the same genomic region but are not organized as an operon (34, 51, 52). *qarB* encodes a
putative cytosolic sensor histidine kinase with conserved PAS and GAF sensory domains
but no transmembrane regions, suggesting it functions as a cytosolic sensor or interacts
with other membrane-bound components for signal transduction through the cell
membrane. *qarC* encodes a standalone response regulator containing a conserved receiver
domain but lacks an output domain—an arrangement common in archaea but rare in
bacteria (42). Both *qarB* and *qarC* encode the essential phosphorylatable residues H347
and D58 respectively, as well as other residues important for histidine kinase and REC
domain functions (Fig. S1a,b)

To define their roles in motility and cell shape, deletion mutants were generated and
compared to wild type and the Δ*qarA* strain. While Δ*qarB* and Δ*qarC* showed slightly lower
OD_{600} values (Fig. S2a), the viable counts at comparable growth times remained similar.
However, motility phenotypes of these mutants were striking: both were completely non-
motile on soft agar with or without 25% CM (Fig. 3a,b). Complementation with plasmid-
borne *qarB* or *qarC* restored motility to wild-type or near wild-type levels (Fig. S3b),
confirming that both genes are essential for motility.

Microscopy imaging revealed that in early log phase, Δ*qarB* and Δ*qarC* cells were
rod-shaped, yet significantly shorter than wild type and transitioned to disk morphology
earlier in growth (Fig. 5a,b). Although still rod-shaped, live-cell microscopy showed that
these rods were non-motile under conditions where wild-type and Δ*qarA* strains were
motile (Fig. 4a,b). When grown in the presence of CM, Δ*qarB* and Δ*qarC* cultures were fully
disk-shaped (Fig. 5c,d). Together, these results show that *qarB* and *qarC* are both required
for motility and may promote the motile state in response to environmental cues such as
DFS concentration.

qarA influences transcription of motility, cell-shape, and TCS genes

To identify genes differentially transcribed in the absence of the putative DNA-
binding response regulator QarA, we compared RNA-seq profiles of wild-type and Δ*qarA*
Hfx. volcanii at early-log and late-log phases (full datasets in Tables S2–S5).

Early-log phase: In Δ*qarA*, 60 differentially expressed genes (DEGs) were identified
(Fig. 6a; Table S3). Thirteen DEGs were upregulated, including genes encoding archaella
components *arlA1* (*HVO_1210*) and *arlA2* (*HVO_1211*), motor complex components *arlH*
(*HVO_1216*) and *arlI* (*HVO_1217*), and the chemotaxis adaptor *cheF* (HVO_1219), all of
which are important for motility (33, 53). Genes encoding proteins involved in rod
formation, *rdfA* (*HVO_2174*) and *sph3* (*HVO_2175*), were also upregulated (31). *HVO_2649*,
identified in our 25% CM screen as well as the previous hypermotility screen (34), was
similarly upregulated. The most highly upregulated genes were *HVO_B0192* and
HVO_B0193, encoding a putative transmembrane protein and transcriptional regulator
with a domain of unknown function (DUF7521), respectively.

In contrast, 47 genes were downregulated in the Δ*qarA* mutant. A notable finding was the decreased expression of *qarC*. Another gene in this region, *HVO_1359*, encoding a CPxCG zinc-finger protein, was also downregulated. The most strongly downregulated gene overall was *HVO_B0276*, encoding a predicted drug/metabolite transporter (DMT) superfamily protein. Functional enrichment analysis of early-log DEGs revealed, next to motility, an overrepresentation of genes involved in signal transduction (Fig. 6c).

Late-log phase: RNA-seq analysis identified 41 DEGs in the Δ*qarA* strain, including 3 upregulated and 38 downregulated genes (Fig. 6b; Table S5). A key observation in late-log was the pronounced downregulation of *qarC* and *HVO_1359*, the latter of which was more strongly down-regulated at increased significance compared to the early-log dataset. *HVO_B0276*, the most strongly downregulated gene in early-log, is also highly downregulated in late-log but drops in relative rank, highlighting the increasing prominence of potential TCS and regulatory components in late-log DFS responses.

The remaining late-log-specific DEGs included upregulation of *HVO_2375* (encoding PstS family transporter), *HVO_2063*, and *HVO_A0133*—encoding a predicted glycine zipper protein with a TAT secretion signal previously reported as highly abundant in both CM-grown and rod-shaped cells (20, 31). Other downregulated late-log-specific genes included *HVO_A0350* (encoding TAT lipoprotein) and *HVO_A0351*. Notably, two mutants with disruptions in *HVO_A0350* were identified in our CM screen (Table 1), supporting a potential role for this region in DFS-dependent phenotypes. Interestingly, our transcriptomics also identified several genes encoding surface-associated PQQ-like β-propeller repeat proteins (*HVO_2606*, *HVO_2607*, *HVO_B0138*, *HVO_B0139*), highlighting potential changes in adhesion, extracellular interactions, or signaling among late-log DEGs. Like in the early-log dataset, functional enrichment analysis found enrichment of genes with roles in signal transduction, but now also carbohydrate metabolism and transport, likely due to surface-associated proteins like the PQQ-like β-propeller repeat proteins (Fig. 6d).

Several genes showed consistent changes across both early- and late-log datasets, suggesting they contribute to a common regulatory program in response to loss of QarA. These included *HVO_1310*, *HVO_0036 and HVO_2879* encoding a probable thioredoxin, probable transmembrane protein, and ornithine cyclodeaminase, respectively. In addition, multiple genes within or near the *HVO_2013–HVO_2020* locus, whose proteins were previously implicated in cell shape determination, were differentially expressed (31). Specifically, *HVO_2020* (encoding a membrane protein; DUF502 family) and *HVO_2024* were downregulated in both datasets, whereas *HVO_2014* and *HVO_2017*, encoding a zinc finger domain protein and a pentapeptide repeat protein, respectively, were identified only in the early-log dataset.

To validate our RNA-seq results, real-time quantitative reverse transcriptase PCR (qRT-PCR) was performed to compare transcript levels between the wild-type and Δ*qarA* strains, focusing on representative genes identified as differentially expressed in the

transcriptomic analysis. Consistent with the RNA-seq data, *arlA1* was significantly upregulated in Δ*qarA*, whereas *HVO_B0276* was downregulated (Fig. 7a). Expression of *rdfA* also showed an overall increase in Δ*qarA* (mean relative expression = 1.32); however, variability among biological replicates was high under the tested conditions, and the difference was not statistically significant. Attempts to quantify *qarC* expression were inconclusive due to apparent low transcript abundance (data not shown). Given that Δ*qarC* displays opposite phenotypes to that of Δ*qarA*, and that deletion of *qarA* restores motility in the Δ*qarC* background, we also examined the expression of these same genes in Δ*qarC* to assess potential reciprocal regulation. As expected, *arlA1* and *rdfA* were downregulated in Δ*qarC*, with *arlA1* showing little to no expression, consistent with its non-motile phenotype (Fig. 7b). *HVO_B0276* exhibited a non-significant increase in expression (mean relative expression = 1.4). Strikingly, *qarA* transcript levels were elevated approximately 4 times in Δ*qarC* relative to the wild type. Transcriptomic observations that QarA promotes *qarC* expression, and that *qarA* is overexpressed in cells lacking *qarB* are consistent with spontaneous motile suppressors of Δ*qarB* and Δ*qarC* we occasionally observed emerging from both non-motile strains (Fig S5). Whole-genome sequencing of three suppressors from both deletion strain backgrounds revealed mutations in *qarA* (Table S1), suggesting that QarA is functionally linked to QarB and QarC as its possible partners in this TCS pathway.

DISCUSSION

In this study, we developed a genetic screen to identify components of the DFS-mediated QS network in *Haloferax volcanii*. Through reverse genetics, microscopy, and transcriptomic analyses, we uncovered a three-gene signaling module—*qarABC*—that regulates cell shape and motility in response to DFS. Bioinformatic predictions indicate that *qarA* encodes a multidomain DNA-binding response regulator, *qarB* a cytosolic histidine kinase, and *qarC* a standalone REC-domain protein, all together forming a TCS-like signaling architecture (Fig. 2, Fig. S1).

Functional analyses revealed that QarA acts as the principal effector of this pathway. Deletion of *qarA* caused a pronounced hypermotile phenotype (Fig. 3a) that does not arise from altered swimming behavior (Fig. 4) but rather from a shift in the DFS-dependent rod-to-disk morphological transition (Fig. 5b). Δ*qarA* cells require substantially higher DFS levels to convert to disk-shaped cells compared to wild-type cells, maintaining a larger fraction of rod-shaped, motile cells (Fig. 5e). This reduced DFS sensitivity likely explains the hypermotility observed on soft-agar plates: local DFS accumulation within the halo is less effective at inducing the non-motile state, leaving a higher fraction of motile cells and thereby increasing halo migration compared to wild type. Similarly, the DFS concentration present in 25% CM soft agar plates is likely below the threshold needed to fully convert Δ*qarA* cells to non-motile disks. Consistent with our view that QarA functions as a motility repressor, plasmid-based overexpression of QarA not only reduced the ability of Δ*qarA* to

swim in the presence of DFS but also reduced motility below the plasmid-bearing wild-type
strain (Fig. S3a).

In contrast to the hypermotile Δ*qarA* strain*,* deletion of *qarB* or *qarC* resulted in
non-motile short rod-shaped cells (Fig. 3a; Fig. 4) that also transitioned prematurely to
disk-shaped cells (Fig. 5b). These reciprocal phenotypes, together with the observation that
motile suppressors of Δ*qarB* and Δ*qarC* carry secondary mutations in *qarA* (Table S1)
support a model in which QarB and QarC act upstream to inhibit QarA activity under low-
DFS conditions.

We propose that QarABC form a DFS-responsive signaling cascade that governs the
transition between motile and sessile states (Fig. 8). Based on their predicted domain
architectures, QarB and QarC likely function as a histidine kinase–response regulator pair,
forming the upstream module of a two-component–like system. Under low DFS conditions,
QarB and QarC maintain the motile, rod-shaped state by repressing QarA activity. As DFS
accumulates, QarB and QarC become inactivated, relieving repression of QarA. Activated
QarA then downregulates rod-associated genes such as *rdfA*, *sph3*, and *arlA1*—which are
upregulated in the Δ*qarA* mutant (Fig. 6a, 7a; Table S3)—thereby promoting conversion to
the non-motile, disk-shaped state.

Consistent with this model, the abundance of QarB decreases under high-DFS
conditions (20), suggesting that QarB becomes inactive as DFS levels rise, permitting
activation of QarA. Because QarC lacks a canonical output domain (Fig. 2), it is plausible
that it transmits this regulatory signal to QarA through direct protein–protein interactions
rather than through transcriptional control.

Interestingly, we also observed that *qarA* expression is elevated in Δ*qarC* cells (Fig.
7b), suggesting an additional regulatory link between QarC and QarA . This could reflect
indirect control via other transcriptional regulators of *qarA*, or possibly autoregulation by
QarA itself, a mechanism commonly observed among response regulators (36). If QarA
indeed activates its own expression, this feedback could amplify its activity once repression
by QarB/C is lost. Such a model would explain the reciprocal phenotypes between the
Δ*qarB*/Δ*qarC* strains and Δ*qarA*: in both cases, premature QarA activation—driven by loss
of upstream inhibition and potential positive feedback—would promote early transition to
the non-motile, disk-shaped state even under low DFS conditions. Notably, QarA also
promotes expression of *qarC* (Fig. 6a,b), forming a regulatory loop in which QarA activates
its own repressor—an arrangement that may serve to fine-tune or stabilize the transition
between motile and non-motile states.

Finally, although QarA contains a conserved REC domain suggestive of
phosphorylation, we detected no mutations within this region in our suppressor analyses.
Thus, the functional significance of QarA phosphorylation remains unresolved, but it may
represent an additional regulatory layer within the DFS-responsive QarABC signaling
cascade.

Transcriptomic profiling also revealed that QarA governs a broader regulatory
network linking DFS signaling to motility, morphology, and small-molecule transport.
Among the most strongly downregulated genes in ΔqarA was *HVO_B0276* (Fig. 6a,b; Table
S3,5), encoding a predicted DMT superfamily transporter previously identified as a
regulatory target of the metabolic regulator TbsP, which indirectly influences cell-shape
transitions in *Hfx. volcanii* (54). Given that the mechanism of DFS secretion remains
unknown, *HVO_B0276* represents a compelling candidate for mediating DFS trafficking.

Two additional genes, *HVO_B0192* and *HVO_B0193*, were strongly upregulated in the
qarA knockout during early-log phase (Fig. 6a). Both genes are repressed by the cell
division regulator CdsR, suggesting that loss of QarA disrupts early shape-related
transcriptional programs (55). This finding supports the idea that the QarABC system
functions early in growth to sense and respond to low levels of accumulating DFS. It is
likely that at higher cell density surrounding nutrients are limited and the disk shape
promotes better nutrient uptake. Thus, the QarABC system may allow cells to initiate the
rod-to-disk transition ensuring they are fully disk-shaped by later growth phases when
nutrients become limiting. Consistent with this model, ΔqarB and ΔqarC mutants form
short rods during early-log phase (Fig. 5a) but are non-motile (Fig. 3a; Fig. 4), suggesting
that the shape-transition program has already been triggered—suppressing motility before
morphological conversion to disks is complete.

Beyond these DFS pathway candidates, we also identified a consistent group of
differentially expressed genes across both early- and late-log datasets (Fig. 6a,b), including
HVO_1310, HVO_0036, and *HVO_2879,* encoding a probable thioredoxin, predicted
transmembrane protein,and ornithine cyclodeaminase, respectively, and the *HVO_2013–*
HVO_2020 region, which encodes multiple shape-related factors in *Hfx. volcanii* (31).
Another intriguing candidate is *HVO_1359*, located immediately downstream of *qarABC* and
is co-transcribed with *qarC* and *HVO_1360* (Fig. 2) (51, 52). Although *HVO_1359* shows
modest differential regulation in early-log phase, it becomes one of the most strongly
downregulated genes in late-log phase, suggesting a link to DFS accumulation (Fig. 6a,b).
Consistent with this, *HVO_1359* abundance is elevated in cells grown in the presence of high
DFS concentrations (20). *HVO_1359* encodes a small CPxCG-type zinc finger protein, part of
a conserved but poorly understood family implicated in haloarchaeal stress and signaling
regulation (56-58). Notably, *HVO_1359* shares homology and genomic context with the Brz
protein of *Hbt. salinarum*, located adjacent to the Bat regulator, with which it is thought to
function as a co-regulator of bacterio-opsin transcription (59, 60). This connection suggests
that *HVO_1359* may also act as a co-regulator with QarA in DFS signaling. Moreover, zinc
finger proteins and AAA-type ATPases, such as *cdc48a* identified in our CM QS bypass
screen (Table 1), frequently co-occur with Bat-type proteins in haloarchaeal genomes (60,
61), further suggesting that *HVO_1359* and *cdc48a* may also be components of the DFS
regulatory network.

The overall findings described in this work define a DFS-responsive signaling
module that integrates environmental and community-derived cues to coordinate
collective behavior in *Hfx. volcanii*. The QarABC system represents the first experimentally
characterized TCS in Archaea with demonstrated roles in QS, exhibiting parallels to both
bacterial architectures and archaeal-specific innovations such as standalone REC-domain
proteins.

Future studies dissecting the mechanisms of DFS sensing, QarA signaling, and DFS
production will deepen understanding of archaeal QS-like systems and their possible
conservation across lineages. While the QarABC system initiates early shape transitions,
the ability of Δ*qarA* cells to form disks at high DFS levels indicates additional pathways act
downstream. Together, these directions will illuminate how archaeal communities use
small-molecule signaling to regulate collective behaviors, for which the foundations have
been laid in the current study.

METHODS

Plasmids, primers, and strains. The plasmids, primers, and strains generated through
molecular cloning are listed in Table S6. Summarized sequencing results of strains
generated through motility screens can be found in Table S1.

Growth conditions. *Hfx. volcanii* strains were grown at 45 °C in liquid (orbital shaker,
250 rpm, 2.54-cm orbital) or on solid semi-defined Casamino Acids (Hv-Cab) agar (1.5%
[wt/vol]) medium (27), supplemented with tryptophan and uracil (50 µg mL^{-1} each). Uracil
was omitted for strains carrying pTA963-based plasmids. Solid plates were incubated in
sealable bags to prevent drying. *Escherichia coli* strains were grown at 37 °C in NZCYM
medium, either in liquid or solid agar (1.5% [wt/vol]) supplemented with ampicillin
(100 µg mL^{-1}). Growth of liquid cultures was monitored via OD$_{600}$ (Thermo Spectronic
20+).

Preparation of conditioned medium (CM) and CM-agar plates. CM and CM-agar plates
were prepared as previously described (20). Briefly, a 1L culture of *Hfx. volcanii* was grown
to OD$_{600}$ 1.5 and cells pelleted by centrifugation (Beckman Coulter Avanti J-25I, JA-10 rotor,
8,000 x *g*, 30 min). The resulting supernatant (CM) was then filtered (Millipore Stericup,
0.22 µm PVDF membrane) and stored at −20°C. Soft agar motility plates contained 0.35%
[wt/vol] agar of Hv-Cab medium supplemented with CM to a final concentration of 25%.

Conditioned media motility screens. Colonies of wild type *Hfx. volcanii* were inoculated
into 5 mL liquid medium and grown at 45 °C with continuous shaking until mid-log phase
(OD$_{600}$ ~0.5). Isolates from the MG series (Table S1) were stab-inoculated halfway into the
agar with a sterile toothpick, then dragged horizontally within the agar to form a groove.

For SB series isolates (Table S1) , 20 µL culture was spotted onto CM soft-agar plates and a
line streaked across the surface with a sterile toothpick. One or two lines were streaked per
plate, followed by incubation at 45 °C for 4–5 days in a plastic box with wet paper towels,
changed daily, to prevent drying. Cells from the edge of well-separated outgrowth flares
were sampled with sterile toothpicks and streaked on Hv-Cab plates to isolate individual
suppressor strains. Colonies were re-stabbed on 25% CM soft-agar plates to confirm the
CM-swimming phenotype. Plates were imaged with the lid removed and photographing
from above or using a document scanner.

Whole-genome sequencing of deletion and suppressor strains. Genomic DNA was
extracted using the GeneJET Genomic DNA Purification Kit (Thermo Scientific) and
quantified with a Qubit 3.0 Fluorometer and BR dsDNA kit (Invitrogen). Illumina whole-
genome sequencing and variant calling were performed by SeqCenter (Pittsburgh, PA,
USA). Libraries were prepared using the tagmentation-based Illumina DNA Prep kit with
custom IDT 10 bp unique dual indices and a target insert size of 280 bp. Sequencing was
performed on a NovaSeq X Plus, generating 2 × 150 bp paired-end reads. Demultiplexing,
quality control, and adapter trimming were performed with bcl-convert (Illumina) (MG4-
10 v3.9.3; MG15-33 v4.0.3; SB strains v4.1.5). Variant calling was carried out with BreSeq
(MG4-10 v0.36.1; MG15-33 v0.37.1; SB strains v0.38.1) (62) using *Hfx. volcanii* DS2
(GCF_000025685.1) as a reference. Predicted mutations were compared to the
Pohlschröder H53 lab strain (https://doi.org/10.5281/zenodo.17419175) background to
identify newly mutated genes. For suppressor strains with no predicted mutations, the
"Unassigned missing coverage," "Unassigned new junction," and marginal prediction files
were manually inspected, prioritizing DEGs identified in our RNA-seq experiments.
Manually predicted mutations were confirmed by mapping reads to *Hfx. volcanii* DS2
(GCF_000025685.1) using the Geneious mapper (Geneious Prime v.2025.2.2; structural
variants, short insertions/deletions of any size, medium/low sensitivity). Strains with no
identified mutations were further screened for large insertions using ISCompare v1.0.5
with default parameters, with none being detected (63).

Plasmid preparation and transformation. The pTA131 plasmid was used to generate
chromosomal deletions (64), and the pTA963 plasmid was used for complementation
experiments (65). To prevent restriction of methylated DNA by *Hfx. volcanii*, plasmids were
first transformed into *E. coli* NEB 5-alpha, then into homemade *E. coli* DAM⁻ strain DL739,
before transformation into *Hfx. volcanii*. Plasmid preparations for both *E. coli* strains were
performed using the PureLink Quick Plasmid Miniprep Kit (Invitrogen). *Hfx. volcanii*
transformations were carried out using the polyethylene glycol method (66), and *E. coli*
transformations by heat shock (67).

Generation of chromosomal deletions in *Hfx. volcanii*. Chromosomal deletions were generated using the pop-in/pop-out method following transformation with pTA131-based plasmids, as previously described (64). Constructs for deletion (pJC1, pJC2, pJC3) were generated by amplifying ~750 nt flanking each gene via PCR, followed by overlap PCR (primer sets 3–4, 7–8, 11–12). The fused upstream and downstream fragments were cloned into pTA131 at XbaI and XhoI sites. Following transformation and selection, deletions were confirmed by colony PCR using primers flanking and internal to the deleted gene (primer sets 5–6, 9–10, 13–14). Deletion strains were further validated by whole-genome sequencing and variant calling to confirm complete gene deletion and assess secondary genome alterations.

Construction of complementation plasmids. The *HVO_1356*-StrepII expression plasmid pJC4 was generated by PCR amplification of *HVO_1356* from *Hfx. volcanii* genomic DNA using primer set 15 and cloning into pTA963. The corresponding complementation plasmid, *HVO_1356*-StrepII-Myc (pJC6), was created by adding a Myc tag to pJC4 via site-directed mutagenesis (New England BioLabs Q5 kit) with primer set 16. Similarly, *HVO_1357*-StrepII (pJC5) was amplified with primer set 17 and cloned into pTA963, and the complementation plasmid *HVO_1357*-StrepII-FLAG (pJC7) was obtained by introducing a FLAG tag to pJC5 using site-directed mutagenesis with primer set 18. HA-StrepII-*HVO_1358* (pJC8) was synthesized by Twist Bioscience (San Francisco, CA, USA).

Growth curves. Colonies of each strain were inoculated into 5 mL liquid medium and grown at 45 °C with continuous shaking until mid-log phase (OD_{600} 0.3–0.5). Cultures were diluted to OD_{600} 0.01 in 200 μL fresh medium per well of a non-treated, flat-bottom 96-well plate (Corning). Growth was measured using a BioTek Epoch 2 microplate reader with Gen6 software (Agilent v1.04). Four biological replicates were used per strain. To minimize evaporation, empty wells were filled with 200 μL water, and a 2 °C gradient was applied to the 45 °C incubator setting. Readings were taken every 30 min at 600 nm, with continuous double-orbital shaking (335 rpm, 4 mm) between measurements, for a total of 72 h.

Determination of viable cell counts. Cultures for each strain were grown to mid-log phase (OD_{600} 0.3–0.5). Cultures were diluted to OD_{600} 0.3 and serially ten-fold diluted. 30μL of 10^{-3} to 10^{-5} dilutions were spread onto Hv-Cab solid agar plates and incubated for 3-5 days. Colonies were manually counted and reported as log_{10} CFU/mL of the original culture.

Motility assays and motility halo quantification. Motility was assessed on 0.35% [wt/vol] agar Hv-Cab medium. Colonies were inoculated into 5 mL liquid medium and grown at 45 °C with continuous shaking until mid-log phase (OD_{600} 0.3–0.5). Cultures were diluted to the lowest OD_{600} among the strains tested and stab-inoculated into the agar with

a toothpick. Plates were incubated upright at 45 °C in a plastic box with wet paper towels, changed daily, to maintain humidity. After two days, plates were removed and imaged following at least one day at room temperature to allow halos to darken. Images were acquired using a document scanner, with lids removed. Halo areas were quantified as previously described (54) using FIJI (68): the scale was calibrated to plate diameter, halos were segmented with the threshold tool, and areas measured using the Analyze Particles function.

Cell shape microscopy. *Hfx. volcanii* strains were inoculated from single colonies into 5 mL Hv-Cab liquid medium; for CM experiments, 50 µL CM (1% [vol/vol]) was added. Cultures were grown to early-log (OD_{600} 0.045–0.050), mid-log (OD_{600} 0.30), or stationary phase (72 h, OD_{600} ~1.9–3.0). For early- and mid-log cultures, 1 mL aliquots were centrifuged at 4,900 × g for 6 min and pellets resuspended in their supernatant (~5 µL for early-log, ~50 µL for mid-log). For imaging, 1.5 µL of resuspended pellet or stationary culture was placed on glass slides (Fisher Scientific) and covered with No. 1 18 × 18 mm coverslips (Globe Scientific Inc.), gently pressed to maintain a single cell layer. Images were acquired on a Leica DMi8 inverted microscope with DFC9000 GT camera using Leica Application Suite X (v3.6.0.20104). DIC and brightfield images were captured with a 100× Plan Apochromat objective (NA 1.4, WD 0.09 mm). Brightfield images for quantification were background-subtracted in FIJI (ImageJ v2.16.0/1.54p)(68) and analyzed in CellProfiler (v4.2.8)(69). The FIJI macro and CellProfiler pipeline used are available at https://doi.org/10.5281/zenodo.17419175.

CM titration microscopy. *Hfx. volcanii* strains were inoculated from single colonies into 5 mL Hv-Cab liquid medium containing 0.2, 0.6, 1, 2, or 4µL/mL CM. Cultures were grown to early-log phase (OD_{600} 0.045–0.050) and imaged and analyzed using FIJI (ImageJ v2.16.0/1.54p)(68) and CellProfiler (v4.2.8)(69) as described above. Fraction of disk-shaped cells was calculated by finding the ratio of cells with the CellProfiler eccentricity values below 0.9.

Motility chamber construction. Motility chambers were assembled using 24 × 50 mm (Fisher Scientific) and 18 × 18 mm (Corning) No. 1 coverslips. Coverslips were cleaned in a saturated KOH/ethanol bath for 30 min, rinsed with deionized water, and immersed in 1% solution of polyvinylpyrrolidone K30 containing 0.01% Tween-20 for 1 h. Coverslips were removed slowly, air-dried vertically, and baked at 120 °C for 1 h. Chambers were constructed by placing two double layers of parafilm (~5 × 25 mm) ~8 mm apart on a large prepared coverslip on a 95 °C hotplate, followed by covering them with a small prepared coverslip to form an ~8 × 18 mm chamber with a height of ~200 µm.

Live-cell tracking and analysis. Cultures were grown to early-log phase (OD_{600} 0.045–0.050) and diluted 1:5 (wild type, Δ*qarA*) or 1:2 (Δ*qarB*, Δ*qarC*) in pre-heated (45°C) motility buffer (Hv-Cab medium with 1% [wt/vol] BSA). Motility chambers were partially filled (~2/3) from one end with pre-heated motility buffer to further passivate glass surfaces, then fully filled with diluted cells. Chamber ends were sealed with liquid VALAP wax and imaged immediately. Swimming behavior was recorded on a Nikon TI2-E microscope with a 20× CFI60 Plan Apochromat Lambda objective (NA 0.75, WD 1.0 mm) under brightfield illumination. Videos were captured at 5 fps for 5 min using a Hamamatsu Orca-Fusion Gen-III sCMOS camera using NIS-Elements (6.10.01). Cells were tracked near the bottom coverslip. Videos were processed in FIJI, and trajectories were automatically analyzed using a custom macro with TrackMate plugins (70, 71). For determination of non-motile vs motile cells, $1\mu m$/s was used as a threshold based on observations of non-motile objects in our motility chambers.

RNA extraction and purification. Liquid cultures (5 mL) of each strain were grown at 45 °C with shaking to OD_{600} 0.05 (early-log) or 0.9 (late-log). Early-log (4–5 mL) and late-log (0.5 mL) cultures were harvested and pelleted at 4,900 × g for 6 min. RNA was extracted using the Qiagen RNeasy Plus Micro Kit, with lysates homogenized via QIAshredder columns. For late-log RNA-seq and qRT-PCR samples, DNase digestion was performed using the RNase-free DNase kit (Qiagen) followed by RNeasy MinElute Cleanup (Qiagen). RNA concentration was measured with Qubit RNA Broad Range Assay Kits (Invitrogen), and integrity assessed using RNA ScreenTape on a 4200 Tapestation (Agilent). All RNA-seq and qRT-PCR samples had RINe values above 8.9 and 7.0, respectively.

RNA sequencing and data analysis. Cell pellets were collected and RNA extracted as described previously. For early-log and late-log samples, four and three biological replicates were collected, respectively. Purified RNA was sent to SeqCenter LLC (Pittsburgh, PA, USA) for library preparation and sequencing. RNA was DNase-treated (Invitrogen, RNase-free) prior to library preparation using Illumina Stranded Total RNA Prep Ligation with Ribo-Zero Plus kit and 10 bp unique dual indices. Additional rRNA depletion was performed with *Hfx. volcanii*-specific probes from Rados *et al.* (72). Sequencing was performed on a NovaSeq X Plus with 2 × 150 bp paired-end reads. Demultiplexing, quality control, and adapter trimming were done with bcl-convert (v4.2.4). Paired-end RNA-seq reads were quality-checked, rRNA/tRNA reads were removed, and the remaining reads were aligned to the *Hfx. volcanii* DS2 reference genome (GCF_000025685.1) using minimap2 (73), with alignments processed using samtools (74). Gene-level counts were obtained with featureCounts from the Rsubread R package (75). Differential expression between Δ*qarA* and wild type was analyzed using DESeq2 (76); genes with <10 total counts were excluded. Log_2 fold changes were estimated with the Wald test and shrinkage-adjusted using apeglm (77). Gene annotations, including product

descriptions, arCOG assignments, and COG functional classes, were obtained from RefSeq and locus tag reference tables. Genes with adjusted p-values (Benjamini–Hochberg) <0.01 and $|\log_2 FC| > 0.58$ were considered differentially expressed. PCA and volcano plots were generated with ggplot2 and plotly to visualize sample clustering and differential expression. Functional enrichment of significant DEGs was assessed by comparing the distribution of gene function categories in each DEG set to the genome-wide background using Fisher's exact tests. The pipeline used to perform RNA-seq analyses can be found at https://doi.org/10.5281/zenodo.17419175

qPCR analysis. Three biological replicates of Δ*qarA* and Δ*qarC* were collected as previously described for early-log RNA samples, along with two sets of three biological replicates of wild-type controls per comparison. cDNA was synthesized from 1 µg RNA using the ProtoScript II First Strand cDNA Synthesis Kit (New England BioLabs) with random primers; parallel negative controls without reverse transcriptase (-RT) were prepared.

Primers for *Hfx. volcanii* genes *eif1a, arlA1, rdfA, HVO_1357, HVO_1358,* and *HVO_B0276* were designed using Primer3 and synthesized as RxnReady™ primers. qRT-PCR reactions (20 µL) contained 2× iTaq Universal SYBR Green Supermix (Bio-Rad), 5% DMSO, 300 nM primers, and 8 µL of 1:125 diluted cDNA. No-template (NTC) and -RT controls were included; reactions were considered acceptable if ΔCt ≤10 between sample and control. Primer efficiencies determined via standard curves ranged from 85.5–99.5%. Each reaction included three biological replicates, each with three technical replicates, run on a QuantStudio 3 Real-Time PCR System (Applied Biosystems).

Thermal cycling conditions were: 94 °C for 2 min, then 40 cycles of 95 °C for 15 s and 60 °C for 30 s (*eif1a*, *HVO_1357*, *HVO_B0276*) or 58 °C for 30 s (*arlA1*, *rdfA, HVO_1358)*. SYBR Green fluorescence was recorded at each cycle's end, and the default melt curve program following the qPCR run verified primer specificity. Relative expression was calculated in Excel using the Pfaffl method (78), normalizing to *eif1a*, and compared to wild-type controls to determine differential expression.

DATA AVAILABILITY

The raw RNA-seq data generated in this study have been deposited in the NCBI Sequence Read Archive (SRA) under BioProject accession number PRJNA1347779. Additional materials are available at https://doi.org/10.5281/zenodo.17419175, including: whole-genome sequencing and variant-calling data for mutant, knockout, and background strains; images, scripts, and processed data for cell-shape analyses; images and data for soft-agar motility assays; live-cell tracking data; pipelines and key outputs from RNA-seq analyses; and raw and processed qPCR data.

ACKNOWLEDGEMENTS

We acknowledge Kiara Escudero, Shalaya Brown, and Sophia Sun for their contributions to phenotypic assays and CM screens, respectively. We also thank Dr. Alexandre Bisson for his guidance with RNA-seq analysis and for providing rDNA-depletion probes. We gratefully acknowledge Dr. Friedhelm Pfeiffer, Dr. Fevzi Daldal, Dr. Yirui Hong, and Everlyne Mutua for their insightful feedback on the manuscript. This work was supported by the National Science Foundation (UPenn NSF-MBC2222076) and the University of Pennsylvania Research Fund. J.C. was supported by the National Science Foundation Graduate Research Fellowship. P.C. was supported by a National Institutes of Health Training Grant (UPenn T32 GM007229) as well as a National Institutes of Health Ruth L. Kirschstein National Research Service Award (1F31AI181536-01). M.G. was supported by a Merck/PennFERBS fellowship. A.M. acknowledges funding from the Charles E. Kaufman Foundation (Early Investigator Research Award KA2022-129523; New Initiative Research Award KA2024-144001) and the National Science Foundation (UPenn MRSEC DMR-2309043).

REFERENCES

1. Miller MB, Bassler BL. 2001. Quorum Sensing in Bacteria. Annual Review of Microbiology 55:165–199.
2. Bassler BL, Losick R. 2006. Bacterially Speaking. Cell 125:237–246.
3. Reading NC, Sperandio V. 2006. Quorum sensing: the many languages of bacteria. FEMS Microbiology Letters 254:1–11.
4. West SA, Winzer K, Gardner A, Diggle SP. 2012. Quorum sensing and the confusion about diffusion. Trends in Microbiology 20:586–594.
5. Soto-Aceves MP, Diggle SP, Greenberg EP. 2023. Microbial Primer: LuxR-LuxI Quorum Sensing. Microbiology 169.
6. Giannakara M, Koumandou VL. 2022. Evolution of two-component quorum sensing systems. Access Microbiology 4.
7. Hornby JM, Jensen EC, Lisec AD, Tasto JJ, Jahnke B, Shoemaker R, Dussault P, Nickerson KW. 2001. Quorum Sensing in the Dimorphic Fungus*Candida albicans* Is Mediated by Farnesol. Applied and Environmental Microbiology 67:2982–2992.
8. Erez Z, Steinberger-Levy I, Shamir M, Doron S, Stokar-Avihail A, Peleg Y, Melamed S, Leavitt A, Savidor A, Albeck S, Amitai G, Sorek R. 2017. Communication between viruses guides lysis–lysogeny decisions. Nature 541:488–493.
9. Pfeifer K, Ergal İ, Koller M, Basen M, Schuster B, Rittmann SKMR. 2021. Archaea Biotechnology. Biotechnology Advances 47:107668.
10. Aparici-Carratalá D, Esclapez J, Bautista V, Bonete M-J, Camacho M. 2023. Archaea: current and potential biotechnological applications. Research in Microbiology 174:104080.
11. Eme L, Spang A, Lombard J, Stairs CW, Ettema TJG. 2017. Archaea and the origin of eukaryotes. Nature Reviews Microbiology 15:711–723.

12.   López-García P, Moreira D. 2020. The Syntrophy hypothesis for the origin of
eukaryotes revisited. Nature Microbiology 5:655–667.
13.   Eme L, Tamarit D, Caceres EF, Stairs CW, De Anda V, Schön ME, Seitz KW,
Dombrowski N, Lewis WH, Homa F, Saw JH, Lombard J, Nunoura T, Li W-J, Hua Z-S,
Chen L-X, Banfield JF, John ES, Reysenbach A-L, Stott MB, Schramm A, Kjeldsen KU,
Teske AP, Baker BJ, Ettema TJG. 2023. Inference and reconstruction of the
heimdallarchaeial ancestry of eukaryotes. Nature 618:992–999.
14.   Chaban B, Ng SYM, Jarrell KF. 2006. Archaeal habitats — from the extreme to the
ordinary. Canadian Journal of Microbiology 52:73–116.
15.   Offre P, Spang A, Schleper C. 2013. Archaea in Biogeochemical Cycles. Annual
Review of Microbiology 67:437–457.
16.   Baker BJ, De Anda V, Seitz KW, Dombrowski N, Santoro AE, Lloyd KG. 2020.
Diversity, ecology and evolution of Archaea. Nature Microbiology 5:887–900.
17.   Kim J-G, Gazi KS, Awala SI, Jung M-Y, Rhee S-K. 2021. Ammonia-oxidizing archaea in
biological interactions. Journal of Microbiology 59:298–310.
18.   Bang C, Schmitz RA. 2015. Archaea associated with human surfaces: not to be
underestimated. FEMS Microbiology Reviews 39:631–648.
19.   Moissl-Eichinger C, Pausan M, Taffner J, Berg G, Bang C, Schmitz RA. 2018. Archaea
Are Interactive Components of Complex Microbiomes. Trends in Microbiology
26:70–85.
20.   Chatterjee P, Consoli CE, Schiller H, Winter KK, McCallum ME, Schulze S,
Pohlschroder M. 2025. Quorum sensing mediates morphology and motility
transitions in the model archaeon Haloferax volcanii. mBio 16:e0090625.
21.   Paggi RA, Madrid EA, D'Alessandro CP, Cerletti M, De Castro RE. 2010. Growth
phase-dependent biosynthesis of Nep, a halolysin-like protease secreted by the
alkaliphilic haloarchaeon Natrialba magadii. Letters in Applied Microbiology 51:36–
41.
22.   Charlesworth J, Kimyon O, Manefield M, Beloe CJ, Burns BP. 2020. Archaea join the
conversation: detection of AHL-like activity across a range of archaeal isolates.
FEMS Microbiology Letters 367.
23.   Liao Y, Williams TJ, Ye J, Charlesworth J, Burns BP, Poljak A, Raftery MJ, Cavicchioli
R. 2016. Morphological and proteomic analysis of biofilms from the Antarctic
archaeon, Halorubrum lacusprofundi. Scientific Reports 6:37454.
24.   Zhang G, Zhang F, Ding G, Li J, Guo X, Zhu J, Zhou L, Cai S, Liu X, Luo Y, Zhang G, Shi
W, Dong X. 2012. Acyl homoserine lactone-based quorum sensing in a
methanogenic archaeon. The ISME Journal 6:1336–1344.
25.   Li J, Zheng X, Guo X, Qi L, Dong X. 2014. Characterization of an archaeal two-
component system that regulates methanogenesis in Methanosaeta harundinacea.
PLoS One 9:e95502.
26.   Pohlschroder M, Schulze S, Pfeiffer F, Hong Y. 2025. *Haloferax volcanii*: a
versatile model for studying archaeal biology. Journal of Bacteriology 207:e00062–
25.

27.   de Silva RT, Abdul-Halim MF, Pittrich DA, Brown HJ, Pohlschroder M, Duggin IG.
794   2021. Improved growth and morphological plasticity of Haloferax volcanii.
795   Microbiology 167.

796   28.   Nußbaum P, Ithurbide S, Walsh JC, Patro M, Delpech F, Rodriguez-Franco M, Curmi
797   PMG, Duggin IG, Quax TEF, Albers S-V. 2020. An Oscillating MinD Protein
798   Determines the Cellular Positioning of the Motility Machinery in Archaea. Current
799   Biology 30:4956–4972.e4.

800   29.   Li Z, Kinosita Y, Rodriguez-Franco M, Nußbaum P, Braun F, Delpech F, Quax TEF,
801   Albers S-V. 2019. Positioning of the Motility Machinery in Halophilic Archaea. mBio
802   10:10.1128/mbio.00377–19.

803   30.   Duggin IG, Aylett CH, Walsh JC, Michie KA, Wang Q, Turnbull L, Dawson EM, Harry
804   EJ, Whitchurch CB, Amos LA, Löwe J. 2015. CetZ tubulin-like proteins control
805   archaeal cell shape. Nature 519:362–5.

806   31.   Schiller H, Hong Y, Kouassi J, Rados T, Kwak J, DiLucido A, Safer D, Marchfelder A,
807   Pfeiffer F, Bisson A, Schulze S, Pohlschroder M. 2024. Identification of structural
808   and regulatory cell-shape determinants in Haloferax volcanii. Nature
809   Communications 15:1414.

810   32.   Makarova KS, Galperin MY, Koonin EV. 2017. Proposed Role for KaiC-Like ATPases
811   as Major Signal Transduction Hubs in Archaea. mBio 8:10.1128/mbio.01959–17.

812   33.   Chatterjee P, Garcia MA, Cote JA, Yun K, Legerme GP, Habib R, Tripepi M, Young C,
813   Kulp D, Dyall-Smith M, Pohlschroder M. 2024. Involvement of ArlI, ArlJ, and CirA in
814   archaeal type IV pilin-mediated motility regulation. Journal of Bacteriology
815   206:e00089–24.

816   34.   Collins M, Afolayan S, Igiraneza AB, Schiller H, Krespan E, Beiting DP, Dyall-Smith
817   M, Pfeiffer F, Pohlschroder M. 2021. Mutations Affecting HVO_1357 or HVO_2248
818   Cause Hypermotility in Haloferax volcanii, Suggesting Roles in Motility Regulation.
819   Genes 12:58.

820   35.   Stock AM, Robinson VL, Goudreau PN. 2000. Two-Component Signal Transduction.
821   Annual Review of Biochemistry 69:183–215.

822   36.   Groisman EA. 2016. Feedback Control of Two-Component Regulatory Systems.
823   Annu Rev Microbiol 70:103–24.

824   37.   Gao R, Bouillet S, Stock AM. 2019. Structural Basis of Response Regulator Function.
825   Annu Rev Microbiol 73:175–197.

826   38.   Yamamoto K, Hirao K, Oshima T, Aiba H, Utsumi R, Ishihama A. 2005. Functional
827   Characterization in Vitro of All Two-component Signal Transduction Systems from
828   Escherichia coli*. Journal of Biological Chemistry 280:1448–1456.

829   39.   Mizuno T. 1997. Compilation of All Genes Encoding Two-component
830   Phosphotransfer Signal Transducers in the Genome of Escherichia coli. DNA
831   Research 4:161–168.

832   40.   Najnin T, Siddiqui KS, Taha, Elkaid N, Kornfeld G, Curmi PMG, Cavicchioli R. 2016.
833   Characterization of a temperature-responsive two component regulatory system
834   from the Antarctic archaeon, Methanococcoides burtonii. Scientific Reports
835   6:24278.

836   41.   Rudolph J, Tolliday N, Schmitt C, Schuster SC, Oesterhelt D. 1995. Phosphorylation
837   in halobacterial signal transduction. The EMBO Journal 14:4249–4257.
838   42.   Galperin Michael Y, Makarova Kira S, Wolf Yuri I, Koonin Eugene V. 2018. Phyletic
839   Distribution and Lineage-Specific Domain Architectures of Archaeal Two-
840   Component Signal Transduction Systems. Journal of Bacteriology
841   200:10.1128/jb.00681–17.
842   43.   Baliga NS, Kennedy SP, Ng WV, Hood L, DasSarma S. 2001. Genomic and genetic
843   dissection of an archaeal regulon. Proceedings of the National Academy of
844   Sciences 98:2521–2525.
845   44.   Gropp F, Betlach MC. 1994. The bat gene of Halobacterium halobium encodes a
846   trans-acting oxygen inducibility factor. Proceedings of the National Academy of
847   Sciences 91:5475–5479.
848   45.   Leong D, Boyer H, Betlach M. 1988. Transcription of genes involved in bacterio-
849   opsin gene expression in mutants of a halophilic archaebacterium. Journal of
850   Bacteriology 170:4910–4915.
851   46.   Heikaus CC, Pandit J, Klevit RE. 2009. Cyclic Nucleotide Binding GAF Domains from
852   Phosphodiesterases: Structural and Mechanistic Insights. Structure 17:1551–1557.
853   47.   Abdul Halim MF, Karch KR, Zhou Y, Haft DH, Garcia BA, Pohlschroder M. 2015.
854   Permuting the PGF Signature Motif Blocks both Archaeosortase-Dependent C-
855   Terminal Cleavage and Prenyl Lipid Attachment for the Haloferax volcanii S-Layer
856   Glycoprotein. J Bacteriol 198:808–15.
857   48.   Patro M, Duggin IG, Albers SV, Ithurbide S. 2023. "Influence of plasmids, selection
858   markers and auxotrophic mutations on Haloferax volcanii cell shape plasticity".
859   Front Microbiol 14:1270665.
860   49.   Wolfe AJ, Berg HC. 1989. Migration of bacteria in semisolid agar. Proc Natl Acad Sci
861   U S A 86:6973–7.
862   50.   Colin R, Ni B, Laganenka L, Sourjik V. 2021. Multiple functions of flagellar motility
863   and chemotaxis in bacterial physiology. FEMS Microbiology Reviews 45.
864   51.   Laass S, Monzon VA, Kliemt J, Hammelmann M, Pfeiffer F, Förstner KU, Soppa J.
865   2019. Characterization of the transcriptome of Haloferax volcanii, grown under four
866   different conditions, with mixed RNA-Seq. PLoS One 14:e0215986.
867   52.   Babski J, Haas KA, Näther-Schindler D, Pfeiffer F, Förstner KU, Hammelmann M,
868   Hilker R, Becker A, Sharma CM, Marchfelder A, Soppa J. 2016. Genome-wide
869   identification of transcriptional start sites in the haloarchaeon Haloferax volcanii
870   based on differential RNA-Seq (dRNA-Seq). BMC Genomics 17:629.
871   53.   Albers S-V, Jarrell KF. 2018. The Archaellum: An Update on the Unique Archaeal
872   Motility Structure. Trends in Microbiology 26:351–362.
873   54.   Hackley RK, Hwang S, Herb JT, Bhanap P, Lam K, Vreugdenhil A, Darnell CL, Pastor
874   MM, Martin JH, Maupin-Furlow JA, Schmid AK. 2024. TbsP and TrmB jointly regulate
875   gapII to influence cell development phenotypes in the archaeon Haloferax volcanii.
876   Molecular Microbiology 121:742–766.
877   55.   Liao Y, Vogel V, Hauber S, Bartel J, Alkhnbashi OS, Maaß S, Schwarz TS, Backofen R,
878   Becher D, Duggin IG, Marchfelder A. 2021. CdrS Is a Global Transcriptional
879   Regulator Influencing Cell Division in Haloferax volcanii. mBio 12:e0141621.

880   56.   Nagel C, Machulla A, Zahn S, Soppa J. 2019. Several One-Domain Zinc Finger μ-
881   Proteins of Haloferax Volcanii Are Important for Stress Adaptation, Biofilm
882   Formation, and Swarming. Genes (Basel) 10.

883   57.   McMillan LJ, Hwang S, Farah RE, Koh J, Chen S, Maupin-Furlow JA. 2018. Multiplex
884   quantitative SILAC for analysis of archaeal proteomes: a case study of oxidative
885   stress responses. Environ Microbiol 20:385–401.

886   58.   Üresin D, Schulte J, Morgner N, Soppa J. 2024. C(P)XCG Proteins of Haloferax
887   volcanii with Predicted Zinc Finger Domains: The Majority Bind Zinc, but Several Do
888   Not. Int J Mol Sci 25.

889   59.   Tarasov VY, Besir H, Schwaiger R, Klee K, Furtwängler K, Pfeiffer F, Oesterhelt D.
890   2008. A small protein from the bop-brp intergenic region of Halobacterium
891   salinarum contains a zinc finger motif and regulates bop and crtB1 transcription.
892   Mol Microbiol 67:772–80.

893   60.   Tarasov V, Schwaiger R, Furtwängler K, Dyall-Smith M, Oesterhelt D. 2011. A small
894   basic protein from the brz-brb operon is involved in regulation of bop transcription
895   in Halobacterium salinarum. BMC Mol Biol 12:42.

896   61.   Sharma AK, Walsh DA, Bapteste E, Rodriguez-Valera F, Ford Doolittle W, Papke RT.
897   2007. Evolution of rhodopsin ion pumps in haloarchaea. BMC Evolutionary Biology
898   7:79.

899   62.   Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved
900   microbes from next-generation sequencing data using breseq. Methods Mol Biol
901   1151:165–88.

902   63.   Mogro EG, Ambrosis NM, Lozano MJ. 2021. Easy identification of insertion sequence
903   mobilization events in related bacterial strains with ISCompare. G3
904   Genes|Genomes|Genetics 11.

905   64.   Allers T, Ngo H-P, Mevarech M, Lloyd RG. 2004. Development of Additional
906   Selectable Markers for the Halophilic Archaeon *Haloferax volcanii* Based on
907   the *leuB* and *trpA* Genes. Applied and Environmental Microbiology
908   70:943–953.

909   65.   Allers T, Barak S, Liddell S, Wardell K, Mevarech M. 2010. Improved Strains and
910   Plasmid Vectors for Conditional Overexpression of His-Tagged Proteins in
911   *Haloferax volcanii*. Applied and Environmental Microbiology 76:1759–1769.

912   66.   Dyall-Smith M. 2008. The Halohandbook: protocols for haloarchaeal genetics.
913   Haloarchaeal Genetics Laboratory, Melbourne 14.

914   67.   Sambrook J, Russell DW. 2006. Preparation and Transformation of Competent E.
915   coli Using Calcium Chloride. CSH Protoc 2006.

916   68.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch
917   S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,
918   Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image
919   analysis. Nature Methods 9:676–682.

920   69.   Stirling DR, Swain-Bowden MJ, Lucas AM, Carpenter AE, Cimini BA, Goodman A.
921   2021. CellProfiler 4: improvements in speed, utility and usability. BMC
922   Bioinformatics 22:433.

70.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch
S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K,
Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image
analysis. Nat Methods 9:676–82.

71.   Ershov D, Phan M-S, Pylvänäinen JW, Rigaud SU, Le Blanc L, Charles-Orszag A,
Conway JRW, Laine RF, Roy NH, Bonazzi D, Duménil G, Jacquemet G, Tinevez J-Y.
2022. TrackMate 7: integrating state-of-the-art segmentation algorithms into
tracking pipelines. Nature Methods 19:829–832.

72.   Rados T, Andre K, Cerletti M, Bisson A. 2023. A sweet new set of inducible and
constitutive promoters in Haloferax volcanii. Frontiers in Microbiology Volume 14 -
2023.

73.   Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics
34:3094–3100.

74.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
Durbin R. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics
25:2078–9.

75.   Liao Y, Smyth GK, Shi W. 2019. The R package Rsubread is easier, faster, cheaper
and better for alignment and quantification of RNA sequencing reads. Nucleic Acids
Res 47:e47.

76.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biology 15:550.

77.   Zhu A, Ibrahim JG, Love MI. 2018. Heavy-tailed prior distributions for sequence
count data: removing the noise and preserving large differences. Bioinformatics
35:2084–2092.

78.   Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time
RT–PCR. Nucleic Acids Research 29:e45–e45.

FIGURES

Figure 1. Isolation of Quorum sensing bypass mutants. Cultures were streaked on 25%
CM soft-agar plates and incubated for 4–5 days. For better feature observation, this
representative plate was stored at room temperature for one week to allow cells to darken.
(a) Non-motile cells from the initial streak. (b) Motile flare containing a spontaneous QS
bypass mutant

965
966
Figure 2. Genomic context, domains, and spontaneous mutations of *qarA*. Arrows
representing genes surrounding *qarA* are annotated with locus tag numbers above and
protein product names below. Grey arrows indicate the transcript map reported by (51);
scale bar and nucleotide numbering are shown below. The types and positions of *qarA*
mutations in this study and by Collins *et al.* (34) are indicated by symbols above *qarA* (key
right). White boxes within gene arrows denote protein domains identified by InterProScan
in the Interpro database for each gene: HVO_1356 (QarB), histidine kinase (D4GXP2);
HVO_1357 (QarA), BAT-like response regulator (D4GXP4); HVO_1358 (QarC), response
regulator (D4GXP7). Domains include GAF (IPR029016), PAS (IPR035965), HIS_KIN
(IPR005467), Receiver (IPR001789), HalX (IPR013971), BAT (IPR031803), and HTH-10
(IPR007050). Purple lines indicate approximate position of phosphorylatable residues Asp-
67 (QarA), Asp-58 (QarC), His-243 (QarB) within their Receiver or HIS_KIN domains
respectively.

Figure 3. Deletion of *qarA* causes hypermotility while Δ*qarB* and Δ*qarC* abolish motility. Cells were stab-inoculated into 0.35% (w/vol) agar plates containing Hv-Cab medium (a) or Hv-Cab medium supplemented with 25% CM (b), incubated for 2 days at 45 °C, then stored 1-3 days at room temperature before imaging. Halo area (mm^2) was normalized to wild type. Representative halo images are shown above their corresponding quantification. Scale bar represents 1cm. Error bars indicate SD from twelve biological replicates per strain over three independent experiments; dots represent individual halos values. Statistical analysis was performed using nested one-way ANOVA on non-

Figure 4. Δ*qarB* and Δ*qarC* are non-motile in early-log liquid culture while Δ*qarA* cells display motility behaviors comparable to wild type. Cells were grown to early-log phase (OD_{600} = 0.045–0.050) and diluted 1:5 (wild type, Δ*qarA*) or 1:2 (Δ*qarB*, Δ*qarC*) in pre-heated motility buffer. Diluted cultures were loaded into pre-prepared motility chambers (see Methods) and imaged immediately under brightfield illumination. Swimming behavior was recorded at 5 fps for 5 min, and trajectories were extracted in FIJI and analyzed using a custom TrackMate-based macro. This resulted in >2000 trajectories for each strain from two independent experiments. (a) Representative single-cell trajectories for each strain. (b) Fraction of motile cells (average swimming speed >1μm/s) within each population, error bars represent SD. (c) Distribution of average swimming speeds for motile strains; average speeds listed above. (d) Relative frequency of instantaneous cell velocities.

Figure 5. Δ*qarA* cells show reduced sensitivity to DFS signals. Cultures were grown to OD_{600} 0.045–0.050 (early log), 0.30 (mid log), or stationary phase (~72 h; OD_{600} 1.8–3.0). Differential interference contrast images are shown for (a) early-log cultures without additive and (c) with 1% [vol/vol] CM. Scale bar is 10μm. Cell shape quantification was performed on three biological replicates per strain in three independent experiments for each growth phase without additive (b) or at early-log with 1% [vol/vol] CM (d). Aspect ratio (major axis divided by minor axis) was calculated using CellProfiler; values < 2

(dashed line) indicate disks, and values > 2 indicate rods, with larger ratios reflecting more
elongated rods. Statistical analysis performed via nested one-way ANOVA with multiple
comparisons to wild type. ns $P > 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$. (e) Wild-
type and $\Delta qarA$ cultures grown to early-log with varying CM concentrations were analyzed
for the resulting fraction of disk-shaped cells using CellProfiler eccentricity values (foci of
ellipse-shaped object divided by major axis); values <0.9 indicate disks, and >0.9 indicate
rods. Error bars represent SD of three biological replicates per strain across three
independent experiments.

1053

Figure 6. QarA regulates expression of motility and cell-shape determinant genes. Volcano plots depict differential gene expression in *Hfx. volcanii* between Δ*qarA* and wild-type strains during (a) early-log (motile condition) and (b) late-log (non-motile condition) growth. Each point represents a single gene, with genes of interest annotated. Genes with |log$_2$ fold-change| > 0.58 and adjusted p-value (Benjamini–Hochberg) < 0.01 were considered significantly differentially expressed. *qarA* was excluded from plots to better visualize other DEGs. Bubble plots show significantly enriched COG functional categories among DEGs identified in the Δ*qarA* vs. wild type comparison during early-log (c) and late-log (d) growth. The size of each bubble corresponds to the number of DEGs in the category, and bubble color indicates enrichment significance which was determined by Fisher's exact test. Categories were considered enriched if they had an unadjusted p-value < 0.05. For significantly enriched categories, the COG function, p-value, adjusted p-value (Benjamini–Hochberg), and odds ratio are reported.

1067

1068

Figure 7. qRT-PCR analysis of Δ*qarA* confirms differential regulation of DEGs identified by RNA-seq while Δ*qarC* displays reciprocal regulation of key DEGs. Reactions were performed using primers for *eif1a*, *arlA1*, *rdfA*, *qarA*, and *HVO_B0276* under the following conditions: (a) wild type vs. Δ*qarA*, and (b) wild type vs. Δ*qarC*. Relative expression levels were calculated using the Pfaffl method, normalized to eif1a, and expressed relative to wild-type controls. Error bars indicate SD of three biological replicates, each with three technical replicates. Statistical significance was determined by $\log_2$-transforming relative expression values and performing unpaired t-tests with Welch's correction. ns P > 0.05, * P < 0.05, ** P < 0.01 *** P < 0.001.

Figure 8. Proposed model of the DFS-responsive QarABC regulatory system. RNA-seq and qRT-PCR data reveal QarA regulatory targets (solid lines), while other phenotypic, genetic, and transcriptomic observations support putative interactions (dashed lines).

1113

1114

Figure S1. Multiple sequence alignment of *Hfx. volcanii* histidine kinase and response regulator proteins (QarABC) with representative archaeal and Escherichia coli sequences. Alignments were visualized in Jalview, with shading based on the default ClustalX coloring scheme. A conservation histogram and consensus sequence are shown below each alignment. In panel (a), key catalytic residues of *E. coli* CheY (D12–D13, D57, T/S87, Y106, K109; UniProt: P0AE67) are indicated above the alignment, with residue numbering referenced to CheY. In panel (b), conserved motifs of *E. coli* EnvZ (H-box, N, G1, F, and G2 boxes; UniProt: P0AEJ4) are highlighted above the alignment, with residue numbering referenced to EnvZ. CheY D57 corresponds to D67 in QarA and D58 in QarC, and EnvZ H243 corresponds to H347 in QarB.

1125

1126

1127

1128

1129

1130 **Figure S2. Growth and viability of QarABC mutants.** (a) Growth curves of wild-type and
1131 QarABC mutants cultured in liquid medium at 45 °C for 72h. Data represent the mean and
1132 SD of four biological replicates. (b) Viable cell counts ($\log_{10}$ CFU/mL) of mid-log-phase
1133 cultures determined by serial dilution and plating on Hv-Cab agar. Error bars represent SD
1134 of five biological replicates from five independent experiments.

 Fig. S3. *qarA* **overexpression represses motility;** *qarABC* **are complementable** *in*
 trans. Cells were stab-inoculated into 0.35% (w/v) agar plates containing (a) Hv-Cab
 medium with 25% CM or (b) Hv-Cab medium alone, incubated for 2 days at 45 °C, and
 stored for 1–3 days at room temperature before imaging. Motility was quantified by (a)
 halo diameter (mm) or (b) halo area (mm^2) normalized to the wild type. Error bars indicate
 SD from (a) four biological replicates per strain or (b) twelve biological replicates across
 three independent experiments; dots represent individual halo values. Statistical analysis
 was performed using nested one-way ANOVA comparing complementation strains to their
 respective knockouts and to wild type. Non-normalized halo areas were used for analysis in
 (b). ns > 0.05, *** P < 0.001, **** P < 0.0001.

1175

1176

Fig. S4. Δ*qarA* **appears to have a similar reversal frequency to wild type.** Cells were
grown to early-log phase (OD_{600} = 0.045–0.050) and diluted 1:5. Diluted cultures were
loaded into pre-prepared motility chambers (see Methods) and imaged immediately under
brightfield illumination. Swimming behavior was recorded at 5 fps for 5 min, and
trajectories were extracted in FIJI and analyzed using a custom TrackMate-based macro.
This resulted in >2000 trajectories for each strain from two independent experiments.
Reversal frequency was determined by the average time between two subsequent >90°
turns for each tracked cell over its entire track. Error bars represent SEM.

1200

1201 **Fig. S5. Representative soft-agar plate displaying motile suppressors of Δ*qarB* and**
1202 **Δ*qarC*.** Cells were stab-inoculated into 0.35% (w/vol) agar plates containing Hv-Cab
1203 medium and incubated for 2 days at 45 °C, then stored 1-3 days at room temperature
1204 before imaging. White arrows represent spontaneous suppressor mutants from non-motile
1205 strains.

1221

1222

TABLES

1223

1224

Table 1. Genome alterations in *Hfx. volcanii* mutants less responsive to DFS-mediated motility inhibition.

1225
1226

1227

Isolate[§]	Affected Gene(s)[†]	Mutation	Position on gene	Secondary Mutation(s)[‡]	Gene Product
MG32	*HVO_1212* (*cirA*)	P168L	505/741 nt	Yes	KaiC-type circadian clock protein CirA
SB1	*HVO_1212* (*cirA*)	Δ1 bp	657/741 nt		KaiC-type circadian clock protein CirA
SB7	*HVO_1212* (*cirA*)	+1 bp	657/741 nt		KaiC-type circadian clock protein CirA
SB13	*HVO_1212* (*cirA*)	F133S	399/741 nt		KaiC-type circadian clock protein CirA
SB15	*HVO_1212* (*cirA*)	Y43D	128/741 nt		KaiC-type circadian clock protein CirA
MG6	*HVO_1357* (*qarA*)	L529P	1586/1749 nt		Receiver/bat box HTH-10 family transcription regulator
SB17	*HVO_1357* (*qarA*)	Δ261 bp	1412-1672/1749 nt		Receiver/bat box HTH-10 family transcription regulator
SB18	*HVO_1357* (*qarA*)	+46bp	1,424/1,749 nt		Receiver/bat box HTH-10 family transcription regulator
SB19	*HVO_1357* (*qarA*)	+46bp	1,424/1,749 nt		Receiver/bat box HTH-10 family transcription regulator
MG4	*HVO_2248*	Δ4 bp	803-806/816 nt	Yes	Uncharacterized protein
MG7	*HVO_2239−HVO_2298*	Δ67,424 bp			66 Genes including HVO_2248 (Uncharacterized protein)
SB2	*HVO_A0350**	K199N	597/642 nt	Yes	Uncharacterized protein
SB20	*HVO_A0350**	K199N	597/642 nt		Uncharacterized protein
MG5	*HVO_1136* (*pgsA1*)*	Δ9 bp	409-417/825 nt	Yes	CDP-alcohol 1-archaetidyltransferase
MG8	*HVO_2176* (*ddfA*)	+138bp	142/201 nt		Disk-determining factor A
MG10	*HVO_2001−HVO_2069*	Δ69,438 bp			72 genes
MG15	*HVO_2885* (*gltP2*)*	A352T	1,056/1,305 nt		SDF family transport protein (probable substrate glutamate/asparta
MG27	*HVO_2830* (*cdc48a*)	D339G	1,016/2,232 nt		AAA-type ATPase (CDC48 subfamily)
SB3	*HVO_2649*	T→A	-35 nt upstream		Uncharacterized protein

1228

[§]Isolate names starting with MG indicate those from the first screen while SB indicates
those from the second (see methods). [†]Mutations predicted by BreSeq (62) and not found
in Pohlschröder lab H53 strain, sorted by frequency. *Manual assertion using marginal
evidence for isolates where BreSeq did not predict a mutation. [‡]BreSeq-predicted
mutations in addition to primary affected gene. Full details for each isolate are provided in
Table S1. The BreSeq report for the Pohlschröder lab H53 strain background can be found
at https://doi.org/10.5281/zenodo.17419175.