Structural insights into cobalamin loading and reactivation of human methionine synthase

Douglas S. M. Ferreira[1,2], Katie McLennan[2], Calum Diamond[2], Melanie Vollmar[3,5], Wasim Kiyani[3], D. Sean Froese[3,6], Jola Kopec[3,7], Henry J. Bailey[3,8], Rod Chalk[1], Arnaud Baslé[2], Jonathan M. Elkins[1], Jesse A Coker[4], Wyatt W. Yue[1,2]*, Thomas J. McCorvie[2]*

[1]Centre for Medicines Discovery, Nuffield Department of Clinical Medicine, University of Oxford, Oxford, OX3 7DQ, UK

[2]Biosciences Institute, The Medical School, Newcastle University, Newcastle upon Tyne, NE2 4HH, UK

[3]Structural Genomics Consortium, University of Oxford, Oxford, OX3 7DQ, UK

[4]Center for Therapeutics Discovery, Lerner Research Institute, Cleveland Clinic, Cleveland, Ohio, 44195, United States.

[5]Present address: European Bioinformatics Institute (EMBL-EBI), Main Building, A2-34, Wellcome Genome Campus, Hinxton, CB10 1SD, United Kingdom

[6]Present address: Division of Metabolism and Children's Research Center, University Children's Hospital Zurich, University of Zurich, Zurich 8008, Switzerland

[7]Present address: Viska AI, Stockholm, Sweden

[8]Present address: Institute of Biochemistry II, Medical Faculty, Goethe-University, Frankfurt am Main and Buchmann Institute for Molecular Life Sciences, Frankfurt am Main, Germany

*Correspondence e-mail: wyatt.yue@newcastle.ac.uk; thomas.mccorvie@newcastle.ac.uk

Human methionine synthase (MTR) is an essential enzyme of one carbon metabolism. Consisting of a catalytic N-half and a cobalamin binding C-half, MTR utilises this intricate organometallic cofactor in the methyl transfer from methyltetrahydrofolate to homocysteine producing methionine. Cobalamin loading into MTR, and its subsequent activation, requires methylmalonic aciduria and homocystinuria Type D (MMADHC) protein and methionine synthase reductase (MTRR), respectively. However, the molecular basis of cobalamin binding and activation of human MTR aided by MMADHC and MTRR remains unknown. Here, using cryo-electron microscopy, we determined structures of human MTR in its apo, and cobalamin bound states. Apo MTR adopts a conformation where the two halves of the enzyme act independently with the C-half posed to bind cobalamin. Binding of cobalamin and its activation causes conformational changes in MTR that result in a flexible catalytically active state. AlphaFold predictions, validated by interaction studies, show that MMADHC interacts with the C-half of apo MTR to facilitate cobalamin loading. Unexpectedly we found that MTRR interacts at two distinct sites within the C-half of MTR which may aid in activation. Collectively these findings lay the groundwork to uncover the mechanisms through how MMADHC and MTRR coordinate cobalamin loading and activation of human MTR.

Introduction

Methionine   synthase   (EC   2.1.1.13;   5-methyltetrahydrofolate-homocysteine
methyltransferase, MTR) forms a crucial node connecting the methionine and folate cycles in
one-carbon metabolism. MTR catalyses the methylation of homocysteine (Hcy) using 5-
methyltetrahydrofolate (MeTHF) as the methyl group donor, to generate methionine (Met) and
tetrahydrofolate (THF)[1]. A key function of MTR is to ensure a supply of Met precursor to
synthesise the global methyl donor *S*-adenosyl-L-methionine (SAM) involved in a myriad of
epigenetic methylation reactions[2]. MTR also serves as an important regulator of the folate
cycle which carries one-carbon units for purine and pyrimidine biosynthesis[3,4]. In humans, the
primary circulating folate species is MeTHF[5], and MTR is the only enzyme that can metabolise
MeTHF to regenerate THF, the active form required for subsequent one-carbon transfer
reactions and DNA synthesis[6].

MTR is a ~140 kDa monomeric protein comprising a catalytic N-half and a cofactor
binding C-half (Fig. 1a). The N-half consists of two rigidly linked (Linker 1) TIM barrels, known
as a homocysteine (Hcy) domain and a folate (Fol) domain, which exclusively bind the
substrates Hcy and MeTHF, respectively. Connected to the N-half by a flexible linker (Linker
2), the C-half consists of a four-helical bundle Cap domain, a cobalamin binding domain (Cob),
and a SAM binding activation domain (Act)[7]. Notably, MTR is one of only two human enzymes
that uses cobalamin (Cbl) as a cofactor[7].

Cbl, also known as vitamin B_{12}, is a large 1.3 kDa molecule consisting of a centrally
bound hexacoordinated cobalt ion within a corrin ring, a ribose group tethered 5,6-dimethyl-
benzimidazole (DMB) tail that can coordinate as a lower α-axial ligand ('base-on') and an
upper β-axial ligand. This upper ligand can vary with the most common being a cyano or
hydroxy group, whereas adenosyl or methyl group are the biological active forms in humans[7].
Within MTR the Cbl cofactor binds with the DMB group disengaged ('base-off') where in some
instances a conserved histidine residue can bind as the lower axial ligand (His-off or His-on).
During MTR's catalytic cycle, the bound Cbl undergoes different cobalt oxidation and
coordination   states,   such   as   alternating   between   the   oxidised   hexacoordinated
methylcob(III)alamin form and the reduced tetracoordinated cob(I)alamin form. Cob(I)alamin
is a strong nucleophile capable of extracting the methyl group from MeTHF, however it is also
highly susceptible to oxidative inactivation, resulting in a cob(II)alamin form that renders MTR
inactive. This necessitates the enzyme to enter a reactivation cycle where a reductive
methylation process regenerates the functional methyl(III)cobalamin form from the inactive
cob(II)alamin[8]. Previous structural studies using isolated domains of the bacterial orthologue,
MetH, have suggested that for MTR to function large conformational changes are required in
order for the Cob domain to interact with the Hcy and Fol domains within the rigid N-half during
catalysis but also with the Act domain in the C-half during reactivation[9-11].

Differentiating from its bacterial counterpart, human MTR has been shown to interact
with three eukaryote-only proteins, namely methylmalonic aciduria and homocystinuria type C
protein (MMACHC), methylmalonic aciduria and homocystinuria type D protein (MMADHC),
and methionine synthase reductase (MTRR)[12]. MMACHC and MMADHC are known Cbl
processing enzymes where MMACHC has Cbl reductase and dealkylase activities, to process
various Cbl forms by removing the upper axial ligand to eventually produce cob(II)alamin[13].
Recent studies have shown that MMADHC can form a complex with MMACHC[14,15], and can
load Cbl onto MTR through forming a rare cobalt-sulfur bond donating a cysteine residue as
one of the axial ligands[16,17]. MTRR is a multi-domain diflavin oxidoreductase that contains at
its N-terminus a flavin mononucleotide (FMN) binding domain, flexibly linked to a C-terminal
module that contains of a connecting (Conn) domain, a flavin adenine dinucleotide (FAD)
binding domain, and a nicotinamide adenine dinucleotide phosphate (NADPH) binding
domain[18]. MTRR plays a key role in reactivating MTR for catalysis, working in concert with the
SAM-bound Act domain of MTR. MTRR donates electrons from NADPH to reduce and
methylate   the   inactive   cob(II)alamin   bound   to   MTR,   to   produce   functional
methylcob(III)alamin[19,20]. This is a multistep sequential process where electrons from NADPH
are initially transferred to FAD and then FMN before ultimately being transferred to the bound

Cbl[7]. In bacteria this function is carried out by two separate proteins, flavodoxin and flavodoxin
reductase[21]. Significantly, the essential roles of MTR and its interaction partners in human
health are stressed by reported pathogenic mutations in their corresponding genes, that are
associated with a collection of rare inherited metabolic disorders[22]. Altered Cbl and folate
metabolism is also known to be linked to an increased risk of neural tube defects[23].
Little is known on how human MTR binds Cbl and if it shares the same conformational
flexibility as its bacterial orthologues. Even less is known on how human MTR interacts with
its eukaryote specific partner proteins that support its functionality. To address these
questions, we have used cryo-electron microscopy (cryo-EM) together with AlphaFold 3 (AF3)
structure predictions, to investigate how human MTR binds Cbl and how it interacts with its
functional protein partners MMACHC, MMADHC, and MTRR.
108
Results
Human MTR can be purified without bound cobalamin
We expressed and purified apo full-length human MTR (MTR^{FL}) with a C-terminal His$_6$
and FLAG tag from insect cells. Though milligrams of recombinant protein could be obtained,
size-exclusion gel filtration demonstrated that the majority of MTR^{FL} formed aggregates with
some monomeric protein (Supplementary Fig.1a). Negative stain electron microscopy (NS-
EM) confirmed the presence of homogenous and monodisperse particles from the monomeric
peak (Supplementary Fig. 1b). The molecular weight was determined to be 142.69 kDa by
native mass-spectrometry, agreeing with the calculated molecular weight of 143.85 kDa (with
tag) and confirming the purified monomeric protein was in the apo form without any detectable
bound Cbl ligands (Supplementary Fig.1c) as previously reported[24,25].
Using an absorbance-based activity assay[26] monomeric MTR^{FL} was determined to be
catalytically active (Supplementary Fig. 1d). To further assess functionality, we turned to
thermal shift experiments testing MTR^{FL} along with purified MTR^{N-Half} (Hcy and Fol domains)
and MTR^{C-Half} (Cap, Cob, and Act domains) constructs. Here we determined the melting
temperature (T$_m$) of these constructs as purified along with being in the presence of relevant
substrates and different cobalamins. In these experiments, MTR^{FL} was stabilised by MeTHF
with a 3.3 ± 0.3 °C melting temperature increase (ΔT$_m$) (Supplementary Fig. 1e, g). MeTHF
also induced a ΔT$_m$ increase of 4.8 ± 0.4 °C for MTR^{N-Half}, whereas no statistically significant
change in T$_m$ was observed for MTR^{C-Half} as expected (Supplementary Fig.1e, g). In contrast,
both Hcy and SAM showed no T$_m$ changes for any of the three constructs (Supplementary
Fig. 1g). We then tested hydroxycobalamin (HOCbl), cyanocobalamin (CNCbl), and
methylcobalamin (MeCbl) against the three constructs. As expected, no T$_m$ changes were
observed for MTR^{N-Half}, whereas all three Cbl ligands stabilised both MTR^{FL} and MTR^{C-Half} with
MeCbl eliciting the largest T$_m$ shifts (Supplementary Fig. 1f, g). Overall, these experiments
demonstrate that we can successfully purify functional apo full-length and constructs
corresponding to each half of human MTR.
136
Apo MTR is in a reactivation conformation for cobalamin loading
Having successfully purified apo MTR^{FL} and encouraged by our NS-EM results, we
attempted to determine a high-resolution structure of this as-purified, Cbl free state by cryo-
EM. Initial screening resulted in 2D classes representing only the C-half of MTR likely in the
reactivation conformation where the Act domain is engaged with the Cob domain
(Supplementary Fig. 2a). Unfortunately, after many attempts we could not produce conditions
that yielded reliable good 2D classes representing the entirety of full-length MTR, likely due to
instability and air-water denaturation. To overcome this, we screened and collected data in
the presence of MeTHF, reasoning that it would stabilise MTR^{FL} as found in our thermal shift
experiments (Supplementary Fig 1e, g). Initial screening resulted in separate 2D classes of
both halves of MTR (Supplementary Fig. 2b). Despite multiple processing strategies no
classes showed the concurrent presence of both halves of MTR. A larger dataset was then
collected which allowed us to determine a 3.1 Å resolution structure of the 70 kDa N-half

bound to MeTHF and a 2.8 Å resolution structure of the 68 kDa apo C-half (Fig. 1,
Supplementary Fig. 3, 9, 10).
The N-half structure of MTR[FL] confirmed that the Hcy and Fol domains are TIM barrel
folds with the short Linker 1 (Pro347 to Phe371) that tightly locks the domains side by side
(Fig. 1b). This arrangement implies that large conformational changes are required for the Cob
domain to interact with the N-half domains during catalysis as has been suggested for bacterial
MetH[7,9]. The Hcy domain showed weak density for a disulfide bond between Cys260 and
Cys323 with no other apparent density for substrate (Fig 1b, Supplementary Fig. 4a). On the
other hand, the Fol domain showed clear density for the MeTHF ligand (Fig. 1c) making
interactions with the conserved residues Asn470, Asp537, and Asn579 through its pterin ring
containing the methyl group. Additionally, the p-aminobenzoic acid (PABA) side chain of
MeTHF forms hydrogen bonds with the residues Arg384, Arg585, and Arg591 (Fig. 1c). This
MeTHF bound structure is highly similar to our previously deposited 2.7 Å crystal structure of
the MTR[N-Half] construct alone bound to THF (PDB 4CCZ), showing that the binding of MeTHF
is near identical to THF, with a RMSD of 0.68 Å (Supplementary Fig. 4a). Comparing our
human N-half cryo-EM structure to the reported N-half sections from *Thermus filiformis* (PDB
8G3H) and *Thermus thermophilus* HB8 (PDB 8SSC) MetH structures demonstrates how this
tight arrangement between the Hcy and THF domains is evolutionarily conserved
(Supplementary Fig. 4b)[24,27].
The Cbl free C-half structure of MTR[FL] was found to be in a conformation where the
Cap domain is displaced (Cap-off) from the Cob domain, allowing free access to the Cbl
binding pocket and engagement with the Act domain (Fig. 1d). No density was apparent for
the Cbl cofactor as expected (Fig. 1e). This domain arrangement is similar to that reported in
crystal structures of the isolated C-half from bacteria MetH bound to cob(II)alamin, which has
been suggested to be the reactivation conformation productive for reductive methylation[10,28,29]
(Supplementary Fig. 4c). Structural alignment against the *Escherichia coli* C-half structure
bound to cob(II)alamin (PDB 3BUL) and the apo C-half section from *T. thermophilus* (PDB
8SSC) shows only minor differences in the relative position of the Cap to the other domains
(Supplementary Fig. 4c). The expected lower axial ligand for the Cbl cofactor, His785, is in a
His-off conformation and the corresponding histidine sidechain adopts variable positions in
the bacteria MetH structures of the His-off state (Fig 1e, Supplementary Fig. 4c). Tyr1177,
which promotes the tetracoordinated cob(I)alamin form during MTR reactivation[29], is
positioned 3.7 Å closer to the putative position of the cobalt atom, when compared to the *apo*
MeTH structure from *T. thermophilus* (Supplementary Fig. 4c)[24].
184

Fig. 1 | Cryo-EM analysis of MTR[FL] bound to MeTHF. a, Cartoon schematic of the domain
organization of MTR. Regions resolved of MTR[FL] by cryo-EM are indicated. Representative
2D classes for each MTR map show the presence of just one half of the full-length enzyme.
b, The 3.1 Å resolution map and model of N-half bound to MeTHF. Domains are coloured as
in Fig. 1a and MeTHF is coloured magenta. **c,** Overlay of the map and model of the MeTHF
binding pocket in the N-half of MTR[FL], highlighting key residues and the ligand. **d,** The 2.8 Å
resolution map and model of the C-half with no bound Cbl cofactor. Domains are coloured as
in Fig. 1a. **e,** Overlay of the map and model of the cobalamin binding pocket in the C-half of
MTR[FL], highlighting key residues in the region and showing no detectable density
corresponding to Cbl.

Binding of HOCbl to MTR causes local structural changes
The conformation of the C-half of MTR[FL] suggests that the apo enzyme forms this
"reactivation-like" conformation before the binding of free Cbl from solution or Cbl loading
assisted by MMADHC. Therefore, to determine the conformational changes upon initial
binding of Cbl, we collected cryo-EM data of MTR[FL] in the presence of MeTHF and HOCbl.
SAM was also added in an attempt to determine what conformational changes could occur
before methylation of the bound Cbl. As with the MeTHF dataset, only separate 2D classes of
the N-half and C-half alone could be determined, suggesting their independence when MTR[FL]
is in the presence of these ligands. Processing of the N-half particles generated a 2.9 Å
resolution structure (Supplementary Fig. 5, 9, 10), identical to that with MeTHF alone (RMSD
0.356 Å) (Supplementary Fig. 6a).
Processing of the C-half particles revealed two states with an overall conformation as
the Cbl free form (Supplementary Fig. 5, 9, 10). Inspection of the Cob domain in both states
showed clear density for the bound cofactor, HOCbl (Fig. 2a, b, Supplementary Fig. 6b);
however, the density for SAM in the Act domain was discontinuous and did not allow for
modelling, suggesting flexibility or low occupancy of this ligand (Supplementary Fig. 6c). The
first state, determined to a resolution of 2.9 Å, is representative of the majority of the C-half
particles. In this state, density was apparent for HOCbl bound in a cob(II)alamin, base-off, His-
off state (Fig. 2a, Supplementary Fig. 6b). Density for the dimethylbenzimidazole (DMB) tail
bound in the pocket of the Cob domain was clear, interacting with conserved residues Ser830
and Gly860 (Supplementary Fig. 6b). The corrin ring showed less apparent density,
suggesting flexibility with Leu832 in the location of the lower axial ligand (Fig. 2c,
Supplementary Fig. 6b).
The second state was determined to a resolution of 3.2 Å (Fig. 2b, Supplementary Fig.
6b) and is representative of the minority of the particles. Similar to the His-off state described
above, density was apparent for HOCbl, however its corrin ring has shifted by 2.8 Å (Fig. 2c,
d) and is now in a His-on state, where His785 becomes the lower axial ligand (Fig. 2c).
Furthermore, conformational changes are apparent in the Cob domain when comparing the
His-off and His-on structures (Fig. 2d). These changes occur throughout the entire Cob
domain, involving a shift away from the Act domain during the transition from the His-off to the
His-on state. The density of the helical Linker 4 between the Cob and Act domains is also less
defined (Fig. 2b), suggesting a degree of flexibility in this state.
In the C-half maps of both states, the Cap domain showed less detail than the other
domains, suggesting heterogeneity possibly due to flexibility. Interested in analysing this
further, we turned to 3D variability analysis (3DVA)[30] where we compared the apo C-half map
against the Cbl bound C-half map with both His-off and His-on particles pooled together
(Supplementary Fig. 7, Supplementary Movies 1-4). For both the apo and Cbl bound states,
flexing of the Cap domain was evident where it moves towards the Act domain.
Simultaneously, the Cob domain appears to be moving away from the Act domain
(Supplementary Fig. 7a, b, Supplementary Movies 1, 3). Other movements were observed in
the Cbl bound state where the Cbl cofactor transitions from His-off to His-on correlating with
larger movements of the Cob domain away from the Act domain (Supplementary Fig. 7b,
Supplementary Movie 3) along with some rotation towards the Cap (Supplementary Fig. 7d,

Supplementary Movie 4). We believe that this displacement of the Cob domain is
representative of the conformational changes that occur when MeCbl is formed via reductive
methylation and the subsequent change from a five-coordinate His-off Cap-off state to a six-
coordinate His-on Cap-on state[31]. As no electron source was added to our sample, reductive
methylation cannot occur, thereby allowing us to trap a small percentage of the MTR protein
bound to HOCbl in this putative prelude state (His-on, Cap-off) before entering the catalytic
cycle, similar to what has been suggested for a crystal structure of the His-on HOCbl bound
state of the C-half from E. *coli* (PDB 3IV9)[29].
250

251
252
Fig. 2 | Cryo-EM analysis of the C-half of MTR[FL] bound to HOCbl. a, The 2.9 Å resolution
map and model of the C-half of MTR[FL] bound to HOCbl in a His-off state. Domains are
coloured as in Fig. 1a and HOCbl is coloured cyan. **b,** The 3.2 Å resolution map and model of
the C-half of MTR[FL] bound to HOCbl in a His-on state. Domains are coloured as in Fig. 1a and
HOCbl is coloured cyan. **c,** Positioning of HOCbl and His785 in the His-off state (left, beige)
and His-on state (right, cyan). **d,** Comparison of the movements of HOCbl and the Cob domain
between the His-off and His-on states. The HOCbl and Cob domain in the His-off state are
shown in beige, while HOCbl in the His-on state is shown in cyan, and the Cob domain in the
His-on state is shown in red.
262
Binding of MeCbl remodels the C-half of MTR[FL]
After Cbl loading and reductive methylation aided by MTRR, MTR is activated and
transforms into a six-coordinate His-on MeCbl-bound conformation, poised to begin the
catalytic cycle[32]. Recent studies have suggested that *T. filiformis* and *T. thermophilus* MetH
enters a so-called "resting" or "gating" state in the presence of Cbl and without substrates. In
this state the N-half (Hcy and Fol domains) stably interacts with the Cap and Cob domains of
the C-half in a Cap-on and a disengaged Act domain conformation[27,33]. Interested in capturing
a similar Cap-on state of human MTR without substrate, we prepared MTR[FL] in the presence
of MeCbl for cryo-EM analysis. MeCbl has been shown to readily bind to the Cob domain from
solution[17] and also stabilised the protein in our thermal shift experiments (Supplementary Fig.
1e, g).
Preliminary screening of MTR[FL] in the presence of MeCbl resulted in discernible 2D
classes of the 70 kDa N-half only, and unlike in our previous datasets no apparent 2D classes
could be determined corresponding to the C-half. However, as this sample of MTR[FL] lacked
any substrates, a larger dataset was then collected that resulted in a 2.7 Å resolution structure
of the apo N-half of MTR[FL] (Fig. 3b, Supplementary Fig. 8-10). Comparing this structure with
our MeTHF- and THF-bound structures shows little conformational changes in both Hcy and

Fol domains (Fig. 3c). No other domains were present in the map, and despite further
processing no other states containing the N-half could be found unlike what has been reported
for structures of thermophilic MTR homologues[27,33]. In these structures a loop located in the
Fol domain changes conformation due to the absence of folate, and was suggested to play a
role in altering the conformation of Linker 2 which connects the two halves of the enzyme,
resulting in the "resting/gating" state[27,33]. However, unlike thermophilic MetH, the folate
sensing loop in human MTR (residues 613-619) shows no structural changes due to the
absence of folate and is positioned in a similar manner as the ligand bound states (Fig. 3d).

The lack of a C-half structure from this dataset could have been due to a combination
of intrinsic flexibility and air-water denaturation. In theory the binding of MeCbl should induce
large conformational changes in the C-half, shifting MTR into a catalytical competent state and
displacing the Act domain from the Cob domain to allow interactions of the bound Cbl with the
N-half domains and Cap during catalysis. Further careful processing of our MeCbl-bound
dataset revealed classes equating to the Act domain of MTR^{FL} (≈ 38 kDa), showing its
disengagement from the Cob domain (Fig. 3e). Despite its small size, further processing
resulted in a 3.1 Å resolution map (Fig. 3e, Supplementary Fig. 8-10). Comparison of this Act
domain-alone structure with the Act domain in the context of our C-half structures (i.e. together
with Cob and Cap domains) shows little structural changes within the Act domain
(Supplementary Fig. 11a). Comparison to the crystal structure of an isolated Act domain
construct (PDB 2O2K)[34] showed that only a few residues in flexible regions displayed any
significant conformational differences, likely resulting from crystal packing (Supplementary
Fig. 11b). Therefore, we interpret that our MeCbl-bound MTR^{FL} dataset is likely in a His-on,
Cap-on state consisting of three flexible independent modules, namely the N-half (≈ 70 kDa),
the Cap and Cob domains (≈ 28 kDa), and the Act domain (≈ 38 kDa). Despite considerable
effort in data processing, we failed to find any 2D classes equating to the MeCbl-bound Cap-
Cob domains, likely due to its small size and its position between the N-half and Act domain.
Indeed, inspecting the 2D classes and low pass filtering our Act domain map show the
presence of blurry low-resolution density at the expected location of these two domains,
suggesting their presence (Supplementary Fig. 11c, d). Although the difference between our
MeCbl-bound structure and the reported structures of thermophilic MetH bound to inactive
Cbl[27,33] forms may be due to variations in sample preparation, it suggests that MTR can adopt
more conformations than has been previously suggested (Fig. 3f).

Fig. 3 | Cryo-EM analysis of MTR bound to MeCbl. a, Cartoon schematic of the domain
organization of MTR. Domains resolved by cryo-EM are indicated. **b,** The 2.6 Å resolution
map and model of the N-half of MTR[FL] in the apo state. Domains are coloured as in Fig. 3a.
c, Structural alignment of our apo (yellow) and folate bound (cyan) structures focused on the
folate binding site. Little conformational changes occur due to folate binding with only a slight
shift away from the active site for the small loop containing Ala381. **d,** Structural alignment of
our apo N-half structure (yellow) and thermophilic MetH structure (8G3H, light green) focused
on the folate binding site. Unlike the position of residues 605-611 in thermophilic MetH,
corresponding to a putative folate sensing loop, no structural changes occur in residues 613-
619 in human MTR due to the absence of folate. **e,** The 3.1 Å resolution map and model of
the MTR[FL] Act domain. The 2D classes shown are representative of the Act domain as
observed in the cryo-EM data analysis. In the model the missing linkers, Cap, and Cob
domains are also shown as a transparent model arranged as found in our C-half structures. **f,**
Proposed domain arrangements as cartoon representations of MeCbl-bound active human
MTR and thermophilic HOCbl-bound inactive MetH, without substrate. In human MTR MeCbl
binding results in a conformation with the N-half, Cap-Cob domains, and Act domain acting as
three independent modules. In contrast, HOCbl binding to thermophilic MetH results in a two-
module conformation with the N-half plus Cap-Cob domains, and the Act domain acting
independently.
333
MMADHC, not MMACHC, binds MTR directly to load cobalamin
The cobalamin chaperone enzymes MMACHC and MMADHC can form a heterodimer
in the presence of Cbl[14], and previous studies have implied a stable complex of MTR with
MMACHC and MMADHC[12,35]. Additionally, Cbl transfer from MMADHC to MTR has been
monitored through size-exclusion chromatography[17]. However, no structural insights into their
complex formation have been determined experimentally. Therefore, we turned to AlphaFold3
(AF3)[36] modelling to predict how both MMACHC and MMADHC could interact with MTR to
load Cbl.
AF3 predicted modelling of the MTR binary complexes was initially carried out
using the sequences of the full-length proteins. This was then followed by using only the
sequences of the N- and C-halves of MTR to confirm which domains were involved in protein-
protein interactions. AF3 models for the MTR[FL]-MMACHC[FL], MTR[N-Half]-MMACHC[FL], and
MTR[C-Half]-MMACHC[FL] complexes demonstrated random spatial positions of MMACHC on
MTR (Supplementary Fig 12, 14a, b). This combined with their low ipTM scores[36] (~ 0.2),
which is a confidence measure of the accuracy of predicted complexes, suggest that
MMACHC may not form a complex with MTR. In contrast the AF3 models for the MTR[FL]-
MMADHC[FL] complex (Fig. 4a, Supplementary Fig. 13) and the MTR[C-Half]-MMADHC[FL]
subcomplex (Fig. 4b, Supplementary Fig. 14d) resulted in highly similar and consistent
predicted models with correspondingly moderate to high ipTM scores[36] above 0.5. On the
other hand, the MTR[N-Half]-MMADHC[FL] subcomplex showed random spatial positions of each
protein with a corresponding low ipTM score suggesting that MMADHC does not interact with
either the Hcy or Fol domains of MTR (Supplementary Fig. 14c).
In both sets of predictions with either MTR[FL] or MTR[C-Half], MMADHC is straddled above
the Cob domain of MTR in the reactivation conformation, forming potential interactions with
all three domains of the C-half of MTR (Fig. 4a, b, Supplementary Fig. 13, 14a, b). This
interface results in the loop aa 255-266 from MMADHC to point towards the Cbl-binding pocket
in the Cob domain of MTR. This loop harbours the residues Cys261 and Cys262, which are
responsible for the binding of Cbl and its loading into MTR[17]. Superimposing the predicted
models with our HOCbl-bound C-half structure of MTR[FL] and the crystal structure of MMADHC
coordinating a thiolato-cobalamin ligand (PDB 6X8Z)[37] suggests that some conformational
changes are required for Cbl loading. The Cbl cofactor needs to move 4.3 Å towards the
binding pocket within the Cob domain of MTR when being loaded by MMADHC
(Supplementary Fig 15).
To verify MTR's predicted lack of interaction with MMACHC and its predicted positive
interaction with MMADHC we performed affinity-based pulldowns using MTR[FL] against a GST

fusion of MMADHC$^{\Delta N}$ (MMADHC lacking the first 123 residues that were predicted to be
disordered) and a GST fusion of full-length MMACHC (Fig. 4c, d, Supplementary Fig. 16a).
These pulldowns show that MMACHC weakly interacts with MTR^{FL} (lane 7) whereas
MMADHC appears to strongly interact (lane 8). To further confirm which halves of MTR are
involved we carried out similar affinity-based pulldowns using MTR^{N-Half} and MTR^{C-Half}
constructs (Fig. 4c, e). These pulldowns confirmed the weak, to potentially absent, interaction
of MMACHC against the MTR^{N-Half} and MTR^{C-Half} constructs (Fig. 4e, Supplementary Fig. 16a),
in which no positive pulldowns were observed (lanes 9, 11). Similar to previous published
analyses on a C-terminal construct of MTR[17], we observed that MMADHC$^{\Delta N}$ interacts with
MTR^{C-Half} (lane 12) but not with MTR^{N-Half} (lane 10). Overall, these collection of pulldowns are
consistent with their corresponding AF3 predictions.
We next used size-exclusion gel filtration to determine the loading of HOCbl onto
MTR^{FL} by MMADHC$^{\Delta N}$ (without a GST tag), through monitoring absorbance at 350 nm,
indicative of the presence of Cbl. MTR^{FL} displays no absorbance at 350 nm, indicating that it
is not bound to cobalamin (Fig. 4f upper left panel) agreeing with our native mass spectrometry
measurements (Supplementary Fig. 1c). In the absence of MMADHC, low amounts of HOCbl
binds MTR (Fig. 4f upper right panel) whereas MeCbl binds more readily (Fig. 4f lower left
panel), agreeing with a previous analysis on a C-terminal construct of MTR[17]. When
preincubated with MMADHC$^{\Delta N}$, HOCbl is loaded into MTR^{FL} through MMADHC$^{\Delta N}$, suggesting
a direct interaction (Fig 4f lower right panel). Together our pulldown and Cbl loading
experiments suggest that MTR not only interacts with MMADHC for Cbl loading but also
appears to interact with MMADHC when Cbl is not present.
391

Fig. 4 | MMADHC forms a complex with MTR to load cobalamin. a, Representative predicted AF3 model of the MTR-MMADHC complex. The arrangement of the full-length complex is shown as a cartoon depicted with the domain colours as shown in Fig. 4a. The zoom-in cartoon shows only MTR^{C-Half} with MMADHC; MMADHC is a ribbon model whereas MTR is shown as a surface model. **b,** Structural overlay of the predicted MTR^{C-Half}-MMADHC AF structure against our cryo-EM derived model of His-off MTR^{C-Half}. The zoom-in panel focuses on the MTR-MMADHC interface showing the relative location of the functionally important MMADHC residues C261 and C262. **c**, Constructs and domain arrangements of MTR, MMACHC, and MMADHC used in cobalamin transfer and pulldown assays. Affinity tagged constructs are indicated. **d,** GS resin affinity pulldown of MTR^{FL} against GST-MMACHC and GST-MMADHC$^{\Delta N}$. Lanes 1–3 represent the protein inputs used for the assay. Lanes 4–6 represent the elution fractions of each individual protein incubated with the resin to verify their ability to bind the resin. Lanes 7 and 8 represent the pulldown of MTR^{FL} with GST-MMACHC and the pulldown of MTR^{FL} with GST-MMADHC, respectively. **e,** GS resin affinity pulldown of MTR^{N-Half} and MTR^{C-Half} against the GST-MMACHC and GST-MMADHC$^{\Delta N}$. Lanes 1–4 represent the protein inputs used for the assay. Lanes 5–8 represent the elution fractions of each individual protein incubated with the resin to verify their ability to bind the resin. Lanes 9 and 10 represent the pulldown of MTR^{N-Half} with GST-MMACHC and GST-MMADHC, respectively. Lanes 11 and 12 represent the pulldown of MTR^{C-Half} with GST-

414   MMACHC and GST-MMADHC, respectively. **f,** Size-exclusion chromatography of cobalamin
415   loading assay showing MTR^{FL}, MTR^{FL} plus HOCbl, MTR^{FL} plus MeCbl, and MTR^{FL} plus
416   HOCbl-loaded MMADHC$^{\Delta N}$ monitored at 280 nm (blue) and 350 nm (black for HOCbl or red
417   for MeCbl).
418

MTRR interacts with MTR through multiple domains

For the reactivation cycle, MTR requires the diflavin oxidoreductase MTRR to reduce
the bound cob(II)alamin for methylation by SAM[19]. Beyond reports that the FMN domain of
MTRR can interact with the Act domain of MTR[38], little is known on how MTRR interacts with
MTR to transfer electrons from NADPH. To gain structural insights into their interactions we
turned to AF3 to predict possible models of the MTR-MTRR complex.
AF3 modelling of the complex between MTR^{FL} and MTRR^{FL} produced predicted
models with moderate ipTM scores[36], in which the MTR N-half location varies between the
individual predictions and the MTR C-half adopts a reactivation conformation (Fig. 5a,
Supplementary Fig 17). Consistent across the five predicted structures is the relative location
of MTRR where the MTRR FMN domain is modelled to sit atop the MTR Cob domain and
appears to interact with the Act domain (Fig. 5a). This agrees with previous biophysical studies
demonstrating a physical interaction between the MTRR FMN domain and MTR Act domain[38].
To our surprise, the models reveal that in addition to the FMN domain, the other regions of
MTRR (i.e. Conn domain, FAD domain, and NADPH domain; referred thereafter as MTRR
ΔFMN) can also interact with MTR which was consistent across each prediction. This
interaction occurs through two interfaces of MTRR ΔFMN, one involving the Conn domain that
interacts with Linker 4 of MTR, and the other involving the NADPH domain that interacts with
the MTR Cob domain. (Fig. 5a, 6a, Supplementary Fig. 17). This previously unidentified
interaction was also consistently replicated in the AF3 modelling of complexes between the
MTR C-half and the MTRR ΔFMN (Supplementary Fig. 18a). On the contrary, no consistent
interactions with MTRR were predicted when the human MTR C-half sequence is replaced
with bacterial MetH C-half sequences, suggesting that the interaction between the MTR C-
half and MTRR ΔFMN is unique to eukaryotes (Supplementary Fig. 18b-d). Together, these
predictions suggest that human MTRR interacts with human MTR, involving not only the FMN
domain of MTRR, but also its C-terminal Conn, FAD, and NADPH domains.
To confirm these interactions *in vitro*, we used a combination of recombinant co-
expression and reconstitution pulldown experiments. Firstly, co-expressing MTR^{FL} with
MTRR^{FL} in insect cells and purifying using the FLAG tag on MTR confirmed the presence of a
complex without Cbl (Supplementary Fig. 19a). Additionally, in blue native PAGE experiments
MTR^{FL} co-expressed with MTRR^{FL} ran at a larger molecular weight than purified MTR^{FL} alone
showing complex formation (Supplementary Fig. 19b). We then employed talon resin-based
affinity pulldowns using purified MTRR^{FL} as bait against the purified MTR^{FL}, MTR^{N-Half}, and
MTR^{C-Half} (Fig. 5b, Supplementary Fig. 19c), where MTRR^{FL} showed an interaction with MTR^{FL}
(Fig. 5c lane 9) and MTR^{C-Half} (lane 11), but not MTR^{N-Half} (lane 10). Importantly, these pulldown
experiments confirm that the interaction between MTR and MTRR occurs through MTR^{C-Half},
even in the absence of Cbl or other ligands. Analytical size exclusion assays further confirmed
that both full-length MTR and MTRR can form a complex (Fig. 5d left) specifically mediated
by the C-half of MTR (right), but not its N-half (middle).
458

460
461
Fig. 5 | MTRR forms a stable complex with MTR. a, Constructs and domain arrangements
of MTR and MTRR used in pull-down and gel filtration assays. Affinity tagged constructs are
indicated. **b**, Representative predicted AF3 structure of the MTR-MTRR complex. The
arrangement of the complex is shown as a cartoon depicted with the domain colours as shown
in Fig. 5a. **c,** Talon resin affinity pulldown of the constructs shown in Fig. 5b. Lanes 1–4
represent the protein inputs used for the assay. Lanes 5–8 represent the elution fractions of
each individual protein incubated with the resin to verify their ability to bind the resin. Lanes
9–11 represent the pulldown of the different protein combinations (His-MTRR^FL with MTR^FL,
MTR^{N-Half}, or MTR^{C-Half}, respectively). **d,** Analytical size-exclusion chromatography of MTRR^FL
in complex with MTR^FL (left), MTR^{N-half} (middle), or MTR^{C-half} (right). Top panels show the
normalised spectrum at 280 nm and bottom panels show the corresponding SDS-PAGE of
the eluted fractions.
474
To confirm the putative interaction between MTRR ΔFMN and MTR, we expressed and
purified a GST tagged construct of MTRR^{ΔFMN} (Fig. 6b, Supplementary Fig. 19d) for interaction
studies. In GST pulldown experiments this construct showed an interaction with both MTR^FL
(Fig. 6c lane 9) and MTR^{C-Half} (lane 11), but not MTR^{N-Half} (lane 10). Furthermore, analytical
size exclusion using MTRR^{ΔFMN} without the GST tag confirmed the interactions of this region
of MTRR with MTR^FL (Fig. 6d left) and MTR^{C-Half} (right), but not with MTR^{N-Half} (middle). Overall,
our co-expression, pulldown, and gel filtration experiments are consistent with the AF3
predictions of the complex between MTR and MTRR.
483

485
486
Fig. 6 | MTRR without the FMN domain can interact with MTR. a, Constructs and domain
arrangements of MTR and MTRR used in pull-down and gel filtration assays. Affinity tagged
constructs are indicated. **b**, Representative predicted AF3 structure of the MTR^C-half-MTRR^ΔFMN
complex. The arrangement of the complex is shown as a cartoon depicted with the domain
colours as shown in Fig. 6a. **c,** GS resin affinity pulldown of the constructs shown in Fig. 6b.
Lanes 1–4 represent the protein inputs used for the assay. Lanes 5–8 represent the elution
fractions of each individual protein incubated with the resin to verify their ability to bind the
resin. Lanes 9–11 represent the pulldown of the different protein combinations (GST-
MTRR^ΔFMN with MTR^FL, MTR^N-Half, or MTR^C-Half, respectively). **d,** Analytical size-exclusion
chromatography of MTRR^ΔFMN in complex with MTR^FL (left), MTR^N-half (middle), or MTR^C-half.
Top panels show the normalised spectrum at 280 nm and bottom panels the corresponding
SDS-PAGE analyses of the fractions indicated.

499
Discussion
Utilising vitamin B12 as a cofactor, MTR coordinates folate and methionine metabolism
by performing three methylation reactions through its catalytic and reactivation cycles. Early
x-ray crystallography and biochemical studies[9-11,39,40] on the bacterial homologue MetH
demonstrated that the enzyme has structural modularity in order to carry out these reactions.
Similarly, our cryo-EM data reveals human MTR to be modular comprising either two or three
independent modules linked by flexible linkers, representing inactive and active states
respectively. Studies on MetH have suggested that when the enzyme is active, large
conformational changes occur as the Cob domain shuttles between the Fol and Hcy domains
to transfer a methyl group from MeTHF to Hcy using MeCbl during catalysis[9]. Every ~2000

reactions the bound cob(I)alamin intermediate can become oxidised to inactive
cob(II)alamin[32]. Reactivation requires another large conformational change that repositions
the Act domain to interact with the Cob domain to allow methylation of the inactive Cbl by
SAM[7]. These conformational changes are highly likely to be regulated and to be conserved in
human MTR. Although Cbl oxidation state and substrates have been suggested to be
important contributors to conformational changes[40], little is known on how and when they are
triggered.
Recent MetH structures together with our findings on human MTR are beginning to
suggest a model for how these domain movements are regulated (Figure 7). Though revealing
different conformational states, as will be discussed below, our structures are consistent with
recent reports of a tetra-domain structure of *T. filiformis* MetH[27], along with a full-length
structure[24] and multiple crystal structures of different constructs of *T. thermophilus* MetH
bound to various ligands[33] (Supplementary Fig. 20). The collection of thermophilic MetH
structures has revealed long sought snapshots of the full-length enzyme, and of the Cob
domain engaged with either the Hcy or Fol domains, giving insights into catalysis[33].
Surprisingly a new conformation in MetH was initially reported by cryo-EM termed the
"resting state" - a hydroxycob(II)alamin bound, substrate free conformation where the Act
domain is disengaged, the Cap-on Cob module intimately interacts with the N-half, and a folate
sensing loop is positioned into the folate binding site (PDB 8G3H)[27]. This resting state was
replicated by crystallisation with a MetH construct without the Act domain and bound to
propylcob(III)alamin (PDB 9CBP). Two additional crystal structures of a MetH Fol-Cap-Cob
construct bound to MeCbl in a Cap-on conformation were also determined (PDB 9CBO)[33].
These three states were subsequently termed "gating states", as they are interpreted to be
the "Hcy-gate" (PDB 9CBP), "Fol-gate" (PDB 9CBO), and "Act-gate" (PDB 9CBO) before the
requisite conformational change that leads to the Hcy-on, Fol-on, and Act-on states
respectively[33]. However, both "resting state" and "Hcy-gate" structures are of inactive Cbl
forms without any bound substrates, suggesting that they are more likely representative of the
enzyme inactivated by oxidation and of the cob(I)alamin state before the binding of MeTHF.
Therefore the resting state is more likely the shared conformation between the catalytic and
reactivation cycles[27] for both the "Act-gate" (i.e. before reactivation) and "Fol-gate" (i.e. before
catalysis), depending on the oxidation state of Cbl.
Favouring this alternative interpretation is that *E. coli* MetH shows ordered ping-pong
kinetics where MeTHF is the first substrate to bind and THF is the last product to leave[41],
meaning that the resting state/Hcy-gate structures could not be substrate bound intermediates
during catalysis. Additionally, it is likely that the probability of oxidation of the bound
cob(I)alamin is increased when MeTHF is limited, and a recent study showed that the rate of
reactivation for human MTR is increased in the presence of both the substrate MeTHF and
the product THF[42]. This increased reactivation likely occurs due to substrate MeTHF binding
to the Fol domain and displacing the folate sensing loop[27] allowing the formation of a Cap-off
Act-on state productive for reactivation. A similar conformational change due to MeTHF
binding to the cob(I)alamin state would occur resulting in a Fol-on conformation. In the MeCbl-
bound "Fol-gate" and "Act-gate" crystal structures of MetH, the folate sensing loop is not
displaced into the folate binding site suggesting that folate binding has little to no influence in
any structural transitions in these states[33]. These two structures also have different relative
orientations of the Fol domain to the Cap-on Cob domain module that is suggestive of flexibility
(Supplementary Fig. 20). These structures also agree with our MeCbl-bound cryo-EM data of
human MTR, where the folate sensing loop has not moved into the folate binding site, and the
Act domain is disengaged from the highly flexible Cap-on Cob domain module. This
conformation seen in our MTR structure, along with the MetH crystal structures, should be
more aptly termed the "Hcy-gate" as the enzyme is bound to MeCbl, a Cbl form that is
productive for methyl transfer to the substrate Hcy. The MeCbl-bound enzyme without bound
folate adopts this more flexible state, instead of the more compact "resting state", likely due to
the distance that the Cob domain has to travel between the Fol and Hcy domains (~50 Å)
during the catalytic cycle.

564 Despite sharing a catalytic mechanism with MetH that exploits conformational
565 dynamics between domains, human MTR has built in distinctive features, not present in MetH,
566 to accommodate human Cbl metabolism. For example, while both MetH and MTR share
567 similar reactivation conformations prior to receiving Cbl, MetH does not require a highly
568 specialised chaperone to load the cofactor[43]. On the contrary, MMADHC has a central role in
569 Cbl trafficking in eukaryotes and is known to direct the co-factor to either methylmalonyl-CoA
570 mutase (MMUT) in the mitochondria or MTR in the cytosol[14]. AlphaFold predictions suggest
571 that MMADHC engages the exposed Cbl-binding pocket via all three domains in the C-half of
572 MTR when in a reactivation conformation. This interaction model aligns with experimental data
573 on Cbl loading, suggesting a structural rationale for how MMADHC and MTR form a complex.
574 Cbl is loaded as cob(I)alamin onto MTR, but most of it is readily oxidised into
575 cob(II)alamin[17] which cannot support catalysis. The reductive methylation of cob(II)alamin,
576 required for the reactivation, is assisted by electron transfer from the essential flavoprotein
577 MTRR[20]. MTRR is an unusual member of the diflavin reductase family where it has a much
578 longer hinge region linking the N-terminal FMN domain with the rest of the enzyme, than other
579 family members[44]. Previous studies revealed that the FMN domain of MTRR engages with
580 MTR to reduce the cobalamin, which is subsequently methylated to MeCbl[19]. Our AF3
581 modelling and biophysical analyses of the MTR-MTRR complex revealed previous unknown
582 interacting interfaces between these proteins and explains its long hinge region. AF3 predicted
583 models suggested that MTRR can also interact with MTR through its NADPH domain and
584 Conn domain when the FMN domain is engaged. This unexpected additional interaction was
585 confirmed by our biophysical assays using truncated constructs. The additional interfaces
586 could indicate a mechanism of MTR to optimize the efficiency of the reductive methylation by
587 forming a stable complex with MTRR even prior to entering the reactivation cycle. Expectedly,
588 this interaction is not predicted between MTRR and MetH, the latter relying on a different
589 reactivation system supported by flavodoxin[21]. The fact that MTR cannot be reactivated by
590 flavodoxin, and conversely MetH cannot be reactivated by MTRR, suggests that this newly
591 identified interaction may be essential for Cbl reduction[25]. This lack of cross-reactivity in
592 reactivation could be due to the lower sequence conservation in the region of MTR that is
593 responsible for forming this extended interaction with MTRR (Supplementary Fig. 21 α-helix
594 35).
595 Figure 7 integrates our cryo-EM snapshots, AF3 predictions, biophysical data, and
596 interpretation of previous studies along with other structures, into a proposed composite
597 mechanism for Cbl loading, reactivation, and catalysis. During cobalamin loading (i), apo MTR
598 is poised in the Cap-off reactivation conformation, ready to interact with MMADHC for
599 receiving Cbl. MMADHC loads cob(I)alamin which without SAM can be readily oxidised to
600 cob(II)alamin. Therefore, during the reactivation cycle (ii) MTR still in the reactivation
601 conformation is ready to interact with MTRR, for the reductive methylation of the bound Cbl
602 using SAM. This interaction is secured by two interfaces and allows an efficient electron
603 transfer from the FMN domain of MTRR to the Cbl. Following Cbl methylation, MTR undergoes
604 a large conformational change as the bound Cbl goes from His-off to His-on, triggering MTRR
605 unbinding. This results in the flexible MeCbl-bound state (Hcy-gate), where the Act and Cob
606 domains disengage, and the Cap domain takes its position atop the Cbl to protect it from
607 oxidation before beginning catalysis. This primes MTR within the catalytic cycle (iii) ready to
608 engage with the Hcy domain to carry out a methyl transfer reaction to form Met. After catalysis,
609 the products THF and Hcy leave MTR and result in a "Fol-gate" Cap-on conformation to protect
610 the cob(I)alamin intermediate. If oxidation has occurred that results in cob(II)alamin, MTR
611 would also form the same Cap-on conformation. In these two scenarios, binding of the
612 substrate MeTHF would elicit different responses. When MTR is bound with cob(I)alamin,
613 MeTHF binding to the Fol domain disrupts its interaction with the Cap and Cob domains to
614 promote a "Fol-on" conformation to produce MeCbl and continue another catalytic cycle (iii).
615 When MTR is bound with cob(II)alamin, MeTHF binding to the Fol domain also disrupts its
616 interaction with the Cap and Cob domains but promotes the Act-on reactivation conformation
617 which allows MTR to proceed with the reactivation cycle (ii).
618

620
621
Fig. 7 | Proposed mechanism of cobalamin loading, reactivation, and catalysis of MTR.
Model showing various cartoons of proposed conformations and complexes for human MTR
with MMADHC and MTRR to illustrate three major aspects of its function: (i) cobalamin
loading, (ii) reactivation cycle, and (iii) catalytic cycle. Please refer to the main text for more
details for each aspect. Different Cbl oxidation states are stated and represented as light
orange for Cbl(II) (cob(II)alamin), red for Cbl(III) (cob(III)alamin), and dark cyan for Cbl(I)
(cob(I)alamin). Domains are coloured as previously depicted in previous figures. This
proposed mechanism is not comprehensive. Therefore, other conformations and complexes
may also occur.

631
While our proposed mechanism attempts to show how MTR traverses from a Cbl-free
apo state to a MeCbl-bound active state, there are a number of caveats and unanswered
questions. While our cryo-EM structures show for the first time Cbl binding of human MTR and
its intrinsic flexibility, these structures are determined from samples treated with free Cbl in
solution and thus may not be fully representative of when Cbl is loaded by MMADHC and
reactivated by MTRR. Additionally, we have carried out two-way interaction studies between
MTR and its proposed interacting partners without presence of any ligands or additional
proteins, which may further modulate how these proteins interact. For example, MMACHC
may stably interact with MTR when MMADHC is present as a ternary complex[14,15] with
possible consequences for cofactor loading. The findings that both MMADHC and MTRR can
bind to MTR without any bound Cbl also suggests that these proteins may have further roles
beyond Cbl loading and reactivation. Supporting this hypothesis is a previous report which
suggested that MTRR could act as a chaperone of apo MTR as it increases MTR protein
stability[25]. A recent study also proposed a chaperone role of MMADHC for MTR[42]. These
studies point to the possibility of ternary complexes where a Cbl-free complex of MTR and
MTRR may interact with Cbl-free or Cbl bound MMADHC. This is also underlined by reports

of a quaternary complex of all four proteins namely MTR, MTRR, MMACHC and MMADHC[12].
Another report suggested a direct interaction of MMACHC with MTR[35]. However, this does not
appear to be supported by our AF3 predictions and pulldown assays where MMACHC shows
little to no interaction. Clearly, more studies delving into three/four-way complexes are
warranted in future.
Overall, our findings expand current knowledge on human MTR and contextualize
previous reported findings on both human and bacterial orthologues. Importantly we reveal
insights into eukaryote-unique aspects in terms of the complexes that MTR forms to support
its function. Given that pathogenic mutations in the MTR, MTRR, MMACHC, and MMADHC
genes are linked to Cbl deficiency, it is likely that some variants effect residues at protein-
protein interfaces of these MTR complexes. Indeed, disease-causing variants of MMADHC
have been suggested to weaken its interaction with MTR hindering Cbl loading[17].
Consequently, future experimental structure determination of the MTR complexes with
MMADHC and MTRR will be essential in understanding how pathogenic mutants alter their
complex formation. Such molecular insights could inform the development of targeted
therapies towards the treatment of these rare metabolic disorders.
664

Methods
Cloning, expression, and purification of MTR constructs
Full-length MTR (MTR^{FL}) for cryo-EM, activity, and thermal stability assays was cloned into
pFB-CT10HF-LIC encoding for a C-terminal TEV cleavable His-tag and FLAG-tag for insect
cell expression. MTR^{FL} was expressed in Sf9 cells infected with P3 baculovirus and grown in
Sf-900 III SFM (Life Technologies) for 72 hrs. Cell pellets were harvested and homogenized
in lysis buffer (50 mM HEPES pH 7.5, 500 mM NaCl, 5% glycerol, 0.5 mM Tris(2-
carboxyethyl)phosphine hydrochloride (TCEP) and 20 mM imidazole) using sonication, and
insoluble material was removed by centrifugation. MTR was then purified by affinity (Ni-
sepharose; Cytiva) and size-exclusion (Superdex 200; Cytiva) chromatography to separate
monomeric MTR from aggregates. Purified MTR^{FL} was then concentrated to 3–10 mg/ml,
aliquoted, and stored in storage buffer (50 mM HEPES pH 7.5, 500 mM NaCl, 5% glycerol,
0.5 mM TCEP) at −80 °C.
Full-length MTR (MTR^{FL}) for pull-down and gel filtration assays, was cloned into pNIC-
CT10HSII encoding for a C-terminal TEV cleavable His-tag and tandem StrepII-tag for
bacterial expression. MTR^{FL} was expressed in ArcticExpress DE3 cells. Cells were cultured in
Terrific Broth (Formedium) at 30 °C in Terrific Broth and expression was carried out for 60 hrs
at 10 °C induced with 0.1 mM isopropyl-β-D-thiogalactopyranoside (IPTG). Cell pellets were
harvested and homogenized in lysis buffer (50 mM sodium phosphate pH 7.5, 500 mM NaCl,
5% glycerol, 0.5 mM TCEP, 1.0 % triton X-100, and 20 mM imidazole) using sonication, and
insoluble material was removed by centrifugation. MTR^{FL} was then purified from the
supernatant by tandem affinity using Ni-sepharose followed by Streptactin XT resin (Cytiva).
MTR^{FL} was then treated with His-tagged TEV protease overnight at 4 °C and passed over Ni-
sepharose resin to remove the TEV protease and any uncleaved protein. A final size-exclusion
(Superdex 200; Cytiva) chromatography step was used to separate monomeric MTR^{FL} from
degradation products. Purified MTR^{FL} was then concentrated to 3–10 mg/ml, aliquoted, and
stored in storage buffer (50 mM HEPES pH 7.5, 500 mM NaCl, 5% glycerol, 0.5 mM TCEP) at
−80 °C.
The N-terminal construct of MTR, residues 16-657 (MTR^{N-Half}), was cloned into pNIC-
Bsa4 encoding for a N-terminal TEV cleavable His-tag. The C-terminal construct of MTR,
residues 661-1265, (MTR^{C-Half}) was cloned and into pNIC-CH as a TEV cleavable His-GST N-
terminally tagged for bacterial expression (Twist Bioscience). A stop codon was added at the
end of the MTR^{C-Half} coding sequence before the vector encoded His-tag. Both constructs were
expressed in Rosetta DE3 cells. Cells were cultured in Terrific Broth (Formedium) at 37 °C
and expression was induced overnight at 18 °C with 0.2 mM IPTG. Pellets were harvested,
homogenized in lysis buffer (50 mM HEPES pH 7.5, 500 mM NaCl, 5% glycerol, 0.5 mM TCEP
and 20 mM imidazole) by sonication, and insoluble material was removed by centrifugation.
The supernatant was subjected to affinity (Ni-sepharose; Cytiva) and the eluted protein

applied to size-exclusion (Superdex 200; Cytiva) chromatography. Each construct was then
treated with His-tagged TEV protease overnight at 4 °C and passed over Ni-sepharose resin
to remove the TEV protease and uncleaved protein. In some incidences, an additional size-
exclusion step was carried out to remove residual TEV protease. Purified MTR^{N-Half} and MTR^{C-Half}
proteins were concentrated to 10–20 mg/ml and stored in storage buffer (50 mM HEPES
pH 7.5, 500 mM NaCl, 5% glycerol, 0.5 mM TCEP) at −80 °C.

Cloning, expression, and purification of MTRR constructs

Full-length codon optimised MTRR (MTRR^{FL}) was cloned into pFastBac (Genscript) encoding
for a TEV cleavable N-terminal His-tag and was expressed in Sf9 cells grown in Sf-900 III
SFM (Life Technologies) for 72 hrs. Cell pellets were harvested and resuspended in lysis
buffer (50 mM sodium phosphate pH 7.5, 500 mM NaCl, 5% glycerol, 0.5 mM TCEP, 1.0%
triton x-100, 100 µM flavin mononucleotide). The resuspended pellet was incubated at room
temperature for 30 minutes with spinning, followed by sonication, and then subsequent
centrifugation to remove insoluble material. MTRR^{FL} was then purified from the soluble fraction
by affinity (Ni-sepharose; Cytiva). Purified MTRR^{FL} was then diluted in low salt buffer (50 mM
HEPES pH 7.5, 50 mM NaCl, 5% glycerol, 0.5 mM TCEP) and applied to a HiTrap Q column
(Cytiva). A gradient from 50 mM to 1M NaCl was used to elute MTRR^{FL}. Appropriate fractions
were pooled, concentrated, and buffer exchanged into storage buffer (50 mM HEPES pH 7.5,
500 mM NaCl, 5% glycerol, 0.5 mM TCEP.)

The MTRR construct without the FMN domain, residues 217-698, (MTRR$^{\Delta FMN}$) was
cloned from a codon optimised sequence (Twist Bioscience) into pGTVL2 to produce a TEV
cleavable His-GST N-terminally tagged protein for bacterial expression. MTRR$^{\Delta FMN}$ was
expressed in Rosetta DE3 cells. Cells were cultured in Terrific Broth (Formedium) at 37 °C
and expression was induced overnight at 18 °C with 0.2 mM IPTG. Pellets were harvested,
homogenized in lysis buffer (50 mM sodium phosphate pH 7.5, 500 mM NaCl, 5% glycerol,
0.5 mM TCEP, 1.0% triton X-100, and 20 mM imidazole) by sonication, and insoluble material
was removed by centrifugation. To produce tagged protein for pull-down assays the
supernatant was subjected to affinity (Ni-sepharose; Cytiva) and the eluted protein was further
purified by anion exchange using a HiTrap Q column (Cytiva). Clean fractions containing the
GST tagged construct were pooled and further treated to size-exclusion (Superdex 200;
Cytiva) chromatography. To produce cleaved MTRR$^{\Delta FMN}$ protein for analytical gel filtration
assays the supernatant was subjected to affinity (Ni-sepharose; Cytiva), and the eluted protein
incubated with His-tagged TEV protease overnight at 4 °C. The cleaved protein was then
subjected to size-exclusion (Superdex 200; Cytiva) chromatography. Peak fractions of cleaved
MTRR$^{\Delta FMN}$ were pooled and passed over Ni-sepharose resin to remove any remaining TEV
protease. Both the tagged and cleaved MTRR$^{\Delta FMN}$ proteins were concentrated to 5-10 mg/ml
and stored in storage buffer (50 mM HEPES pH 7.5, 500 mM NaCl, 5% glycerol, 0.5 mM
TCEP) at −80 °C.

Co-expression, and purification of the MTR-MTRR complex

Using the biGBac cloning system the genes for full-length MTR and full-length MTRR were
cloned into pBIG1a. The MTR^{FL} gene encoded for a C-terminal TEV cleavable His and FLAG
whereas the MTRRFL gene encoded a N-terminal His-tag. The MTR^{FL}-MTRR^{FL} complex was
expressed in Sf9 cells grown in Sf-900 III SFM (Life Technologies) for 72 hrs. Cell pellets were
harvested and homogenized in lysis buffer (50 mM HEPES pH 7.5, 500 mM NaCl, 5% glycerol,
0.5 mM TCEP) using sonication, and insoluble material was removed by centrifugation. The
complex was then purified by affinity (Ni-sepharose; GE Healthcare) and size-exclusion
(Superdex 200; GE Healthcare) chromatography. Fractions corresponding to the complex
were pooled, concentrated to 5–10 mg/ml and stored in storage buffer (50 mM HEPES pH 7.5,
500 mM NaCl, 5% glycerol, 0.5 mM TCEP) at −80 °C.

Cloning, expression, and purification of MMACHC and MMADHC$^{\Delta N}$

The cloning, expression, and purification of MMACHC and MMADHC$^{\Delta N}$ (residues 124-296)
has been described in detail elsewhere[13,14]. In brief His-GST tagged MMACHC and

MMADHC$^{\Delta N}$ were expressed in Rosetta DE3 cells. Cells were cultured in Terrific Broth
(Formedium) at 37 °C and expression was induced overnight at 18 °C with 0.1 mM IPTG.
Pellets were harvested, homogenized in lysis buffer (50 mM HEPES pH 7.5, 500 mM NaCl,
5% glycerol, 0.5 mM TCEP and 20 mM imidazole) by sonication, and insoluble material was
removed by centrifugation. The supernatant was subjected affinity (Ni-sepharose; Cytiva) and
the eluted protein applied to size-exclusion (Superdex 200; Cytiva) chromatography. To
produce cleaved MMADHC$^{\Delta N}$ the His-GST tagged protein after size exclusion was treated
with His-tagged TEV protease overnight at 4 °C and passed over Ni-sepharose resin to
remove the TEV protease and uncleaved protein. Purified tagged MMACHC, tagged
MMADHC$^{\Delta N}$, and cleaved MMADHC$^{\Delta N}$ proteins were concentrated to 10–20 mg/ml and stored
in storage buffer (50 mM HEPES pH 7.5, 500 mM NaCl, 5% glycerol, 0.5 mM TCEP) at −80 °C.

769
Native mass spectrometry
By dialysis, purified MTR^{FL} was exchanged from into 50 mM ammonium acetate, pH 6.5. The
sample was loaded into Quantitative Time-of-Flight (Q-TOF) 6530 mass spectrometer
attached to a standard electrospray ionisation source (Agilent Technologies), with 77 a
constant flow rate of 6 µl min^{-1}, using a 1.0-ml gas-tight positive displacement syringe
(Hamilton) and PEEK capillary tubing with an inner diameter of 0.005 inches.

776
Enzyme activity assay
MTR activity assay was adapted from the previously described cuvette-based nonradioactive
assay[26]. The current assay was miniaturised for a U-bottom 96-well plate. The following
reagents were added to each well in the mentioned order: 24.4 µL of 100 mM phosphate
buffered saline (PBS) pH 7.5, 2 µL of 500 mM dithiothreitol (DTT), 2 µL of 3.8 mM S-
adenosylmethionine (SAM), 2 µL of 100 mM L-homocysteine (Hcy), 5 µL of 4 µM MTRF^{L}, and
4 µL of 500 µM HOCbl. The plate was then incubated for 5 minutes at 37 °C, allowing DTT
and HOCbl to create a semi-anaerobic environment by reducing O_2 into H_2O_2. After this time,
2.4 µL of 6.6 mM 5-methyltetrahydrofolate (MeTHF) was added to initiate the catalytic cycle
of MTR, thus producing tetrahydrofolate. After 10 minutes at 37 °C, the reaction was quenched
by adding 10 µL of 5 M HCl in aqueous 60% formic acid and immediately heated at 80 °C for
10 minutes to convert the produced tetrahydrofolate to methenyltetrahydrofolate, which
concentration can be determined using $\Delta\varepsilon_{350} = 26{,}500$ M^{-1} cm^{-1}. Absorbance was recorded
on a OmegaSTAR (BMG Biotech), which were pathlength corrected. The data was analysed
and plotted on GraphPad Prism 10.

792
Thermal shift assay
MTR proteins were diluted in thermal shift buffer (25 mM HEPES, pH 7.5, 200 mM NaCl) to
0.4 mg/ml with SYPRO-Orange (Invitrogen) diluted 1000X in a total volume of 20 µl. When
appropriate, ligands were added at a final concentration of 1 mM, except for HOCbl, CNCbl,
and MeCbl which were added at a final concentration of 50 µM. A QuantStudio 3 RT-PCR
machine (Thermo Fisher Scientific) was used to monitor protein melting across 25 °C – 95 °C
with a heating rate of 0.5 °C per min. Data were analysed in GraphPad Prism software, and
the thermal melting temperatures (T_m) were determined using a nonlinear Boltzmann
equation.

802
Negative stain electron microscopy
Purified recombinant monomeric MTR^{FL} from size exclusion chromatography was diluted to a
final concentration of 40 nM in buffer (25 mM HEPES pH 7.5, 250 mM NaCl). 10 µl sample
was added to copper grids with carbon coating (300 mesh, Quantifoil Micro Tools GmbH),
plasma treated for 1 min using PELCO easiGlow™ (Ted Pella, Inc). After 30 seconds excess
sample was dried off using filter paper (Whatman), and the grid was placed on a drop of filtered
1.5 % uranyl acetate (Sigma-Aldrich) for 30 seconds. Grids were dried again using filter paper
before visualisation. Imaging was performed using a Hitachi HT7800 microscope operating at
120 kV and with a magnification of 60k, located at the Electron Microscopy Research Services
(EMRS) facility at Newcastle University.

813
Grid preparation and cryo-electron microscopy
MTR^{FL} was diluted to 0.5 mg/ml into 25 mM HEPES, pH 7.5, 250 mM NaCl with either 1 mM
MeTHF, 1 mM MeTHF, 1 mM SAM and 50 µM HOCbl, or 50 µM MeCbl. Grids were prepared
using a FEI Vitrobot Mark III (Thermo Fisher Scientific) at 4 °C and 100% humidity. 3 µl of
sample was applied to a plasma treated R1.2/1.3 300 mesh UltrAuFoil (Quantifoil) grid, with
a blot force of 0, a blot time of 3 seconds and a wait time of 10 seconds. All grids were initially
screened by collecting a small ≈ 1,000 micrograph dataset on a 200 kV Glacios microscope
equipped with a Falcon 4 direct electron detector (ThermoFisher Scientific) at York Structural
Biology Laboratory. Data was collected at a magnification of 240k with a pixel size of 0.574 Å
and a total dose of 50 e$^-$/Å^2. Glacios data was processed in cryoSPARC[45] and only grids that
resulted in clear 2D classes were sent for larger data collections on a 300 keV Krios
microscope.
All high-resolution data collection was carried out at Electron Bio-Imaging Centre
(eBIC, Diamond Light Source, UK). For MTR^{FL} with MeTHF, and MTR^{FL} with MeTHF, SAM
and HOCbl were collected on a 300 kV Titan Krios equipped with a K3 direct electron detector
and energy filter (Gatan). K3 data was collected at a magnification of 165k equating to a pixel
size of 0.508 Å (super resolution mode 0.254 Å) with an exposure time of 1.00 s, and a total
dose of 65.20 e$^-$/Å^2. For MTR^{FL} with MeCbl data was collected on a 300 kV Titan Krios
equipped with a Falcon 4i direct electron detector and selectris X energy filter (ThermoFisher
Scientific). Falcon 4i data was collected at a magnification of 215k equating to a pixel size of
0.576 Å with an exposure time of 2.46 s and a total dose of 70 e$^-$/Å^2.
Data processing for each dataset followed the same following general process: All
movies were imported into cryoSPARC[45] where they were subjected to patch motion
correction and patch CTF estimation. Micrographs were into 2,000 micrograph subsets for
particle picking, extraction, and 2D classification. All 2D classifications used the standard
preset parameters expect that force over max poses/shifts was turned off, the number of
online-EM iterations was increased from 20 to 40, the number of full EM iterations was
increased from 1 to 3, and the batch size was increased to 400. Initially blob picking was used
on each micrograph subset. Particles were extracted in a 400-pixel box Fourier cropped to
200 pixels and subjected to multiple rounds of 2D classification to remove bad picks and junk.
Once good 2D classes were determined it was clear that two types of particles were present
in each dataset. Separate 2D classes for these particle types were used for separate template-
based picking which were then again extracted in a 400-pixel box Fourier cropped to 200
pixels and subjected to rounds of 2D classification. Finally for both particle types, these
particles were used in combination with Topaz[46] to pick particles that were missed or poorly
centred by blob and template picking. Again, these particles were extracted in a 400-pixel box
Fourier cropped to 200 pixels and subjected to rounds of 2D classification. For each particle
type particles from blob, template, and Topaz[46] based picking were separately pooled,
duplicates removed and subjected to another round of 2D classification. Ab-initio with three to
six classes followed by heterogenous refinement was used to determine the best particles that
were then subjected to non-uniform refinement followed by local refinement. Reference based
motion correction was then used followed by another round of ab-initio, heterogenous, non-
uniform, and local refinement. In the case of the MTR^{C-Half} maps 3D variability analysis[30] was
carried out with 3 components filtered to 5.0 Å. For the MTR^{C-Half} map from the MeTHF, SAM,
and HOCbl dataset 3D variability was done in cluster mode followed by local refinement. One
cluster equated to the His-off state whereas one cluster equated to the His-on state. As none
of the maps reached Nyquist the extracted particle images were left Fourier cropped meaning
that maps from the K3 datasets have a pixel size of 1.016 Å and the Falcon 4i maps have a
pixel size of 1.152 Å. More details are shown in the supplementary information
(Supplementary Fig. 3, 5, and 8). Local resolution, conical FSC area ratio (cFR), and sampling
compensation factor (SCF) were determined within cryoSPARC[45] for each map
(Supplementary Fig. 9).
866
Atomic model fitting, refinement, and validation

For all MTR structures described we used the predicted AlphaFold structure of MTR^{FL}
([Q99707](#))[47] as an initial starting model. For each map the appropriate domains were manually
fitted and docked using ChimeraX[48]. Isolde[49] global flexible fitting was then carried out followed
by real space refinement as implemented in Phenix[50]. Ramachandran outliers were inspected
and manually corrected in Coot[51] followed by a final Phenix real space refinement. For the
ligands methyltetrahydofolate and cobalamin, Phenix eLBOW[52] was used to produce their
restraints. Each ligand was manually docked into their appropriate density using Coot and
were subjected to rounds of manual refinement and Phenix real space refinement. All models
were validated using Molprobity[53]. Model fit was assessed by model to map FSC determination
in Phenix[50] and using the Q-score[54] implementation in ChimeraX[48] (Supplementary Fig. 10)
878

Structure prediction using AlphaFold3
The AlphaFold3 server[36] was used to predict the complex structures of MTR ([Q99707-1](#)),
MTRR ([Q9UBK8-2](#)), MMACHC ([Q9Y4U1](#)), and MMADHC ([Q9H3L0](#)) using their amino acid
sequences as input. The sequences of MTR from *Thermus thermophilus* ([Q9RA53](#)), *T.*
filiformis ([A0A0A2XCD7](#)), and *Escherichia coli* ([P13009](#)) were also used for the AF3 complex
predictions. In some incidences the sequences corresponding to the boundaries of MTR^{N-Half},
MTR^{C-Half}, and MTRR^{ΔFMN} were used. All five output models for each prediction were manually
inspected in ChimeraX[48]. pLDDT[55] values from the server were used to analyse the confidence
of modelling for individual proteins. Both pTM and ipTM scores calculated from the server
were used to judge the confidence of each predicted complex structure[36,56,57].
889

890 **Blue native-PAGE**
891 Blue native-PAGE was carried out according to manufacturer's instructions (Life
892 Technologies). Both MTR^{FL} and the MTR^{FL}-MTRR^{FL} complex expressed and purified from
893 insect cells were diluted to 1.0 mg/ml in 25 mM HEPES, pH 7.5, 250 mM NaCl.
894

Talon pull-down assay
The bait, MTRR^{FL} with a N-terminal His-tag at 0.5 mg ml^{-1} was mixed with either prey proteins
MTR^{FL}, MTR^{N-Half} or MTR^{C-Half} without His-tags at 0.5 mg ml^{-1} in a total volume of 100 µl diluted
in binding buffer (25 mM HEPES, pH 7.5, 100 mM NaCl, 1 mM TCEP, 0.2% Tween-20).
Protein mixes were incubated with rotation at room temperature for 30 min. Next, 80 µl of a
50% slurry of talon resin (Clontech) in binding buffer was added, followed by incubation for a
further 30 min at room temperature. The resin was washed with 2 ml of binding buffer with 10
mM imidazole and eluted with 40 µl 4× sodium dodecyl sulfate (SDS)–PAGE sample buffer.
Controls consisted of adding prey to resin without bait. All samples were run on SDS–PAGE
and stained with Coomassie blue. Each experiment was replicated two times.
905

Glutathione pull-down assay
The bait proteins, MMACHC, MMADHC^{ΔN} or MTRR^{ΔFMN} each with a N-terminal GST-tag at
0.3 mg ml^{-1} were mixed with either prey proteins MTR^{FL}, MTR^{N-Half} or MTR^{C-Half} without GST-
tags at 0.3 mg ml^{-1} in a total volume of 100 µl diluted in binding buffer (25 mM HEPES, pH 7.5,
100 mM NaCl, 1 mM TCEP, 0.2% Tween-20). Protein mixes were incubated with rotation at
room temperature for 30 min. Next, 200 µl of a 50% slurry of glutathione sepharose 4 fast flow
resin (Cytiva) in binding buffer was added, followed by incubation for a further 60 min at room
temperature. The resin was washed with 2 ml of binding buffer and eluted with 40 µl 4× sodium
dodecyl sulfate (SDS)–PAGE sample buffer. Controls consisted of adding prey to resin without
bait. All samples were run on SDS–PAGE and stained with Coomassie blue. Each experiment
was replicated two times.
917

Cobalamin loading assay
Cobalamin loading from MMADHC to MTR was carried out as similar to described[17]. 100 µM
of HOCbl was mixed with 120 µM of MMADHC^{ΔN} in KCl Buffer (100 mM HEPES, 150 mM KCl,
10% glycerol) for a final volume of 40 µL. The reaction was incubated at room temperature for
1 hour, and the UV-Vis spectrum was recorded every 3 minutes to check the reaction. To

detect the cobalamin transfer from MMADHC$^{\Delta N}$ to MTR^{FL}, MMADHC$^{\Delta N}$ bound was
concentrated by multiple rounds of centrifugation (4000 xg, 4 °C) in Amicon® concentrators
with a 10 kDa molecular weight cut-off (Sigma Aldrich) followed by adding more KCl buffer to
remove free HOCbl. 15 µM MMADHC$^{\Delta N}$ -HOCbl was then incubated with 8.8 µM of MTR in a
final volume of 100 µL of KCl buffer for 1 hour at room temperature. After this time, the sample
was loaded into a Superose® 6 10/300 GL gel filtration column (Cytiva) equilibrated with KCl
buffer at 4 °C at a flow rate of 0.2 ml/min. The elution profile was monitored at 280 and 350
nm to detect the protein and cobalamin respectively. For binding of HOCbl and MeCbl, 8.8 µM
of MTR^{FL} was incubated with 100 µM Cbl for and the applied onto the gel filtration column. All
experiments were performed twice except for MeCbl binding which was carried out only once.

Analytical size exclusion chromatography
Analytical size exclusion chromatography was carried out using a 10/300 GL S200 Increase
column (Cytiva) equilibrated in 25 mM HEPES, pH 7.5, 150 mM NaCl. Proteins were mixed
as appropriate in 25 mM HEPES, pH 7.5, 150 mM NaCl to a final concentration of each protein
at 1.5 mg/ml in a total volume of 250 ul. After 30 minutes incubation rotating at room
temperature each were loaded onto the size-exclusion column and eluted with a flow rate of
0.3 ml/min. 500 µl fractions were collected for analysis by SDS-PAGE.

Data availability
The authors declare that the main data supporting the findings of this study are available within
the article and Supplementary Information. EM maps and models generated in this study of
Human Methionine Synthase With Methyltetrahydrofolate, N-Half From Full-Length (EMDB-
55190, PDB 9SSP), Human Methionine Synthase With Methyltetrahydrofolate, C-Half From
Full-Length (EMDB-55191, PDB 9SSQ), Human Methionine Synthase With
Methyltetrahydrofolate, Hydroxocobalamin, and SAM, N-Half From Full-Length (EMDB-
55192, PDB 9SSR), Human Methionine Synthase With Methyltetrahydrofolate,
Hydroxocobalamin, and SAM, C-Half His-ON From Full-Length (EMDB-55193, PDB 9SSS),
Human Methionine Synthase With Methyltetrahydrofolate, Hydroxocobalamin, and SAM, C-
Half His-OFF From Full-Length (EMDB-55194, PDB 9SST), Human Methionine Synthase With
Methylcobalamin, N-Half From Full-Length (EMDB-55195, PDB 9SSU) and Human
Methionine Synthase With Methylcobalamin, Activation Domain From Full-Length (EMDB-
55196, PDB 9SSV), have been deposited to the Electron Microscopy Data Bank (EMDB) and
Protein Data Bank (PDB) Other structures referenced in this article are indicated, including
PDB ID 4CCZ, 8G3H, 8SSC, 9CBO, 9CBP, 9CBQ, and 9CBR. Source data are provided with
this paper.

Acknowledgements
Negative stain electron microscopy was done at Newcastle University EM Research Services,
and we are grateful to Tracey Davey for technical support. The Newcastle University EM
Research Services is funded by the BBSRC (grant number BB/R013942/1). We thank Johan
Turkenburg and Sam Hart for their assistance in collecting Glacios data at York Structural
Biology Laboratory (YSBL). The YSBL is funded by the BBSRC, the Wellcome Trust (grant
number 206161/Z/17/Z), Tony Wild, and the Wolfson Foundation. We acknowledge the
Diamond Light Source for access and support to the UK's Electron Bio-imaging Centre (eBIC,
under BAG proposal BI34172) funded by the Wellcome Trust, MRC, and BBRSC. We
specifically want to thank David Owen and David Farmer for their assistance in data collecting.
D.S.M.F. was supported by the Nuffield Department of Medicine Prize Studentship at the
University of Oxford. T.J.M. received cryo-EM training through the Wellcome/MRC funded
programme (218785/Z/19/Z). Initial work on this project was carried out at the Structural
Genomics Consortium, a registered charity (Number 1097737) that receives funds from
AbbVie, Bayer Pharma AG, Boehringer Ingelheim, Canada Foundation for Innovation,
Eshelman Institute for Innovation, Genome Canada, Innovative Medicines Initiative
(EU/EFPIA) [ULTRA-DD grant no. 115766], Janssen, Merck & Co., Novartis Pharma AG,

Author contributions

D.S.M.F., J.M.E., J.A.C., W.W.Y., and T.J.M. designed the experiments. D.S.M.F. expressed and purified MTR and MTRR constructs, carried out biochemical experiments, screened, and collected EM data, analysed, and refined all cryo-EM structures. K.M., C.D., H.J.B., and J.K. purified MTR constructs and did preliminary experiments. W.K. expressed, purified, and crystallized the N-terminal MTR construct. M.V. solved and refined the N-terminal MTR construct crystal structure. D.S.F. expressed and purified MMADHC and MMACHC. T.J.M. expressed and purified MMACHC, MTRR and MTR proteins along with guiding cryo-EM screening, data collection, data processing, AF3 predictions, and biochemical experiments. R.C. carried out the native mass spectrometry. A.B. supplied computational resources for cryo-EM data processing. D.S.M.F., W.W.Y., and T.J.M. supervised the work, carried out data analysis, and wrote the manuscript with contributions from all authors.

Competing interests

The authors declare no competing interests.

Correspondence and requests for materials should be addressed to Wyatt W. Yue or Thomas J. McCorvie

References

1. Banerjee, R.V. & Matthews, R.G. Cobalamin-dependent methionine synthase. *The FASEB journal* **4**, 1450-1459 (1990).
2. Chiang, P.K. et al. S-Adenosylmethionine and methylation. *FASEB J* **10**, 471-80 (1996).
3. Ghergurovich, J.M. et al. Methionine synthase supports tumour tetrahydrofolate pools. *Nat Metab* **3**, 1512-1520 (2021).
4. Sullivan, M.R. et al. Methionine synthase is essential for cancer cell proliferation in physiological folate environments. *Nat Metab* **3**, 1500-1511 (2021).
5. Wright, A.J., Dainty, J.R. & Finglas, P.M. Folic acid metabolism in human subjects revisited: potential implications for proposed mandatory folic acid fortification in the UK. *Br J Nutr* **98**, 667-75 (2007).
6. Froese, D.S., Fowler, B. & Baumgartner, M.R. Vitamin B12, folate, and the methionine remethylation cycle—biochemistry, pathways, and regulation. *Journal of inherited metabolic disease* **42**, 673-685 (2019).
7. McCorvie, T.J., Ferreira, D., Yue, W.W. & Froese, D.S. The complex machinery of human cobalamin metabolism. *Journal of Inherited Metabolic Disease* **46**, 406-420 (2023).
8. Matthews, R.G., Koutmos, M. & Datta, S. Cobalamin-dependent and cobamide-dependent methyltransferases. *Current opinion in structural biology* **18**, 658-666 (2008).
9. Evans, J.C. et al. Structures of the N-terminal modules imply large domain motions during catalysis by methionine synthase. *Proceedings of the National Academy of Sciences* **101**, 3729-3736 (2004).
10. Bandarian, V. et al. Domain alternation switches B12-dependent methionine synthase to the activation conformation. *Nature structural biology* **9**, 53-56 (2002).
11. Drennan, C.L., Huang, S., Drummond, J.T., Matthews, R.G. & Ludwig, M.L. How a protein binds B12: a 3.0 Å X-ray structure of B12-binding domains of methionine synthase. *Science* **266**, 1669-1674 (1994).

1029   12.   Bassila, C. et al. Methionine synthase and methionine synthase reductase interact with
1030   MMACHC and with MMADHC. *Biochimica et Biophysica Acta (BBA)-Molecular Basis*
1031   *of Disease* **1863**, 103-112 (2017).

1032   13.   Froese, D.S. et al. Structure of MMACHC reveals an arginine-rich pocket and a
1033   domain-swapped dimer for its B12 processing function. *Biochemistry* **51**, 5083-5090
1034   (2012).

1035   14.   Froese, D.S. et al. Structural insights into the MMACHC-MMADHC protein complex
1036   involved in vitamin B12 trafficking. *Journal of Biological Chemistry* **290**, 29167-29177
1037   (2015).

1038   15.   Mascarenhas, R. et al. An unusual Co-S bond links B(12) chaperones in an interprotein
1039   complex. *Proc Natl Acad Sci U S A* **122**, e2507700122 (2025).

1040   16.   Li, Z. et al. An interprotein Co–S coordination complex in the B12-trafficking pathway.
1041   *Journal of the American Chemical Society* **142**, 16334-16345 (2020).

1042   17.   Mascarenhas, R. et al. Cobalt–Sulfur Coordination Chemistry Drives B12 Loading onto
1043   Methionine Synthase. *Journal of the American Chemical Society* **145**, 24678-24689
1044   (2023).

1045   18.   Wolthers, K.R., Basran, J., Munro, A.W. & Scrutton, N.S. Molecular dissection of
1046   human methionine synthase reductase: determination of the flavin redox potentials in
1047   full-length enzyme and isolated flavin-binding domains. *Biochemistry* **42**, 3911-3920
1048   (2003).

1049   19.   Wolthers, K.R. & Scrutton, N.S. Cobalamin uptake and reactivation occurs through
1050   specific protein interactions in the methionine synthase–methionine synthase
1051   reductase complex. *The FEBS Journal* **276**, 1942-1951 (2009).

1052   20.   Olteanu, H. & Banerjee, R. Human methionine synthase reductase, a soluble P-450
1053   reductase-like dual flavoprotein, is sufficient for NADPH-dependent methionine
1054   synthase activation. *Journal of Biological Chemistry* **276**, 35558-35563 (2001).

1055   21.   Hall, D.A. et al. Mapping the interactions between flavodoxin and its physiological
1056   partners flavodoxin reductase and cobalamin-dependent methionine synthase. *Proc*
1057   *Natl Acad Sci U S A* **98**, 9521-6 (2001).

1058   22.   Froese, D.S. & Gravel, R.A. Genetic disorders of vitamin B12 metabolism: eight
1059   complementation groups–eight genes. *Expert reviews in molecular medicine* **12**, e37
1060   (2010).

1061   23.   Copp, A.J., Stanier, P. & Greene, N.D. Neural tube defects: recent advances, unsolved
1062   questions, and controversies. *Lancet Neurol* **12**, 799-810 (2013).

1063   24.   Mendoza, J., Purchal, M., Yamada, K. & Koutmos, M. Structure of full-length
1064   cobalamin-dependent methionine synthase and cofactor loading captured in crystallo.
1065   *Nature Communications* **14**, 6365 (2023).

1066   25.   Yamada, K., Gravel, R.A., Toraya, T. & Matthews, R.G. Human methionine synthase
1067   reductase is a molecular chaperone for human methionine synthase. *Proceedings of*
1068   *the National Academy of Sciences* **103**, 9476-9481 (2006).

1069   26.   Drummond, J.T., Jarrett, J., Gonzalez, J.C., Huang, S. & Matthews, R.G.
1070   Characterization of nonradioactive assays for cobalamin-dependent and cobalamin-
1071   independent methionine synthase enzymes. *Analytical biochemistry* **228**, 323-329
1072   (1995).

1073   27.   Watkins, M.B., Wang, H., Burnim, A. & Ando, N. Conformational switching and
1074   flexibility in cobalamin-dependent methionine synthase studied by small-angle X-ray
1075   scattering and cryoelectron microscopy. *Proceedings of the National Academy of*
1076   *Sciences* **120**, e2302531120 (2023).

1077   28.   Datta, S., Koutmos, M., Pattridge, K.A., Ludwig, M.L. & Matthews, R.G. A disulfide-
1078   stabilized conformer of methionine synthase reveals an unexpected role for the
1079   histidine ligand of the cobalamin cofactor. *Proceedings of the National Academy of*
1080   *Sciences* **105**, 4115-4120 (2008).

1081   29.   Koutmos, M., Datta, S., Pattridge, K.A., Smith, J.L. & Matthews, R.G. Insights into the
1082   reactivation of cobalamin-dependent methionine synthase. *Proceedings of the*
1083   *National Academy of Sciences* **106**, 18527-18532 (2009).

30. Punjani, A. & Fleet, D.J. 3D variability analysis: Resolving continuous flexibility and discrete heterogeneity from single particle cryo-EM. *J Struct Biol* **213**, 107702 (2021).

31. Fleischhacker, A.S. & Matthews, R.G. Ligand trans influence governs conformation in cobalamin-dependent methionine synthase. *Biochemistry* **46**, 12382-12392 (2007).

32. Jarrett, J.T., Huang, S. & Matthews, R.G. Methionine synthase exists in two distinct conformations that differ in reactivity toward methyltetrahydrofolate, adenosylmethionine, and flavodoxin. *Biochemistry* **37**, 5372-5382 (1998).

33. Mendoza, J., Yamada, K., Castillo, C., Wilhelm, C.A. & Koutmos, M. Orchestrating Improbable Chemistries: Structural Snapshots of B12-Dependent Methionine Synthase's Catalytic Choreography. *bioRxiv*, 2024.08.29.610163 (2024).

34. Wolthers, K.R. et al. Crystal structure and solution characterization of the activation domain of human methionine synthase. *The FEBS Journal* **274**, 738-750 (2007).

35. Fofou-Caillierez, M.a.B. et al. Interaction between methionine synthase isoforms and MMACHC: characterization in cblG-variant, cblG and cblC inherited causes of megaloblastic anaemia. *Human molecular genetics* **22**, 4591-4601 (2013).

36. Abramson, J. et al. Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature*, 1-3 (2024).

37. Li, Z. et al. An Interprotein Co–S Coordination Complex in the B_{12}-Trafficking Pathway. (2020).

38. Wolthers, K.R. & Scrutton, N.S. Protein interactions in the human methionine synthase− methionine synthase reductase complex and implications for the mechanism of enzyme reactivation. *Biochemistry* **46**, 6696-6709 (2007).

39. Jarrett, J.T., Hoover, D.M., Ludwig, M.L. & Matthews, R.G. The mechanism of adenosylmethionine-dependent activation of methionine synthase: a rapid kinetic analysis of intermediates in reductive methylation of cob (II) alamin enzyme. *Biochemistry* **37**, 12649-12658 (1998).

40. Bandarian, V., Ludwig, M.L. & Matthews, R.G. Factors modulating conformational equilibria in large modular proteins: a case study with cobalamin-dependent methionine synthase. *Proceedings of the National Academy of Sciences* **100**, 8156-8163 (2003).

41. Banerjee, R.V., Frasca, V., Ballou, D.P. & Matthews, R.G. Participation of cob(I) alamin in the reaction catalyzed by methionine synthase from Escherichia coli: a steady-state and rapid reaction kinetic analysis. *Biochemistry* **29**, 11101-9 (1990).

42. Guha, A. & Banerjee, R. The common homocystinuria-associated P1173L variant of human methionine synthase impairs reductive methylation. *J Biol Chem* **301**, 110366 (2025).

43. Suormala, T. et al. The cblD defect causes either isolated or combined deficiency of methylcobalamin and adenosylcobalamin synthesis. *Journal of Biological Chemistry* **279**, 42742-42749 (2004).

44. Haque, M.M. et al. Distinct conformational behaviors of four mammalian dual-flavin reductases (cytochrome P450 reductase, methionine synthase reductase, neuronal nitric oxide synthase, endothelial nitric oxide synthase) determine their unique catalytic profiles. *FEBS J* **281**, 5325-40 (2014).

45. Punjani, A., Rubinstein, J.L., Fleet, D.J. & Brubaker, M.A. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nature methods* **14**, 290-296 (2017).

46. Bepler, T. et al. Positive-unlabeled convolutional neural networks for particle picking in cryo-electron micrographs. *Nat Methods* **16**, 1153-1160 (2019).

47. Varadi, M. et al. AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res* **50**, D439-D444 (2022).

48. Pettersen, E.F. et al. UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein science* **30**, 70-82 (2021).

49.   Croll, T.I. ISOLDE: a physically realistic environment for model building into low-
resolution electron-density maps. *Acta Crystallographica Section D: Structural Biology*
74, 519-530 (2018).
50.   Terwilliger, T.C. et al. Iterative model building, structure refinement and density
modification with the PHENIX AutoBuild wizard. *Acta Crystallographica Section D:*
Biological Crystallography **64**, 61-69 (2008).
51.   Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta*
Crystallogr D Biol Crystallogr **60**, 2126-32 (2004).
52.   Moriarty, N.W., Grosse-Kunstleve, R.W. & Adams, P.D. electronic Ligand Builder and
Optimization Workbench (eLBOW): a tool for ligand coordinate and restraint
generation. *Acta Crystallogr D Biol Crystallogr* **65**, 1074-80 (2009).
53.   Chen, V.B. et al. MolProbity: all-atom structure validation for macromolecular
crystallography. *Acta Crystallogr D Biol Crystallogr* **66**, 12-21 (2010).
54.   Pintilie, G. et al. Measurement of atom resolvability in cryo-EM maps with Q-scores.
Nat Methods **17**, 328-334 (2020).
55.   Jumper, J. et al. Highly accurate protein structure prediction with AlphaFold. *Nature*
596, 583-589 (2021).
56.   Zhang, Y. & Skolnick, J. Scoring function for automated assessment of protein
structure template quality. *Proteins* **57**, 702-10 (2004).
57.   Xu, J. & Zhang, Y. How significant is a protein structure similarity with TM-score = 0.5?
Bioinformatics **26**, 889-95 (2010).
1158