Biological sex affects human islet gene expression and mitochondrial function in type 2

diabetes

Sing-Young Chen[1,* †], Haoning Howard Cen[1], Charlotte F. Chao [1], Andrew R. Pepper[2], James D.

Johnson[1,†], Elizabeth J. Rideout[1, †]

[1] Life Sciences Institute, Department of Cellular and Physiological Sciences, The University of

British Columbia, Vancouver, BC, Canada

[2] Department of Surgery, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, AB,

Canada.

* first author

† co-corresponding authors

Email addresses:

Sing-Young Chen: singyoung.chen@ubc.ca, ORCiD: 0000-0001-5258-8254

James D. Johnson: james.d.johnson@ubc.ca, ORCiD: 0000-0002-7523-9433

Elizabeth J. Rideout: elizabeth.rideout@ubc.ca, ORCiD: 0000-0003-0012-2828

Keywords: pancreatic islet, type 2 diabetes, sex differences, beta-cell, alpha-cell, insulin

Grants/fellowships supporting the writing of this paper: S-Y.C. was supported by the J.

Andrew McKee Postdoctoral Fellowship (JDRF and Stem Cell Network), Canadian Institute of

Health Research (CIHR) Postdoctoral Fellowship, and Michael Smith Health Research BC

Trainee Award. H.H.C. was supported by a Breakthrough T1D Postdoctoral Fellowship. S-Y.C.,

H.H.C., and J.D.J. acknowledge support from grant #3-COE-2022-1103-M-B from the

Breakthrough T1D Canucks for Kids Fund Centre of Excellence at UBC. E.J.R. gratefully

acknowledges funding from the Canadian Institutes of Health Research (GS4-171365),

Diabetes Canada (OG-3-22-5646-ER), and J.D.J. and E.J.R. acknowledge team grant funding

Disclosure summary: All authors declare no competing interests.

Abstract

The clinical characteristics of type 2 diabetes (T2D) differ between the sexes. For example, the

risk of T2D is higher in males than in premenopausal females, whereas the risk of T2D-

associated cardiovascular disease is higher in females. However, the sex-dependent

mechanisms of T2D pathogenesis remain incompletely understood. Publicly available human

islet datasets, such as HPAP and Humanislets.com, offer a valuable tool for uncovering the

impact of biological sex on islet structure, gene expression, and function at a scale that was not

previously possible. We performed an integrated analysis of data from publicly available

sources to identify sex differences in baseline islet characteristics in donors without diabetes

and subsequently examined these features in donors who lived with T2D. Among donors

without diabetes, female islets had a greater proportion of alpha-cells compared with male islets

and showed enriched expression of ribosomal and mitochondrial pathways in both beta- and

alpha-cells. Measurements of mitochondrial function in female islets revealed lower spare

respiratory capacity compared to male islets. Male and female islets had distinct changes in

gene and protein expression in the context of T2D with female islets having greater preservation

of insulin content and fewer defects in islet function. Together, these data show female islets

have fewer islet impairments in T2D. This highlights the need for detailed mechanistic studies in

both sexes to support effective and sex-informed interventions for T2D.

Introduction

Biological sex influences type 2 diabetes (T2D) incidence, complications, and treatment efficacy. There is a higher diabetes risk in men until approximately 50-60 years of age [1–4]. This male bias has been observed across all sociodemographic indices [5], though we note a female bias in some populations [6–8]. Relative protection against diabetes in pre-menopausal women has been attributed in part to the protective effects of estrogen [9,10]. Despite the lower incidence of T2D in premenopausal women, women who live with diabetes are at a higher risk of most T2D complications (e.g. cardiovascular and renal) compared to men [11–18]. Men and women with T2D also show differential responses to commonly used medications: male sex predicts better HbA1c-lowering efficacy of sulfonylureas, which stimulate beta-cell insulin secretion [19,20], and SGLT2 inhibitors, which decrease renal glucose reabsorption [20,21], whereas insulin-sensitizing thiazolidinediones and GLP-1 receptor agonists (GLP-1RAs) have greater HbA1c-lowering effects in females [20–24].

Sex differences in clinical observations are likely due to a combination of differences in insulin sensitivity and pancreatic beta-cell function. Greater insulin sensitivity in females is well-established and conserved from humans [25] to mice [26] to fruit flies [27]. Recent studies suggest that sex differences in pancreatic islets may also play a role. In mice, female islets show greater resilience to endoplasmic reticulum stress than male islets and have higher expression of genes involved in protein synthesis [28,29]. Sex differences in electrical activity have also been observed in mouse islets: glucose stimulation in female beta-cells results in lower potassium currents leading to a more depolarized membrane potential, more frequent burst-type action potentials, and smaller glucose-induced transient calcium increases [30,31]. HFD feeding in mice also causes sex-dependent transcriptomic changes [28], and female mice are better able to adapt to this metabolic stress through alpha-to-beta cell communication to maintain beta-cell calcium dynamics [32]. In humans, as in mice, female islets show enriched expression of pathways related to protein synthesis [29,33]. Beta-cell gene expression is closely linked to epigenetic modifications,

which differ between sexes on both sex chromosomes and autosomes, including at T2D

candidate genes [34,35]. *Ex vivo* insulin secretion studies have also suggested that female islets

may secrete more insulin than male islets in response to high glucose [29,33,34]. T2D causes

profound changes in gene expression among islets from both sexes, and these changes are

also partly sex-dependent [33,35,36]. However, our understanding of the molecular and functional

changes that occur with T2D in female and male islets remains far from complete.

The Human Pancreas Analysis Program (HPAP) consolidates data from donor human

islets across multiple institutes. The data collected through this program are uploaded to the

publicly available PancDB database, and include measurements of human islet calcium imaging,

mass cytometry, islet perifusion, oxygen consumption, and scRNA-seq, among others [37,38].

Humanislets.com is a repository with functional and omics data on a largely distinct pool of

donors, including a wider range of perifusion experiments and a large human islet proteomics

dataset [39]. Importantly, the datasets have enough donors to test for sex differences in many islet

and islet cell attributes.

Our analysis of HPAP and Humanislets.com revealed pronounced sex differences in islet

composition, RNA and protein levels, and mitochondrial function in islets from donors without

diabetes. We further find that gene and protein expression are differentially altered in T2D

depending on biological sex, and that male islets exhibited mitochondrial defects that were

absent in female islets among donors with T2D. Together, these data demonstrate clear

differences in diverse islet characteristics between male and female islets among individuals

without T2D, and show sex-dependent changes to islet biology in individuals who lived with T2D.

Materials and Methods

Data sources

Publicly available data were downloaded from the Human Pancreas Analysis Program (HPAP)

Database, consortia under Human Islet Research Network (RRID:SCR_014393,

https://hpap.pmacs.upenn.edu/) [37,38] (NIH grant numbers UC4-DK112217 and UC4-DK112232,

RRID:SCR_016202), and Humanislets.com [39], an initiative of the Alberta Diabetes Institute

IsletCore funded by the Canadian Institutes of Health Research, Breakthrough T1D Canada,

and Diabetes Canada (5-SRA-2021-1149-S-B/TG 179092). Specific datasets used are

described in detail below. To study sex differences within the healthy islet state, we specifically

focused on donors without diabetes and aged 15-39 years. This age bracket was chosen to

determine sex differences prior to the onset of menopause [40] and was used to compare male

and female control (non-diabetic) donors only. However, when comparing across control and

T2D for males and females, donors of all ages were included. For age-matched data, donors

with T2D were matched to donors without diabetes with the closest possible age. Donors with

type 1 diabetes were not included in our analyses due to insufficient donor numbers to support

sex-specific analysis. Histograms displaying key metadata variables in each dataset are shown

in Figure S1.

Cell type composition

Islet endocrine cell type composition data were available from both HPAP and Humanislets.com.

CyToF data [41] estimating cell counts for each cell type were obtained from donor metadata

available on HPAP, from which endocrine cell type proportions were calculated as described in

the CyTOF Staining Workflow and Protocol on HPAP

(https://hpap.pmacs.upenn.edu/explore/workflow/imaging-mass-cytometry?protocol=1, protocol

accessed 23 September 2025, last updated 11 January 2021). Donors with a total endocrine

cell count less than 30-fold lower than the geometric mean of all donors were excluded, as in [42].

On Humanislets.com, endocrine cell type proportions were estimated based on deconvolution of

whole islet proteomics data [39] and were downloaded directly.

Gene expression analysis

Gene expression data from HPAP and Humanislets.com were analyzed to compare the sexes.

Transcriptomic data from HPAP included scRNAseq data. To analyze scRNAseq data, whole

islet, beta-cell, and alpha-cell expression were first pseudobulked to generate one gene

expression profile per donor per cell type using the *Seurat* package [43]. Pseudobulking

overcomes the confounding effects of uneven cell numbers per donor and appropriately treats

individual donors, rather than cells, as independent observations. For Humanislets.com, only

bulk islet RNAseq was analyzed by sex due to low n's available for scRNAseq data. Bulk islet

proteomics data [44] were also downloaded and analyzed from Humanislets.com. Three

comparisons were performed: female vs male among donors without diabetes aged 15-39,

female control (without diabetes) vs female T2D across all ages, and male control vs male T2D

across all ages. For all differential expression analyses, age was included as a covariate.

For whole islet transcriptomics, HPAP pseudobulk data and Humanislets.com bulk RNAseq

data were combined and analyzed using *limma* [45], with age and dataset as covariates. For

HPAP pseudobulk beta-cell, HPAP pseudobulk alpha-cell, and Humanislets.com proteomics

data, *limma* [45] was used with age as a covariate. Gene Set Enrichment Analysis (GSEA) was

performed on differential expression analysis output to identify key pathways using the

clusterProfiler package [46], in which the direction signed -log10 *p*-values were used as the rank

scores of the genes. To improve pathway visualization, pathways were considered redundant if

their core enrichment genes exceeded a Jaccard index of 0.6. When selecting pathways for

visualization, redundant pathways were first refined by omitting the pathway with fewer gene

components. For visualization of pathways, we grouped redundant or similar pathways together

by using Leiden clustering[47] with Jaccard index as similarity measurement between pathways.

For comparisons between female and male control donors aged 15-39, pathways with adjusted

p-value < 0.05 were included in visualization. For comparisons between control and T2D donors,

pathways with adjusted *p*-value < 0.0001 were included in visualization, but all pathways with

adjusted *p*-value < 0.05 were included in tables.

Insulin content

Islet insulin content data were downloaded from perifusion datasets on HPAP. There were two

sets of islet perifusion data with corresponding insulin content data. The data from the University

of Pennsylvania (UPenn) were presented per islet using a lysis buffer containing EDTA and

detergent [48,49]. The data from Vanderbilt University were presented per islet equivalent (IEQ),

using acid-ethanol extraction to lyse cells and isolate insulin. Acid-ethanol extraction is widely

used in the field to extract insulin from biological samples at maximum yields [50–54], leveraging

the strong acid to avoid the insulin precipitation pH range of ~4.5-6.5 [51,55] and ethanol to

minimize the contaminating influences of exocrine impurities [56,57] while retaining insulin in

solution [58]. Humanislets.com also used acid-ethanol extraction and presented insulin content

data per IEQ. These data were obtained from the provided metadata. Given the consistent

methods and units, the Vanderbilt and Humanislets.com data were pooled for combined

analyses with dataset as an additional covariate.

Oxygen consumption

Islet oxygen consumption in response to stimuli was available from both HPAP and

Humanislets.com. On Humanislets.com, oxygen consumption data were obtained using a

Seahorse Bioanalyzer (Agilent) and calculated parameters were directly downloaded (spare

respiratory capacity = maximum uncoupled respiration – baseline, maximum respiratory

capacity = maximum uncoupled respiration – minimum non-mitochondrial respiration, response

to high glucose = mean respiration in 16.7 mM glucose – mean respiration in 3 mM glucose,

baseline = last basal respiration) [39]. As three readings per donor were available, outliers were

identified using Grubb's test and the average per donor was calculated. On HPAP, these data

were obtained using a chemical oxygen probe (Oxyphor G3). Raw data were downloaded and

parameters were calculated using the same formulae as used on Humanislets.com. The HPAP

and Humanislets.com data were combined to assess potential sex differences among donors

aged 15-39 without diabetes. Due to the low number of T2D donors in the Humanislets.com

dataset, only the HPAP dataset was used for assessing the impact of T2D status.

Intracellular calcium

Intracellular calcium imaging data using the ratiometric dye Fura-2 were available only from

HPAP. Raw data were downloaded and processed as follows: 1) only regions that responded to

high glucose were included (response counted if 16.7 mM glucose stimulated an increase in

baseline-corrected calcium signal of at least 50%); 2) after filtering for glucose response, signals

from multiple regions for each run were averaged; 3) runs without a specified time of first

stimulus addition or without high glucose were excluded; 4) to account for different time periods

between stimuli, only the first x minutes after stimulus addition were counted for each run,

where x is the minimum period of time between that stimulus and the next, for any of the runs; 5)

responses were averaged for each individual donor and summary statistics were calculated.

Dynamic hormone secretion

HPAP provides dynamic hormone secretion data from islet perifusion experiments performed at

two sites: UPenn and Vanderbilt. For the UPenn data, areas under the curve (AUC) were

calculated for the periods of perifusion during which islets were exposed to each stimulus, for

both insulin and glucagon secretion. For the Vanderbilt data, there was a visible delay between

stimulus addition and response. Therefore, the AUC calculated for each stimulus was

determined based on the time course tracings: 12-39 min for 16.7 mM glucose, 63-87 min for

16.7 mM glucose + IBMX, 93-111 min for 1.7 mM glucose + epinephrine, and 123-138 min for

KCl. Humanislets.com contains islet perifusion data measuring insulin secretion in response to

glucose, leucine, and a mix of the free fatty acids oleate and palmitate [44]. For each

Humanislets.com perifusion experiment, AUC were calculated for the periods of perifusion

during which islets were exposed to each stimulus. As the HPAP UPenn and Humanislets.com

datasets used equivalent units and similar glucose concentrations (3 mM for low glucose), these

were combined for analyses and summary statistics were taken, including mean insulin

secretion at 3 mM glucose, peak insulin secretion at high glucose (15 mM for Humanislets.com,

16.7 mM for HPAP UPenn), peak insulin secretion at 30 mM KCl, and stimulation index (ratio of

peak secretion at high glucose over mean secretion at 3 mM glucose).

Statistics

Unless otherwise specified, to determine statistical differences due to sex and disease,

ANCOVA testing was performed with age as covariate. When multiple datasets were combined,

dataset was also included as a covariate. Multiple comparisons were performed using Tukey's

correction and the *emmeans* R package. For age-matched data, donors with T2D were matched

to donors without diabetes with the closest possible age within each sex using the *MatchIt*

package, and statistical comparisons were performed to assess the effect of T2D status only. As

some significant age differences remained in some cases, age was still included as a covariate

for statistical analyses of age-matched data. For comparing distributions in Figure S1, the two-

sample Kolgomorov-Smirnov test was used. Unless otherwise mentioned, a *p*-value threshold of

0.05 (or adjusted *p*-valued threshold where relevant) was used. All relevant scripts for data

processing and graphing are available at https://github.com/singyoungchen/sex-differences-

human-islet-characteristics-.

Results

Female islets contain a greater proportion of alpha-cells and a smaller proportion of beta-cells

We first examined islet cell type composition in male and female donors without T2D aged 15-39 years. Female islets had a significantly higher alpha-cell proportion compared to male islets in both the HPAP and Humanislets.com datasets (Figure 1A-B). Consistent with increased alpha-cell proportion, beta-cell proportion was lower in female islets compared with male islets, a difference that was statistically significant in the Humanislets.com data and which showed a non-significant but similar trend (p=0.11) in the HPAP data (Figure 1C-D). The proportion of other endocrine cells was significantly higher in males than females in HPAP (Figure 1E) but not Humanislets.com (Figure 1F). When donors of all ages were considered together, however, the trends we observed were not consistent between the HPAP CyToF results and the Humanislets.com cell type proportion estimates. The HPAP data suggested that, among donors without diabetes of all ages, female islets showed higher alpha-cell (p<0.05, Figure 1G), lower beta-cell (p=0.072, Figure 1H), and lower non-alpha, non-beta cell (p=0.083, Figure 1I) proportions, consistent with observations from donors in the 15-39 age bracket. In contrast, when including data for donors of all ages from Humanislets.com, we observed no sex difference in endocrine cell proportions between female and male donors without T2D (Figure 1J-L).

We next considered endocrine cell proportions in donors who lived with T2D. The HPAP data showed that T2D was associated with a trend toward increased alpha-cell proportion (p=0.08) and a significant decrease in beta-cell proportion (p<0.05) in males but not females (Figure 1G-H). The HPAP data also showed a T2D-associated increase in the proportion of other endocrine cells in females but not males (Figure 1I). In contrast, the Humanislets.com data showed an increase in alpha-cell proportion and a decrease in beta-cell proportion in female, but not male, donors who lived with T2D (Figure 1J-K). In the Humanislets.com dataset, the proportion of other endocrine cells was significantly increased with T2D in males only (Figure 1L). These

trends remained in age-matched data (Figure S2). While the reason for this discrepancy is

unclear, it may reflect the different techniques used by each team to obtain cell proportion

estimates (antibody-based identification in HPAP, estimation based on marker protein

expression data in Humanislets.com). Thus, while alpha-cell proportion was elevated among

females in young donors without diabetes, we observed no consistent sex difference in donors

who lived with T2D across datasets.

Islets from young female donors show higher expression of protein synthesis genes

We next checked for sex differences in baseline gene expression among islets from donors

aged 15-39 without diabetes by combining pseudobulk scRNAseq data from HPAP with bulk

islet RNAseq from Humanislets.com. After conducting differential expression analysis to

compare female and male donors (Table 1, S1), we performed GSEA. We found 23 female-

biased pathways and 16 male-biased pathways (Figure 2A, Table S2). Pathways related to

ribosomal biogenesis and protein synthesis showed a strong female bias (Figure 2A, Table S2).

The potential for enhanced protein synthesis capacity in female islets has previously been

shown by us [29], and others [33], and is likely associated with the effects of estrogen in alleviating

ER stress and preserving protein synthesis [59–62]. Mitochondrial pathways were also female-

enriched (Figure 2A, Table S2). Male-biased pathways included those related to cell division

and histone modification (Figure 2A, Table S2). The HPAP scRNAseq data also allowed us to

perform sex-based analysis of beta-cell- and alpha-cell-specific pseudobulk data. Both cell

types showed a female-biased enrichment of ribosomal and mitochondrial gene pathways and a

male-biased enrichment of mitosis and histone modification pathways (Figure S3, Table S2).

Humanislets.com includes, to our knowledge, the largest open-source whole islet proteomics

dataset currently available. We conducted differential expression analysis to compare protein

levels in female and male donors without diabetes aged 15-39 (Table S3) followed by GSEA.

We found female-biased abundance of proteins related to ribosomes and mitochondria, as well

as lipid transport and the plasma membrane (Figure 2B, Table S4). Male-biased pathways

included those related to chromatin modification and transcriptional regulation (Figure 2B, Table

S4). Female-biased and male-biased pathways identified in proteomics included many similar

pathways as those identified in transcriptomics (Table 2). Thus, the female bias in expression of

proteins related to protein synthesis and mitochondrial function was highly consistent between

our RNAseq and proteomics data.

T2D is associated with differential shifts in gene and protein expression depending on sex

We next compared islet gene expression between donors with and without diabetes to evaluate

any sex bias in gene expression changes associated with T2D. In the combined transcriptomics

dataset, 3 genes were significantly altered with T2D in females and 431 in males (Table 3, S5-6).

GSEA identified more pathways significantly altered with T2D in males than in females (Table 4,

S7-8). Among the pathways increased with T2D in female islets were pathways related to

transcriptional regulation and histone modification (Figure 3A, Table S7). In male islets,

pathways related to ribosomes and inflammation were increased with T2D (Figure 3B, Table S8).

Pathways decreased with T2D in female islets were dominated by pathways such as ribosomal

and mitochondrial gene pathways (Figure 4A, Table S7). In males, T2D was associated with a

decrease in pathways related to mitochondria, vesicle processing and secretion, and

microtubules (Figure 4B, Table S8). Our data therefore suggest that there were distinct T2D-

associated changes in gene expression between the sexes in whole islets.

We similarly analyzed beta-cell and alpha-cell pseudobulk scRNAseq data from HPAP. In

females, T2D was associated with positive enrichment of transcriptional regulation pathways

and negative regulation of mitochondrial and ribosomal pathways in both cell types (Figure S4A-

B, Table S7). In males, T2D was associated with an increase in RNA splicing and ribosome

pathways, and a decrease in angiogenesis and extracellular matrix pathways in beta-cells

(Figure S4C, Table S8). In alpha-cells among males, no significant pathways were detected at

the $p < 0.0001$ threshold (Figure S4D), suggesting that T2D may be associated with greater

changes in alpha-cell gene expression in female islets than in male islets.

At the protein level, T2D was associated with altered expression of 671 proteins in female

islets and 712 proteins in male islets (Table 3, Table S9-10). GSEA showed that T2D was

associated with enrichment of pathways related to cell adhesion in both sexes (Figure 3A-B,

Table S11-12). In female islets, T2D was also associated with enrichment of pathways related to

the inflammatory response and cytoskeleton organization (Figure 3A, Table S11). In male islets,

T2D was associated with an enrichment of cytosolic ribosomal pathways (Figure 3B, Table S12),

consistent with RNAseq data (Figure 3B, Table S8). Pathways that were decreased with T2D

were similar between males and females, and included those related to vesicle trafficking and

exocytosis, as well as mitochondrial components (Figure 4A-B, Table S11-12). Unlike in non-

T2D islets, pathways identified based on protein expression largely did not overlap with

pathways from our gene expression data, highlighting the importance of collecting multiple data

types. Taken together, our analysis reveals sex differences in both islet gene and protein

expression in donors without diabetes and individuals who lived with T2D.

Female islets may show greater preservation of insulin content than male islets in T2D

Healthy beta-cells have a remarkable capacity to rapidly increase insulin biosynthesis upon

glucose stimulation [63]. In the context of T2D, islet insulin content has often [64], but not always [44,65],

been reported to decrease. Given the sex differences in ribosome- and protein synthesis-related

pathways, we compared islet insulin content in male and female islets.

The Humanislets.com and Vanderbilt datasets both used the acid-ethanol method to obtain

insulin content data, including from detergent resistant granules, and normalized to IEQ;

therefore, these two datasets were analyzed both together and independently. The UPenn

dataset involved a detergent-based method to obtain insulin content, possibly reflecting more

recently synthesized insulin and less granule-stored insulin, and normalized per islet; this was

analyzed alone. Among donors without diabetes aged 15-39, no significant sex difference in

insulin content was observed either in the combined Humanislets.com and Vanderbilt data

(Figure 5A), or when these datasets were analyzed independently (Figure S5A-B). The UPenn

data showed a trend (p=0.068) toward higher insulin content in female islets (Figure 5B). Among

donors without diabetes of all ages, Humanislets.com and Vanderbilt data found no sex

difference in insulin content (Figure 5C, S5C-D), whereas UPenn insulin content data showed

significantly higher islet insulin content in females (Figure 5D). Thus, insulin solubilized using

the detergent method showed higher insulin content in female islets among donors without

diabetes, but not insulin solubilized by acid-ethanol.

With respect to the effect of T2D on islet insulin content, data from Vanderbilt and

Humanislets.com data showed that T2D was associated with increased insulin content in

females (Figure 5C, S5C-D), which remained the case when data were age-matched for

Vanderbilt, though not Humanislets.com (Figure S6A-F). This finding was not reproduced in the

UPenn data, in which T2D was associated with decreased insulin content among female donors

(Figure 5D, S6G-H). Male donors who lived with T2D did not have significantly altered islet

insulin content compared to male donors without diabetes across all three datasets (Figure 5C-

D, S5C-D, S6). Therefore, T2D may be associated with increased acid-ethanol-solubilized

insulin but decreased detergent-solubilized insulin specifically in female islets.

Islets from female donors who lived with T2D have preserved mitochondrial activity

We next examined whether sex differences were present in islet oxygen consumption rate (OCR)

data, which reflects mitochondrial function [66–68]. For donors without diabetes aged 15-39, we

combined HPAP and Humanislets.com OCR data (Figure 6A-B) to analyze key parameters

related to mitochondrial function. Neither basal nor maximal respiration was different between

the sexes (Figure 6C-D). Spare respiratory capacity refers to the difference between maximal

respiratory capacity and basal respiration and reflects cells' ability to ramp up respiration in

response to increased energy demand. Female islets had lower spare respiratory capacity than

male islets (Figure 6E). Similar trends were observed when datasets were analyzed individually

(Figure S7).

We expanded our analysis to include donors of all ages, including donors who lived with T2D,

focusing on the HPAP dataset due to the small number of donors who lived with T2D in the

Humanislets.com dataset (Figure 6F-J). In male islets, T2D was associated with a strong

impairment of mitochondrial function. Strikingly, male islets from donors with T2D showed a

drastic increase in basal OCR that was not observed in female islets (Figure 6G), with no

significant sex difference in maximal respiratory capacity (Figure 6H). In male islets, but not

female islets, spare respiratory capacity was also significantly decreased with T2D (Figure 6I),

driven by the increase in basal respiration (Figure 6J). Male islets from donors who lived with

T2D also demonstrated a significantly impaired ability to increase OCR in response to high

glucose; a finding we did not reproduce in female islets (Figure 6J). These T2D-associated

defects in spare respiratory capacity and high glucose-stimulated respiration were also

observed specifically in male islets in age-matched data, although the increase in basal

respiration was no longer significant (Figure S8). Thus, our data indicate that T2D is associated

with fewer perturbations in islet mitochondrial parameters in females compared with males.

Biological sex did not significantly affect intracellular calcium in whole islets

Intracellular calcium triggers insulin exocytosis [69]. There were no significant differences in

intracellular calcium following stimulation with the amino acid mixture, low glucose (3 mM), high

glucose (16.7 mM), or KCl (30 mM) between the sexes, regardless of age or diabetes status

(Figure S9A-E; S10A-E).

T2D is associated with impaired hormone secretion in response to stimuli in both sexes

Dynamic insulin secretion protocols were different with respect to stimulus and exposure time

across all three sites. Among donors without diabetes aged 15-39, no statistically significant sex

differences were observed in insulin secretion in either UPenn or Vanderbilt data (Figure S11A-

L). In the same age range, perifusion results from Humanislets.com showed higher insulin

secretion from male islets compared to female islets (Figure S11M-Q). We next considered

donors of all ages and focused on the UPenn and Humanislets.com perifusions (Figure 7A-B),

as these data were normalized to islet number rather than IEQ and used the same basal

glucose concentration (3 mM), though we note amino acids were present in the UPenn

perifusions. Combining these datasets, we found that T2D was associated with decreased

insulin secretion at 3 mM glucose in males, a trend that was not significant in females (Figure

7C). In both sexes, T2D was associated with decreased insulin secretion in response to high

glucose (15 or 16.7 mM, Figure 7D) and 30 mM KCl (Figure 7E). The stimulation index was

decreased with T2D in both sexes (Figure 7F). These trends were unchanged when age-

matched data were analyzed (Figure S12). When datasets were analyzed individually, there

was high variability within each dataset, reflecting the heterogeneity of nutrient-induced insulin

secretion in human islets [44,70,71], but the T2D-related impairment was apparent in most

parameters (Figure S13-14). Overall, these data indicate that T2D was associated with

impairment of high glucose-induced insulin secretion in both sexes.

Humanislets.com was the only database with data on insulin secretion in response to

leucine and a fatty acid mixture. Consistent with results from the glucose-focused perifusion in

this donor pool (Figure S11N, S13N), among donors without diabetes, male islets secreted more

insulin than female islets at baseline and in response to leucine or fatty acids in both the 15-39

age bracket and across all ages (Figures S15-16).

Perifusion experiments at UPenn and Vanderbilt also measured glucagon secretion and

content. Among donors without diabetes aged 15-39, there were no significant sex differences

in glucagon secretion or glucagon content in either dataset (Figure S17). Among donors of all

ages (Figure 18A-J), UPenn data showed that T2D was associated with a general decrease in

glucagon secretion in female, but not male islets, in terms of basal (Figure S18C), AAM-

stimulated (Figure S18E), and KCl-stimulated secretion (Figure S18I), but this was not apparent

in the Vanderbilt data (Figure S18D, F, J). In both sexes, T2D was associated with decreased

islet glucagon content in the UPenn data (Figure S18K) but not the Vanderbilt data (Figure

S18L). These trends were retained when age-matched data were used (Figure S19). Therefore,

glucagon measurements during the perifusion studies at UPenn suggest that T2D is associated

with greater abnormalities in alpha-cell glucagon secretion in female islets than male islets, but

this was not observed in the Vanderbilt data.

Discussion

The goals of the present study were to (1) identify sex differences in human islet characteristics

in the non-T2D state and (2) assess how these characteristics were altered in female and male

donors who lived with T2D. By performing integrated analysis of several outcomes and using

two independent databases, we discovered robust sex differences across molecular, cellular,

and functional phenotypes in both baseline and T2D contexts. Importantly, T2D was associated

with distinct changes in gene and protein expression in female islets compared to male islets.

This suggests sex-dependent mechanisms of pathogenesis, demonstrating the importance of

considering biological sex in developing prevention and treatment strategies for T2D.

One key finding from our work was that female islets may be less susceptible to T2D-

associated mitochondrial impairments compared to male islets. Mitochondrial respiration is not

only central to GSIS [69,72], but also predicts transplantation success [73–75]. Among young donors

without diabetes, mitochondrial pathways were among the top female-biased pathways,

consistent with prior studies in humans [35] and rats [76]. While lower spare respiratory capacity was

observed in female islets from young donors without diabetes, this was not associated with

defects in insulin secretion. Additionally, although T2D was associated with a female-specific

decrease in mitochondrial gene and protein expression, in line with a previous report [35], this did

not translate to a functional impairment in mitochondrial respiration in islets from female donors

with T2D. A potential explanation for the absence of mitochondrial impairment in female islets is

the higher baseline expression of these genes, which may provide greater capacity for

adaptation, possibly contributing to the lower risk of T2D [1–4] and later age of diagnosis [77] in

women than in men.

Another prominent finding from our analysis was that ribosomal pathways were among

the top gene and protein pathways enriched in healthy female islets. A female bias in protein

synthesis pathways has been observed in several previous studies in humans [29,33,35], and

elevated expression of ribosomal proteins in females has been found across species, including

rodent islets [28,29] and most *Drosophila* cells [78]. The female-biased enrichment of pathways

related to both protein synthesis and ATP generation may support the energy-intensive process

of synthesizing insulin [79]. While our analysis revealed no consistent sex bias in insulin content or

insulin secretion to accompany this higher expression of genes related to protein synthesis and

mitochondrial function, this lack of effect may be due to the lower beta-cell proportion in female

islets. Further work will also be needed to understand why T2D was associated with a positive

enrichment of cytosolic ribosome genes in male islets. While these pathways may be expected

to be beneficial under conditions of high insulin demand, as in T2D, it is possible that male islets

have insufficient mitochondrial activity to support increased levels of protein synthesis.

We further show that alpha-cells may play a greater role in supporting female islets

under physiological conditions and may be more perturbed under T2D conditions. Consistent

with previous analyses in human islets [42,70], we confirmed that female islets had higher

proportions of alpha-cells and lower proportions of beta-cells and non-alpha, non-beta

endocrine cells compared to male islets. Importantly, increased alpha-cell proportion in females

was observed in both the HPAP dataset and Humanislets.com datasets, which used vastly

different methods to estimate this parameter. The HPAP dataset used an antibody-based CyToF

method to obtain counts, which carries the caveat that islets must first be dispersed to single

cells, possibly leading to losses of more fragile cell populations. The Humanislets.com dataset

estimated cell type proportions based on marker protein expression in whole islet proteomics;

these results may be affected by differences in cell identity. A female bias for increased alpha-

cell proportion has also been observed in mice [32], in which alpha-cells support beta-cell function

by paracrine signaling [80–83]. Under high-fat diet stress, the greater alpha-cell proportion in

females may promote beneficial intra-islet structural rearrangements that preserve beta-cell

calcium dynamics [32]. Because alpha-cell proportion further increases in female islets during

pregnancy, this suggests these cells play an important role during times of additional metabolic

stress [84]. An important role for alpha-cells in supporting female metabolic health is further

indicated by our finding that T2D-related changes in alpha-cell gene expression were greater in

females compared to males. Indeed, the HPAP UPenn perifusion data suggested that T2D may

be associated with greater changes in alpha-cell function in females than males. Together, these

results highlight the importance of characterizing potential sex differences in alpha-cells, which

have been understudied in the literature.

Despite multiple sex differences in islet gene and protein expression, and in cellular

processes, perifusion data did not demonstrate consistent sex differences in insulin secretion.

While prior analyses of earlier iterations of the HPAP islet perifusion data observed increased

insulin secretion in female islets compared to male islets [29,33,34], inclusion of new donors shows

the levels of heterogeneity were too high to detect a statistical sex difference. This is consistent

with a recent analysis of insulin secretion data from 576 donors from an independent donor pool

showing high heterogeneity and no measurable impact of donor sex on insulin secretion

patterns[71]. However, humans demonstrate sex differences in insulin secretion *in vivo* [85], and

acute assays in isolated human donor islets may not capture chronic changes in islet physiology

before, during, and after T2D onset. More rigorous investigation of clinical data, with

stratification and analysis by sex, will be required to determine how sex differences in islets at

the molecular and cellular level contribute to T2D pathogenesis and treatment outcomes.

Additional clinical characteristics may also affect islet-level outcomes, including current

medication and potential history of gestational diabetes. Although some medication history was

available in HPAP metadata, it was insufficient to perform stratified analysis. Gestational

diabetes drastically increases the risk of future T2D [86] and is likely relevant to sex-specific

disease characteristics. While there was no record of gestational diabetes among the female

donors included in this study, this may be an avenue for future investigation. More work will also

be needed to capture the effects of genetic ancestry on islet biology. The clinical features of T2D,

including BMI at onset [87–92], clinical presentation [93–96], and genetic risk [97–99], can vary widely

across geography and genetic ancestry. Human islet characteristics, including insulin secretion

[70,100] and gene expression [35] also vary by race. Therefore, while we illustrate sex-specific

differences in islet physiology, we also acknowledge that this is only a first step towards

improving health equity.

In conclusion, by systematic sex-based analysis of multiple data types, we identified key

sex differences in composition, gene expression, and function among donors without diabetes.

These differences likely contribute to sex differences in the function of islets from donors who

lived with T2D, as well as potentially sex-dependent pathways of T2D onset. Elucidating these

sex-dependent disease progression pathways will be critical for developing therapeutic

interventions that are effective in both sexes.

Acknowledgements

Data from this article were downloaded from publicly available databases. HPAP data were

obtained from PancDB, repository of the HPAP Database, consortia under Human Islet

Research Network (RRID:SCR_014393, https://hpap.pmacs.upenn.edu/) (NIH grant numbers

UC4-DK112217 and UC4-DK112232, RRID:SCR_016202). Humanislets.com data and analyses

were obtained from Humanislets.com, funded by the Canadian Institutes of Health Research,

JDRF Canada, and Diabetes Canada (5-SRA-2021-1149-S-B/TG 179092) with data from islets

isolated by the Alberta Diabetes Institute IsletCore with the support of the Human Organ

Procurement and Exchange (HOPE) program, Trillium Gift of Life Network (TGLN). We thank

members of the Rideout and Johnson labs for valuable feedback and advice. We thank

members of the Doliba, MacDonald and Pepper labs for clarification regarding HPAP and

Humanislets.com datasets. We acknowledge that our research takes place on the traditional,

ancestral, and unceded territory of the Musqueam people, and Treaty 6, 7 and 8 territories, a

traditional gathering place for diverse Indigenous peoples including the Cree, Blackfoot, Métis,

Nakota Sioux, Iroquois, Dene, Ojibway/ Saulteaux/Anishinaabe, Inuit, and many others.

Data availability
Original data generated and analyzed during this study are included in this published article or

in the data repositories listed in References.

Supplementary tables and R scripts are available at https://github.com/singyoungchen/sex-

differences-human-islet-characteristics-.

References

1. Wild S, Roglic G, Green A, Sicree R, King H. Global Prevalence of Diabetes: Estimates for
the year 2000 and projections for 2030. *Diabetes Care*. 2004;27(5):1047-1053.
doi:10.2337/diacare.27.5.1047

2. Anjana RM, Deepa M, Pradeepa R, et al. Prevalence of diabetes and prediabetes in 15
states of India: results from the ICMR–INDIAB population-based cross-sectional study.
Lancet Diabetes Endocrinol. 2017;5(8):585-596. doi:10.1016/S2213-8587(17)30174-2

3. Luk AOY, Ke C, Lau ESH, et al. Secular trends in incidence of type 1 and type 2 diabetes
in Hong Kong: A retrospective cohort study. *PLoS Med*. 2020;17(2):e1003052.
doi:10.1371/journal.pmed.1003052

4. Sun H, Saeedi P, Karuranga S, et al. IDF Diabetes Atlas: Global, regional and country-level
diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes Res Clin Pract*.
2022;183:109119. doi:10.1016/j.diabres.2021.109119

5. Xie J, Wang M, Long Z, et al. Global burden of type 2 diabetes in adolescents and young
adults, 1990-2019: systematic analysis of the Global Burden of Disease Study 2019. *BMJ*.
2022;379:e072385. doi:10.1136/bmj-2022-072385

6. Savage PJ, Bennett PH, Gordon Senter R, Miller M. High Prevalence of Diabetes in Young
Pima Indians: Evidence of Phenotypic Variation in a Genetically Isolated Population.
Diabetes. 1979;28(10):937-942. doi:10.2337/diab.28.10.937

7. Matsumoto CL, Tobe S, Schreiber YS, et al. Diabetes prevalence and demographics in 25
First Nations communities in northwest Ontario (2014–2017). *Can J Rural Med*.
2020;25(4):139. doi:10.4103/CJRM.CJRM_99_19

8. Hare MJL, Zhao Y, Guthridge S, et al. Prevalence and incidence of diabetes among
Aboriginal people in remote communities of the Northern Territory, Australia: a
retrospective, longitudinal data-linkage study. *BMJ Open*. 2022;12(5):e059716.
doi:10.1136/bmjopen-2021-059716

9. Louet JF, LeMay C, Mauvais-Jarvis F. Antidiabetic actions of estrogen: Insight from human
and genetic mouse models. *Curr Atheroscler Rep*. 2004;6(3):180-185. doi:10.1007/s11883-
004-0030-9

10. Salpeter SR, Walsh JME, Ormiston TM, Greyber E, Buckley NS, Salpeter EE. Meta-
analysis: effect of hormone-replacement therapy on components of the metabolic
syndrome in postmenopausal women. *Diabetes, Obesity and Metabolism*. 2006;8(5):538-
554. doi:10.1111/j.1463-1326.2005.00545.x

11. Villar E, Chang SH, McDonald SP. Incidences, Treatments, Outcomes, and Sex Effect on
Survival in Patients With End-Stage Renal Disease by Diabetes Status in Australia and
New Zealand (1991–2005). *Diabetes Care*. 2007;30(12):3070-3076. doi:10.2337/dc07-
0895

12.   Carrero JJ, de Mutsert R, Axelsson J, et al. Sex differences in the impact of diabetes on
mortality in chronic dialysis patients. *Nephrol Dial Transplant.* 2011;26(1):270-276.
doi:10.1093/ndt/gfq386

13.   Kautzky-Willer A, Harreiter J, Pacini G. Sex and Gender Differences in Risk,
Pathophysiology and Complications of Type 2 Diabetes Mellitus. *Endocr Rev.*
2016;37(3):278. doi:10.1210/er.2015-1137

14.   The Emerging Risk Factors Collaboration. Diabetes mellitus, fasting blood glucose
concentration, and risk of vascular disease: a collaborative meta-analysis of 102
prospective studies. *Lancet.* 2010;375(9733):2215-2222. doi:10.1016/S0140-
6736(10)60484-9

15.   Harjutsalo V, Maric-Bilkan C, Forsblom C, Groop PH. Impact of Sex and Age at Onset of
Diabetes on Mortality From Ischemic Heart Disease in Patients With Type 1 Diabetes.
Diabetes Care. 2013;37(1):144-148. doi:10.2337/dc13-0377

16.   Peters SAE, Huxley RR, Woodward M. Diabetes as a risk factor for stroke in women
compared with men: a systematic review and meta-analysis of 64 cohorts, including 775
385 individuals and 12 539 strokes. *The Lancet.* 2014;383(9933):1973-1980.
doi:10.1016/S0140-6736(14)60040-4

17.   Sattar N, Rawshani A, Franzén S, et al. Age at Diagnosis of Type 2 Diabetes Mellitus and
Associations With Cardiovascular and Mortality Risks. *Circulation.* 2019;139(19):2228-
2237. doi:10.1161/CIRCULATIONAHA.118.037885

18.   Kaptoge S, Seshasai SRK, Sun L, et al. Life expectancy associated with different ages at
diagnosis of type 2 diabetes in high-income countries: 23 million person-years of
observation. *Lancet Diabetes Endocrinol.* 2023;11(10):731-742. doi:10.1016/S2213-
8587(23)00223-1

19.   Dennis JM, Henley WE, Weedon MN, et al. Sex and BMI alter the benefits and risks of
sulfonylureas and thiazolidinediones in type 2 diabetes: A framework for evaluating
stratification using routine clinical and individual trial data. *Diabetes Care.*
2018;41(9):1844-1853. doi:10.2337/dc18-0344

20.   Dennis JM, Young KG, Cardoso P, et al. A five-drug class model using routinely available
clinical features to optimise prescribing in type 2 diabetes: a prediction model development
and validation study. *The Lancet.* 2025;405(10480):701-714. doi:10.1016/S0140-
6736(24)02617-5

21.   Cardoso P, Young KG, Nair ATN, et al. Phenotype-based targeted treatment of SGLT2
inhibitors and GLP-1 receptor agonists in type 2 diabetes. *Diabetologia.* 2024;67(5):822-
836. doi:10.1007/s00125-024-06099-3

22.   Aroda VR, Brett J, Khutoryansky N, Ratner R. Identifying Predictors of Response to
Liraglutide in Type 2 Diabetes Using Recursive Partitioning Analysis. *Can J Diabetes.*
2012;36(5):S45. doi:10.1016/j.jcjd.2012.07.358

23.   Durden E, Lenhart G, Lopez-Gonzalez L, Hammer M, Langer J. Predictors of glycemic
control and diabetes-related costs among type 2 diabetes patients initiating therapy with

liraglutide in the United States. *J Med Econ*. 2016;19(4):403-413. doi:10.3111/13696998.2015.1131703

24. Hemmer A, Maiter D, Buysschaert M, Preumont V. Long-term effects of GLP-1 receptor agonists in type 2 diabetic patients: A retrospective real-life study in 131 patients. *Diabetes Metab Syndr*. 2019;13(1):332-336. doi:10.1016/j.dsx.2018.09.007

25. Gado M, Tsaousidou E, Bornstein SR, Perakakis N. Sex-based differences in insulin resistance. *Journal of Endocrinology*. 2024;261(1). doi:10.1530/JOE-23-0245

26. Racine KC, Iglesias-Carres L, Herring JA, et al. The high-fat diet and low-dose streptozotocin type-2 diabetes model induces hyperinsulinemia and insulin resistance in male but not female C57BL/6J mice. *Nutr Res*. 2024;131:135-146. doi:10.1016/j.nutres.2024.09.008

27. Biswas P, Bako JA, Liston JB, et al. Insulin/insulin-like growth factor signaling pathway promotes higher fat storage in Drosophila females. *Cell Rep*. 2025;44(7). doi:10.1016/j.celrep.2025.115915

28. Liu G, Li Y, Zhang T, et al. Single-cell RNA Sequencing Reveals Sexually Dimorphic Transcriptome and Type 2 Diabetes Genes in Mouse Islet β Cells. *Genomics, Proteomics & Bioinformatics*. 2021;19(3):408-422. doi:10.1016/j.gpb.2021.07.004

29. Brownrigg GP, Xia YH, Chu CMJ, et al. Sex differences in islet stress responses support female beta cell resilience. *Mol Metab*. 2023;69:101678. doi:10.1016/j.molmet.2023.101678

30. Mastrolia V, Flucher SM, Obermair GJ, et al. Loss of α2δ-1 Calcium Channel Subunit Function Increases the Susceptibility for Diabetes. *Diabetes*. 2017;66(4):897-907. doi:10.2337/db16-0336

31. Jacobo-Piqueras N, Theiner T, Geisler SM, Tuluc P. Molecular mechanism responsible for sex differences in electrical activity of mouse pancreatic β cells. *JCI Insight*. 2024;9(6):e171609. doi:10.1172/jci.insight.171609

32. Visa M, Berggren PO. Sex-dependent intra-islet structural rearrangements affecting alpha-to-beta cell interactions lead to adaptive enhancements of Ca2+ dynamics in prediabetic beta cells. *Diabetologia*. 2024;67(8):1663-1682. doi:10.1007/s00125-024-06173-w

33. Yong HJ, Toledo MP, Nowakowski RS, Wang YJ. Sex Differences in the Molecular Programs of Pancreatic Cells Contribute to the Differential Risks of Type 2 Diabetes. *Endocrinology*. 2022;163(11):bqac156. doi:10.1210/endocr/bqac156

34. Hall E, Volkov P, Dayeh T, et al. Sex differences in the genome-wide DNA methylation pattern and impact on gene expression, microRNA levels and insulin secretion in human pancreatic islets. *Genome Biol*. 2014;15(12):522. doi:10.1186/s13059-014-0522-z

35. Qadir MMF, Elgamal RM, Song K, et al. Sex-specific regulatory architecture of pancreatic islets from subjects with and without type 2 diabetes. *EMBO J*. Published online November 20, 2024:1-19. doi:10.1038/s44318-024-00313-z

36. Van Anh N, Gil HW, Park S, Ryu S. Exploring Sex Differences in Type 2 Diabetes via a
Male-Dominant Beta-Cell Cluster from Single-Cell Pancreatic Sequencing of Public
Datasets. *Endocrinol Metab (Seoul).* Published online May 19, 2025.
doi:10.3803/EnM.2025.2297

37. Kaestner KH, Powers AC, Naji A, HPAP Consortium, Atkinson MA. NIH Initiative to
Improve Understanding of the Pancreas, Islet, and Autoimmunity in Type 1 Diabetes: The
Human Pancreas Analysis Program (HPAP). *Diabetes.* 2019;68(7):1394-1402.
doi:10.2337/db19-0058

38. Shapira SN, Naji A, Atkinson MA, Powers AC, Kaestner KH. Understanding islet
dysfunction in type 2 diabetes through multidimensional pancreatic phenotyping: The
Human Pancreas Analysis Program - Type 2 Diabetes (HPAP-T2D). *Cell Metab.*
2022;34(12):1906-1913. doi:10.1016/j.cmet.2022.09.013

39. Ewald JD, Lu Y, Ellis CE, et al. HumanIslets.com: Improving accessibility, integration, and
usability of human research islet data. *Cell Metab.* 2025;37(1):7-11.
doi:10.1016/j.cmet.2024.09.001

40. Takahashi TA, Johnson KM. Menopause. *Med Clin North Am.* 2015;99(3):521-534.
doi:10.1016/j.mcna.2015.01.006

41. Wang YJ, Golson ML, Schug J, et al. Single-cell mass cytometry analysis of the human
endocrine pancreas. *Cell Metab.* 2016;24(4):616-626. doi:10.1016/j.cmet.2016.09.007

42. Toledo MP, Xie G, Wang YJ. Comprehensive Characterization of Islet Remodeling in
Development and in Diabetes Using Mass Cytometry. *Endocrinology.*
2024;165(9):bqae094. doi:10.1210/endocr/bqae094

43. Hao Y, Stuart T, Kowalski MH, et al. Dictionary learning for integrative, multimodal and
scalable single-cell analysis. *Nat Biotechnol.* 2024;42(2):293-304. doi:10.1038/s41587-
023-01767-y

44. Kolic J, Sun WG, Cen HH, et al. Proteomic predictors of individualized nutrient-specific
insulin secretion in health and disease. *Cell Metab.* 2024;36(7):1619-1633.e5.
doi:10.1016/j.cmet.2024.06.001

45. Ritchie ME, Phipson B, Wu D, et al. limma powers differential expression analyses for
RNA-sequencing and microarray studies. *Nucleic Acids Res.* 2015;43(7):e47.
doi:10.1093/nar/gkv007

46. Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R Package for Comparing Biological
Themes Among Gene Clusters. *OMICS.* 2012;16(5):284-287. doi:10.1089/omi.2011.0118

47. Traag VA, Waltman L, van Eck NJ. From Louvain to Leiden: guaranteeing well-connected
communities. *Sci Rep.* 2019;9(1):5233. doi:10.1038/s41598-019-41695-z

48. Doliba NM, Liu Q, Li C, et al. Inhibition of cholinergic potentiation of insulin secretion from
pancreatic islets by chronic elevation of glucose and fatty acids: Protection by casein
kinase 2 inhibitor. *Mol Metab.* 2017;6(10):1240-1253. doi:10.1016/j.molmet.2017.07.017

49.   Doliba NM, Rozo AV, Roman J, et al. α Cell dysfunction in islets from nondiabetic, glutamic
685   acid decarboxylase autoantibody-positive individuals. *J Clin Invest*. 2022;132(11):e156243.
686   doi:10.1172/JCI156243

687   50.   Banting FG, Best CH, Collip JB, Campbell WR, Fletcher AA. Pancreatic Extracts in the
688   Treatment of Diabetes Mellitus. *Can Med Assoc J*. 1922;12(3):141-146.

689   51.   Scott DA, Fisher AM. Crystalline insulin. *Biochem J*. 1935;29(5):1048-1054.
690   doi:10.1042/bj0291048

691   52.   Romans RG, Scott DA, Fisher AM. Preparation of Crystalline Insulin. *Ind Eng Chem*.
692   1940;32(7):908-910. doi:10.1021/ie50367a010

693   53.   Randall SS. The small-scale preparation of crystalline insulin. *BBA General Subjects*.
694   1964;90(3):472-476. doi:10.1016/0304-4165(64)90226-0

695   54.   Shapcott D, O'Brien D. A Method for the Isolation of Insulin from Single Human Pancreas.
696   *Diabetes*. 1970;19(11):831-836. doi:10.2337/diab.19.11.831

697   55.   Hallas-Møller K, Petersen K, Schlichtkrull J. Crystalline and Amorphous Insulin-Zinc
698   Compounds with Prolonged Action. *Science*. 1952;116(3015):394-398.

699   56.   Scott EL. On the influence of intravenous injections of an extract of the pancreas on
700   experimental pancreatic diabetes. *Am J Physiol*. 1912;29(3):306-310.
701   doi:10.1152/ajplegacy.1912.29.3.306

702   57.   Collip JB. III. The Preparation of Pancreatic Extracts containing Insulin. 2. The Preparation
703   of the Extracts as used in the first Clinical Cases. In: *Proceedings and Transactions of the
704   Royal Society of Canada*. Vol 16. 3. Royal Society of Canada; 1922:28-29.

705   58.   Kimball CP, Allen RS, Piper HA. Precipitation reactions of insulin. *Proc Soc Exp Biol Med*.
706   1923;20(7):414-416. doi:10.3181/00379727-20-202

707   59.   Zhou Z, Ribas V, Rajbhandari P, et al. Estrogen receptor α protects pancreatic β-cells from
708   apoptosis by preserving mitochondrial function and suppressing endoplasmic reticulum
709   stress. *J Biol Chem*. 2018;293(13):4735-4751. doi:10.1074/jbc.M117.805069

710   60.   Gatward LD, Doherty L, Khot S, Gonzalez DC, Ruent S, King A. Sex hormone contribution
711   to sex differences in a mouse model of beta cell endoplasmic reticulum (ER) stress. In:
712   *Endocrine Abstracts*. Vol 94. Bioscientifica; 2023. doi:10.1530/endoabs.94.OC6.3

713   61.   Xu B, Allard C, Alvarez-Mercado AI, et al. Estrogens Promote Misfolded Proinsulin
714   Degradation to Protect Insulin Production and Delay Diabetes. *Cell Rep*. 2018;24(1):181-
715   196. doi:10.1016/j.celrep.2018.06.019

716   62.   De Paoli M, Shah D, Zakharia A, et al. Investigating the Role of 17-Beta Estradiol in the
717   Regulation of the Unfolded Protein Response (UPR) in Pancreatic Beta Cells. *Int J Mol Sci*.
718   2024;25(3):3. doi:10.3390/ijms25031816

63. Schuit FC, In't Veld PA, Pipeleers DG. Glucose stimulates proinsulin biosynthesis by a
dose-dependent recruitment of pancreatic beta cells. *Proc Natl Acad Sci U S A*.
1988;85(11):3865-3869.

64. Marchetti P, Del Guerra S, Marselli L, et al. Pancreatic Islets from Type 2 Diabetic Patients
Have Functional Defects and Increased Apoptosis That Are Ameliorated by Metformin. *J*
Clin Endocrinol Metab. 2004;89(11):5535-5541. doi:10.1210/jc.2004-0150

65. Deng S, Vatamaniuk M, Huang X, et al. Structural and Functional Abnormalities in the
Islets Isolated From Type 2 Diabetic Subjects. *Diabetes*. 2004;53(3):624-632.
doi:10.2337/diabetes.53.3.624

66. Chance B, Williams GR. A Simple and Rapid Assay of Oxidative Phosphorylation. *Nature*.
1955;175(4469):1120-1121. doi:10.1038/1751120a0

67. Brand MD, Nicholls DG. Assessing mitochondrial dysfunction in cells. *Biochem J*.
2011;435(Pt 2):297-312. doi:10.1042/BJ20110162

68. Schmidt CA, Fisher-Wellman KH, Neufer PD. From OCR and ECAR to energy:
Perspectives on the design and interpretation of bioenergetics studies. *J Biol Chem*.
2021;297(4):101140. doi:10.1016/j.jbc.2021.101140

69. Henquin JC, Ravier MA, Nenquin M, Jonas JC, Gilon P. Hierarchy of the β-cell signals
controlling insulin secretion. *Eur J Clin Invest*. 2003;33(9):742-750. doi:10.1046/j.1365-
2362.2003.01207.x

70. Evans-Molina C, Pettway YD, Saunders DC, et al. Heterogeneous endocrine cell
composition defines human islet functional phenotypes. *bioRxiv*. Preprint posted online
August 25, 2025:2024.11.20.623809. doi:10.1101/2024.11.20.623809

71. Gnanasekar P, Saponaro C, Mühlemann M, et al. Standardized dynamic glucose testing
reveals donor-dependent heterogeneity in human islet function. *Cell Transplant*.
2025;34:09636897251350654. doi:10.1177/09636897251350654

72. Affourtit C, Alberts B, Barlow J, Carré JE, Wynne AG. Control of pancreatic β-cell
bioenergetics. *Biochem Soc Trans*. 2018;46(3):555-564. doi:10.1042/BST20170505

73. Pepper AR, Hasilo CP, Melling CWJ, et al. The Islet Size to Oxygen Consumption Ratio
Reliably Predicts Reversal of Diabetes Posttransplant. *Cell Transplant*. 2012;21(12):2797-
2804. doi:10.3727/096368912X653273

74. Papas KK, Bellin MD, Sutherland DER, et al. Islet Oxygen Consumption Rate (OCR) Dose
Predicts Insulin Independence in Clinical Islet Autotransplantation. *PLoS One*.
2015;10(8):e0134428. doi:10.1371/journal.pone.0134428

75. Sweet IR, Gilbert M, Scott S, et al. Glucose-Stimulated Increment in Oxygen Consumption
Rate as a Standardized Test of Human Islet Quality. *Am J Transplant*. 2008;8(1):183-192.
doi:10.1111/j.1600-6143.2007.02041.x

76.   Li T, Jiao W, Li W, Li H. Sex effect on insulin secretion and mitochondrial function in
756   pancreatic beta cells of elderly Wistar rats. *Endocr Res.* 2016;41(3):167-179.
757   doi:10.3109/07435800.2015.1124437

758   77.   Wright AK, Welsh P, Gill JMR, et al. Age-, sex- and ethnicity-related differences in body
759   weight, blood pressure, HbA1c and lipid levels at the diagnosis of type 2 diabetes relative
760   to people without diabetes. *Diabetologia.* 2020;63(8):1542-1553. doi:10.1007/s00125-020-
761   05169-6

762   78.   Pal S, Avellaneda J, Cherian CM, et al. Cell type specific allometry controls sex-differences
763   in Drosophila body size. *bioRxiv.* Published online August 25, 2025:2025.08.25.671808.
764   doi:10.1101/2025.08.25.671808

765   79.   Schuit FC, Kiekens R, Pipeleers DG. Measuring the balance between insulin synthesis and
766   insulin release. *Biochem Biophys Res Commun.* 1991;178(3):1182-1187.
767   doi:10.1016/0006-291X(91)91017-7

768   80.   Holter MM, Saikia M, Cummings BP. Alpha-cell paracrine signaling in the regulation of
769   beta-cell insulin secretion. *Front Endocrinol.* 2022;13. doi:10.3389/fendo.2022.934775

770   81.   Fouque A, Oshima M, Mode N, et al. Constructing chimeric mouse islets to study alpha-
771   and delta-cell influence on beta-cell feature. *Mol Metab.* Published online September 1,
772   2025:102245. doi:10.1016/j.molmet.2025.102245

773   82.   Balasenthilkumaran NV, Bombek LK, Ramirez D, et al. Role of GLP1-receptor-mediated α-
774   β-cell communication in functional β-cell heterogeneity. *bioRxiv.* Preprint posted online
775   April 27, 2025:2025.04.25.650736. doi:10.1101/2025.04.25.650736

776   83.   Ho KH, Barmaver SN, Gibson SE, et al. Paracrine Hormonal Signals From Islet α-Cells
777   Regulate Microtubule Dynamics in β-Cells to Promote Insulin Secretion in Mouse and
778   Human Islets. *Diabetes.* Published online January 9, 2026:db241025. doi:10.2337/db24-
779   1025

780   84.   Seedat F, Holden K, Davis S, et al. A new paradigm of islet adaptations in human
781   pregnancy: insights from immunohistochemistry and proteomics. *Nat Commun.*
782   2025;16(1):6687. doi:10.1038/s41467-025-61852-5

783   85.   Couchepin C, Lê KA, Bortolotti M, et al. Markedly Blunted Metabolic Effects of Fructose in
784   Healthy Young Female Subjects Compared With Male Subjects. *Diabetes Care.*
785   2008;31(6):1254-1256. doi:10.2337/dc07-2001

786   86.   Vounzoulaki E, Khunti K, Abner SC, Tan BK, Davies MJ, Gillies CL. Progression to type 2
787   diabetes in women with a known history of gestational diabetes: systematic review and
788   meta-analysis. *BMJ.* 2020;369:m1361. doi:10.1136/bmj.m1361

789   87.   Resnick HE, Valsania P, Halter JB, Lin X. Differential effects of BMI on diabetes risk among
790   black and white Americans. *Diabetes Care.* 1998;21(11):1828-1835.
791   doi:10.2337/diacare.21.11.1828

88. Huxley R, James WPT, Barzi F, et al. Ethnic comparisons of the cross-sectional
relationships between measures of body size with diabetes and hypertension. *Obes Rev*.
2008;9(s1):53-61. doi:10.1111/j.1467-789X.2007.00439.x

89. Chiu M, Austin PC, Manuel DG, Shah BR, Tu JV. Deriving Ethnic-Specific BMI Cutoff
Points for Assessing Diabetes Risk. *Diabetes Care*. 2011;34(8):1741-1748.
doi:10.2337/dc10-2300

90. Ntuk UE, Gill JMR, Mackay DF, Sattar N, Pell JP. Ethnic-Specific Obesity Cutoffs for
Diabetes Risk: Cross-sectional Study of 490,288 UK Biobank Participants. *Diabetes Care*.
2014;37(9):2500-2507. doi:10.2337/dc13-2966

91. Tillin T, Sattar N, Godsland IF, Hughes AD, Chaturvedi N, Forouhi NG. Ethnicity-specific
obesity cut-points in the development of Type 2 diabetes – a prospective study including
three ethnic groups in the United Kingdom. *Diabet Med*. 2015;32(2):226-234.
doi:10.1111/dme.12576

92. Caleyachetty R, Barber TM, Mohammed NI, et al. Ethnicity-specific BMI cutoffs for obesity
based on type 2 diabetes risk in England: a population-based cohort study. *Lancet
Diabetes Endocrinol*. 2021;9(7):419-426. doi:10.1016/S2213-8587(21)00088-7

93. Mifsud F, Saint-Martin C, Dubois-Laforgue D, Bouvet D, Timsit J, Bellanné-Chantelot C.
Monogenic diabetes in adults: A multi-ancestry study reveals strong disparities in diagnosis
rates and clinical presentation. *Diabet Res Clin Pract*. 2022;188.
doi:10.1016/j.diabres.2022.109908

94. Banerji MA, Chaiken RL, Huey H, et al. GAD Antibody Negative NIDDM in Adult Black
Subjects with Diabetic Ketoacidosis and Increased Frequency of Human Leukocyte
Antigen DR3 and DR4: Flatbush Diabetes. *Diabetes*. 1994;43(6):741-745.
doi:10.2337/diab.43.6.741

95. Mauvais-Jarvis F, Sobngwi E, Porcher R, et al. Ketosis-Prone Type 2 Diabetes in Patients
of Sub-Saharan African Origin: Clinical Pathophysiology and Natural History of β-Cell
Dysfunction and Insulin Resistance. *Diabetes*. 2004;53(3):645-653.
doi:10.2337/diabetes.53.3.645

96. Tuomi T, Santoro N, Caprio S, Cai M, Weng J, Groop L. The many faces of diabetes: a
disease with increasing heterogeneity. *The Lancet*. 2014;383(9922):1084-1094.
doi:10.1016/S0140-6736(13)62219-9

97. Mahajan A, Spracklen CN, Zhang W, et al. Multi-ancestry genetic study of type 2 diabetes
highlights the power of diverse populations for discovery and translation. *Nat Genet*.
2022;54(5):560-572. doi:10.1038/s41588-022-01058-3

98. Srinivasan S, Liju S, Sathish N, et al. Common and Distinct Genetic Architecture of Age at
Diagnosis of Diabetes in South Indian and European Populations. *Diabetes Care*.
2023;46(8):1515-1523. doi:10.2337/dc23-0243

99. Smith K, Deutsch AJ, McGrail C, et al. Multi-ancestry polygenic mechanisms of type 2
diabetes. *Nat Med*. 2024;30(4):1065-1074. doi:10.1038/s41591-024-02865-3

831 100. Doliba NM, Roman J, Rozo AV, et al. 253-LB: Ethnic Differences in Pancreatic Hormone
832    Secretion in Health and T2D. *Diabetes*. 2022;71(Supplement_1):253-LB.
833    doi:10.2337/db22-253-LB

Figures

Figure 1. Islet endocrine cell proportions.

Proportions were determined by CyToF in the HPAP dataset or deconvolution from whole islet

proteomics in the Humanislets.com dataset. For islets from donors without diabetes aged 15-39,

alpha-cell proportions from HPAP (A) and Humanislets.com (B), beta-cell proportions from

HPAP (C) and Humanislets.com (D), and non-alpha, non-beta endocrine cell proportions from

HPAP (E) and Humanislets.com (F). For islets from donors without diabetes and donors with

T2D of all ages, alpha-cell (G), beta-cell (H), and non-alpha, non-beta endocrine cell (I)

proportions from HPAP and alpha-cell (J), beta-cell (K), and non-alpha, non-beta endocrine cell

(L) from Humanislets.com. * indicates $p<0.05$, ns = not significant, $p≥0.05$.

Figure 2. Gene and protein expression analysis of islets from female versus male donors without diabetes aged 15-39.

Significantly altered pathways according to sex in GSEA analysis among combined transcriptomics datasets (HPAP pseudobulk

scRNAseq and Humanislets.com bulk islet RNAseq) (A). Top 60 GO pathways that were significantly altered by sex in GSEA analysis

in Humanislets.com bulk islet proteomics (B). n = 7 female, 17 male for HPAP scRNAseq, 8 female, 18 male for Humanislets.com

bulk RNAseq, and 12 female, 21 male for Humanislets.com proteomics. NES = normalized enrichment score. Pink dots indicate

pathways associated with protein synthesis and processing, red dots indicate pathways associated with mitochondrial regulation and

function, blue dots indicate pathways associated with chromatin and chromatin remodeling, green dots indicate pathways associated

with DNA repair, yellow dots indicate pathways associated with synapses and synapse formation, and purple dots indicate pathways

associated with cell growth and proliferation.

Figure 3. GO pathways upregulated with T2D in islets from female and male donors of all ages.

Pathways that were significantly increased with T2D in female islets in GSEA analysis at $p<0.0001$ among combined transcriptomics

datasets (HPAP pseudobulk scRNAseq and Humanislets.com bulk islet RNAseq) and Humanislets.com proteomics (A). Top 60

pathways that were significantly increased with T2D in male islets in GSEA analysis at $p<0.0001$ among combined transcriptomics

datasets and Humanislets.com proteomics (B). Venn diagrams show common and distinct pathways that were significantly increased

with T2D at $p<0.05$ in the combined transcriptomics versus Humanislets.com proteomics datasets. For HPAP scRNAseq, n = 16

female control, 11 female T2D, 24 male control, and 7 male T2D donors. For Humanislets.com bulk RNAseq, n = 36 female control, 5

female T2D, 66 male control, and 10 male T2D donors. For Humanislets.com proteomics, n = 40 female control, 7 female T2D, 77

male control, and 10 male T2D donors. NES = normalized enrichment score.

Figure 4. GO pathways downregulated with T2D in islets from female and male donors of all ages.

Pathways that were significantly decreased with T2D in female islets in GSEA analysis at $p<0.0001$ among combined transcriptomics

datasets (HPAP pseudobulk scRNAseq and Humanislets.com bulk islet RNAseq) and Humanislets.com proteomics (A). Pathways

that were significantly decreased with T2D in male islets in GSEA analysis at $p<0.0001$ among combined transcriptomics datasets

and Humanislets.com proteomics (B). Venn diagrams show common and distinct pathways that were significantly decreased with

T2D at $p<0.05$ in the combined transcriptomics versus Humanislets.com proteomics datasets. For HPAP scRNAseq, n = 16 female

control, 11 female T2D, 24 male control, and 7 male T2D donors. For Humanislets.com bulk RNAseq, n = 36 female control, 5 female

T2D, 66 male control, and 10 male T2D donors. For Humanislets.com proteomics, n = 40 female control, 7 female T2D, 77 male

control, and 10 male T2D donors. NES = normalized enrichment score.

Donors without diabetes aged 15-39

**Donors without diabetes and donors
who lived with T2D of all ages**

Figure 5. Islet insulin content.

Data shown from combined Humanislets.com and HPAP Vanderbilt (A), and HPAP UPenn (B)

for donors without diabetes aged 15-39, and from combined Humanislets.com and HPAP

Vanderbilt (C) and HPAP UPenn (D) for donors without diabetes and donors with T2D of all

ages. * indicates $p<0.05$, ns = not significant, $p≥0.05$.

Donors without diabetes aged 15-39

Donors without diabetes and donors who lived with T2D of all ages

Figure 6. Oxygen consumption rate in whole islets from HPAP and Humanislets.com.

For islets from donors without diabetes aged 15-39, average OCR ± SD tracings from HPAP (A)

and Humanislets.com (B). Baseline respiration (C), maximal respiratory capacity (D), and spare

respiratory capacity (E) for donors without diabetes aged 15-39 from combined HPAP and

Humanislets.com data. For islets from donors of all ages without diabetes or with T2D, average

OCR tracings ± SD for female and male islets from HPAP (F). Summary data for baseline OCR

(G), maximal respiratory capacity (H), spare respiratory capacity (I), and high glucose-

stimulated OCR (J) from donors of all ages without diabetes or with T2D, from HPAP. AAM =

amino acid mix (0.44 mM alanine, 0.19 mM arginine, 0.038 mM aspartate, 0.094 mM citrulline,

0.12 mM glutamate, 0.30 mM glycine, 0.077 mM histidine, 0.094 mM isoleucine, 0.16 mM

leucine, 0.37 mM lysine, 0.05 mM methionine, 0.70 mM ornithine, 0.08 mM phenylalanine, 0.35

mM proline, 0.57 mM serine, 0.27 mM threonine, 0.073 mM tryptophan, and 0.20 mM valine, 2

mM glutamine), G = glucose, FCCP = carbonyl cyanide-p-trifluoromethoxyphenylhydrazone

(mitochondrial uncoupler), NaN3 = sodium azide (cytochrome oxidase inhibitor), AntA =

antimycin A (complex III inhibitor). * indicates $p<0.05$, ns = not significant, $p \geq 0.05$.

904
Figure 7. HPAP UPenn and Humanislets.com dynamic insulin secretion data following glucose stimulation for islets from

donors without diabetes and donors with T2D of all ages.

Perifusion curves for HPAP UPenn (A) and Humanislets.com (B) data. Average insulin secretion at 3 mM glucose for combined

HPAP UPenn (AAM also present) and Humanislets.com data (C). Peak insulin secretion at high glucose for combined HPAP UPenn

(16.7 mM glucose plus AAM) and Humanislets.com data (15 mM glucose) (D). Peak insulin secretion at 30 mM KCl for combined

HPAP UPenn and Humanislets.com data (E). Stimulation index calculated as peak high glucose-stimulated secretion divided by

average secretion at 3 mM glucose ± AAM for combined HPAP UPenn and Humanislets.com data (F). AAM = amino acid mix (0.44

mM alanine, 0.19 mM arginine, 0.038 mM aspartate, 0.094 mM citrulline, 0.12 mM glutamate, 0.30 mM glycine, 0.077 mM histidine,

0.094 mM isoleucine, 0.16 mM leucine, 0.37 mM lysine, 0.05 mM methionine, 0.70 mM ornithine, 0.08 mM phenylalanine, 0.35 mM

proline, 0.57 mM serine, 0.27 mM threonine, 0.073 mM tryptophan, and 0.20 mM valine, 2 mM glutamine). G = glucose. IBMX = 3-

isobutyl-1-methylxanthine at 0.1 mM. Perifusion curves show mean ± SD. * indicates $p<0.05$, ns = not significant, $p≥0.05$.

Tables

Table 1. Total genes or proteins significantly altered according to biological sex in differential
expression analysis among islets from donors without diabetes aged 15-39.

	Number of female-biased genes/proteins	Number of male-biased genes/proteins
Combined transcriptomics (HPAP pseudobulk scRNAseq and Humanislets.com bulk RNAseq)	1	17
HPAP alpha-cell scRNAseq	1	10
HPAP beta-cell scRNAseq	1	12
Humanislets.com proteomics	0	0

923 **Table 2.** Categories of female-biased and male-biased GO pathways identified by GSEA in
924 islets from donors without diabetes aged 15-39. Values indicate percentage of significant
925 pathways in that category (significant pathways in category/total significant pathways).

	Combined transcriptomics	Proteomics
Female-biased pathways		
Mitochondria	47.8% (11/23)	13.2% (12/91)
Ribosome and protein synthesis	21.7% (5/23)	15.4% (14/91)
Metabolic process	8.7% (2/23)	19.8% (18/91)
Oxidative stress response	4.4% (1/23)	4.4% (4/91)
Metabolite transport	0	23.1% (21/91)
Plasma membrane	0	13.2% (12/91)
Sex chromosome	8.7% (2/23)	0
Wound healing	0	2.2% (2/91)
Endoplasmic reticulum and Golgi	0	6.6% (6/91)
Chromatin	8.7% (2/23)	0
Melanosome	0	2.2% (2/91)
Male-biased pathways		
Chromatin	6.3% (1/16)	21.0% (25/119)
Cell growth and proliferation	68.9% (11/16)	6.7% (8/119)
DNA repair	12.5% (2/16)	9.2% (11/119)
Synapse	6.3% (1/16)	18.5% (22/119)
Cell regulation	6.3% (1/16)	0.8% (1/119)
Transcriptional regulation	0	28.6% (34/119)
Protein synthesis, modification, and degradation	0	9.2% (11/119)
Lysosome	0	0.8% (1/119)
ATPase and GTPase	0	3.4% (4/119)

928 **Table 3.** Total genes or proteins significantly altered according to T2D status in differential
929 expression in islets from donors of all ages.

	Number of genes/proteins increased with T2D in females	Number of genes/proteins increased with T2D in males	Number of genes/proteins decreased with T2D in females	Number of genes/proteins decreased with T2D in males
Combined transcriptomics (HPAP pseudobulk scRNAseq and Humanislets.com bulk RNAseq)	0	310	3	121
HPAP alpha-cell scRNAseq	0	0	0	0
HPAP beta-cell scRNAseq	0	6	0	8
Humanislets.com proteomics	371	299	300	413

932   **Table 4.** Total GO pathways significantly altered according to T2D status in GSEA in islets from
933   donors of all ages.

	Number of GO pathways increased with T2D in females	Number of GO pathways increased with T2D in males	Number of GO pathways decreased with T2D in females	Number of GO pathways decreased with T2D in males
Combined transcriptomics (HPAP pseudobulk scRNAseq and Humanislets.com bulk RNAseq)	399	1430	370	170
HPAP alpha-cell scRNAseq	49	1	252	23
HPAP beta-cell scRNAseq	158	101	193	333
Humanislets.com bulk proteomics	544	269	225	332