Biological sex affects human islet gene expression and mitochondrial function in type 2
diabetes

Sing-Young Chen[1,* †], Haoning Howard Cen[1], James D. Johnson[1,†], Elizabeth J. Rideout[1, †]

[1] Life Sciences Institute, Department of Cellular and Physiological Sciences, The University of

British Columbia, Vancouver, BC, Canada

* first author
[†] co-corresponding authors

Email addresses:

Sing-Young Chen: singyoung.chen@ubc.ca, ORCiD: 0000-0001-5258-8254

James D. Johnson: james.d.johnson@ubc.ca, ORCiD: 0000-0002-7523-9433

Elizabeth J. Rideout: elizabeth.rideout@ubc.ca, ORCiD: 0000-0003-0012-2828

Keywords: pancreatic islet, type 2 diabetes, sex differences, beta-cell, alpha-cell, insulin-

Disclosure summary: All authors declare no competing interests.

Abstract

The clinical characteristics of type 2 diabetes (T2D) differ between the sexes. For example, the

risk of T2D is higher in males than in premenopausal females, whereas the risk of T2D-

associated cardiovascular disease is higher in females. The sex-dependent mechanisms of T2D

pathogenesis remain incompletely understood. Publicly available human islet datasets, such as

HPAP and Humanislets.com, offer a valuable tool for uncovering the impact of biological sex on

islet structure, gene expression, and function at a scale that was not previously possible. We

performed an integrated analysis of data from publicly available sources to identify sex

differences in baseline islet characteristics in donors without diabetes and subsequently

examined these features in donors who lived with T2D. Among donors without diabetes, female

islets had a greater proportion of alpha-cells compared with male islets and showed enriched

expression of ribosomal and mitochondrial pathways in both beta- and alpha-cells.

Measurements of mitochondrial function in female islets revealed lower spare respiratory

capacity compared to male islets due to higher basal respiration. Male and female islets had

distinct changes in gene and protein expression in the context of T2D with female islets having

greater preservation of insulin content and fewer defects in islet function. Together, these data

show female islets have fewer islet impairments in T2D. This highlights the need for detailed

mechanistic studies in both sexes to support effective and sex-informed interventions for T2D.

Introduction

Biological sex influences type 2 diabetes (T2D) incidence, complications, and treatment efficacy.

There is a higher diabetes risk in men until approximately 50-60 years of age [1–4]. This male bias

has been observed across all sociodemographic indices [5], though we note a female bias in

some populations [6–8]. Relative protection against diabetes in pre-menopausal women has been

attributed in part to the protective effects of estrogen [9,10]. Despite the lower incidence of T2D in

premenopausal women, women who live with diabetes are at a higher risk of most T2D

complications (e.g. cardiovascular and renal) compared to men [11–18]. Men and women with T2D

also show differential responses to commonly used medications: male sex predicts better

HbA1c-lowering efficacy of sulfonylureas, which stimulate beta-cell insulin secretion [19,20], and

SGLT2 inhibitors, which decrease renal glucose reabsorption [20,21], whereas insulin-sensitizing

thiazolidinediones and GLP-1 receptor agonists (GLP-1RAs) have greater HbA1c-lowering

effects in females [20–24].

Sex differences in clinical observations are likely due to a combination of differences in

insulin sensitivity and pancreatic beta-cell function. Improved insulin sensitivity in females is

well-established and conserved from humans [25] to mice [26] to fruit flies [27]. Recent studies

suggest that sex differences in pancreatic islets may also play a role. In mice, female islets

show greater resilience to endoplasmic reticulum stress than male islets and have higher

expression of genes involved in protein synthesis [28,29]. Sex differences in electrical activity have

also been observed in mouse islets: glucose stimulation in female beta-cells results in lower

potassium currents leading to a more depolarized membrane potential, more frequent burst-type

action potentials, and smaller glucose-induced transient calcium increases [30,31]. HFD feeding in

mice also causes sex-dependent transcriptomic changes [28], and female mice are better able to

adapt to this metabolic stress through alpha-to-beta cell communication to maintain beta-cell

calcium dynamics [32]. In humans, as in mice, female islets show enriched expression of

pathways related to protein synthesis [29,33]. Beta-cell gene expression is closely linked to

epigenetic modifications, which differ between sexes on both sex chromosomes and autosomes,

including at T2D candidate genes [34,35]. *Ex vivo* insulin secretion studies have also suggested

that female islets may secrete more insulin than male islets in response to high glucose [29,33,34].

T2D causes profound changes in gene expression among islets from both sexes, and these

changes are also partly sex-dependent [33,35,36]. However, our understanding of the molecular and

functional changes that occur with T2D in female and male islets remains far from complete.

76   The Human Pancreas Analysis Program (HPAP) consolidates data from donor human

islets across multiple institutes. The data collected through this program are uploaded to the

publicly available PancDB database, and include measurements of human islet calcium imaging,

mass cytometry, islet perifusion, oxygen consumption, and scRNA-seq, among others [37,38].

Humanislets.com is a repository with functional and omics data on a largely distinct pool of

donors, including a wider range of perifusion experiments and a large human islet proteomics

dataset [39]. Importantly, the datasets have enough donors to test for sex differences in many

available outcomes.

84   Our analysis of HPAP and Humanislets.com revealed pronounced sex differences in islet

composition, RNA and protein levels, and mitochondrial function in islets from donors without

diabetes. We further find that gene and protein expression are differentially altered in T2D

depending on biological sex, and that male islets exhibited mitochondrial defects that were

absent from female islets among donors with T2D. Together, these data demonstrate clear

differences in diverse islet characteristics between male and female islets among individuals

without T2D, and show sex-dependent changes to islet biology in individuals who lived with T2D.

Materials and Methods

Data sources

Publicly available data were downloaded from the Human Pancreas Analysis Program (HPAP)

Database, consortia under Human Islet Research Network (RRID:SCR_014393,

https://hpap.pmacs.upenn.edu/) [37,38] (NIH grant numbers UC4-DK112217 and UC4-DK112232,

RRID:SCR_016202), and Humanislets.com [39], an initiative of the Alberta Diabetes Institute

IsletCore funded by the Canadian Institutes of Health Research, Breakthrough T1D Canada,

and Diabetes Canada (5-SRA-2021-1149-S-B/TG 179092). Specific datasets used are

described in detail below. To study sex differences within the healthy islet state, we specifically

focused on donors without diabetes and aged 15-39 years. This age bracket was chosen to

determine sex differences prior to the onset of menopause [40] and was used to compare male

and female control (non-diabetic) donors only. However, when comparing across control and

T2D for males and females, donors of all ages were included. Donors with type 1 diabetes were

not included in our analyses due to insufficient donor numbers to support sex-specific analysis.

Histograms displaying key metadata variables in each dataset are shown in Figure S1.

Cell type composition

Islet endocrine cell type composition data were available from both HPAP and Humanislets.com.

CyToF data [41] estimating cell counts for each cell type were obtained from donor metadata

available on HPAP, from which endocrine cell type proportions were calculated as described in

the CyTOF Staining Workflow and Protocol on HPAP

(https://hpap.pmacs.upenn.edu/explore/workflow/imaging-mass-cytometry?protocol=1, protocol

accessed 23 September 2025, last updated 11 January 2021). Donors with a total endocrine

cell count less than 30-fold lower than the geometric mean of all donors were excluded, as in [42].

On Humanislets.com, endocrine cell type proportions were estimated based on deconvolution of

whole islet proteomics data [39] and were downloaded directly.

Gene expression analysis

Gene expression data from HPAP and Humanislets.com were analyzed to compare the sexes.

Transcriptomic data from HPAP included scRNAseq data. To analyze scRNAseq data, beta-cell

and alpha-cell expression were first pseudobulked to generate one gene expression profile per

donor per cell type using the *Seurat* package [43]. Pseudobulking overcomes the confounding

effects of uneven cell numbers per donor and appropriately treats individual donors, rather than

cells, as independent observations. For Humanislets.com, only bulk islet RNAseq was analyzed

by sex due to low n's available for scRNAseq data. Bulk islet proteomics data [44] were also

downloaded and analyzed from Humanislets.com. Three comparisons were performed: female

vs male among donors without diabetes aged 15-39, female control (without diabetes) vs female

T2D across all ages, and male control vs male T2D across all ages. For all differential

expression analyses, age was included as a covariate. The *ComplexUpset* R package was used

for the UpSet plot.

DESeq2 [45] was used to analyze HPAP pseudobulk beta-cell and pseudobulk alpha-cell with

apeglm shrinkage of effect size [46]. For Humanislets.com RNAseq data, pre-processed data

were used, which already had batch-correction, and comparisons were performed using

ANCOVA with age as covariate and Benjamini-Hochberg correction for multiple testing. For

Humanislets.com proteomics data, *limma* [47] was used to compare female and male control

donors aged 15-39, correcting for age. Gene Set Enrichment Analysis (GSEA) was performed

on differential expression analysis output to identify key pathways using the clusterProfiler

package [48], in which the direction signed -log10 p-values were used as the rank scores of the

genes. To improve pathway visualization, pathways were considered redundant if their core

enrichment genes exceeded a Jaccard index of 0.6. When selecting top pathways, redundant

pathways were first refined by omitting the pathway with fewer gene components.

Insulin content

Islet insulin content data were downloaded from perifusion datasets on HPAP. There were two

sets of islet perifusion data with corresponding insulin content data. The data from the University

of Pennsylvania (UPenn) were presented per islet using a lysis buffer containing EDTA and

detergent [49,50]. The data from Vanderbilt University were presented per islet equivalent (IEQ),

using acid-ethanol extraction to lyse cells and isolate insulin. Acid-ethanol extraction is widely

used in the field to extract insulin from biological samples at maximum yields [51–55], leveraging

the strong acid to avoid the insulin precipitation pH range of ~4.5-6.5 [52,56] and ethanol to

minimize the contaminating influences of exocrine impurities [57,58] while retaining insulin in

solution [59]. Humanislets.com also used acid-ethanol extraction and presented insulin content

data per IEQ. These data were obtained from the provided metadata.

Oxygen consumption

Islet oxygen consumption in response to stimuli was available from both HPAP and

Humanislets.com. On Humanislets.com, oxygen consumption data were obtained using a

Seahorse Bioanalyzer (Agilent) and calculated parameters were directly downloaded (spare

respiratory capacity = maximum uncoupled respiration – minimum non-mitochondrial respiration,

response to high glucose = mean respiration in 16.7 mM glucose – mean respiration in 3 mM

glucose, baseline = last basal respiration) [39]. As three readings per donor were available,

outliers were identified using Grubb's test and the average per donor was calculated. On HPAP,

these data were obtained using a chemical oxygen probe (Oxyphor G3). Raw data were

downloaded and parameters were calculated using the same formulae as used on

Humanislets.com.

Intracellular calcium

Intracellular calcium imaging data using the ratiometric dye Fura-2 were available only from

HPAP. Raw data were downloaded and processed as follows: 1) only regions that responded to

high glucose were included (response counted if 16.7 mM glucose stimulated an increase in

baseline-corrected calcium signal of at least 50%); 2) after filtering for glucose response, signals

from multiple regions for each run were averaged; 3) runs without a specified time of first

stimulus addition or without high glucose were excluded; 4) to account for different time periods

between stimuli, only the first x minutes after stimulus addition were counted for each run,

where x is the minimum period of time between that stimulus and the next, for any of the runs; 5)

responses were averaged for each individual donor and summary statistics were calculated.

Dynamic hormone secretion

HPAP provides dynamic hormone secretion data from islet perifusion experiments performed at

two sites: UPenn and Vanderbilt. For the UPenn data, areas under the curve (AUC) were

calculated for the periods of perifusion during which islets were exposed to each stimulus, for

both insulin and glucagon secretion. For the Vanderbilt data, there was a visible delay between

stimulus addition and response. Therefore, the AUC calculated for each stimulus was

determined based on the time course tracings: 12-39 min for 16.7 mM glucose, 63-87 min for

16.7 mM glucose + IBMX, 93-111 min for 1.7 mM glucose + epinephrine, and 123-138 min for

KCl. Humanislets.com contains islet perifusion data measuring insulin secretion in response to

glucose, leucine, and a mix of the free fatty acids oleate and palmitate [44]. For each database,

AUC were calculated for the periods of perifusion during which islets were exposed to each

stimulus.

Statistics

Unless otherwise specified, to determine statistical differences due to sex and disease,

ANCOVA testing was performed with age as covariate. Multiple comparisons were performed

using Tukey's correction and the *emmeans* R package. For comparing distributions in Figure S1,

the two-sample Kolgomorov-Smirnov test was used. Correlation of two numerical variables was

performed using Pearson's correlation. Unless otherwise mentioned, a *p*-value threshold of 0.05

(or adjusted *p*-valued threshold where relevant) was used. All relevant script for data processing

and graphing are available at https://github.com/singyoungchen/sex-differences-human-islet-

characteristics-.

Results

Female islets contain a greater proportion of alpha-cells and a smaller proportion of beta-cells

We first examined islet cell type composition in male and female donors without T2D aged 15-

39 years. Female islets had a significantly higher alpha-cell proportion compared to male islets

in both the HPAP and Humanislets.com datasets (Figure 1A-B). Beta-cell proportion, in contrast,

was lower in female islets compared with male islets, a difference that was statistically

significant in the Humanislets.com data and which showed a non-significant but similar trend

(*p*=0.11) in HPAP (Figure 1A-B). When we considered donors of all ages, however, the trends

we observed were not consistent between the HPAP CyToF results and the Humanislets.com

cell type proportion estimates. The HPAP data suggested that, among donors without diabetes

of all ages, female islets showed higher alpha-cell (*p*<0.05) and lower beta-cell (*p*=0.072)

proportions, consistent with observations from donors in the 15-39 age bracket (Figure 1C). In

contrast, when including data for donors of all ages from Humanislets.com, we observed no sex

difference in endocrine cell proportions between female and male donors without T2D (Figure

1D).

We next considered endocrine cell proportions in donors who lived with T2D. The HPAP data

showed that T2D was associated with a trend for an increase in alpha-cell proportion (*p*=0.08)

and a significant decrease in beta-cell proportion (*p*<0.05) in males but not females (Figure 1C).

In contrast, the Humanislets.com data showed an increase in alpha-cell proportion and a

decrease in beta-cell proportion in female, but not male, donors who lived with T2D (Figure 1D).

While the reason for this discrepancy is unclear, it may reflect the different techniques used by

each team to obtain cell proportion estimates (antibody-based identification in HPAP, estimation

based on marker protein expression data in Humanislets.com). Thus, while alpha-cell proportion

was elevated among females in young donors without diabetes, we observed no consistent sex

difference in donors who lived with T2D.

Islets from young female donors show higher expression of protein synthesis genes

We next checked for sex differences in baseline gene expression among islets from donors

aged 15-39 without diabetes. Three RNAseq datasets were analyzed: pseudobulk beta-cell

scRNAseq from HPAP, pseudobulk alpha-cell scRNAseq from HPAP, and bulk islet RNAseq

from Humanislets.com. After conducting differential expression analysis to compare female and

male donors (Table 1, S1), we performed GSEA. We found many female-biased pathways but

relatively few male-biased pathways (Figure 2A-B, Table S2). In particular, pathways related to

ribosomal biogenesis and protein synthesis showed a strong female bias in beta-cells and

whole islets (Figure 2A, Table S2). The potential for enhanced protein synthesis capacity in

female islets has previously been shown by us [29], and others [33], and is likely associated with the

effects of estrogen in alleviating ER stress and preserving protein synthesis [60–63]. Mitochondrial

pathways were also female-enriched (Figure 2A, Table S2), with very few male-biased pathways

identified (Figure 2B, Table S2). The HPAP scRNAseq data also allowed us to perform sex-

based analysis of alpha-cell transcriptomes. As with beta-cells, alpha-cells showed a female-

biased enrichment of ribosomal and mitochondrial gene pathways (Figure 2A). "Histone

demethylase activity" was the only significantly enriched pathway in male alpha-cells compared

with female alpha-cells (Figure 2B).

Humanislets.com includes, to our knowledge, the largest open-source whole islet proteomics

dataset currently available. We conducted differential expression analysis to compare protein

levels in female and male donors without diabetes aged 15-19 (Table S3) followed by GSEA.

Wfound female-biased abundance of proteins related to ribosomes and mitochondria, as well as

lipid transport and the plasma membrane (Figure 2C, Table S4). Male-biased pathways included

those related to chromatin modification and transcriptional regulation (Figure 2C, Table S4).

Thus, the female bias in expression of proteins related to protein synthesis and mitochondrial

function was highly consistent between our RNAseq and proteomics data.

T2D is associated with differential shifts in gene and protein expression depending on sex

We next compared islet gene expression between donors with and without diabetes to evaluate

any sex bias in gene expression changes associated with T2D. In the HPAP beta-cell

scRNAseq dataset, 13 genes were significantly altered with T2D in females and 34 in males

(Table 2, S5-6). In the Humanislets.com bulk RNAseq dataset, 0 genes were significantly

altered with T2D in females and 583 in males (Table 2, S5). GSEA identified more pathways

significantly altered with T2D in males than in females in beta-cells and whole islets (Table 3,

S7). Among the pathways increased with T2D in female islets were pathways related to

transcriptional regulation and histone modification (Figure 3A, Table S7; Figure 2B-C). In male

islets, pathways related to cytosolic ribosomes increased with T2D in beta-cells and whole islets

(Figure 3B, Table S8). Pathways decreased with T2D in female islets were dominated by

pathways such as ribosomal and mitochondrial gene pathways (Figure 3C, Table S7). In males,

T2D was associated with a decrease in beta-cell pathways related to vesicle processing and

secretion, the extracellular matrix, and cell differentiation (Figure 3D, Table S8). Our data

therefore suggest that there were distinct T2D-associated changes in gene expression between

the sexes in beta-cells and whole islets.

The ability to distinguish beta-cell and alpha-cell gene expression in the HPAP scRNAseq

dataset provides additional cell type-specific insights. GSEA showed that T2D was associated

with significant enrichment of 345 pathways in females and 75 pathways in males (Table 3, S7-

8). Among top pathways across the 3 transcriptomic datasets in female islets, T2D was

associated with an increase in pathways associated with transcriptional regulation and a

decrease in ribosomal and mitochondrial pathways in alpha-cells, similar to changes observed

in beta-cells (Figure 3A,C). In males, a smaller proportion of top pathways altered in T2D were

identified in the pseudobulk alpha-cell scRNAseq dataset, suggesting T2D was associated with

greater changes in alpha-cell gene expression in female islets than in male islets.

At the protein level, T2D was associated with altered expression of 441 proteins in female

islets and 482 proteins in male islets, among which 230 were common between sexes and

changed in the same direction (Figure 4A, Table S9-10). While no proteins changed in opposite

directions, many were only significantly altered with T2D in only one sex (Figure 4A). GSEA

showed that T2D was associated with enrichment of pathways related to cell adhesion and cell

death in both sexes (Figure 4B-C, Table S11-12). In female islets, T2D was also associated with

enrichment of pathways related to the inflammatory response and cytoskeleton organization

(Figure 4B, Table S11). In male islets, T2D was associated with an enrichment of cytosolic

ribosomal pathways (Figure 4C, Table S12), consistent with RNAseq data (Figure 3C, Table S8).

Pathways that were decreased with T2D were similar between males and females, and included

those related to vesicle trafficking and exocytosis, as well as mitochondrial components (Figure

4B-C, Table S11-12). Taken together, our analysis reveals sex differences in both islet gene and

protein expression in donors without diabetes and individuals who lived with T2D.

Female islets may show greater preservation of insulin content than male islets in T2D

Healthy beta-cells have a remarkable capacity to rapidly increase insulin biosynthesis upon

glucose stimulation [64]. In the context of T2D, islet insulin content has often [65], but not always [44,66],

been reported to decrease. Given the sex differences in ribosome- and protein synthesis-related

pathways, we compared islet insulin content in male and female islets.

Among donors without diabetes aged 15-39, no significant sex difference was observed in

the Vanderbilt and Humanislets.com datasets, which used the acid-ethanol method to obtain

insulin content data, including from detergent resistant granules (Figure 5A-B). In the UPenn

data, however, a detergent-based method was used to obtain insulin content, possibly reflecting

more recently synthesized insulin and less granule-stored insulin. The UPenn data showed a

trend (p=0.068) toward higher insulin content in female islets (Figure 5C). Among donors

without diabetes of all ages, Vanderbilt and Humanislets.com data found no sex difference in

insulin content (Figure 5D-E), whereas UPenn insulin content data showed significantly higher

islet insulin content in females (Figure 5F). Thus, insulin solubilized using the detergent method

showed higher insulin content in female islets among donors without diabetes, but not insulin

solubilized by acid-ethanol.

With respect to the effect of T2D on islet insulin content, data from Vanderbilt and

Humanislets.com data showed that T2D was associated with increased insulin content in

females (Figure 5D-E). This finding was not reproduced in the UPenn data, in which T2D was

associated with decreased insulin content among female donors (Figure 5F). Male donors who

lived with T2D did not have significantly altered islet insulin content compared to male donors

without diabetes across all three datasets (Figure 5D-F). Therefore, T2D is associated with

increased acid-ethanol-solubilized insulin but decreased detergent-solubilized insulin specifically

in female islets.

Islets from female donors who lived with T2D have preserved mitochondrial activity

We next examined whether sex differences were present in islet oxygen consumption rate (OCR)

data, which reflects mitochondrial function [67–69]. Consistent with higher mitochondrial gene

expression in female islets, both HPAP and Humanislets.com showed that female islets from

donors aged 15-39 without diabetes had a non-significant trend toward higher basal respiration

than male islets (Figure 6A-B, E-F). Maximal respiratory capacity was not different between the

sexes (Figure 6C, G). Spare respiratory capacity refers to the difference between maximal

respiratory capacity and basal respiration and reflects cells' ability to ramp up respiration in

response to increased energy demand. Measurements of spare respiratory capacity showed

that female islets had non-significant trends for lower spare respiratory capacity than male islets

in both HPAP (p=0.055, Figure 6D) and Humanislets.com (p=0.08, Figure 6H).

We expanded our analysis to include donors of all ages, including donors who lived with T2D

(Figure 6I-M). In male islets, T2D was associated with a strong impairment of mitochondrial

function. Strikingly, male islets from donors with T2D showed a drastic increase in basal OCR

that was not observed in female islets (Figure 6J), with no significant sex difference in maximal

respiratory capacity (Figure 6K). In male islets, but not female islets, spare respiratory capacity

was also significantly decreased with T2D (Figure 6L), driven by the increase in basal

respiration (Figure 6J). Male islets from donors who lived with T2D also demonstrated a

significantly impaired ability to increase OCR in response to high glucose; a finding we did not

reproduce in female islets (Figure 6M). Thus, our data indicate that T2D is associated with fewer

perturbations in islet mitochondrial parameters in females compared with males.

Biological sex did not significantly affect intracellular calcium in whole islets

Intracellular calcium triggers insulin exocytosis [70]. There were no significant differences in

intracellular calcium following stimulation with the amino acid mixture, low glucose (3 mM), high

glucose (16.7 mM), or KCl (30 mM) in donors without diabetes, regardless of age or diabetes

status (Figure S2A-E; S3A-E).

T2D is associated with impaired hormone secretion in response to stimuli in both sexes

Dynamic insulin secretion protocols were different with respect to stimulus and exposure time

between all three sites. Among donors without diabetes aged 15-39, no statistically significant

sex differences were observed in insulin secretion in either UPenn or Vanderbilt data (Figure

S4A-L). In the same age range, perifusion results from Humanislets.com showed higher insulin

secretion from male islets compared to female islets (Figure S4M-Q). Across donors of all ages

(Figure 7A-C), T2D was associated with decreased insulin secretion at 0 mM and 3 mM glucose

in females but not males in the UPenn perifusion dataset (Figure 7D-E). Amino acid-stimulated

insulin secretion was impaired with T2D in both sexes in the UPenn data (Figure 7F). In contrast,

Vanderbilt data showed no effect of sex or T2D on insulin secretion at 5.6 mM glucose (Figure

7G), while the Humanislets.com data showed higher insulin secretion at baseline (3 mM glucose)

and in 6 mM glucose in islets from male donors without diabetes compared to female donors

without diabetes (Figure 7H-I). In all three datasets, T2D was associated with impaired high

glucose-stimulated insulin secretion in both sexes (Figure 7J-L). T2D was also associated with

impaired IBMX-stimulated secretion in the UPenn and Vanderbilt data (Figure 7M-N). KCl-

stimulated insulin secretion was impaired with T2D in both sexes in the UPenn dataset (Figure

7O), males in the Vanderbilt dataset (Figure 7P), and neither sex in the Humanislets.com

dataset (Figure 7Q). Hence, T2D was associated with impairment of high glucose-induced

insulin secretion in both sexes, and we detected no sex differences that were consistent

between the datasets. We note high variability within each dataset, however, reflecting the

heterogeneity of nutrient-induced insulin secretion in human islets [44,71,72].

Humanislets.com was the only database with data on insulin secretion in response to

leucine and a fatty acid mixture. Consistent with results from the glucose-focused perifusion in

this donor pool (Figure S5M-Q), among donors without diabetes, male islets secreted more

insulin than female islets at baseline and in response to leucine or fatty acids in both the 15-39

age bracket and across all ages (Figures S5-7). T2D was associated with significantly impaired

insulin secretion in response to 5 mM leucine + 6 mM glucose in males; the same trend did not

reach statistical significance in females (Figure S6D). T2D was associated with impaired insulin

secretion in response to a mixture of oleate and palmitate combined with 6 mM glucose in both

sexes (Figure S6I).

Perifusion experiments at UPenn and Vanderbilt also measured glucagon secretion and

content. Among donors without diabetes aged 15-39, there were no significant sex differences

in glucagon secretion or glucagon content in either dataset (Figure S7). Among donors of all

ages (Figure S8A-B), UPenn data showed that T2D was associated with decreased baseline

glucagon secretion in female but not male islets (Figure S8C), but there were no sex differences

in the Vanderbilt data (Figure S8D). The UPenn data also showed that AAM-stimulated

glucagon secretion was decreased in donors who lived with T2D in both sexes, showing a

significant reduction in females and a non-significant trend (p=0.09) in males (Figure S8E). In

the Vanderbilt dataset, glucagon secretion in response to 1.7 mM glucose and epinephrine was

not affected by sex or T2D status (Figure S8F). Neither dataset showed any significant effect of

sex or T2D status on IBMX-stimulated glucagon secretion (Figure S8G-H). T2D was associated

with decreased KCl-stimulated glucagon secretion in female but not male islets in the UPenn

dataset (Figure S8I), and neither sex in the Vanderbilt dataset (Figure S8J). T2D was associated

with decreased islet glucagon content in the UPenn data (Figure S8K) but not the Vanderbilt

data (Figure S8L). Therefore, glucagon measurements during the perifusion studies at UPenn

suggest that T2D is associated with greater abnormalities in alpha-cell glucagon secretion in

female islets than male islets, but this was not observed in the Vanderbilt data.

Discussion

The goals of the present study were to (1) identify sex differences in human islet characteristics

in the non-T2D state and (2) assess how these characteristics were altered in female and male

donors who lived with T2D. By performing integrated analysis of several outcomes and using

two independent databases, we discovered robust sex differences across molecular and

functional phenotypes in both baseline and T2D contexts. Importantly, T2D was associated with

distinct changes in gene and protein expression in female islets compared to male islets. This

suggests sex-dependent mechanisms of pathogenesis, demonstrating the importance of

considering biological sex in developing prevention and treatment strategies for T2D.

One key finding from our work was that female islets may have greater mitochondrial

activity at baseline and are less susceptible to T2D-associated mitochondrial impairments

compared to male islets. Mitochondrial respiration is not only central to GSIS [70,73], but also

predicts transplantation success [74–76]. Among young donors without diabetes, mitochondrial

pathways were among the top female-biased pathways, consistent with prior studies in humans

[35] and rats [77]. Enrichment of mitochondrial pathways may support the higher basal respiration in

female islets, which has also been observed in non-human primates [78]. While this higher basal

respiration contributed to lower spare respiratory capacity in female islets, this was not

associated with defects in insulin secretion. Additionally, although we observed a female-specific

decrease in mitochondrial gene and protein expression in islets from donors who lived with T2D,

in line with a previous report [35], this did not translate to a functional impairment in mitochondrial

respiration in islets from female donors with T2D. A potential explanation for the ability of female

islets to maintain function despite changes to mitochondrial gene expression is the higher

baseline expression of these genes, providing greater capacity for adaptation, and possibly

contributing to the lower risk of T2D [1–4] and later age of diagnosis [79] in women than in men.

Another prominent finding from our analysis was that ribosomal pathways were among

the top gene and protein pathways enriched in healthy female islets. A female bias in protein

synthesis pathways has been observed in several previous studies in humans [29,33,35], and

elevated expression of ribosomal proteins in females has been found across species, including

rodent islets [28,29] and most *Drosophila* cells [80]. The female-biased enrichment of pathways

related to both protein synthesis and ATP generation may support the energy-intensive process

of synthesizing insulin [81]. While our analysis revealed no consistent sex bias in insulin content or

insulin secretion to accompany this higher expression of genes related to protein synthesis and

mitochondrial function, this lack of effect may be due to the lower beta-cell proportion in female

islets. Further work will also be needed to understand why T2D was associated with an

enrichment of cytosolic ribosome genes in male islets. While these pathways were identified as

female-biased among young donors without diabetes, and would therefore be expected to

promote islet resilience, it is possible that male islets have insufficient mitochondrial activity to

support such high levels of protein synthesis.

We further show that alpha-cells may play a greater role in supporting female islets

under physiological conditions and may be more perturbed under T2D conditions. Consistent

with previous analyses in human islets [42,71], we confirmed that female islets had higher

proportions of alpha-cells and lower proportions of beta-cells compared to male islets. The

same trend has also been observed in mice [32], in which alpha-cells support beta-cell function by

paracrine signaling [82–84]. Under high-fat diet stress, the greater alpha-cell proportion in females

may promote beneficial intra-islet structural rearrangements that preserve beta-cell calcium

dynamics [32]. Because alpha-cell proportion further increases in female islets during pregnancy,

this suggests these cells play an important role during times of additional metabolic stress [85]. An

important role for alpha-cells in supporting female metabolic health is further indicated by our

finding that T2D-related changes in alpha-cell gene expression were greater females compared

to males. Indeed, the HPAP UPenn perifusion data suggested that T2D may be associated with

greater changes in alpha-cell function in females than males. Together, these results highlight

the importance of characterizing potential sex differences in alpha-cells, which have been

understudied in the literature.

Despite multiple sex differences in islet gene and protein expression, and in cellular

processes, perifusion data did not demonstrate consistent sex differences in insulin secretion.

While prior analyses of earlier iterations of the HPAP islet perifusion data observed increased

insulin secretion in female islets compared to male islets [29,33,34], inclusion of new donors shows

the levels of heterogeneity were too high to detect a statistical sex difference. This is consistent

with a recent analysis of insulin secretion data from 576 donors from an independent donor pool

showing high heterogeneity and no measurable impact of donor sex on insulin secretion

patterns[72]. However, humans demonstrate sex differences in insulin secretion *in vivo* [86], acute

assays in isolated human donor islets may not capture chronic changes in islet physiology

before, during, and after T2D onset. More rigorous investigation of clinical data, with

stratification and analysis by sex, will be required to determine how sex differences in islets at

the molecular and cellular level contribute to T2D pathogenesis and treatment outcomes. More

work will also be needed to capture the effects of genetic ancestry on islet biology. The clinical

features of T2D, including BMI at onset [87–92], clinical presentation [93–96], and genetic risk [97–99], can

vary widely across geography and genetic ancestry. Human islet characteristics, including

insulin secretion [71,100] and gene expression [35] also vary by race. Therefore, while we illustrate

sex-specific differences in islet physiology, we also acknowledge that this is only a first step

towards improving health equity.

In conclusion, by systematic sex-based analysis of multiple data types, we identified key

sex differences in composition, gene expression, and function among donors without diabetes.

These differences likely contribute to sex differences in the function of islets from donors who

lived with T2D, as well as potentially sex-dependent pathways of T2D onset. Elucidating these

sex-dependent disease progression pathways is critical for developing interventions that are

effective in both sexes.

Acknowledgements

Data from this article were downloaded from publicly available databases. HPAP data were

obtained from PancDB, repository of the HPAP Database, consortia under Human Islet

Research Network (RRID:SCR_014393, https://hpap.pmacs.upenn.edu/) (NIH grant numbers

UC4-DK112217 and UC4-DK112232, RRID:SCR_016202). Humanislets.com data and analyses

were obtained from Humanislets.com, funded by the Canadian Institutes of Health Research,

JDRF Canada, and Diabetes Canada (5-SRA-2021-1149-S-B/TG 179092) with data from islets

isolated by the Alberta Diabetes Institute IsletCore with the support of the Human Organ

Procurement and Exchange (HOPE) program, Trillium Gift of Life Network (TGLN), and other

Canadian organ procurement organizations with written informed donor consent as approved by

the Human Research Ethics Board at the University of Alberta (Pro00013094). We thank

members of the Rideout and Johnson labs for valuable feedback. We acknowledge and thank

the organ donors and their families. We acknowledge that our research takes place on the

traditional, ancestral, and unceded territory of the Musqueam people, a privilege for which we

are grateful.

Funding

S-Y.C. was supported by the J. Andrew McKee Postdoctoral Fellowship (JDRF and Stem Cell

Network), Canadian Institute of Health Research (CIHR) Postdoctoral Fellowship, and Michael

Smith Health Research BC Trainee Award. H.H.C. was supported by a Breakthrough T1D

Postdoctoral Fellowship. S-Y.C., H.H.C., and J.D.J. acknowledge support from grant #3-COE-

2022-1103-M-B from Breakthrough T1D. E.J.R. gratefully acknowledges funding from the

Canadian Institutes of Health Research (GS4-171365), Diabetes Canada (OG-3-22-5646-ER),

and J.D.J. and E.J.R. acknowledge team grant funding from Breakthrough Type 1 Diabetes and

the Canadian Institutes of Health Research (AWD-024144 BRET1DCA 2022).

Disclosures

All authors declare no competing interests.

Data availability

Data download sources and dates, code for analysis, and supplementary tables are available at

https://github.com/singyoungchen/sex-differences-human-islet-characteristics-.

References

1.　Wild S, Roglic G, Green A, Sicree R, King H. Global Prevalence of Diabetes: Estimates for
the year 2000 and projections for 2030. *Diabetes Care*. 2004;27(5):1047-1053.
doi:10.2337/diacare.27.5.1047

2.　Anjana RM, Deepa M, Pradeepa R, et al. Prevalence of diabetes and prediabetes in 15
states of India: results from the ICMR–INDIAB population-based cross-sectional study.
Lancet Diabetes Endocrinol. 2017;5(8):585-596. doi:10.1016/S2213-8587(17)30174-2

3.　Luk AOY, Ke C, Lau ESH, et al. Secular trends in incidence of type 1 and type 2 diabetes
in Hong Kong: A retrospective cohort study. *PLoS Med*. 2020;17(2):e1003052.
doi:10.1371/journal.pmed.1003052

4.　Sun H, Saeedi P, Karuranga S, et al. IDF Diabetes Atlas: Global, regional and country-level
diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes Res Clin Pract*.
2022;183:109119. doi:10.1016/j.diabres.2021.109119

5.　Xie J, Wang M, Long Z, et al. Global burden of type 2 diabetes in adolescents and young
adults, 1990-2019: systematic analysis of the Global Burden of Disease Study 2019. *BMJ*.
2022;379:e072385. doi:10.1136/bmj-2022-072385

6.　Savage PJ, Bennett PH, Gordon Senter R, Miller M. High Prevalence of Diabetes in Young
Pima Indians: Evidence of Phenotypic Variation in a Genetically Isolated Population.
Diabetes. 1979;28(10):937-942. doi:10.2337/diab.28.10.937

7.　Matsumoto CL, Tobe S, Schreiber YS, et al. Diabetes prevalence and demographics in 25
First Nations communities in northwest Ontario (2014–2017). *Can J Rural Med*.
2020;25(4):139. doi:10.4103/CJRM.CJRM_99_19

8.　Hare MJL, Zhao Y, Guthridge S, et al. Prevalence and incidence of diabetes among
Aboriginal people in remote communities of the Northern Territory, Australia: a
retrospective, longitudinal data-linkage study. *BMJ Open*. 2022;12(5):e059716.
doi:10.1136/bmjopen-2021-059716

9.　Louet JF, LeMay C, Mauvais-Jarvis F. Antidiabetic actions of estrogen: Insight from human
and genetic mouse models. *Curr Atheroscler Rep*. 2004;6(3):180-185. doi:10.1007/s11883-
004-0030-9

10.　Salpeter SR, Walsh JME, Ormiston TM, Greyber E, Buckley NS, Salpeter EE. Meta-
analysis: effect of hormone-replacement therapy on components of the metabolic
syndrome in postmenopausal women. *Diabetes, Obesity and Metabolism*. 2006;8(5):538-
554. doi:10.1111/j.1463-1326.2005.00545.x

11.　Villar E, Chang SH, McDonald SP. Incidences, Treatments, Outcomes, and Sex Effect on
Survival in Patients With End-Stage Renal Disease by Diabetes Status in Australia and
New Zealand (1991–2005). *Diabetes Care*. 2007;30(12):3070-3076. doi:10.2337/dc07-
0895

12.   Carrero JJ, de Mutsert R, Axelsson J, et al. Sex differences in the impact of diabetes on
542   mortality in chronic dialysis patients. *Nephrol Dial Transplant*. 2011;26(1):270-276.
543   doi:10.1093/ndt/gfq386

544   13.   Kautzky-Willer A, Harreiter J, Pacini G. Sex and Gender Differences in Risk,
545   Pathophysiology and Complications of Type 2 Diabetes Mellitus. *Endocr Rev*.
546   2016;37(3):278. doi:10.1210/er.2015-1137

547   14.   The Emerging Risk Factors Collaboration. Diabetes mellitus, fasting blood glucose
548   concentration, and risk of vascular disease: a collaborative meta-analysis of 102
549   prospective studies. *Lancet*. 2010;375(9733):2215-2222. doi:10.1016/S0140-
550   6736(10)60484-9

551   15.   Harjutsalo V, Maric-Bilkan C, Forsblom C, Groop PH. Impact of Sex and Age at Onset of
552   Diabetes on Mortality From Ischemic Heart Disease in Patients With Type 1 Diabetes.
553   *Diabetes Care*. 2013;37(1):144-148. doi:10.2337/dc13-0377

554   16.   Peters SAE, Huxley RR, Woodward M. Diabetes as a risk factor for stroke in women
555   compared with men: a systematic review and meta-analysis of 64 cohorts, including 775
556   385 individuals and 12 539 strokes. *The Lancet*. 2014;383(9933):1973-1980.
557   doi:10.1016/S0140-6736(14)60040-4

558   17.   Sattar N, Rawshani A, Franzén S, et al. Age at Diagnosis of Type 2 Diabetes Mellitus and
559   Associations With Cardiovascular and Mortality Risks. *Circulation*. 2019;139(19):2228-
560   2237. doi:10.1161/CIRCULATIONAHA.118.037885

561   18.   Kaptoge S, Seshasai SRK, Sun L, et al. Life expectancy associated with different ages at
562   diagnosis of type 2 diabetes in high-income countries: 23 million person-years of
563   observation. *Lancet Diabetes Endocrinol*. 2023;11(10):731-742. doi:10.1016/S2213-
564   8587(23)00223-1

565   19.   Dennis JM, Henley WE, Weedon MN, et al. Sex and BMI alter the benefits and risks of
566   sulfonylureas and thiazolidinediones in type 2 diabetes: A framework for evaluating
567   stratification using routine clinical and individual trial data. *Diabetes Care*.
568   2018;41(9):1844-1853. doi:10.2337/dc18-0344

569   20.   Dennis JM, Young KG, Cardoso P, et al. A five-drug class model using routinely available
570   clinical features to optimise prescribing in type 2 diabetes: a prediction model development
571   and validation study. *The Lancet*. 2025;405(10480):701-714. doi:10.1016/S0140-
572   6736(24)02617-5

573   21.   Cardoso P, Young KG, Nair ATN, et al. Phenotype-based targeted treatment of SGLT2
574   inhibitors and GLP-1 receptor agonists in type 2 diabetes. *Diabetologia*. 2024;67(5):822-
575   836. doi:10.1007/s00125-024-06099-3

576   22.   Aroda VR, Brett J, Khutoryansky N, Ratner R. Identifying Predictors of Response to
577   Liraglutide in Type 2 Diabetes Using Recursive Partitioning Analysis. *Can J Diabetes*.
578   2012;36(5):S45. doi:10.1016/j.jcjd.2012.07.358

579   23.   Durden E, Lenhart G, Lopez-Gonzalez L, Hammer M, Langer J. Predictors of glycemic
580   control and diabetes-related costs among type 2 diabetes patients initiating therapy with

liraglutide in the United States. *J Med Econ*. 2016;19(4):403-413.
doi:10.3111/13696998.2015.1131703

24.   Hemmer A, Maiter D, Buysschaert M, Preumont V. Long-term effects of GLP-1 receptor
agonists in type 2 diabetic patients: A retrospective real-life study in 131 patients. *Diabetes*
Metab Syndr. 2019;13(1):332-336. doi:10.1016/j.dsx.2018.09.007

25.   Gado M, Tsaousidou E, Bornstein SR, Perakakis N. Sex-based differences in insulin
resistance. *Journal of Endocrinology*. 2024;261(1). doi:10.1530/JOE-23-0245

26.   Racine KC, Iglesias-Carres L, Herring JA, et al. The high-fat diet and low-dose
streptozotocin type-2 diabetes model induces hyperinsulinemia and insulin resistance in
male but not female C57BL/6J mice. *Nutr Res*. 2024;131:135-146.
doi:10.1016/j.nutres.2024.09.008

27.   Biswas P, Bako JA, Liston JB, et al. Insulin/insulin-like growth factor signaling pathway
promotes higher fat storage in Drosophila females. *Cell Rep*. 2025;44(7).
doi:10.1016/j.celrep.2025.115915

28.   Liu G, Li Y, Zhang T, et al. Single-cell RNA Sequencing Reveals Sexually Dimorphic
Transcriptome and Type 2 Diabetes Genes in Mouse Islet β Cells. *Genomics, Proteomics*
& Bioinformatics. 2021;19(3):408-422. doi:10.1016/j.gpb.2021.07.004

29.   Brownrigg GP, Xia YH, Chu CMJ, et al. Sex differences in islet stress responses support
female beta cell resilience. *Mol Metab*. 2023;69:101678.
doi:10.1016/j.molmet.2023.101678

30.   Mastrolia V, Flucher SM, Obermair GJ, et al. Loss of α2δ-1 Calcium Channel Subunit
Function Increases the Susceptibility for Diabetes. *Diabetes*. 2017;66(4):897-907.
doi:10.2337/db16-0336

31.   Jacobo-Piqueras N, Theiner T, Geisler SM, Tuluc P. Molecular mechanism responsible for
sex differences in electrical activity of mouse pancreatic β cells. *JCI Insight*.
2024;9(6):e171609. doi:10.1172/jci.insight.171609

32.   Visa M, Berggren PO. Sex-dependent intra-islet structural rearrangements affecting alpha-
to-beta cell interactions lead to adaptive enhancements of Ca2+ dynamics in prediabetic
beta cells. *Diabetologia*. 2024;67(8):1663-1682. doi:10.1007/s00125-024-06173-w

33.   Yong HJ, Toledo MP, Nowakowski RS, Wang YJ. Sex Differences in the Molecular
Programs of Pancreatic Cells Contribute to the Differential Risks of Type 2 Diabetes.
Endocrinology. 2022;163(11):bqac156. doi:10.1210/endocr/bqac156

34.   Hall E, Volkov P, Dayeh T, et al. Sex differences in the genome-wide DNA methylation
pattern and impact on gene expression, microRNA levels and insulin secretion in human
pancreatic islets. *Genome Biol*. 2014;15(12):522. doi:10.1186/s13059-014-0522-z

35.   Qadir MMF, Elgamal RM, Song K, et al. Sex-specific regulatory architecture of pancreatic
islets from subjects with and without type 2 diabetes. *EMBO J*. Published online November
20, 2024:1-19. doi:10.1038/s44318-024-00313-z

36.   Van Anh N, Gil HW, Park S, Ryu S. Exploring Sex Differences in Type 2 Diabetes via a
Male-Dominant Beta-Cell Cluster from Single-Cell Pancreatic Sequencing of Public
Datasets. *Endocrinol Metab (Seoul)*. Published online May 19, 2025.
doi:10.3803/EnM.2025.2297

37.   Kaestner KH, Powers AC, Naji A, HPAP Consortium, Atkinson MA. NIH Initiative to
Improve Understanding of the Pancreas, Islet, and Autoimmunity in Type 1 Diabetes: The
Human Pancreas Analysis Program (HPAP). *Diabetes*. 2019;68(7):1394-1402.
doi:10.2337/db19-0058

38.   Shapira SN, Naji A, Atkinson MA, Powers AC, Kaestner KH. Understanding islet
dysfunction in type 2 diabetes through multidimensional pancreatic phenotyping: The
Human Pancreas Analysis Program - Type 2 Diabetes (HPAP-T2D). *Cell Metab*.
2022;34(12):1906-1913. doi:10.1016/j.cmet.2022.09.013

39.   Ewald JD, Lu Y, Ellis CE, et al. HumanIslets.com: Improving accessibility, integration, and
usability of human research islet data. *Cell Metab*. 2025;37(1):7-11.
doi:10.1016/j.cmet.2024.09.001

40.   Takahashi TA, Johnson KM. Menopause. *Med Clin North Am*. 2015;99(3):521-534.
doi:10.1016/j.mcna.2015.01.006

41.   Wang YJ, Golson ML, Schug J, et al. Single-cell mass cytometry analysis of the human
endocrine pancreas. *Cell Metab*. 2016;24(4):616-626. doi:10.1016/j.cmet.2016.09.007

42.   Toledo MP, Xie G, Wang YJ. Comprehensive Characterization of Islet Remodeling in
Development and in Diabetes Using Mass Cytometry. *Endocrinology*.
2024;165(9):bqae094. doi:10.1210/endocr/bqae094

43.   Hao Y, Stuart T, Kowalski MH, et al. Dictionary learning for integrative, multimodal and
scalable single-cell analysis. *Nat Biotechnol*. 2024;42(2):293-304. doi:10.1038/s41587-
023-01767-y

44.   Kolic J, Sun WG, Cen HH, et al. Proteomic predictors of individualized nutrient-specific
insulin secretion in health and disease. *Cell Metab*. 2024;36(7):1619-1633.e5.
doi:10.1016/j.cmet.2024.06.001

45.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol*. 2014;15(12):550. doi:10.1186/s13059-014-
0550-8

46.   Zhu A, Ibrahim JG, Love MI. Heavy-tailed prior distributions for sequence count data:
removing the noise and preserving large differences. *Bioinformatics*. 2019;35(12):2084-
2092. doi:10.1093/bioinformatics/bty895

47.   Ritchie ME, Phipson B, Wu D, et al. limma powers differential expression analyses for
RNA-sequencing and microarray studies. *Nucleic Acids Res*. 2015;43(7):e47.
doi:10.1093/nar/gkv007

48.   Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R Package for Comparing Biological
Themes Among Gene Clusters. *OMICS*. 2012;16(5):284-287. doi:10.1089/omi.2011.0118

49.   Doliba NM, Liu Q, Li C, et al. Inhibition of cholinergic potentiation of insulin secretion from
pancreatic islets by chronic elevation of glucose and fatty acids: Protection by casein
kinase 2 inhibitor. *Mol Metab*. 2017;6(10):1240-1253. doi:10.1016/j.molmet.2017.07.017

50.   Doliba NM, Rozo AV, Roman J, et al. α Cell dysfunction in islets from nondiabetic, glutamic
acid decarboxylase autoantibody-positive individuals. *J Clin Invest*. 2022;132(11):e156243.
doi:10.1172/JCI156243

51.   Banting FG, Best CH, Collip JB, Campbell WR, Fletcher AA. Pancreatic Extracts in the
Treatment of Diabetes Mellitus. *Can Med Assoc J*. 1922;12(3):141-146.

52.   Scott DA, Fisher AM. Crystalline insulin. *Biochem J*. 1935;29(5):1048-1054.
doi:10.1042/bj0291048

53.   Romans RG, Scott DA, Fisher AM. Preparation of Crystalline Insulin. *Ind Eng Chem*.
1940;32(7):908-910. doi:10.1021/ie50367a010

54.   Randall SS. The small-scale preparation of crystalline insulin. *BBA General Subjects*.
1964;90(3):472-476. doi:10.1016/0304-4165(64)90226-0

55.   Shapcott D, O'Brien D. A Method for the Isolation of Insulin from Single Human Pancreas.
Diabetes. 1970;19(11):831-836. doi:10.2337/diab.19.11.831

56.   Hallas-Møller K, Petersen K, Schlichtkrull J. Crystalline and Amorphous Insulin-Zinc
Compounds with Prolonged Action. *Science*. 1952;116(3015):394-398.

57.   Scott EL. On the influence of intravenous injections of an extract of the pancreas on
experimental pancreatic diabetes. *Am J Physiol*. 1912;29(3):306-310.
doi:10.1152/ajplegacy.1912.29.3.306

58.   Collip JB. III. The Preparation of Pancreatic Extracts containing Insulin. 2. The Preparation
of the Extracts as used in the first Clinical Cases. In: *Proceedings and Transactions of the
Royal Society of Canada*. Vol 16. 3. Royal Society of Canada; 1922:28-29.

59.   Kimball CP, Allen RS, Piper HA. Precipitation reactions of insulin. *Proc Soc Exp Biol Med*.
1923;20(7):414-416. doi:10.3181/00379727-20-202

60.   Zhou Z, Ribas V, Rajbhandari P, et al. Estrogen receptor α protects pancreatic β-cells from
apoptosis by preserving mitochondrial function and suppressing endoplasmic reticulum
stress. *J Biol Chem*. 2018;293(13):4735-4751. doi:10.1074/jbc.M117.805069

61.   Gatward LD, Doherty L, Khot S, Gonzalez DC, Ruent S, King A. Sex hormone contribution
to sex differences in a mouse model of beta cell endoplasmic reticulum (ER) stress. In:
Endocrine Abstracts. Vol 94. Bioscientifica; 2023. doi:10.1530/endoabs.94.OC6.3

62.   Xu B, Allard C, Alvarez-Mercado AI, et al. Estrogens Promote Misfolded Proinsulin
Degradation to Protect Insulin Production and Delay Diabetes. *Cell Rep*. 2018;24(1):181-
196. doi:10.1016/j.celrep.2018.06.019

63.   De Paoli M, Shah D, Zakharia A, et al. Investigating the Role of 17-Beta Estradiol in the
Regulation of the Unfolded Protein Response (UPR) in Pancreatic Beta Cells. *Int J Mol Sci*.
2024;25(3):1816. doi:10.3390/ijms25031816

64.   Schuit FC, In't Veld PA, Pipeleers DG. Glucose stimulates proinsulin biosynthesis by a
dose-dependent recruitment of pancreatic beta cells. *Proc Natl Acad Sci U S A*.
1988;85(11):3865-3869.

65.   Marchetti P, Del Guerra S, Marselli L, et al. Pancreatic Islets from Type 2 Diabetic Patients
Have Functional Defects and Increased Apoptosis That Are Ameliorated by Metformin. *J
Clin Endocrinol Metab*. 2004;89(11):5535-5541. doi:10.1210/jc.2004-0150

66.   Deng S, Vatamaniuk M, Huang X, et al. Structural and Functional Abnormalities in the
Islets Isolated From Type 2 Diabetic Subjects. *Diabetes*. 2004;53(3):624-632.
doi:10.2337/diabetes.53.3.624

67.   Chance B, Williams GR. A Simple and Rapid Assay of Oxidative Phosphorylation. *Nature*.
1955;175(4469):1120-1121. doi:10.1038/1751120a0

68.   Brand MD, Nicholls DG. Assessing mitochondrial dysfunction in cells. *Biochem J*.
2011;435(Pt 2):297-312. doi:10.1042/BJ20110162

69.   Schmidt CA, Fisher-Wellman KH, Neufer PD. From OCR and ECAR to energy:
Perspectives on the design and interpretation of bioenergetics studies. *J Biol Chem*.
2021;297(4):101140. doi:10.1016/j.jbc.2021.101140

70.   Henquin JC, Ravier MA, Nenquin M, Jonas JC, Gilon P. Hierarchy of the β-cell signals
controlling insulin secretion. *Eur J Clin Invest*. 2003;33(9):742-750. doi:10.1046/j.1365-
2362.2003.01207.x

71.   Evans-Molina C, Pettway YD, Saunders DC, et al. Heterogeneous endocrine cell
composition defines human islet functional phenotypes. *bioRxiv*. Preprint posted online
August 25, 2025:2024.11.20.623809. doi:10.1101/2024.11.20.623809

72.   Gnanasekar P, Saponaro C, Mühlemann M, et al. Standardized dynamic glucose testing
reveals donor-dependent heterogeneity in human islet function. *Cell Transplant*.
2025;34:09636897251350654. doi:10.1177/09636897251350654

73.   Affourtit C, Alberts B, Barlow J, Carré JE, Wynne AG. Control of pancreatic β-cell
bioenergetics. *Biochem Soc Trans*. 2018;46(3):555-564. doi:10.1042/BST20170505

74.   Pepper AR, Hasilo CP, Melling CWJ, et al. The Islet Size to Oxygen Consumption Ratio
Reliably Predicts Reversal of Diabetes Posttransplant. *Cell Transplant*. 2012;21(12):2797-
2804. doi:10.3727/096368912X653273

75.   Papas KK, Bellin MD, Sutherland DER, et al. Islet Oxygen Consumption Rate (OCR) Dose
Predicts Insulin Independence in Clinical Islet Autotransplantation. *PLoS One*.
2015;10(8):e0134428. doi:10.1371/journal.pone.0134428

76.   Sweet IR, Gilbert M, Scott S, et al. Glucose-Stimulated Increment in Oxygen Consumption
730   Rate as a Standardized Test of Human Islet Quality. *Am J Transplant.* 2008;8(1):183-192.
731   doi:10.1111/j.1600-6143.2007.02041.x

732   77.   Li T, Jiao W, Li W, Li H. Sex effect on insulin secretion and mitochondrial function in
733   pancreatic beta cells of elderly Wistar rats. *Endocr Res.* 2016;41(3):167-179.
734   doi:10.3109/07435800.2015.1124437

735   78.   Carroll DT, Elsakr JM, Miller A, et al. Maternal Western-style diet in nonhuman primates
736   leads to offspring islet adaptations including altered gene expression and insulin
737   hypersecretion. *Am J Physiol Endocrinol Metab.* 2023;324(6):E577-E588.
738   doi:10.1152/ajpendo.00087.2023

739   79.   Wright AK, Welsh P, Gill JMR, et al. Age-, sex- and ethnicity-related differences in body
740   weight, blood pressure, HbA1c and lipid levels at the diagnosis of type 2 diabetes relative
741   to people without diabetes. *Diabetologia.* 2020;63(8):1542-1553. doi:10.1007/s00125-020-
742   05169-6

743   80.   Pal S, Avellaneda J, Cherian CM, et al. Cell type specific allometry controls sex-differences
744   in Drosophila body size. *bioRxiv.* Published online August 25, 2025:2025.08.25.671808.
745   doi:10.1101/2025.08.25.671808

746   81.   Schuit FC, Kiekens R, Pipeleers DG. Measuring the balance between insulin synthesis and
747   insulin release. *Biochem Biophys Res Commun.* 1991;178(3):1182-1187.
748   doi:10.1016/0006-291X(91)91017-7

749   82.   Holter MM, Saikia M, Cummings BP. Alpha-cell paracrine signaling in the regulation of
750   beta-cell insulin secretion. *Front Endocrinol.* 2022;13. doi:10.3389/fendo.2022.934775

751   83.   Fouque A, Oshima M, Mode N, et al. Constructing chimeric mouse islets to study alpha-
752   and delta-cell influence on beta-cell feature. *Mol Metab.* Published online September 1,
753   2025:102245. doi:10.1016/j.molmet.2025.102245

754   84.   Balasenthilkumaran NV, Bombek LK, Ramirez D, et al. Role of GLP1-receptor-mediated α-
755   β-cell communication in functional β-cell heterogeneity. *bioRxiv.* Preprint posted online
756   April 27, 2025:2025.04.25.650736. doi:10.1101/2025.04.25.650736

757   85.   Seedat F, Holden K, Davis S, et al. A new paradigm of islet adaptations in human
758   pregnancy: insights from immunohistochemistry and proteomics. *Nat Commun.*
759   2025;16(1):6687. doi:10.1038/s41467-025-61852-5

760   86.   Couchepin C, Lê KA, Bortolotti M, et al. Markedly Blunted Metabolic Effects of Fructose in
761   Healthy Young Female Subjects Compared With Male Subjects. *Diabetes Care.*
762   2008;31(6):1254-1256. doi:10.2337/dc07-2001

763   87.   Resnick HE, Valsania P, Halter JB, Lin X. Differential effects of BMI on diabetes risk among
764   black and white Americans. *Diabetes Care.* 1998;21(11):1828-1835.
765   doi:10.2337/diacare.21.11.1828

88. Huxley R, James WPT, Barzi F, et al. Ethnic comparisons of the cross-sectional
relationships between measures of body size with diabetes and hypertension. *Obes Rev*.
2008;9(s1):53-61. doi:10.1111/j.1467-789X.2007.00439.x

89. Chiu M, Austin PC, Manuel DG, Shah BR, Tu JV. Deriving Ethnic-Specific BMI Cutoff
Points for Assessing Diabetes Risk. *Diabetes Care*. 2011;34(8):1741-1748.
doi:10.2337/dc10-2300

90. Ntuk UE, Gill JMR, Mackay DF, Sattar N, Pell JP. Ethnic-Specific Obesity Cutoffs for
Diabetes Risk: Cross-sectional Study of 490,288 UK Biobank Participants. *Diabetes Care*.
2014;37(9):2500-2507. doi:10.2337/dc13-2966

91. Tillin T, Sattar N, Godsland IF, Hughes AD, Chaturvedi N, Forouhi NG. Ethnicity-specific
obesity cut-points in the development of Type 2 diabetes – a prospective study including
three ethnic groups in the United Kingdom. *Diabet Med*. 2015;32(2):226-234.
doi:10.1111/dme.12576

92. Caleyachetty R, Barber TM, Mohammed NI, et al. Ethnicity-specific BMI cutoffs for obesity
based on type 2 diabetes risk in England: a population-based cohort study. *Lancet
Diabetes Endocrinol*. 2021;9(7):419-426. doi:10.1016/S2213-8587(21)00088-7

93. Mifsud F, Saint-Martin C, Dubois-Laforgue D, Bouvet D, Timsit J, Bellanné-Chantelot C.
Monogenic diabetes in adults: A multi-ancestry study reveals strong disparities in diagnosis
rates and clinical presentation. *Diabet Res Clin Pract*. 2022;188.
doi:10.1016/j.diabres.2022.109908

94. Banerji MA, Chaiken RL, Huey H, et al. GAD Antibody Negative NIDDM in Adult Black
Subjects with Diabetic Ketoacidosis and Increased Frequency of Human Leukocyte
Antigen DR3 and DR4: Flatbush Diabetes. *Diabetes*. 1994;43(6):741-745.
doi:10.2337/diab.43.6.741

95. Mauvais-Jarvis F, Sobngwi E, Porcher R, et al. Ketosis-Prone Type 2 Diabetes in Patients
of Sub-Saharan African Origin: Clinical Pathophysiology and Natural History of β-Cell
Dysfunction and Insulin Resistance. *Diabetes*. 2004;53(3):645-653.
doi:10.2337/diabetes.53.3.645

96. Tuomi T, Santoro N, Caprio S, Cai M, Weng J, Groop L. The many faces of diabetes: a
disease with increasing heterogeneity. *The Lancet*. 2014;383(9922):1084-1094.
doi:10.1016/S0140-6736(13)62219-9

97. Mahajan A, Spracklen CN, Zhang W, et al. Multi-ancestry genetic study of type 2 diabetes
highlights the power of diverse populations for discovery and translation. *Nat Genet*.
2022;54(5):560-572. doi:10.1038/s41588-022-01058-3

98. Srinivasan S, Liju S, Sathish N, et al. Common and Distinct Genetic Architecture of Age at
Diagnosis of Diabetes in South Indian and European Populations. *Diabetes Care*.
2023;46(8):1515-1523. doi:10.2337/dc23-0243

99. Smith K, Deutsch AJ, McGrail C, et al. Multi-ancestry polygenic mechanisms of type 2
diabetes. *Nat Med*. 2024;30(4):1065-1074. doi:10.1038/s41591-024-02865-3

100. Doliba NM, Roman J, Rozo AV, et al. 253-LB: Ethnic Differences in Pancreatic Hormone
Secretion in Health and T2D. *Diabetes*. 2022;71(Supplement_1):253-LB.
doi:10.2337/db22-253-LB

Figures

Figure 1. Alpha-cell and beta-cell proportions among total endocrine cells.

Proportions were determined by CyToF in the HPAP dataset (A) or deconvolution from whole

islet proteomics in the Humanislets.com dataset (B) in islets from donors without diabetes aged

15-39. Proportions determined by CyToF in the HPAP dataset (C) or deconvolution from whole

islet proteomics in the Humanislets.com dataset (D) in islets from donors without diabetes and

donors with T2D of all ages.

821
Figure 2. Gene and protein expression analysis of islets from female versus male donors without diabetes aged 15-39.

Top 30 GO pathways that were significantly altered according to sex in GSEA analysis among all transcriptomics datasets (HPAP

pseudobulk beta-cell scRNAseq, HPAP pseudobulk alpha-cell scRNAseq, and Humanislets.com bulk islet RNAseq), including (A)

female-biased pathways (positively enriched in female) and (B) male-biased pathways (positively enriched in male). Top 60 GO

pathways that were significantly altered by sex in GSEA analysis in Humanislets.com proteomics (C). n = 7 female, 17 male for HPAP

scRNAseq, 8 female, 18 male for Humanislets.com bulk RNAseq, and 12 female, 21 male for Humanislets.com proteomics. NES:

normalized enrichment score.

Figure 3. Transcriptomic analysis of islets from donors without diabetes versus donors who lived with T2D in females and

males of all ages.

Transcriptomics results are shown from HPAP pseudobulk beta-cell scRNAseq, HPAP pseudobulk alpha-cell scRNAseq, and

Humanislets.com bulk islet RNAseq. Top 30 GO pathways across all datasets that were significantly altered according to T2D status

in GSEA analysis, including those increased with T2D in females (A) and males (B) and those decreased with T2D in females (C)

and males (D). For HPAP scRNAseq, n = 16 female control, 11 female T2D, 24 male control, and 7 male T2D donors. For

Humanislets.com bulk RNAseq, n = 36 female control, 5 female T2D, 66 male control, and 10 male T2D donors. NES: normalized

enrichment score.

Figure 4. Differential expression and GSEA analysis results comparing the proteomes of islets from donors without

842 **diabetes to donors who lived with T2D in females and males of all ages.**

843 Proteins with significantly altered expression depending on T2D status in male islets, female islets, or both (A). Top 60 GO pathways

844 that were significantly altered according to T2D status in GSEA analysis in females (A) and males (B). n = 40 female control, 7

845 female T2D, 77 male control, and 10 male T2D donors. NES: normalized enrichment score.

Donors without diabetes aged 15-39

**Donors without diabetes and donors
who lived with T2D of all ages**

Figure 5. Islet insulin content.

Data shown from Humanislets.com (A), HPAP Vanderbilt (B), and HPAP UPenn (C) for donors

without diabetes aged 15-39, and from Humanislets.com (D), HPAP Vanderbilt (E) and HPAP

UPenn (F) for donors without diabetes and donors with T2D of all ages.

Figure 6. Oxygen consumption rate in whole islets from HPAP and Humanislets.com.

For islets from donors without diabetes aged 15-39, average OCR ± SD tracings from HPAP (A)

and Humanislets.com (B). Baseline respiration from HPAP (C) and Humanislets.com (D),

maximal respiratory from HPAP (E) and Humanislets.com (F), and spare respiratory capacity

from HPAP (G) and Humanislets.com (H) for donors without diabetes aged 15-39. For islets

from donors of all ages without diabetes or with T2D, average OCR tracings ± SD for female

and male islets from HPAP (I). Summary data for baseline OCR (J), maximal respiratory

capacity (K), spare respiratory capacity (L), and high glucose-stimulated OCR (M) from donors

of all ages without diabetes or with T2D, from HPAP. AAM = amino acid mix (0.44 mM alanine,

0.19 mM arginine, 0.038 mM aspartate, 0.094 mM citrulline, 0.12 mM glutamate, 0.30 mM

glycine, 0.077 mM histidine, 0.094 mM isoleucine, 0.16 mM leucine, 0.37 mM lysine, 0.05 mM

methionine, 0.70 mM ornithine, 0.08 mM phenylalanine, 0.35 mM proline, 0.57 mM serine, 0.27

mM threonine, 0.073 mM tryptophan, and 0.20 mM valine, 2 mM glutamine), G 3 = 3 mM

glucose, G 16.7 = 16.7 mM glucose, FCCP = carbonyl cyanide-p-

trifluoromethoxyphenylhydrazone (mitochondrial uncoupler), NaN3 = sodium azide (cytochrome

oxidase inhibitor), AntA = antimycin A (complex III inhibitor).

Figure 7. HPAP and Humanislets.com dynamic insulin secretion data following glucose stimulation for islets from donors

without diabetes and donors with T2D of all ages.

Perifusion curves for HPAP UPenn (A), HPAP Vanderbilt (B), and Humanislets.com (C) data. Area under the curve for baseline (0

mM glucose, D), 3 mM glucose-induced (E), and amino acid-induced (F) insulin secretion for HPAP UPenn data. Area under the

curve for baseline (5.6 mM glucose) insulin secretion for HPAP Vanderbilt data (G). Area under the curve for baseline (3 mM glucose,

H) and 6 mM glucose-induced (I) insulin secretion for Humanislets.com data. Area under the curve for high glucose-induced insulin

section for HPAP UPenn (16.7 mM glucose, J), HPAP Vanderbilt (16.7 mM glucose, K), and Humanislets.com (15 mM glucose, L).

Area under the curve for high glucose and IBMX-induced insulin secretion for HPAP UPenn (M) and HPAP Vanderbilt (N). Area under

the curve for potassium-induced insulin secretion for HPAP UPenn (30 mM KCl, O), HPAP Vanderbilt (20 mM KCl, P), and

Humanislets.com (30 mM KCl, Q). AAM = amino acid mix (0.44 mM alanine, 0.19 mM arginine, 0.038 mM aspartate, 0.094 mM

citrulline, 0.12 mM glutamate, 0.30 mM glycine, 0.077 mM histidine, 0.094 mM isoleucine, 0.16 mM leucine, 0.37 mM lysine, 0.05

mM methionine, 0.70 mM ornithine, 0.08 mM phenylalanine, 0.35 mM proline, 0.57 mM serine, 0.27 mM threonine, 0.073 mM

tryptophan, and 0.20 mM valine, 2 mM glutamine). G = glucose. IBMX = 3-isobutyl-1-methylxanthine at 0.1 mM. Perifusion curves

show mean ± SD.

Tables

Table 1. Total genes significantly altered according to biological sex in differential expression
analysis among islets from donors without diabetes aged 15-39.

	Number of female-biased genes	Number of male-biased genes
HPAP alpha-cell scRNAseq	5	17
HPAP beta-cell scRNAseq	1	10
Humanislets.com bulk RNAseq	0	0

891 Table 2. Total genes significantly altered according to T2D status in differential expression
892 analysis for the 3 transcriptomics datasets (HPAP alpha-cell and beta-cell scRNAseq and
893 Humanislets.com bulk RNAseq) in islets from donors of all ages.

	Number of genes increased with T2D in females	Number of genes increased with T2D in males	Number of genes decreased with T2D in females	Number of genes decreased with T2D in males
HPAP alpha-cell scRNAseq	0	0	1	0
HPAP beta-cell scRNAseq	7	8	6	26
Humanislets.com bulk RNAseq	0	375	0	209

895 Table 3. Total GO pathways significantly altered according to T2D status in GSEA for the 3
896 transcriptomics datasets (HPAP alpha-cell and beta-cell scRNAseq and Humanislets.com bulk
897 RNAseq) and the Humanislets.com proteomics dataset in islets from donors of all ages.

	Number of GO pathways increased with T2D in females	Number of GO pathways increased with T2D in males	Number of GO pathways decreased with T2D in females	Number of GO pathways decreased with T2D in males
HPAP alpha-cell scRNAseq	90	23	255	52
HPAP beta-cell scRNAseq	251	176	196	379
Humanislets.com bulk RNAseq	54	557	84	178
Humanislets.com bulk proteomics	544	269	225	332