Title: Phosphorus stress and spatial confinement lower the threshold for quorum-sensing activation of

redox-active metabolite production in *Pseudomonas synxantha*

Authors: R. E. Alcalde[1], H. Jeckel[1], O. Zhang[2], R. L. Avila[1], N. Dalleska[3], D. Mavrodi[4], C. Yang[5], D. K.

Newman[1,6,a]

[1]Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA

91125

[2]Department of Physics & Astronomy, Texas A&M University, College Station, TX 77843

[3]Global Environmental Center, California Institute of Technology, Pasadena, CA 91125

[4]USDA, Agricultural Research Service, Wheat Health, Genetics and Quality Research Unit, Pullman,

Washington, USA

[5]Division of Engineering and Applied Science, California Institute of Technology, Pasadena, CA 91125

[6]Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA 91125

[a]To whom correspondence may be addressed. Email: dkn@caltech.edu

Abstract:

Bacteria often coordinate collective behaviors such as biofilm formation and secondary metabolite

production through quorum sensing (QS), a regulatory system traditionally linked to high cell density.

However, in environments like soil, where microbial populations are spatially fragmented, sparse,

nutrient-limited, and subject to mass transport, the mechanisms that enable QS-dependent processes

remain incompletely understood. Here, we investigate the regulation of a secreted redox-active

metabolite, phenazine-1-carboxylic acid (PCA), in *Pseudomonas synxantha* 2-79, a model

rhizobacterium, under phosphorus (P) limitation, a persistent stress in many soils. Using a combination of

microscopy and molecular genetic approaches, we show that P limitation sensitizes the QS activation

threshold by an order of magnitude, enabling phenazine induction at relatively low population densities in

comparison to P replete conditions. This induction is abolished in QS-deficient mutants and restored by

the addition of exogenous acyl-homoserine lactone (AHL), demonstrating that quorum sensing remains

essential but its threshold becomes environmentally tuned. Under P limitation, spatial confinement and

pore saturation levels further shape the timing and location of induction, illustrating how physical

structure and nutrient stress can modulate bacterial activities. Moreover, phosphorus stress confers both

collaborative and competitive advantages, enabling *P. synxantha* to undergo low-cell density AHL cross-

talk between related *Pseudomonas* spp. and to suppress other rhizobacteria. Lastly, on plant roots, PCA

genes are more predominantly induced under P limitation. These findings illustrate how the nutrient status

of an environment can modulate the onset of quorum sensing, enabling quorum-regulated behaviors to

activate at lower thresholds.

Introduction:

Soil microbes sustain ecosystems. Through their metabolism, they drive nutrient cycling, modulate plant

health, and influence mineral weathering. Many of these functions depend not only on individual

physiology but on the coordinated activity of microbial populations and communities, often mediated

through the release of small molecules that act as public goods, chemical signals, or antibiotics. Among

these, phenazines—redox-active metabolites produced by diverse soil bacteria that support a variety of

biological functions[1–4]—are particularly notable. Through their redox chemistry, they promote

phosphorus[5,6] and iron availability,[7–10] shape microbial competition,[11–13] and affect plant-microbe

interactions[14–16] in the rhizosphere.

Phenazine biosynthesis is typically controlled by quorum sensing (QS), a gene regulatory system in which

bacteria produce and respond to diffusible signals, such as acyl-homoserine lactones (AHLs), to

coordinate group-level behaviors.[17,18] Once a threshold concentration of signal accumulates, often

reflected by high cell density, QS activates gene expression for costly cooperative functions like biofilm

formation, virulence, or secondary metabolite production.[19–25] This framework is well supported in

laboratory systems. However, in natural environments like soil, microbial populations are often spatially

fragmented, sparse, and physically confined within pore structures,[26–28] conditions that can shift signal

accumulation and QS activation. Cluster analyses in structured soils report that larger clusters (>100 cells)

become sparse as water content decreases, implying many neighborhoods fall below 100 cells.[26] Despite

this, phenazines[29,30,4], and other QS-regulated metabolites[31–33] are commonly detected in the rhizosphere.

How, then, are these behaviors triggered in environments where signal accumulation may be restricted?

Several models emphasize that QS dynamics also depend on environmental structure and signal transport.

Diffusion sensing and efficiency sensing, alternative descriptions of quorum sensing, describe how mass

transfer, spatial confinement, and local cell organization determine whether signal concentrations reach

activation thresholds.[34–37] In structured habitats like biofilms and soils, spatial confinement and mass

transfer limitation may allow signals to accumulate locally even when bulk population density is low.[28,38–]

[41] Consistent with this view, a few single cells confined in picoliter microfluidic chambers initiate "high-

density" QS programs.[38] Likewise, flow cell experiments show that advection suppresses QS in exposed

regions of a biofilm while sheltered niches maintain activation.[42]

Beyond physical structure and transport, rhizosphere chemistry and biology, such as nutrient limitation,

physicochemical stress, and host-derived cues can also tune QS induction[43]. Here we focus on

phosphorus limitation in *Pseudomonas synxantha* (formerly *P. fluorescens*) 2-79, a plant growth-

promoting rhizobacterium that produces phenazine-1-carboxylic acid (PCA).[44,45] In this strain, PCA

production is governed by a single AHL-based quorum-sensing system, PhzIR; deletion of either *phzI* or

phzR abolishes PCA production.[45–47] Although PCA increases under phosphorus limitation,[6,48] whether

this induction is mediated by the PhzIR circuit, arises via cross-talk with other QS pathways, or is driven

independently of QS remains unknown. This organism-level question is motivated by the broader soil

context: freely available inorganic phosphate is extremely low because much of it is immobilized through

sorption to mineral surfaces, [49–51] and in *Pseudomonas aeruginosa*, phosphate (and iron) limitation is

known to influence secondary metabolism through multiple QS pathways (e.g., PQS, IQS, RhlR)[52–58] via

the PhoBR two component that governs system central phosphate sensing and response.[59–61]

We set out to determine whether phosphorus stress alters QS dynamics in *P. synxantha* 2-79. Specifically,

we asked whether phosphorus-limited induction of PCA is QS-dependent or QS-independent. We

postulated that phosphorus stress would trigger AHL-based PhzIR circuits, thereby lowering the effective

QS cell density threshold because PCA production under P limitation occurs at lower cell densities in

comparison to standard conditions.[48] We further hypothesized that structured environments act with

phosphorus levels to shape the onset, location, and ecological consequences of QS-regulated behaviors.

To test these predictions, we integrated genetic, optical, and ecological approaches to better understand

the rules governing microbial coordination in nature.

Results

Phosphorus limited environments enhance phenazine gene expression of *Pseudomonas synxantha* in

plant root microcosms

To begin exploring how phenazine biosynthesis responds to phosphorus availability in the rhizosphere,

we imaged *Pseudomonas synxantha* 2-79 colonizing *Brachypodium distachyon* Bd21-3 roots under

phosphorus-replete (P-rep; excess inorganic phosphate in medium), phosphorus-limited (P-lim; no

inorganic phosphate in medium), and phosphate-sequestering (P-HFO; phosphate bound to iron, i.e., not

bioavailable) conditions using light sheet fluorescence microscopy (LSFM). In many Fe-oxide–rich,

acidic to neutral soils, phosphate is strongly retained by iron oxides (e.g., hydrous ferric oxide,

HFO),[50,62,63] so our P–HFO treatment approximated this sequestration. A dual fluorescent reporter was

used to track phenazine biosynthesis gene expression (P_{phzA}-GFP, green channel) and colonization density

readout (constitutive, P_{tac}-mCherry, red channel). Fluorescence intensity was quantified from maximum

intensity projections (MIPs) of LSFM images. For each object within a region of interest, the mean

intensity was calculated as integrated density divided by object area. This metric approximates

fluorescence per unit area in 2D projections, serving as a readout for relative reporter expression and cell

density in colonized regions. Bar plots and Empirical Cumulative Distribution Functions (ECDFs)

revealed condition-specific differences in both fluorescence magnitude and distribution (Fig. 1A–B).

Induction of phenazine biosynthesis (P_{phzA}) was highest under the P-HFO condition (mean = 0.138),

followed by P-lim (mean = 0.061), while remaining minimal under P-replete conditions (mean = 0.016).

Although the P-lim condition exhibited lower average expression than P-HFO, its distribution was

markedly right-skewed and heavy-tailed, indicating rare but strong induction events. In contrast, P-HFO

displayed a broader, moderate distribution, while P-rep remained narrowly distributed around low

fluorescence values (Fig. 1B). Representative LSFM images corroborated these findings, revealing robust

activation of the P_{phzA} reporter under P-HFO but not under P-replete conditions (Fig. 1C). Interestingly,

phenazine expression did not directly correlate with cell density. The constitutive reporter was highest in

P-HFO and P-rep (means = 0.117 and 0.115, respectively) and lower under P-lim (mean = 0.051), yet

only P-lim and P-HFO conditions induced the phenazine reporter.

These observations indicate that at the time of measurement, the cell density was insufficient to trigger

phenazine gene expression under nutrient-replete conditions. How, then, does phosphorus limitation

stimulate phenazine gene induction at local colonization levels that fail to trigger expression under

phosphorus-replete conditions? This observation suggests that canonical QS induction of phenazines may

be altered under nutrient stress, but whether this results from a bypass of QS regulation, an increased

sensitivity to quorum signals, or integration with parallel stress-response pathways was unclear. We set

out to disentangle these possibilities.

Figure 1: Phenazine gene expression of *Pseudomonas synxantha* 2-79 on plant roots under different phosphorus conditions. (A) Bar plots showing mean ± 95% CI of normalized red channel (constitutive; P_{tac}) and green channel (phenazine biosynthesis; P_{phzA}) fluorescence intensities under phosphorus-replete (P-rep), phosphorus-limited (P-lim), and phosphate-sorbing hydrous ferric oxide (P-HFO) conditions. Statistical comparisons were performed using two-sided Mann–Whitney tests; significance is indicated by asterisks (ns, not significant; **** $p < 0.0001$). **(B)** Empirical cumulative distribution functions (ECDFs) of normalized fluorescence intensities from the same dataset shown in (A), illustrating the distribution of object-level fluorescence values across experimental conditions. **(C)** Representative maximum intensity projection (MIP) images acquired by light sheet fluorescence microscopy (LSFM) of *P. synxantha* colonizing plant roots under P-rep, P-lim and P-HFO conditions. Red channel images (left) are shown with identical intensity thresholds to compare red fluorescence. Composite images (right) overlay red and green channels using the same fixed thresholds across conditions, revealing differences in P_{phzA} reporter induction. Scale bars = 500 μm. See also Movie S1 for a 3D reconstruction, Table S1 for summary statistics, and Figure S1 for object-level strip plot.

Quorum sensing is required for phenazine production under phosphorus limitation, and

phosphorus stress sensitizes the QS system to AHL.

To determine whether phenazine production under phosphorus limitation requires the *phzIR* quorum

sensing (QS) pathway, we examined wild type and mutants defective in either the phenazine production

genes (*phzA-G*), the QS signal synthase (*phzI*), or the phosphorus starvation response regulator (*phoB*)

under P-rep (7 mM Pi) or P-lim (100 µM Pi) conditions in 96-well plate liquid cultures (Fig. 2A). The

wild type produced phenazines under both conditions but showed markedly higher induction per cell

under P-lim. In contrast, the Δ*phzI* mutant failed to produce phenazines under P-lim and P-rep,

confirming that QS is required for induction under P-lim. Similarly, the Δ*phoB* mutant did not produce

phenazines under P-lim but retained production under P-rep, indicating that phosphorus stress signaling

via PhoBR is also required in the P-lim response.

We next asked whether exogenous addition of the QS signal produced by PhzI would complement Δ*phzI*

and whether this response is specific to phosphorus depletion. We performed a dose–response assay by

supplementing the Δ*phzI* mutant with the N-(3-hydroxy-hexanoyl)-L-homoserine lactone (3-OH-C6-

HSL) (Fig. 2B). *P. synxantha* produces at least six AHLs, including three of the hydroxy form, with 3-

OH-C6-HSL having the highest sensitivity to the PhzR.[46] To restrict growth and isolate regulatory

effects, we used a defined buffer medium lacking either nitrogen (No-N) or both nitrogen and phosphorus

(No-N/P). After 20 h, PCA production was absent in No-N cultures, even at high QS signal

concentrations, but was strongly induced in No-N/P cultures in a dose-dependent manner.

We further asked whether phosphorus limitation increases sensitivity to QS molecules. We therefore

performed a second 3-OH-C6-HSL dose–response assay (Fig. 2C), this time allowing the Δ*phzI* mutant to

grow under P-rep (7 mM) and P-lim (100 µM) conditions. After 20 h, PCA production was present in

both conditions in a dose-dependent manner, but under P-lim, lower QS signal concentrations were

sufficient to induce phenazine production. Fitting a two-parameter Hill (log-logistic) function revealed

half-maximal effective concentrations (EC50) of 0.51 µM (P-rep) and 0.05 µM (P-lim), indicating ~10-

fold greater sensitivity under phosphorus stress.

Lastly, to verify QS signal production, we directly quantified AHLs (i.e., 3OH-C6-HSL, 3OH-C8-HSL,

and 3OH-C10-HSL) by LC-MS under P-rep, P-lim, and N-lim (nitrogen limited) conditions for both the

wild-type strain and Δ*phzI*. Results are shown for WT (Fig. 2D). For the wild-type strain, QS signals were

produced under both P-lim and P-rep but were not detected under N-lim. No AHLs were detected with the

Δ*phzI* strain. 3OH-C10-HSL was not detected under any conditions. Our limit of detection was 10 nM for

all AHLs.

Together, these results demonstrate that phenazine production under P-lim is QS-dependent and requires

the PhoBR two-component system. Moreover, phosphorus stress sensitizes QS by lowering the AHL

activation threshold, enabling induction at levels that are non-inducing under P-rep or N-lim.

Figure 2: Quorum sensing requirement and reduced AHL concentration threshold for phenazine
production in *Pseudomonas synxantha* 2-79 under phosphorus limitation. (A) Phenazine-associated
absorbance normalized to cell density ($ABS_{367} \cdot OD_{600}^{-1}$) in *P. synxantha* strains under phosphorus-replete
(P-rep) and phosphorus-limited (P-lim) medium at 20 h post-inoculation. Values were baseline-corrected

using the $\Delta phzA$-G mutant mean for each condition. Bars show mean ± SD of replicates; points represent
individual measurements; n=4. **(B)** Dose–response of the $\Delta phzI$ mutant strain to exogenous 3-OH-C6-
HSL under nitrogen-free (No-N) and combined nitrogen- and phosphorus-free (No-N, P) conditions at 20
h post-inoculation. Responses were baseline-corrected by subtracting the mean signal at 0 µM 3-OH-C6-
HSL within each condition. Points show per-dose means; shaded regions indicate mean ± SD of
replicates; n=5. Curves show four-parameter logistic model fits. Insert in legend: mean ± SD of OD_{600} at
the assay timepoint. **(C)** Dose–response of the $\Delta phzI$ mutant strain to exogenous 3-OH-C6-HSL at 20 h
under P-rep and P-lim medium at 20 h post-inoculation. Responses were baseline-corrected and scaled to
an empirical E_{max} (median of the two highest doses) to yield 0–1 normalization. Points show per-dose
means; shaded bands indicate ±SD across replicates; n=3. Curves are two-parameter Hill fits; Insert in
legend: mean ± SD of OD_{600} at the assay timepoint and EC_{50} values from the fits. **(D)** AHL concentration
normalized to cell density ($\mu M \cdot OD_{600}^{-1}$)) for 3OH-C6-HSL and 3OH-C8-HSL across P-rep, P-lim, and
nitrogen-limited (N-lim) conditions. Bar plots show mean ± SD. Values above each bar indicate the
corresponding optical density (OD_{600}) at sampling time (22 h). "ND" denotes values below the detection
limit.

Phosphorus limitation enables quorum sensing activation at low cell densities in planktonic culture

Because QS is required for phenazine production under P-lim, we inferred that phosphorus stress

decreases the cell-density threshold for QS activation. To test this, we used reporter strains to track QS-

dependent induction in wild type *P. synxantha* 2-79 grown in well-mixed liquid culture across phosphate

regimes (25, 100, and 7000 µM Pi) with or without exogenous 3-OH-C6-HSL (AHL) (Fig. 3A-B). Our

previously constructed P_{phoA} and P_{phzA} reporter strains[48] served to monitor phosphorus stress (activation of

PhoBR) and phenazine production, respectively.

The *phoA* reporter was induced in mid-exponential phase at 25 and 100 µM Pi (P-lim), with both

induction timing and cell density increasing with phosphorus availability. It was not detected at 7000 µM

Pi (P-rep). Without added AHL, the *phzA* reporter was induced in late exponential/early stationary phase

under all conditions, but earlier and at a substantially lower cell density in P-lim compared to P-rep

(OD_{600} at induction: 25 µM, 0.0846 ± 0.001; 100 µM, 0.219 ± 0.005; 7000 µM, 0.806 ± 0.006; mean ±

SD), representing ~10-fold and ~4-fold reductions at 25 and 100 µM, respectively. Addition of exogenous

AHL advanced induction time and lowered the density threshold further; under P-lim. Moreover, *phzA*

onset aligned more closely with *phoA*, consistent with phosphorus stress priming the QS response while

endogenous signal accumulation is rate-limiting when AHL is not supplied. It is also noteworthy that

even when AHL is exogenously added in excess, gene expression still requires the cells to be under P

limitation or relatively high cell density under P-rep for induction. That is, it seems that beyond requiring

high AHL concentrations the cells need to be in a state of physiologically readiness (e.g., P stress, other

stress responses at high cell density).

These results indicate that phosphorus stress reduces the cell-density requirement for quorum activation

and speeds QS-dependent gene expression in well-mixed cultures, motivating tests in structured

environments that better simulate rhizosphere matrices, where mass transfer and bacterial confinement

could further tune this sensitization.

Figure 3: Induction benchmarks and growth dynamics under phosphorus regimes of planktonic and sessile *Pseudomonas synxantha* 2-79 cultures. (A) Induction curves for phosphorus-stress (*phoA*) and phenazine biosynthesis (*phzA*) reporters in liquid cultures across phosphate regimes (Pi = 25, 100, 7000 μM). Raw fluorescence intensity values are shown. **(B)** Growth curves in liquid cultures as cell density (OD_{600}). **(C)** Induction curves for *phoA* and *phzA* in agar biofilms under P-lim and P-rep conditions in either 5 mL or 2 mL agar volumes. Raw fluorescence intensity values are shown. **(D)** Growth curves in agar biofilms represented by the constitutive reporter fluorescence intensity (cell

density readout). **(E)** Representative images of reporter induction time course for biofilms grown under P-
lim in a 2 mL agar volume. Intensity of fluorescence is depicted by red to yellow hue. Gray shading
denotes the area coverage of aggregates. Legend: WT P_{phoA} without AHL addition (green), WT P_{phzA}
without AHL addition (orange), P_{phzA} with AHL addition (blue). Curves denote mean; shading denotes ±
SD; n = 3. Error not seen is smaller than the line. Dashed vertical lines mark reporter induction times.
Arrows on third graph correspond to images in (E)

Phosphorus limitation also lowers the cell density threshold for quorum sensing activation in sessile

populations

We employed an early-stage colony–biofilm assay using our previously developed imaging system[48] to

quantify growth and reporter expression in *P. synxantha*. In this diffusion-dominated context, AHL

concentrations in the biofilm arise from the balance between signal production, diffusive loss into the

underlying agar (which serves as a sink), and retention within the biofilm matrix. Reporters on

phosphorus stress (P_{phoA}) and phenazine production (P_{phzA}) tracked induction, and a constitutive reporter

($P_{PA10403}$) provided a readout for cell density (Fig. 3C-E). By varying nutrient status, agar volume, and

AHL supplementation, we modulated regulatory sensitivity, sink capacity (5 mL vs 2 mL agar volumes),

and signal availability, respectively, to reveal how these factors shape QS activation.

5 mL agar: Under P-lim, *phoA* was detected early (~7.5 hr), but *phzA* induction was markedly delayed

(~22.5 hr). Under P-rep, *phoA* did not induce (as expected), whereas *phzA* induced at 19.5 h. This pattern

is opposite to planktonic cultures, where P-lim induces earlier than P-rep, indicating that in this

heterogeneous, diffusion-dominated context, QS-signal accumulation is delayed under P-lim.

Notwithstanding the delay, the biofilm density required for *phzA* activation was ~9-fold lower under P-

lim than P-rep (constitutive reporter [a.u.]: P-lim 27,852 ± 1,207; P-rep 248,856 ± 13,115; mean ± SD),

which is consistent with planktonic trends.

2 mL agar: Under P-lim, *phoA* was, again, detected early (~9.5 hr), *phzA* induction occurred (~18 hr;

earlier than 5 mL condition), and before P-rep (~20 hr), suggesting that QS-signal thresholds were

achieved more rapidly when sink capacity was reduced (i.e., lower agar volume). Interestingly, the

biofilm density at the time of *phzA* induction was nearly identical between the 2 mL and 5 mL conditions,

for both P-lim and P-rep, indicating that biomass accumulation was faster under 2 mL, likely contributing

to the earlier induction. Lastly, as in the 5 mL agar experiments, P-lim, again, required ~10-fold lower

cell density than P-rep for *phzA* activation.

The addition of exogenous AHL caused P-lim colonies to induce *phzA* expression rapidly, coincident with

the onset of phosphorus stress and bypassing the need for signal accumulation through growth. AHL

rescue in P-rep colonies also accelerated the induction time but still required a higher cell density than P-

lim. This is again consistent with liquid cultures. Exogenous AHL accelerated induction but did not

bypass the requirement for phosphorus limitation or a high biofilm density.

Together, these results show that QS activation in colony biofilms reflects an interplay between nutrient

status and the balance of AHL signal accumulation versus diffusive loss in the surrounding agar.

Phosphorus limitation lowers the biofilm density required for QS activation, but induction still depends

on local signal concentration, which is influenced by retention within the biofilm versus loss into the agar.

Phosphorus stress and spatial confinement accelerate quorum sensing in soil-like

microenvironments

To examine how the physical structure of the environment shapes local signal accumulation and QS

activation, we used a previously designed microfluidic platform that mimics soil-like porous media[64] (Fig.

4A). Low-density cultures (OD_{600} = 0.05) of a dual reporter strain (*P. synxantha* 2-79 P_{tac}-P_{phzA}) were

inoculated and air was introduced to generate a landscape of water pockets that varied in size,

connectivity, saturation levels (Fig. 4B). Across roughly 50 regions of interest (ROIs) imaged in each

experiment, we tracked biomass (P_{tac}) and *phz* induction (P_{phzA}) (Fig. 4C, Fig. S2).

Comparing P-lim and P-rep conditions revealed that phosphorus stress strongly advanced QS activation

and lowered the biomass required for *phz* induction (Fig. 4D,E). Under P-lim, ROIs typically induced

within a few hours (mean induction time ~5 hr) and did so at relatively low biomass, whereas under P-rep

conditions induction was markedly delayed (mean ~32 hr) and required substantially higher biomass at

the time of activation (on average 5-fold greater intensity). Induction times were also much more

heterogeneous across ROIs under P-rep conditions. Because this reactor is not well mixed, additional

transport and resource gradients almost certainly emerge that affect the observed timing that we do not

control for (e.g., oxygen depletion in larger pores). In this more realistic geometry, the contrast between

P-lim and P-rep induction time appears amplified relative to previous experiments (Figures 2 and 3).

We next asked whether QS induction under P limitation depends on initial cell density within individual

water pockets. To quantify this feature, we used the ratio of cell area to water area at $t = 1$ hr for each ROI

as a metric for initial cell density and related it to the time of *phz* induction for each ROI. Across P-lim

experiments, induction time decreased systematically with increasing starting density, following a non-

linear, approximately power-law relationship ($R^2 = 0.65$, Fig. 4F, Fig. S3). Pearson and Spearman

correlations gave similar results ($r = -0.771$, $p = 4.14e-20$; $\rho = -0.839$, $p = 1.59e-26$). Therefore, most of

the variation in induction timing is explained by initial cell density, supporting a view in which

phosphorus stress and spatial confinement allow cells concentrated in pore water to accelerate phosphate

depletion and AHL accumulation to facilitate QS activation.

Taken together, these data demonstrate that the combination of phosphorus stress and spatial confinement

enables QS activation in microenvironments that would otherwise restrict signal buildup. We next asked

whether lowered QS concentration thresholds could provide an ecological advantage in a community

setting.

Figure 4: Relationship between cell density, water saturation, and *phz* induction time in microfluidic microenvironments. (A) Schematic (left) and scanning electron micrograph (right) of the synthetic porous platform used to create spatially structured microenvironments. The reactor consists of a 2 by 1 cm pore network containing an array of evenly spaced cylindrical posts 300 μm in diameter, a

depth of 10 µm, and a porosity of 0.39. **(B)** Representative images of pore-scale water distribution at high
saturation and low saturation levels, with pillars shown in green, water-filled pores in gray, and air-filled
pores in red. **(C)** Representative composite images of reporter induction (constitutive: red, *phzA*: green)
over time. **(D)** Boxplots summarizing *phz* reporter induction times for individual ROIs across replicate P-
lim (P-lim 1–2) and P-rep (P-rep 1–2) experiments. Boxes show medians and interquartile ranges;
whiskers extend to 1.5× the interquartile range. See Figure S2 for induction time estimation. **(E)** Boxplots
summarizing red-channel integrated fluorescence per ROI (density proxy) at the time of *phz* reporter
induction across replicate P-lim (P-lim 1–2) and P-rep (P-rep 1–2) experiments. Boxes show medians and
interquartile ranges, and whiskers extend to 1.5× the interquartile range. **(F)** Distribution of induction
times under P-lim. Relationship between cell density in pore water (x-axis; area fraction of cells in water-
filled regions at t = 1 hr) and *phz* induction time (y-axis) for P-lim 1 (circles) and P-lim 2 (squares). Points
are colored by water saturation (fraction of pore volume filled with water), as indicated by the color bar.
The dashed line shows a power-law fit to all data. Data represent pooled technical replicates from regions
in the reactor (P-lim 1, n = 44 ROIs; P-lim 2, n = 52 ROIs).

Phosphorus limitation confers cooperative and antagonistic quorum-sensing-driven community

dynamics

Under nutrient stress, microbial communities can flip between cooperative information exchange (e.g.,

quorum signals) and antagonistic interference (e.g., antibiotics), reshaping which partners win or lose.

To determine whether phosphorus limitation broadly enhances AHL production and promotes cross-talk

among *Pseudomonas* species, we assembled a panel of rhizosphere-associated strains representing both

close relatives of *P. synxantha* and more distantly related pseudomonads. Each strain was grown under N-

lim, P-lim, or P-rep conditions, cell-free supernatants were then collected to challenge the *P. synxantha*

Δ*phzI*–P$_{phzA}$ reporter, which cannot synthesize AHLs but fluoresces upon QS-activated *phzA* expression

(Fig. 5A). Supernatants from multiple donors activated the reporter, with the strongest and broadest

induction under P-lim and P-rep, and little to none under N-lim. As expected, controls with the Δ*phzI*

donor did not induce, whereas the wild-type donor triggered a robust response. Notably, across the panel,

strains carrying homologous *phzIR* operons consistently induced *phzA* under P-lim, suggesting that

phosphorus stress broadly elevates AHL output among *Pseudomonas* species. In contrast, *P. aeruginosa*

PA14 failed to induce the reporter, consistent with its production of 3-oxo-acyl-HSLs rather than the 3-

hydroxy-acyl-HSL(s) recognized by PhzR in *P. synxantha*.

Next, we asked whether QS activation under P-lim alters microbial interactions within competitive

communities under oxic conditions where phenazines primarily function as antibiotics.[11] We performed

co-culture assays between *P. synxantha* (wild type and Δ*phzA-G*, a phenazine-null mutant) and

Pedobacter kyonggii, a common rhizosphere isolate. When P-limited, the wild-type strain significantly

suppressed *P. kyonggii* growth, whereas the Δ*phzA-G* mutant did not (Fig. 5B). This phenazine-linked

advantage was absent under N-lim in the wild-type strain and, interestingly, under P-rep conditions as

well. Although the wild type produced PCA under P-replete conditions, induction occurred much later

(~35 h vs ~16 h in P-lim; see Fig. S5), by which time the co-cultures had already completed most of their

growth, leaving little opportunity for inhibition to manifest. Thus, the phenotype likely reflects the timing

of phenazine onset relative to culture growth.

These findings indicate that phosphorus limitation broadly elevates AHL signaling across *Pseudomonas*

spp., priming QS in *P. synxantha*. Moreover, phenazine production under phosphorus stress suppresses

growth of *P. kyonggii*.

A

B

Figure 5: (A) *Pseudomonas* **spp. supernatant cross-induction QS assay.** Heatmap of fluorescence induction in *P. synxantha* Δ*phzI*-P$_{phzA}$ across supernatants of selected strains and media conditions at 24 h. Rows represent media (N-lim, P-lim, P-rep); columns are strain supernatants. Each cell shows the fold-

change of the chosen metric relative to the reference strain, *P. synxantha* Δ*phzI*, within the same medium.
Where shown, numeric labels are fold-change values rounded to two decimals; the *P. synxantha* Δ*phzI*
column equals 1 by definition. Fluorescence is normalized by OD600. (mean, $n = 6$) **(B) Competition**
assays between *P. synxantha* against *Pedobacter kyonggii* under various nutrient regimes. Bar plots
show OD600 at 72 h (mean ± SD, $n = 6$) for four pairings. The first name indicates the strain measured;
the second is its pairing partner. Results are shown separately for N-lim, P-lim, P-rep media. *P. synxantha*
wild type (2-79 WT) and *P. synxantha* Δ*phzA-G* (2-79 Δ*phzA-G*). Statistical comparisons use Welch's *t*-
test; brackets denote significance (ns, not significant; ** $p < 0.01$).

Discussion

Our study reveals that phosphorus limitation rewires the quorum-sensing (QS) circuitry of *Pseudomonas*

synxantha 2-79, lowering the effective signal concentration required for autoinduction and enabling

phenazine biosynthesis at conditions corresponding to much lower densities (OD ≈ 0.08 in liquid

cultures). Phosphorus stress reduces the acyl-homoserine lactone (AHL) concentration required for

activation by roughly an order of magnitude under our tested conditions, thereby heightening cellular

sensitivity to QS signals. This response depends on the PhoBR two-component system: inactivation of

PhoBR abolishes phenazine induction even when AHL levels are high, demonstrating that phosphorus

stress provides a necessary regulatory input. In structured habitats, where mass transfer alters QS

molecule accumulation, phosphorus limitation emerges as a key modulator of QS activation. In colony

biofilms, phenazine induction times under P limitation are strongly affected by agar volumes and biofilm

density, implicating biofilm retention and diffusive losses in achieving threshold AHL concentrations.

Exogenous AHL addition collapses this concentration requirement but still requires a physiological state

primed for induction (i.e., P limitation or higher cell densities under P replete). Confinement in

micropores creates microenvironments that retain signals and enable localized QS activation, again

showing that physical structure and nutrient state jointly set QS thresholds. These environmentally-tuned

threshold shifts have ecological consequences. Under phosphorus limitation, *P. synxantha* inhibits growth

of the rhizosphere isolate *Pedobacter kyonggii* through phenazine production, a benefit lost in the

phenazine-null mutant and under nitrogen limitation or phosphorus-replete conditions. Early phenazine

synthesis may thus serve dual functions in resource competition and phosphorus acquisition, coordinated

through stress-responsive signaling.

Our data argue against "concentration-only" or "density-only" views and instead support an AND-gate

model for QS activation: (i) sufficient signal accumulation (production versus diffusion, advection,

sorption, and decay) to cross a threshold and (ii) a physiological readiness state that licenses

responsiveness. Density remains a useful proxy because per-cell production scales with biomass and high

density often coincides with a physiological readiness, but other regulatory programs can shift the

effective threshold, explaining low-density induction under nutrient stress. In our system, QS signal

responsiveness is tunable by nutrient status, PhoBR under phosphorus limitation, so signal is necessary

but not sufficient. This framework reconciles our observations: under P-replete + excess AHL, low-

density cultures fail to induce despite ample signal, indicating physiological unreadiness; responsiveness

emerges by mid- to late-exponential growth as global regulators (e.g., Gac/Rsm) and stress programs rise.

Under P limitation, induction synchronizes with phosphorus stress. Exogenous AHL enables induction at

comparatively lower cell densities, with the activation density tunable by the imposed Pi level. Again,

though, induction did not occur until P limitation was achieved.

This model aligns with prior work. Overactivation of the stringent response via RelA in *P. aeruginosa*

prematurely activates RpoS, LasR, RhlR and autoinducer synthesis, driving LasB production independent

of density, i.e., physiological readiness upstream of signal thresholds.[65] Exogenous PQS (an autoinducer)

addition can override cell-density dependence but not growth-phase dependence, underscoring the

physiological readiness requirement.[66] More broadly, nutrient limitation and growth rate jointly modulate

extracellular signal accumulation across species, with phosphorus and sulfur limitation disproportionately

elevating alkylquinolones even at lower OD.[67] Our findings extend this signal-accumulation +

physiological readiness concept by showing that native PhoBR directly sensitizes an unmodified AHL-

based QS network in a rhizosphere pseudomonad, reframing QS as an environmentally tuned

concentration-sensing mechanism whose effective activation threshold is set jointly by autoinducer

concentration and regulatory state. Related frameworks[43] reach similar conclusions; nevertheless,

proliferating new labels for QS risks semantic confusion that can imply a single dominant driver.[36]

Instead, we favor a generalizable AND-gate view that explains QS outputs in sparse, fragmented soils and

highlights practical levers (e.g., nutrient availability, confinement, and mass transfer) for engineering

compatible microbial communities and biosensors in heterogeneous environments.

Methods:

Bacteria. The primary strain was *Pseudomonas synxantha* (formerly *P. fluorescens*) 2-79 WT. Isogenic

derivatives included Δ*phzI* (AHL synthase null), Δ*phzA-G* (phenazine-null), and Δ*phoB* (phosphorus-

stress response regulator null). Reporters carrying transcriptional fusions included P*phzA*–mNeonGreen

(QS/phenazine biosynthesis), P*phoA*–mNeonGreen (phosphorus stress), P*phzA*–GFP, P*tac*–mCherry

(QS/phenazine biosynthesis, density metric) and P*PA10403*-mNeonGreen (density metric). Donor/competitor

strains for crosstalk and competition assays included *Pseudomonas aureofaciens* 30-84, *P.*

brassicacearum 106, *P. libanensis* 84, *P. orientalis*, *P. aeruginosa* Δ*phz1/2* and *Pedobacter kyonggii*.

Sources, identifiers, and genotypes are listed in Table S2.

Culture conditions. Unless noted, overnight cultures of *Pseudomonas* strains were propagated at 30 °C,

250 rpm in LB (Miller) for 16-18 h; *E. coli* DH10B/SM10 were grown at 37 °C. Experimental assays

used defined MOPS medium (below). Antibiotics were used only for selection (gentamicin 30 μg mL^{-1},

tetracycline 15 μg mL^{-1}). All experiments with reporters were antibiotic-free. Strains were stored as −80

°C glycerol stocks (20% v/v) and streaked fresh for each experiment.

Media compositions and nutrient-regime definitions

MOPS defined medium. Per liter: 25 mM MOPS (pH 7.0; adjusted with 1 M NaOH), 55.5 mM glucose,

16 mM NH₄Cl, 7 mM KH₂PO₄, 0.41 mM MgSO₄·7H₂O, 0.68 mM CaCl₂·2H₂O, Aquil-based trace

metals[68] with Fe 10 μM and EDTA 100 μM. KH₂PO₄ 7 mM (P-rep/N-lim) or 25–100 μM (P-lim); NH₄Cl

16 mM (P-rep/P-lim) or 2 mM (N-lim). No-N and No-N/P buffers omitted NH_4Cl and NH_4Cl/KH_2PO_4,

respectively. MOPS agar consisted of MOPS medium with 1.0% (w/v) noble agar.

Plant medium. 0.5× MS (Sigma-Aldrich M5519), 0.6% (w/v) Phytagel (Sigma-Aldrich P8169), pH 5.8

(KOH); P-lim omits KH_2PO_4.

Phosphate-sorbing (P-HFO) condition. Hydrous ferric oxide (HFO) was synthesized, phosphate was

then sorbed and sequestered (P-HFO). P-HFO was added to phosphorus deficient plant medium (see

below) at 1.6 g/L. See Note S1 for P-HFO preparation.

MES defined medium. Per L: 50 mM MES pH 5.8 (KOH), 50 mM sucrose, salts as above. KH_2PO_4 7

mM (P-rep/N-lim) or 100 µM (P-lim); NH_4Cl 16 mM (P-rep/P-lim) or 2 mM (N-lim).

Reporter and mutant generation and verification

Transcriptional fusions were constructed using the mini-Tn7 system.[69] Promoter (P_{phzA}) was cloned

upstream of mNeonGreen in pJM220 by Gibson assembly and transformed into *E. coli* DH10B.

Gentamicin-resistant transformants (30 µg mL^{-1}) were verified by colony PCR and Sanger sequencing.

Mini-Tn7 elements and pTNS1 helper (TnsABCD; delivered from *E. coli* SM10/pTNS1) were introduced

into electrocompetent *P. synxantha* 2-79. Transconjugants were selected on LB-gentamicin and screened

by junction PCR for single-copy insertion at attTn7 (downstream of *glmS*). Reporter function was

validated in plate-reader assays (P_{phzA} induction under QS-permissive conditions). Plasmids and primers

are listed in Tables S3–S4.

In-frame deletion mutants were constructed using a suicide plasmid. Approximately 1-kb regions

upstream and downstream of *phzA-G* and *phzI* were PCR-amplified and assembled with SmaI-digested

pEX18Gm by Gibson assembly. The plasmid was electroporated into *E. coli* DH10B, and transformants

were selected on LB agar with gentamicin (30 µg mL^{-1}). Correct constructs were verified by PCR and

introduced into *P. synxantha 2-79* wild-type strains via triparental conjugation. Merodiploids were

selected on Vogel–Bonner minimal medium (VBMM) containing gentamicin (30 µg mL^{-1}), followed by

counter-selection on sucrose medium (5 g L^{-1}yeast extract, 10 g L^{-1}tryptone, 100 g L^{-1}sucrose). Colonies

in which homologous recombination was successful were identified by PCR. Deletions were confirmed

by diagnostic PCR across the targeted loci and by phenotype: Δ*phzA-G* lacked ABS367nm (PCA readout)

signal and lacked phenazine product, verified via LC-MS analysis. Δ*phzI* required exogenous AHL for

induction and lacked AHL production, verified via LC-MS/MS analysis.

Plant-microbe reporter induction assay

Brachypodium distachyon Bd21-3 seeds were dehusked, sterilized with chlorine gas (see Note S2), plated

on 0.6% Phytagel with 0.5× MS, and vernalized (4 °C, 4 d). Seedlings were grown 2 d (16 h light/8 h

dark, 25 °C), then immersed 30 min in 0.5× MS lacking phosphate (P-lim) containing *P. synxantha* 2-79

at OD$_{600}$ = 0.1. Seedlings were transferred into 1 cc syringes filled with 0.6% Phytagel prepared with 0.5×

MS under the indicated phosphorus condition (P-lim, P-rep, or P-HFO) and incubated vertically for 2 d

prior to imaging.

Seedlings embedded in phytagel were mounted so that roots+phytagel were submerged in sterile 0.5× MS

buffer. Imaging used a Zeiss Lightsheet Z.1 with EC Plan-Neofluar 5×/0.16 air objective. Dual-channel

acquisition captured GFP/mNeonGreen (488 nm ex, 498 nm em) and mCherry (561 nm ex, 571 nm em)

on separate cameras with a dual-band emission filter (Zeiss LBF 488/561). Z-stacks were collected at 5

μm step, XY pixel 0.93 μm, 100 ms exposure per slice; laser power was held constant across conditions

(20% at 488 and 561 nm). Maximum intensity projections (MIPs) were generated for analysis.

All analysis used Fiji/ImageJ with a custom macro. Red channel (P$_{tac}$–mCherry) was median-filtered (r =

4), background-subtracted (rolling ball 50 px), converted to 8-bit, and locally thresholded (Bernsen, r =

15) to build masks. Objects were defined by particle analysis (min size 20 px) and applied to original

red/green images to extract integrated density (IntDen) and area. Mean intensity per object was computed

as IntDen/area for both channels. CSVs were exported for Python analysis.

Planktonic 96-well kinetic experiments

Overnight cultures of *P. synxantha* 2-79 (WT, $\Delta phzA$-G, $\Delta phzI$, $\Delta phoB$, $\Delta phzI$–P_{phzA}) were washed 3× in MOPS containing 10 µM phosphate, and inoculated into 96-well plates at $OD_{600} = 0.01$. Plates were incubated in a Tecan Spark 10M (30°C; orbital shaking amplitude 2.5 mm, 216 rpm) with reads every 15 min for 24 h. OD_{600} was read at 600 nm; ABS367 was read at 367nm for PCA-associated absorbance; mNeonGreen at Ex/Em 485/535 nm. Induction time was approximated from fluorescence time series by manual curation. ODs at induction were obtained by linear interpolation of OD_{600} (t) at that time.

For the AHL-supplemented dose-response assays, 3-OH-C6-HSL (Adipogen; ≥98%) stock was prepared with methanol (final solvent ≤0.02% v/v; vehicle controls showed no effect from solvent carryover). Dose ranges spanned sub-nM to tens of µM (specified per figure).

For the no-growth supplementation assay, $\Delta phzI$ overnight culture was washed (3×) in the respective treatment buffer (No-N or No-N/P) were inoculated at $OD_{600} = 0.1$. For the growth-permissive supplementation assays, $\Delta phzI$ overnight culture was washed (3×) in MOPS containing 10 µM phosphate and inoculated at $OD_{600} = 0.01$. Responses were baseline-corrected to 0 µM within condition and fit with a two-parameter Hill (log-logistic) model (top = 1, bottom = 0) to estimate EC_{50} with 95% CIs.

AHL quantification by LC–MS/MS

Overnight cultures of *P. synxantha* 2-79 (WT, $\Delta phzI$) were washed 3× in MOPS, then grown in glass culture tubes (30 °C, 250 rpm) under P-rep, P-lim, and N-lim conditions for 22 h. Cultures were then pelleted, supernatants were collected, filtered through a 0.22 µm PES filter, diluted 1:1 in methanol, incubated for 1 h at -20°C, pelleted, then supernatants were collected before analyzing.

Analytical standards (3-OH-C6-HSL, 3-OH-C8-HSL, 3-OH-C10-HSL, and 3-oxo-C8-HSL; ≥95–98% purity; Adipogen/Cayman) were prepared by solubilizing powders in methanol as 1-10 g L^{-1} primary

frozen stocks (-20 °C). A 100 µM all analyte mix was made in methanol with 0.1% formic acid.

Calibration standards were prepared by diluting the analyte mix in methanol with 0.1% formic acid.

A Waters (Milford, MA) Acquity Premier UPLC was fitted with a HSS T3 column and coupled to a Xevo

Q-TOF G2S. Mobile phase A was 0.1% formic acid in water and mobile phase B was 100% acetonitrile.

The initial mobile phase composition was 2.5 % B for 0.5 minutes with a linear gradient to 95% B ending

at 5 minutes. After a 0.5 minute hold (5.5 min) the composition was returned rapidly to the initial

conditions (5.6 min) and held for an additional 0.25 minutes (to 5.74 minutes). The gradient was cycled

without interruption during any acquisition. The flow rate was held constant throughout at 0.3 mL/min,

the samples held at 10 °C, the column temperature maintained at 40 °C, and the injection volume was 4

µL. The 2.1 mm × 50 mm × 1.8 µm column was used without a guard column.

The ion source block was held at 120 °C, capillary voltage was 3 kV, desolvation gas flow and

temperature were 600 lph and 500 °C, the cone gas flow as 15 lph, and the source offset was 80 V.

Separate mass spectral data sets were collected in both MS mode from 50-1200 m/z and using scheduled

MS2 for parallel reaction monitoring of the individual targeted analytes. MS data was acquired in positive

sensitivity mode (approximately 18,000 resolution). The mass axis was calibrated against sodium formate

and was confirmed daily. Leucine enkephalin was used as lock mass reference during acquisition.

Targeted analysis was performed using the Quanlynx program within the Masslynx software suite used to

operate the instrument.

Sessile early-stage colony biofilm experiments

Overnight cultures of *P. synxantha* 2-79 (WT, P_{phoA}, P_{phzA}, $P_{PA10403}$) were washed 3× in MOPS containing

7 mM phosphate and adjusted to OD_{600}= 0.05. A 5 µL droplet was spotted onto agar plates, air-dried 1 h,

and placed in a custom humidity-controlled platform. The agar plates contained either 2 mL or 5 mL of

agar. For P-lim plates, phosphorus was omitted from the MOPS agar; the only source of phosphorus was

the 5 µL inoculation drop that contained 7 mM phosphate. P-rep consisted of MOPS agar at 7 mM.

Images were acquired hourly for 36 h.

Time-lapse imaging of bacteria was performed using a previously reported complex-field and

fluorescence microscopy system based on the aperture scanning technique (CFAST).[48] Samples were

mounted in a 12-well plate and positioned on a motorized XY scanning stage (Thorlabs MLS203-1) to

enable automated, parallel scanning of multiple wells.

To minimize evaporation of the growth medium during the two-day imaging experiments, the 12-well

plate was placed in a custom 3D-printed holder containing wet paper towels to maintain high local

humidity. To maintain focus while minimizing photobleaching and phototoxicity, 0.5-µm red fluorescent

beads (580/605 nm, ThermoFisher F8812) were pre-mixed with the bacterial suspension at a $1:10^5$

dilution from stock. These beads served as markers for focal plane tracking. The beads were excited with

a 568 nm laser (Coherent Sapphire 568 LP), and the emitted fluorescence was filtered using a dichroic

mirror (Semrock Di01-R488/561) and a bandpass filter (Semrock FF01-R523/610).

A 488 nm laser (Thorlabs LP488-SF20G) was used both for complex-field illumination and for excitation

of mNeonGreen fluorescence. The complex field data comprising both amplitude and quantitative

phase was used to quantify the area coverage of bacterial colonies. Fluorescence intensity was extracted

from the mNeonGreen images and subsequently normalized using brightfield images to compensate for

variations in excitation power due to differences in medium thickness across the sample.

Microfluidic soil-analog reactor experiments

Devices followed a silicon-etched porous design[64] and fabrication process (see Note S3). Overnight

culture of *P. synxantha* 2-79 (dual reporter, P_{phzA}-P_{tac}) was washed 3× in MOPS P-lim (100 µM) or P-rep

medium and adjusted to OD_{600}= 0.05 for inoculation into the reactor. Cells were inoculated at a flow rate

of 0.1 mL min^{-1} for 3 minutes. Air was then injected and generated heterogeneous water saturation

(connected high-saturation domains vs confined pockets). After initial displacement, air was continuously

injected at a flow rate of 0.01 mL min^{-1}. Imaging was conducted for 16-40 h depending on experiment.

All imaging was performed on a Nikon Ti2 widefield epifluorescence microscope. Time-lapse series used

a Plan Apo 20×/0.75 dry objective. GFP was excited with a 488-nm LED and collected with a FITC filter

set; mCherry was excited with a 561-nm LED and collected with a TRITC filter set. Images were

collected every hour from multiple stage positions.

Custom MATLAB code was used to segment bacterial cells based on the red channel (constitutive

reporter) and pockets of air based on the green channel (induction of phenazine production). To identify

cells, background subtraction was applied to the original image followed by thresholding using a fixed

intensity value. Objects smaller than 5px were discarded. To segment pockets of air, we first identified

and removed signal from induction, which would otherwise interfere with the detection. For this we

segmented fluorescing cells by simple thresholding, expanded their area by dilation (10px) and then

replaced intensity values within this area with the background intensity. Following this, air pockets were

segmented by smoothing and background subtraction followed by thresholding. Areas smaller than

1000px were discarded and holes within air pockets smaller than 50000px were closed. When necessary,

segmentation of air pockets was corrected manually. Intensity ratios were based on background-corrected

intensities. Induction time vs cell density was fit to a power-law model using log–log linear regression;

goodness-of-fit (R^2) and CI on parameters are reported.

Supernatant cross-induction assay

Overnight cultures of *Pseudomonas* strains were washed 3× in MOPS (P-lim and N-lim) medium and

inoculated at OD_{600} =0.01 into glass culture tubes (30 °C, 250 rpm) under P-rep, P-lim, and N-lim

conditions for 22 h. Cultures were then pelleted, and supernatants were collected and filtered through a

0.22 μm PES filter. The cell-free supernatant was then mixed 1:1 with appropriate medium (i.e., P-rep, P-

lim, N-lim) for cross-induction assay.

An overnight culture of *P. synxantha* 2-79 Δ*phzI*-P$_{phzA}$ was washed 3× in MOPS (P-lim and N-lim) and

inoculated at OD$_{600}$ =0.01. A 96-well plate assay was then conducted with the cell-free supernatant

mixture following the "Planktonic 96-well kinetic experiments" parameters. Fold-change for RFU/OD$_{600}$

relative to the reference strain *P. synxantha* Δ*phzI* within each medium was then computed. The resulting

scores were plotted as a single heatmap.

Pairwise competition assays

Co-culture growth and induction dynamics were monitored using a two-chamber co-culture system that

allows fluidic exchange while physically separating populations (Duet Co-Culture System, Cerillo). *P.*

synxantha (WT or Δ*phzA-G*) and *P. kyonggii* overnight cultures were washed 3× in MES (P-lim and N-

lim) medium. Cells were then loaded to either side of the chamber at a starting OD$_{600}$ of 0.01 under N-lim,

P-lim, or P-rep media. Duets have the footprint of a 96-well plate, thus the assay was conducted following

"Planktonic 96-well kinetic experiments" instrument parameters. Phenazine-dependent suppression was

assessed by comparing WT vs Δ*phzA-G*. Statistics used Welch's t-test with two-sided α = 0.05. Note that

competition assays were run in MES-defined medium (pH 5.8) with sucrose as the carbon source. Sucrose

was chosen because *P. synxantha* 2-79 does not utilize it; on glucose, *P. synxantha* rapidly overgrows

Pedobacter kyonggii, obscuring interaction effects. Using sucrose maintains competitive viability of *P.*

kyonggii and shifts *P. synxantha* toward relying on metabolites released by P. *kyonggii*, enabling

resolution of phenazine-dependent outcomes. The pH was set to 5.8 (buffered with MES) to approximate

mildly acidic rhizosphere conditions and because PCA is more membrane-permeant—and thus more

potent—at lower pH.

Acknowledgements:

We acknowledge the Resnick Ecology and Biosphere Engineering Facility and Caltech Biological

Imaging Center for the training and use of confocal and light sheet fluorescence microscopes. We are

grateful to the Resnick Sustainability Institute and National Institute of Health (2R01AI127850-06A1) for

enabling resources and financial support. R.E.A. was supported by an NSF Postdoctoral Research

Fellowship in Biology (Grant No. 2209379). H.J. was supported by a Helen Hay Whitney Postdoctoral

Research Fellowship.

Author contributions: R.E.A., and D.K.N. designed research; R.E.A., H.J., O.Z., R.L.A., and N.D.

performed research; R.E.A., H.J., O.Z., R.L.A., and N.D. analyzed data; and R.E.A., H.J., O.Z., R.L.A.,

N.D., D.M., C.Y., and D.K.N. wrote the paper.

References:

1.   Mavrodi, D.V., Blankenfeldt, W., Mavrodi, O.V., Weller, D.M., and Thomashow, L.S. (2025).
Microbial phenazines: biosynthesis, structural diversity, evolution, regulation, and biological
significance. Microbiology and Molecular Biology Reviews *0*, e00147-23.
https://doi.org/10.1128/mmbr.00147-23.

2.   Dahlstrom, K.M., McRose, D.L., and Newman, D.K. (2020). Keystone metabolites of crop
rhizosphere microbiomes. Current Biology *30*, R1131–R1137.
https://doi.org/10.1016/j.cub.2020.08.005.

3.   Zhou, Y., Wang, H., Sun, J., Wicaksono, W.A., Liu, C., He, Y., Qin, Y., Berg, G., Li, L., Lin, H., et
al. (2025). Phenazines contribute to microbiome dynamics by targeting topoisomerase IV. Nat
Microbiol *10*, 2396–2411. https://doi.org/10.1038/s41564-025-02118-0.

4.   Dar, D., Thomashow, L.S., Weller, D.M., and Newman, D.K. (2020). Global landscape of phenazine
biosynthesis and biodegradation reveals species-specific colonization patterns in agricultural soils
and crop microbiomes. eLife *9*, e59726. https://doi.org/10.7554/eLife.59726.

5.   Menezes-Blackburn, D., Giles, C., Darch, T., George, T.S., Blackwell, M., Stutter, M., Shand, C.,
Lumsdon, D., Cooper, P., Wendler, R., et al. (2018). Opportunities for mobilizing recalcitrant
phosphorus from agricultural soils: a review. Plant Soil *427*, 5–16. https://doi.org/10.1007/s11104-
017-3362-2.

6.   McRose, D.L., and Newman, D.K. (2021). Redox-active antibiotics enhance phosphorus
bioavailability. Science *371*, 1033–1037. https://doi.org/10.1126/science.abd1515.

7.　Cox, C.D. (1986). Role of pyocyanin in the acquisition of iron from transferrin. Infection and
Immunity *52*, 263–270. https://doi.org/10.1128/iai.52.1.263-270.1986.

8.　Hernandez, M.E., Kappler, A., and Newman, D.K. (2004). Phenazines and other redox-active
antibiotics promote microbial mineral reduction. Appl Environ Microbiol *70*, 921–928.
https://doi.org/10.1128/AEM.70.2.921-928.2004.

9.　Wang, Y., Wilks, J.C., Danhorn, T., Ramos, I., Croal, L., and Newman, D.K. (2011). Phenazine-1-
Carboxylic Acid Promotes Bacterial Biofilm Development via Ferrous Iron Acquisition. Journal of
Bacteriology *193*, 3606–3617. https://doi.org/10.1128/jb.00396-11.

10.　LeTourneau, M.K., Marshall, M.J., Grant, M., Freeze, P.M., Strawn, D.G., Lai, B., Dohnalkova,
A.C., Harsh, J.B., Weller, D.M., and Thomashow, L.S. (2019). Phenazine-1-Carboxylic Acid-
Producing Bacteria Enhance the Reactivity of Iron Minerals in Dryland and Irrigated Wheat
Rhizospheres. Environ. Sci. Technol. *53*, 14273–14284. https://doi.org/10.1021/acs.est.9b03962.

11.　Price-Whelan, A., Dietrich, L.E.P., and Newman, D.K. (2006). Rethinking "secondary" metabolism:
physiological roles for phenazine antibiotics. Nat Chem Biol *2*, 71–78.
https://doi.org/10.1038/nchembio764.

12.　Meirelles, L.A., and Newman, D.K. (2018). Both toxic and beneficial effects of pyocyanin contribute
to the lifecycle of Pseudomonas aeruginosa. Molecular Microbiology *110*, 995–1010.
https://doi.org/10.1111/mmi.14132.

13.　Perry, E.K., and Newman, D.K. (2022). Prevalence and Correlates of Phenazine Resistance in
Culturable Bacteria from a Dryland Wheat Field. Applied and Environmental Microbiology *88*,
e02320-21. https://doi.org/10.1128/aem.02320-21.

14.　Thomashow, L.S., and Weller, D.M. (1988). Role of a phenazine antibiotic from Pseudomonas
fluorescens in biological control of Gaeumannomyces graminis var. tritici. J Bacteriol *170*, 3499–
3508. https://doi.org/10.1128/jb.170.8.3499-3508.1988.

15.　Chin-A-Woeng, T.F.C., Bloemberg, G.V., and Lugtenberg, B.J.J. (2003). Phenazines and their role in
biocontrol by Pseudomonas bacteria. New Phytologist *157*, 503–523. https://doi.org/10.1046/j.1469-
8137.2003.00686.x.

16.　LeTourneau, M.K., Marshall, M.J., Cliff, J.B., Bonsall, R.F., Dohnalkova, A.C., Mavrodi, D.V.,
Devi, S.I., Mavrodi, O.V., Harsh, J.B., Weller, D.M., et al. (2018). Phenazine-1-carboxylic acid and
soil moisture influence biofilm development and turnover of rhizobacterial biomass on wheat root
surfaces. Environmental Microbiology *20*, 2178–2194. https://doi.org/10.1111/1462-2920.14244.

17.　Fuqua, W.C., Winans, S.C., and Greenberg, E.P. (1994). Quorum sensing in bacteria: the LuxR-LuxI
family of cell density-responsive transcriptional regulators. J Bacteriol *176*, 269–275.
https://doi.org/10.1128/jb.176.2.269-275.1994.

18.　Venturi, V. (2006). Regulation of quorum sensing in Pseudomonas. FEMS Microbiol Rev *30*, 274–
291. https://doi.org/10.1111/j.1574-6976.2005.00012.x.

19.　Miller, M.B., and Bassler, B.L. (2001). Quorum Sensing in Bacteria. Annual Review of
Microbiology *55*, 165–199. https://doi.org/10.1146/annurev.micro.55.1.165.

20. Waters, C.M., and Bassler, B.L. (2005). QUORUM SENSING: Cell-to-Cell Communication in
Bacteria. Annual Review of Cell and Developmental Biology *21*, 319–346.
https://doi.org/10.1146/annurev.cellbio.21.012704.131001.

21. Parsek, M.R., and Greenberg, E.P. (2005). Sociomicrobiology: the connections between quorum
sensing and biofilms. Trends Microbiol *13*, 27–33. https://doi.org/10.1016/j.tim.2004.11.007.

22. Papenfort, K., and Bassler, B.L. (2016). Quorum sensing signal–response systems in Gram-negative
bacteria. Nat Rev Microbiol *14*, 576–588. https://doi.org/10.1038/nrmicro.2016.89.

23. Abisado, R.G., Benomar, S., Klaus, J.R., Dandekar, A.A., and Chandler, J.R. (2018). Bacterial
Quorum Sensing and Microbial Community Interactions. mBio *9*, e02331-17.
https://doi.org/10.1128/mBio.02331-17.

24. Whiteley, M., Diggle, S.P., and Greenberg, E.P. (2017). Bacterial quorum sensing: the progress and
promise of an emerging research area. Nature *551*, 313–320. https://doi.org/10.1038/nature24624.

25. Mukherjee, S., and Bassler, B.L. (2019). Bacterial quorum sensing in complex and dynamically
changing environments. Nat Rev Microbiol *17*, 371–382. https://doi.org/10.1038/s41579-019-0186-5.

26. Raynaud, X., and Nunan, N. (2014). Spatial Ecology of Bacteria at the Microscale in Soil. PLOS
ONE *9*, e87217. https://doi.org/10.1371/journal.pone.0087217.

27. Wang, G., and Or, D. (2010). Aqueous films limit bacterial cell motility and colony expansion on
partially saturated rough surfaces. Environmental Microbiology *12*, 1363–1373.
https://doi.org/10.1111/j.1462-2920.2010.02180.x.

28. Tecon, R., and Or, D. (2017). Biophysical processes supporting the diversity of microbial life in soil.
FEMS Microbiol Rev *41*, 599–623. https://doi.org/10.1093/femsre/fux039.

29. Mavrodi, D.V., Mavrodi, O.V., Parejko, J.A., Bonsall, R.F., Kwak, Y.-S., Paulitz, T.C., Thomashow,
L.S., and Weller, D.M. (2012). Accumulation of the Antibiotic Phenazine-1-Carboxylic Acid in the
Rhizosphere of Dryland Cereals. Applied and Environmental Microbiology *78*, 804–812.
https://doi.org/10.1128/AEM.06784-11.

30. Mavrodi, D.V., Mavrodi, O.V., Elbourne, L.D.H., Tetu, S., Bonsall, R.F., Parejko, J., Yang, M.,
Paulsen, I.T., Weller, D.M., and Thomashow, L.S. (2018). Long-Term Irrigation Affects the
Dynamics and Activity of the Wheat Rhizosphere Microbiome. Front. Plant Sci. *9*.
https://doi.org/10.3389/fpls.2018.00345.

31. Schuhegger, R., Ihring, A., Gantner, S., Bahnweg, G., Knappe, C., Vogg, G., Hutzler, P., Schmid, M.,
Van Breusegem, F., Eberl, L., et al. (2006). Induction of systemic resistance in tomato by N-acyl-L-
homoserine lactone-producing rhizosphere bacteria. Plant Cell Environ *29*, 909–918.
https://doi.org/10.1111/j.1365-3040.2005.01471.x.

32. DeAngelis, K.M., Firestone, M.K., and Lindow, S.E. (2007). Sensitive Whole-Cell Biosensor
Suitable for Detecting a Variety of N-Acyl Homoserine Lactones in Intact Rhizosphere Microbial
Communities. Appl Environ Microbiol *73*, 3724–3727. https://doi.org/10.1128/AEM.02187-06.

33. DeAngelis, K.M., Lindow, S.E., and Firestone, M.K. (2008). Bacterial quorum sensing and nitrogen
cycling in rhizosphere soil. FEMS Microbiol Ecol *66*, 197–207. https://doi.org/10.1111/j.1574-
6941.2008.00550.x.

34. Redfield, R.J. (2002). Is quorum sensing a side effect of diffusion sensing? Trends in Microbiology
10, 365–370. https://doi.org/10.1016/S0966-842X(02)02400-9.

35. Hense, B.A., Kuttler, C., Müller, J., Rothballer, M., Hartmann, A., and Kreft, J.-U. (2007). Does
efficiency sensing unify diffusion and quorum sensing? Nat Rev Microbiol *5*, 230–239.
https://doi.org/10.1038/nrmicro1600.

36. Platt, T.G., and Fuqua, C. (2010). What's in a name? The semantics of quorum sensing. Trends
Microbiol *18*, 383–387. https://doi.org/10.1016/j.tim.2010.05.003.

37. West, S.A., Winzer, K., Gardner, A., and Diggle, S.P. (2012). Quorum sensing and the confusion
about diffusion. Trends in Microbiology *20*, 586–594. https://doi.org/10.1016/j.tim.2012.09.004.

38. Boedicker, J.Q., Vincent, M.E., and Ismagilov, R.F. (2009). Microfluidic Confinement of Single
Cells of Bacteria in Small Volumes Initiates High-Density Behavior of Quorum Sensing and Growth
and Reveals Its Variability. Angewandte Chemie International Edition *48*, 5908–5911.
https://doi.org/10.1002/anie.200901550.

39. Darch, S.E., Simoska, O., Fitzpatrick, M., Barraza, J.P., Stevenson, K.J., Bonnecaze, R.T., Shear,
J.B., and Whiteley, M. (2018). Spatial determinants of quorum signaling in a Pseudomonas
aeruginosa infection model. Proceedings of the National Academy of Sciences *115*, 4779–4784.
https://doi.org/10.1073/pnas.1719317115.

40. Stewart, P.S. (2003). Diffusion in Biofilms. Journal of Bacteriology *185*, 1485–1491.
https://doi.org/10.1128/jb.185.5.1485-1491.2003.

41. Parsek, M.R., and Greenberg, E.P. (2005). Sociomicrobiology: the connections between quorum
sensing and biofilms. Trends Microbiol *13*, 27–33. https://doi.org/10.1016/j.tim.2004.11.007.

42. Kim, M.K., Ingremeau, F., Zhao, A., Bassler, B.L., and Stone, H.A. (2016). Local and global
consequences of flow on bacterial quorum sensing. Nat Microbiol *1*, 15005.
https://doi.org/10.1038/nmicrobiol.2015.5.

43. Hense, B.A., Müller, J., Kuttler, C., and Hartmann, A. (2012). Spatial Heterogeneity of Autoinducer
Regulation Systems. Sensors *12*, 4156–4171. https://doi.org/10.3390/s120404156.

44. Thomashow, L.S., and Weller, D.M. (1988). Role of a phenazine antibiotic from Pseudomonas
fluorescens in biological control of Gaeumannomyces graminis var. tritici. J Bacteriol *170*, 3499–
3508. https://doi.org/10.1128/jb.170.8.3499-3508.1988.

45. Mavrodi, D.V., Ksenzenko, V.N., Bonsall, R.F., Cook, R.J., Boronin, A.M., and Thomashow, L.S.
(1998). A Seven-Gene Locus for Synthesis of Phenazine-1-Carboxylic Acid by Pseudomonas
fluorescens 2-79. J Bacteriol *180*, 2541–2548.

46. Khan, S.R., Mavrodi, D.V., Jog, G.J., Suga, H., Thomashow, L.S., and Farrand, S.K. (2005).
Activation of the phz Operon of Pseudomonas fluorescens 2-79 Requires the LuxR Homolog PhzR,

N-(3-OH-Hexanoyl)-l-Homoserine Lactone Produced by the LuxI Homolog PhzI, and a cis-Acting phz Box. J Bacteriol *187*, 6517–6527. https://doi.org/10.1128/JB.187.18.6517-6527.2005.

47. Khan, S.R., Herman, J., Krank, J., Serkova, N.J., Churchill, M.E.A., Suga, H., and Farrand, S.K. (2007). N-(3-Hydroxyhexanoyl)-l-Homoserine Lactone Is the Biologically Relevant Quormone That Regulates the phz Operon of Pseudomonas chlororaphis Strain 30-84. Applied and Environmental Microbiology *73*, 7443–7455. https://doi.org/10.1128/AEM.01354-07.

48. Zhang, O., Alcalde, R.E., Zhou, H., Yin, S., Newman, D.K., and Yang, C. (2024). Investigating 3D microbial community dynamics of the rhizosphere using quantitative phase and fluorescence microscopy. Proceedings of the National Academy of Sciences *121*, e2403122121. https://doi.org/10.1073/pnas.2403122121.

49. Holford, I.C.R. (1997). Soil phosphorus: its measurement, and its uptake by plants. Soil Res. *35*, 227–240. https://doi.org/10.1071/s96047.

50. Hinsinger, P. (2001). Bioavailability of soil inorganic P in the rhizosphere as affected by root-induced chemical changes: a review. Plant and Soil *237*, 173–195. https://doi.org/10.1023/A:1013351617532.

51. Ducousso-Détrez, A., Fontaine, J., Lounès-Hadj Sahraoui, A., and Hijri, M. (2022). Diversity of Phosphate Chemical Forms in Soils and Their Contributions on Soil Microbial Community Structure Changes. Microorganisms *10*, 609. https://doi.org/10.3390/microorganisms10030609.

52. Whooley, M.A., and McLoughlin, A.J. (1982). The regulation of pyocyanin production in Pseudomonas aeruginosa. European J. Appl. Microbiol. Biotechnol. *15*, 161–166. https://doi.org/10.1007/BF00511241.

53. Jensen, V., Löns, D., Zaoui, C., Bredenbruch, F., Meissner, A., Dieterich, G., Münch, R., and Häussler, S. (2006). RhlR Expression in Pseudomonas aeruginosa Is Modulated by the Pseudomonas Quinolone Signal via PhoB-Dependent and -Independent Pathways. JB *188*, 8601–8606. https://doi.org/10.1128/JB.01378-06.

54. Oglesby, A.G., Farrow, J.M., Lee, J.-H., Tomaras, A.P., Greenberg, E.P., Pesci, E.C., and Vasil, M.L. (2008). The influence of iron on Pseudomonas aeruginosa physiology: a regulatory link between iron and quorum sensing. J Biol Chem *283*, 15558–15567. https://doi.org/10.1074/jbc.m707840200.

55. Lee, J., Wu, J., Deng, Y., Wang, J., Wang, C., Wang, J., Chang, C., Dong, Y., Williams, P., and Zhang, L.-H. (2013). A cell-cell communication signal integrates quorum sensing and stress response. Nat Chem Biol *9*, 339–343. https://doi.org/10.1038/nchembio.1225.

56. Nguyen, A.T., Jones, J.W., Ruge, M.A., Kane, M.A., and Oglesby-Sherrouse, A.G. (2015). Iron Depletion Enhances Production of Antimicrobials by Pseudomonas aeruginosa. J Bacteriol *197*, 2265–2275. https://doi.org/10.1128/JB.00072-15.

57. Welsh, M.A., and Blackwell, H.E. (2016). Chemical Genetics Reveals Environment-Specific Roles for Quorum Sensing Circuits in *Pseudomonas aeruginosa*. Cell Chemical Biology *23*, 361–369. https://doi.org/10.1016/j.chembiol.2016.01.006.

58. Meng, X., Ahator, S.D., and Zhang, L.-H. (2020). Molecular Mechanisms of Phosphate Stress Activation of Pseudomonas aeruginosa Quorum Sensing Systems. mSphere *5*, 10.1128/msphere.00119-20. https://doi.org/10.1128/msphere.00119-20.

59. Monds, R.D., Newell, P.D., Schwartzman, J.A., and O'Toole, G.A. (2006). Conservation of the Pho
regulon in Pseudomonas fluorescens Pf0-1. Appl Environ Microbiol *72*, 1910–1924.
https://doi.org/10.1128/AEM.72.3.1910-1924.2006.

60. Hsieh, Y.-J., and Wanner, B.L. (2010). Global Regulation by the Seven-component Pi Signaling
System. Curr Opin Microbiol *13*, 198–203. https://doi.org/10.1016/j.mib.2010.01.014.

61. Santos-Beneit, F. (2015). The Pho regulon: a huge regulatory network in bacteria. Front. Microbiol.
6. https://doi.org/10.3389/fmicb.2015.00402.

62. Goldberg, S., and Sposito, G. (1984). A Chemical Model of Phosphate Adsorption by Soils: I.
Reference Oxide Minerals. Soil Science Soc of Amer J *48*, 772–778.
https://doi.org/10.2136/sssaj1984.03615995004800040015x.

63. Dzombak, D.A., and Morel, F. (1990). Surface complexation modeling: hydrous ferric oxide (Wiley).

64. Boyd, V., Yoon, H., Zhang, C., Oostrom, M., Hess, N., Fouke, B., Valocchi, A.J., and Werth, C.J.
(2014). Influence of Mg2+ on CaCO3 precipitation during subsurface reactive transport in a
homogeneous silicon-etched pore network. Geochimica et Cosmochimica Acta *135*, 321–335.
https://doi.org/10.1016/j.gca.2014.03.018.

65. van Delden, C., Comte, R., and Bally, A.M. (2001). Stringent Response Activates Quorum Sensing
and Modulates Cell Density-Dependent Gene Expression inPseudomonas aeruginosa. Journal of
Bacteriology *183*, 5376–5384. https://doi.org/10.1128/jb.183.18.5376-5384.2001.

66. Diggle, S.P., Winzer, K., Chhabra, S.R., Worrall, K.E., Cámara, M., and Williams, P. (2003). The
Pseudomonas aeruginosa quinolone signal molecule overcomes the cell density-dependency of the
quorum sensing hierarchy, regulates rhl-dependent genes at the onset of stationary phase and can be
produced in the absence of LasR. Molecular Microbiology *50*, 29–43. https://doi.org/10.1046/j.1365-
2958.2003.03672.x.

67. Dubern, J.-F., Halliday, N., Cámara, M., Winzer, K., Barrett, D.A., Hardie, K.R., and Williams, P.
(2023). Growth rate and nutrient limitation as key drivers of extracellular quorum sensing signal
molecule accumulation in Pseudomonas aeruginosa. Microbiology *169*, 001316.
https://doi.org/10.1099/mic.0.001316.

68. W. G. Sunda, N. M. Price, and F. M. M. Morel (2005). Algal culturing techniques R. A. Andersen,
ed. (Elsevier/Academic Press).

69. Choi, K.-H., and Schweizer, H.P. (2006). mini-Tn 7 insertion in bacteria with single att Tn 7 sites:
example Pseudomonas aeruginosa. Nat Protoc *1*, 153–161. https://doi.org/10.1038/nprot.2006.24.

811