Tunneling nanotubes propagate a BMP-dependent preneoplastic state.

Cristina Cuella Martin[1,2,3], Thanh Thuy Nguyen *[1,2,3], Alexia Bertin *[1,2,3], Kevin Geistlich[1,2,3,4], Boris Guyot[1,2,3], Sylvain Lefort[1,2,3], Emmanuel Delay[1,2,3,4], Charline Dalverny[3,5], Alice Boissard[6], Cécile Henry[6], François Guillonneau[6,7,8], Eve-Isabelle Pécheur §*[1,2,3] and Véronique Maguer-Satta §*[1,2,3,4]

1. CNRS UMR5286, Centre de Recherche en Cancérologie de Lyon, 69008 Lyon, France;

2. Inserm U1052, Centre de Recherche en Cancérologie de Lyon, 69008 Lyon, France;

3. Université Claude Bernard Lyon 1, Lyon, France;

4. Centre Léon Bérard, 69008 Lyon, France;

5. Centre Technologique des Microstructures (CTµ), 69622 Villeurbanne, France;

6. Prot'ICO proteomics facility, Institut de Cancérologie de l'Ouest, 49055 Angers, France,

7. Nantes Université, Inserm, CNRS, CRCI2NA, U1307, UMR6075, 44000 Nantes, France;

8. Equipe Labellisée Ligue Contre le Cancer, Paris, France.

* These authors contributed equally; § Co-corresponding authors

Co-corresponding authors: Eve-Isabelle PÉCHEUR and Véronique MAGUER-SATTA, Centre de Recherche en Cancérologie de Lyon-CRCL, UMR CNRS 5286/Inserm U1052, 28 rue Laënnec, 69373 Lyon Cedex 08, France; 33-4-7878-2907 or 33-4-7878-2806.

Emails: eve-isabelle.pecheur@inserm.fr ; veronique.maguer-satta@lyon.unicancer.fr

Keywords: Tunneling nanotubes, cell communication, preneoplasia, BMP, luminal breast tumor initiation, stem cells.

Word count : 3321.

Abstract.

Tunneling nanotubes (TNTs) are thin, actin-based structures allowing long-distance communication between cells through the transfer of molecules and organelles. Although well-known to contribute to cancer progression, their role in early tumor initiation remains as yet unexplored. Here we address this question by using normal human mammary primary cells and an in-house human breast cell model recapitulating early luminal breast transformation. TNTs become more abundant and elongated during this process, and preferentially connect transformed donor cells to non-transformed acceptor cells. We show that this long-range directional communication involves the transfer of the signaling receptor BMPR1b. Within days, this transfer induces gene expression changes in acceptor cells, consistent with early transformation programs. Functional analyses of these acceptor cells revealed phenotypic changes, including anchorage-independent growth. In particular, the transferred BMPR1b receptor sensitized acceptor cells to BMP2 signals present in the microenvironment, amplifying their transformation potential. Hence, by tracking the earliest molecular responses in acceptor cells, we deciphered the initial steps of a transformation cascade triggered by TNT-mediated transfer. These findings uncover a BMP-dependent mechanism by which transformed cells propagate a preneoplastic state to adjacent and distant cells at the very onset of transformation, offering new perspectives on how epithelial transformation arises and spreads to neighboring cells.

With 2.3 million women affected worldwide and 700,000 deaths per year, breast cancer is the leading cause of cancer-related death in women, with luminal types representing 70% of all cases (*1*), and ductal carcinoma *in situ* (DCIS) the earliest non-invasive stage. The Bone Morphogenetic Proteins (BMP) pathway is a key actor altered during luminal breast tumorigenesis (*2–4*). Together with an interleukin-6 (IL6)-mediated inflammation, enhanced BMP2 secretion by the niche surrounding mammary stem cells fuels their transformation into cancer stem cells *via* amplification of their response to the kinase BMP receptor type 1b (BMPR1b), without detectable genetic alterations (*3, 5*). Cancer stem cells can drive tumorigenesis and chemoresistance through exogenous signaling pathways that maintain their stemness and promote survival (*6, 7*). BMPR1b-silenced transformed mammary cells have drastically decreased capacities to initiate tumor formation *in vivo* (*3*). However, the early dynamics of BMP-mediated transformation and the evolution of non-transformed mammary cells towards tumor cells remain unexplored.

Tunneling nanotubes (TNTs) are thin, transient actin-based structures enabling long-range intercellular communication (*8, 9*), *via* the transfer of organelles, small molecules, proteins and nucleic acids. TNTs have been implicated in tumor progression, metastasis, and chemoresistance. Cancer cells linked to neurons by TNTs exhibited enhanced metastatic capacities (*10–12*). Yet the role of TNTs at earliest stages of tumorigenesis is unknown.

While BMPs and TNTs have independently been associated with breast cancer, their potential interaction is unexplored. Also, the nature, directionality, and functional consequences of cell/cell communications at the tumor initiation stage remain ill-defined.

In this work, we aimed to delineate the interplay between the BMP pathway and TNT-mediated communication, to shed light on the earliest steps of transformation of (immature) mammary cells.

Results.

TNT formation during mammary epithelial stem cells transformation.

Early BMP-driven transformation dynamics was explored by using our MCF10A-derived stem

cell models obtained by chronic exposure to IL6 and BMP2 (MC26 and M1B26; Fig. 1A) (*3*,

5). Sorting of BMPR1b-expressing cells prior to cytokine treatment, followed by soft agar

clones selection, led to quicker establishment of transformed cells (M1B26), with greater *in*

vitro anchorage-independent growth and *in vivo* engraftment than in non BMPR1b-sorted cells

(MC26). These models display DCIS molecular features and robustly recapitulate the *in vivo*

BMP pathway alterations (*3, 5)*. Confocal microscopy observations of non-transformed control

(MCF10A-CT) and transformed cells (MC26, M1B26) revealed long-range contacts *via* long

thin F-actin-rich TNTs (*13*) (Fig. 1B). Although TNT quantification requires low-confluence

cultures (*14*), long-range TNTs were also detectable at high cell density (Fig. S1A).

Transformed cells exhibited over twice as many TNTs as non-transformed cells (Fig. 1C, left).

The proportion of long TNTs increased ~fivefold with transformation (Fig. 1C, right). However,

TNT average length did not differ between cell lines (Fig. S1B). Neither TNT number nor length

varied in non-transformed cells exposed to conditioned medium of transformed ones (Fig. S1B-

D). TNTs in transformed cells were often branched, with ramifications and bulges (Fig. 1D

black arrow) (*8, 13*). While most TNTs were open-ended, allowing a cell-cell cytoplasmic

continuum (Fig. 1E), some close-ended TNTs featuring a Gap junction at one tip were seen

(Fig. S1E). TNTs did not adhere to the substrate and were observed within 3D structures such

as acini, basic functional units of breast glandular tissue (Fig. 1F; Fig. S1F). TNTs also

contained microtubules (*8, 15*) (Fig. S1G), and mitochondria (*16*) (Fig. 1G; Fig. S1H).

Importantly, transformed cells could also establish TNT connections with non-transformed

primary fibroblasts (Fig. 1H) and epithelial mammary cells (Fig. 1I) isolated from healthy

human breast samples.

This indicates that early mammary cell transformation leads to an enhanced ability of

transformed cells to connect to their neighbors through TNTs.

Directional transfer of cellular material *via* TNTs from transformed to non-transformed

cells. To investigate how TNTs connect cells, we cocultured mCherry-expressing MCF10A-CT

with GFP-expressing M1B26 cells, seeded at low density (*14*). Both M1B26 transformed (Tf)

and MCF10A-CT non-transformed (NoTf) cells formed TNT structures, connecting Tf to Tf,

NoTf to NoTf and Tf to NoTf cells (Fig. 2A). Quantification of TNT contacts revealed that

connections were more frequent and abundant from Tf towards NoTf cells than in the opposite

direction (Fig. 2B,C). Fluorescent material from GFP-tagged Tf cells accumulated within the

cytoplasm of NoTf cells (Fig. 2D), suggesting a directional transfer *via* TNTs, Tf cells as donors

and NoTf cells as acceptors (Fig. 2E; Movie S1; Fig. S2A). To assess the proportion of cellular

exchange between cells, mCherry-NoTf cells were cocultured with GFP-Tf cells for 8 days.

Flow cytometry analysis revealed a higher proportion of mCherry-NoTf cells exhibiting green

fluorescence uptake compared to GFP-Tf cells acquiring red fluorescence (Fig. S2B), indicating

a preferential Tf-to-NoTf directional transfer (Fig. 2F-H). Switching fluorophores between Tf

and NoTf cells yielded similar results (Fig. S2C,D), and this directional transfer increased over

time (Fig. S2E-I). When using normal human primary mammary cells as NoTf cell sources, we

also observed a Tf-to-NoTf directional material transfer in coculture (Fig. 2J,K), as with

MCF10A-CT cells (Fig. S2J). To evaluate the contribution of cell-cell communication through

secretion of soluble compounds and extracellular vesicles, we performed cultures of normal

human primary cells: (i) with Tf cells separated by a physical barrier (Transwell™), to impede

TNT formation; (ii) in Tf cells conditioned medium (CM, schematic Fig. 2I). In the former

condition, the number of acceptor cells dropped in both transfer directions (Fig. 2J,K; Fig. S2J).

In the latter, no fluorescence exchange *via* soluble components was observed (Fig. S2K,L).

Therefore, directional material transfer preferentially occurs *via* TNTs from Tf to NoTf cells,

and rather than through other forms of cell-cell communication.

TNT-mediated transfer modulates proteomic features of acceptor cells.

Fluorescently-labeled cells cultured for 8 days (Fig. 3A), were sorted by flow cytometry into

four populations: non-transformed (NoTf), transformed (Tf), non-transformed cells that

received fluorescent material (NoTf accept), and non-transformed cells from the same coculture

that did not receive any material (NoTf no accept). Unbiased proteomic screening of each

population followed by gene enrichment pathway analysis revealed that most enriched

pathways in NoTf accept, compared to NoTf no accept of the same coculture, were related to

transport (organic/nitrogen compounds, protein, transmembrane, intracellular), and to

localization of intracellular organelles, macromolecules and proteins (Fig. 3B). Other pathways

pointed to mitochondrion organization and activity. Volcano plot analysis revealed that NoTf

accept were statistically distinct from NoTf no accept cells of the same coculture, and from

NoTf in monoculture (Fig. 3C; Fig. S3A). Of note, the most overexpressed proteins in the

former phenotype were involved in lipid metabolism and transport (*ATP8B1, RAB6B, CES1 –*

EST1, *MEST*, *PIGA*), intracellular membrane and secretory vesicle trafficking (*RAB6B,*

RAB11B, SYT8, CAV2, TMEM167A), cell growth and morphology maintenance (*TRIQK*), membrane protrusion formation (*ATP8B1*) and actin filament bundling (*LCP1* – Plastin-2) (Table S1). Other overexpressed proteins were linked to mitochondrial activities (*COX7C, MALSU1*), possibly related to the presence of mitochondria within TNTs (Fig. 1G; Figs. S1H, S3B).

Six proteins, EST1, Plastin-2, Rab-6B, TRIQK, MEST and MASU1, were markedly overexpressed in NoTf accept compared to NoTf no accept cells from same cocultures (Fig. 3D; Fig. S3C). Corresponding RNA levels were also enhanced in Tf compared to NoTf cells (Fig. S3D). The TNT-mediated transfer of material therefore induces in acceptor cells transcriptional and translational upregulation of key actors involved in progression towards a transformed-like phenotype. This aligns with the downregulation of DAP1 (*DAP*) and thymosin β-10 (*TMSB10*), respectively involved in ribosome hibernation and actin polymerization inhibition (Table S1).

When TNT formation was physically impeded, acceptor cells showed phenotypic differences compared to those in direct coculture (Table S2), but remained comparable to NoTf cells in monoculture (Fig. S3E,F). Notably, NoTf accept cells from direct cocultures exhibited a higher expression of early inflammatory markers (*SERPINA3, CD14*) and motility-related genes (*IGFBP6*) than their Transwell™ counterparts. These findings indicate that TNT-mediated direct contacts with transformed cells induce profound changes in the proteomic landscape of non-transformed acceptor cells, similar to those observed in early transformed cells.

Ras-driven oncogenic transformation being a well-established process, we next investigated whether it could also promote the formation of TNTs. Proteomics revealed a marked increase in H-Ras and K-Ras oncoproteins in NoTf accept cells compared to NoTf cells in monoculture (Fig. 3E). We then examined two additional models involving Ras-mediated transformation of mammary epithelial cells. First, in the Weinberg's model of human mammary epithelial cells immortalized with SV40-L antigen and hTERT, then transformed with H-Ras (*17*), TNTs were present in malignant and non-malignant cells, with shorter TNTs in the former (Fig. 3F,H). Second, in H-Ras-transformed immature MCF10A-derived AT1 and DCIS cells, malignant cells exhibited fewer TNTs than non-malignant ones, without length difference (Fig. 3G,I). Hence, Ras-induced transformation does not enhance long-distance TNT-mediated communication, contrary to clinically relevant inflammation- and BMP-driven mechanisms (*3*).

TNT-mediated transfer impacts the BMP pathway of recipient cells.

Since transmembrane and protein transports were the most impacted processes (Fig. 3B), and given that the BMP pathway is a key driver of luminal transformation (*3*, *4*) (Fig. S4A,B), we examined whether this pathway was affected in non-transformed acceptor cells. We focused on the canonical SMAD cascade, and among non-canonical MAPK downstream cascades, on p38 (MAPK14), because the p-p38/p38 ratio exhibited the highest level of expression in Tf cells (Fig. S4A,B). Eight target or regulator proteins of these cascades were extracted from our proteomic screening (Table S3), and assessed for their expression across sorted cell populations post-TNT connections (Fig. 3A). Several proteins involved in the BMP pathway — including ID1, Nup214 (*18*), spartin, and USP15 (*19*) — were consistently up- or down-regulated in NoTf acceptor cells, reaching levels comparable to those observed in Tf cells, exclusively after TNT transfer (Fig. 4A). ATF2, a transcription regulator positioned at the BMP pathway/p38 cascade interface, was underexpressed in NoTf acceptor cells, at levels comparable to those in Tf cells. Components of the p38 cascade were markedly altered following TNT-mediated transfer: MAPKAPK2, a kinase activated by p38 phosphorylation, was underexpressed in NoTf acceptor cells, consistent with the role of p38 inhibition in facilitating mammary tumorigenesis (*20*). Conversely, Rac1—part of a complex involved in actin bundling and stabilization, regulated by p38—was overexpressed in NoTf cells that received material *via* TNTs (Fig. 4A). BMPR1b expression increased with transformation, and the receptor was localized within TNTs, particularly in bulges, in normal primary and model mammary cells (Fig. 4B). This was concomitant to the activation of both canonical p-SMAD1/5/8 / SMAD1/5/8, and non-canonical p-p38/p38 and p-ERK/ERK cascades (Fig. S4A,B). Correlative light (scanning) electron microscopy (CLEM) enables to distinguish intra- from extra-cellular epitopes by overlaying optical and scanning electron images of the same region. Confocal imaging revealed a continuum of BMPR1b in cell bodies, as well as in or on TNTs. However, electron microscopy did not reveal any extracellular BMPR1b epitopes at TNTs (Fig. 4C). This suggests an extracellular localization in the cell body and an intraluminal presence in TNTs, consistent with BMPR1b trafficking through/within TNTs. Although incomplete (Fig. 4D,E top), *BMPR1b* silencing in Tf cells drastically reduced TNT number, and abolished the formation of long TNTs (Fig. 4E bottom). Silencing of G protein-coupled estrogen receptor 1 (*GPER*) was used as control, since no report linked GPER to TNT formation/function. Though more efficient than *BMPR1b* silencing (Fig. 4E top), *GPER* knockdown had no effect on TNT number or length (Fig. 4D,E bottom). Therefore, BMPR1b contributes to the formation and elongation of (long) TNTs.

To evaluate whether BMPR1b transfer *via* TNTs could sensitize NoTf cells to a BMP2/IL6-

enriched microenvironment, the four sorted cell populations (Figs. 3A, 4F) were analyzed for

BMPR1b mRNA expression. It was found ~8-fold higher in NoTf accept cells previously

connected *via* TNTs, compared to an only ~2-fold increase when TNT formation was blocked

(Fig. 4G). This supports a role for a TNT-mediated activation of the BMP pathway.

These findings underscores BMP pathway activation in NoTf cells acceptor of material from

Tf cells, *via* TNT-mediated transfer. BMPR1b plays a dual role by promoting TNT formation

and being transferred to acceptor cells.

TNT-trafficked BMPR1b sensitizes acceptor cells to a transforming microenvironment.

Long-term impact of BMPR1b transfer was assessed on sorted cells, cultured for up to 16 weeks

with or without BMP2/IL6, mimicking *in vivo* luminal breast tumor conditions (*2*) (Fig. 4F).

BMPR1b protein levels rose between weeks 6 and 10 before stabilizing, in NoTf accept cells

exposed to cytokines after sorting (Fig. 5A left), reaching the highest expression among all

populations by week 14 (Fig. 5A right; Fig. S4C,D). Accordingly, p-SMAD1/5/8 / SMAD1/5/8

and p-p38/p38 cascades were activated upon 4 to 6 week-exposure to BMP2/IL6 in NoTf accept

cells (Fig. 5B,C; Fig. S4E,F). While the SMAD cascade did not return to baseline over weeks

(Fig. 5B), the p38 cascade, after a strong activation as early as 4 weeks with cytokines post-

sorting, reached baseline levels after 10 weeks (Fig. 5C).

We then cross-referenced our proteomic and transcriptomic datasets (*5*), and identified three

tumor-associated genes significantly upregulated during transformation: *HOXB6* (*21*), *CCND2*

(*22*), and *EREG* (*23*) (Fig. 5D). Their mRNA levels were increased in NoTf acceptor cells

connected through TNTs with Tf cells, compared to NoTf cells in monoculture, and to NoTf

acceptor cells for which TNT formation was blocked (Fig. 5E). This supports the notion that

TNT-mediated transfer may promote a shift towards a preneoplastic state. HoxB6, Cyclin D2

and (pro)-epiregulin protein levels also tended to increase with transformation (Fig. S5A).

HRAS was overexpressed in NoTf acceptor cells, further substantiating a shift towards a

preneoplastic status (Fig. 4E). However, *HRAS* was neither involved in TNT formation, nor in

TNT-mediated preneoplastic switch of acceptor cells (Fig. 3F-I). As a negative control, we

selected *UFM1*, a gene involved in endoplasmic reticulum stress, a cellular insult suggested to

contribute to TNT development (*24*). *UFM1* transcript levels remained unchanged regardless

of transformation status or material uptake (Fig. S5B,C), indicating that although ER stress may

be present, it is not a key driver of the cellular behavior described here.

NoTf acceptor cells grown with cytokines up to 16 weeks post-TNT transfer and sorting (Fig. S6A), displayed anchorage-independent growth at earlier time points than other types of sorted cells, and with a number of clones obtained comparable to that of Tf cells (Fig. 5F; Fig. S6B). Potential cross-contamination between Tf and NoTf sorted cell samples was ruled out based on the specific fluorescence of derived soft agar clones (Fig. S6C). All these features are hallmarks of a preneoplastic transformation, observed exclusively in cells that had established TNT contacts with transformed cells prior to sorting, enabling the transfer of transforming information. BMPR1b, trafficked *via* TNTs from Tf to NoTf cells, thus emerges as a key mediator of this process.

Hence, these findings unravel the key role of the BMP pathway in driving the preneoplastic switch in NoTf cells receiving material from Tf cells through TNT-mediated transfer. TNT communication would prime non-transformed cells to respond to a tumor-promoting microenvironment, ultimately contributing to the propagation of transformation (Fig. 5G).

Discussion

Our findings reveal that TNT-mediated transfer of BMPR1b from transformed to non-transformed cells activates BMP signaling in acceptor cells, sensitizing them to a tumor-promoting microenvironment and initiating early features of transformation. The TNT-mediated transfer of mitochondria from stromal to cancer (stem) cells has been well documented (*25*), and recently, TNTs were shown to enhance the metastatic capacity of cancer cells in primary tumors, receiving mitochondria from neurons (*10*).

In contrast, the delivery of material from transformed cells to non-transformed neighbors, remains unexplored at very early stages of transformation. Our study is the first to investigate the functional impact of TNT-mediated material exchange between cells of identical phenotype (i.e. epithelial cells), on the predisposition of non-transformed cells to adopt a preneoplastic status. Although donor-to-acceptor material transfer through secretion of soluble compounds and extracellular vesicles occurred, it remained low to negligible, indicating that the processes mainly relied on physical TNT-mediated communications. This directional transfer from transformed to non-transformed cells leads within days to the overexpression of tumor-related genes and of BMPR1b itself, evocative of a possible feedback loop that maintains transforming conditions (Fig. 5G). Non-transformed cells also undergo profound changes in their proteomic landscape, linked to transport processes, and within weeks acquire phenotypic traits of preneoplasia, such as anchorage-independent growth, sustained BMPR1b overamplification and related mounting of a non-canonical signaling cascade. The earliest events, within one week, comprised the overexpression of *HOXB6*, *CCND2* and *EREG* by non-transformed acceptor cells, after TNT contacts with transformed (donor) cells. Interestingly, upregulation of these genes upon transformation has already been linked to other cancers: (i) *HOXB6*, encoding a homeobox domain-containing transcription factor involved in development, is upregulated in acute myeloid leukemia and osteosarcoma (*21*, *26*), with a possible role in cancer initiation (*27*); (ii) *CCND2*, encoding cyclin D2, involved in DNA replication, cell division and cell cycle regulation, was dysregulated in gastric, thyroid and colorectal cancers (*28*), and its overexpression in breast cancer promotes MAPK-mediated cell proliferation and tumor growth (*22*, *29*); (iii) upregulation of *EREG,* encoding the secreted protein ligand of the EGF receptor (EGFR) epiregulin, is linked to breast cancer progression and resistance to tamoxifen by EGFR activation (*23*, *30*).

Following TNT material transfer, including BMPR1b, *BMPR1B* becomes overexpressed in acceptor cells within a week. The receptor molecule was detected within TNTs, overexpressed in transformed cells, and structurally contributing to TNT elongation. As BMPR are known to be located in lipid rafts, BMPR1b could induce TNT formation *via* its interaction with these microdomains (*31*), from which TNTs might arise (*32*, *33*). These cholesterol-rich microdomains provide high curvature and elasticity necessary for membrane deformation linked to TNT formation. Cholesterol levels together with lipid rafts increase upon transformation, as evidenced in prostate and breast cancer cell lines (*34*), further substantiated by our proteomic data with acceptor cells displaying enhanced expression of EST1, a carboxylesterase hydrolyzing cholesterol esters into free cholesterol. Additionally, BMP2 was reported to promote actin polymerization via PI3Kα activation (*35*). We herein observed that *LCP1* was upregulated upon transformation, concomitantly to its protein Plastin-2, in acceptor cells following TNT transfer. Plastin-2 is a key actin-bundling protein involved in the formation of cellular protrusions (*36*), overexpressed in epithelial-derived cancers including breast (*37*). Its overexpression correlated with cancer cell migration, invasion, metastasis (*38*) and chemoresistance (*39*). Plastin-2 activation depends on serine 5 phosphorylation *via* ERK/MAPK, PI3K, and PKA/PKC pathways, enabling its recruitment to actin filaments and promoting bundling and protrusion elongation, as shown in breast carcinoma (*40*). This agrees with findings from the Zurzolo's group, showing that TNT induction by Eps8 involves its actin-bundling activity (*41*). Altogether, this might explain the higher number of TNTs observed upon transformation.

At the functional level, the analysis of acceptor cells sorted post-TNT transfer, revealed their gradual transformation over weeks, faster and more extensively than non-acceptor cells, concomitant to a progressive but sustained activation of the BMP pathway. This induced an initial burst of p38 downstream response, followed by the prolonged activation of the canonical SMAD cascade, only observed when TNT contacts existed. These recipient cells then exhibited preneoplastic traits. Experiments on TNT-forming capacities of mammary cell models, where transformation of either normal primary (*17*) or MCF10A cells (*42*) is « switched on » by the H-Ras oncoprotein, showed fewer and shorter TNTs connecting cells. *HRAS* is overexpressed in invasive breast tumors, but the prevalence of *HRAS* mutations in luminal types is negligible (*1*, *43*). These mammary models do not faithfully recapitulate the earliest stages of non-invasive luminal breast tumors triggered, for example, by exposure to carcinogenic cues such as environmental pollutants, endocrine disruptors or alcohol (*3*, *44–46*).

Here, we unveil for the first time that TNTs transfer a signaling receptor, BMPR1b. This transfer

sensitizes acceptor cells to BMP2 signals of the microenvironment, rapidly triggering a cascade

of molecular events that culminate in phenotypic changes, with anchorage-independent growth

of previously non-transformed cells (Fig. 5G). Combined with the recent demonstration that

TNTs can transfer material *in vivo* (*47*), our work provides the foundation for a new concept:

the propagation, through TNTs, of a preneoplastic state within a population of initially non-

transformed mammary cells. This process could precede the development of luminal-type

tumors, and identify BMPR1b as a major early contributor of the BMP-dependent preneoplastic

switch of acceptor cells.

Data availability statement. The mass spectrometry proteomics data have been deposited to
the ProteomeXchange Consortium via the PRIDE (https://www.ebi.ac.uk/pride/) (*48, 49*)
partner repository, with the dataset identifier PXD060023. Transcriptomic data have been
deposited on the Gene Expression Omnibus repository, GSE18673411.

Acknowledgements. We thank P. Battiston-Montagne, M. Gautier and C. Vanbelle (CRCL,
platform PIC), and D. Ressnikoff (SFR Santé Lyon Est, CIQLE). We thank the "AniRA
lentivectors production facility" from the CELPHEDIA Infrastructure and SFR Biosciences,
UAR3444/CNRS, US8/Inserm, ENS de Lyon, UCBL, especially G. Froment, D. Nègre and C.
Costa. We thank J. Brocard, head of the LYMIC-PLATIM UAR 3444 / US8 microscopy
platform (ENS Lyon, Univ Lyon 1), and E. Chatre. We thank B. Manship for english
proofreading, P. Mehlen, P. Juin, R. Tomasini, C. Vérollet and B. Bartosch for inspiring
discussions. Some illustrations were created in https://BioRender.com.
This work was supported by Agence Nationale de la Recherche (VMS, ANR-10-LABX-0061,
ANR-CESA-018-04 and Convergence PLAsCAN ANR-17-CONV-0002); Ligue Nationale
contre le Cancer (CCM, doctoral grant), Dept Ain, Rhône (VMS), Saône et Loire (EIP) ;
Fondation ARC (CCM, 4th year doctoral grant; VMS, SFI20111203500, PJA20171206331) ;
the ERiCAN program of Fondation MSD-Avenir (VMS/EIP, DS-2018-0015) and Associations
[Déchaîne Ton Cœur (VMS); Ruban Rose Prix Avenir 2021 (VMS) ; Comité féminin pour le
dépistage du cancer du sein 74 (VMS) ; Symatese Fondation (VMS), and Fondation pour la
Recherche Médicale (équipe FRM EQU202203014695 VMS).

Contributions: V.M.S. and E-I.P designed and supervised the study, conceived methodological
developments and acquired funding. V.M.S. provided administrative support. C.C.M., S.L. and
B.G. collaborated in study design, contributed to methodological developments and data
interpretations. K.G. performed RNA-seq and proteomic bioinformatic data analyses, and
statistical gene-expression analyses. C.C.M., T.T.N., A.B. and E-I.P. performed and interpreted
the experiments. C.D. designed and performed SEM and CLEM microscopy experiments. F.G.,
A.B. and C.H. performed proteomic analyses, and F.G. interpreted proteomic data. E.D.
provided primary breast samples, and assisted with their processing. C.C.M., V.M.S and E-I.P.
wrote the manuscript. All authors reviewed the manuscript and approved the final version for
publication.

Competing interests. The authors declare no competing interests.

References.

1. Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumors. *Nature* **490**, 61–70 (2012).

2. M. Zabala, N. A. Lobo, J. Antony, L. S. Heitink, G. S. Gulati, J. Lam, N. Parashurama, K. Sanchez, M. Adorno, S. S. Sikandar, A. H. Kuo, D. Qian, T. Kalisky, S. Sim, L. Li, F. M. Dirbas, G. Somlo, A. Newman, S. R. Quake, M. F. Clarke, LEFTY1 Is a Dual-SMAD Inhibitor that Promotes Mammary Progenitor Growth and Tumorigenesis. *Cell Stem Cell* **27**, 284-299.e8 (2020).

3. M. Chapellier, E. Bachelard-Cascales, X. Schmidt, F. Clément, I. Treilleux, E. Delay, A. Jammot, C. Ménétrier-Caux, G. Pochon, R. Besançon, T. Voeltzel, C. Caron de Fromentel, C. Caux, J.-Y. Blay, R. Iggo, V. Maguer-Satta, Disequilibrium of BMP2 levels in the breast stem cell niche launches epithelial transformation by overamplifying BMPR1B cell response. *Stem Cell Rep.* **4**, 239–254 (2015).

4. W.-J. Jiang, T.-H. Zhou, H.-J. Huang, L.-S. Li, H. Tan, R. Zhang, Q.-S. Wang, Y.-M. Feng, Breast Cancer Subtype–Specific Organotropism Is Dictated by FOXF2-Regulated Metastatic Dormancy and Recovery. *Cancer Res.* **85**, 644–659 (2025).

5. B. Guyot, F. Clément, Y. Drouet, X. Schmidt, S. Lefort, E. Delay, I. Treilleux, J.-P. Foy, S. Jeanpierre, E. Thomas, J. Kielbassa, L. Tonon, H. H. Zhu, P. Saintigny, W.-Q. Gao, A. de la Fouchardiere, F. Tirode, A. Viari, J.-Y. Blay, V. Maguer-Satta, An Early Neoplasia Index (ENI10), Based on Molecular Identity of CD10 Cells and Associated Stemness Biomarkers, is a Predictor of Patient Outcome in Many Cancers. *Cancer Res. Commun.* **3**, 1966–1980 (2023).

6. F. Ju, M. M. Atyah, N. Horstmann, S. Gul, R. Vago, C. J. Bruns, Y. Zhao, Q.-Z. Dong, N. Ren, Characteristics of the cancer stem cell niche and therapeutic strategies. *Stem Cell Res. Ther.* **13**, 233 (2022).

7. D. Bayik, J. D. Lathia, Cancer stem cell-immune cell crosstalk in tumour progression. *Nat. Rev. Cancer* **21**, 526–536 (2021).

8. N. Resnik, A. Erman, P. Veranič, M. E. Kreft, Triple labelling of actin filaments, intermediate filaments and microtubules for broad application in cell biology: uncovering the cytoskeletal composition in tunneling nanotubes. *Histochem. Cell Biol.* **152**, 311–317 (2019).

9. A. Rustom, R. Saffrich, I. Markovic, P. Walther, H.-H. Gerdes, Nanotubular highways for intercellular organelle transport. *Science* **303**, 1007–1010 (2004).

10. G. Hoover, S. Gilbert, O. Curley, C. Obellianne, M. T. Lin, W. Hixson, T. W. Pierce, J. F. Andrews, M. F. Alexeyev, Y. Ding, P. Bu, F. Behbod, D. Medina, J. T. Chang, G. Ayala, S. Grelet, Nerve-to-cancer transfer of mitochondria during cancer metastasis. *Nature* **644**, 252–262 (2025).

11. G. Pinto, C. Brou, C. Zurzolo, Tunneling Nanotubes: The Fuel of Tumor Progression? *Trends Cancer* **6**, 874–888 (2020).

12. M.-L. Vignais, A. Caicedo, J.-M. Brondello, C. Jorgensen, Cell Connections by Tunneling Nanotubes: Effects of Mitochondrial Trafficking on Target Cell Metabolism, Homeostasis, and Response to Therapy. *Stem Cells Int.* **2017**, 6917941 (2017).

13. L. Huang, J. Zhang, Z. Wu, L. Zhou, B. Yu, Y. Jing, D. Lin, J. Qu, Revealing the structure and organization of intercellular tunneling nanotubes (TNTs) by STORM imaging. *Nanoscale Adv.* **4**, 4258–4262 (2022).

14. I. Sáenz-de-Santa-María, J. M. Henderson, A. Pepe, C. Zurzolo, Identification and Characterization of Tunneling Nanotubes for Intercellular Trafficking. *Curr. Protoc.* **3**, e939 (2023).

15. D. Cordero Cervantes, C. Zurzolo, Peering into tunneling nanotubes-The path forward. *EMBO J.* **40**, e105789 (2021).

16. G. Pinto, I. Saenz-de-Santa-Maria, P. Chastagner, E. Perthame, C. Delmas, C. Toulas, E. Moyal-Jonathan-Cohen, C. Brou, C. Zurzolo, Patient-derived glioblastoma stem cells transfer mitochondria through tunneling nanotubes in tumor organoids. *Biochem. J.* **478**, 21–39 (2021).

17. B. Elenbaas, L. Spirio, F. Koerner, M. D. Fleming, D. B. Zimonjic, J. L. Donaher, N. C. Popescu, W. C. Hahn, R. A. Weinberg, Human breast cancer cells generated by oncogenic transformation of primary mammary epithelial cells. *Genes Dev.* **15**, 50–65 (2001).

18. G. Sapkota, C. Alarcón, F. M. Spagnoli, A. H. Brivanlou, J. Massagué, Balancing BMP signaling through integrated inputs into the Smad1 linker. *Mol. Cell* **25**, 441–454 (2007).

19. M. Inui, A. Manfrin, A. Mamidi, G. Martello, L. Morsut, S. Soligo, E. Enzo, S. Moro, S. Polo, S. Dupont, M. Cordenonsi, S. Piccolo, USP15 is a deubiquitylating enzyme for receptor-activated SMADs. *Nat. Cell Biol.* **13**, 1368–1375 (2011).

20. H.-C. Wen, A. Avivar-Valderas, M. S. Sosa, N. Girnius, E. F. Farias, R. J. Davis, J. A. Aguirre-Ghiso, p38α Signaling Induces Anoikis and Lumen Formation During Mammary Morphogenesis. *Sci. Signal.* **4**, ra34 (2011).

21. N. A. Fischbach, S. Rozenfeld, W. Shen, S. Fong, D. Chrobak, D. Ginzinger, S. C. Kogan, A. Radhakrishnan, M. M. Le Beau, C. Largman, H. J. Lawrence, HOXB6 overexpression in murine bone marrow immortalizes a myelomonocytic precursor in vitro and causes hematopoietic stem cell expansion and acute myeloid leukemia in vivo. *Blood* **105**, 1456–1466 (2005).

22. J. Ouyang, Z. Liu, X. Yuan, C. Long, X. Chen, Y. Wang, L. Liu, S. Liu, H. Liang, LncRNA PRNCR1 Promotes Breast Cancer Proliferation and Inhibits Apoptosis by Modulating microRNA-377/CCND2/MEK/MAPK Axis. *Arch. Med. Res.* **52**, 471–482 (2021).

23. M. He, Q. Jin, C. Chen, Y. Liu, X. Ye, Y. Jiang, F. Ji, H. Qian, D. Gan, S. Yue, W. Zhu, T. Chen, The miR-186-3p/EREG axis orchestrates tamoxifen resistance and aerobic glycolysis in breast cancer cells. *Oncogene* **38**, 5551–5565 (2019).

24. Y. Wang, J. Cui, X. Sun, Y. Zhang, Tunneling-nanotube development in astrocytes depends on p53 activation. *Cell Death Differ.* **18**, 732–742 (2011).

25. J. Nakhle, K. Khattar, T. Özkan, A. Boughlita, D. Abba Moussa, A. Darlix, F. Lorcy, V. Rigau, L. Bauchet, S. Gerbal-Chaloin, M. Daujat-Chavanieu, F. Bellvert, L. Turchi, T. Virolle, J.-P. Hugnot, N. Buisine, M. Galloni, V. Dardalhon, A.-M. Rodriguez, M.-L. Vignais, Mitochondria Transfer from Mesenchymal Stem Cells Confers Chemoresistance to Glioblastoma Stem Cells through Metabolic Rewiring. *Cancer Res. Commun.* **3**, 1041–1056 (2023).

26. H. Huang, D. Lu, K. Li, M. Zheng, X. Qin, X. Cui, Y. Chen, C. Chen, N. Huang, L. Zheng, J. Zhao, B. Zhu, Hsa_circ_0007031 promotes the proliferation and migration of osteosarcoma cells by sponging miR-196a-5p to regulate the HOXB6. *Biochem. Pharmacol.* **214**, 115667 (2023).

27. B. Z. Vider, A. Zimber, E. Chastre, C. Gespach, M. Halperin, P. Mashiah, A. Yaniv, A. Gazit, Deregulated expression of homeobox-containing genes, HOXB6, B8, C8, C9, and Cdx-1, in human colon cancer cell lines. *Biochem. Biophys. Res. Commun.* **272**, 513–518 (2000).

28. Z.-Y. Ding, R. Li, Q.-J. Zhang, Y. Wang, Y. Jiang, Q.-Y. Meng, Q.-L. Xi, G.-H. Wu, Prognostic role of cyclin D2/D3 in multiple human malignant neoplasms: A systematic review and meta-analysis. *Cancer Med.* **8**, 2717–2729 (2019).

29. H. Zhong, J. Yang, B. Zhang, X. Wang, L. Pei, L. Zhang, Z. Lin, Y. Wang, C. Wang, LncRNA GACAT3 predicts poor prognosis and promotes cell proliferation in breast cancer through regulation of miR-497/CCND2. *Cancer Biomark. Sect. Dis. Markers* **22**, 787–797 (2018).

30. M. C. Abba, R. Canzoneri, A. Gurruchaga, J. Lee, P. Tatineni, H. Kil, E. Lacunza, C. M. Aldaz, LINC00885 a Novel Oncogenic Long Non-Coding RNA Associated with Early Stage Breast Cancer Progression. *Int. J. Mol. Sci.* **21**, 7407 (2020).

31. T. Hirschhorn, M. Levi-Hofman, O. Danziger, N. I. Smorodinsky, M. Ehrlich, Differential molecular regulation of processing and membrane expression of Type-I BMP receptors: implications for signaling. *Cell. Mol. Life Sci. CMLS* **74**, 2645–2662 (2017).

32. J. Reindl, M. Shevtsov, G. Dollinger, S. Stangl, G. Multhoff, Membrane Hsp70-supported cell-to-cell connections via tunneling nanotubes revealed by live-cell STED nanoscopy. *Cell Stress Chaperones* **24**, 213–221 (2019).

33. V. Thayanithy, V. Babatunde, E. L. Dickson, P. Wong, S. Oh, X. Ke, A. Barlas, S. Fujisawa, Y. Romin, A. L. Moreira, R. J. Downey, C. J. Steer, S. Subramanian, K. Manova-Todorova, M. A. S. Moore, E. Lou, Tumor exosomes induce tunneling nanotubes in lipid raft-enriched regions of human mesothelioma cells. *Exp. Cell Res.* **323**, 178–188 (2014).

34. Y. C. Li, M. J. Park, S.-K. Ye, C.-W. Kim, Y.-N. Kim, Elevated levels of cholesterol-rich lipid rafts in cancer cells are correlated with apoptosis sensitivity induced by cholesterol-depleting agents. *Am. J. Pathol.* **168**, 1107–1118; quiz 1404–1405 (2006).

35. C. Gamell, N. Osses, R. Bartrons, T. Rückle, M. Camps, J. L. Rosa, F. Ventura, BMP2 induction of actin cytoskeleton reorganization and cell migration requires PI3-kinase and Cdc42 activity. *J. Cell Sci.* **121**, 3960–3970 (2008).

36. H. Ishida, K. V. Jensen, A. G. Woodman, M. E. Hyndman, H. J. Vogel, The Calcium-Dependent Switch Helix of L-Plastin Regulates Actin Bundling. *Sci. Rep.* **7**, 40662 (2017).

37. C. S. Lin, T. Park, Z. P. Chen, J. Leavitt, Human plastin genes. Comparative gene structure, chromosome location, and differential expression in normal and neoplastic cells. *J. Biol. Chem.* **268**, 2781–2792 (1993).

38. S. M. Riplinger, G. H. Wabnitz, H. Kirchgessner, B. Jahraus, F. Lasitschka, B. Schulte, G. van der Pluijm, G. van der Horst, G. J. Hämmerling, I. Nakchbandi, Y. Samstag, Metastasis of prostate cancer and melanoma cells in a preclinical in vivo mouse model is enhanced by L-plastin expression and phosphorylation. *Mol. Cancer* **13**, 10 (2014).

39. B. Janji, L. Vallar, Z. Al Tanoury, F. Bernardin, G. Vetter, E. Schaffner-Reckinger, G. Berchem, E. Friederich, S. Chouaib, The actin filament cross-linker L-plastin confers resistance to TNF-alpha in MCF-7 breast cancer cells in a phosphorylation-dependent manner. *J. Cell. Mol. Med.* **14**, 1264–1275 (2010).

40. R. A. C. Machado, D. Stojevski, S. De Landtsheer, P. Lucarelli, A. Baron, T. Sauter, E. Schaffner-Reckinger, L-plastin Ser5 phosphorylation is modulated by the PI3K/SGK pathway and promotes breast cancer cell invasiveness. *Cell Commun. Signal. CCS* **19**, 22 (2021).

41. E. Delage, D. C. Cervantes, E. Pénard, C. Schmitt, S. Syan, A. Disanza, G. Scita, C. Zurzolo, Differential identity of Filopodia and Tunneling Nanotubes revealed by the opposite functions of actin regulatory complexes. *Sci. Rep.* **6**, 39632 (2016).

42. P. J. Dawson, S. R. Wolman, L. Tait, G. H. Heppner, F. R. Miller, MCF10AT: a model for the evolution of cancer from proliferative breast disease. *Am. J. Pathol.* **148**, 313–319 (1996).

43. S.-J. Kim, S. Saeidi, N.-C. Cho, H.-B. Lee, W. Han, K. P. Kim, Y.-J. Surh, Cooperation between oncogenic NRF2 and wild-type HRAS promotes progression of breast cancer. *Cell. Signal.* **134**, 111920 (2025).

44. I. Romieu, C. Scoccianti, V. Chajès, J. de Batlle, C. Biessy, L. Dossus, L. Baglietto, F. Clavel-Chapelon, K. Overvad, A. Olsen, A. Tjønneland, R. Kaaks, A. Lukanova, H. Boeing, A. Trichopoulou, P. Lagiou, D. Trichopoulos, D. Palli, S. Sieri, R. Tumino, P. Vineis, S. Panico, H. B. A. Bueno-de-Mesquita, C. H. van Gils, P. H. Peeters, E. Lund, G. Skeie, E. Weiderpass, J. R. Quirós García, M.-D. Chirlaque, E. Ardanaz, M.-J. Sánchez, E. J. Duell, P. Amiano, S. Borgquist, E. Wirfält, G. Hallmans, I. Johansson, L. M. Nilsson, K.-T. Khaw, N. Wareham, T. J. Key, R. C. Travis, N. Murphy, P. A. Wark, P. Ferrari, E. Riboli, Alcohol intake and breast cancer in the European prospective investigation into cancer and nutrition. *Int. J. Cancer* **137**, 1921–1930 (2015).

45. C. Giampiccolo, B. Fervers, T. Coudon, D. Praud, A. Vigneron, L. Grassot, B. Mercoeur, E. Faure, P. Frenoy, M. Génard-Walton, F. Couvidat, G. Severi, F. R. Mancini, P. Roy, A. Amadou, Statistical approaches to analyse the combined effect of seven air pollutants

and breast cancer risk: a case–control study nested in the French E3N-Generations cohort. *BMC Med.* **24**, 112 (2026).

46. F. Clément, X. Xu, C. F. Donini, A. Clément, S. Omarjee, E. Delay, I. Treilleux, B. Fervers, M. Le Romancer, P. A. Cohen, V. Maguer-Satta, Long-term exposure to bisphenol A or benzo(a)pyrene alters the fate of human mammary epithelial stem cells in response to BMP2 and BMP4, by pre-activating BMP signaling. *Cell Death Differ.* **24**, 155–166 (2017).

47. O. Korenkova, S. Liu, I. Prlesi, A. Pepe, S. Albadri, F. Del Bene, C. Zurzolo, Tunneling nanotubes enable intercellular transfer in zebrafish embryos. *Dev. Cell*, S1534-5807(24)00635-X (2024).

48. Y. Perez-Riverol, Q.-W. Xu, R. Wang, J. Uszkoreit, J. Griss, A. Sanchez, F. Reisinger, A. Csordas, T. Ternent, N. Del-Toro, J. A. Dianes, M. Eisenacher, H. Hermjakob, J. A. Vizcaíno, PRIDE Inspector Toolsuite: Moving Toward a Universal Visualization Tool for Proteomics Data Standard Formats and Quality Assessment of ProteomeXchange Datasets. *Mol. Cell. Proteomics MCP* **15**, 305–317 (2016).

49. Y. Perez-Riverol, C. Bandla, D. J. Kundu, S. Kamatchinathan, J. Bai, S. Hewapathirana, N. S. John, A. Prakash, M. Walzer, S. Wang, J. A. Vizcaíno, The PRIDE database at 20 years: 2025 update. *Nucleic Acids Res.* **53**, D543–D553 (2025).

Figure Legends.

Fig. 1 │ Structural characterization of TNTs upon mammary cell transformation. A, Schematic of our MCF10A-derived cell lines *(5)*. B, Immunofluorescence images of actin (red) in TNTs, observed by Airyscan confocal microscopy with 40x oil objective. Nuclei, Hoechst labeling (blue). Scale bars, 20 μm. C, TNT number (> 8 μm) quantification *per* 100 cells (left, n = 9 biological replicates), and % of long TNTs (> 50 μm; right, n = 4 biological replicates). Ten Z-stacks of 10 sections each were taken *per* condition, and 100 cells minimum *per* experiment were counted. TNTs were counted and measured in each Z-stack. Error bars depict SD. P-values were calculated with one-way ANOVA with post-hoc Tukey test (ns, not significant; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$). D, Scanning electron microscopy images of TNTs of MCF10A-CT, MC26 and M1B26 cells. Black arrow, TNT bulge. Scale bars MCF10A-CT: 40 and 50 μm; MC26: 50 and 5 μm; M1B26: from left to right: 20, 3, 20, 3 μm. E, Transmission electron microscopy images of an open-ended TNT linking two M1B26 cells, with mitochondria (red asterisks). Scale bars, 5 μm (low mag), 500 nm (high mag). F, Immunofluorescence images of mitochondria (green) in MCF10A-CT and M1B26 cells. Actin, red; nuclei, blue. Scale bars, 20 μm. G, Immunofluorescence images of actin (red) in mammary acini grown from MC26 and M1B26 cells. Scale bars, 5 μm. Right, 3D-projection of a TNT linking two M1B26 cells, obtained with Zeiss ZENlite3.6 software. Actin, red; nuclei, blue. H, Images of TNTs linking M1B26 (GFP, green, and transmitted light) to normal human primary mammary fibroblasts (transmitted light). Nuclei, blue. Scale bars, 20 μm. I, Images of TNTs linking M1B26 (GFP, green) to normal human primary mammary cells. Actin, red; nuclei, blue. Asterisks: emerging from M1B26 (green); between both cell types (red). Scale bars: top, 20 μm, bottom, 10 μm.

Fig. 2 │ TNTs link transformed and non-transformed mammary cells, favoring a directional transfer of information. A, Fluorescence images of GFP-labeled-M1B26 (transformed - Tf; green) and mCherry-labeled MCF10A (non-transformed - NoTf; red) cells, after 48 h in coculture. Scale bars, 5 μm. B,C, Relative number of TNTs linking NoTf to Tf, or Tf to NoTf cells (B) and TNT ratio defined as $\frac{NoTf>Tf}{NoTf-NoTf}$ (black arrows *vs* red lines) or $\frac{Tf>NoTf}{Tf-Tf}$ (black arrows *vs* green lines) after 48h in coculture (C). Z stacks of 10 sections each were taken *per* condition and 100 cells minimum were counted. TNTs were counted and measured in each Z-stack. Error bars depict SD. Statistical significance was calculated using an unpaired Student's two-sided t test with Welch's correction (n = 5 biological replicates; *** $p < 0.001$). D,

Fluorescence images of material transfer (green dots) from Tf (green) to NoTf cells.

Membranes, red; nuclei, blue. Scale bars: 100 μm, in zoom 20 μm. E, Time-lapse snapshots of

green material trafficking within a TNT bulge (*), from a (green) Tf to NoTf cells after 48 h in

coculture. Membranes, red; nuclei, blue. Scale bar, 100 μm; images recorded for 75 min, with

one image every 18 min (see Supplementary Movie 1). F, Flow cytometry analysis of an 8-day

coculture of green Tf and red NoTf cells, with a schematic of NoTf accepting material from Tf

cells (top: NoTf accept green, red cell with green dots), and Tf cells accepting material from

NoTf cells (bottom: Tf accept red, green cell with red dots). G,H, Quantification of the

percentage of exchange (G; n = 8 biological replicates) and fluorescence intensity (H; n = 7

biological replicates), for indicated transfer. Error bars depict SD. Statistical significance was

calculated using an unpaired Student's two-sided t test with Welch's correction (*** $p < 0.001$).

I, Schematic of the experimental design of coculture experiments. J,K, Quantification of the

number of acceptor cells *per* 10,000 cells (J), and fluorescence intensity (K), in standard or

Transwell™ cocultures, for indicated transfer involving normal human primary mammary

(normal) and M1B26 cells (Tf). Error bars depict SD. Statistical significance was calculated

using an unpaired Student's two-sided t test with Welch's correction (n = 3 biological replicates;

ns, not significant; * $p < 0.05$; ** $p < 0.01$).

Fig. 3 │ TNT-dependent phenotypic changes of non-transformed cells acceptor of material from

transformed cells. A, Schematic of the experimental design: 4 cell populations, sorted by flow

cytometry (NoTf, Tf, NoTf in coculture without fluorescent transfer [non-transformed non-

acceptor cells - NoTf no accept], NoTf in coculture with fluorescent transfer [non-transformed

acceptor cells - NoTf accept]), underwent proteomic analyses and RT-qPCR. B,C, Volcano plot

(B) showing Log_2 enrichment of individual proteins in NoTf accept compared to NoTf no

accept cells from the same coculture (see schematic). P-value threshold was set at 0.05 and 0.8

for Log_2 fold change. C, To obtain a larger pool of hits, it was lowered to 0.5, generating a list

of 193 proteins upregulated in NoTf accept cells, used to generate the GOBP subset with

STRING (https://string-db.org). D,E, Box and whiskers plots of Log_2LFQ values (Protein

Levels) of EST1 (*CES1*), Plastin-2 (*LCP1*), Rab-6B (*RAB6B*), TRIQK (*TRIQK*) (D), and H-

Ras and K-Ras (E), for indicated sorted cell populations. Statistical significance was calculated

using one-way ANOVA with Dunnett's multiple comparison test (n = 5 biological replicates; *

$p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$). For clarity, comparisons with Tf cells

were omitted, and in (E), only significant comparison are shown. F,G, Immunofluorescence

images of actin (red) and tubulin (green) in (F) HMLE and HMLER cells, and (G) MCF10A,

MCF10.AT1 and MCF10.DCIS cells. Nuclei, blue. Scale bars = 20 μm. H,I, From left to right:

quantification of TNT number *per* cell, average length, and percentage of long TNTs in

indicated cell lines. Ten Z-stacks of 10 sections each were taken *per* condition and 100 cells

minimum *per* experiment were counted. TNTs were counted and measured in each Z-stack.

Statistical significance was calculated using an unpaired Student's two-sided t test with Welch's

correction (n = 8 and n = 5 biological replicates for HMEC-derived and MCF10A-derived cell

lines, respectively; ns, not significant; ** $p < 0.01$).

Fig. 4 │ Expression of actors of the BMP signaling pathway is enhanced in acceptor cells after

TNT transfer. A, Box and whiskers plots of Log_2LFQ values (Protein Levels) of ID1, Nup214,

Spartin, USP15, ATF2, MAPKAPK2 and Rac1, for indicated sorted cell populations. Statistical

significance was calculated using one-way ANOVA with Dunnett's multiple comparison test (n

= 5 biological replicates; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$). For clarity,

only comparison between NoTf and NoTf accept cells is shown. B, Immunofluorescence

images of actin (red) and BMPR1b (green) in indicated cells. Nuclei (blue). Scale bars, 20 μm.

C, Correlative light scanning electron microscopy (SEM) of BMPR1b, immunostained with

Alexa Fluor®488-Fluoronanogold™ (green) in M1B26 cells. From left to right: confocal,

scanning electron microscopy, zooms, correlative (overlay). Scale bars, 10 μm, unless

otherwise indicated. D, Immunofluorescence of actin (green), after transfection with siRNA

pools targeting *BMPR1b* (siBMPR1b) or *GPER* (siGPER). Non-targeting siRNA: siCTRL. E,

BMPR1B and *GPER* transcript and protein expression upon their silencing (top),with siCTRL

set at 1 (n = 4), and (bottom) TNT number (> 8 μm) *per* cell and % of long TNTs (> 50 μm)

upon indicated silencing, normalized against siCTRL. Error bars depict SD. Statistical

significance was calculated using an unpaired Student's two-sided t test with Welch's correction

(n = 4 biological replicates, *** $p < 0.001$). F, Schematic of experimental design: after sorting,

cell populations were either lysed and subjected to RT-qPCR for selected transcript analyses, or

grown for several weeks with or without BMP2/IL6, and immunoblot analyses performed at

end points. G, RT-qPCR measurement of BMPR1b mRNA in indicated cell populations, the

expression in NoTf cells (monocultures) set at 1. Statistical significance was calculated using

one-way ANOVA with post-hoc Tukey test (n = 5 biological replicates). For clarity, only

comparison between NoTf accept and NoTf cells is shown (**** $p < 0.0001$).

Fig. 5 | Non-transformed cells acceptor of information from transformed cells *via* TNTs

display preneoplastic features. A, Left, quantification of BMPR1b immunoblots performed on

NoTf accept cells grown in the presence of BMP2/IL6, for indicated weeks post sorting. Box

and whiskers plot of 6 biological replicates. Statistical significance was calculated using a one-

way ANOVA with non-parametric Kruskal-Wallis test, and for clarity, only p-values ≤ 0.05 are

indicated. Right, quantification of BMPR1b immunoblots performed on indicated cells, 14

weeks post sorting, with NoTf conditions set at 1 (n = 6 biological replicates). Statistical

significance was calculated using a one-way ANOVA with post-hoc Tukey test (*** $p < 0.001$).

B,C, Quantification of p-SMAD1/5/8 / SMAD1/5/8 (B) and p-p38/p38 (C) immunoblots

performed on NoTf accept cells grown in the presence of BMP2/IL6, for indicated weeks post

sorting. Box and whiskers plot of 6 biological replicates. Statistical significance was calculated

using a one-way ANOVA with non-parametric Kruskal-Wallis test, and for clarity, only p-values

≤ 0.05 are indicated. D, RNA-sequencing results of transcriptomic expression levels of *HOXB6,*

CCND2, EREG and *HRAS* in indicated cell lines (n = 3 biological replicates). Data deposited

on the Gene Expression Omnibus repository, GSE186734 *(5)*. Error bars depict SD. Statistical

significance was calculated using a one-way ANOVA with post-hoc Tukey test (* $p < 0.05$; **

$p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$). E, RT-qPCR measurements of HOXB6, CCND2,

EREG and HRAS mRNA in indicated cell populations, with the expression in NoTf cells set at

to 1. Error bars depict SD. Statistical significance was calculated using a one-way ANOVA with

post-hoc Tukey test (biological replicates: n = 8 for *HOXB6*, n = 5 for *CCND2*, n = 7 for *EREG*,

n = 8 for *HRAS*). For clarity, only comparisons between NoTf, NoTf Transwell and NoTf accept

cells are shown (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$). F, Quantification of

soft agar clones of indicated sorted cells, after 8, 10, 14 and 16 weeks in culture with or without

BMP2/IL6 (n = 3 biological replicates). Error bars represent SD. Statistical significance was

calculated using a one-way Welch and Brown-Forsythe ANOVA test (* $p < 0.05$; ** $p < 0.01$;

*** $p < 0.001$; **** $p < 0.0001$). For clarity, only p-values ≤ 0.05 are indicated, and

comparisons with Tf cells are omitted. G, Model depicting how TNTs might be involved in the

propagation of a BMP-dependent preneoplastic state. Fibrils and fibers depict extracellular

matrix components.

Cuella Martin *et al* Figure 3

1   **Supplementary materials & methods to :**
2
3
4
5   **Tunneling nanotubes propagate a BMP-dependent preneoplastic state.**
6
7

8   Cristina Cuella Martin[1,2,3], Thanh Thuy Nguyen * [1,2,3], Alexia Bertin * [1,2,3], Kevin
9   Geistlich[1,2,3,4], Boris Guyot[1,2,3], Sylvain Lefort[1,2,3], Emmanuel Delay[1,2,3,4], Charline
10   Dalverny[3,5], Alice Boissard[6], Cécile Henry[6], François Guillonneau[6,7,8], Eve-Isabelle Pécheur
11   §* [1,2,3] and Véronique Maguer-Satta §* [1,2,3,4]
12
13
1, CNRS UMR5286, Centre de Recherche en Cancérologie de Lyon, 69008 Lyon, France;
2, Inserm U1052, Centre de Recherche en Cancérologie de Lyon, 69008 Lyon, France;
3, Université Claude Bernard Lyon 1, Lyon, France;
4, Centre Léon Bérard, 69008 Lyon, France;
5, Centre Technologique des Microstructures (CTµ), 69622 Villeurbanne, France;
6, Prot'ICO proteomics facility, Institut de Cancérologie de l'Ouest, 49055 Angers, France,
7, Nantes Université, Inserm, CNRS, CRCI2NA, U1307, UMR6075, 44000 Nantes, France;
8, Equipe Labellisée Ligue Contre le Cancer, Paris, France.
22
23
* These authors contributed equally; § Co-corresponding authors

Materials & Methods.

Cell lines. MCF10A cells were purchased from ATCC (ATCC Batch#7635052). The MC26 cell

line was obtained upon MCF10A chronic exposure to BMP2 and IL6, while M1B26 came from

MCF10A first sorted on BMPR1b overexpression, then chronically exposed to BMP2 and IL6,

and derived from a soft agar colony (*1*). MCF10A, MCF10A.DCIS, MC26 and M1B26 were

cultured in phenol-red free Dulbecco's Modified Eagle Medium Ham's F12 (DMEM-F12,

Gibco), supplemented with 2% horse serum (Gibco), 10,000 U/ml - penicillin and 10,000 µg/ml

- streptomycin (Gibco) (P/S), 10 µg/ml insulin (NovoRapid CLB pharmacy), 0.5 µg/ml

hydrocortisone (Sigma), 100 ng/ml cholera toxin (Sigma), 20 ng/ml epithelial growth factor

(Sigma). MCF10A.AT1 and MCF10A.DCIS, obtained from ATCC, were cultured with the

same medium supplemented with 5% horse serum. MC26 and M1B26 culture medium was

supplemented with 5 ng/ml BMP2 (R&D) and 10 ng/ml IL6 (Peprotech). HMEC cells were

obtained from R. Weinberg's group (MIT, Massachussetts, USA). Cells were grown in 1:1

MEGM Complete Medium [Lonza/Clonetics, CC-3150] plus DMEM/F12 (1:1) with

supplements : 10 ng/ml hEGF, 0.5 µg/ml hydrocortisone, 10 µg/ml insulin, 10,000 U/ml -

penicillin and 10,000 µg/ml - streptomycin (Gibco). Cells were passed twice a week at 70% of

confluence. Cells washed twice with Phosphate Buffer Saline (PBS, Life Technologies) were

treated with 0.48 mM Versene (ThermoFischer, #15040066) for 20 min, or with trypsin 0.05%

for 10 min at 37°C. After collecting cells, neutralization was achieved with PBS/20% FBS.

Plated cells were incubated at 37°C under 5% atmospheric CO_2.

Fluorescent cell lines generation. Fluorescent cells were obtained by viral transduction of

parental ones with lentivirus produced in 293T cells by transfection of three plasmids encoding

respectively for the viral envelope G of VSV (vesicular stomatitis virus), the structural and

enzymatic proteins Gag and Pol of the lentivirus, and for a reporter gene mCherry or GFP (in

collaboration with AniRA lentivectors production facility from the CELPHEDIA Infrastructure

and SFR Biosciences, UAR3444/CNRS, US8/Inserm, ENS de Lyon/UCBL). Of note, these

viruses are defective, meaning that no viral protein or particle are produced by cells after

transduction.

Coculture of MCF10A-CT and M1B26 cells. Cells were cocultured at a 50% ratio for up to 8

days in MCF10A complete medium. Fresh medium was added every 3-4 days. For the

conditioned medium assay, the M1B26 green or MCF10A-CT green supernatant was collected,

centrifuged for 5 min at 800xg, and filtered on a 20 µm-pore filter. Monocultures of fluorescent

cells were grown for 8 days in this medium, and fresh one was added after 3-4 days. For
cocultures in Transwell, a Transwell system with pores of 0.4 μm was used (Merck Millipore,
#PHP01250). Four and ½ ml of MCF10A complete medium was added at the bottom, and 450
μl of MCF10A complete medium at the top of the insert. Cells were cocultured in the Transwell
system for 8 days, and fresh complete medium was added after 3-4 days.

Acini formation. The protocol for 3D-acini formation was derived from Vidi *et al* (*2*), using
the 3D-embedded culture process.

Gene silencing with siRNA. Silencing of BMPR1b and GPER was performed using small
interfering RNAs, purchased at Horizon Discovery (ON-TARGETplus siRNA, SMARTpool
siRNA). Cells were transfected through electroporation by a Neon Electroporator Device.

Western-Blotting. Cells were harvested, pelleted, and lysed in CHAPS buffer (40 mM Hepes
pH 7.4, 120 mM NaCl, 2 mM EDTA, 0.3% CHAPS) and protease/phosphatase inhibitors
(Roche Complete inhibitor cocktail #4693132001). Proteins were then assayed by
spectrophotometry, along with a BSA standard curve, using the Pierce™ Coomassie Protein
Assay Kit (Thermo Fisher Scientific). Twenty to 30 μg of proteins were used and mixed with
1X Laëmmli and H_2O, and samples boiled for 5 min at 95°C. The mixture was loaded onto 10%
acrylamide gels, and run in 25 mM Tris, 192 mM glycine, 0.1% SDS. After transfer onto PVDF
membranes previously activated in ethanol and transfer buffer (Biorad™ Transfer buffer, 20 %
ethanol) using the Transblot Turbo™ from BioRad. Membranes were then incubated for 1h
with TBST [Tris-buffered saline 1X (Euromedex) containing 0.1% Tween-20] with 5% fat-free
BSA for membrane saturation at room temperature. Primary antibodies were incubated at 4°C
overnight. Primary antibodies used were directed against BMPR1b (1:300 Abcam, ab78417 and
1:300 Santa Cruz, sc-515886), GAPDH (1:10,000 Cell Signaling Technology, #8884 and
1:50,000, Proteintech #600004-1), p-SMAD1/5/8 (1:500, Cell signalling, #13820), Smad1/5/8
(1:500, Santa Cruz, sc-6031-R), p-ERK (1:2,000, Cell Signalling Technology, #4370S), ERK
(1:1,500, Cell Signalling Technology #4695S), p-p38 (1:2,000, BD Bioscience #612288), p38
(1:2,000, BD Biosciences, #612168), GPER (1:500 Novus Biologicals, #NLS4290 or 1:500,
Invitrogen, #PA5-28647), EREG (1:400, abcam, #1041688-1), Cyclin-D2 (1:3,000,
Proteintech, #67048-i-IG), or HoxB6 (1:1,500, Santa Cruz, sc-166950), in 5% BSA. After
TBST buffer washes, membranes were incubated at room temperature for 1h with horseradish
peroxydase (HRP)-conjugated secondary antibodies goat anti-rabbit, goat anti-mouse or rabbit

anti-goat secondary antibodies (Jackson, #111-035-003, #115-035-003, #305-035-003) at

1:10,000 in TBST/3% BSA. After extensive washings with TBST buffer, revelation was

achieved through Clarity™ Western ECL Substrate (Biorad) and Chemidoc™ Imaging System

(Biorad). Protein bands were quantified using the ImageLab™ software.

Immunofluorescence. Cells were seeded on coverslips and grown at 60% of confluence. For

staining BMPR1b and actin staining, the following protocol was used : after washing with PBS,

fixation was performed for 20 min in 4% final paraformaldehyde (PFA) at room temperature

(RT). After PBS washing, permeabilization and saturation were performed for 45 min with

PBS/3% BSA/0.3% Triton X-100 at RT. After PBS washing, primary antibody against BMPR1b

(1:300 #sc-293428 Santa Cruz) and its corresponding isotype were incubated overnight at 4°C.

Coverslips were washed three times with PBS/3% BSA/0.1% Tween-20 for 10 min. Secondary

antibody goat anti-mouse Dylight 488 diluted 1:1,000 was incubated for 45 minutes at RT, with

1:50,000 of Hoechst 33342 (Invitrogen #H3570) and Phalloidin AlexaFluor-Plus 647

(Invitrogen, #A30107). Then, cells were washed twice with PBS/3% BSA/0.1% Tween-20 for

5 minutes and twice with PBS for 5 minutes. Coverslips were then mounted using the Dako

mounting medium (Agilent, ref S3023). For mitochondria staining, MitoTracker™ deep red or

green were used (Thermofischer, #M22426 or #M7514, respectively). Microscopic

observations were performed using the x40 oil immersion objective of a confocal Zeiss

LSM980 microscope, with Airyscan mode. This was done at the LYMIC-PLATIM, Imaging

and Microscopy Core Facility of SFR Biosciences (UMS3444), Lyon, France. Images were

taken in 16-bit, and formatted via the Fiji software. For immunofluorescence staining of actin

(Phalloidin) and tubulin, the following protocol optimized by the Dr. C. Zurzolo's team was

used to identify and characterize TNTs. Subconfluent cells were fixed for 15min at RT, with a

4% PFA/0.05% glutaraldehyde mixture in PBS. The fixation buffer was removed, and and cells

incubated with 100 mM NH_4Cl for 10min at RT to quench the PFA-related autofluorescence.

After PBS washing, cells were saturated with 2% BSA for 20 min at RT. The primary antibody

against tubulin was diluted at 1:500 (Abcam #ab6046) in 2% BSA, and incubated for 1h at RT,

or OVN at 4°C. After PBS washing, cells were incubated with a secondary goat anti-rabbit

antibody-488nm, diluted at 1:1,000 in 2% BSA, and incubated for 40 min in the dark at RT.

After PBS washing, cells were finally incubated with phalloidin AlexaFluor Plus-647

(Invitrogen, #A30107) diluted at 1:1,000, and with Hoechst 33342 (Invitrogen, #H3570),

diluted at 1:50,000 in 1% BSA, for 30 min in the dark. After PBS washing, slides were mounted

using the Dako mounting medium (Agilent, S3023). Microscopic observations were performed

with the objective 40x oil of a confocal Zeiss LSM980 microscope, with Airyscan mode. This

was performed at the LYMIC-PLATIM, Imaging and Microscopy Core Facility of SFR

Biosciences (UMS3444), Lyon, France, and at the imaging quantitative center of Lyon Est

(CIQLE). Images were taken in 16-bit, and formatted *via* the Fiji software. TNTs were finally

manually counted with the Fiji software.

Live imaging. Live imaging was performed after 48h of coculture in µslide 8-well plate with

thin glass bottom (Ibidi, #80827), adequate for imaging. After PBS washing, cells were stained

with Cell Mask Deep Red (at 1:500, #C10046, Invitrogen) and Hoechst (Invitrogen, #H3570,

1:500), diluted in complete medium, for 10 min at 37ºC. After PBS washing, complete fresh

medium was added. Observations were performed using the Opera Phenix HCS microscope

with the 40x water immersion objective.

RT-qPCR. For RT-qPCR analyses, pelleted cells were resuspended with 300µl Trizol reagent

(Life Technologies™). Total RNA purification was achieved using direct-zol RNA MicroPrep

w/ Zymo-Spin IC Columns extraction kit (Ozyme). For sorted fluorescent cells after coculture,

RNA extraction was performed using the AllPrep DNA/RNA Micro Kit (Qiagen, #80284). C-

DNA was then obtained using PrimeScript™ RT Reagent Kit (Takara kit), and qPCR then

performed using the SyberGreen Premix Ex Taq™ (Takara kit) as the fluorescent DNA

intercalant, and amplicons read by a Light Cycle 480 (Roche, LC480). Results were normalized

to housekeeping genes : TATA-binding protein (*TBP*) and hypoxanthineguanine phosphoribosyl

transferase (*HPRT*) (see Table S4 for primers used). An additional normalization to control cells

(MCF10A or MCF10A-CT) was also performed.

Proteomic analyses. *Sample preparation for proteome analysis*. Proteome analyses were

performed at the Prot'ICO proteomics facility: FACS-sorted samples (FACS-Melody) were

quickly thawed and proteins equivalent to 66,000 cells were concomitantly extracted and

denatured while thiol groups were chemically reduced in 46 µl 5% SDS, containing 5 mM

TCEP, 10 mM chloroacetamide, heated in a thermostatic shaker set at 95°C under 500xg for 30

min. Sonication was performed twice 30s using an ultrasonic sonicator bath (ThermoFisher

Scientific). S-Trap Micro Spin Column was used according to the manufacturer's protocol

(Protifi, Farmingdale, NY, USA), to trap precipitated proteins under acidic and hydrophobic

conditions to remove detergents, and 1µg of porcine trypsin (Sciex) was used at 37°C overnight

to cleave peptide bonds at carbonyl-side of lysine and arginine to generate bottom-up LC MS-

compatible peptides. Resulting peptides were then cleared from potential insoluble material by

centrifugation. Peptides were then speed-vacuum dried and resuspended in 0.1% formic acid

(FA) before injection in LC-MS/MS with DIA-PASEF mode using a TIMS-TOF Pro 2 mass

spectrometer (Bruker, Bremen, Germany).

LC-MS/MS setup. Each sample was loaded and separated on a C18 reverse phase column

(Aurora Ultimate series 1.7µm particles size, 75µm inner diameter and 25cm length, IonOptics,

Fitzroy, Australia) using a NanoElute 2 LC system with Hystar 6.3.1.8 software (Bruker,

Bremen, Germany). Eluates were electrosprayed for a 60 min-long duration of the

hydrophobicity gradient ranging from 99% of solvent A containing 0.1% FA in milliQ-grade

H_2O, to 27% of solvent B containing 80% acetonitrile containing 0.1% FA in mQ-H_2O. The

mass spectrometer acquired data throughout the elution process and operated in data-

independent analysis (DIA) with PASEF-enabled method, using the TIMS-Control software

v.5.1.8 (Bruker). Samples were injected in batch replicates in order to circumvent possible

interferences of technical biases. The m/z acquisition range was 250–1201 with an ion mobility

range of 0.7–1.25 1/K0 [V s/cm^2], which corresponded to an estimated cycle time of one second.

DIA-PASEF windows, and collision energy were also left to default with a base of 0.85 1/K0

[V s/cm^2] set at 20 eV and 1.30 1/K0 [V s/cm^2] set at 59 eV. TOF mass calibration and TIMS

ion mobility 1/K0 were performed linearly using three reference ions at 622, 922, and 1,222

m/z (Agilent Technology, Santa Clara, CA, USA)

LC-MS/MS raw data analysis. The raw data were extracted, normalized and analyzed using

Spectronaut® 19.0.24 (Biognosys A.G. Schlieren, Switzerland) in DirectDIA+ mode, which

modelized elution behavior, mobility and MS/MS events based on the "Uniprot/SwissprotKB

model organism_9606_2024_04_03" FASTA sequence database of 45,514 human proteins and

381 entries of common contaminant proteins. Protein identification false discovery rate (FDR)

was restricted to 1% maximum, and inter-injection data normalization. The enzyme's cleavage

specificity was that of trypsin. The precursor and fragment mass tolerances were set to default.

Acetyl (protein N-term) and oxidation of methionines were set as variable modifications, while

thiol groups from cysteines were considered completely alkylated by carbamidomethylation. A

minimum of two ratios of peptides was required for relative quantification between groups.

Protein quantification analysis was performed using Label-Free Quantification (LFQ)

intensities. Spectronaut also linked Gene Ontology terms to identified protein using the EBI

GO database, and its statistical tools were used to visualize data, assess reproducibility and

quality to allow quantitative dataset comparisons. The mass spectrometry proteomics data have

been deposited to the ProteomeXchange Consortium via the PRIDE (*3, 4*) partner repository with the dataset identifier PXD060023.

Volcano plots were generated with the package EnhancedVolcano (https://bioconductor.org/packages/release/bioc/html/EnhancedVolcano.html), with a p-value threshold set at 0.05 and a LogFC threshold set at either 0.8 or 0.95. Pathway analysis were performed using STRING (https://string-db.org/) with the Biological Process category from the Gene Ontology collection. For the search of BMP and P38 pathways targets, a combination of several pathways from different collections have been compiled (GOBP_POSITIVE_REGULATION_OF_BMP_SIGNALING_PATHWAY ; GOBP_REGULATION_OF_BMP_SIGNALING_PATHWAY ; GOBP_RESPONSE_TO_BMP ; PID_BMP_PATHWAY ; BIOCARTA_P38MAPK_PATHWAY ; GOBP_P38MAPK_CASCADE ; GOBP_REGULATION_OF_P38MAPK_CASCADE ; REACTOME_P38MAPK_EVENTS ; WP_P38_MAPK_SIGNALING).

Flow cytometry and cell sorting. After cell detachment as previously described, cells were resuspended in PBS/1% FBS. Sorting of cocultured cells was performed using the BD FACSMelody™ Cell Sorter, at the CRCL Flow cytometry platform, SFR Lyon-EST (CYLE). Results were analyzed with the FlowJo™ software.

Soft agar colony formation assay. Each well of a 6-well plate was coated with agarose 1.5% mixed with 2X complete medium. After polymerisation at RT, a top layer was added, containing agarose 0.9%, mixed with 2X complete medium containing 13,000 cells. PBS was added between wells, so that the agar would not dry out. Clone counting was performed manually after 3-4 weeks.

Electron microscopies. *Transmission EM.* Cell samples in Petri dishes were washed 3 times 15min in sodium cacodylate buffer 0.2M pH 7.3, and fixed for 1h in glutaraldehyde 2% in the same buffer at 0.1M. After washing in buffer, samples were prepared for TEM observations after Epoxy resin embedding and microtome ultrathin sectioning at the Center for Microstructure Technology (CTµ, Univ Lyon 1). Observations were performed with a Jeol 1400 Flash microscope operated at 200 kV, and equipped with a Gatan Rio 16 digital camera.

Correlative light scanning electron microscopy (CLEM). CLEM was performed in collaboration with the Center for Microstructure Technology (CTµ, Univ Lyon 1). Cells were

plated on standard square coverslips ITO (22x22mm), and fixed with PFA 4% 20 min. After

three washes with phosphate buffer 0.2M pH 7.2 for 20min, saturation was achieved using

phosphate buffer with 3% BSA and 0.3% Triton X-100 for 45min. Incubation with primary

antibodies and washings were performed as mentioned in the section "Immunofluorescence".

AlexaFluor-488 Fluoronanogold (FluoroNanogold™, Nanoprobes, #7201) was diluted at 1:100

in phosphate buffer with 1% BSA and 0.2% Tween-20, and incubated for 1h at RT. After

washings, coverslips were mounted on a "Shuttle and Find" Zeiss device, and the observation

was done with a Zeiss confocal LSM800 microscope. After confocal observation, coverslips

were washed three times with H_2O. Then silver enhancement was performed, using the HQ

silver enhancement kit (Nanoprobes, #2012) for 5 minutes. After H_2O washings, cells were

incubated with 1% final OsO4 in PBS 0.1M pH 7.2, for 45min. After H_2O washings, cells were

incubated overnight with 20% acetone in H_2O. The following day, 100% acetone was added for

1h, followed by critical point drying for 2h. SEM imaging was then performed at the Center for

Microstructure Technology (CTμ, Univ Lyon 1), using a scanning electron microscope Zeiss

Merlin Compact VP.

Statistical methods. Normality test was performed using the Shapiro-Wilk test. Mean

comparisons of two samples were performed with an unpaired two-sided Student's t-test with

Welch's correction. One-way Anova with post-hoc Tukey test, or with Dunnett's correction test,

or with non-parametric Kruskal-Wallis test, or a one-way Welch and Brown-Forsythe ANOVA

were used, when multiple comparisons were performed. Statistical analyses were performed

using GraphPad Prism software (La Jolla, CA). Significant p values are indicated by asterisks

in the figures (* $p<0.05$; ** $p<0.01$; *** $p<0.001$, **** $p<0.0001$).

References to Supplementary Materials & Methods, Figures and Tables.

1. B. Guyot, F. Clément, Y. Drouet, X. Schmidt, S. Lefort, E. Delay, I. Treilleux, J.-P. Foy, S. Jeanpierre, E. Thomas, J. Kielbassa, L. Tonon, H. H. Zhu, P. Saintigny, W.-Q. Gao, A. de la Fouchardiere, F. Tirode, A. Viari, J.-Y. Blay, V. Maguer-Satta, An Early Neoplasia Index (ENI10), Based on Molecular Identity of CD10 Cells and Associated Stemness Biomarkers, is a Predictor of Patient Outcome in Many Cancers. *Cancer Res. Commun.* **3**, 1966–1980 (2023).

2. P.-A. Vidi, M. J. Bissell, S. A. Lelièvre, Three-dimensional culture of human breast epithelial cells: the how and the why. *Methods Mol. Biol.* **945**, 193–219 (2013).

3. Y. Perez-Riverol, Q.-W. Xu, R. Wang, J. Uszkoreit, J. Griss, A. Sanchez, F. Reisinger, A. Csordas, T. Ternent, N. Del-Toro, J. A. Dianes, M. Eisenacher, H. Hermjakob, J. A. Vizcaíno, PRIDE Inspector Toolsuite: Moving Toward a Universal Visualization Tool for Proteomics Data Standard Formats and Quality Assessment of ProteomeXchange Datasets. *Mol. Cell. Proteomics MCP* **15**, 305–317 (2016).

4. Y. Perez-Riverol, C. Bandla, D. J. Kundu, S. Kamatchinathan, J. Bai, S. Hewapathirana, N. S. John, A. Prakash, M. Walzer, S. Wang, J. A. Vizcaíno, The PRIDE database at 20 years: 2025 update. *Nucleic Acids Res.* **53**, D543–D553 (2025).

Supplementary figures and tables to :

Tunneling nanotubes propagate a BMP-dependent preneoplastic state.

Cristina Cuella Martin[1,2,3], Thanh Thuy Nguyen * [1,2,3], Alexia Bertin * [1,2,3], Kevin Geistlich[1,2,3,4], Boris Guyot[1,2,3], Sylvain Lefort[1,2,3], Emmanuel Delay[1,2,3,4], Charline Dalverny[3,5], Alice Boissard[6], Cécile Henry[6], François Guillonneau[6,7,8], Eve-Isabelle Pécheur [§*][1,2,3] and Véronique Maguer-Satta [§*][1,2,3,4]

1. CNRS UMR5286, Centre de Recherche en Cancérologie de Lyon, 69008 Lyon, France;
2. Inserm U1052, Centre de Recherche en Cancérologie de Lyon, 69008 Lyon, France;
3. Université Claude Bernard Lyon 1, Lyon, France;
4. Centre Léon Bérard, 69008 Lyon, France;
5. Centre Technologique des Microstructures, 69622 Villeurbanne, France;
6. Prot'ICO proteomics facility, Institut de Cancérologie de l'Ouest, 49055 Angers, France,
7. Nantes Université, Inserm, CNRS, CRCI2NA, U1307, UMR6075, 44000 Nantes, France;
8. Equipe Labellisée Ligue Contre le Cancer, Paris, France.

* These authors contributed equally; [§] Co-corresponding authors

Cuella Martin *et al*

Figure S1

Figure S1. A, Representative immunofluorescence images of actin (red) and tubulin (green) in MCF10A-CT and M1B26 cells, plated at sub-confluent cell densities (20, 40 and 60K, respectively 20,000;40,000 or 60,000 cells/cm^2), and observed by Airyscan confocal microscopy. Nuclei, Hoechst labeling (blue). Scale bars, 20 µm. B,C,D, Quantification of TNT average length (B, n = 4 biological replicates), TNT number *per* 100 cells (C, n = 9 biological replicates), and % of long TNTs (D; > 50 µm; n = 4 biological replicates) for indicated cell conditions (MCF10A-CT+M1B26 CM : MCF10A-CT cells incubated for 48h with conditioned medium of M1B26 cells). Ten Z-stacks of 10 sections each were taken *per* condition, and 100 cells minimum *per* experiment were counted. TNTs were counted and measured in each   Z-stack. Error bars depict SD. Statistical significance was calculated using a one-way ANOVA with post-hoc Tukey test (B) or an unpaired two-sided Student's t test with Welch's correction (C,D). Ns, not significant. E, TEM images of a close-ended TNT linking two M1B26 cells via a Gap junction. Scale bars, 2 µm. F, Immunofluorescence images of actin (red) in acini grown from indicated cell lines. See Methods for acini formation. Nuclei, blue. Scale bars, 50 µm. Bottom right, acini area (in µm^2). Statistiscal significance was calculated using a one-way ANOVA with post-hoc Tukey test (only *** p < 0.001 is indicated). G, Immunofluorescence images of actin (green) and tubulin (red) in indicated cell lines. Nuclei, blue. Scale bars, 20 µm. H, Immunofluorescence images of mitochondria (green, Mitotracker™), and actin (grey) in indicated cell lines. Nuclei, blue. Scale bars, 20 µm. All fluorescence observations were performed with a confocal Airyscan Zeiss microscope.

Figure S2

Figure S2. A, Time-lapse snapshots of green material trafficking within a TNT bulge (*), from GFP (green) M1B26 to MCF10A-CT cells after 48h in coculture. Nuclei, turquoise blue. Scale bar, 100 µm. Images were recorded for 75 min, with one image every 18 min. For movie source data, see Movie S1. B, Flow cytometry analysis of an 8-day coculture of red M1B26 and green MCF10A-CT cells. C,D, Quantification of the percentage of exchange (C) and fluorescence intensity (D), of the indicated transfer. Error bars depict SD. Statistical significance was calculated using an unpaired Student's two-sided t test with Welch's correction (n = 7 biological replicates; *** p < 0.001; **** p < 0.0001). E,F,G, Flow cytometry analysis of 2 (E), 5 (F) or 8 (G) days of coculture of green M1B26 and red MCF10A-CT cells. H,I, Percentage of material exchange (H) and fluorescence intensity (I), of the indicated transfer after 2, 5 or 8 days of coculture. Error bars are SD. Statistical significance was calculated using a one-way ANOVA with post-hoc Tukey test (n = 6 biological replicates; * p < 0.05; ** p < 0.01). For clarity, only statistical significance is shown. J, Quantification of the number of acceptor cells *per* 10,000 cells, in standard (10 biological replicates) or Transwell cocultures (5 biological replicates), or in conditioned medium (5 biological replicates), for the indicated transfer involving MCF10A-CT (NoTf) and M1B26 (Tf) cells. Error bars are SD. Statistical significance was calculated using an unpaired Student's two-sided t test with Welch's correction (ns, not significant; ** p < 0.01). K,L, Quantification of the number of acceptor cells *per* 10,000 cells (K), and fluorescence intensity (L), in cocultures or with conditioned medium, for the indicated transfer involving normal human primary mammary (normal) and M1B26 cells (Tf). Error bars are SD. Statistical significance was calculated using an unpaired Student's two-sided t test with Welch's correction (n = 3 biological replicates; ns, not significant; * p < 0.05; ** p < 0.01).

Figure S3

Figure S3. A, Volcano plot showing Log_2 enrichment of individual proteins in non-transformed acceptor cells (NoTf accept) in direct coculture, compared to non transformed (NoTf) cells in monoculture. The threshold for the P-value was set at 0.05 and 0.8 for Log2 fold change. The Log_2 fold change threshold was lowered to 0.5 in order to obtain a larger pool of hits, with a minimum of differences between groups. Proteins with a magenta frame are also overexpressed in NoTf accept cells compared to non transformed non acceptor cells (NoTf no accept), in cocultures (see Fig. 3C). B, From a list of 193 proteins upregulated in NoTf accept cells (in coculture) compared to NoTf cells (in monoculture), STRING (https://string-db.org) was used to generate the gene ontology analysis with the biological process subset. C, Box and whiskers plots of Log_2LFQ values (Protein Levels) of MEST and MASU1, for indicated sorted cell populations. Statistical significance was calculated using one-way ANOVA with Dunnett's multiple comparison test (n = 5 biological replicates; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$). For clarity, only significant comparisons are shown. D, RNA-sequencing results showing the transcriptomic expression levels of *LCP1, RAB6B, CES1, TRIQK, MEST and MALSU1* in MCF10A-CT (NoTf) and M1B26 (Tf) (n = 3 biological replicates). Data deposited on the Gene Expression Omnibus repository, GSE186734 *(1)*. Statistical significance was calculated using an unpaired two-sided Student's t test with Welch's correction (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$). E,F, Volcano plots showing Log_2 enrichment of individual proteins in NoTf accept in Transwell coculture compared to NoTf cells in monoculture (E), and in NoTf accept cells in direct coculture compared to NoTf accept cells in Transwell coculture (F). Proteins with a magenta frame are also overexpressed in NoTf accept cells compared to NoTf no accept cells, in cocultures (see Fig. 3B), and the protein with a green frame is found overexpressed only in NoTf accept cells in direct coculture compared to NoTf accept cells in Transwell coculture. The threshold for the P-value was set at 0.05 and 0.8 for Log2 fold change. The Log_2 fold change threshold was lowered to 0.5 in order to obtain a larger pool of hits, with a minimum of differences between groups.

Figure S4

Figure S4. A, From left to right : Immunoblots of indicated proteins, in MCF10A-CT, MC26 and M1B26 cells. B, From left to right : quantification of immunoblots shown in A. Error bars are SD. Statistical significance was calculated using a one-way ANOVA with post-hoc Tukey test (biological replicates : n = 3 for BMPR1b, n = 9 for p-SMAD1/5/8 / SMAD1/5/8, n = 5 for p-p38/p38 and p-ERK/ERK ; ns, not significant, * p < 0.05 ; ** p < 0.01). Data are mean ± SD. C, Quantification of BMPR1b immunoblots, from lysates of cells grown for indicated time post flow cytometry sorting (see schematic Fig. 4G). Violin plots of 3 to 6 biological replicates, with NoTf and NoTf 2-6, MCF10A-CT cells grown in the absence or presence of BMP2/IL6, respectively; NoTf accept and NoTf accept 2-6, MCF10A cells grown in coculture with M1B26 cells, and cultured in the absence or presence of BMP2/IL6 post-sorting, respectively; Tf, M1B26 cells. NoTf conditions are set at 1. Statistical significance was calculated using a two-way ANOVA with Tukey's multiple comparison test. For clarity, only significant comparisons are shown. D, Quantification of BMPR1b immunoblots performed on indicated cells, 14 weeks post sorting, with NoTf conditions set at 1. Violin plot of 5 biological replicates. Statistical significance was calculated using a one-way ANOVA with post-hoc Tukey test, with only significant comparisons shown (* p < 0.05; ** p < 0.01; *** p < 0.001). Legend as in C, with : Transwell, coculture in Transwell; CM, NoTf (MCF10A-CT) cells grown in conditioned medium of M1B26 cells, sorted and grown in the absence or presence of BMP2/IL6. E,F, Quantification of p-SMAD1/5/8/SMAD1/5/8 (E) and p-p38/p38 (F) immunoblots, on same samples as in B.

A

Figure S5. A, Quantification of immunoblots of HoxB6, Cyclin D2 and (pro)-epiregulin in MCF10A-CT, MC26 and M1B26 cells, with normalization against MCF10A-CT levels. Data are mean ± SEM (Biological replicates : n = 8 for HoxB6; n = 11 for cyclin D2; n = 4 for (pro)epiregulin). Statistical significance was calculated using a one-way ANOVA with post-hoc Tukey test (ns, not significant; * p < 0.05). B, RNA-sequencing results showing the transcriptomic expression levels of UFM1 in indicated cell lines (n = 3 biological replicates). Data deposited on the Gene Expression Omnibus repository, GSE186734 *(1)*. Error bars represent SD. Statistical significance was calculated using a one-way ANOVA with post-hoc Tukey test (ns, not significant). C, RT-qPCR measurements of UFM1 mRNA in indicated cell populations, with the expression in NoTf cells set at to 1. Error bars represent SD. Statistical significance was calculated using a one-way ANOVA with post-hoc Tukey test (n = 5 biological replicates; ns, not significant).

Figure S6

Figure S6. A, Schematic of long-term soft agar colony formation assays. B, Representative bright-field images of soft agar clones after indicated weeks in the absence or presence of BMP2/IL6. Pictures with a dark frame are representative of big or small clones observed for the indicated cell condition. Scale bar represents 200 μm. C, Representative images of soft agar clones obtained from indicated cells and conditions, and observed by epifluorescence microscopy. After sorting by flow cytometry, cells were grown in the absence or presence of BMP2/IL6 for several weeks (see Fig. 5B), and at given time points, collected to perform soft agar colony formation assay (see panel A). Clones shown here were formed from cells grown for 16 weeks post sorting. From left to right : bright field, mCherry, GFP. Scale bars are 50 μm.

Table S1. The ten most differentially expressed genes (from proteomic analyses) between NoTf accept and NoTf no accept cells, in direct coculture conditions, showing Log$_2$ enrichment of individual proteins (LogFC) and corresponding p-value (see legend to Fig. S3A for more details).

UP				DOWN		
Protein	LogFC	P value		Protein	LogFC	P value
CES1	0.889122589477469	0.0141628941286904		ABHD14B	-1.00315026985703	0.000980307865197619
MEST	0.932759743282637	0.00188098627865132		PHPT1	-1.01510146569795	0.00356732439665008
RAB6B	0.933811759708842	0.0109448236612979		ENSA	-1.02370657619597	0.0161174249871283
CAV2	0.99732226024381	0.0408269875434742		JPT2	-1.10698538807139	0.00935740869041231
COX7C	1.07266576845297	0.010209697566701		JPT1	-1.10835843431927	0.00793600149391597
LCP1	1.09840574413178	0.00154609002947331		OGFOD2	-1.1132683686952	0.0415033799952983
SYT8	1.17672071307107	0.043954377019624		TMSB10	-1.14967105039942	0.00540682517524317
RAB11B	1.3714054705064	0.000626507681921401		ZNF706	-1.15566233765069	0.00111603705555586
TMEM167A	1.45520736261194	0.0219143303939675		EEF1AKMT2	-1.27111422803223	0.00946055818632448
ATP8B1	2.09279285090789	0.000411271663574841		DAP	-1.40988711973543	0.0455217421389704

Table S2. The ten most differentially expressed genes (from proteomic analyses) between NoTf accept in direct coculture and NoTf accept in Transwell coculture conditions, showing Log$_2$ enrichment of individual proteins (LogFC) and corresponding p-value (see legend to Fig. S3A for more details).

UP				DOWN		
Protein	LogFC	P value		Protein	LogFC	P value
CD14	1.75212297141304	0.000191011658635565		TYMS	-3.27748800106074	0.0134778717596743
IGFBP6	1.76618122025904	0.0139229645066205		UBE2T	-3.13040276466167	0.023473217872104
SOD2	1.77111253782108	0.000523439457102529		KPNA2	-2.98962869371859	0.0202660294833902
LAMB3	1.78650743040783	0.00216132909781278		UBE2S	-2.78386369927229	0.0408842282474856
H1-0	1.85763513264292	0.00242570948335453		CDK1	-2.70314055151148	0.0017239046757659
MYO1C	2.0046600551578	0.0143021854781629		HMGCS1	-2.26815542277271	0.003911088063124
MACROD1	2.04828979371154	0.00240255771521197		DHFR	-2.15668666737884	0.0015530984577745
CHI3L1	2.18947902717664	0.00727017238146173		PBK	-2.15384894310531	0.00508833653275616
LAMC2	2.3234530990449	0.00671123053131829		PCNA	-2.10549674961135	0.00314193588565591
SERPINA3	2.52280933849719	0.00535426160167644		PLK1	-2.09690800691763	0.0106792923363495

Table S3. Pathway analyses were performed using STRING (https://string-db.org/) with the Biological Process category from the Gene Ontology collection. For the search of BMP and p38 pathways targets, a combination of several pathways from different collections have been compiled (GOBP_POSITIVE_REGULATION_OF_BMP_SIGNALING_PATHWAY ; GOBP_REGULATION_OF_BMP_SIGNALING_PATHWAY ; GOBP_RESPONSE_TO_BMP ; PID_BMP_PATHWAY ; BIOCARTA_P38MAPK_PATHWAY ; GOBP_P38MAPK_CASCADE ; GOBP_REGULATION_OF_P38MAPK_CASCADE ; REACTOME_P38MAPK_EVENTS ; WP_P38_MAPK_SIGNALING).

#Category	Term ID	Term description	Observed gene counts	Background gene counts	Strength	Signal	False discovery rate (FDR)	Matching proteins in network (IDs)	Matching proteins in network (labels)
GO Process	GO:0007166	Cell surface receptor signaling pathway	7	2040	0.93	0.67	0.0068	9606.ENSP00000264110,9606.ENSP0000280377,9606.ENSP000003391 91,9606.ENSP0000034861,9606.ENSP0000356070,9606.ENSP0000037 9975,9606.ENSP00000414147	ATF2,USP15,CAV1,RAC1,MAPKAPK2,ILK,SPART
GO Process	GO:0071363	Cellular response to growth factor stimulus	5	473	1.42	1.01	0.0068	9606.ENSP00000264110,9606.ENSP0000280377,9606.ENSP000003391 91,9606.ENSP0000356070,9606.EN SP00000414147	ATF2,USP15,CAV1,MAPKAPK2,SPART
GO Process	GO:0007167	Enzyme-linked receptor protein signaling pathway	5	641	1.28	0.92	0.0075	9606.ENSP00000264110,9606.ENSP0000280377,9606.ENSP000003484 61,9606.ENSP0000356070,9606.EN SP00000414147	ATF2,USP15,RAC1,MAPKAPK2,SPART
GO Process	GO:0006468	Protein phosphorylation	5	736	1.22	0.82	0.0119	9606.ENSP00000264110,9606.ENSP0000280377,9606.ENSP000003391 91,9606.ENSP0000356070,9606.EN SP00000379975	ATF2,USP15,CAV1,MAPKAPK2,ILK
GO Process	GO:0030509	BMP signaling pathway	3	92	1.9	1.03	0.0156	9606.ENSP00000264110,9606.ENSP0000280377,9606.ENSP000000414147	ATF2,USP15,SPART
GO Process	GO:0030510	Regulation of BMP signaling pathway	3	104	1.85	0.97	0.0191	9606.ENSP00000339191,9606.ENSP00000379975,9606.ENSP000000414147	CAV1,ILK,SPART
GO Process	GO:0038066	p38MAPK cascade	2	13	2.58	1.01	0.0223	9606.ENSP00000264110,9606.ENSP0000356070	ATF2,MAPKAPK2
GO Process	GO:0071310	Cellular response to organic substance	6	2019	0.86	0.51	0.0356	9606.ENSP00000264110,9606.ENSP0000280377,9606.ENSP000003391 91,9606.ENSP0000356070,9606.EN SP0000379975,9606.ENSP0000004 4147	ATF2,USP15,CAV1,MAPKAPK2,ILK,SPART

Table S4. Primers for RT-qPCR used in this study.

HPRT FW	TGACCTTGATTTATTTTGCATACC
HPRT RV	CGAGCAAGACGTTCAGTCCT
TBP FW	TGTATCCACAGTGAATCTTGGTTG
TBP RV	GGTTCGTGGCTCTCTTATCCTC
BMPR1B FW	CTGTGGTCACTTCTGGTTGC
BMPR1B RV	TTCCTTTCTGTGCAGCATTC
EREG FW	TCTGCCTGGGTTTCCATCTT
EREG RV	ATTGACACTTGAGCCACACG
HOXB6 FW	CGTGCAACAGTTCCTCCTTT
HOXB6 RV	GCGTCAGGTAGCGATTGTAGT
CCND2 FW	TGCTGGAGTGGGAACTGGTG
CCND2 RV	ACACAGAGCAATGAAGGTCT
UFM1 FW	AAAGTTCCTGCTGCAACAAGTGC
UFM1 RV	ACACGATCTCTAGGAATAATCCGC
HRAS FW	ACGCACTGTGGAATCTCGGCAG
HRAS RV	TCACGCACCAACGTGTAGAAGG