Cell-type-resolved *NRXN1* isoforms across human brain tissues and hiPSC organoids

Lei Cao[1*], Yu Fan[1*], Sadaf Ghorbani[2*], Jessica Mariani[2], Yanchun Zhang[1], Michael B. Fernando[3], Jaroslav Bendl[1,4], John Fullard[1,4], Susana Isabel Ramos[5], Edward A. Mead[1], Gintaras Deikus[1], Kristin G Beaumont[1], Robert Sebra[1], Nadejda Tsankova[5], Panos Roussos[1,4,6], Kristen J. Brennand[2#] and Gang Fang[1#]

[1] *Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY 10029*
[2] *Department of Psychiatry, Yale School of Medicine, New Haven, CT, 06520*
[3] *Nash Family Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York, NY, 10029*
[4] *Center for Disease Neurogenomics; & Friedman Brain Institute; & Department of Psychiatry, New York, NY, USA*
[5] *Department of Pathology, Molecular, and Cell-Based Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA*
[6] *Center for Precision Medicine and Translational Therapeutics, Mental Illness Research Education and Clinical Center (MIRECC), James J. Peters VA Medical Center, New York, 10468, USA*

[*] *These authors contributed equally.*

[#] *Correspondence: kristen.brennand@yale.edu, gang.fang@mssm.edu*

ABSTRACT

NRXN1 undergoes extensive alternative splicing that generates a highly diverse repertoire of isoforms, diversifying protein–protein interactions, shaping synaptic specialization, and contributing to neuropsychiatric disease when disrupted. However, the splicing landscape of *NRXN1* across distinct human brain cell types remains poorly defined. Because *NRXN1* expression level is relatively low in adult brains and human induced pluripotent stem cell (hiPSC) derived neurons, single-cell Iso-seq RNA sequencing often provides insufficient coverage to capture its full isoform diversity, particularly across heterogeneous cell populations. To address this gap, we developed an integrative sequencing strategy that combines single-cell transcriptomics, targeted enrichment of *NRXN1* transcripts, and long-read sequencing. This approach reveals a comprehensive catalog of cell-type resolved *NRXN1* isoforms across adult and fetal human postmortem brains as well as hiPSC-derived cortical organoids. From the adult prefrontal cortex (PFC) region, our analyses reveal distinct splicing programs across interneuron

subtypes, pyramidal neurons, and glial lineages. Comparisons between prenatal and adult brains

indicate that *NRXN1* isoform profiles are established during early development and remain stable

throughout neuronal maturation. In hiPSC organoid models, splicing profiles partially mirror both

developmental and mature brain patterns, with a subset of splice sites exhibiting cell type- and

stage-specific divergence. In the cerebellum of an autism case and in organoids derived from

schizophrenia patients, carrying non-recurrent heterozygous *NRXN1* deletions, we identify

disrupted isoform expression and mutant *NRXN1α* isoforms and characterize their distribution

across diverse cell types. These findings help understand the complex alternative splicing of

NRXN1 in adult human brain and hiPSC-derived models and establish a framework for decoding

cell type-specific isoform diversity for important genes with relatively low gene expression levels.

INTRODUCTION

Alternative splicing (AS) is a key post-transcriptional process in the human brain that generates

diverse mRNA isoforms, contributing to the brain's functional complexity[1–4]. Presynaptic cell-

adhesion molecules known as neurexins (*NRXN1-3*) stand out as canonical examples of AS-

driven synaptic diversity, playing central roles in synapse formation, specification, and plasticity[5–

9]. The *NRXN1* gene generates an exceptional number of isoforms through six major alternative

splice sites (SS1-6)[10–12], which are essential for fine-tuning synaptic properties and diversification.

For instance, alternative splicing at splice site 4 (SS4) is known to reshape ligand binding[13–16],

while differences at SS3 in hippocampal GABAergic interneurons, specifically parvalbumin-

positive (*PV+*) and cholecystokinin-positive (*CCK+*) subtypes, have been linked to distinct

presynaptic calcium channel usage and modulatory receptors[17,18]. Aberrant alternative splicing

due to *NRXN1*[+/−] deletions disrupt the excitatory–inhibitory balance by weakening excitatory

transmission in glutamatergic neurons and enhancing inhibitory transmission in GABAergic

neurons[19], underscoring how precise isoform regulation is critical for maintaining synaptic

function.

A complete, cell-type-resolved isoform catalog of *NRXN1* remained elusive, particularly due to

the technological limitations for detection of the immense number of neurexin isoforms and their

extensive cellular diversity in the human brain. Methods such as patch pipette-based qPCR[20] and

next generation sequencing[20] can examine various insertions at alternatively spliced segments

but fail to assess the critical combinatorial nature of splicing events on a single mRNA molecule

and lack the throughput to capture the *NRXN1* splicing patterns in large populations of diverse
cells. Long-read platforms, coupled with single-cell RNA barcoding, represent a major
development to study splicing of complex genes such as *NRXN1*. However, due to the low
expression level of *NRXN1*, these transcriptomic approaches provide insufficient depth to
comprehensively assess the large repertoire of *NRXN1* isoforms.

To overcome these technical barriers, we developed an integrative sequencing strategy that
combines (1) probe-based targeted capture[21,22] with (2) single-cell/single-nucleus RNA isoform
sequencing for in-depth *NRXN1* isoform profiling, complemented with (3) an optimized RACE-seq
[23,24] (Rapid Amplification of cDNA Ends) based enrichment module to focus on specific mutant
isoforms of interest. As the core idea of this integrative strategy, single-cell transcriptomics is used
to provide cell-type deconvolution, while targeted capture is used to enrich *NRXN1* transcripts for
in-depth analysis; long read sequencing is then employed to achieve the crucial full-length isoform
resolution. The use of both probe-based targeting and RACE-seq (PCR-based) enrichment
complement each other in recovering the diversity of *NRXN1* isoforms (the strength of probes)
and detecting rare or low-abundance mutant isoforms (the strength of PCR). Using this integrative
strategy, we mapped comprehensive catalog of cell-type-specific *NRXN1* isoforms across
diverse neuronal and glial populations. From the PFC region of the human adult brain, we
revealed distinct splicing profiles across GABAergic interneuron, pyramidal neurons and glial
lineages. Interneurons derived from the same developmental origin exhibited similar AS patterns,
whereas those derived from different origins showed divergent profiles. Analyses of the AS
pattern in fetal brain suggest that *NRXN1* splicing profiles are defined in early development and
remain largely consistent during neural maturation. Using hiPSC cortical organoids derived from
patients with *NRXN1*[+/−] deletions and controls, we observed that glial progenitor cells and upper-
layer excitatory neurons (Ex-L2/3) are enriched for the mutant isoforms, and we found that the
organoids partially recapitulate the *NRXN1* splicing patterns of both developmental and adult
brains. In the postmortem brain tissue from an autistic patient case carrying a heterozygous
NRXN1 deletion, we found the mutant isoforms transcribed from deletion allele were enriched in
molecular layer interneurons (MLI1/2) and astroglia cells, and decreased the *CBLN1*, a *NRXN1*
ligand, expression in granule neurons. These findings provide critical insights into the complex
alternative splicing of *NRXN1* and establish a framework for decoding cell type-specific isoform
diversity for important genes with low gene expression levels.

RESULTS

An integrated strategy reveals cell-type catalog of *NRXN1* isoforms across human brain and hiPSC cortical o

rganoids

Single-cell long-read transcriptomics provides a unique ability to resolve full-length *NRXN1* isoforms across cell types, but whole transcriptomic profiling does not provide sufficient depth of *NRXN1* due to its very low expression level. From the published human postmortem brain single-nuclei Iso-seq data[25], *NRXN1* reads constituted only 0.09-0.2% of the total reads across different cell types, underscoring the need for a targeted enrichment strategy (**Fig. 1a, upper panel**). To maximize isoform recovery while preserving cell-type resolution, we used a targeted sequencing strategy with two complementary components (**Fig. 1b**). To generate a comprehensive isoform catalog, we designed capture probes tiled across all *NRXN1* exons at 1× density, which achieved a ~36-96-fold enrichment of *NRXN1* reads over the non-targeted sequencing method (**Fig. 1a, lower panel**). To recover cell type-specificity for isoforms of interest, particularly the mutant isoforms transcribed from the allele carrying exonic deletions, we adapted the RACE-seq (Rapid Amplification of cDNA Ends) method from short-read and optimized for long-read sequencing to achieve a more sensitive detection of low-abundance isoforms. To link isoforms to specific cell types, we matched the UMIs and cell barcodes on the *NRXN1* reads (from *NRXN1*-targeted long-read sequencing data) with the UMIs from the cell cluster information by short-read single-cell RNA-seq data (**Methods**). This design leveraged the sequencing depth and accuracy of short reads together with full-length isoform resolution by long reads, enabling precise assignment of complex *NRXN1* isoforms to distinct cell types.

We applied this workflow to single-cell or single-nucleus cDNA from multiple sources, including the PFC of adult brains, gestational cortical samples and cerebellar tissue from an *NRXN1*[+/–] autism patient. In addition, cortical organoids derived from schizophrenia patients harboring unique *NRXN1*[+/–] deletions were profiled to assess the impact of these variants on splicing regulation. After isoform identification and quality filtering, using long-read Capture-seq, we identified 50 distinct *NRXN1* isoforms across all datasets, including 27 previously annotated (five α, five β, five γ, and twelve known others) and 23 novel isoforms not represented in Ensembl[26]. Expression of these novel isoforms was validated across our in-house datasets and confirmed using independent long-read transcriptomes from postmortem human brains reported by others[25,27] (**Fig. 1c, Supplementary Fig. 1**). Among all the *NRXN1* isoforms, **alpha** isoforms

constitute the largest and most structurally complex class. We applied long-read RACE-seq, a
more sensitive method to single-cell or single-nucleus cDNA of all samples and identified 103
alpha isoforms including 57 that has been reported previously. Together, our targeted sequencing
strategy will provide a comprehensive underrating of cell-type-specific *NRXN1* alpha isoforms
(**Supplementary Fig. 1**).

NRXN1 **exhibits distinct splicing patterns across cell types in adult human brains**

To understand the *NRXN1* alternative splicing patterns across mature neurons in the human
brain, we applied the probe-based long-read Capture-seq on the single-nucleus cDNA from the
PFC region of 61 adult donors[28] (**Fig. 2a, b; Supplementary Fig. 2**). We generated 2.43M
NRXN1 reads, among which 1.13M had 10x single cell barcodes, and 1.03M reads sharing the
barcodes with short-read data. By integrating the barcode information from short- and long-read
data, a total of 125,199 cells were identified across all donors. From the long-read Capture-seq
of the single nuclei cDNA pools, we identified 46 isoforms with confident cell barcodes and UMIs
matched to short-read data, including 6 alpha isoforms not annotated but reported previously, and
14 isoforms not annotated in Ensembl[26] but later validated by our in-house brain data set (**Fig.**
1c, Supplementary Fig. 1).

The brain comprises a great diversity of cell types, and studying neurons with shared
developmental markers offers a means to uncover how *NRXN1* alternative splicing contributes to
this cellular and functional diversity. Based on developmental markers[28], neurons were
categorized as Neurod2-expressing pyramidal cells[29] (PC), *Lhx6*-expressing interneurons[30]
(*Lhx6*+-INs originated from medial ganglionic eminence, MGE), *Prox1*-expressing interneurons[31]
(*Prox1*+-INs originated from caudal ganglionic eminence, CGE), and *Lamp5*+ *Lhx6*+ GABAergic
neurons (*Prox1*+ labeled as *Lamp5*). *Lhx6*+-INs can be further subclassified into *Pvalb*+ and *Sst*+
INs, while *Prox1*+-INs are divided into *Lamp5*+, *Sncg*+, and *Vip*+ INs. GABAergic INs are a highly
heterogeneous collection of cell types that are characterized by their unique spatial and temporal
capabilities to influence neuronal circuits[32–34]. We observed that GABAergic interneurons showed
the greatest *NRXN1* splicing diversity (**Fig. 2c, top**). Splicing patterns were more variable at SS1,
SS3, and SS4, whereas SS2 and SS6 were invariably excluded and SS5 was consistently
included. Within the *Prox1*+-INs, *the Lamp5*+, *Sncg*+ and *Vip*+ subtypes share similar patterns
across all 6 alternative splicing sites, while *Pvalb*+ and *Sst*+, subclasses of *Lhx6*+-INs, differed at

SS1. We also identified a *Lamp5⁺ Lhx6⁺* subclass that shared splicing patterns with the *Prox1⁺*-

INs and *Sst⁺* of *Lhx6⁺*- INs in (**Fig. 2c**). In PC neurons (**Fig. 2c, middle**), we found: (1) across

cortical layers, exons at SS1, SS3 and SS5 were dominantly spliced in ("in"; upward bars), while

exons at ASS6 and SS4 are dominantly spliced out. (2) layer-specific differences at SS2, where

exon 7 was often spliced out in PCs of layers L2/3-IT, L5-IT, and L6-IT-Car3, but spliced into most

PCs of layers L4-IT, L5/6-NP, L6B, L6-CT, and L6-IT. Among glial cells (astrocytes,

oligodendrocyte progenitor cells (OPCs), oligodendrocytes), exon 17 at SS6 was consistently

spliced out, whereas exon 23 at SS5 was spliced in. Oligodendrocytes showed spliced-in for

exon1 and exon5 at SS1, which were dominantly spliced-out in astrocytes and OPCs. At ASS4,

exon 21 was included in astrocytes but not in OPCs or oligodendrocytes, consistent with prior

studies showing that astrocytic neurexins splice SS4 at high ratios compared to excitatory

neuronal neurexins[35,36]. Building on the striking patterns of *NRXN1* alternative splicing observed

in the human brain, we conducted principal component analysis (PCA) of splicing levels across

cell types, which separately clustered *Prox1⁺*-INs, *Lhx6⁺*-INs and PCs classes (**Fig. 2d**). Overall,

these findings suggest that *NRXN1* splicing profiles exhibit substantial differences across various

neuronal populations and glial lineages. Within GABAergic neurons, interneurons derived from

the CGE (*Prox1⁺*-INs subtypes) are closely clustered by the splicing patterns, yet distinct from INs

originating from the MGE (*Lhx6⁺*-INs subtypes).

***NRXN1* splicing profiles remain consistent across cells sharing a neurogenic origin during**

neural maturation

To determine whether the *NRXN1* splicing patterns of adult brain are defined as early as in the

human developing neocortex, we generated a snRNA-seq dataset from two unfixed snap-frozen

neocortex of postmortem samples obtained from the second and third trimesters of gestation (24

and 33 gestational weeks, gw). After quality control, we obtained the transcriptomic data from

52,426 nuclei, with a median of 2,910 genes, 6,156 UMIs from each nucleus. We identified 10

primary cell types from the single cell data: five neuronal clusters, including excitatory neuronal

subpopulations expressing layer-specific cortical marker genes (EX-L2/3, EX-L4/5/6); inhibitory

neuronal subpopulations (IN-CGE, IN-MGE and IN-Reelin); and glial cell populations, radial glia

and astrocyte (RG/AC) clusters, astrocytes, OPCs, transient glial intermediate progenitor cell

(gIPC) population, microglia and *FLT1*+ blood vessel cells (BVCs)[37] (**Fig. 3a**). We further validated

our cell annotations by comparing top differentially expressed genes (DEGs) per cell cluster to

published canonical lineage and proliferation markers[37,38] (**Fig. 3c**). Next, we analyzed how cell
type proportion varied between prenatal neocortex and adult PFC. The 33 gw neocortex sample
had more glial, INs-MGE and INs-CGE, whereas the 23 gw sample had more INs-Reelin and
excitatory neurons EX-L4/5/6 (**Fig. 3b**). Neuronal populations constituted most cells in the fetal
neocortex, whereas glial transient intermediate progenitor cells were exclusively present in fetal
neocortex and subsequently disappeared or transformed along with the cortical development in
adult PFC, which is consistent with the stages of neurogenesis in human cerebral cortex[39,40]
(**Supplementary Fig.3**).

To determine *NRXN1* splicing patterns by cell type in the fetal neocortex, we applied the probe-
based long-read Capture-seq on the single nuclei cDNA. Based on regional marker genes, exon
23 at SS5 was predominantly spliced in across all cell types, whereas exons at other splice sites
showed variable inclusion. In most (>60%) PCs across all layers, exons at SS1 and SS3 were
consistently spliced in, exon 21 at SS4 was spliced out, and exon 17 at SS6 showed variable
usage. In contrast, exon 7 at SS2 was differentially spliced between upper-layer (L2/3) and
deeper-layer (L4/5/6) PCs. Similarly, in GABAergic interneurons (INs), exons at SS1 and SS3
were largely spliced in, while exons at SS2, SS4, and SS6 were spliced out. Notably, the
proportion of IN-*Lhx6*$^+$ cells was higher than that of IN-*Prox1*$^+$ cells at SS3 (spliced in) and SS4
(spliced out). The glial cells were variably spliced at SS1, SS2, SS6 and SS4, but consistently
spliced-in at SS3 and SS5 (**Fig. 3d**). We next compared *NRXN1* splicing in matched neuronal
populations between prenatal neocortex and adult PFC. Exon inclusion level across 6 splicing
sites were highly correlated between prenatal neocortex and adult PFC for Lhx6$^+$-INs (*Pearson's*
r = 0.82, *p* = 0.01), *Prox1*$^+$-INs (*Pearson's r* = 0.87, *p* = 0.005), PCs (*Pearson's r* = 0.87, *p* = 0.005)
and glial populations (OPC, *Pearson's r* = 0.9, *p* = 0.003; RG/AC, *Pearson's r* = 0.82, *p* = 0.01)
(**Fig. 3e, Supplementary Fig. 3**). These findings suggest that *NRXN1* splicing profiles are defined
in early development and remain largely consistent during neural maturation.

***NRXN1*$^{+/-}$ hCOs splicing profiles recapitulate developmental and mature features and**
identify cell types expressing the mutant isoforms.

From hiPSC-derived 2D neurons with a 3′-*NRXN1* deletion, we previously uncovered dozens of
mutant isoforms exclusively from the deletion allele (mainly affects alpha and beta isoforms)[12].
Later, we demonstrated that this exonic deletions at the 3' site disrupted normal *NRXN1* splicing
and isoform expression, leading to distinct phenotypes in iGLUT and iGABA neurons, primarily

reflected in altered synaptic frequency[19,41]. Building on these observations, we sought to explore

how the deletions affect the splicing events across cell types in a more complex

neurodevelopmental system. We generated human cortical organoids (hCOs) from the control

line and schizophrenia probands with the 3'deletion (3'-del) in parallel. hCOs were guided toward

the pallium of dorsal forebrain[42] (**Methods, Supplementary Fig. 4**). To profile cell types in the

hCOs, we performed single cell RNA-seq on hCOs from two 3'-del cases (line 581 and line 641)

and one control line (line 690) at day 150 with the 10x NextGEM 5pv2 kit, when neural activity is

well characterized[43–45]. After quality control data processing steps, we merged high-quality

scRNA-seq data from 3 cortical organoid samples into a core dataset of 27,325 cells. Following

unsupervised clustering, we identified and annotated major cell types using an extensive curated

list of known marker genes[19,42], which consisted of transit-amplifying cell (TAC), neuronal

intermediate progenitor cells (nIPCs) giving rise to distinct subpopulations of glutamatergic

excitatory neurons (Ex-L2/3, Ex-L4/5/6) and GABAergic inhibitory neurons (INs, IN-SST,IN-

Reelin); gIPCs leading to the subpopulations of radial glia and astrocyte clusters (RG/AC); as well

as BVCs (**Fig. 4a**). We further validated our cell annotations by comparing gene expression

signatures of our scRNAseq cell clusters to published brain organoid datasets as reference (**Fig.**

4b). We then compared cortical organoids derived from control iPSC line with the prenatal

neocortex in terms of cell-type composition. We found gIPCs (OR = 2.06, *Fisher's test*) and Ex-

L2/3 (OR = 1.67, *Fisher's test*) were significantly enriched in hCOs from the 3'del lines than in

control lines, while cell number of IN-Reelin were significantly low in 3'-del lines (OR = 0.03,

Fisher's test) (**Fig. 4c**).

To more comprehensively understand the greater splicing patterns of *NRXN1* across the cortical

organoid cell types, we applied the targeted Capture-seq on the full-length single-cell cDNA. From

the control lines and those with the 3' deletion, we generated 317,521 *NRXN1* reads, of which

144,450 with 10x single cell barcodes and 120,873 reads sharing the barcodes with short-read

data. Integrating the barcode information across short- and long-read data, we identified 23,187

cells from all cortical organoid samples (**Supplementary Fig. 4**). Using these targeted single cell

data, we first analyzed the *NRXN1* AS patterns in excitatory neurons, inhibitory neurons and non-

neuronal cells for both 3'del lines and control lines at the advanced maturation stage. Within the

glutamatergic excitatory neurons (the major population in hCOs), *NRXN1* splicing patterns were

highly similar between control lines and 3'del lines, except for differences in exon inclusion level

at SS1 (exon 3) and SS3 (exon 12) (**Fig. 4d,e**). The ~136-kb deletion impacting SS4 and SS5

(exons 21-23)[12,19] slightly reduced exon inclusion at SS4 in the 3'del lines (**Fig. 4e**).

As the deletion in the 3'-region of *NRXN1* generates mutant alpha and mutant beta isoforms, we

sought to know which cell types are enriched for these mutant isoforms. As some of the mutant

isoforms are low in abundance and the Capture-seq works better with abundant isoforms, we

employed long-read 5'RACE-seq to resolve alpha isoforms with single cell barcode information

from the 5'GEX single cell cDNA (Methods). We identified 21 α-isoforms across all cortical

organoids, of which 18 were not detected by long-read Capture-seq, including six mutant isoforms

from the two 3′-del *NRXN1*$^{+/-}$ organoids (**Fig. 4f**). In addition, we identified seven β-isoforms,

including one mutant β-isoform (**Supplementary Fig. 4**). Compared with the control lines, the

mutant α-isoforms were expressed widely across all cell types in both lines with 3′-*NRXN1*$^{+/-}$

(except for *DLX1/2*-INs in 641). In contrast, the mutant β-isoform was detected in fewer cell

clusters, showing consistent expression in excitatory neurons in both 3′-*NRXN1*$^{+/-}$ lines, and

exclusively in TACs in line 581 and RG/AC cells in line 641 (**Fig. 4g**). The composition of *NRXN1*

isoform by cell type demonstrated that mutant isoforms from the allele bearing 3'-deletion are

expressed in all cell types, contributing to altered synaptic activity in excitatory neurons and

DLX1/2-INs.

With *NRXN1* splicing profiles available across multiple developmental stages, we next asked

which stage the control hCOs most closely resemble in terms of *NRXN1* splicing– the fetal

neocortex or the adult PFC. We therefore compared the exon inclusion level across splicing sites

(SS1-6) for *NRXN1* in three major cell classes that appear in all samples, *i.e.,* GABAergic

inhibitory neurons, excitatory neurons, and RG/AC (**Fig. 4h**). In GABAergic neurons, exon

inclusion levels were highly correlated between organoids and the fetal neocortex (*Pearson's r =*

0.97, p = 3.82E-04) and between cortical organoids and the adult PFC (*Pearson's r = 0.88, p =*

0.008), although exons at SS2, SS3, and SS5 displayed differential splicing. Similarly, significant

agreement was observed between cortical organoids and the fetal neocortex in glutamatergic

excitatory (Glu-EX; *Pearson's r = 0.73, p = 0.04*) and radial glia/astrocyte (RG/AC; *Pearson's r =*

0.78, p = 0.04) populations, and between organoids and the adult PFC in Glu-EX (*Pearson's r =*

0.88, p = 0.004) and RG/AC (*Pearson's r = 0.92, p = 0.003*) populations (**Supplementary Fig. 4**).

Together, these findings indicate that cortical organoid splicing profiles partially recapitulate both

developmental and mature splicing patterns, with a subset of splice sites showing cell type- and

stage-specific divergence (**Fig.4h, Supplementary Fig. 4**).

***Exonic* deletion impacts *NRXN1* splicing in cerebellum from an ASD brain**

NRXN1 deletions are also strong risk factors for autism spectrum disorder (ASD)[46,47]. The
cerebellum shows consistent structural and functional alterations in ASD, and *NRXN1α* isoform
complexity is significantly different in the cerebellum compared to cortex. To uncover how *NRXN1*
isoform diversity contributes to ASD, we applied our probe-based long-read Capture-seq to the
postmortem cerebellum of an *NRXN1*[+/-] ASD patient with an exon 9-13 deletion (**Supplementary**
Fig. 5). We performed single-nucleus RNA-seq on this sample and obtained 9,689 high-quality
nuclei. Unsupervised clustering and annotation identified eight cerebellar cell classes: excitatory
granule cells; inhibitory neurons (Purkinje cells; two types of molecular layer interneurons MLI1,
MLI2; and Purkinje-layer interneurons); astrocytes; Bergmann glia; oligodendrocytes; and OPCs
(**Fig. 5a**). These annotations matched expected marker gene expression[48] (**Fig. 5b**).

When applying the probe-based long-read Capture-seq to the single nuclei cDNA of cerebellum,
we identified 9 *NRXN1* isoforms with cell barcodes matched to short-read data, including two
mutant isoforms associated with the exon 9-13 deletion (**Fig. 5c)**. Among all the *NRXN1* reads
with cell barcodes matched to short-read data, the mutant isoform transcript 3312 and transcript
4131 accounted for ~35% of reads (**Fig. 5d**). When assigning isoforms to cell types, we found the
two mutant isoforms were significantly enriched in MLI1 (*Fisher's Exact test*, *p* =5.72E-18), MLI2
(Fisher's Extact *test*, *p* =0.04), and astroglia cells (*Fisher's Extact test*, *p* =51.49E-13) compared
to wild-type *NRXN1* isoforms (**Fig. 5e**). Next, we examined *NRXN1* splicing at six alternative
splicing sites in these cell types. We found that: (1) over 70% of granule cells are spliced-in at
SS1, SS3, SS4 and SS5, while more than half spliced-in at SS2; granule cells predominantly
spliced out at SS6; (2) The two MLI subtypes showed different splicing trends at SS1, SS2 and
SS5; (3) within the glia lineage, OPC and oligodendrocytes are uniformly spliced-in at SS5 but
spliced-out at ASS6. As the exonic deletion expanded the SS3 (exon12) and produced mutant
alpha isoforms, we asked whether mutant isoform expression correlated with synaptic ligands.
Among granule neurons, cells expressing mutant-type *NRXN1* isoforms showed significantly
decreased expression of cerebellin genes than cells expressing wildtype isoforms. Because
granule cells regulate Purkinje cells via the *CBLN1–NRXN1* complex that organizes and
maintains Purkinje cell synapses[49], our results link *NRXN1* deletions to abnormal Purkinje cell
signaling and highlight *NRXN1*'s functional role in ASD.

DISCUSSION

We applied an integrative isoform sequencing strategy to generate a cell-type-resolved catalog
of *NRXN1* isoform diversity across human cortical organoids, prenatal cortex, adult PFC, and
cerebellum samples. *NRXN1* demonstrated extensive cell type- and cell subtype specificity. For
example, in hCO organoids and adult PFC, *NRXN1* splicing displayed apparent cell-type
specificity, while the cerebellum of an ASD case exhibited distinct deletion-associated isoforms
enriched in molecular-layer interneurons and astroglia. Together, these results establish a
framework for decoding *NRXN1* alternative splicing.

Our integrative pipeline, which combines probe-based capture, single-cell barcoding, and long-
read sequencing, overcomes major obstacles in detecting low-abundance, full-length *NRXN1*
transcripts. Targeted enrichment increased on-target recovery by ~50-fold on average compared
with untargeted long-read methods, while preserving single-cell resolution. The workflow links full-
length isoforms to their cell identity, enabling quantitative mapping of *NRXN1* splicing patterns
across neuronal and glial populations, linking the genetic variations with functional phenotypes.
RACE-seq and Capture-seq represent complementary strategies for characterizing transcript
isoforms. Capture-seq employs probe-based hybridization to enrich targeted transcripts from
barcoded cDNA, enabling detection of novel isoforms (alternative transcription start sites/
transcription termination sites) for many gene targets in one assay across cell types. This method
preserves relative transcript abundance and is compatible with single-cell barcoding[50], making it
well-suited for analyzing splicing events at defined sites. However, Capture-seq preferentially
detects abundant isoforms and shows reduced sensitivity for low-abundance or long α-isoforms.
In contrast, long-read RACE-seq selectively amplifies transcripts from defined 5′ or 3′ ends,
allowing accurate reconstruction of specific full-length isoforms, including those spanning deletion
junctions in mutant alleles. Although long-read RACE-seq is less quantitative and lower
throughput, it offers greater sensitivity and completeness for targeted transcripts, while being
more susceptible to RNA degradation. Together, these approaches provide a comprehensive
view of transcript diversity: long-read Capture-seq defining global splicing architecture and long-
read RACE-seq validating and extending isoform-level resolution in selected contexts, though
both are limited by RNA integrity and read length, particularly in postmortem brain. This approach
is applicable to other synaptic genes with extensive splicing, bridging the gap between single-cell
transcriptomics and isoform-level resolution.

As the dominant population in the cortex, glutamatergic excitatory neurons demonstrated
remarkable concordance of *NRXN1* splicing patterns between prenatal neocortex and adult PFC.
Likewise, glutamatergic neurons from iPSC-derived organoids and prenatal cortex shared similar

NRXN1 splicing, indicating that patient-specific *NRXN1[+/-]* organoids captured key developmental

splicing events. GABAergic interneurons in the PFC show extensive molecular and functional

diversity, reflected in their distinct *NRXN1* splicing profiles. Although *Pvalb*-expressing and *Sst*-

expressing interneurons both originate from the MGE (*Lhx6+*), they exhibited subtype-specific

exon usage patterns, consistent with their previously observed phenotypic differences[51,52].

Notably, *NRXN1* splicing programs appear to be established early in development and remain

largely stable, as evidenced by the strong concordance of exon inclusion levels in *Prox1[+]*

interneurons between the prenatal neocortex and adult PFC[20].

Extending this analysis to an *NRXN1[+/−]* cerebellum from an ASD patient revealed unique mutant

isoforms (exon 9-13 deletion) that were highly expressed in molecular-layer interneurons and

astroglia, whereas granule cells retained WT isoforms. Cells expressing mutant *NRXN1* showed

decreased cerebellin ligand expression, implicating the mutant isoforms in disrupted synaptic

adhesion. These findings highlight that *NRXN1* deletions can differentially impact cortical and

cerebellar circuits, aligning with the cerebellar dysfunction frequently reported in ASD[53–55]. In the

cerebellum, granule cells influence Purkinje cells through the *Cbln1-NRXN1* complex, which

organizes and maintains the parallel fiber-Purkinje cell synapse. This trans-synaptic pathway is

essential for forming functional excitatory synapses and inducing long-term depression, a key

form of synaptic plasticity. Our findings suggest that disrupted Purkinje cell signaling may be

linked to *NRXN1* deletions. These findings reveal that *NRXN1* deletions disrupt cerebellar and

cortical circuits through precise, cell-type–specific mechanisms, paralleling the dysfunction

characteristic of ASD and schizophrenia. Given that *NRXN1* isoforms dictate partner specificity

at synapses, the isoform catalog provides a foundation for isoform-targeted interventions, such

as antisense oligonucleotides, to selectively restore normal *NRXN1* splicing and rescue circuit

function*.*

FIGURE CAPTION

Figure1. A targeting strategy reveals cell-type catalog of *NRXN1* isoforms.

(a) *Upper panel* Bar graph showing the percentage of *NRXN1* reads detected by Capture-seq
(orange) and non-targeted iso-seq (blue) across major brain cell classes. GABA, GABAergic
neurons; Glu, glutamatergic neurons; Astro, astrocytes; Micro, microglia; Oligo,
oligodendrocytes; OPC, oligodendrocyte progenitor cells. *Lower panel:* Fold change of *NRXN1*-
mapped reads obtained by Capture-seq relative to the non-targeted single-cell Iso-seq.

(b) Schematic illustrating the comprehensive characterization of cell-type-specific *NRXN1*
isoforms using the targeting strategy. Full-length single-cell/nuclei cDNA is divided into two
parts: one processed for a short-read library using Tn5 transposase and sequenced to identify
cell types; the other subjected to probe-based long-read Capture-seq and RACE-seq to detect
NRXN1 isoforms with matched barcodes and UMIs (see Methods).
(c) Schematic representation of *NRXN1* isoforms identified across samples by single-cell long-
read Capture-seq. Each row represents a unique isoform, color-coded as α, α-mutant, β, γ,
known others (other), and novel isoforms (other-novel). The heatmap displays the expression of
each isoform in adult prefrontal cortex, fetal neocortex, and human cortical organoids.
395

Figure 2. *NRXN1* expression repertoires exhibit distinct splicing patterns across cell
types in human brains.

(a) Workflow overview of long-read Capture-seq applied to single-nucleus cDNA from adult
PFC.
(b) UMAP visualization of putative cell types in the adult PFC. Colored clusters represent
neuronal and glial subtypes.
(c) Bar plots showing normalized *NRXN1* splicing events at SS1–SS6 in GABAergic
interneurons (top), glutamatergic excitatory neurons (middle), and glial lineages (bottom). Each
bar represents a cell class. Within GABAergic interneurons, *Prox1*$^+$-INs are colored in green,
Lhx6$^+$-INs in red, and *Lamp5*$^+$ *Lhx6*$^+$INs in orange. Exon inclusion ("splice-in") is shown upward
and exclusion ("splice-out") downward. Data represent mean ± SEM.
(d) PCA plot based on *NRXN1* splicing patterns across splice sites SS1–SS6, showing that
Prox1$^+$-INs, *Lhx6*$^+$-INs, and PCs form distinct, separately clustered classes. Colors correspond
to cell types as in (c): *Prox1*$^+$- INs (green), *Lhx6*$^+$-INs (red), *Lamp5*$^+$ *Lhx6*$^+$-GABAergic neurons
(orange), glutamatergic neurons (blue), and glial lineages (gray).
411

Figure 3. *NRXN1* splicing profiles remain consistent across cells sharing a neurogenic
origin during neural maturation.

(a)UMAP visualization of putative cell types in the neocortex from two fetal brains. *Left:* merged
UMAP of combined datasets; *Right:* separate UMAPs showing neocortex samples from 23 gw
and 33 gw. Colored clusters represent neuronal and glial subtypes in the fetal neocortex.
Abbreviations: gIPC, glutamatergic intermediate progenitor cells; RG/AC, radial glia/astrocyte
lineage; Ex, excitatory neurons; IN, inhibitory neurons; BVC, blood vessel cells; MG, microglia.
(b) Bar graph showing the proportions of each cell class in 23 gw and 33 gw neocortex samples.
Colored blocks indicate neuronal and glial subtypes.
(c) Heatmap of marker genes expression used for cell-type annotation. Each row represents a
marker gene and each column a cell class; darker red indicates higher expression. Colored
blocks indicate neuronal and glial subtypes in fetal neocortex.
(d) Bar plots showing normalized *NRXN1* splicing events at SS1–SS6 in *Prox1*$^+$-INs (green),
Lhx6$^+$-INs (red), glutamatergic excitatory neurons (blue), and glial lineages (gray). Exon
inclusion ("splice-in") is shown upward and exclusion ("splice-out") downward. Data represent

mean ± SEM.

(e) Pearson's correlation of *NRXN1* exon inclusion levels across splicing sites (SS1–SS6)

between adult PFC and prenatal neocortex in *Lhx6*[+]-INs (left), *Prox1*[+]-INs(middle), and PC

neurons (right).

Figure 4. *NRXN1*[+/−] human cortical organoids (hCOs) splicing profiles recapitulate

developmental and mature features and identify cell types expressing the mutant

isoforms.

(a) *Left:* UMAP of putative cell types in hCOs, with colored clusters representing neuronal and

glial subtypes. *Right:* Proportions of each cell class in organoids derived from control line 690 or

3′del lines (581 and 641). Abbreviations: gIPC, glutamatergic intermediate progenitor cells;

nIPC, neuronal intermediate progenitor cells; TAC, transient amplifying cells; RG/AC, radial

glia/astrocyte lineage; Ex, excitatory neurons; IN, inhibitory neurons; BVC, blood vessel cells.

(b) Heatmap of marker gene expression used for cell-type annotation. Each row represents a

marker gene and each column a cell type; darker red indicates higher expression.

(c) Bar graph showing enrichment of specific cell types in 3′del organoids compared with control

lines.

(d) Bar plots showing normalized *NRXN1* splicing events at SS1–SS6 in all cell classes from

control hCOs (upper) and 3′del hCOs (lower). Exon inclusion is shown upward ("splice-in"),

exclusion downward ("splice-out"). Data represent mean ± SEM.

(e) Pearson's correlation of *NRXN1* exon inclusion levels across splicing sites (SS1–SS6)

between 3′del and control hCOs.

(f) *Left:* Schematic of *NRXN1α* isoform structures identified by long-read RACE-seq. Each row

represents a unique *NRXN1α* isoform, with blue exons indicating inclusion and blanks indicating

exclusion. *Right:* Presence or absence of each *NRXN1α* isoform across 3′del and control hCOs

(green = present; blank = absent).

(g) Bar graph showing the composition of *NRXN1* isoforms identified by long-read RACE-seq

within each cell type across 3′del and control hCOs.

(h) Bar graph showing exon inclusion levels at SS1–SS6 in GABAergic INs (top), glutamatergic

excitatory neurons (middle), and RG/AC (bottom) across hCOs (blue), fetal neocortex (orange),

and adult PFC (green). Statistical comparisons were performed using one-way ANOVA (*, $P <$

0.05; **, $P < 0.01$; ***, $P < 0.001$).Data represent mean ± SEM.

Figure 5. Patient-specific exonic deletion impacts *NRXN1* splicing patterns across cell

types from an ASD cerebellum.

(a) *Left:* UMAP of putative cell types in the cerebellum. *Right:* Proportions of each cell class in

the cerebellum.

(b) Dot plot showing expression of marker genes used for cell-type annotation. Each row

represents a marker gene; cell types are shown along the x-axis.

(c) Schematic of *NRXN1* isoform structures identified by long-read capture-seq, with each row

representing a unique *NRXN1* isoform. The 3′-*NRXN1*[+/-] specific mutant isoforms are highlighted

in red.

(d) Bar graph showing read counts of *NRXN1* isoforms with single-cell barcodes and UMIs by
long-read Capture-seq. The red box highlights 3'-*NRXN1*[+/-] specific mutant isoforms.
(e)Bar graphs show the proportion of cells in each cell class expressing any NRXN1 isoforms,
wild-type isoforms(*NRXN1*-WT) and mutant isoforms (*NRXN1*-MT).
(f) Bar plots showing normalized *NRXN1* splicing events at SS1–SS6 across all cell classes
from *NRXN1*[+/-] cerebellum. Exon inclusion is shown upward ("splice-in") and exclusion
downward ("splice-out"). Data represent mean ± SEM.
(g) Violin plots showing average expression of *NRXN1*-binding ligands in granule cells
expressing wild-type (WT-expressing) versus mutant *NRXN1*(MT-expressing).

METHODS

Data Availability

All sequencing data generated for this study will be uploaded to the Sequence Read Archive
and made publicly available.

Acquisition and processing of human post-mortem samples

The MSSM cohort single nuclei cDNA. The sequencing data and full length single-nucleus
cDNA libraries of 61 individuals from the Mount Sinai School of Medicine cohort were generated
from the previous publication[28]. The leftover barcoded full length single-nucleus cDNA was used
as input for the current study. To obtain 1-1.5 µg of cDNA for the hybridization step, 5-10 ng of
the surplus barcoded 10x cDNA was amplified in two 50 µl reactions, which consists of 25 µL
2X PrimeSTAR GXL premix (Takara, Cat# R051A), 1 µL partial read 1 primer (10 µM, 5'-
CTACACGACGCTCTTCCGATCT-3'), 1 µL partial TSO primer (10 µM, 5'-
AAGCAGTGGTATCAACGCAGAG-3'). The amplification program is set as follows: 98 °C for 30
seconds, 8 cycles of 98 °C for 10 seconds, 65 °C for 15 seconds and 68 °C for 8 minutes, 1
cycle of 68 °C for 10 minutes. The concentration of PCR products was quantified by Qubit, and
additional cycles were added if the yield was lower than 500 ng per reaction. The PCR products
were cleaned up with 0.6× AMPure XP beads.

NRXN1[+/-] **cerebellum of human post-mortem brain.**

Postmortem human cerebellum of the ASD case with a heterozygous *NRXN1*[+/-] mutation were
collected and consented through the Department of Pathology at Western Michigan University
Homer Stryker MD School of Medicine under WCG IRB protocol #20111080 and the other
samples were consented through the National Institute of Mental Health Intramural Research
Program under NIH protocol #90-M-0142, and was acquired by LIBD via material transfer
agreement.

Fetal brain. Two human gestation cortical tissue samples(24 gw and 33 gw) were obtained
from previous study [37] in accordance with the policies and regulations at the Icahn School of
Medicine at Mount Sinai and its Institutional Review Board.

Single cell/nucleus full-length barcode cDNA. 60mg of the frozen post-mortem tissues was
submitted to Yale Center for Genome Analysis for single nuclei isolation. Briefly, single nuclei
were automatically extracted from the frozen tissue using the Singulator system, which includes
washing nuclei suspension in buffer supplemented with sucrose, and nuclei FACS sorting. The
isolated nuclei were then processed using the 10x Genomics 5'GEX V2 to create single nuclei
cDNA and matched short-read RNA-seq libraries per manufacturer's instructions.

Patient-derived organoids

Human induced pluripotent stem cell (hiPSC) lines used were validated using methods as
previously described. Genome-wide SNP genotyping was performed using Illumina genome-wide
GSAMD-24v2-0 SNP microarray at the Children's Hospital of Philadelphia. Cultures were tested
for and maintained Myco-plasma free. A total of 3 control hiPS cell lines derived from fibroblasts
collected from 3 healthy subjects were used for experiments. Approval for this study was obtained
from the Stanford IRB panel, and informed consent was obtained from all subjects.

Stem cell maintenance. Human iPSCs were obtained as previously described in Flaherty et
al., 2019. Under sterile conditions, 6-well plates (Corning 3516) were coated with 60 ug/mL
Geltrex matrix (Thermo Fisher A1413302) diluted in Dulbecco's Modified Eagle Medium
(DMEM, Gibco 11965-092) per well and incubating them for at least 30 min at 37 °C. All hiPSC
lines were seeded on coated plates and maintained in StemFlex (StemFlexTM Basal Medium
(Gibco A3349301) supplemented with StemFlexTM Supplement (Gibco A3349201)) at 37 °C.
After 4 days, hiPSC lines were passaged by removing media washing cells with PBS. Then cells
were incubated in 0.5 μM EDTA in PBS (Invitrogen AM9261) for 4 min at 37 °C. After removing
EDTA, colonies were broken into smaller pieces by triturating using a p1000 pipette and
StemFlex. It's worth noting that very small colonies have a very low survival chance and should
be supplemented with CHR. After dissociation, colonies were distributed in new coated plates
containing StemFlex in a ratio of 1:6 (250K cell per well of a 6 well plate). A complete media
change was performed the following day. Complete media changes were performed as needed
or every other day.

Cortical organoid generation. Cultures of hiPSC were dissociated at 70% confluency and
checked for signs of differentiation. Mature undifferentiated cells were washed with PBS,
followed by incubation with Accutase (Thermo Fisher Scientific) and chroman (CHR) for 10-20
min to achieve complete detachment. Cells were gently triturated to achieve a single-cell
suspension. The resulting cell suspension was transferred into DMEM (Thermo Fisher
Scientific) and centrifuged at 500 × g for 4 min. Cells were resuspended in StemFlex™ medium
(Thermo Fisher Scientific) supplemented with CHR and diluted at a density of 1.5 × 10^6
cells/mL. AggreWell™ plates (STEMCELL Technologies) were prepared with anti-adherent
solution according to the manufacturer's instructions. A volume of 2 mL of the hiPSC
suspension was seeded per well to promote spheroid formation overnight (defined as day 0).
Neural induction was subsequently performed following the human forebrain spheroid protocol
described by Sloan et al. (2018). In brief, medium was replaced daily until day 15, every other
day from days 15–42, and every 4 days from day 43 onward. On day 1, embryoid bodies were
dislodged and transferred to ultra-low adhesion 10 cm dishes containing Neuronal Induction
Media (StemFlex™ medium without StemFlex™ supplement), SMAD inhibitors SB431542 (100
nM; Tocris), LDN193189 (100 nM; Stemgent), XAV (100nM; Stemcell technology). Plates were
placed on an orbital shaker at 53 rpm and maintained at the same speed until day 6. From days
6–25, Neuronal Differentiation Media (Neurobasal-A (Thermo Fisher Scientific), 1x Antibiotic-

Antimycotic, 1x GlutaMAX™, and 1x B27 supplement minus vitamin A (Thermo Fisher
Scientific) was used supplemented with 100 nM Fibroblast Growth Factor (FGF) and 100 nM
Epidermal Growth Factor (EGF). Between days 25–42, Neuronal Differentiation Media was
supplemented with brain-derived neurotrophic factor (BDNF; 100 nM; PeproTech) and
neurotrophin-3 (NT3; 100 nM; PeproTech). From day 42 onwards, organoids were maintained in
Neuronal Differentiation Media.

Cryoprotection and immunocytochemistry. Organoids were washed with PBS and fixed in 4%
paraformaldehyde (PFA)/phosphate buffered saline (PBS) overnight at 4°C. Organoids were then
washed in PBS and transferred to 30% sucrose/PBS for 48 hours. Organoids were embedded in
gelatin/sucrose embedding solution by dissolving 7.5 g gelatin in 100 ml 10% sucrose solution at
37°C to obtain a 7.5% gelatin/10%sucrose embedding solution. For immunofluorescence
staining, 14-16 μm-thick sections were washed with PBS and blocked in 10% donkey serum, and
0.1% Triton-X in PBS for 1 h at room temperature followed by primary (overnight, 4 °C) and
secondary antibody (1–2 h, Jackson ImmunoResearch or ThermoFisher Scientific) incubation.
Slides were then mounted (VECTASHIELD, Vector Labs) and imaged on a Zeiss microscope with
an apotome module quipped with ZEN 3.3 (ZEN pro) software. Antibody list: BCL11B/CTIP2 (rat,
1:500, Abcam), FOXG1 (rabbit, 1:200, Abcam), GAD1/GAD67 (mouse, 1:1000, Millipore), HuC/D
(mouse, 1:500, Invitrogen), KI67 (rabbit, 1:500, Vector Labs), PAX6 (mouse, 1:200, BD
Bioscience), RELN (mouse, 1:100, MBL), SOX1 (goat, 1:20, R&D Systems), TBR1 (rabbit,
1:1000, Abcam).

The sections were then incubated overnight at 4°C with primary antibodies diluted in 1% BSA,
0.1% Triton-X, sodium azide. Cryosections were washed three times and were incubated with
secondary antibodies in 10% donkey serum, 0.1% Tri- ton-X. Aquamount (Polysciences, 18606)
was used to mount cover glass slides. Imaging was performed on a Leica TCS SP8 confocal
microscope or Zeiss upright fluorescent microscope. Images were processed and analyzed
using Fiji (NIH) and IMARIS (Oxford Instruments). To quantify percentage of oligodendrocyte
precursor cells, images of Hoechst and OLIG2+ nuclei were taken. Hoechst images were
converted into 8-bit images, and we performed automatic image threshold (Default, Dark
Background) into a binary image, converted to Mask, performed Watershed (FiJi, NIH). OLIG2
nuclei images were manually thresholded due to inconsistency of background signals by a
researcher blinded to molecular condition. Nucleus from Hoechst and OLIG2 images were
quantified using Analyze Particles with pixel size 30-1000, and the number of OLIG2 nuclei
counts were di- vided by Hoechst nuclei counts to obtain a percentage. Ventricular-like regions
were quantified by examining cross sections of organoids stained with Hoechst and counting
clearly identifiable ventricle-like structures.

Single cell dissociation and RNA-seq library preparation. Organoids with optimal morphology
from each line were selected in days 25, 60, and 120. Five organoids from each line were pooled
at each time point and transferred to a 5 cm plate dish and washed with PBS. Organoids were
chopped into small pieces using a sharp sterile blade. 300 ul of Accumacs was added on top of
the pieces without scattering them. Pieces were collected by a cut tip and were transferred into a
1.5 ml tube. The tubes were incubated at 37°C and 800 g rotation on a thermal shaker for 20 min.
After 20 min, cell suspension was very gently pipetted up to 5 times with a p 1000 tip, to facilitate
the cell dissociation. 500 ul fresh Accumacs was added into the suspension and incubated again
for 20 min. Cell suspension was pipetted up to 5 times more and passed through a filter in a 15
ml sterile tube containing 1 ml of HEPES buffer. The filter was washed with 1 ml of HEPES buffer.
The single cell suspension in a 15 ml tube was centrifuged at 300 g for 4 min to collect cells in a
pellet. After removing the supernatant, the pellet was resuspended in 200 ul HEPES buffer and
cells were counted using a hemocytometer. The suspension was diluted to reach 1200 cell per

ml and was submitted to Yale Genomics Core facility for cDNA library prep using 10x 5' NEXT
GEM V2 according to the manufacturer's instructions.

NGS sequencing of single cell/ nucleus cDNA

603 The prepared 10X Genomics single cell/nucleus libraries were submitted to Yale Center for
604 Genome Analysis and sequenced on an Illumina NovaSeq sequencer with 150-bp paired-end
605 sequencing on an S4 flow cell. For each sample, ~10,000 cells were sequenced at a minimum
606 depth of 50,000 reads per cell.

Probe-panel design

IDT Lockdown probes (Integrated DNA Technologies) are 120-nt long oligos that are
biotinylated at their 5' ends and were designed using the xGen Panel Design Tool
(https://www.idtdna.com/pages/tools/xgen-hyb-panel-design-tool) to ensure robust coverage of
the targeted regions. Briefly, a customized *fasta* file containing all Ensembl[26]-annotated exons
(across all isoforms) of target genes, including UTRs was uploaded as input file. The probe
sequence list was generated at the 1x probe tilling density using "Homo sapiens (Human) NCBI
GRCh37.p13 (hg19)" as the reference (Supplementary table 1).

The probe panel was further synthesized employing TEQUILA-seq[21]. First, the probe panel pool
of 150-nt long oligos, each comprising a 3' end 30-nt universal primer binding sequence (5'-
CGAAGAGCCCTATAGTGAGTCGTATTAGAA-3') followed by a 120-nucleotide target-specific
probe, was synthesized with Twist Bioscience (Supplementary table 1). Then the oligo pools
were amplified and biotin-labeled using nickase-induced linear strand displacement
amplification as described previously. Finally, synthesized probes were purified with 1.8×
volumes of AMPure XP beads and quantified by NanoDrop 2000 Spectrophotometer.

Capture-seq on single cell/ nucleus cDNA

All hybridization and capture steps were performed following the IDT "Hybridization capture of
DNA libraries using xGen Lockdown probes and reagents" protocol. Briefly, 1-1.5 µg of
amplified 10x cDNA together with 1 µL of each blocking oligos (1 mM, Supplementary Table1)
were dried using a speed vac (Thermo, Cat# SPD140P1). Next, the dried sample were re-
hydrated with 8.5 µL of 2× Hybridization Buffer, 2.7 µl Hybridization Buffer Enhancer (xGen IDT
1080577), and 3.8 µL nuclease-free water, followed by denaturation at 95 °C for 10 min. The
denatured mixture was then incubated with 200 ng of TEQUILA probes at 65 °C for 16 h. Next,
the capture step was performed by adding 100 µL of washed M-270 streptavidin beads
(Invitrogen, #65306) to the mixture, followed by incubation at 65 °C for 45 minutes with vortexing
every 10 minutes to ensure thorough mixing. After capture, the mixture was then immediately
washed with high-temperature and room temperature buffers and resuspended in in 40 µL
nuclear-free water.

The streptavidin bead-captured cDNA was amplified using PrimeSTAR GXL premix (Takara,
Cat# R051A) and 10x cDNA amplification primers by incubating at 95 °C for 3 min, followed by
12 cycles of (98 °C for 20 s, 65 °C for 15 s, 68 °C for 8 min), with a final extension at 68 °C for
8 min. Post-capture PCR products were purified using 0.7× volumes of AMPure XP beads, and
quantified by the Qubit dsDNA High Sensitivity assay and size checked by Agilent TapeStation.

Gene-targeted single cell/ nucleus RACE-PCR

To sequence the *NRXN1α* isoforms and 10x single cell barcode from one long-read, the
modified RACE-PCR was employed to amplify the target from the 10x Genomics 3'GEX or
5'GEX cDNA. 5'RACE was used for cDNA generated by 10x 5'GEX kit, while 3'RACE was used
for cDNA generated by 10x 3'GEX kit.

PCR primer design. As the expression level of *NRXN1α* is relatively low in adult brains, nested
PCR would be needed to achieve enough molecules. The gene-specific primers (GSPs) should
meet the criteria: 23-28 nt long to ensure specific annealing, with 50%-70% GC, a T_m 65°C
(best to have a T_m >70°C) and be specific to the target gene (Supplementary Table 2). Universal
primers: The universal primer for first -round PCR should be the partial read1 sequence from
10x cDNA. To increase sensitivity for the second-round PCR, a sequence of
"AATGATACGGCGACCACCGAGATCT" was added to the 5' of the read1 (Supplementary
Table 2), forming the second-round universal PCR primer (UP2) (Supplementary Table 2).

Protocol. The input of 1-50 ng of 10x 5'GEX or 3'GEX cDNA was used for the RACE-PCR. The
first round of RACE-PCR is carried out with 1 µL partial Read1 primer (10 µM) and 4 µL *NRXN1*
-specific primer 1 (10 µM) in a 50 µl-reaction, which consists of additional 25 µL 2X PrimeSTAR
GXL premix (Takara, Cat# R051A), 1- 50 ng of full-length single cell cDNA by 10x 5'GEX or
3'GEX kit. Choose the PCR program based on the GSP T_m . Program 1: when GSP T_m >70°C,
run 5 cycles of 94°C for 30s, 72°C for 8min; 5 cycles of 94°C for 30s, 70°C for 30s and 72°C for
8min; 15-20 cycles of 94°C for 30s, 68°C for 30s and 72°C for 8min. Program 2: when T_m = 60-
70°C, run 15-20 cycles of 94°C for 30s, 68°C for 30s and 72°C for 8min. The concentration of
PCR products was quantified by Qubit, and additional cycles can be added if the concentration
of PCR product is lower than 10 ng/ µL. After the PCR, 3 µL of products were analyzed with 1%
agarose/ GelRed gel (biotium, cat# 41002), while the rest products were purified using 0.7×
volumes of AMPure XP beads and eluted with 30 µL nuclease-free water. If the primary PCR
fails to show the distinct band(s) of interest or produces a smear, a second- round of RCE-PCR
should be performed.

The second-round of RACE-PCR is performed with 2.5 µL of UP2 (10 µM) and 2.5 µL *NRXN1* -
specific primer 2 (10 µM) in a 50 µl-reaction together with 25 µL 2X PrimeSTAR GXL premix
and 10 µL of the purified first-round RACE-PCR products. The PCR program follows the
program 2 and runs for 10-15 cycles based the final concentration of PCR products. After the
PCR, 3 µL of products were analyzed with 1% agarose/ GelRed gel for size check, which is
followed by the purification using 0.7× volumes of AMPure XP beads.

Library preparation for Oxford Nanopore sequencing

Oxford Nanopore SQK-NBD114.24 was used for library preparation. Amplified cDNA from post-
capture of RACE-PCR, approximately 200 ng of PCR products from each sample were used for
library preparation according to the ONT SQK-NBD114.24 protocol. Briefly, cDNA products
were end-repaired and dA-tailed with NEBNext Ultra II End Repair/dA-Tailing Module by
incubating at 20 °C for 30 min and 65 °C for 30 min. The cDNA was then purified with 1× volume
of AMPure XP beads and eluted in 10 µl of nuclease-free water. Native barcode ligation was
then performed using 10 µl of end-prepped DNA, 2.5 µl of native barcodes, and 12.5 µl of
Blunt/TA Ligase Master Mix (NEB, #M0367) by incubating at room temperature for 30 min. After
the ligation step, 2.5 µl of EDTA was added to each tube to inactivate the ligase. Barcoded
cDNA samples were then pooled and purified using 0.4× volume of AMPure XP beads and
eluted in 31 µl of nuclease-free water. Next, ligation of Native Adapter was performed with 30 µl
of pooled barcoded samples, 5 µl of native adapter, 10 µl of 5× NEBNext Quick ligation reaction
buffer, and 5 µl of NEBNext Quick T4 DNA ligase at room temperature for 30 min, followed by

the bead's purification using 0.45× volume of AMPure XP beads and twice wash with short
fragment buffer. The libraries were quantified with Qubit, and 50 fmol of final library was
sequenced with a R10.4.1 MinION or PromethION flow cell for 72 hours.

Single-cell RNA-seq processing and analysis

Raw 10x Genomics single-cell RNA-seq data were processed using Cell Ranger (v7.2.0) with the
GRCh38 reference genome to generate gene-cell UMI count matrices. Downstream analyses
were performed in Seurat (v5.0.3)[56] under R (v4.4.0). Cells with fewer than 200 detected genes,
more than 6,000 detected genes, or mitochondrial transcript fractions exceeding 20% were
excluded. Potential doublets were identified and removed using DoubletFinder[57].

Post-quality control, gene expression values were normalized by scaling each cell's total
expression to 10,000 counts and log-transforming the results. Data were then processed using
SCTransform[56], which performs regularized negative binomial regression to normalize and correct
for sequencing depth and mitochondrial content. Highly variable genes were selected for PCA,
and dimensionality reduction was performed using UMAP.

Clustering was conducted using a shared nearest neighbor (SNN) graph and the Louvain
algorithm to identify transcriptionally distinct populations. Cell type annotation was guided by
canonical marker gene expression and cross-referenced with public human brain (or tissue-
specific) single-cell cataloges. Differential expression analyses between groups or clusters were
performed using the Wilcoxon rank-sum test with Benjamini-Hochberg correction for multiple
testing (FDR < 0.05).

Following dimensionality reduction and clustering, known marker genes were used to guide the
assignment of biological cell identities. Each cluster was annotated based on the expression of
established cell-type-specific markers reported in previous studies.

Long-read isoform sequencing analysis

Full-length transcript isoforms were characterized using long-read RNA sequencing (Oxford
Nanopore or PacBio). Raw reads were quality filtered, adapter-trimmed, and aligned to GRCh38
using minimap2 (v2.28) with spliced alignment parameters.

Isoform reconstruction and quantification were performed using IsoQuant (v3.2) with GENCODE
v43 as reference annotation. IsoQuant[58] clusters reads into high-confidence transcript models,
classifying each as known, novel in catalog (NIC), or novel not in catalog (NNC) based on splice
junction composition. Only isoforms supported by at least three full-length reads were retained.

Integration of long-read isoform data with short-read single-cell transcriptomes

Integration between long-read and short-read data was performed at both the gene and isoform
levels.To associate long-read reads with individual single cells, we applied BLAZE (v1.4.0)[59],
which accurately detects 10x cellular barcodes and unique molecular identifiers (UMIs) directly
from Nanopore cDNA reads. BLAZE searches for barcode sequences within read adapters,
compares them to the 10x whitelist, and assigns valid barcodes (allowing up to one mismatch)
to each read. This enabled recovery of cell identities from long-read data and construction of a
cell × isoform count matrix.

NRXN1 Isoform Quantification using *miniQuant*

For the BA4 samples, *miniQuant*[60] was applied in hybrid mode, integrating both short-read and long-read RNA-seq generated in-house to improve isoform-level quantification. For the FC and hippocampus datasets[25,27], where only long-read RNA-seq data were available, *miniQuant* was applied in long-read-only mode to estimate isoform abundances.

In *miniQuant*, long-read cDNA sequencing provides high-confidence isoform structures and reduces ambiguity in assigning reads to specific isoforms, whereas short-read sequencing improves quantification accuracy for low-abundance transcripts by reducing sampling error. *miniQuant*'s gene-specific weighting model adaptively balances these complementary data sources based on isoform structural complexity (K-value), transcript abundance, and sequencing depth, producing optimized isoform abundance estimates. Isoform expression levels were reported as transcripts per million (TPM) to allow direct comparison across samples.

To ensure quantification reliability, we filtered out low-confidence isoforms following the criteria described in the original *miniQuant* framework, requiring both sufficient long-read support and a minimum expression threshold. Only isoforms with TPM ≥ 1 in at least one sample were retained for downstream analyses.

Data visualization and statistical analysis

All visualizations, including UMAPs, dot plots, violin plots, and bar plots, were generated in *R* or *Python* using the *ggplot2*, *matplotlib*, and *seaborn* libraries. For cell type enrichment analyses, statistical significance was evaluated using a two-tailed Fisher's exact test, with enrichment ratios calculated as the observed versus expected frequencies. For continuous variables, such as comparisons of AS event inclusion levels or isoform expression differences, we applied *Welch's t-test*, which does not assume equal variances between groups.

ACKNOWLEDGEMENTS

We acknowledge Daniel Weinberger and the Lieber Institute for Brain Development at Johns Hopkins School of Medicine, for sharing post-mortem brain tissues. This work is supported by the National Institute of Mental Health grants RO1 MH125579 (GF and KJB) and RO1 MH121074 (KJB and GF), and U01MH116442 (PR) and R01MH125246 (PR). *Research reported in this publication was supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number 1S10OD030363-01A1.* This work was also supported in part through the computational resources and staff expertise provided by the Department of Scientific Computing at the Icahn School of Medicine at Mount Sinai.

REFERENCES

1.   Fagnani, M. *et al.* Functional coordination of alternative splicing in the mammalian
central nervous system. *Genome Biol* **8,** R108 (2007).

2.   Raj, B. & Blencowe, B. J. Alternative splicing in the mammalian nervous system:
recent insights into mechanisms and functional roles. *Neuron* **87**, 14–27 (2015).

3.   Furlanis, E., Traunmüller, L., Fucile, G. & Scheiffele, P. Landscape of ribosome-
engaged transcript isoforms reveals extensive neuronal-cell-class-specific
alternative splicing programs. *Nat Neurosci* **22**, 1709–1717 (2019).

4.   Lipscombe, D. & Soto, E. J. L. Alternative splicing of neuronal genes: new
mechanisms and new therapies. *Curr Opin Neurobiol* **57,** 26–31 (2019).

5.   Gomez, A. M., Traunmüller, L. & Scheiffele, P. Neurexins: molecular codes for
shaping neuronal synapses. *Nature Reviews Neuroscience* vol. 22 137–151 Preprint
at https://doi.org/10.1038/s41583-020-00415-7 (2021).

6.   De Wit, J. & Ghosh, A. Specification of synaptic connectivity by cell surface
interactions. *Nat Rev Neurosci* **17**, 4 (2016).

7.   Südhof, T. C. Synaptic neurexin complexes: a molecular code for the logic of neural
circuits. *Cell* **171,** 745–769 (2017).

8.   Harkin, L. F. *et al.* Neurexins 1-3 each have a distinct pattern of expression in the
early developing human cerebral cortex. *Cerebral Cortex* **27**, 216–232 (2017).

9.   The role of neurexins and neuroligins in the formation, maturation, and function of
vertebrate synapses.

10. Schreiner, D. *et al.* Targeted Combinatorial Alternative Splicing Generates Brain
Region-Specific Repertoires of Neurexins. *Neuron* **84**, 386–398 (2014).

11. Treutlein, B., Gokce, O., Quake, S. R. & Südhof, T. C. Cartography of neurexin
alternative splicing mapped by single-molecule long-read mRNA sequencing. *Proc*
Natl Acad Sci U S A **111**, (2014).

12. Flaherty, E. *et al.* Neuronal impact of patient-specific aberrant NRXN1α splicing. *Nat*
Genet **51**, 1679–1690 (2019).

13. Uemura, T. *et al.* Trans-synaptic interaction of GluRδ2 and Neurexin through Cbln1
mediates synapse formation in the cerebellum. *Cell* **141**, 1068–1079 (2010).

14.   Siddiqui, T. J. *et al*. An LRRTM4-HSPG complex mediates excitatory synapse
791   development on dentate gyrus granule cells. *Neuron* **79**, 680–695 (2013).

792   15.   Boucard, A. A., Chubykin, A. A., Comoletti, D., Taylor, P. & Südhof, T. C. A splice
793   code for trans-synaptic cell adhesion mediated by binding of neuroligin 1 to α-and β-
794   neurexins. *Neuron* **48**, 229–236 (2005).

795   16.   Ko, J., Fuccillo, M. V, Malenka, R. C. & Südhof, T. C. LRRTM2 functions as a neurexin
796   ligand in promoting excitatory synapse formation. *Neuron* **64**, 791–798 (2009).

797   17.   Fuccillo, M. V *et al*. Single-cell mRNA profiling reveals cell-type-specific expression
798   of neurexin isoforms. *Neuron* **87**, 326–340 (2015).

799   18.   Freund, T. F. & Katona, I. Perisomatic inhibition. *Neuron* **56**, 33–42 (2007).

800   19.   Fernando, M. B. *et al*. Phenotypic complexities of rare heterozygous neurexin-1
801   deletions. *Nature* 1–11 (2025).

802   20.   Lukacsovich, D. *et al*. Single-cell RNA-seq reveals developmental origins and
803   ontogenetic stability of neurexin alternative splicing profiles. *Cell Rep* **27**, 3752–3759
804   (2019).

805   21.   Wang, F. *et al*. TEQUILA-seq: a versatile and low-cost method for targeted long-read
806   RNA sequencing. *Nat Commun* **14**, (2023).

807   22.   Ray, T. A. *et al*. Comprehensive identification of mRNA isoforms reveals the diversity
808   of neural cell-surface molecules with roles in retinal development and disease. *Nat
809   Commun* **11**, (2020).

810   23.   Ohara, O., Dorit, R. L. & Gilbert, W. One-sided polymerase chain reaction: the
811   amplification of cDNA. *Proceedings of the National Academy of Sciences* **86**, 5673–
812   5677 (1989).

813   24.   Frohman, M. A., Dush, M. K. & Martin, G. R. Rapid production of full-length cDNAs
814   from rare transcripts: amplification using a single gene-specific oligonucleotide
815   primer. *Proceedings of the National Academy of Sciences* **85**, 8998–9002 (1988).

816   25.   Hardwick, S. A. *et al*. Single-nuclei isoform RNA sequencing unlocks barcoded exon
817   connectivity in frozen brain tissue. *Nat Biotechnol* **40**, 1082–1092 (2022).

818   26.   Dyer, S. C. *et al*. Ensembl 2025. *Nucleic Acids Res* **53**, D948–D957 (2025).

819   27.   Joglekar, A. *et al*. Single-cell long-read sequencing-based mapping reveals
820   specialized splicing patterns in developing and adult mouse and human brain. *Nat
821   Neurosci* **27**, 1051–1063 (2024).

28. Ruzicka, W. B. *et al*. Single-cell multi-cohort dissection of the schizophrenia
transcriptome. *Science (1979)* **384**, eadg5136 (2024).

29. Telley, L. *et al*. Sequential transcriptional waves direct the differentiation of newborn
neurons in the mouse neocortex. *Science (1979)* **351**, 1443–1446 (2016).

30. Liodis, P. *et al*. Lhx6 activity is required for the normal migration and specification of
cortical interneuron subtypes. *Journal of Neuroscience* **27**, 3078–3089 (2007).

31. Miyoshi, G. *et al*. Prox1 regulates the subtype-specific development of caudal
ganglionic eminence-derived GABAergic cortical interneurons. *Journal of*
Neuroscience **35**, 12869–12889 (2015).

32. Tremblay, R., Lee, S. & Rudy, B. GABAergic interneurons in the neocortex: from
cellular properties to circuits. *Neuron* **91**, 260–292 (2016).

33. Cardin, J. A. Inhibitory interneurons regulate temporal precision and correlations in
cortical circuits. *Trends Neurosci* **41**, 689–700 (2018).

34. Huang, Z. J. & Paul, A. The diversity of GABAergic neurons and neural
communication elements. *Nat Rev Neurosci* **20**, 563–572 (2019).

35. Liakath-Ali, K. & Südhof, T. C. The Perils of Navigating Activity-Dependent Alternative
Splicing of Neurexins. Front. Mol. Neurosci. 14, 659681. Preprint at (2021).

36. Baxter, P. S., Dando, O. & Hardingham, G. E. Differential splicing choices made by
neurons and astrocytes and their importance when investigating signal-dependent
alternative splicing in neural cells. *Front Mol Neurosci* **16**, 1214439 (2023).

37. Ramos, S. I. *et al*. An atlas of late prenatal human neurodevelopment resolved by
single-nucleus transcriptomics. *Nat Commun* **13**, 7671 (2022).

38. Wang, L. *et al*. Molecular and cellular dynamics of the developing human neocortex.
Nature 1–10 (2025).

39. Zhu, K. *et al*. Multi-omic profiling of the developing human cerebral cortex at the
single-cell level. *Sci Adv* **9**, eadg3754 (2023).

40. Eze, U. C., Bhaduri, A., Haeussler, M., Nowakowski, T. J. & Kriegstein, A. R. Single-
cell atlas of early human brain development highlights heterogeneity of human
neuroepithelial cells and early radial glia. *Nat Neurosci* **24**, 584–594 (2021).

41. Sebastian, R. *et al*. Schizophrenia-associated NRXN1 deletions induce
developmental-timing-and cell-type-specific vulnerabilities in human brain
organoids. *Nat Commun* **14**, 3770 (2023).

42.   Birey, F. *et al*. Assembly of functionally integrated human forebrain spheroids. *Nature* **545**, 54–59 (2017).

43.   Trujillo, C. A. *et al*. Complex oscillatory waves emerging from cortical organoids model early human brain network development. *Cell Stem Cell* **25**, 558–569 (2019).

44.   Fair, S. R. *et al*. Electrophysiological maturation of cerebral organoids correlates with dynamic morphological and cellular development. *Stem Cell Reports* **15**, 855–868 (2020).

45.   Park, Y. *et al*. Modulation of neuronal activity in cortical organoids with bioelectronic delivery of ions and neurotransmitters. *Cell Reports Methods* **4**, (2024).

46.   Matsunami, N. *et al*. Identification of rare recurrent copy number variants in high-risk autism families and their prevalence in a large ASD population. *PLoS One* **8**, e52239 (2013).

47.   Kim, H.-G. *et al*. Disruption of neurexin 1 associated with autism spectrum disorder. *The American Journal of Human Genetics* **82**, 199–207 (2008).

48.   Kozareva, V. *et al*. A transcriptomic atlas of mouse cerebellar cortex comprehensively defines cell types. *Nature* **598**, 214–219 (2021).

49.   Uemura, T. *et al*. Trans-synaptic interaction of GluRδ2 and Neurexin through Cbln1 mediates synapse formation in the cerebellum. *Cell* **141**, 1068–1079 (2010).

50.   Peng, H. *et al*. Single-cell Rapid Capture Hybridization sequencing reliably detects isoform usage and coding mutations in targeted genes. *Genome Res* **35**, 942–955 (2025).

51.   Lim, L., Mi, D., Llorca, A. & Marín, O. Development and functional diversification of cortical interneurons. *Neuron* **100**, 294–313 (2018).

52.   Nguyen, T.-M. *et al*. An alternative splicing switch shapes neurexin repertoires in principal neurons versus interneurons in the mouse hippocampus. *Elife* **5**, e22757 (2016).

53.   Biswas, M. S., Roy, S. K., Hasan, R. & Pk, M. M. U. The crucial role of the cerebellum in autism spectrum disorder: neuroimaging, neurobiological, and anatomical insights. *Health Sci Rep* **7**, e2233 (2024).

54.   van der Heijden, M. E., Gill, J. S. & Sillitoe, R. V. Abnormal cerebellar development in autism spectrum disorders. *Dev Neurosci* **43**, 181–190 (2021).

885 55. D'Mello, A. M. & Stoodley, C. J. Cerebro-cerebellar circuits in autism spectrum
886   disorder. *Front Neurosci* **9**, 408 (2015).

887 56. Hao, Y. *et al*. Dictionary learning for integrative, multimodal and scalable single-cell
888   analysis. *Nat Biotechnol* **42**, 293–304 (2024).

889 57. McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: doublet detection in
890   single-cell RNA sequencing data using artificial nearest neighbors. *Cell Syst* **8**, 329–
891   337 (2019).

892 58. Prjibelski, A. D. *et al*. Accurate isoform discovery with IsoQuant using long reads. *Nat
893   Biotechnol* **41**, 915–918 (2023).

894 59. You, Y. *et al*. Identification of cell barcodes from long-read single-cell RNA-seq with
895   BLAZE. *Genome Biol* **24**, 66 (2023).

896 60. Li, H. *et al*. Improving gene isoform quantification with miniQuant. *Nat Biotechnol* 1–
897   13 (2025).

An Integrated Targeting Strategy Reveals Cell-Type Atlas of *NRXN1* Isoforms Across Human Brain Development and Genetic Backgrounds

a

Percentage of *NRXN1* reads by cell type with/ without Capture-seq

Fold enrichment of *NRXN1* reads by cell type

b

Workflow for comprehensive study of cell-type specific *NRXN1* isoforms

c

A schematic of the *NRXN1* isoforms identified across samples by single-cell long-read Capture-seq

NRXN1 expression repertoires exhibit distinct splicing patterns across cell types in human brains

a   Resolve cell-type specific *NRXN1* isoforms by probe-based long-read Capture-seq

b   UMAP visualization of cell types in the PFC of adult brains

d   PCA plot of *NRXN1* AS events across cell types

c   Cell-type-specific *NRXN1* splicing patterns of SS1-6 revealed by long-read Capture-seq

NRXN1 splicing profiles remain consistent across cells sharing a neurogenic origin during neural maturation

NRXN1⁺/⁻ hCOs splicing profiles recapitulate developmental and mature features and identify cell types expressing the mutant isoforms

a UMAP visualization of cell types in organoids

b Expression of marker genes used for cell type annotation

c Cell type enrichment for cells in 3'del organoids vs control

d *NRXN1* Alternative splicing levels in control organoids (upper) and 3'del organoids (lower)

e *NRXN1* exon inclusion level across splicing sites between between 3'del and control lines

f *NRXN1α* isoforms identified from organoids by Long-range RACE-seq

g Composition of *NRXN1* isoforms within each cell type

h Exon inclusion levels at splice sites (SS1–SS6) for GABA-IN, Glu-EX, and RG/AC across hCOs, fetal neocortex, and adult PFC

Unique exonic deletion impacts *NRXN1* splicing patterns across cell types in cerebellum of an ASD case

a UMAP visualization of cell types in adult cerebellum

b Expression of marker genes used for cell type annotation

c Catalog of *NRXN1* isoforms (WT and MT) by long-read capture-seq in cerebellum

d Read count for transcripts

e Mutant isoforms enriched in ML1/2 and astroglia

f AS patterns of *NRXN1* in cerebellum across cell types

g *NRXN1* binding ligands expression in granule cells with WT or MT *NRXN1*