1   **Title: Dynamic coupling between visual landmark processing in area 29e and parahippocampal**
2   **path integration circuits of the rodent cortex**
3

Authors: Gorkem Secer[1,2,3,#], Bharath Krishnan[2,3,4,#], Noah J. Cowan[1,3,5,*,†] and James J. Knierim[2,3,6,*,†]

Affiliations:

[1]Laboratory for Computational Sensing and Robotics, Johns Hopkins University; 3400 N. Charles St., Baltimore, MD 21218,
USA

[2]Zanvyl Krieger Mind/Brain Institute, Johns Hopkins University; 3400 N. Charles St., Krieger Hall 338, Baltimore, MD
21218, USA

[3]Kavli Neuroscience Discovery Institute, Johns Hopkins University; 3400 N. Charles St., Clark Hall 316, Baltimore, MD
21218, USA

[4]Department of Biomedical Engineering, Johns Hopkins University; 720 Rutland Avenue., Baltimore, MD 21205, USA

[5]Department of Mechanical Engineering, Johns Hopkins University; 3400 N. Charles St., Latrobe Hall 223, Baltimore, MD
21218, USA

[6]Solomon H. Snyder Department of Neuroscience, Johns Hopkins University; 725 N. Wolfe St., Wood Basic Science
Building 1003, Baltimore, MD 21205, USA

#These authors contributed equally to this work.

†These authors jointly supervised this work.

*Corresponding author. Email: ncowan@jhu.edu, jknierim@jhu.edu
20

Abstract:
External landmarks anchor internal spatial representations. The neural mechanisms underlying
this process and the sources of landmark signals remain unknown. We recorded neurons across
five parahippocampal regions while rats navigated in a virtual reality apparatus, inducing conflict
between self-motion and landmark cues. Area 29e, a parahippocampal field putatively
homologous to primate *area prostriata*, maintained firing coupled to landmarks even when other
regions decoupled from the landmarks. This decoupling was preceded by a decline of gamma-
band influence from 29e to MEC and a concomitant decline of theta-modulated feedback from
MEC to 29e. Area 29e also displayed weak theta modulation, strong gamma rhythmicity, strong
egocentric head-direction tuning, and enhanced landmark contrast sensitivity. These findings
provide strong evidence that 29e serves as a specialized visuospatial hub that provides landmark
signals for anchoring parahippocampal spatial representations to the external world.

33
Main Text:
Many higher-order neural systems display internal dynamics and represent explicit cognitive states in
the absence of overt sensory stimulation. Sensory input does not drive a "blank slate" of quiescent
neurons waiting for input about the external world. Rather, such input modulates a rich set of internally
generated representations that perform important neural computations that are critical for the proper
functioning of the system. Understanding the interplay between external sensory input and endogenous
neural dynamics is key toward understanding the principles of neural computation. Spatial navigation
provides an ideal model for dissecting this interplay. The hippocampal-parahippocampal system,
coordinated by endogenous neural oscillations spanning multiple frequency bands (*1–5*), continuously
updates a position signal on the animal's cognitive map by integrating self-motion-generated cues over
time—a process known as path integration (*6–9*). To prevent error from accumulating in this

incremental estimate of the animal's position, stable landmarks must be utilized to anchor the map to an
external reference frame (*10–14*).

Although the neural correlates of path integration (e.g., grid cells and speed cells) are relatively well
characterized in rodent hippocampal and parahippocampal areas (*9, 15–25*), the neural pathways that
convey visual landmark signals that calibrate and correct this internal computation remain unclear. Area
29e is a relatively understudied region interposed between the dorsal presubiculum/postsubiculum (PR)
and the parasubiculum (PA). Recent work suggests that rodent area 29e is homologous with the primate
area prostriata, which has been implicated in processing stimuli in the peripheral visual field (*26–28*).
Area 29e receives direct inputs from the dorsal lateral geniculate nucleus and V1 and projects to PR and
medial entorhinal cortex (MEC; *29–32*), identifying it as a potential candidate relay of visual landmark
signals to parahippocampal circuits in rodents. Using a virtual reality system, which allowed self-motion
signals to be decoupled from visual landmarks (*12, 33, 34*), and a square open-field arena, we found that
29e contained a heterogeneous population of neurons displaying both spatial and visual firing correlates.
One subpopulation had directionally tuned response curves and firing that remained tightly coupled to
distal visual landmarks under conditions in which the firing fields of neurons in the other regions
became decoupled from these landmarks. Area 29e showed a large proportion of egocentric head-
direction neurons, consistent with retina-centered visual drive, and stronger sensitivity to landmark
contrast compared to other regions. Furthermore, we found strong gamma-mediated feedforward
communication from 29e to MEC and theta-mediated feedback communication from MEC to 29e.
Critically, this feedforward–feedback coupling became disrupted when the landmarks lost control of the
MEC spatial firing. Our findings provide a physiological and functional characterization of area 29e in
freely moving animals. They reveal the neural dynamics through which the parahippocampal system
integrates visual landmark information with internal path-integration computations in a temporally
coordinated loop between visual cortex and the brain's cognitive mapping system.

Results

Area 29e showed spatially correlated activity that was more tightly bound to visual landmarks than
neighboring parahippocampal areas

We performed simultaneous tetrode recordings in area 29e and four other parahippocampal areas (MEC,
PA, PR, and postrhinal cortex [PO]; Fig. 1A and fig. S1-2) as rats (n = 5) navigated freely in our 'Dome'
virtual-reality apparatus, which allowed for precise control of the locations and salience of visual
landmarks during untethered physical locomotion (*12, 33*; Fig. 1B). When the landmarks were
stationary, a comparable proportion of 29e neurons displayed significant spatially correlated activity as
in the other parahippocampal regions (29e: 70/78, MEC: 257/328, PA: 178/232, PO: 87/112, PR: 32/44;
chi-squared test χ^2 (4,796) = 7.64, p = 0.105). Many spatially correlated 29e neurons increased or
decreased their firing rates selectively in the vicinity of these stationary landmarks (Fig. 1C). We defined
two metrics to quantify the strength and between-landmark similarity of such peri-landmark firing-rate
changes: (1) the *z-index* was computed as the mean absolute difference between z-scored firing rates at
track locations closest to landmarks and those midway between landmarks and (2) the *local similarity*
index was computed as the correlation of the firing rate profile of a neuron in the neighborhood of one
landmark with its firing rate near a second landmark, averaged across all pairs of landmarks for a given
neuron. Both metrics were significantly larger in spatially correlated neurons in 29e than those in other
regions (Fig. 1D).

Considering this modulation of 29e neurons' spatially correlated firing near landmarks, we next tested
the extent to which their firing fields were dynamically controlled by landmarks when the entire
landmark constellation was rotated around the track in a behaviorally closed-loop fashion. The rotation
of landmarks was controlled by an experiment gain (G; Fig. 1B) that set the ratio between the rat's travel

distance in the landmark frame of reference to the travel distance in the laboratory frame of reference
(*12*). When G < 1, the landmark array was rotated around the center of the track in the same direction as
the rat, providing the rat with the illusion that it was moving slower than it really was; when G > 1, it
was rotated in the opposite direction, providing the rat with the illusion that it was moving faster than it
really was. In each session, the value of G was initialized at 1 (stationary landmarks), gradually ramped
to its final value, and then held constant at that final gain value for the remainder of the session.

Under these manipulations, spatially correlated neurons across all regions often maintained stable firing
fields in the rotating landmark frame of reference across laps—a manifestation of landmark control over
the internal representations of space in the Dome (Fig. 1E and fig. S3A), as previously demonstrated for
CA1 place cells (*12*) and anterodorsal thalamic nuclei head direction cells (*35*). This lap-to-lap stability,
however, was disrupted in some sessions, typically during the gain-ramp epoch and persisting until the
end of the experiment (See Fig. 1F and fig. S3B for example sessions). To quantify the lap-to-lap
stability of each neuron's firing fields in the rotating landmark frame of reference, we defined a
landmark-locking score based on the spatial autocorrelogram (up to a 9-lap lag) of the neuron's activity
during the last 12 laps of each session (see Methods). A landmark-locking score close to 1 indicates
highly stable lap-to-lap firing of a neuron in the landmark frame (e.g., Fig. 1E bottom inset, all regions),
whereas a score near 0 indicates a lack of this stability (e.g., Fig. 1F bottom inset, non-29e regions).
Averaging landmark-locking scores across all spatially correlated neurons across all regions in each
session (number of neurons recorded per session: mean ± s.d. = 33.0 ± 17.2) revealed a bimodal
distribution (Fig. 1G). An unsupervised clustering algorithm identified two clusters within this
distribution. We refer to the cluster of 13 sessions with higher landmark-locking scores (mean ± s.d. =
0.68 ± 0.08) as *LM-control* sessions, indicating reliable control of the landmarks over firing fields of
spatially correlated neurons from all regions (as in Fig. 1E). By contrast, the remaining cluster of 21
sessions with lower landmark-locking scores (0.27 ± 0.13) are termed *LM-failure* sessions, indicating
the failure of this landmark control over most spatially correlated neurons, particularly those outside 29e
(as in Fig. 1F).

Area 29e neurons were more robustly controlled by the landmarks than the other parahippocampal
regions. In *LM-control sessions*, landmark-locking scores of spatially correlated neurons were strongly
skewed towards 1 in all regions (Fig. 1H, left). In *LM-failure sessions*, however, landmark-locking
scores of spatially correlated neurons in MEC, PA, PO, and PR were skewed towards 0, while those in
29e remained skewed toward 1, indicating greater anchoring of 29e neurons to landmarks (Fig. 1H,
right). Comparing mean landmark-locking scores between simultaneously recorded neurons from 29e
and other regions ensured that these regional differences were not due to session-specific effects or non-
uniform sampling (Fig. 1I). This greater landmark-anchoring by 29e neurons persisted even when
landmark rotation was independent of the rat's movement during open-loop gain manipulations (fig. S4;
see Methods). Collectively, these data suggest that, compared to neighboring parahippocampal regions
specialized for spatial processing with greater reliance on path integration, area 29e may be a
visuospatial region that codes for spatial representations anchored to visual landmarks.

129
Fig. 1. Landmark-anchored spatial tuning in 29e compared to spatial tuning in neighboring parahippocampal regions (MEC,
PA, PO, and PR).
(A) Representative histology section (Nissl-stained, sagittal) showing the recorded parahippocampal regions. Colored
arrowheads demarcate the superficial-layer borders of each region. The same color scheme is used throughout all subsequent
figures. See fig. S1-2 for sections showing actual recording locations within each region.
(B) Schematic of the Dome virtual-reality apparatus and experimental gain manipulation. Top: A large opaque hemispherical

shell (rendered semi-transparent to visualize the interior) displayed visual landmarks. Inside, the rat ran circular laps around
the perimeter of a table. A rotating central pillar carried three radial boom arms: a front arm with a reward spout, a back arm
that guided the recording tether, and a cleaning arm (hardware not shown). Short clear walls attached to the feeding and
tether booms moved with the rat, maintaining a constant enclosure around it. Components that could otherwise provide
angular cues either co-rotated with the animal or were circularly symmetric, making the projected landmarks the dominant
polarizing cues for localization. Bottom left: Illustration of experimental gain manipulation, where visual landmarks were
moved as a function of the rat's movement, controlled by the experimental gain (G; see main text).
(C) Circular rate maps of four example, spatially correlated 29e neurons, with red-to-blue color gradient indicating
maximum-to-minimum firing rates. Within each rate map, rat ID and the tetrode number from which the neuron was recorded
are shown. Dashed lines mark the angular positions of the centers of the three landmarks projected onto the Dome. The first
row shows two neurons that decrease their firing near landmarks, while the second row shows two neurons that increase their
firing near landmarks.
(D) Cumulative distributions of z-index (left) and local-similarity index (right), which quantify the strength and uniformity of
landmark selective firing, respectively. The z-index was significantly larger in 29e than other regions (29e median [IQR]:
0.60 [0.24, 1.02], MEC = 0.33 [0.16, 0.54], PA = 0.40 [0.22, 0.72], PO = 0.44 [0.21, 0.60], PR = 0.33 [0.14, 0.54]; Kruskal-
Wallis test $\chi^2(4) = 22.2$, $p < 0.001$; post-hoc Wilcoxon rank-sum tests with Holm-Bonferroni correction 29e vs. MEC: $Z =$
4.18, $p < 0.001$; 29e vs. PA: $Z = 2.70$, $p = 0.013$; 29e vs. PO: $Z = 2.38$, $p = 0.017$; 29e vs. PR: $Z = 2.87$, $p = 0.012$). The
local-similarity index was also larger in 29e than other regions (29e median [IQR]: 0.05 [–0.24, 0.28], MEC = –0.23 [–0.31,
0.00], PA = –0.23 [–0.32, –0.02], PO = –0.23 [–0.30, –0.01], PR = –0.30 [–0.33, –0.06]; Kruskal-Wallis test $\chi^2(4) = 22.3$, p
< 0.001; post-hoc Wilcoxon rank-sum tests with Holm-Bonferroni correction 29e vs. MEC: $Z = 3.68$, $p < 0.001$; 29e vs. PA:
$Z = 4.06$, $p < 0.001$; 29e vs. PO: $Z = 3.14$, $p < 0.001$; 29e vs. PR: $Z = 3.75$, $p < 0.001$).
(E) Representative closed-loop gain manipulation session showing landmark control across all regions. The top row shows
the experimental gain as a function of laps in the landmark frame (x-axis). The session began with a gain of 1 (i.e., stationary
landmarks). The gain was then gradually ramped to its target value, during which the landmark array rotated at increasing
speeds opposite to the animal's movement direction. The final gain was held constant for the duration of the session. The
bottom row shows the lap-to-lap (with respect to landmarks) firing rate as a function of the animal's angle in the landmark
frame (y-axis) from example neurons, color-coded by region (as in panel A). The intensities were normalized to the
maximum firing rate of the given neuron. In this session, example neurons from 29e (black), MEC (orange), PA (blue) and
PR (green) illustrate that the firing locations of all spatially correlated neurons remained anchored to the rotating landmarks
throughout the session (i.e., they fired at a consistent angle on the track [y-axis] across laps). Inset shows the spatial
autocorrelograms of each representative neuron's activity in the rotating landmark frame, calculated using the last 12 laps for
lags from 0 to 9 laps (ensuring a minimum of three lap-samples per lag for reliable correlation estimates). Note the lap-to-lap
periodicity in the autocorrelation traces for these neurons.
(F) Representative session where most spatially correlated neurons—except in 29e—lost their anchoring during the gain
ramp (same format as panel E). After about lap 20, the firing fields of the MEC (orange), PA (blue), and PO (pink) neurons
became increasingly diffuse. The 29e neuron (black), by contrast, maintained stable firing in the landmark frame throughout
the session. Inset shows the spatial autocorrelograms (calculated as in panel E) of each neuron's activity in the rotating
landmark frame. Note that the lap-to-lap periodicity for the neurons that lost landmark anchoring showed disrupted
autocorrelation structures: either no clear periodicity (e.g., the orange neuron) or a non-unity periodicity (e.g., the blue and
pink neurons), reflecting the continued firing of the neuron at repeated intervals different than a single, landmark-based lap.
(G) Session-level bimodality of neuronal populations' average landmark-locking scores. Sessions were classified as *LM-
failure* (shaded bars) *or LM-control* (unshaded bars).
(H) Distributions of landmark-locking scores for individual spatially correlated neurons. *LM-control sessions* (left) show
statistically comparable landmark-locking scores across all regions (median [IQR]: 29e = 0.73 [0.39, 0.90], MEC = 0.81
[0.52, 0.92], PA = 0.76 [0.46, 0.88], PO = 0.66 [0.39, 0.85], PR = 0.89 [0.23, 0.97]; Kruskal-Wallis test $\chi^2(4) = 4.2$, $p =$
0.376). By contrast, *LM-failure sessions* (right) show significantly larger scores in 29e than other regions (median [IQR]: 29e
= 0.50 [0.25, 0.77], MEC = 0.19 [0.06, 0.41], PA = 0.22 [0.07, 0.50], PO = 0.15 [0.03, 0.34], PR = 0.20 [0.04, 0.44]; Kruskal-
Wallis test $\chi^2(4) = 31.4$, $p < 0.001$; post-hoc Wilcoxon rank-sum tests with Holm-Bonferroni correction 29e vs. MEC: $Z =$
5.13, $p < 0.001$; 29e vs. PA: $Z = 4.11$, $p < 0.001$; 29e vs. PO: $Z = 4.74$, $p < 0.001$; 29e vs. PR: $Z = 3.08$, $p < 0.001$).
(I) Mean landmark-locking scores of spatially correlated 29e neurons within a session (y-axis) versus those of their
simultaneously recorded counterparts in other regions (x-axis) within the same session. In *LM-control* sessions (n = 7), mean
landmark-locking scores did not significantly differ between 29e (mean ± s.e.m. = 0.62 ± 0.04) and other regions (0.66 ±
0.02), whereas in *LM-failure sessions* (n = 8), 29e (0.56 ± 0.05) showed significantly higher mean landmark-locking scores
than other regions (0.21 ± 0.03). In *LM-control sessions* (n = 7; unfilled circles), the normalized difference between scores of
neurons inside 29e vs. outside 29e (i.e., (inside – outside)/(inside + outside)) was not significantly different than zero (one-
sample t-test $t(6) = 0.964$, $p = 0.372$), whereas in *LM-failure* sessions (filled circles), the normalized difference was
significantly positive (one-sample t-test $t(7) = 6.93$, $p < 0.001$), indicating greater anchoring to landmarks in 29e than in other

regions. *LM-failure* sessions showed significantly larger normalized differences than *LM-control* sessions (t-test *LM-failure*
vs. LM-control: $t(13) = 5.7$, $p < 0.001$).

Area 29e was sensitive to sensory aspects of visual landmarks

We next investigated the sensitivity of area 29e to sensory aspects of the landmarks in comparison to
neighboring parahippocampal areas. We first tested the immediate effect of turning off the landmarks
following the gain manipulations (i.e., immediately after Epoch 3). Because the effects of extinguishing
the landmarks would be confounded at the single-unit level by where the rat was located when the
landmarks were turned off (i.e., whether the rat was in a spatial firing field of a particular cell), we
quantified the responses to landmark removal using multi-unit firing rates in five animals. For each
tetrode, we computed a landmark-removal response score as the absolute normalized change in its firing
rate from the 600 ms period immediately preceding landmark removal to the 400 ms immediately
following it (see Methods). Larger positive scores indicate stronger firing-rate changes, regardless of the
direction, whereas scores near zero indicate little or no change. Only tetrodes in 29e and PO showed a
large majority of positive landmark-removal response scores, with 29e exhibiting the largest scores on
average (Fig. 2A). A linear mixed-effects model accounting for dependencies among tetrodes from the
same animal and brain region revealed that 29e showed significantly higher scores than MEC and PA,
but it was not significantly different from PR or PO.

To determine whether individual 29e neurons were tuned to the luminance or contrast of the landmarks,
we performed additional experiments in a subset of animals (n = 2), recording from 29e, MEC, and PR
(no tetrodes were in PA or PO in these animals). As rats ran laps, we pseudo-randomly varied landmark
and background luminance between two levels (i.e., high or low) across six-lap blocks, with the
experimental gain fixed at 1 to keep landmark locations stationary (Fig. 2B). Similar proportions of
neurons with significant luminance sensitivity were identified across all recorded regions (29e: 28/45;
MEC: 14/24; PR: 9/25; chi-squared test $\chi^2(2,94) = 4.66$, p = 0.096). However, the magnitude of
luminance sensitivity was significantly larger in neurons from 29e than those from MEC and PR (Fig.
2C). This heightened effect of landmark and background luminance on 29e single-neuron activity may
reflect their tuning to the total luminance in the environment or to the contrast between the landmark and
background luminance. For each neuron with significant landmark- or background-luminance
sensitivity, we fit a linear model to the peak firing rates on each lap, using landmark contrast and total
luminance as predictors. Two-way ANOVA (region × predictor) on the partial R^2 values—which
quantify the firing-rate variance uniquely explained by each predictor—revealed particularly
pronounced tuning of 29e neurons to landmark contrast, the strength of which significantly exceeded not
only contrast tuning in other regions but also total luminance tuning in any region (Fig. 2D).

Following the luminance-change epochs, we performed closed-loop gain manipulations with large gains
to induce landmark-control failure. Spatially correlated neurons of 29e showed a significant positive
relationship between the strengths of their landmark-contrast tuning and their landmark anchoring (Fig.
2E; fig. S5).

Fig. 2. Greater sensitivity of area 29e to sensory aspects of visual landmarks compared to neighboring parahippocampal areas.

(A) Cumulative distributions of landmark-removal response scores from unique tetrode locations (29e: n = 13, MEC: n = 87, PA: n = 54, PO: n = 19, PR: n = 16; colors as in Panel A). Response scores of tetrodes were significantly positive in 29e (median [IQR] = 0.096 [0.071, 0.129]; Wilcoxon signed-rank test $Z = 2.97$, $p = 0.003$) and PO (0.049 [–0.001, 0.139]; $Z = 2.77$, $p = 0.005$) but not in MEC (0.008 [–0.054, 0.082]; $Z = 1.35$, $p = 0.176$), PA (0.009 [–0.068, 0.061]; $Z = 0.20$, $p = 0.842$), or PR (0.018 [–0.047, 0.053]; $Z = 0.72$, $p = 0.469$). Examining distribution of scores with a linear mixed-effects model showed significant differences across regions ($F(165.2) = 3.24$, $p = 0.014$; see Methods for random-effects structure of the model and further details). The model also showed that 29e exhibited significantly higher scores than MEC ($t(151.3) = 2.4$, $p = 0.018$) and PA ($t(138.0) = 3.2$, $p = 0.002$), but not PR ($t(153.3) = 1.22$, $p = 0.222$) or PO ($t(175.6) = 1.56$, $p = 0.119$).

(B) Protocol for luminance manipulation experiments and an example of a contrast-responsive neuron. Top: Experiment design. Bottom: Example neuron from area 29e showing peak lapwise firing rate changes (black) that more closely followed changes in landmark contrast (C = LM - BG, solid purple) rather than total luminance (TL = LM + BG, dashed purple). Firing rates are normalized to the neuron's peak firing rate in the first lap of the session.

(C) Region-wise cumulative distributions of total η^2 values from two-way ANOVA with background and landmark luminance as factors. Total η^2 was significantly larger in 29e neurons than in MEC and PR neurons (median [IQR] 29e = 0.14 [0.04, 0.32], MEC = 0.06 [0.03, 0.09], PR = 0.04 [0.01, 0.08]; Kruskal-Wallis test $\chi^2(2) = 13.95$, $p < 0.001$; post-hoc Wilcoxon rank-sum tests with Holm-Bonferroni correction 29e vs. MEC: $Z = 2.61$, $p = 0.008$; 29e vs. PR: $Z = 3.30$, $p = 0.001$).

(D) Cumulative distributions of partial R^2 values from linear models predicting neurons' lapwise peak firing rates using landmark contrast (C, solid lines) and total brightness (TL, dashed lines) as predictors. Two-way ANOVA on sqrt-transformed partial R^2 values revealed significant main effects of region ($F(2,96) = 16.3$, $p < 0.001$) and predictor type ($F(1,96) = 7.5$, $p = 0.007$), as well as a significant interaction ($F(2,96) = 9.3$, $p < 0.001$). Post-hoc tests (with Holm-Bonferroni corrections) showed that partial R^2 for landmark contrast in 29e (mean ± s.e.m. = 0.28 ± 0.03) was significantly greater than landmark-contrast partial R^2 in MEC (0.05 ± 0.01) and PR (0.06 ± 0.01), and total-brightness partial R^2 in all recorded regions (29e: 0.07 ± 0.01; MEC: 0.04 ± 0.01; PR: 0.06 ± 0.01); all pairwise comparisons with landmark contrast in 29e: $p < 0.001$.

(E) Significantly positive correlation between the partial R^2 (y-axis) for landmark-contrast tuning of 29e neurons with spatially and their landmark-locking scores (x-axis; Pearson's r = 0.36, $p = 0.027$; landmark-locking scores were normalized with the Fisher-z transform). Each dot corresponds to one neuron.

Area 29e demonstrated tuning to egocentric and allocentric head direction, with egocentric tuning more closely associated with sensory-like responses to landmarks

To further characterize spatially correlated activity in 29e in comparison to neighboring parahippocampal areas, we analyzed single-unit activity obtained as the rats (n = 5) foraged in an open arena after they completed each Dome session (fig. S6A). The arena was in a separate room from the

Dome. We examined allocentric tuning (Fig. 3A; fig. S6B) by fitting a generalized linear model (GLM)
that jointly captures both place and head-direction (HD) tuning of a neuron, thereby minimizing biases
due to behavioral confounds such as uneven sampling of HDs in different locations (*36*, *37*). From the
GLM-derived 2D positional rate map and allocentric HD tuning curve of each neuron, we then
calculated the Skaggs' spatial information (SI) score (*38*) and the Rayleigh score (*39*, *40*), respectively.
SI scores were significantly lower in area 29e than other regions except PR (Fig. 3B top row). By
contrast, Rayleigh scores of allocentric HD tuning curves were significantly larger in area 29e than other
regions (Fig. 3B middle row).

Classic, allocentric HD cells fire maximally in a consistent direction regardless of the animal's location,
yielding parallel preferred HD vectors across the arena (*37*, *41*; fig. S7A). In contrast, egocentric HD
cells demonstrate progressive, location-dependent shifts in their preferred HD, forming vector fields
with rotational, radial (outward/inward-pointing), or hybrid patterns (*42–45*), consistent with tuning for
the egocentric bearing of the head relative to an external anchor point. For example, a cell that fires
whenever a specific location is to the right of the animal's head, regardless of the animal's allocentric
position, would show a vector field with a clockwise rotational flow about that location, and a cell that
fires whenever that location is directly ahead of the rat would show a centripetal, radial flow. Many 29e
neurons displayed preferred HD-vector fields consistent with egocentric HD tuning (Fig. 3C; see also
fig. S7B-C for more examples). To quantify egocentric HD tuning, we developed an egocentric vector-
field (EVF) score that measures the consistency of rotational and radial egocentric patterns within a
neuron's HD vector field across the arena, without relying on an exhaustive search to identify an optimal
anchor point in the environment, as was done in previous studies (*43*, *46–48*) and which introduces
biases in comparing egocentric vs. allocentric tuning curves (*43*; fig. S8; see Methods). The EVF score
ranges from 0 (indicating no egocentric tuning, such as when preferred HD vectors are randomly
oriented or parallel everywhere) to 1 (indicating maximal egocentric tuning). Across all neurons, the
EVF scores had significantly higher values in 29e than other regions (Fig. 3B bottom row).

We used a combination of EVF, Rayleigh, and SI scores, together with a within-session tuning stability
requirement, to classify each cell as an allocentric HD cell; an egocentric HD cell; a nondirectional,
spatially correlated cell; or a nonspatial cell (see Methods). Egocentric HD-tuned neurons were
significantly more prevalent in 29e than other regions (Fig. 3D). Nondirectional place-tuned neurons
were rare in 29e, at a proportion significantly lower than in MEC and PA, but not PO or PR. By contrast,
the prevalence of allocentric HD-tuned neurons was similar across regions.

Because visual sensory cues are initially encoded in egocentric (i.e., retinocentric) frames before being
converted to an allocentric frame in higher-order areas, we hypothesized that egocentric HD-tuned
neurons would be more closely associated with encoding visual landmarks than allocentric ones. Indeed,
compared to 29e neurons classified as allocentric in the open arena, egocentric neurons exhibited, in the
Dome, significantly higher landmark-locking scores (Fig. 3E; fig S9A) and a trend toward greater
sensitivity to landmark contrast (fig. S9B).

Given that egocentric 29e neurons exhibited properties consistent with visual-landmark coding, we
hypothesized that these neurons would be anchored to locations outside the open arena, where salient
landmarks were located. We computed for each egocentric neuron the shortest signed distance from the
arena boundary to its optimal anchor point estimated by maximizing the negative log-likelihood of a
GLM. A negative distance indicates an anchor point inside the arena, while a positive distance indicates
one outside. Median signed distance was significantly positive in 29e, indicating anchoring to locations
beyond the arena boundary. By contrast, the medians were negative in other regions, with only MEC
reaching statistical significance. Cross-regional comparison of signed distance distributions confirmed
that anchoring to locations outside the arena is a distinguishing feature of egocentric neurons in 29e

compared to egocentric neurons of other regions (Fig. 3F). We further investigated the anchoring properties of egocentric 29e neurons by analyzing their preferred bearing angles estimated as described in fig. S8. The distribution of bearing angles showed a significant bias that was contralateral to the recording hemisphere (right hemisphere), with most anchor points concentrated in the animal's left-to-back-left sector, corresponding to the peripheral visual field (Fig. 3G). Supplementary analyses of other regions also showed a contralateral preference in MEC and PA, but not in PO and PR, perhaps due to low statistical power (fig. S9C).

Fig. 3. Two-dimensional spatial representations of Area 29e neurons compared to neurons in neighboring parahippocampal regions (MEC, PA, PO, and PR).

(A) Allocentric tuning of three example neurons from each region. For each neuron, a 2D positional rate map (top) and a 1D allocentric head-direction (HD) tuning curve (bottom) are shown. Rat ID and tetrode number for each neuron are shown above its rate map. The number inset in each plot denotes the peak-firing rate. The y-axis of each HD plot starts at zero.

(B) Region-wise cumulative distributions of SI score (top), Rayleigh score (middle), and egocentric vector field (EVF) score (bottom). All scores differed significantly across regions:

SI scores were significantly lower in 29e (median [IQR] = 0.019 [0.010, 0.045]) than MEC (0.034 [0.016, 0.083]), PA (0.036
[0.016, 0.120]), and PO (0.027 [0.018, 0.054]), but not PR (0.019 [0.009, 0.051]); Kruskal-Wallis omnibus test: $\chi^2(4)$ =
38.25, $p < 0.001$; post-hoc Wilcoxon rank-sum test with Holm-Bonferroni correction 29e vs. MEC: $Z = –4.6$, $p < 0.001$; 29e
vs. PA: $Z = –4.8$, $p < 0.001$; 29e vs. PO: $Z = -2.8$, $p = 0.009$; 29e vs. PR: $Z = 0.0$, $p = 0.961$.Rayleigh scores were
significantly higher in 29e (median [IQR] = 0.165 [0.064, 0.521]) than other regions (MEC = 0.066 [0.033, 0.134]), PA =
0.096 [0.045, 0.295], PO = 0.099 [0.045, 0.264], PR = 0.085 [0.043, 0.396]); Kruskal-Wallis omnibus test $\chi^2(4)$ = 54.37, $p <$
0.001; post-hoc Wilcoxon rank-sum test with Holm-Bonferroni correction 29e vs. MEC: $Z = 6.6$, $p < 0.001$; 29e vs. PA: $Z =$
3.3, $p < 0.001$; 29e vs. PO: $Z = 2.5$, $p = 0.022$; 29e vs. PR: $Z = 2.1$, $p = 0.035$.Similar to Rayleigh scores, EVF scores were
also significantly higher in 29e (median [IQR] = 0.336 [0.195, 0.516]) than other regions (MEC = 0.193 [0.116, 0.282], PA =
0.197 [0.120, 0.338], PO = 0.254 [0.163, 0.395], PR = 0.225 [0.123, 0.353]); Kruskal-Wallis omnibus test $\chi^2(4)$ = 55.73, $p <$
0.001; post-hoc Wilcoxon rank-sum test with Holm-Bonferroni correction 29e vs. MEC: $Z = 7.1$, $p < 0.001$; 29e vs. PA: $Z =$
5.5, $p < 0.001$; 29e vs. PO: $Z = 2.7$, $p = 0.006$; 29e vs. PR: $Z = 3.9$, $p < 0.001$.
(C) Egocentric tuning of example 29e neurons. For each neuron, the left panel shows the HD vector field, with the EVF score
indicated. Arrow direction depicts the neuron's preferred HD within a 15 × 15 cm positional bin centered at the arrow base;
arrow size represents the product of the neuron's total firing rate (summed across head-direction bins) and its mean resultant
length inside the same positional bin (see Methods). Within each HD vector field, red/yellow circles mark anchor locations
inside the arena, while red/yellow arrowheads point to anchors outside the arena boundaries. (Note that the EVF scores are
independent of the anchor locations.) On the right, the top panel shows the allocentric 2D positional rate map and the bottom
panel shows the HD tuning curves in allocentric (gray) and egocentric (black) reference frames. The color scale (blue to
yellow) represents firing rate from minimum to maximum, which are displayed above each rate map as (min, max) Hz.
Numbers in the HD plots denote the mean resultant length of HD tuning in each frame.
(D) Pie charts showing the proportions of egocentric HD-tuned, allocentric HD-tuned, non-directional place-tuned neurons,
and non-spatial neurons. Region IDs and the total neuron counts are given above each chart. There were significant regional
differences in the prevalence of spatial tuning types (Chi-squared test $\chi^2(12,1028)$ = 107.55, p < 0.001). Post-hoc Fisher's
exact test with Holm-Bonferroni correction showed that egocentric HD-tuned neurons were significantly more prevalent in
29e than in MEC ($p < 0.001$), PA ($p < 0.001$), and PO ($p = 0.039$) but not PR ($p = 0.182$). Nondirectional place-tuned neurons
were less prevalent in 29e than in MEC ($p < 0.001$) and PA ($p < 0.001$), but not PO ($p = 0.812$) or PR ($p = 1.000$). The
prevalence of allocentric HD-tuned neurons in 29e was similar across regions (all post-hoc pairwise $p > 0.075$).
(E) Significant difference in the landmark-locking scores between 29e allocentric and egocentric neurons (represented as
dots) during *LM-failure* sessions in the Dome. Bars show population means (allocentric: 0.39; egocentric: 0.60), and error
bars indicate s.e.m. (allocentric: 0.09; egocentric: 0.05); t-test allocentric vs. egocentric: $t(44) = –2.5$, $p = 0.030$.
(F) Cumulative distributions of the signed distances from the estimated anchor points of egocentric neurons to the open-arena
border. We first tested, for each region, whether its signed distances were significantly different from zero. Area 29e showed
significantly positive median distance (median [IQR] = 26 [–14, 289] cm; Wilcoxon's signed-rank test with Holm-Bonferroni
correction: $p = 0.016$). By contrast, the medians were negative in other regions (MEC = –24 [–45, –11], $p = 0.005$; PA = –13
[–28, 8], p = 0.186; PO = –20 [–39, 28], p = 0.892; PR = –24 [–40, 31], p = 0.892), Direct comparison between 29e and other
regions showed that the signed distance was significantly larger in 29e than other regions (Kruskal-Wallis test $\chi^2(4)$ = 22.55,
$p < 0.001$; post-hoc Wilcoxon rank-sum test with Holm-Bonferroni correction 29e vs. MEC: $Z = 4.4$, $p < 0.001$; 29e vs. PA:
$Z = 3.2$, $p = 0.004$; 29e vs. PO: $Z = 2.8$, $p = 0.011$; 29e vs. PR: $Z = 2.5$, $p = 0.012$).
(G) Distribution of preferred bearing angles for egocentric neurons in area 29e. Radial axis indicates the number of neurons.
Significantly more neurons had anchor points on the animal's left (n = 28) than right side (n = 11; binomial test with a null
probability of 0.5, p = 0.009). Furthermore, the overall circular distribution was significantly non-uniform (Rayleigh test: z =
4.02 $p = 0.017$), skewed toward the left-rear quadrant as indicated by the circular mean direction (red arrow).

Area 29e differed from neighboring parahippocampal regions in theta and gamma modulation

Having demonstrated differences between 29e and the other regions in the strength of landmark
anchoring and responses to changes in visual inputs, we next explored physiological characteristics that
might underlie the network mechanisms governing the interactions among these regions. Area 29e
differed from the other regions in both its local field potential (LFP; Fig. 4A) properties and its spike–
LFP coupling. All regions showed significant LFP oscillatory power in the theta (5-10 Hz), beta (20–
35Hz), and gamma (50–120 Hz) bands (Fig. 4B). Given the prominent roles of theta and gamma
rhythms in hippocampal-parahippocampal spatial processing, we focused subsequent analyses on these
two frequency bands. Area 29e had the lowest theta power (mean ± s.e.m. = 0.687 ± 0.072), which was
determined to be significantly different from MEC (1.148 ± 0.028) and PO (1.365 ± 0.058), but not PA

(1.036 ± 0.043) or PR (0.735 ± 0.056) by a linear mixed-effects model accounting for non-independence
among LFP signals from the same rat (fig. S10A). Gamma-band power in 29e was statistically
comparable to other regions (fig. S10B).

These regional differences in LFP were matched at the level of single-neuron firing properties. Mean
firing rates in the Dome were lowest in 29e compared to the other regions (fig. S10C), although the
difference was statistically significant only relative to MEC, PA and PR. Theta rhythmicity (*22, 49–51*)
in the neurons' spike autocorrelograms was significantly weaker in 29e than the other regions (Fig. 4C-
D). In contrast, gamma-band rhythmicity was significantly higher in 29e (Fig. 4E). Spike coupling to
LFP theta oscillations, quantified by the pairwise phase consistency (PPC) of the spike-phase
distribution (*52*), was significantly lower in 29e than other recorded regions (Fig. 4F).

We hypothesized that within area 29e, neurons with stronger sensory-driven, landmark-related
properties would be less modulated by the theta rhythm, whereas other 29e neurons would more closely
resemble the strongly theta-modulated populations in MEC, PA, PO, and PR. To test these hypotheses,
we first focused on *LM-failure* sessions, i.e., the sessions for which we could dissociate landmark-
anchored neurons of 29e from non-anchored ones. Splitting the 29e population into two halves based on
their landmark-locking scores showed that neurons in the top half had significantly lower theta PPC than
those in the bottom half (Fig. 4G; fig. S11A-C). Next, focusing on luminance-manipulation sessions, we
similarly observed that 29e neurons in the top half of the landmark-contrast tuning strength had
significantly lower theta PPC than those in the bottom half (Fig. 4H; fig. S11D-F). Finally, examining
open-arena sessions, we found that 29e neurons classified as egocentric exhibited significantly lower
theta PPC than those classified as allocentric (Fig. 4I; fig. S11G). These results demonstrate that neurons
with more sensory-driven properties prevalent in 29e (landmark locking, landmark-contrast tuning, and
egocentric tuning) were less strongly coupled to LFP theta, the dominant rhythm of the rodent
hippocampal-parahippocampal system.

Fig. 4. LFP-related differences between area 29e and neighboring parahippocampal areas (MEC, PA, PO, and PR).
(A) Representative LFP traces, simultaneously recorded from 29e, MEC, PA, PO, and PR.
(B) Mean ± 2 s.e.m. log-power spectral density (PSD) of LFPs in all recorded regions (unique recording locations for 29e: n
= 20, MEC: n = 104, PA: n = 64, PO: n =22, PR: n = 24). PSDs are computed after correcting the aperiodic 1/f background in
the LFP recordings. Asterisks mark the statistical significance of the power at a particular frequency, as measured by a cluster
permutation test to control familywise error rate at α = 0.05 (*53*).
(C) Population mean ± s.e.m. of the normalized spike autocorrelograms (ACGs) across single neurons in each region (colors
as in panel B). Area 29e shows a monotonic, non-oscillatory decay, whereas other regions exhibit prominent oscillations at
the theta frequency.
(D-F) Cumulative distributions of single-neuron physiological measures across recorded regions (area 29e: n = 123, MEC: n
= 566, PA: n = 414, PO: n = 133, PR: n = 124). *Panel D* shows significantly lower theta rhythmicity of spike ACGs in 29e
(median [IQR] = 2.01 [1.29, 2.89]) than other regions (MEC = 3.91 [2.06, 12.4], PA = 7.08 [2.35, 15.53], PO = 4.67 [1.79,
13.19], and PR = 3.08 [1.44, 10.29]; Kruskal-Wallis test $\chi^2(4) = 100.4$, $p < 0.001$; post-hoc Wilcoxon rank-sum tests with
Holm-Bonferroni correction 29e vs. MEC: $Z = -8.1$, $p < 0.001$; 29e vs. PA: $Z = -9.4$, $p < 0.001$; 29e vs. PO: $Z = -6.2$, $p <$
0.001; 29e vs. PR: $Z = -4.1$, $p < 0.001$). *Panel E* shows significantly larger gamma rhythmicity in 29e (median [IQR] = 0.95
[0.63, 1.69]) than other regions (MEC = 0.76 [0.44, 1.25], PA = 0.46 [0.22, 0.91], PO = 0.87 [0.34, 1.43], PR = 0.69 [0.30,
1.38]; Kruskal-Wallis test $\chi^2(4) = 91.1$, $p < 0.001$; post-hoc Wilcoxon rank-sum tests with Holm-Bonferroni correction 29e
vs. MEC: $Z = 3.40$, $p < 0.001$; 29e vs. PA: $Z = 7.7$, $p < 0.001$; 29e vs. PO: $Z = 2.2$, $p = 0.025$; 29e vs. PR: $Z = 3.2$, $p =$
0.002). *Panel F* shows significantly reduced theta PPC of neurons in 29e (median [IQR] = 0.0016 [0.0004, 0.0052]) than in

other regions (MEC = 0.0496 [0.0119, 0.1629], PA = 0.1040 [0.0192, 0.2773], PO = 0.0376 [0.0046, 0.2066], and PR =
0.0218 [0.0044, 0.1652]; Kruskal-Wallis test $\chi^2(4) = 235.7$, $p < 0.001$; post-hoc Wilcoxon rank-sum tests with Holm-
Bonferroni correction 29e vs. MEC: $Z = -13.7$, $p < 0.001$; 29e vs. PA: $Z = -14.1$, $p < 0.001$; 29e vs. PO: $Z = -8.7$, $p <$
0.001; 29e vs. PR: $Z = -8.4$, $p < 0.001$).
(G-I) Reduced spike–LFP theta coupling in 29e neurons with higher landmark locking (G), stronger landmark-contrast
tuning (H), and egocentric (vs. allocentric) HD tuning (I). Dots indicate single neurons, bars show mean ± s.e.m., *Panel G:*
Spatially correlated 29e neurons recorded in *LM-failure* sessions in the Dome were split at the median landmark-locking
score. The low landmark-locking score group exhibited significantly higher theta PPC than the high landmark-locking score
group (t-test on arcsinh-transformed PPC values; $t(55) = 2.51$, $p = 0.015$). *Panel H*: 29e neurons recorded in luminance-
manipulation sessions were split at the median of their partial R^2 values. The low partial R^2 group showed significantly
higher theta PPC than the high partial R^2 group (t-test on arcsinh-transformed PPC values; $t(43) = 2.94$, $p = 0.005$). *Panel I*:
29e neurons classified as allocentric-HD tuned in open arena sessions had significantly higher theta PPC than their egocentric
counterparts (Wilcoxon's rank-sum test on asinh-transformed PPC values: $Z = 2.63$, $p = 0.009$).

Breakdown of theta- and gamma-mediated interactions between area 29e and MEC LFPs preceded the loss of landmark control across the parahippocampal system

To investigate the interactions and coordination between 29e and MEC networks under conditions of
LM-control and *LM-failure*, we measured coherence and Granger causality between 29e and MEC at
theta and gamma frequencies. During periods of stationary landmarks (i.e., the initial epoch with
experimental gain = 1), average LFP–LFP coherence across 29e–MEC tetrode pairs (n = 32; 4 rats)
exceeded chance levels throughout the theta band and at multiple gamma bands (Fig. 5A), paralleling
the spectral profile of significant oscillatory power in both regions (Fig. 4B). To assess the directionality
of influence between 29e and MEC LFPs, we computed the frequency-resolved Granger causality (GC)
spectra (*54–57*) and calculated a *directed-influence asymmetry index* (DAI) as $(GC_{29e \rightarrow MEC} - GC_{MEC}$
$_{\rightarrow 29e})/(GC_{29e \rightarrow MEC} + GC_{MEC \rightarrow 29e})$ (*58*). A significantly negative DAI indicates a stronger drive from
MEC onto 29e, whereas a significantly positive DAI indicates a stronger drive from 29e to MEC. This
analysis revealed that MEC significantly drove 29e in the theta band, whereas 29e significantly drove
MEC in the gamma band (Fig. 5B).

We quantified, for each 29e-MEC tetrode pair, the changes in LFP–LFP coherence from the initial
epoch with stationary landmarks to the final epoch with rotating landmarks. In *LM-control* sessions,
neither theta nor gamma coherence changed significantly. By contrast, in *LM-failure* sessions, coherence
decreased in both frequency bands, with gamma reaching significance and theta only approaching it
(Fig. 5C). Overall, coherence decreased significantly more in *LM-failure* sessions than in *LM-control*
sessions. This difference between session types was primarily driven by changes in gamma coherence.
These results demonstrate that LM-failure was associated with gamma-band desynchronization between
29e and MEC LFPs, with weaker evidence for a similar trend in the theta band.

LM-failure was also associated with a shift in directionality of 29e-MEC LFP interactions. In *LM-
control* sessions, the DAI curve remained stable from the initial to the final epoch across the whole
frequency range (Fig. 5D-E, left). Thus, even though gain manipulation in the Dome creates a conflict
between path integration and landmarks (*12, 59*), the strength of gamma-mediated, 29e→MEC drive and
the strength of theta-mediated, MEC→29e feedback did not change so long as the landmarks maintained
control over the parahippocampal representations of space. However, in *LM-failure* sessions where
landmark control was lost, the DAI curve shifted downwards (Fig. 5D-E, right), indicating a *decrease* in
the gamma-mediated, 29e→MEC drive and an *increase* in the theta-mediated, MEC→29e drive (note
that a more negative DAI reflects a greater MEC→29e drive). The DAI curve's downward shift was
significant for gamma but only approached significance for theta—a pattern similar to the coherence
results (Fig. 5F; see fig. S12A for further breakdown of the components of the DAI). Overall, DAI
changes in *LM-failure* sessions were significantly different from those in *LM-control* sessions. Thus, the

loss of visual landmark control over the parahippocampal representations during a session was
associated with a breakdown of the normal gamma-band influence from 29e to MEC.

We next asked whether the changes in 29e-MEC LFP interactions emerged before or after the loss of
landmark control in *LM-failure* sessions. To address this question, we quantified the ongoing strength of
landmark control by measuring how consistently the non-29e neuronal population activity vector
repeated across laps in the reference frame of the rotating landmarks (see Methods). The trajectory of
this relative landmark-control strength closely matched the ongoing changes in 29e-MEC LFP coupling
measures (coherence and DAI) in both theta and gamma bands, albeit with some temporal differences
(Fig. 5G). Linear mixed-effects models that account for non-independence of observations within
animals and within tetrode pairs confirmed the statistical significance of these positive associations for
both LFP coupling measures and both frequency bands. Cross-correlation analysis, comparing observed
correlations to a null distribution of correlations from circularly shuffled landmark-control strength
trajectories, revealed significant positive correlations ($\alpha = 0.05$) in 11 of the 18 tetrode pairs for theta-
band coherence, 13 of the 18 for gamma-band coherence, and 10 of the 18 for both theta- and gamma-
band DAI. (all $p < 0.001$ as per binomial tests against a null probability of 0.05).

Analysis of cross-correlation lags using significant tetrode pairs showed that ongoing changes in
landmark-control strength were synchronous with changes in gamma-band coherence and theta-band
DAI across laps but were significantly preceded by changes in theta-band coherence and gamma-band
DAI. A direct comparison between these leading indicators showed that changes in gamma-band DAI
occurred significantly earlier than changes in theta-band coherence ($Z = -2.34$, $p = 0.019$). Taken
together with the existence of landmark-anchored neurons within 29e and the dominant directionality of
gamma drive from 29e to MEC during stable landmark control, this finding suggests reduced 29e-to-
MEC gamma drive as an early indicator of impending landmark-control failure.

Fig. 5. Theta-gamma coordination between area 29e and MEC.

(A) LFP–LFP coherence spectrum between 29e and MEC LFPs when landmarks were stationary. Gray shaded area surrounding the solid black line indicates the mean ± s.e.m. of coherence spectra across 32 tetrode pairs. Asterisks mark the statistical significance of the coherence at a particular frequency, as measured by a hierarchical cluster-based permutation test to control familywise error rate at α = 0.05.

(B) Directed-influence asymmetry index (DAI) between 29e and MEC LFPs when landmarks were stationary. The convention is the same as panel A.

(C) Normalized changes in theta- and gamma-band coherence between 29e and MEC LFPs from the initial epoch to the final epoch across *LM-control* and *LM-failure* sessions. Each data point represents a 29e-MEC tetrode pair; bars represent mean ± s.e.m. The model included fixed effects of session type (*LM-control* vs *LM-failure*), frequency band (theta vs. gamma), and their interaction, and random effects accounting for repeated measurements within rats and tetrode pairs (see Methods for details). Coherence changes in *LM-control* sessions did not differ significantly from zero (theta band: $t(8.3) = -1.6$, $p = 0.300$; gamma band: $t(8.3) = 1.4$, $p = 0.300$), whereas coherence changes in *LM-failure* sessions were negative, reaching statistical significance in the gamma band ($t(9.7) = -3.3$, $p = 0.035$) but not in the theta band ($t(9.7) = -2.8$, $p = 0.062$). Holm-Bonferroni correction was applied to all four comparisons. The model also revealed a significant main effect of session type (LM-*control* vs. *LM-failure*: $F(1, 31.1) = 10.6$, $p = 0.003$) and a significant session type × frequency-band interaction effect ($F(1,33.8) = 5.1$, $p = 0.030$) on within-session coherence changes, but no significant main effect of the frequency band ($F(1,3.3) = 1.4$, $p = 0.313$). Post-hoc pairwise comparisons indicated that the interaction was driven by a difference in gamma-band coherence changes between *LM-control* and *LM-failure* sessions ($t(44.4) = 3.9$, $p = 0.001$; Holm–Bonferroni corrected across four post-hoc comparisons).

(D) Mean DAI spectra across tetrode pairs in the initial (black) and final (grey) epochs of *LM-control* (left) and *LM-failure* (right) sessions. The pattern of frequency-dependent DAI between epochs is similar across initial and final epochs in *LM-control* sessions. By contrast, the DAI in *LM-failure* sessions shifts downward in both theta- and gamma-bands, indicating a shift toward stronger theta drive from MEC onto 29e (in comparison to theta drive in the reciprocal direction) and toward weaker gamma drive from 29e to MEC in the gamma band.

(E) Theta- and gamma-band directed-influence asymmetry index (DAI) values in the initial epoch (x-axis) with stationary landmarks vs. the final epoch (y-axis) with rotating landmarks across *LM-control* (left) and *LM-failure* (right) sessions. Each data point represents a tetrode pair. The diagonal dashed line represents the identity y = x (i.e., initial DAI = final DAI). In *LM-control* sessions (left), points cluster around the identity line, indicating minimal change. In *LM-failure* sessions (right), however, most points fall below the identity line, indicating a systematic decrease in DAI. The within-session DAI decrease has opposite implications for the two frequency bands: in the theta-band, where initial DAI values were predominantly negative, the further decrease points to a shift toward enhanced MEC→29e drive during the loss of landmark control; by contrast, in the gamma-band, where initial DAI values were predominantly positive, the decrease indicates a shift toward diminished 29e→MEC drive during the loss of landmark control.

(F) Initial to final epoch change in theta- and gamma-band DAI (quantified as $DAI_{final} − DAI_{initial}$) between 29e-MEC tetrode pairs across *LM-control* and *LM-failure* sessions. We analyzed data using a linear mixed-effects model that has the same random- and fixed-effects structure as in panel C. Like panel C, the fitted model was first used to test whether DAI changes differed from zero in each of the four conditions (Control–Theta, Control–Gamma, Failure–Theta, Failure–Gamma; Holm–Bonferroni corrected across four tests): DAI changes in *LM-control* sessions were slightly positive but not significant (theta band: $t(7.8) = 1.9$, $p = 0.199$; gamma band: $t(8.3) = 0.5$, $p = 0.660$). By contrast, DAI changes in *LM-failure* sessions were negative, reaching statistical significance in the gamma band ($t(8.7) = −4.4$, $p = 0.008$) and a statistical trend in the theta band ($t(8.7) = −2.9$, $p = 0.058$). The model also revealed a significant main effect of session type (*LM-control* vs. *LM-failure*: $F(1, 43.3) = 20.7$, $p < 0.001$), consistent with an overall decrease in DAI from initial to final epoch in *LM-failure* sessions in contrast to an overall DAI increase in *LM-control* sessions. There was no significant main effect of the frequency band or its interaction with the session type.

(G) Average lapwise trajectories of normalized gain deviation (i.e., the unsigned difference between the current value of the experimental gain and the initial baseline gain of 1) and of 29e–MEC LFP coupling measures (bottom) across 14 tetrode pairs from *LM-failure* sessions, aligned to the point of landmark-control failure (x = 0). The normalized x-axis interval [–1, 1] corresponds to approximately 75 laps. One rat was excluded from this visualization due to instability of firing fields of its non-29e neurons with respect to landmarks during stationary periods, but data from this rat were included in quantitative analyses. For the visualization, all traces were z-scored, smoothed with a moving average filter (half-width = 2 laps), aligned to the point of landmark-control failure (vertical dashed line at 0), time normalized (i.e., temporally scaled to also align the beginning and end points of all experiments), and averaged across tetrode pairs. For each session, the failure point was determined by visual inspection of lapwise firing rate plots of non-29e neurons relative to landmarks (as in Fig. 2D–E). Overall, the plots show that as gain deviation increased (top), landmark-control strength (bottom; blue) progressively declined, with a marked drop occurring at the point of landmark failure. This trend was closely tracked by the trajectories of 29e–MEC LFP coupling measures (theta-band coherence: median [IQR] Pearson's rho = 0.64 [0.28, 0.78]; gamma-band coherence = 0.66 [0.39, 0.79]; theta-band DAI = 0.41 [0.32, 0.53]; gamma-band DAI = 0.41 [0.30, 0.56]), but with apparent temporal (lapwise, x-axis) offsets. Specifically, reduction in gamma DAI—primarily reflecting loss of 29e→MEC gamma drive (see panel D-E)—preceded both the decline in landmark-control strength and changes in other LFP coupling measures. The decline in landmark-locking strength was largely synchronous with the decline in gamma coherence (median [IQR] lag (lap) = –0.2 [–1.4, 1.7]; Wilcoxon sign-rank test $Z = –0.22$, $p = 0.823$) and theta-band DAI (–1.6 [–3.6, 1.5]; $Z = –0.67$, $p = 0.498$), but was significantly preceded by changes in theta-band coherence (–0.6 [–1.6, 0]; $Z = –2.20$, $p = 0.027$) and gamma-band DAI (–3.0 [–4.0, –2.3]; $Z = -2.55$, $p = 0.010$). Further testing with linear mixed-effects models to account for non-independence of observations within animals and within tetrode pairs (see Methods for random-effects structure) confirmed the significance of the these associations between changes in landmark-control strength and changes in both coherence (theta: $β = 0.24$, $t(2.9) = 4.6$, $p = 0.027$); gamma: $β = 0.34$, $t(3.1) = 7.3$, $p = 0.009$) and DAI (theta: $β = 0.37$, $t(3.0) = 7.2$, $p = 0.011$; gamma: $β = 0.31$, $t(2.9) = 3.7$, $p = 0.035$).

Gamma-band feedforward coupling and local circuit connectivity support coordination of landmark signals in 29e with parahippocampal circuits

To determine whether the spiking activity of 29e and MEC neurons were consistent with the LFP interactions reported in Fig. 5, we examined the coupling of spatially correlated 29e and MEC neurons to the other region's LFP in the Dome. Consistent with the LFP–LFP coupling result (Fig. 5A), the spectral profile of cross-regional spike–LFP pairwise phase consistency (PPC) showed clear peaks in the

theta and gamma bands during periods of stationary landmarks in the Dome (Fig. 6A). In the gamma band, the coupling of 29e neurons' firing to MEC was significantly stronger than that of MEC neurons to 29e, whereas in the theta band, the coupling of MEC neurons' firing to 29e was significantly stronger than that of 29e neurons' to MEC. This pattern matched the reversal in the dominant directionality of LFP–LFP Granger Causality influences across theta and gamma bands shown in Fig. 5B. While cross-regional PPC in both bands did not change significantly during gain manipulations in *LM-control* sessions, significant reductions in PPC were present during *LM-failure* sessions, specifically in the coupling of MEC neurons to 29e theta and 29e neurons to MEC gamma (Fig. 6B).

To address whether the disruption of spike–LFP coupling (particularly MEC neurons-to-29e theta and 29e neurons-to-MEC gamma) emerged before or after the loss of landmark control, we analyzed cross-regional PPC on a lap-by-lap basis in *LM-failure* sessions, analogous to our LFP–LFP analysis (Fig. 5G). Ongoing changes in landmark-control strength closely tracked lap-by-lap changes for both coupling measures, albeit with some temporal differences (Fig. 6C). We confirmed the consistency of these positive relationships across animals by linear mixed-effects modeling that account for non-dependencies among observations. Cross-correlation analysis, which accounts for temporal offsets between landmark-control strength and spike-LFP coupling, revealed significant correlations ($\alpha = 0.05$, compared to circularly shuffled distributions) in 17 of 32 neurons from area 29e and 38 out of 95 from MEC (both $p < 0.001$ as per a binomial test against a null probability of 0.05). Analysis of lags at the peak correlation showed that changes in 29e→MEC gamma coupling significantly preceded changes in landmark-control strength, whereas changes in MEC→29e theta coupling were approximately synchronous with landmark-control strength. Direct comparison confirmed that changes in 29e neurons' coupling to MEC gamma occurred significantly earlier than changes in MEC neurons' coupling to 29e theta. This pattern mirrors the LFP–LFP results, where gamma-band DAI—reflecting the strength of 29e-to-MEC drive—was the earliest predictor of landmark-control failure (Fig. 5G). Thus, across both LFP–LFP and spike–LFP measures, the breakdown of gamma-mediated feedforward signaling from 29e to MEC precedes the failure of landmark control.

If 29e neurons provided landmark information to MEC via gamma oscillations, we would expect landmark-anchored 29e neurons to be more strongly coupled to the MEC LFP gamma than the non-anchored neurons. To test this hypothesis, we calculated, separately for 29e and MEC populations recorded in *LM-failure* sessions, the correlation between neurons' landmark-locking scores (measured in the final epoch) and their baseline cross-regional PPC values (measured in the initial epoch with stationary landmarks) at different frequencies (Fig. 6D). 29e neurons displayed a striking frequency-dependent pattern: neurons with high landmark-locking scores showed strong coupling (i.e., high positive PPC-LM score correlation) to MEC gamma (particularly 100–120 Hz) but weak coupling (i.e., high negative PPC-LM score correlation) to MEC theta. This frequency-dependent pattern in 29e was significantly stronger than expected by chance ($p = 0.006$; fig. S12B). A similar analysis of MEC neurons' coupling to 29e LFP showed that MEC neurons with high landmark- locking scores showed moderately strong coupling to 29e gamma (80-95 Hz), while those with weaker anchoring tended to couple more strongly to 29e theta. However, this frequency-dependent difference did not reach statistical significance ($p = 0.078$; fig. S12C). Furthermore, 29e neurons with high landmark-locking scores were more likely than those with low landmark-locking scores to lose their gamma-band coupling to MEC LFP (fig. S12D-E) as MEC spatial maps broke free from landmarks. These results suggest that gamma-band synchrony between 29e's landmark-anchored subpopulation and MEC may support transfer of landmark information to parahippocampal path-integration circuits. Collectively, these findings suggest that landmark-anchored 29e neurons serve as a key source of landmark information for anchoring representations of parahippocampal path-integration circuits to the external world.

Fig. 6. MEC-gamma coupling, local connectivity of landmark-anchored 29e neurons, and a conceptual model of inter-areal communication between area 29e and other parahippocampal areas.

(A) Cross-regional spike–LFP coupling spectrum between 29e and MEC when landmarks were stationary. Shaded area surrounding the solid black (29e neurons' coupling to MEC LFP) and orange (MEC neurons' coupling to 29e LFP) lines indicate the mean ± s.e.m. of the respective spike–LFP spectra. Asterisks mark the statistically significant differences determined by a cluster-based permutation test to control familywise error rate at α =0.05.

(B) Normalized changes in cross-regional spike–LFP coupling ([final – initial]/[final + initial]) for each neuron between 29e and MEC across *LM-control* and *LM-failure* sessions. Each dot represents a single neuron; bars indicate mean ± s.e.m. Asterisks above bars denote significance levels compared to zero; asterisks below brackets denote significance between respective conditions. In *LM-control* sessions, coupling changes did not differ significantly from zero in either theta band (29 neurons-to-MEC theta: median [IQR] = –0.02 [–0.42, 0.44]; Holm-Bonferroni corrected Wilcoxon sign-rank test $Z = 0.2$, $p =$ 1.000; MEC neuron–29e theta: 0.07 [–0.21, –0.32]; $Z = 1.1$, $p = 1.000$) or gamma band (29e neurons-to-MEC gamma: 0.04 [–0.49, 0.36]; $Z = –0.3$, $p = 1.000$; MEC neurons-to-29e gamma: –0.06 [–0.32, 0.26]; $Z = 0.09$, $p = 1.000$). By contrast, *LM-failure* sessions showed significant reductions in couplings of 29e neurons-to-MEC gamma (–0.31 [–0.55, –0.02]; $Z = –3.56$, $p = 0.001$) and of MEC neurons-to-29e theta (–0.32 [–0.52, –0.08]; $Z = –5.21$, $p < 0.001$) but no significant change in couplings of 29e neurons-to-MEC theta (–0.15 [–0.77, 0.41]; $Z = –1.12$, $p = 0.262$) or of MEC neurons-to-29e gamma (–0.16 [–0.51, 0.27]; $Z = -–2.00$, $p = 0.092$). Three-way repeated measures ANOVA (factors: session type, frequency-band, and neuron region-LFP region pair identity) revealed significant main effect of the session type ($F(1,202) = 9.7$, $p = 0.002$) and a

significant three-way interaction ($F_{(1,202)}$ = 4.0, p = 0.048) but no significant main effects of the frequency-band ($F_{(1,202)}$ = 0.3, p = 0.583) or neuron-LFP region pair ($F_{(1,202)}$ = 0.54, p = 0.463), or any two way interaction. Holm-Bonferroni corrected post-hoc pairwise Wilcoxon tests within each frequency band identified two significant differences pertaining to the interaction effect. Within the theta-band, the changes in MEC neuron-29e LFP theta coupling was significantly lower in *LM-failure* sessions than *LM-control* sessions (Z = –4.4, p < 0.001). Within the gamma-band, the changes in 29e neuron-MEC LFP coupling across *LM-failure* sessions was significantly lower than the changes in MEC neuron-29e LFP coupling across *LM-control* sessions (Z = –3.2, p = 0.012).

(C) Average lapwise trajectories of gain deviation (top; defined as in Fig. 5G) and of cross-regional spike–LFP coupling measures across *LM-failure* sessions (bottom). Same format as Fig. 5G. Only neurons that were active in at least 75% of laps and exhibited split-session stability were included (n = 32 from area 29e, n = 95 from MEC) to ensure reliable lap-by-lap estimates of coupling. The same pre-processing steps were applied as in Fig. 5G, except that the smoothing filter used a half-width of 2.5 laps to compensate for the sparser activity of individual neurons, which have spatially localized firing fields, compared to continuously varying LFP signals. Overall, the plots show that both spike–LFP coupling measures declined as experimental gain deviated from 1 (top), closely tracking the progressive decline in landmark-control strength (blue; 29e neurons-to-MEC gamma: median [IQR] Pearson's rho = 0.60 [0.52, 0.70]; MEC neurons-to-29e theta: 0.49 [0.26, 0.69]). Notably, the coupling of 29e neurons to MEC gamma (black) showed a rapid decline before the drop in landmark-control strength (blue; median [IQR] cross-correlation lag (lap) = –3.0 [–4.2, 0]; Wilcoxon signed-rank test Z = –2.61, p = 0.008). By contrast, MEC neurons' coupling to 29e theta (pink) also declined but did so more gradually without showing any clear temporal lead or lag relative to landmark-control strength (0.1 [–3.2, 3.7]; Z = 0.06, p = 0.956). Applying linear mixed-effects modeling that accounted for non-independence of observations within animals and within neurons (see Methods for details), we quantitatively confirmed significant positive associations between ongoing changes in landmark-control strength and changes in both spike–LFP coupling measures (Holm-Bonferroni correction; 29e neurons to MEC gamma: β = 0.37, $t_{(2.9)}$ = 4.4, p = 0.042; MEC neurons to 29e theta: β = 0.22, $t_{(2.7)}$ = 3.55, p = 0.044).

(D) Correlation between baseline cross-regional spike–LFP PPC and landmark-locking score measured during the final 12 laps with rotating landmarks during failure sessions. Spearman's correlation coefficient is shown as a function of frequency (x-axis) for MEC neurons to 29e LFP (orange) and for 29e neurons to MEC LFP (black). Insets show example scatterplots of landmark-locking scores (LM score) versus PPC values (arcsinh-transformed for visualization) at selected frequencies. Asterisks and lighter colored segments indicate frequency ranges with significant nonzero correlations (cluster-based permutation test to control familywise error rate at α = 0.05).

(E) Conceptual model of area-29e–mediated landmark control of path integration. Based on established anatomy and our physiological findings, we propose that gamma-band drive from 29e conveys feedforward landmark information to MEC, whereas theta-band drive from MEC conveys feedback, path-integration–based position estimates to 29e. Area 29e receives low-latency landmark signals via a direct retina→dLGN→29e pathway and is reciprocally connected to downstream areas MEC and PR. We propose that visual landmark signals differentially innervate 29e neurons, yielding the gradient of landmark anchoring observed in our data. For simplicity, the schematic renders this continuum as two subpopulations, landmark-locked (dark gray) and landmark-unlocked (purple). Feedforward output from 29e could arise from the strongly landmark-anchored subpopulation, consistent with their stronger spike–LFP coupling to MEC gamma during landmark control and the large reduction of gamma coupling when landmark control failed. We further hypothesize that MEC feedback preferentially innervates the less landmark-anchored subset, accounting for their stronger theta coupling and functional similarities to neurons in other parahippocampal areas, including MEC. MEC, the putative path-integration hub, is often modeled as a continuous attractor network; during landmark failure, the MEC neuron population remained largely coherent— either landmark-controlled or not—consistent with attractor dynamics. Within 29e, putative monosynaptic connections are biased from landmark-locked to landmark-unlocked neurons (fig. S13), consistent with the skewness in the distribution of (presynaptic – postsynaptic) landmark-locking scores. Consequently, the 29e landmark-unlocked population integrates inputs from locked 29e cells as well as from other parahippocampal areas. An indirect route from 29e to MEC may be relayed via PR, although effective communication between 29e and PR remains to be shown. Together, these interactions form a functional loop in which 29e provides sensory-based, landmark-anchored positional signals to MEC path-integration circuits, while MEC, combining the landmarks with self-motion cues, calculates an internal position signal that feeds back to 29e. Arrows denote established anatomical connectivity; colored waveforms indicate relevant frequency bands.

Discussion

Brains build internal models to predict the world, and because the world is dynamic and predictions can be inaccurate, these models must be continually updated by sensory feedback (*60, 61*). In navigation, the cognitive map embodies an internal model of space, and external landmarks provide allothetic spatial cues to recognize environments and locations within them and to correct accumulated errors in the path-integration-based estimates of position on this map. The neural origins of the landmark-based correction

signal remain unknown in mammals. To address this gap, we performed simultaneous recordings in rats
across five parahippocampal regions in an immersive VR apparatus that preserves naturalistic
locomotion while affording accurate control over the position and salience of visual landmarks (*33*, *62*),
thereby allowing us to create conflicts between internal path integration computations and feedback
from external visual cues. When these conflicts disrupted landmark anchoring in most spatial neurons
across MEC, PA, PO, and PR, many neurons maintained landmark-anchored firing in area 29e. Area 29e
showed weaker theta in the LFP and weaker theta modulation of its neurons' spiking compared to MEC,
PA, PO, and PR. Area 29e also showed stronger egocentric and landmark-contrast tuning than other
regions, and these sensory properties were markedly enhanced in its landmark-anchored subpopulation.
These results reveal physiological and functional dissociations between area 29e and its neighboring
regions, providing the first such characterization of area 29e in rodents. Taken together with the
dynamic functional coupling between 29e and MEC that correlates with the degree of landmark control,
our results identify a visually driven, egocentric-HD population in 29e that relays landmark signals to
parahippocampal circuits supporting path integration.

Historically, anatomists treated area 29e variably: some incorporated it into retrosplenial cortex (*63*), PR
(*64–66*) or PA (*67–69*), while others suggested that it may be a separate region (*70*, *71*). In this work, we
adopted Blackstad's definition of area 29e in rats as a cytoarchitecturally distinct triangular region
interposed between PR and PA (*70*, *72*). Recent work has proposed that area 29e is the rodent homolog
of the primate visual area prostriata (*26*, *29*, *73*), which is located next to the region of striate cortex
(V1) that represents the most extreme periphery of the visual field and is characterized by broad
orientation tuning, large receptive fields, and selectivity for fast-moving stimuli in the visual periphery
(*28*, *27*, *74*, *75*).

Area 29e neurons in the present study displayed hallmarks of distal visual landmark representation—
strong landmark anchoring, contrast sensitivity, and egocentric HD tuning anchored to locations beyond
the open arena where distal cues were situated, with bearing angles preferentially falling in the
peripheral visual field. Distal cues are thought to be a major signal that orients the cognitive map relative
to the external world (*76–78*). Our data from rodent area 29e suggest that the orienting influence of
distal visual cues may be mediated through peripheral vision—consistent with human behavioral studies
reporting that individuals with restricted peripheral vision make greater errors in orientation and distance
when navigating to remembered landmark locations (*79–81*). This peripheral specialization is
anatomically supported by widespread input to 29e from medial V1 (which processes peripheral vision)
and dorsal lateral geniculate nucleus (dLGN; *36*, *37*, *39*). The direct dLGN connection places 29e only
two synapses from the retinal ganglion cells, potentially providing the fastest visual pathway to limbic
circuits and explaining the short-latency responses observed in primates (*28*, *74*). Recent work has
shown that dLGN drives gamma events promoting visual encoding in V1 (*82*) and that dLGN lesions
severely disrupt the anchoring of HD tuning in PR to visual landmarks (*83*), together raising the
possibility that dLGN efferents are the source of gamma-modulated activity characterizing visual
landmark coding within 29e.

Anatomical and functional evidence suggests that 29e transmits these landmark signals to MEC, the
putative hub of path integration (*9*, *84*, *8*, *15–18*, *20*, *85*). Efferents of 29e in rodents target PR and MEC
(albeit more sparsely than PR), in addition to contralateral prostriata and visuomotor areas (*31*, *32*).
MEC is also innervated by PR (*86*, *87*), considered a candidate site for integration of an internal HD
signal with external landmarks (*40*, *88*, *89*, *42*, *44*, *45*, *90*). This anatomical organization suggests two
pathways via which 29e might deliver gamma-modulated visual landmark signals to MEC: a direct
29e→MEC pathway and an indirect 29e→PR→MEC pathway. Our LFP–LFP and spike–LFP coupling
analyses provide functional evidence consistent with such pathways: 29e and MEC showed theta- and

gamma-band synchronization, with a net gamma-band influence of 29e's landmark-anchored population
on MEC and theta-band influence of MEC on 29e. Notably, feedforward gamma-band influence
declined before the failure of landmark control over MEC (as well as other parahippocampal
populations) during large cue conflicts, supporting the idea that 29e transmits the landmark signals that
anchor parahippocampal representations of space to the external world.

The *predictive coding framework* (*91*, *92*) is a general cortical organizing principle in human and
nonhuman animals, proposing that rapidly changing sensory information and prediction errors are
conveyed via feedforward connections, while slower contextual predictions are transmitted via feedback
connections (*93–95*). Consistent with these timescales, feedforward and feedback anatomical pathways
across sensory and frontoparietal networks operate at distinct frequency bands: bottom-up, feedforward
signaling in high-frequency (gamma) and top-down, feedback signaling in low-frequency (alpha/beta)
brain rhythms (*96*, *58*, *97–100*). We found an analogous organization within parahippocampal networks:
area 29e influenced MEC in the gamma band, whereas MEC influenced 29e in the theta band, forming a
gamma–theta-mediated closed loop that signals feedforward landmark information and feedback, path-
integration-based positional predictions between 29e and MEC (Fig. 6E). When path integration and
landmark-based positional cues became increasingly in conflict in our experiments, in many cases these
frequency-specific interactions were disrupted and 29e-MEC rhythms became desynchronized, followed
by a disruption of landmark control across the parahippocampal system. At the level of individual
neurons, 29e neurons that were strongly coupled to MEC theta exhibited comparatively lower landmark
anchoring and contrast sensitivity—similar to MEC neurons and thus presumably reflecting greater
feedback influence from MEC—whereas those strongly coupled to MEC gamma exhibited stronger
landmark anchoring, consistent with a more feedforward role in 29e signaling landmark information to
MEC.

These results demonstrate how a visual area (29e) interacts dynamically with a set of high-order limbic
regions (parahippocampal areas) to bind an internally generated, cognitive representation of a rat's
location to the visual landmarks present in the external world. The cognitive representations in the
parahippocampal regions (MEC, PR, PA, and PO) maintain internal coherence (i.e., the maps remain
stable within their own internal frame of reference), even when the maps become decoupled from the
external sensory world (*85*, *101–105*). The binding of the map to the external world is apparently
accomplished (at least in part) via the gamma-mediated transmission of visual input from 29e to the
parahippocampal regions, with a theta-mediated feedback signal from these regions to 29e that
maintains coherence between the cognitive and sensory processing regions during normal operation.
Under conditions of continual error between visual landmark-derived position information and path-
integration-based updating of the rat's position on its cognitive map, this theta- and gamma-mediated
interareal coupling begins to break down, eventually resulting in the loss of landmark control over the
cognitive map. Consistent with the predictive coding framework, these results suggest a general
organizing framework for interareal coordination of diverse brain regions, necessary to produce
adaptive, coordinated cognition and behavior in a distributed processing system composed of multiple,
anatomically and functionally segregated brain areas.
776
777

References

1. J. Winson, Loss of hippocampal theta rhythm results in spatial memory deficit in the rat. *Science* **201**, 160–163 (1978).

2. G. Buzsáki, Theta oscillations in the hippocampus. *Neuron* **33**, 325–340 (2002).

3. L. L. Colgin, T. Denninger, M. Fyhn, T. Hafting, T. Bonnevie, O. Jensen, M.-B. Moser, E. I. Moser, Frequency of gamma oscillations routes flow of information in the hippocampus. *Nature* **462**, 353–357 (2009).

4. G. Buzsáki, E. I. Moser, Memory, navigation and theta rhythm in the hippocampal-entorhinal system. *Nat. Neurosci.* **16**, 130–138 (2013).

5. L. L. Colgin, Rhythms of the hippocampal network. *Nat. Rev. Neurosci.* **17**, 239–249 (2016).

6. M.-L. Mittelstaedt, H. Mittelstaedt, Homing by path integration in a mammal. *Naturwissenschaften* **67**, 566–567 (1980).

7. A. S. Etienne, K. J. Jeffery, Path integration in mammals. *Hippocampus* **14**, 180–192 (2004).

8. F. Savelli, J. J. Knierim, Origin and role of path integration in the cognitive representations of the hippocampus: computational insights into open questions. *J. Exp. Biol.* **222** (2019).

9. B. L. McNaughton, F. P. Battaglia, O. Jensen, E. I. Moser, M. B. Moser, Path integration and the neural basis of the "cognitive map." *Nat. Rev. Neurosci.* **7**, 663–678 (2006).

10. J. Knierim, H. Kudrimoti, B. McNaughton, Place cells, head direction cells, and the learning of landmark stability. *J. Neurosci.* **15**, 1648–1659 (1995).

11. J. J. Knierim, H. S. Kudrimoti, B. L. McNaughton, Interactions between idiothetic cues and external landmarks in the control of place cells and head direction cells. *J. Neurophysiol.* **80**, 425–446 (1998).

12. R. P. Jayakumar, M. S. Madhav, F. Savelli, H. T. Blair, N. J. Cowan, J. J. Knierim, Recalibration of path integration in hippocampal place cells. *Nature* **566**, 533–537 (2019).

13. K. M. Gothard, W. E. Skaggs, B. L. McNaughton, Dynamics of mismatch correction in the hippocampal ensemble code for space: interaction between path integration and environmental cues. *J. Neurosci.* **16**, 8027–8040 (1996).

14. K. J. Jeffery, Learning of landmark stability and instability by hippocampal place cells. *Neuropharmacology* **37**, 677–687 (1998).

15. T. Hafting, M. Fyhn, S. Molden, M. Moser, E. I. Moser, Microstructure of a spatial map in the entorhinal cortex. *Nature* **436**, 801–806 (2005).

16. T. Solstad, C. N. Bocarra, E. Kropff, M. B. Moser, E. Moser, Representation of geometric borders in the entorhinal cortex. *Science* **322**, 1861–1865 (2008).

17. F. Savelli, D. Yoganarasimha, J. J. Knierim, Influence of boundary removal on the spatial representations of the medial entorhinal cortex. *Hippocampus* **18**, 1270–1282 (2008).

18. E. Kropff, J. E. Carmichael, M.-B. Moser, E. I. Moser, Speed cells in the medial entorhinal cortex. *Nature* **523**, 419–424 (2015).

819
19.   J. R. Hinman, M. P. Brandon, J. R. Climer, G. W. Chapman, M. E. Hasselmo, Multiple Running
Speed Signals in Medial Entorhinal Cortex. *Neuron* **91**, 666–679 (2016).

20.   F. Sargolini, M. Fyhn, T. Hafting, B. L. McNaughton, M. P. Witter, M.-B. Moser, E. I. Moser,
Conjunctive representation of position, direction, and velocity in entorhinal cortex. *Science* **312**,
758–762 (2006).

21.   L. M. Giocomo, T. Stensola, T. Bonnevie, T. Van Cauter, M.-B. Moser, E. I. Moser, Topography
of head direction cells in medial entorhinal cortex. *Curr. Biol.* **24**, 252–262 (2014).

22.   C. N. Boccara, F. Sargolini, V. H. Thoresen, T. Solstad, M. P. Witter, E. I. Moser, M.-B. Moser,
Grid cells in pre- and parasubiculum. *Nat. Neurosci.* **13**, 987–994 (2010).

23.   H. Stensola, T. Stensola, T. Solstad, K. Frøland, M.-B. Moser, E. I. Moser, The entorhinal grid
map is discretized. *Nature* **492**, 72–78 (2012).

24.   J. Taube, R. Muller, J. Ranck, Head-direction cells recorded from the postsubiculum in freely
moving rats. I. Description and quantitative analysis. *J. Neurosci.* **10**, 420–435 (1990).

25.   J. Taube, R. Muller, J. Ranck, Head-direction cells recorded from the postsubiculum in freely
moving rats. II. Effects of environmental manipulations. *J. Neurosci.* **10**, 436–447 (1990).

26.   S. Ding, Comparative anatomy of the prosubiculum, subiculum, presubiculum, postsubiculum,
and parasubiculum in human, monkey, and rodent. *J. Comp. Neurol.* **521**, 4145–4162 (2013).

27.   K. Mikellidou, J. W. Kurzawski, F. Frijia, D. Montanaro, V. Greco, D. C. Burr, M. C. Morrone,
Area prostriata in the human brain. *Curr. Biol.* **27**, 3056-3060.e3 (2017).

28.   H.-H. Yu, T. A. Chaplin, A. J. Davies, R. Verma, M. G. P. Rosa, A specialized area in limbic
cortex for fast analysis of peripheral vision. *Curr. Biol.* **22**, 1351–1357 (2012).

29.   W. Lu, S. Chen, X. Chen, J. Hu, A. Xuan, S. Ding, Localization of area prostriata and its
connections with primary visual cortex in rodent. *J. Comp. Neurol.* **528**, 389–406 (2020).

30.   J.-M. Hu, C.-H. Chen, S.-Q. Chen, S.-L. Ding, Afferent projections to area prostriata of the
mouse. *Front. Neuroanat.* **14**, 605021 (2020).

31.   C.-H. Chen, J.-M. Hu, S.-Q. Chen, S.-M. Liu, S.-L. Ding, Homotopic commissural projections of
area prostriata in rat and mouse: comparison with presubiculum and parasubiculum. *Front.*
Neural Circuits **14**, 605332 (2020).

32.   C.-H. Chen, J.-M. Hu, S.-Y. Zhang, X.-J. Xiang, S.-Q. Chen, S.-L. Ding, Rodent area prostriata
converges multimodal hierarchical inputs and projects to the structures important for visuomotor
behaviors. *Front. Neurosci.* **15**, 772016 (2021).

33.   M. S. Madhav, R. P. Jayakumar, S. G. Lashkari, F. Savelli, H. T. Blair, J. J. Knierim, N. J.
Cowan, The dome: a virtual reality apparatus for freely locomoting rodents. *J. Neurosci.*
Methods **368**, 109336 (2022).

34.   M. S. Madhav, R. P. Jayakumar, B. Y. Li, S. G. Lashkari, K. Wright, F. Savelli, J. J. Knierim, N.
J. Cowan, Control and recalibration of path integration in place cells using optic flow. *Nat.*
Neurosci. **27**, 1599–1608 (2024).

35.   R. P. Jayakumar, Y. Sueoka, M. K. Ferreyros, B. Y. Li, M. S. Madhav, X. Chen, J. J. Knierim,
N. J. Cowan, Synchronized path-integration recalibration but distinct landmark-control dynamics
in head direction and CA1 place cells. Neuroscience [Preprint] (2025).
https://doi.org/10.1101/2025.10.08.681135.

36. F. Cacucci, C. Lever, T. J. Wills, N. Burgess, J. O'Keefe, Theta-modulated place-by-direction
cells in the hippocampal formation in the rat. *J. Neurosci.* **24**, 8265–8277 (2004).

37. N. Burgess, F. Cacucci, C. Lever, J. O'Keefe, Characterizing multiple independent behavioral
correlates of cell firing in freely moving animals. *Hippocampus* **15**, 149–153 (2005).

38. W. Skaggs, B. McNaughton, K. Gothard, "An information-theoretic approach to deciphering the
hippocampal code" in *Advances in Neural Information Processing Systems*, S. Hanson, J.
Cowan, C. Giles, Eds. (Morgan-Kaufmann, 1992)vol. 5.

39. E. Batschelet, *Circular Statistics in Biology* (Academic Press, 1981)*Mathematics in biology*.

40. J. P. Goodridge, J. S. Taube, Interaction between the postsubiculum and anterior thalamus in the
generation of head direction cell activity. *J. Neurosci.* **17**, 9315–9330 (1997).

41. J. S. Taube, R. U. Muller, J. B. Ranck, Head-direction cells recorded from the postsubiculum in
freely moving rats. I. Description and quantitative analysis. *J. Neurosci.* **10**, 420–435 (1990).

42. A. Peyrache, N. Schieferstein, G. Buzsáki, Transformation of the head-direction signal into a
spatial code. *Nat. Commun.* **8**, 1752 (2017).

43. C. Wang, X. Chen, H. Lee, S. S. Deshmukh, D. Yoganarasimha, F. Savelli, J. J. Knierim,
Egocentric coding of external items in the lateral entorhinal cortex. *Science* **362**, 945–949
(2018).

44. P. A. LaChance, T. P. Todd, J. S. Taube, A sense of space in postrhinal cortex. *Science* **365**,
eaax4192 (2019).

45. A. S. Alexander, L. C. Carstensen, J. R. Hinman, F. Raudies, G. W. Chapman, M. E. Hasselmo,
Egocentric boundary vector tuning of the retrosplenial cortex. *Sci. Adv.* **6**, eaaz2322.

46. A. Sarel, A. Finkelstein, L. Las, N. Ulanovsky, Vectorial representation of spatial goals in the hi
ppocampus of bats. *Science* **355**, 176–180 (2017).

47. L. Kunz, A. Brandt, P. C. Reinacher, B. P. Staresina, E. T. Reifenstein, C. T. Weidemann, N. A.
Herweg, A. Patel, M. Tsitsiklis, R. Kempter, M. J. Kahana, A. Schulze-Bonhage, J. Jacobs, A
neural code for egocentric spatial maps in the human medial temporal lobe. *Neuron* **109**, 2781-
2796.e10 (2021).

48. J. Ormond, J. O'Keefe, Hippocampal place cells have goal-oriented vector fields during
navigation. *Nature* **607**, 741–746 (2022).

49. A. Jeewajee, C. Barry, J. O'Keefe, N. Burgess, Grid cells and theta as oscillatory interference:
electrophysiological data from freely moving rats. *Hippocampus* **18**, 1175–1185 (2008).

50. S. S. Deshmukh, D. Yoganarasimha, H. Voicu, J. J. Knierim, Theta modulation in the medial and
the lateral entorhinal cortices. *J. Neurophysiol.* **104**, 994–1006 (2010).

51. M. M. Yartsev, M. P. Witter, N. Ulanovsky, Grid cells without theta oscillations in the entorhinal
cortex of bats. *Nature* **479**, 103–107 (2011).

52. M. Vinck, M. Van Wingerden, T. Womelsdorf, P. Fries, C. M. A. Pennartz, The pairwise phase
consistency: a bias-free measure of rhythmic neuronal synchronization. *Neuroimage* **51**, 112–122
(2010).

53. E. Maris, R. Oostenveld, Nonparametric statistical testing of EEG- and MEG-data. *J. Neurosci.*
Methods **164**, 177–190 (2007).

54. C. W. J. Granger, Testing for causality. *J. Econom. Dynam. Control* **2**, 329–352 (1980).

55. J. Geweke, Measurement of linear dependence and feedback between multiple time series. *J. Am.*
Stat. Assoc. **77**, 304–313 (1982).

56. M. Dhamala, G. Rangarajan, M. Ding, Analyzing information flow in brain networks with
nonparametric granger causality. *Neuroimage* **41**, 354–362 (2008).

57. Y. B. Saalmann, M. A. Pinsk, L. Wang, X. Li, S. Kastner, The pulvinar regulates information
transmission between cortical areas based on attention demands. *Science* **337**, 753–756 (2012).

58. A. M. Bastos, J. Vezoli, C. A. Bosman, J.-M. Schoffelen, R. Oostenveld, J. R. Dowdall, P. De
Weerd, H. Kennedy, P. Fries, Visual areas exert feedforward and feedback influences through
distinct frequency channels. *Neuron* **85**, 390–401 (2015).

59. G. Secer, J. J. Knierim, N. J. Cowan, Explicit error coding can mediate gain recalibration in
continuous bump attractor networks. *Nat Commun* **16**, 9215 (2025).

60. D. M. Wolpert, R. C. Miall, M. Kawato, Internal models in the cerebellum. *Trends in Cognitive*
Sciences **2**, 338–347 (1998).

61. I. Groves, S. L. Grella, C. W. Harley, O. Hardt, L. Nadel, How prediction error drives memory
updating: role of locus coeruleus–hippocampal interactions. *Trends in Neurosciences* **48**, 865–
876 (2025).

62. R. P. Jayakumar, M. S. Madhav, F. Savelli, H. T. Blair, N. J. Cowan, J. J. Knierim, Recalibration
of path integration in hippocampal place cells. *Nature* **566**, 533–537 (2019).

63. K. Brodmann, *Vergleichende Lokalisationslehre Der Grosshirnrinde : In Ihren Prinzipien*
Dargestellt Auf Grund Des Zellenbaues (Barth, Leipzig, 2 unveränderte Aufl., 1909).

64. W. J. S. Krieg, Connections of the cerebral cortex. I. The albino rat. A. Topography of the
cortical areas. *J. Comp. Neurol.* **84**, 221–275 (1946).

65. W. J. S. Krieg, Connections of the cerebral cortex. I. The albino rat. B. Structure of the cortical
areas. *J. Comp. Neurol.* **84**, 277–323 (1946).

66. J. E. Rose, C. N. Woolsey, Structure and relations of limbic cortex and anterior thalamic nuclei
in rabbit and cat. *J. Comp. Neurol.* **89**, 279–347 (1948).

67. R. W. Smith, L. E. White Jr., The fiberarchitectonics of the cat hippocampal formation. *J. Comp.*
Neurol. **123**, 11–27 (1964).

68. H. Stephan, Ed., *Allocortex* (Springer Berlin Heidelberg, Berlin, Heidelberg, 1975)vol. 4/9 of
Handbuch der mikroskopischen Anatomie des Menschen.

69. N. L. M. Cappaert, N. M. Van Strien, M. P. Witter, "Hippocampal formation" in *The Rat*
Nervous System (Elsevier, 2015;
https://linkinghub.elsevier.com/retrieve/pii/B9780123742452000206), pp. 511–573.

70. T. W. Blackstad, Commissural connections of the hippocampal region in the rat, with special
reference to their mode of termination. *J. Comp. Neurol.* **105**, 417–537 (1956).

71. F.-M. Š. Haug, *Sulphide Silver Pattern and Cytoarchitectonics of Parahippocampal Areas in the*
Rat (Springer Berlin Heidelberg, Berlin, Heidelberg, 1976; http://link.springer.com/10.1007/978-
3-642-66448-9).

72. A. V. Ferreira, The cortical areas of the albino rat studied by silver impregnation. *J. Comp.*
Neurol. **95**, 177–243 (1951).

73. S. Chen, C. Chen, X. Xiang, S. Zhang, S. Ding, Chemoarchitecture of area prostriata in adult and
developing mice: comparison with presubiculum and parasubiculum. *J. Comp. Neurol.* **530**,
2486–2517 (2022).

74. K. S. Rockland, Visual system: prostriata — a visual area off the beaten path. *Curr. Biol.* **22**,
R571–R573 (2012).

75. M. Tamietto, D. A. Leopold, Visual cortex: the eccentric area prostriata in the human brain.
Curr. Biol. **28**, R17–R19 (2018).

76. B. L. McNaughton, C. A. Barnes, J. L. Gerrard, K. Gothard, M. W. Jung, J. J. Knierim, H.
Kudrimoti, Y. Qin, W. E. Skaggs, M. Suster, K. L. Weaver, Deciphering The Hippocampal
Polyglot: the Hippocampus as a Path Integration System. *J Exp Biol* **199**, 173–185 (1996).

77. J. J. Knierim, D. A. Hamilton, Framing Spatial Cognition: Neural Representations of Proximal
and Distal Frames of Reference and Their Roles in Navigation. *Physiological Reviews* **91**, 1245–
1279 (2011).

78. Y.-Q. Zhou, V. Puliyadi, X. Chen, J. L. Lee, L.-Y. Zhang, J. J. Knierim, Vector coding and place
coding in hippocampus share a common directional signal. *Nat Commun* **15**, 10630 (2024).

79. F. C. Fortenbaugh, J. C. Hicks, L. Hao, K. A. Turano, Losing sight of the bigger picture:
Peripheral field loss compresses representations of space. *Vision Research* **47**, 2506–2520
(2007).

80. F. C. Fortenbaugh, J. C. Hicks, K. A. Turano, The Effect of Peripheral Visual Field Loss on
Representations of Space: Evidence for Distortion and Adaptation. *Invest. Ophthalmol. Vis. Sci.*
49, 2765–2772 (2008).

81. E. M. Barhorst-Cates, K. M. Rand, S. H. Creem-Regehr, The Effects of Restricted Peripheral
Field-of-View on Spatial Learning while Navigating. *PLOS ONE* **11**, e0163785 (2016).

82. Q. Perrenoud, A. H. de O. Fonseca, A. Airhart, J. Bonanno, R. Mao, J. A. Cardin, Flexible
perceptual encoding by discrete gamma events. *Nature* **647**, 962–969 (2025).

83. J. S. Street, K. J. Jeffery, The dorsal thalamic lateral geniculate nucleus is required for visual
control of head direction cell firing direction in rats. *The Journal of Physiology* **602**, 5247–5267
(2024).

84. E. I. Moser, E. Kropff, M.-B. Moser, Place cells, grid cells, and the brain's spatial representation
system. *Annu. Rev. Neurosci.* **31**, 69–89 (2008).

85. R. J. Gardner, E. Hermansen, M. Pachitariu, Y. Burak, N. A. Baas, B. A. Dunn, M.-B. Moser, E.
I. Moser, Toroidal topology of population activity in grid cells. *Nature* **602**, 123–128 (2022).

86. C. B. Canto, N. Koganezawa, P. Beed, E. I. Moser, M. P. Witter, All layers of medial entorhinal
cortex receive presubicular and parasubicular inputs. *J. Neurosci.* **32**, 17620–17631 (2012).

87. P. Preston-Ferrer, S. Coletta, M. Frey, A. Burgalossi, Anatomical organization of presubicular
head-direction circuits. *eLife* **5**, e14592 (2016).

88. R. M. Yoder, B. J. Clark, J. S. Taube, Origins of landmark encoding in the brain. *Trends*
Neurosci. **34**, 561–571 (2011).

89. A. S. Alexander, D. A. Nitz, Retrosplenial cortex maps the conjunction of internal and external spaces. *Nat. Neurosci.* **18**, 1143–1151 (2015).

90. P. A. LaChance, J. Graham, B. L. Shapiro, A. J. Morris, J. S. Taube, Landmark-modulated directional coding in postrhinal cortex. *Sci. Adv.* **8**, eabg8404 (2022).

91. R. P. N. Rao, D. H. Ballard, Predictive coding in the visual cortex: a functional interpretation of some extra-classical receptive-field effects. *Nat Neurosci* **2**, 79–87 (1999).

92. K. Friston, A theory of cortical responses. *Phil. Trans. R. Soc. B* **360**, 815–836 (2005).

93. A. M. Bastos, W. M. Usrey, R. A. Adams, G. R. Mangun, P. Fries, K. J. Friston, Canonical microcircuits for predictive coding. *Neuron* **76**, 695–711 (2012).

94. S. Shipp, Neural Elements for Predictive Coding. *Front. Psychol.* **7** (2016).

95. G. B. Keller, T. D. Mrsic-Flogel, Predictive Processing: A Canonical Cortical Computation. *Neuron* **100**, 424–435 (2018).

96. T. J. Buschman, E. K. Miller, Top-Down Versus Bottom-Up Control of Attention in the Prefrontal and Posterior Parietal Cortices. *Science* **315**, 1860–1862 (2007).

97. G. Michalareas, J. Vezoli, S. van Pelt, J.-M. Schoffelen, H. Kennedy, P. Fries, Alpha-beta and gamma rhythms subserve feedback and feedforward influences among human visual cortical areas. *Neuron* **89**, 384–397 (2016).

98. J. D. Semedo, A. I. Jasper, A. Zandvakili, A. Krishna, A. Aschner, C. K. Machens, A. Kohn, B. M. Yu, Feedforward and feedback interactions between visual cortical areas use different population activity patterns. *Nat. Commun.* **13**, 1099 (2022).

99. T. Van Kerkoerle, M. W. Self, B. Dagnino, M.-A. Gariel-Mathis, J. Poort, C. Van Der Togt, P. R. Roelfsema, Alpha and gamma oscillations characterize feedback and feedforward processing in monkey visual cortex. *Proc. Natl. Acad. Sci.* **111**, 14332–14341 (2014).

100. A. M. Bastos, M. Lundqvist, A. S. Waite, N. Kopell, E. K. Miller, Layer and rhythm specificity for predictive routing. *Proceedings of the National Academy of Sciences* **117**, 31459–31469 (2020).

101. E. L. Hargreaves, D. Yoganarasimha, J. J. Knierim, Cohesiveness of spatial and directional representations recorded from neural ensembles in the anterior thalamus, parasubiculum, medial entorhinal cortex, and hippocampus. *Hippocampus* **17**, 826–841 (2007).

102. R. J. Gardner, L. Lu, T. Wernle, M.-B. Moser, E. I. Moser, Correlation structure of grid cells is preserved during sleep. *Nat Neurosci* **22**, 598–608 (2019).

103. S. G. Trettel, J. B. Trimper, E. Hwaun, I. R. Fiete, L. L. Colgin, Grid cell co-activity patterns during sleep reflect spatial overlap of grid fields during active behaviors. *Nat Neurosci* **22**, 609–617 (2019).

104. N. Almog, G. Tocker, T. Bonnevie, E. I. Moser, M.-B. Moser, D. Derdikman, During hippocampal inactivation, grid cells maintain synchrony, even when the grid pattern is lost. *eLife* **8**, e47147 (2019).

105. T. Waaga, H. Agmon, V. A. Normand, A. Nagelhus, R. J. Gardner, M.-B. Moser, E. I. Moser, Y. Burak, Grid-cell modules remain coordinated when neural activity is dissociated from external sensory cues. *Neuron* **110**, 1843-1856.e6 (2022).

106. M. Quigley, K. Conley, B. Gerkey, J. Faust, T. Foote, J. Leibs, R. Wheeler, A. Y. Ng, "ROS: an
open-source robot operating system" in *ICRA Workshop on Open Source Software* (Kobe, 2009)
vol. 3, p. 5.

107. B. P. Vagvolgyi, R. P. Jayakumar, M. S. Madhav, J. J. Knierim, N. J. Cowan, Wide-angle,
monocular head tracking using passive markers. *J. Neurosci. Methods* **368**, 109453 (2022).

108. P. Andrey, Y. Maurin, *Free-D*: an integrated environment for three-dimensional reconstruction
from serial sections. *J. Neurosci. Methods* **145**, 233–244 (2005).

109. C. N. Boccara, L. J. Kjonigsen, I. M. Hammer, J. G. Bjaalie, T. B. Leergaard, M. P. Witter, A
three-plane architectonic atlas of the rat hippocampal region. *Hippocampus* **25**, 838–857 (2015).

110. J. E. Chung, J. F. Magland, A. H. Barnett, V. M. Tolosa, A. C. Tooker, K. Y. Lee, K. G. Shah, S.
H. Felix, L. M. Frank, L. F. Greengard, A fully automated approach to spike sorting. *Neuron* **95**,
1381-1394.e6 (2017).

111. G. Secer, B. Krishnan, N. J. Cowan, J. J. Knierim, "Automated vs. manual spike sorting:
assessing the quality of automated spike sorting using grid cells" in *2023 Neuroscience Meeting
Planner* (Society for Neuroscience, Washington, D.C.).

112. A. P. Buccino, C. L. Hurwitz, S. Garcia, J. Magland, J. H. Siegle, R. Hurwitz, M. H. Hennig,
SpikeInterface, a unified framework for spike sorting. *eLife* **9**, e61834 (2020).

113. T. Chen, C. Guestrin, "XGBoost: a scalable tree boosting system" in *Proceedings of the 22nd
ACM SIGKDD International Conference on Knowledge Discovery and Data Mining* (ACM, San
Francisco California USA, 2016; https://dl.acm.org/doi/10.1145/2939672.2939785), pp. 785–
794.

114. S. Royer, B. V. Zemelman, A. Losonczy, J. Kim, F. Chance, J. C. Magee, G. Buzsáki, Control of
timing, rate and bursts of hippocampal place cells by dendritic and somatic inhibition. *Nat.
Neurosci.* **15**, 769–775 (2012).

115. J. Platt, Probabilistic outputs for support vector machines and comparisons to regularized
likelihood methods. *Adv, Large Margin Classif,* **10** (2000).

116. A. Niculescu-Mizil, R. Caruana, "Predicting good probabilities with supervised learning" in
Proceedings of the 22nd International Conference on Machine Learning (Association for
Computing Machinery, New York, NY, USA, 2005)*ICML '05*, pp. 625–632.

117. D. H. Wolpert, Stacked generalization. *Neural Networks* **5**, 241–259 (1992).

118. M. Zaheer, S. Kottur, S. Ravanbakhsh, B. Poczos, R. R. Salakhutdinov, A. J. Smola, "Deep sets"
in *Advances in Neural Information Processing Systems* (Curran Associates, Inc., 2017)vol. 30.

119. C. Chow, On optimum recognition error and reject tradeoff. *IEEE Trans, Inf, Theor,* **16**, 41–46
(1970).

120. R. El-Yaniv, Y. Wiener, On the foundations of noise-free selective classification. *J. Mach.
Learn. Res.* **11**, 1605–1641 (2010).

121. K. Miettinen, *Nonlinear Multiobjective Optimization* (Springer US, Boston, MA, 1998;
http://link.springer.com/10.1007/978-1-4615-5563-6)vol. 12 of *International Series in
Operations Research & Management Science*.

122. R. T. Marler, J. S. Arora, Survey of multi-objective optimization methods for engineering.
Structural and Multidisciplinary Optimization **26**, 369–395 (2004).

123. A. A. Emondi, S. P. Rebrik, A. V. Kurgansky, K. D. Miller, Tracking neurons recorded from
tetrodes across time. *Journal of Neuroscience Methods* **135**, 95–105 (2004).

124. S. S. Shapiro, M. B. Wilk, An analysis of variance test for normality (complete samples).
Biometrika **52**, 591–611 (1965).

125. S. S. Shapiro, R. S. Francia, An approximate analysis of variance test for normality. *J. Am. Stat.*
Assoc. **67**, 215–216 (1972).

126. J. P. Royston, An extension of shapiro and wilk's W test for normality to large samples. *Appl.*
Stat. **31**, 115 (1982).

127. P. Royston, Approximating the shapiro-wilk W-test for non-normality. *Stat. Comput.* **2**, 117–119
(1992).

128. S. G. Luke, Evaluating significance in linear mixed-effects models in R. *Behav Res* **49**, 1494–
1502 (2017).

129. A. Kuznetsova, P. B. Brockhoff, R. H. B. Christensen, **lmerTest** Package: Tests in Linear Mixed
Effects Models. *J. Stat. Soft.* **82** (2017).

130. T. Donoghue, M. Haller, E. J. Peterson, P. Varma, P. Sebastian, R. Gao, T. Noto, A. H. Lara, J.
D. Wallis, R. T. Knight, A. Shestyuk, B. Voytek, Parameterizing neural power spectra into
periodic and aperiodic components. *Nat. Neurosci.* **23**, 1655–1665 (2020).

131. G. Kühn, H. E. Rashidy, F. Calegari, S. Dhingra, Aperiodicity in mouse CA1 and DG power
spectra. bioRxiv [Preprint] (2025). https://doi.org/10.1101/2025.01.30.635678.

132. M. Lundqvist, P. Herman, M. R. Warden, S. L. Brincat, E. K. Miller, Gamma and beta bursts
during working memory readout suggest roles in its volitional control. *Nat Commun* **9**, 394
(2018).

133. S. Abbaspoor, A. T. Hussin, K. L. Hoffman, Theta- and gamma-band oscillatory uncoupling in
the macaque hippocampus. *eLife* **12**, e86548 (2023).

134. J. Daume, J. Kamiński, A. G. P. Schjetnan, Y. Salimpour, U. Khan, M. Kyzar, C. M. Reed, W. S.
Anderson, T. A. Valiante, A. N. Mamelak, U. Rutishauser, Control of working memory by
phase–amplitude coupling of human hippocampal neurons. *Nature* **629**, 393–401 (2024).

135. P. Hegedüs, J. Heckenast, B. Hangya, Differential recruitment of ventral pallidal e-types by
behaviorally salient stimuli during pavlovian conditioning. *iScience* **24** (2021).

136. R. Oostenveld, P. Fries, E. Maris, J.-M. Schoffelen, FieldTrip: open source software for
advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput. Intell.*
Neurosci. **2011**, 156869 (2011).

137. J. D. Monaco, G. Rao, E. D. Roth, J. J. Knierim, Attentive scanning behavior drives one-trial
potentiation of hippocampal place fields. *Nat. Neurosci.* **17**, 725–731 (2014).

138. A. V. Olypher, P. Lánský, R. U. Muller, A. A. Fenton, Quantifying location-specific information
in the discharge of rat hippocampal place cells. *J. Neurosci. Methods* **127**, 123–135 (2003).

139. C. E. Shannon, A mathematical theory of communication. *Bell Syst. Tech. J.* **27**, 379–423
(1948).

140. B. C. Souza, R. Pavão, H. Belchior, A. B. L. Tort, On information metrics for spatial coding.
Neuroscience **375**, 62–73 (2018).

141.   M. Ester, H.-P. Kriegel, J. Sander, X. Xu, "A density-based algorithm for discovering clusters in
large spatial databases with noise" in *Proceedings of the Second International Conference on
Knowledge Discovery and Data Mining* (AAAI Press, Portland, Oregon, 1996)*KDD'96*, pp.
226–231.

142.   K. Hardcastle, N. Maheswaranathan, S. Ganguli, L. M. Giocomo, A multiplexed, heterogeneous,
and adaptive code for navigation in medial entorhinal cortex. *Neuron* **94**, 375-387.e7 (2017).

143.   G. Carter, C. Knapp, A. Nuttall, Estimation of the magnitude-squared coherence function via
overlapped fast fourier transform processing. *IEEE Trans. Audio Electroacoust.* **21**, 337–344
(1973).

144.   M. R. Jarvis, P. P. Mitra, Sampling properties of the spectrum and coherency of sequences of
action potentials. *Neural Comput.* **13**, 717–749 (2001).

145.   G. Nolte, O. Bai, L. Wheaton, Z. Mari, S. Vorbach, M. Hallett, Identifying true brain interaction
from EEG data using the imaginary part of coherency. *Clin. Neurophysiol.* **115**, 2292–2307
(2004).

146.   N. Karalis, C. Dejean, F. Chaudun, S. Khoder, R. R. Rozeske, H. Wurtz, S. Bagur, K.
Benchenane, A. Sirota, J. Courtin, C. Herry, 4-hz oscillations synchronize prefrontal–amygdala
circuits during fear behavior. *Nat. Neurosci.* **19**, 605–612 (2016).

147.   X. Xu, L. Song, R. Kringel, I. L. Hanganu-Opatz, Developmental decrease of entorhinal-
hippocampal communication in immune-challenged DISC1 knockdown mice. *Nat Commun* **12**,
6810 (2021).

148.   Y.-N. Chen, J. K. Kostka, S. H. Bitzenhofer, I. L. Hanganu-Opatz, Olfactory bulb activity shapes
the development of entorhinal-hippocampal coupling and associated cognitive abilities. *Current
Biology* **33**, 4353-4366.e5 (2023).

149.   A. Fernández-Ruiz, A. Oliva, M. Soula, F. Rocha-Almeida, G. A. Nagy, G. Martin-Vazquez, G.
Buzsáki, Gamma rhythm communication between entorhinal cortex and dentate gyrus neuronal
assemblies. *Science* **372**, eabf3119 (2021).

150.   V. Saravanan, G. J. Berman, S. J. Sober, Application of the hierarchical bootstrap to multi-level
data in neuroscience. *Neurons Behav. DATA Anal. Theory* **3**, 0–29 (2020).

151.   A. M. Bastos, J.-M. Schoffelen, A tutorial review of functional connectivity analysis methods
and their interpretational pitfalls. *Front. Syst. Neurosci.* **9**, 0–23 (2016).

152.   J. F. Geweke, Measures of conditional linear dependence and feedback between time series. *J.
Am. Stat. Assoc.* **79**, 907–915 (1984).

153.   M. Ding, Y. Chen, S. L. Bressler, "Granger causality: basic theory and application to
neuroscience" in *Handbook of Time Series Analysis*, B. Schelter, M. Winterhalder, J. Timmer,
Eds. (Wiley, ed. 1, 2006; https://onlinelibrary.wiley.com/doi/10.1002/9783527609970.ch17), pp.
437–460.

154.   Y. Chen, S. L. Bressler, M. Ding, Frequency decomposition of conditional granger causality and
application to multivariate neural field potential data. *J. Neurosci. Methods* **150**, 228–237 (2006).

155.   R. Kobayashi, S. Kurita, A. Kurth, K. Kitano, K. Mizuseki, M. Diesmann, B. J. Richmond, S.
Shinomoto, Reconstructing neuronal circuitry from parallel spike trains. *Nat. Commun.* **10**, 4468
(2019).

156.   D. Endo, R. Kobayashi, R. Bartolo, B. B. Averbeck, Y. Sugase-Miyamoto, K. Hayashi, K. Kawano, B. J. Richmond, S. Shinomoto, A convolutional neural network for estimating synaptic connectivity from spike trains. *Sci. Rep.* **11**, 12087 (2021).

Acknowledgments

We thank Rahul Jakati, Ravikrishnan Jayakumar, Shahin Lashkari, Francesco Savelli, Manu Madhav, Katelan Balkissoon, Bethel Gebreselassie and Makenna Dixon for useful discussions and assistance with the experiments. We thank Daeyeol Lee and Sung Soo Kim for providing valuable comments on the manuscript.

Funding:

National Institutes of Health grant R01 NS102537 (NJC, JJK)
National Institutes of Health grant R01 MH118926 (JJK, NJC)
Johns Hopkins Kavli Neuroscience Discovery Institute (NDI) Distinguished Postdoctoral Fellowship (GS)
Johns Hopkins Kavli Neuroscience Discovery Institute (NDI) Distinguished Doctoral Fellowship (BK)
National Institutes of Health grant T32GM007057 (BK)

Author contributions:

Conceptualization: GS, BK, NJC, JJK
Experiments: GS, BK
Data analysis: GS, BK
Writing: GS, BK, NJC, JJK
Funding acquisition: GS, BK, NJC, JJK
Supervision: NJC, JJK

Competing interests: Authors declare that they have no competing interests.

Materials and Methods

Subjects

Five adult Long–Evans rats (Envigo Harlan), 3 M (994, 995, and 1041) and 2 F (1061 and 1062), served as subjects. At the time of surgery, males weighed 440–470 g and were 6–9 months old, whereas females weighed 315–325 g and were 15–18 months old. Animals were housed individually in a temperature- and humidity-controlled room on a 12 h:12 h light/dark cycle; all behavioral training and electrophysiological recordings were carried out during the dark phase. Rats were handled nearly daily throughout habituation, training, surgery, and data-collection periods. Data from all five animals were pooled for all subsequent analyses, but individual rat data are shown in supplementary tables. Potential sex differences were confounded by non-uniform sampling of cell layers across animals, and small sample sizes for each sex prevented meaningful assessment of se as a biological variable. All procedures conformed to the National Institutes of Health guidelines and were approved by the Johns Hopkins University Institutional Animal Care and Use Committee.

The Dome apparatus

The virtual-reality Dome (*12*, *33*) is a planetarium-style setup built around a hemispherical fiberglass shell with a 2.3-m inner diameter. A 15-cm aperture at the apex allowed light emitted from a ceiling-mounted projector (Sony VPL-FH30) fitted with a long-throw lens (Navitar ZM 70–125 mm). A 25-cm hemispherical first-surface mirror (JR Cumberland), mounted on a central pillar at the geometric center, reflected the projected image onto the inner surface of the Dome to create full-field visual stimulation. Visual scenes were rendered in C++ using OpenGL and pre-warped to compensate for the projection optics before display. To provide uniform, rotationally symmetric background illumination that did not overlap the landmarks, a narrow overhead ring was projected continuously, including during landmark-absent epochs. Behavior was monitored with an overhead near-infrared camera providing real-time, top-down video (see "*Experiment control and behavior tracking*").

Rats locomoted along the outer edge of an annular table (outer diameter 152.4 cm, inner diameter 45.7 cm) positioned at the center of the Dome. An enclosure subtending 45° around the rat was attached to a motorized central pillar via radial boom arms to ensure that the animal remained near the table's outer rim and to carry the transmitter unit of the neural recording system (in wireless recording sessions) or the headstage and Litz tethers (for wired recording sessions). The motorized central pillar incrementally advanced the enclosure to keep the rat near a predefined setpoint within the enclosure; it engaged only when the rat was within a defined window around that setpoint, ensuring the enclosure moved along with the rat. Liquid rewards (Ensure diluted 50% with water) were delivered pseudo randomly onto the table via a feed tube mounted on a boom arm diametrically opposite the enclosure; delivery timing and volume were controlled by a micro-peristaltic pump (Takasago fluidics) on the central pillar using custom software (*33*). This boom arm also held a plastic spreader and paper towels to disperse urine and food odors and to remove feces, minimizing the salience and stability of local olfactory cues. After each session the track was thoroughly wiped and then rotated by a multiple of 45° (one of eight possible orientations), altering the spatial relationship of any tactile cues to the rest of the apparatus. This prevented local tactile features on the annular track from becoming stable global landmarks.

Closed-loop gain manipulation experiments in the Dome

During closed-loop gain manipulation sessions, the locations of visual landmarks (3 or 4 landmarks, depending on the session; Supplementary table 1) were updated in closed loop with the rat's movement according to an experimental gain, G, defined as the ratio between landmark-frame and laboratory-frame speed of the animal. During these manipulations, landmarks were moved in unison, maintaining their relative configuration, and rotating as a coherent ensemble. Landmarks were stationary when G = 1.

They were rotated along the circular track in the same direction as the animal when $0 < G < 1$ or in the
opposite direction when $G > 1$.

Every gain-manipulation session comprised three epochs.

• **Epoch 1**: $G = 1$ (veridical condition). During this Epoch, the animal ran 8–15 laps with
stationary landmarks.

• **Epoch 2**: G was ramped linearly at a rate of 1/52 per lap until it reached its final value, G_{final}.

• **Epoch 3**: $G = G_{final}$ remained constant.

After the final epoch (Epoch 3), all landmarks were extinguished, leaving only a dim, non-polarizing
ring for illumination.

The final gain value, G_{final}, took the form of $1 \pm n/13$ ($n = 0, 1, 2, 3, 4\ldots$), so that positions in the
laboratory and landmark frames coincided only once every 13 laps when $n > 0$. Each rat first completed
one to three baseline sessions with $G_{final} = 1$; thereafter the gain was systematically altered in following
sessions. Sessions with more extreme values ($G_{final} \gg 1$ or $G_{final} \ll 1$) often showed landmark-control
failure. Thus, to maximize the number of sessions that retained strong landmark control, more extreme
values of G_{final} were chosen progressively across days and were seldom repeated on consecutive
sessions. Experimenters were aware of the gain during data collection and analysis, and sample sizes
were determined empirically without formal power calculations.

Luminance manipulation experiments in the Dome

Landmarks and the background were implemented as separate OpenGL objects, allowing independent
control of their luminance by scaling object intensities. Intensities were normalized to the projector's
dynamic range (0–1). Landmark (LM) luminance took values 1.00 (High) and 0.55 (Low); background
(BG) luminance took values 0.47 (High) and 0.01 (Low). By providing greater LM luminance than BG
luminance, these values ensured no "negative contrast" conditions (i.e., landmarks being darker than the
background).

During luminance manipulation blocks consisted of equal-length epochs where landmarks were
stationary (experiment gain $G = 1$). Within a given epoch, LM and BG luminance were held constant.
The four illumination conditions (2 LM luminance levels × 2 BG luminance levels) were presented
across 6-lap epochs in pseudorandom order predetermined by the experimenter, with the constraint that
no condition was repeated in consecutive epochs and all conditions were experienced an equal number
of times. Across a session, the luminance conditions changed only at epoch boundaries, and each of the
four conditions were presented at least three times.

At the end of the luminance epochs, the experiment gain was abruptly changed to an extreme value (i.e.,
much larger or smaller than 1), which induced landmark failure, allowing us to test the landmark
anchoring strength for all cells, as described in the "Landmark-locking score" subsection of Methods.

Open-loop gain manipulation experiments in the Dome

To dissociate landmark motion from the animal's self-motion in open loop experiments, visual
landmark movement was controlled by a pre-recorded animal motion trajectory obtained from a
different session. The reference trajectory (recorded as a ROS .bag file) contained the animal's angular
position along the circular track as a function of time. During open-loop sessions, this trajectory was
replayed in real time and used to control landmark motion via the same experiment gain (G) parameter
used in closed-loop experiments. If the open-loop session duration outlasted the reference trajectory

duration, the recorded animal trajectory was looped from its beginning without interruption to provide continuous control of landmark position.

Each open-loop session followed a standardized gain protocol.

- **Epoch 1:** G = 1, during which landmarks were stationary for at least 8 laps (same as closed-loop).

- **Epoch 2:** G was ramped from 1 to a target value, as in closed-loop sessions. However, unlike closed-loop sessions where G was applied to the animal's own movement, here G was applied to a pre-recorded trajectory from a reference session, thereby dissociating landmark motion from the animal's own movement in the current session.

- **Epoch 3:** G underwent a series of step changes ("jumps") to non-unity gain values across 2–4 sub-epochs, each sub-epoch lasting at least 8 laps. The set of gains across sub-epochs always included at least one value G > 1 and at least one value G < 1, ensuring the rat experienced landmarks moving in both directions relative to its own movement. As in Epoch 2, G was applied to the pre-recorded trajectory to move landmarks independently of the animal's actual motion in the current session. Landmarks were extinguished at the end of the session.

Open arena apparatus

During open-field foraging sessions, each rat foraged for randomly distributed precision chocolate pellets (Bio-Serv Dustless Precision Pellets, Purified), on a square platform that measured 137 × 137 cm and was surrounded by opaque walls 36.5 cm high. Experiments were conducted in two different recording rooms. In the larger room (approximately 330 × 700 cm) neural activity was transmitted wirelessly via a FreeLynx device, whereas in the second room (292 × 292 cm) signals were carried through a tether connected to a DigitalLynx SX system and Saturn commutator (Neuralynx, Inc.). Both rooms were dimly illuminated by overhead white LEDs and contained fixed distal cues—such as shelving, computers, and other laboratory equipment—that were kept in the same positions across days to provide stable landmarks for the animals. After every session the arena floor was wiped with 70 % ethanol. Any urine or feces were removed immediately during the session once the rat had moved away from that spot. Throughout each 30-40 min session the experimenter walked outside the arena while dispensing the chocolate precision pellets.

Training

Rats were handled daily for 3-5 days initially to habituate them to human contact, then placed on a controlled feeding schedule to reduce body weight to approximately 80% of their *ad libitum* weight.

Dome training: Rats were first trained on a circular table in a room separate from the Dome apparatus. The table matched the Dome table in diameter but lacked the enclosure and automation (see "*Dome apparatus*"). Reward droplets (1:1 mixture of water and chocolate Ensure) were placed manually at arbitrary locations on the track, and the interval between rewards gradually increased to sustain running and delay satiation. Training continued until the rats consistently completed at least 80 laps in the same direction without experimenter intervention, typically after 3–4 weeks. The rats were then introduced to the Dome, which contained stationary visual landmarks, and were trained there until their lap counts matched those achieved on the open table, usually within one additional week. After hyperdrive implantation, the rats were retrained in the Dome and transitioned to automated reward delivery, in which drop timing varied to encourage continuous forward running with minimal pauses. Experiments began once the tetrodes reached the target recording regions and the rats consistently ran many laps.

Open-arena training: We acclimated rats to the open arena by giving them 2–3 consecutive days of free foraging for Bio-Serv Dustless Precision Pellets, intended to promote spontaneous exploration. After 1–2 weeks they explored the entire arena readily and uniformly. After hyperdrive implantation, training continued until the rat, with the transmitter in place, traversed the arena broadly and almost continuously, pausing only briefly—a criterion typically achieved within 7–14 days after surgery. Experiments began once this behavioral criterion, the Dome-training criterion, and accurate tetrode placement had all been met.

Surgery and electrode implantation

After behavioral training in the Dome and open arena achieved desired behavioral performance, rats were implanted with custom hyperdrives carrying 18 independently drivable tetrodes. Each tetrode was composed of four twisted, heat-bonded nichrome wires (0.0017″ diameter, VG/HML insulation; California Fine Wire Co., CA, USA). Hyperdrives were built in-house using a custom design around a 72-channel interface board (EIB-72-QC; Neuralynx, Bozeman, MT, USA). During the 5–6 days preceding surgery, tetrode tips were gold-plated daily with a nanoZ system (White Matter LLC, Seattle, WA, USA) to reduce and stabilize impedance from 2–3 MΩ to ~130 kΩ. Each assembled hyperdrive was sterilized in ethylene-oxide gas (12-h cycle) and allowed to aerate for 36 h before implantation.

Anesthesia prior to surgery was induced with 4% isoflurane in oxygen (1 L/min) followed by ketamine (80 mg/kg) and xylazine (10 mg/kg); anesthesia was maintained with isoflurane as needed. Local analgesia was provided by applying 0.15 mL of bupivacaine (Marcaine) or ropivacaine directly to the skull. Postoperatively, meloxicam (1–2 mg/kg; Ostilox, Meloxidyl, or Metacam) was administered orally for two days. Postoperative antibiotics (30 mg tetracycline and 0.15 mL of a 22.7% enrofloxacin solution) were given orally once daily for 10 days post-surgery. Under isoflurane anesthesia, a craniotomy was made over the right hemisphere, the rostral edge of the transverse sinus was exposed, and the dura was removed. The hyperdrive was positioned so that the posterior-most central tetrode entered the brain 350–530 μm anterior to the rostral edge of the transverse sinus and 4.30–4.55 mm lateral to bregma, yielding a mediolateral recording span of 3.4–5.45 mm lateral to bregma across animals. To increase effective tetrode travel within MEC, the hyperdrive was tilted 10° anteriorly in the sagittal plane. The craniotomy was then sealed with Kwik-Sil (World Precision Instruments). Hyperdrives were anchored to the skull with jeweler's screws; a screw over the cerebellum served as the recording ground. The implant was secured with a three-layer build of OptiBond (Kerr Dental), Charisma (Kulzer), and dental cement.

Neural recordings

Wideband signals were digitized at 30 kHz and acquired on a computer running Neuralynx Cheetah software using either a wireless FreeLynx (Neuralynx) transmitter or, for wired sessions, a Litz tether connected to a Digital Lynx SX system by a unity gain 72 channel headstage (HS72QC, Neuralynx) attached to the implanted hyperdrive. Both Dome and open arena experiments were run with either wired or wireless configurations of this recording system. TTL pulses from the behavioral control computer (see Experimental control) were recorded as timestamped events on the acquisition system to enable post hoc synchronization of behavioral and neural data. Raw wideband data were saved as Neuralynx .nrd files and used for spike sorting.

Experiment control and behavior tracking

Dome sessions were run using three computers: (1) an experiment-control PC, (2) a neural-acquisition PC running Cheetah (Neuralynx, Inc.), and (3) a video-tracking PC. Task logic and device control were implemented as Robot Operating System (*106*) nodes communicating continuously with a master node on the experiment-control PC. Synchronization was achieved by sending pseudorandom TTL pulses

from the control PC to the acquisition system; these were time-stamped in Cheetah and used to align
data streams across machines. Hardware/software integration and control details are described in a
previous publication (*33*). Head position and 3D head orientation were estimated in real time at 30-45
fps using a single-camera, ROS-based tracker that used custom software and fiducial markers mounted
on the hyperdrive in a known 3D configuration (*107*).

Open-arena sessions typically used two computers: one for neural acquisition with FreeLynx/Cheetah
and one for behavior tracking running the single-camera 3D tracker. Synchronization between the neural
acquisition computer and the behavior tracking computer was implemented using the Domain Time II
software (Greyware Automation Products, Inc.). In a subset of sessions recorded with the wired Digital
Lynx SX system, a single computer handled both neural acquisition and behavioral tracking through the
Cheetah software (Neuralynx, Inc.); in those sessions, head position and 2D head orientation were
extracted from two LEDs mounted on the headstage.

Histological identification

After the final recording sessions, rats were transcardially perfused with 3.7% formalin. To improve the
quality of tetrode tracks, the brain was partially exposed and soaked in PFA or formalin overnight with
the hyperdrive attached before tetrodes were retracted. The brain was extracted and stored in 30%
sucrose-formalin solution until fully submerged, and sectioned sagittally at 40 µm intervals. The
sections were mounted and stained with 0.1% cresyl violet.

For tetrode track reconstruction, Nissl-stained serial sections were imaged on a light microscope, and
whole-section images were acquired with IC Capture 2.1. Images were imported into Free-D (*108*) for
manual tracing of tetrode tracks across consecutive sections and 3D reconstruction of each tetrode's
trajectory. Reconstructed trajectories were visualized from the top-down and registered to the known
hyperdrive bundle geometry to assign tetrode identities. For each identified tetrode, the 3D reconstructed
trajectory was exported as an STL file for further analysis in MATLAB to represent the track as a 3D
curve and to compute arc length (distance along the curve) at each point.

Recording depths were logged for each tetrode in each session as cumulative turns of a 0-80 hyperdrive
screw from its initial position at implantation. To map these depths onto the reconstructed tracks for
each tetrode, we used the most ventral point of each reconstructed track as a reference corresponding to
the deepest recording depth, defining arc length as the distance measured along the track from this point.
Screw-turn values of each tetrode were converted to expected tetrode displacement using the 0-80 screw
pitch (317.5 µm/turn) and adjusted for ~20% tissue shrinkage, yielding an arc-length position along the
reconstructed trajectory for each session's recording depth. For each tetrode, these arc-length positions
were then converted to a specific histological section and an (x, y) coordinate within that section using
the microscope's pixel-to-micron scale (µm/pixel) for that imaging magnification.

For each tetrode, the regional identity of the recording location in each session was determined by
examining the corresponding (x, y) location within the identified histological slice under high
magnification based on cytoarchitectonic features and anatomical (e.g., sulcal) landmarks. Borders
involving area 29e, PR and PA were identified using Haug's Silver-sulfide stained sagittal sections (*71*)
as a reference, whereas borders involving MEC, PA, and PO were identified following previously
published parahippocampal and entorhinal anatomical boundaries (*22, 109*) and a Long-Evans rat atlas
from Menno Witter and colleagues.

High-resolution images of histological sections (Fig. 1A; fig. S1) were acquired on a KEYENCE BZ-
X1000 fluorescence microscope, capturing six brightfield tiles (2×3) with a 4x objective and stitching
them into a single mosaic with the BZ-X Analyzer stitching & multi-point imaging module software.

Spike sorting

Wideband extracellular recordings were band-pass filtered between 300 and 3000 Hz to isolate spiking activity, whitened, and artifact-masked before automatic spike detection and clustering with MountainSort4 (*110*). Outlier events within each cluster were identified in the peak-amplitude feature space of the spike waveforms using MATLAB's *robustcov* function and removed, thereby reducing residual contamination from other sources (*111*). The cleaned clusters were transferred to *Phy* via the *SpikeInterface* framework (*112*) for further manual curation based on spike waveforms, amplitude stability over time, and the distribution of waveforms in principal-component feature space. Inter-spike-interval histograms and pairwise cross-correlograms were also examined to decide whether clusters should be merged or split: clusters were split when mixed waveform populations were evident and merged when two clusters displayed nearly identical waveforms together with a pronounced refractory trough in their spike cross-correlogram. Units showing clear amplitude drift or more than 1% of inter-spike intervals shorter than a 1.5 ms refractory period were discarded. The remaining well-isolated, temporally stable clusters were retained for all subsequent analyses. In a subset of sessions, MountainSort4 units were compared to manually sorted clusters using custom software (WinClust, J. J. Knierim) and were found to achieve comparable isolation quality, as indicated by similar waveform features and refractory-period violations.

Selection criteria for independent neuron and LFP samples

To avoid including the same neuron more than once in an analysis, we employed a deliberately conservative, depth-based duplicate-control procedure. Two clusters recorded on different days were judged to be distinct only if the tetrode had been advanced by at least three ⅛-turn screw steps(~119 µm). For each analysis, clusters from all sessions were first pooled by rat and tetrode; the largest subset of clusters that satisfied this constraint was then chosen through an optimization algorithm. The same depth-based constraint was applied to LFP signals: LFPs from the same region were considered as different samples only if they were recorded at least 119 µm apart. Mixed-effect models were used to account for the nonindependence of LFP signals from the same animal. See relevant sections for details.

Machine learning-assisted physiological recording region identification for ambiguous histological cases

Pipeline architecture

In cases where the histological brain-region assignment of a tetrode was ambiguous or infeasible due to histological imperfections, we developed a machine-learning (ML) pipeline to classify the recording region, based on the firing properties of all neurons recorded on that tetrode (fig. S2B). The pipeline was designed to ensure high confidence assignments (i.e., to minimize incorrect assignments to a particular brain region) at the possible expense of classifying neurons as ambiguous. Because histologically ambiguous tetrode recording sites were most often located near the borders of area 29e or PR with neighboring regions, we trained two separate binary classification pipelines: area 29e vs. rest (i.e., MEC, PA, PO and PR) and PR vs. rest (i.e., 29e, MEC, PA, PO). For each pipeline, the "positive class" was defined as neurons that were unambiguously assigned to the target region (area 29e or PR), while the "negative class" included the remaining neurons unambiguously assigned to all other regions. Training data for all pipeline components consisted exclusively of neurons from recording sites with unambiguous histological region assignments (i.e., ground-truth data); neurons from histologically ambiguous recording sites were never used for model development or its performance verification.

The first layer of each pipeline (i.e., 29e vs. rest and the PR vs. rest) comprised an XGBoost model (a boosted decision-tree algorithm; *83*) that performed brain-region classification using the physiological features of single neurons. Features included: (i) spike-waveform characteristics: the peak-aligned

normalized waveform, spike width, and spike asymmetry (trough-to-peak amplitude ratio); (ii) theta-
modulation measures: the mean resultant length (MRL) of spike–theta phases (*50*) and three metrics
quantifying theta rhythmicity (*50, 51, 36*); (iii) firing-pattern features: mean firing rate, burst ratio
(proportion of inter-spike intervals < 8 ms), burst index and refractory period (*114*); and (iv) spike-
autocorrelogram (ACG) properties: mean and mode lag of the ACG, and the power spectrum of the
spike ACG (1–90 Hz). After eliminating highly correlated features in this set, a four-fold cross
validation routine was employed for final feature selection and optimization of model hyperparameters
within the training data. Classification probabilities from the resulting XGBoost model were then
calibrated with Platt scaling (*115, 116*) using an additional cross-validation procedure within the training
data.

The second layer employed stacked generalization (*117*), a meta-model that classified the brain region
of a tetrode by combining the single-neuron probability outputs from the first classification layer across
all simultaneously recorded neurons on that tetrode. To ensure the meta-model was invariant to the
presentation order of neurons, we first transformed single-neuron probabilities from the first layer into a
permutation-invariant feature representation using symmetric aggregations. e.g., sum, sum-of-cosines
(*118*). These permutation-invariant representations served as input to a logistic regression meta-model.
We trained this meta-model with a synthetic dataset generated by random subsampling of the positive
and negative classes in the training data for each pipeline (i.e., the pipeline for 29e vs. rest or PR vs. rest
classification). Hyperparameter and Platt scaling-based probability calibration for the meta model were
optimized using cross-validation on this synthetic training dataset.

The final layer applied independent confidence thresholds to the output probabilities of the meta-model
in each pipeline for both positive ("29e" or "PR") and negative ("not 29e" or "not PR") class
assignments—a technique known as classification with a reject option (*119, 120*). Region assignments
were accepted only when the output probability for either class exceeded its respective threshold;
otherwise, no assignments were made to ensure that only high-confidence assignments were accepted.
Thresholds for each pipeline were optimized via cross-validation on the training data: after meta-model
fitting on each fold's training data, a range of threshold values was scanned on validation data to
quantify both coverage (the fraction of accepted assignments) and accuracy among accepted
assignments. The resulting trade-off between coverage and accuracy was visualized as a Pareto front
(*121, 122*) from which threshold values were manually selected to provide an optimal balance between
these metrics for both positive and negative class assignments.

Pipeline verification

To assess pipeline performance on unseen data, we performed tetrode-stratified, five-fold cross-
validation (CV) on the histological ground-truth dataset. We refer to this as external CV to distinguish it
from the internal CV procedures used by the pipeline for tuning its parameters within the training data,
as described above. In each fold of the external, tetrode-stratified CV, the dataset was split into training
and test sets in a "tetrode-aware" fashion, i.e., by leaving out all data from selected tetrodes. This split
ensured that neurons recorded on the same tetrode did not appear in both sets, preventing potential data
leakage due to the correlated nature of single units recorded on the same tetrode. All tuning and
calibration procedures of the pipeline (e.g., feature selection, hyperparameter optimization, probability
calibration, and reject-option threshold selection) described above were carried out using internal CV
restricted to the training set within each fold of the external CV.

We then evaluated the resulting pipeline on the test set of the external CV, comprising neurons from
held-out tetrodes. To quantify performance as a function of the number of neurons available for tetrode-
level regional classification, we generated synthetic test samples by pooling neurons across held-out
tetrodes within each class (positive or negative) and then randomly subsampling N neurons (N=1, 2,

…,7) from the appropriate pool. This approach provided sufficient test cases at each N, allowing us to
quantify each pipeline's sensitivity and specificity for identifying the respective positive class ("area
29e" or "PR") as a function of N (fig. S2D). The sensitivity and specificity of both 29e and PR models
increased as N increased.

Pipeline application

After verifying pipeline performance via tetrode-stratified cross-validation (fig. S2D), we trained the
final versions of each pipeline (29e vs. rest and PR vs. rest) on the complete histologically ground-truth
dataset. For these final models, all tuning procedures—feature selection, hyperparameter optimization,
probability calibration, and reject-option threshold selection—were performed via internal cross-
validation on this full dataset, following the same methodology described under *pipeline architecture*.
Note that performance metrics reported in fig. S2D were obtained exclusively from the tetrode-stratified
verification CV, not from this final refit. The finalized pipelines were then applied to classify the
recording regions of tetrodes with ambiguous or infeasible histological localization.

Since classification performance (sensitivity and specificity) improved as the number of simultaneously
recorded neurons increased (fig. S2D), we set minimum thresholds for the number of simultaneously
recorded cells per tetrode based to ensure that the pipeline was applied only under conditions of high
reliability (both sensitivity and specificity ≥ 0.85). Specifically, we applied the 29e vs. rest pipeline to
ambiguous tetrodes with at least 5 neurons, and the PR vs. rest pipeline to ambiguous tetrodes with at
least 7 neurons. Assignments were accepted only when the model's output probability exceeded the
confidence threshold for either the positive or negative class; otherwise, no assignment was made. All
ML-based region assignments for ambiguous tetrodes were finalized prior to downstream data analyses,
ensuring that classification decisions remained blinded to experimental outcomes.

Inclusion criteria

During preprocessing, we enforced a common set of inclusion criteria across analyses. Unless otherwise
specified, spikes were analyzed only when the animal's running speed exceeded 4 cm/s and landmarks
were visible. Units were included if they produced ≥ 100 spikes within the relevant analysis epochs.
Only clusters that could be unequivocally assigned to MEC, PA, PO, PR, or 29e—by histological
examination, or, when histology was ambiguous, by our physiological-feature classifier—were retained
(see "*Brain region classification from single-cell physiological features*").

For tetrode-level landmark-OFF response analyses (Fig. 3A, B), tetrodes were included only if the
population mean firing rate was ≥ 2.5 Hz during the analyzed period. For cross-regional LFP interaction
analyses, a tetrode was included only if at least one neuron with spatially correlated activity was
recorded during the period with stationary landmarks in the Dome VR. For monosynaptic-connection
analyses, putative monosynaptic pairs were considered only if both presynaptic and postsynaptic units
had average firing rates ≥ 0.2 Hz.

For open-arena analyses were restricted to sessions that were ≥ 20 min. To ensure sufficient coverage
during running, we required that in every square bin of a coarse 4 × 4 grid tiling the arena, the animal
had spent at least 75 seconds over the full session and at least 30 seconds within each half of the session.

For cross-regional LFP–LFP and spike–LFP coupling analyses (see below), we used tetrodes that had at
least one neuron with spatially correlated activity, detected based on the initial epoch (G = 1) of the
Dome VR experiments with stationary landmarks.

Neuron matching between Dome and open arena recording sessions

Putative neurons (clusters) were matched across Dome and open arena recording sessions performed on
the same day based on the similarity of their spike waveform features, as in (*123*). Peak waveform
voltage features from the tetrode channels were extracted for each cluster and modeled as multivariate
Gaussian distributions. The 'cluster distance' between each pair of clusters across sessions was
quantified as the minimum of the two asymmetric Kullback-Leibler divergence values between their
Gaussian distributions. For each cluster in a session (i.e., Dome or open arena), we selected the cluster in
the other session with the minimum cluster distance as its putative match. To ensure confidence in this
match, we checked whether any other candidate cluster had a distance falling within 0.5 of this
minimum distance: if so, we treated the match as ambiguous and discarded it, leaving the cluster without
a match in the other recording session. For each cluster that retained a putative match after this
confidence check, we additionally compared the putative match's cluster distance against a threshold to
reject poor matches. To optimize this threshold, we followed a procedure which divided sessions in half
to create ground-truth same-unit pairs and then searched for a threshold that balanced matching accuracy
with rejection rate. This procedure identified a threshold of 8, which we applied to reject all Dome–
open-arena cluster pairs with distances exceeding this value. Finally, we only kept matches that were
bidirectional, i.e. each cluster had to choose the other as its unambiguous match.

Statistics

Statistical analyses were performed in MATLAB (MathWorks, Inc.). All tests were two-sided and
evaluated at a familywise $\alpha = 0.05$ unless otherwise indicated. Whenever multiple comparisons were
performed, *p*-values were adjusted using the Holm–Bonferroni step-down method to control the
familywise error rate.

Nonparametric tests (Kruskal-Wallis test if number of groups > 2, Wilcoxon rank-sum test otherwise)
were used for group comparisons unless the data satisfied the following parametric assumptions.
Normality was assessed with the Shapiro–Wilk test (*124*), except for groups with sample kurtosis > 3,
where we used the Shapiro–Francia test, which has greater power to detect non-normal distributions
with more extreme values or heavy tails (*125–127*). Homogeneity of variances across groups was
examined with both Levene's test (absolute-deviation variant) and Bartlett's χ^2 test (MATLAB's
vartestn with options 'LeveneAbsolute' and 'Bartlett'). If every group passed the normality test and
either one of the tests for equal variances were non-significant ($p > 0.05$), we treated the data as suitable
for parametric analysis. We ran a one-way ANOVA to test for overall group differences, checked that
the ANOVA residuals were still normal, and then compared groups pairwise with standard two-sample
t-tests. For two group unpaired comparisons, we used the nonparametric Wilcoxon rank-sum test unless
the data met parametric assumptions. If both groups were normally distributed, we tested equality of
variances with the two-sample F-test; when variances were equal we used the standard two-sample t-
test, and when variances differed we used the Welch's t-test. For paired data, we analyzed the within-
pair differences: if these difference scores were normally distributed, we applied a paired t-test;
otherwise, we used the Wilcoxon signed-rank test.

To account for non-independence among LFP measurements from the same animal (and, where
applicable, repeated measurements from the same tetrode pair, neuron, or brain region), we employed
linear mixed-effects models. Following established practices for inference in linear mixed-effects
models (*128, 129*), models were fit by restricted maximum likelihood (MATLAB's *fitlme*) and statistical
significance of fixed effects was evaluated using Satterthwaite-approximated denominator degrees of
freedom (MATLAB's *anova* and *coefTest* functions on fitted mixed-effects models). This approach
yields accurate *p*-values, when the number of animal is relatively small and/or when the effective sample
size is closer to the number of animals than to the total number of observations, because repeated
measurements from the same animal are correlated. In contrast, alternative approaches that assume

large-sample asymptotics (e.g., Wald/z-tests) or use residual degrees of freedom (e.g., classical
regression/t-tests) can yield inaccurate, optimistically low p-values by overstating the amount of
independent evidence under these conditions.

Using Satterthwaite-approximated degrees of freedom, we evaluated statistical significance of fixed
effects as follows: Fixed effects spanning more than two conditions (e.g., differences among more than
two brain regions) or interaction terms were assessed via F-tests on the fitted model (MATLAB's
anova). All other tests on fixed effects, including single coefficients vs. zero (e.g., a condition mean or
the slope relating two variables) and post-hoc pairwise comparisons, were evaluated via t-tests on linear
contrasts of the fixed-effects coefficients from the fitted model (MATLAB's *coefTest*). Model-specific
details are provided in the relevant sections below.

Power spectral analysis of LFP

1/f background correction

The wavelet transform was applied to each tetrode's LFP signal using Morlet wavelets spanning 1–140
Hz (Buzsáki Lab, GitHub: https://github.com/buzsakilab/buzcode). Instantaneous power was computed
from the wavelet coefficients and averaged across the session. To isolate oscillatory power from the
aperiodic background in each LFP power spectrum, we used the FOOOF algorithm (*130*) with the
following parameters: knee mode, peak threshold = 2.5, peak width = 2–45 Hz, and max_n_peaks = 4.
Starting from the original spectrum, FOOOF was run iteratively as suggested in (*131*). In each iteration,
FOOOF took the spectrum from the previous iteration as an input and fitted a parametric model for the
oscillatory peaks and the aperiodic (1/f) background. Subtracting the identified peaks from this spectrum
resulted in a refined spectrum, which was then used as input for the next iteration. This process
continued until no oscillatory peaks greater than 1% above the aperiodic background remained, or until a
maximum of ten iterations. The final peak-removed spectrum was then fit with a simple log–linear
model to robustly estimate the aperiodic background. This fitted background was subtracted from the
log-transformed original spectrum to obtain the 1/f-corrected (flattened) log power spectrum for each
tetrode. Positive values in the 1/f-corrected log power spectrum indicate frequencies where the power
exceeded the modeled aperiodic background, suggesting presence of oscillatory components, while
values near or below zero indicate frequencies without a detectable oscillatory excess.

Cluster-based permutation testing of significant oscillatory power across frequencies

We assessed the statistical significance of oscillatory components in the LFP signal across tetrodes
within each brain region while controlling for familywise error (due to multiple comparisons). A naïve
Bonferroni correction would not account for the continuous (correlated) nature of the frequency
spectrum, so we opted for a cluster-based permutation test (*53, 132–134*). For each permutation, the sign
of the entire 1/f-corrected log power spectrum from each tetrode was randomly flipped for a given
session. At each frequency, a one-sample, one-sided t-test was performed across tetrodes for both the
observed and permuted values. Clusters were defined as contiguous frequencies ranges over which there
were significant positive deviations from zero according to $p < 0.05$ (one-sided t-test). The size of each
cluster was calculated as the sum of t-statistics within the cluster. A null distribution was created from
the maximal cluster size from each of the 1000 permuted data sets. Each cluster in the observed data was
then compared to this null distribution, yielding permutation-based cluster-level *p*-values. Clusters that
satisfied $p < 0.05$ for this permutation test were considered significant.

Mixed effects-model of theta- and gamma-band LFP power

Average powers in theta (5–10 Hz) and gamma (50–120 Hz) bands were calculated from the 1/f-
corrected log-power spectrum of each tetrode to test for regional differences. We used 1/f-corrected

power spectra rather than raw power due to potential differences in aperiodic background activity across
days and/or animals, which were not perfectly balanced across recorded regions. To also account for
non-independence of LFP signals within animals when comparing band-limited power across regions,
we used linear mixed-effects modeling.

For each frequency band (theta and gamma), data were arranged as one observation per tetrode location
with the following variables: band-limited 1/f-corrected power, region identity (categorical variable with
five values: 29e, MEC, PA, PO, and PR), and rat identity (categorical variable). Region identity was
included as a fixed effect to test for systematic differences in power across regions. The model included
two random-intercept terms: a rat-specific random intercept (1|rat) to capture correlation among LFP
observations from the same animal, and a rat-by-region specific random intercept (1|rat:region) to
capture additional correlation among LFP observations from the same region within the same animal
(i.e., allowing each rat to have a region-specific baseline). For each frequency band, we then fit the
following mixed-effects model for each band (theta or gamma):

$$\text{power} \sim \text{region} + (1|\text{rat}) + (1|\text{rat:region}).$$

Each power band model was fit by restricted maximum likelihood (using MATLAB's *fitlme*) and the
statistical significance of fixed effects (i.e., both omnibus test of region and post-hoc pairwise contrasts
between regions) was evaluated using Satterthwaite-approximated denominator degrees of freedom.
Omnibus effect of region was assessed with F-test using MATLAB's *anova* function on the fitted
mixed-effect models. When this omnibus effect was significant, we performed post-hoc pairwise
comparisons between each region and 29e using t-tests on linear contrasts of fixed effects' coefficients
from the fitted model (implemented via MATLAB's *coefTest*). *P*-values across the four region-versus-
29e contrasts for each model were adjusted for multiple comparisons using the Holm–Bonferroni
correction.

Theta and gamma rhythmicity of single neurons

1644 Rhythmicity was assessed from power spectrum of each neuron's spike autocorrelogram as follows: the
1645 theta index was calculated as the mean power in 5–10 Hz divided by the mean power in 1–50 Hz (*50*),
1646 and the gamma index as the mean power in 50–120 Hz divided by the mean power in 5–250 Hz (*135*).

Coupling of single neurons to LFP theta rhythm

Each tetrode's LFP signal was band-pass filtered between 5–10 Hz (4[th] order Butterworth filter). The
Hilbert transform was estimated using MATLAB's *hilbert* command, which returns a complex signal
called the analytic signal. Instantaneous theta-band power and phase were extracted from the magnitude
and angle of the analytic signal, respectively. For each neuron, the spike-phase distribution was obtained
by circularly interpolating the instantaneous theta phase from each tetrode's LFP signal at the neuron's
spike times. Pairwise phase consistency (PPC) was then computed for each neuron's spike-phase
distribution relative to each tetrode using FieldTrip (*136*). The highest PPC across tetrodes was used to
quantify each neuron's coupling to the LFP theta rhythm.

Spatially correlated activity in the Dome

A circular firing-rate map was created for each neuron to quantify its spatial tuning in the Dome when
landmarks were stationary (i.e., experimental gain, G = 1). We assigned both the angular position of
each spike and the animal's position at each behavioral time point to 36 angular bins (10° per bin)
spanning all locations across the circular track. We then computed the firing rate in each bin by dividing
the number of spikes by the amount of time spent in that bin. The resulting rate maps were smoothed
with a circular Gaussian kernel (σ = 15°). The smoothed circular rate map was then used to calculate
each neuron's Skaggs' spatial information (SI) score (in bits per spike; *38*):

$$\text{SI} = \sum_i p_i \frac{r_i}{r} \log_2 \frac{r_i}{r},$$

1664

where p_i is the fraction of total session time the animal spent in i^{th} angular bin, r_i is the neuron's firing
rate in the same bin, and r is the neuron's overall mean firing rate; the summation is over the bins, $i =$
$1, \dots, 36$. Because this score (originally developed for 2D firing-rate maps) can produce both false-
positive and false-negative errors when applied to data from 1D circular tracks (*137*), we also quantified
spatial tuning using two alternative SI metrics, the Olypher SI score and the mutual SI score, defined
below.

The Olypher SI score is defined as the maximum Kullback-Leibler divergence between the neuron's
spike count distribution at any given position and its overall spike count distribution (*138*):

1673
$$\text{Olypher SI} = \max_i \sum_{k \geq 0} P(k|i) \log_2 \frac{P(k|i)}{P(k)},$$

1674 where $P(k|i)$ is the probability of observing k spikes within a 100 ms time-segment when the animal
1675 occupies positional bin i, and $P(k)$ is the marginal probability of observing k spikes, regardless of the
1676 animal's position.

1677 The mutual SI score is defined as the mutual information between the neuron's spike count and the
1678 animal's position (*139, 140*):

1679
$$\text{Mutual SI} = \sum_{i,k \geq 0} P(k,i) \log_2 \frac{P(k,i)}{P(k)P(i)},$$

where $P(i)$ is the marginal probability of the animal occupying the positional bin i within a 100 ms time
segment (independent of the number of spikes), $P(k)$ is the marginal probability of observing k spikes
within a 100 ms time segment (independent of occupancy), and $P(k,i)$ is the joint probability of
observing k spikes and the animal being in positional bin i within the same 100 ms time segment.

To assess significance of each SI score for each neuron, we generated a null distribution by circularly
shifting the neuron's spike train relative to behavioral data by a randomly chosen time offset (uniform
distribution between 25 and T – 25 seconds, where T is the total duration of the first experimental
epoch) and recalculating the SI scores 100 times. A neuron was considered to have significant spatially
correlated activity if at least one of its observed SI scores exceeded the 99$^{\text{th}}$ percentile of the respective
shuffled SI distribution. We refer to such neurons as "spatially correlated".

Landmark-selective firing modulation in the Dome

Z-index

The circular firing-rate map was linearly interpolated to 1° resolution (0–359°) and normalized to z-
scores across angular bins. For each landmark location, we identified the angular bin corresponding to
the circular track position closest to the landmark. We then computed the mean z-scored firing rate
within a symmetric, 24° angular window centered on that bin (i.e., ±12° around the landmark angle,
wrapping circularly as needed). For comparison, we identified midpoint angles halfway between each
pair of adjacent landmarks and computed the mean z-scored firing rate within an identically defined 24°
window centered on each midpoint. The landmark selectivity z-index was then defined as the absolute
difference between the mean z-scored firing rate averaged across all landmark-centered windows and
that across all midpoint-centered windows.

Local similarity index

Each neuron's circular firing rate map was linearly interpolated with 1° resolution and partitioned into
sectors around each landmark. Each landmark sector was defined as the arc spanning between the
midpoints of that landmark and its two neighboring landmarks. For example, if there were three equally
spaced landmarks, there would be three 120° sectors, one centered on each landmark; however, if the
landmarks were unevenly spaced (as was typical), the sectors would have different widths. For the given
neuron and for each pair of landmarks, the Pearson correlation coefficient was computed between the
peri-landmark firing-rate profiles, using only the angular range relative to the center of the landmarks
shared by both sectors (i.e., their intersection). This shared range varied from as little as ±20° up to ±60°
for different pairs, depending on the number and spatial arrangement of landmarks within the Dome
(Fig. 1B). For two landmarks, there was one such pairwise correlation, for three landmarks there were
three pairwise correlations, and four landmarks there were six pairwise correlations, based on the
formula *N(N-1)/2* where *N* is the number of landmarks. The local similarity index for each neuron was
then defined as the average of all such pairwise correlations across landmark pairs.

Landmark-locking score

For each neuron, we computed a lapwise firing-rate map in the landmark frame by binning spikes and
the animal's behavioral data within each lap into 10° angular bins, dividing the spike count in each bin
by the time spent in that bin, and smoothing with a Gaussian kernel with $\sigma = 15°$. Using the last 12 laps
in each session of this firing-rate map, we calculated the spatial autocorrelation function for spatial lags
between 0 and 9 laps. (Thus, the minimum overlap in the autocorrelation calculation was 3 laps,
corresponding to a lag of 9 laps within the last 12 laps.) Neurons with stable lapwise firing exhibited
autocorrelation functions with prominent oscillations at integer-lap periods, indicating consistent
periodicity across laps (Fig. 1E; fig. S2A), whereas neurons that broke free from landmarks displayed
autocorrelation functions with weak or irregular oscillatory structure (Fig. 1F; fig. S2B).

To quantify the lap-to-lap periodicity of each neuron's autocorrelation function, we first computed all
possible pairwise Pearson correlations among circularly seven shifted versions of the 9-lap
autocorrelation function at full-lap lags (circular shifts of 0, 3, 4, 5, 6, 7, and 8 laps, leading to 21
pairwise Pearson correlation values amongst these autocorrelograms) as well as all possible correlations
between these full-lap shifted versions and the half-lap rotated versions (five circular shifts from 3.5 to
7.5 laps, in 1-lap steps, leading to 35 pairwise Pearson correlation values between full-lap and half-lap
shifted autocorrelograms). Neurons with stable lapwise periodicity are expected to show higher
correlations among the full-lap rotations than between the full-lap and half-lap groups.

Accordingly, the landmark-locking score was defined as a normalized difference measure: the sum of all
Pearson correlations within the full-lap rotation group minus the sum of all correlations between the full-
lap and half-lap groups, divided by the total number of correlations (Fig. 2; fig. S2). A score close to 1
indicates strong, consistent periodicity at the spatial scale of one lap, while a score near zero indicates
little or no periodic firing structure.

Spatial tuning in stationary and mobile segments of open-loop sessions

The decoupling of landmark rotation from animal movement in open-loop experiments created periods
where landmarks rotated while the animal remained stationary. Supplementary analysis (fig. S4)
examined whether neural firing patterns relative to landmarks were preserved during these *stationary*
periods. Accordingly, we defined two conditions from data where the animal's speed relative to
landmarks was > 4 cm/s: *mobile* and *stationary. The mobile* condition corresponded to periods when the
animal's lab-frame speed was also ≥ 4 cm/s (regardless of landmark motion), whereas
the *stationary* condition corresponded to periods when the animal's lab-frame speed was < 4 cm/s
(meaning landmarks moved relative to the stationary animal).

The discontinuous nature of *stationary* segments precluded use of the landmark-locking score—which requires continuous spatial sampling across laps—to quantify the extent of landmark-anchoring during these segments. Instead, we assessed spatial tuning during *stationary* segments by comparing landmark-referenced firing patterns between *stationary* and *mobile* segments. We restricted this analysis to "landmark-anchored" neurons—defined as those with landmark-locking scores > 0.52 (the threshold separating *LM-control* and *LM-failure* sessions in Fig. 1G)—to ensure that anchoring to landmarks during stationary segments were evaluated only for neurons with stably anchored firing fields to the rotating landmarks during *mobile* periods.

For each landmark-anchored neuron, we computed the 1D circular firing rate map as a function of angular position relative to landmarks (10° bins) for both *mobile* and *stationary* periods. Pearson's correlation between the two rate maps quantified the *mobile–stationary* similarity of the rate maps. To determine the statistical significance of the observed similarity for each neuron, we circularly shifted the *stationary* rate map by 2 to 34 bins and computed its Pearson correlation relative to the *mobile* rate map, yielding a shuffled distribution of correlation coefficients. Landmark-anchored neurons with observed correlations exceeding the 95th percentile of its shuffle distribution were considered statistically significant in terms of similarity of its rate maps across *mobile* and *stationary* conditions. Due to limited number of landmark-anchored neurons from MEC, PA, PO, and PR in these experiments, neurons from these regions were pooled into a single "non-29e" group for comparison with area 29e.

Unsupervised identification of LM-control and LM-failure session types

To identify session-wise differences in landmark control over neuronal populations, we first computed the average landmark-locking score across all recorded neurons from all regions in each session. Sessions were then clustered based on their respective average landmark-locking score using an enhanced density-based clustering algorithm (MATLAB's *dbscan*) procedure that set MinPts = 4 and automatically tuned the neighborhood radius ε through k-distance heuristic with k = MinPts (*141*). Briefly, the absolute difference between the average landmark-locking score of each session and that of all remaining sessions was calculated. We then measured, for each session, the distance to its k^{th}-nearest neighbor session. Measurement of this distance for each session led to the so-called k-distance graph. The neighborhood radius ε was set as the distance corresponding to the knee point of this graph. DBSCAN was run with this ε and the corresponding MinPts. Any session labeled as noise (i.e., left out of clustering) was assigned to the cluster whose centroid was nearest. This unsupervised approach identified two clusters which we defined as LM-control and LM-failure. See Fig. 1G.

Landmark removal score

Multi-unit, unclustered spike trains from each tetrode were binned at 10 ms resolution. To quantify the neural response to landmark removal, we aligned spike counts to the time of landmark extinction and extracted a window from –1 to +1 seconds relative to this event for each session.

The pre-removal baseline was defined as the period from –0.6 to 0 seconds before landmark removal, and the post-removal response period as 0 to +0.4 seconds after removal (the larger pre-removal time period was used to get a better estimate of the intrinsic variability before removal; the result does not change when symmetric 400 ms pre-and post-removal time periods were used). For each time bin, we calculated the absolute deviation from baseline as $d(i) = |$ spike count in bin i – mean baseline spike count $|$. The landmark removal response score was then computed as:

$$(\text{mean}(|d_{\text{post}}|) - \text{mean}(|d_{\text{pre}}|)) / (\text{mean}(|d_{\text{post}}|) + \text{mean}(|d_{\text{pre}}|)),$$

where the means are taken over all 10ms time bins in the pre- or post-removal period; thus, mean($|d_{\text{pre}}|$)
and mean($|d_{\text{post}}|$) represent the mean absolute deviations from baseline during pre- and post-removal
periods, respectively.

This absolute deviation approach treats increases, decreases, and even biphasic responses (i.e., a
combination of increase and decrease) equivalently. The normalized score ranges from –1 to +1, with
positive values indicating that post-removal firing deviates more from baseline than expected from pre-
removal fluctuations (suggesting a response to landmark removal), while values near zero indicate
deviations comparable to baseline fluctuations.

Single neuron responses to landmark and background luminance manipulations

Lapwise firing rate maps for each neuron were computed as before (see Landmark-locking score
section). Landmark and background luminance were each set to one of two levels (high/low), yielding
four landmark–background luminance combinations (2 × 2 factorial design).

Luminance sensitivity analysis

We fitted a two-way ANOVA to each neuron's lapwise peak firing rates, using lapwise landmark
luminance (LM) and background luminance (BG) as factors. The effect size for each factor X $\in$ {LM,
BG} was quantified by

1805
$$\eta_X^2 = \frac{SS_X}{SS_{Total}},$$

where SS_X denotes the sum of the squares in peak firing rate explained by the variation in the factor X
and SS_{Total} denotes the total sum of squares in the peak firing rates. Because these luminance factors
were independently manipulated and uncorrelated, we quantified their combined overall effect size as
the sum $\eta_{overall}^2 = \eta_{LM}^2 + \eta_{BG}^2$.

Contrast analysis

Although landmark and background luminance were independently manipulated by a two-factorial
design, the resulting contrast (C=LM – BG) and total luminance (TL=LM + BG) values were linearly
dependent and did not obey a fully factorial design, precluding the use of ANOVA to analyze firing-rate
dependence on (C, TL). Accordingly, a linear model was fit with C and TL as predictors and each
neuron's peak firing rates as the dependent variable. The unique contribution of each predictor X $\in$ {C,
TL} was quantified using partial R²:

1817
$$R_{\text{partial}}^2(X) = \frac{(\text{SSE}_{\text{reduced}} - \text{SSE}_{\text{full}})}{\text{SSE}_{\text{reduced}}} = \frac{(R_{\text{full}}^2 - R_{\text{reduced}}^2)}{\left(1 - R_{\text{reduced}}^2\right)}$$

Here, SSE is the sum of squared errors, "full" refers to the model with both predictors, and "reduced"
refers to the model with X removed. Partial R² thus quantifies the additional variance in firing rate
explained by a predictor, after accounting for the presence of the other, even when the predictors are
correlated.

Allocentric place and head direction tuning in the open arena

For robust characterization of allocentric place and HD tuning scores despite differences in behavioral
sampling of place and HD, we used a Poisson generalized linear model (GLM), as in previous work (*44*,
142). Briefly, each neuron's spike train was binned into dt = 125 ms segments. For each segment t_i, the
expected spike count was modeled as

1827
$$\lambda(t_i) = dt \cdot \exp\left[X_p(t_i)\, w_p + X_h(t_i) w_h\right],$$

where X_p and X_h are one-hot encoded row vectors representing, respectively, the animal's (x,y) position
(in 7×7 cm^2 bins) and HD (in 8° angular bins) at the center of that segment. The column vectors w_p and
w_h denote the predictive weights for position and HD (analogous to tuning curves). Fitting of these
weights was performed by maximizing an objective function that combined the Poisson log-likelihood
of the observed spike counts with a smoothing penalty to discourage large weight differences between
adjacent bins, as previously described (44, 142). Optimization was performed using the MATLAB's
fminunc search routine , aided by analytically computed gradients. Using the optimized weights $\hat{w}_p$ and
$\hat{w}_h$ for each neuron, we constructed a 2D spatial firing rate map and a 1D HD tuning curve, each
marginalized over the other behavioral variable:

1837
$$r_p(X_p) = \exp[X_p \, \hat{w}_p] \frac{1}{N_h} \sum_{X_h} \exp[X_h \, \hat{w}_h]$$

1838
$$r_h(X_h) = \exp[X_h \, \hat{w}_h] \frac{1}{N_p} \sum_{X_p} \exp[X_p \, \hat{w}_p],$$

Here, $N_h = 72$ (the number of 5° HD bins) and $N_p = 35 \times 35$ (the number of 4×4 cm^2 positional bins).
These marginal rate maps were then used to calculate the Skaggs' SI score in bits per spike as a measure
of each neuron's place-tuning strength and Rayleigh score (i.e., mean resultant length) as a measure of
its allocentric-HD tuning strength.

Testing significance and stability of allocentric place tuning

We circularly shifted each neuron's spike train by a randomly chosen time offset (between 25 and T –
25 seconds, where T is the total session duration) 100 times, refitting a GLM to each shuffled spike train
to generate a null distribution for both SI and Rayleigh score.

For tuning stability of each neuron, we first computed the conventional (i.e., without using a GLM) 2D
place rate maps and HD polar tuning curves from the first and second halves of the session, separately.
The split-session stability of place tuning was then quantified as the Pearson correlation between the
split-session rate maps, while the split-session stability of HD tuning was quantified as the difference in
mean angular direction between the first and second halves of the session.

A neuron was considered significantly tuned to place if its GLM-derived SI exceeded the 99[th] percentile
relative to the shuffled distribution of GLM-derived SI scores and if its split-session stability exceeded
0.5. A neuron was considered significantly tuned to allocentric HD if its GLM-derived allocentric HD
tuning curve passed the Rayleigh test of uniformity at $\alpha = 0.01$, if its GLM-derived Rayleigh score
exceeded the 99[th] percentile of the respective GLM-based shuffled distribution, and if its split-session
mean directional difference was less than 45° as previously used (45).

Egocentric head-direction tuning in the open arena

Because egocentric coding can in principle be anchored to any arbitrary point in an environment that is
unknown to the investigator, quantifying a neuron's egocentric tuning curve typically involves an
exhaustive search to determine its optimal anchor point. In contrast, allocentric coding is characterized
by a single possible tuning curve per neuron, requiring no anchor-point search. If one compares the
single allocentric tuning curve to the most specifically tuned egocentric tuning curve derived from the
anchor-point search, this asymmetry creates a bias in favor of the egocentric model over the allocentric
model when evaluating a neuron's preferred frame of reference (43). Model selection metrics criteria
(e.g., AIC/BIC) partly compensate for this bias but rely on assumptions that poorly capture the
complexity of the anchor search and can still reward overfitting. More rigorous controls (e.g., surrogate

tests) would help but are computationally prohibitive because each surrogate demands a new exhaustive
anchor search.

To circumvent these issues, we developed an *egocentric vector-field* (EVF) score that quantifies
egocentric tuning without requiring specification of an anchor point. The EVF score is computed as
follows: We constructed a 2D HD vector field for each neuron by dividing the arena into a uniform 9×9
grid of (x, y) locations with each spatial bin being 15×15 cm². We identified all time points when the
animal's position fell within the bin and assigned the corresponding HDs to 12 angular bins (30° each).
If any angular bin at a grid location had less than 0.2 seconds of occupancy, the spatial bin size at that
location was increased iteratively until every HD bin reached this sampling threshold.

For the local HD tuning curve constructed at each (x, y) grid location, we computed its complex-valued
first harmonic vector,

1879
$$v_{hd} = \sum_{k=1}^{12} \mathrm{FR}_k e^{i\theta_k},$$

where FR_k is the firing rate in HD bin k associated with the angle θ_k. Note that the angle of this vector is
the same as the mean angular direction of the local HD tuning curve, while its magnitude is equal to the
product of the summed firing rate within the local HD tuning curve and the MRL of the local HD-tuning
curve. This vector is equal to the first (i.e., fundamental) Fourier component of the local HD-tuning
curve, which defines the single sinusoid that best fits the tuning curve in the least-squares sense. The
collection of these vectors across all grid locations comprised the 2D HD vector field for the neuron.

Egocentric vector field (EVF) score

To quantify egocentric tuning without specifying an anchor point, we analyzed the spatial derivatives of
each neuron's local HD vectors (i.e., how these vectors vary along the x and y axes of the arena).
Specifically, we computed two classical vector calculus measures, *curl* and *divergence,* for each spatial
bin (x, y) of the open arena (fig. S8).

The *Curl* of a vector field captures local rotational (swirling) structure of the HD vectors:

1892
$$\mathrm{curl}(x, y) \triangleq \frac{\partial Q(x, y)}{\partial x} - \frac{\partial P(x, y)}{\partial y}$$

1893 The *Divergence* captures the local inward/outward flow structure of the HD vectors:

1902
$$\mathrm{div}(x, y) \triangleq \frac{\partial P(x, y)}{\partial x} + \frac{\partial Q(x, y)}{\partial y}$$

1903
where, $P(x, y)$ and $Q(x, y)$, are the *x*- and *y*-components of the two-dimensional HD vector-field
(corresponding to real and imaginary components of v_{hd} at each x,y location). These partial derivatives
were estimated using centered-difference approximations between bin centers. Neurons lacking any
systematic HD tuning and neurons with allocentric HD tuning (characterized by parallel vectors
throughout the arena) exhibited near-zero curl and divergence everywhere (these neurons can be
distinguished from each other by measuring the sharpness and significance of their allocentric
directional tuning curves). By contrast, neurons with strong egocentric tuning generated spatially
consistent, high curl and/or divergence values (fig. S8).

We combined the curl and divergence values at each spatial bin into a 2D vector. The curl was assigned
as the horizontal component of this vector, whereas the divergence was assigned as its vertical
component. This approach merged the separate curl and divergence fields into a new vector field that
captures both rotational and flow characteristics of the preferred HD throughout the arena.

We then calculated the Egocentric Vector Field (EVF) score as the MRL of all divergence-curl vectors across the arena. This score ranges from 0 to 1, where values close to 0 indicate random or inconsistent curl/divergence patterns, whereas values close to 1 indicate highly consistent patterns characteristic of strong egocentric tuning.

Generalized linear models for egocentric versus allocentric preference

To dissociate each HD-tuned neuron's preferred frame of reference (egocentric vs. allocentric) and to assess split-session stability of egocentric tuning, we used a Poisson GLM framework (*43–45*). Specifically, for each neuron, we fitted two separate GLMs to its spike train, binned into 125 ms time segments: one GLM used allocentric head direction (HD) as a predictor, while the other used egocentric HD, computed relative to the (x, y) coordinates of a candidate anchor point. For the egocentric model, the optimal anchor point was identified by grid search: the GLM was re-fit at each candidate anchor location spaced every 5 cm across the arena in both x and y directions, and the location yielding the largest log-likelihood was selected.

This GLM-based anchor optimization served two purposes: (1) comparing predictive performance of allocentric versus egocentric models, and (2) identifying an anchor point to assess split-session stability of egocentric tuning (see below) and to compare arena-to-anchor point distance across regions (see Fig. 3G). Note that statistical significance of egocentric tuning was assessed separately using the anchor-free EVF score (below), which avoids the statistical biases inherent in anchor optimization as described above.

To compare the predictive performance of the two models while accounting for the increased complexity of the egocentric GLM ($K_{ego} = 2$ additional parameters corresponding to the anchor–point optimization), we calculated the difference in Bayesian information criterion

$$\Delta\text{BIC} = 2\ln\hat{L}_{allo} - 2\ln\hat{L}_{ego} + K_{ego}\ln N,$$

where $\hat{L}_{ego}$, $\hat{L}_{allo}$ are the optimal log-likelihoods of GLMs with egocentric HD and allocentric HD variables, respectively, and N is the number of bins (time segments). A positive ΔBIC suggests that the allocentric model provides a better fit, whereas a negative ΔBIC suggests a better fit for the egocentric model.

Testing significance and stability

To assess statistical significance of egocentric tuning, we generated two null distributions for the anchor-free EVF score of each neuron. Similar to other spatial tuning measures, one null distribution was obtained by circularly shifting the neuron's spike train relative to the animal's behavioral data by a single, randomly chosen time offset (between 25 seconds and T−25 seconds, where T is the total session duration), and recomputing the EVF score 100 times. The second null distribution was generated by disrupting the relationship between position and head-direction vectors in the HD vector field 1,000 times: for each iteration, we circularly translated the entire HD vector field along both the x and y axes by random amounts and then recomputed the EVF score.

To assess split-session stability of egocentric tuning, we constructed conventional (i.e., without using a GLM) egocentric-HD polar tuning curves from the first and second halves of the session. Both tuning curves were referenced to the optimal anchor point identified by the GLM grid search on the full session data (described above). Stability was quantified as the absolute angular difference between the preferred directions of the tuning curves from the two halves.

1949   A neuron was considered significantly tuned to egocentric HD if its observed EVF score exceeded the
1950   99[th] percentile of both null distributions and the split-session mean directional difference was less than
1951   45°.

Classification of spatial and directional tuning of cells in open field

1953   All neurons were evaluated for allocentric HD, egocentric HD, and allocentric place tuning as described
1954   above. Neurons meeting criteria for only egocentric-HD or only allocentric-HD tuning were classified
1955   accordingly. For neurons meeting criteria for both, we compared their egocentric and allocentric GLMs
1956   (i.e., the egocentric one corresponding to the optimal anchor point and the allocentric one) and classified
1957   each neuron according to the GLM that yielded the lower Bayesian Information Criterion score (BIC).
1958   Results were unchanged when Akaike Information Criterion was used instead. Neurons failing to meet
1959   either HD tuning criterion were classified as nondirectional and screened for allocentric place tuning.

LFP–LFP coherence

1961   LFP signals were segmented into 1-second candidate windows with 95% overlap (0.05 s step size).
1962   These windows were evaluated sequentially from the start of the recording. Any window in which the
1963   animal's velocity was below 4 cm/s for more than 25% of the time, or where the maximum velocity did
1964   not reach 10 cm/s, was discarded. If a window satisfied both speed criteria, it was retained for analysis.
1965   Other candidate windows that overlapped in time with this accepted window were excluded from further
1966   consideration. This process was repeated iteratively until no windows remained, resulting in a set of
1967   non-overlapping, speed filtered 1-second windows for the subsequent spectral analysis.

1968   Within each retained window, LFP signals were determined. Spectral analysis was performed using the
1969   multi-taper Fourier transform with Slepian (DPSS) tapers via FieldTrip (*136*) for frequencies between 0
1970   and 140 Hz with a frequency resolution of 1 Hz (as determined by the 1-second window size). For
1971   frequencies below 25 Hz, three tapers were used (yielding a frequency smoothing half-width of 2 Hz);
1972   for frequencies above 25 Hz, six tapers were used (smoothing half-width 3.5 Hz). For each window and
1973   frequency, the complex-valued cross-spectral density (CSD) matrix was computed from all
1974   simultaneously recorded tetrode pairs.

Baseline coherence levels

1976   For each tetrode pair, we estimated the complex coherency (*C*) based on averaged bivariate CSD
1977   matrices across all retained, non-overlapping windows within the initial stationary-landmark epoch (G =
1978   1). The magnitude of the resulting complex coherency estimate was biased upward due to a limited
1979   number of windows and has a variable variance that again depends on the number of windows. Under
1980   the null hypothesis of zero coherence, both bias and variance have analytically known forms (*143*, *144*)
1981   which can be used for bias correction and variance normalization as follows:

1982
$$coh = \frac{C}{|C|} \sqrt{\frac{|C|^2 - \frac{1}{\nu - 2}}{\frac{1}{\nu - 2}}}.$$

1983   Here, ν denotes the degrees of freedom (equal to twice the product of the number of tapers and the
1984   number of windows), *C* is the raw coherency estimate, and *coh* is its bias-corrected and variance
1985   normalized version. To capture true (i.e., time-lagged, non-spurious) coupling between LFP signals
1986   while remaining insensitive to instantaneous shared signals arising from volume conduction or common
1987   reference effects, we focused on the imaginary part of this corrected coherency (i.e., *coh*), following
1988   previous work (*145–149*). Specifically, we took the absolute value of the imaginary component,

1989   normalized it by $\sqrt{1 - \mathrm{Re(coh)}^2}$—the theoretically maximum possible value of the imaginary
1990   component given the observed real component—and applied a Fisher z-transform, yielding what we
1991   refer to as the "z-imaginary coherence spectrum" (zCoh):

$$1992 \qquad \mathrm{zCoh} = \tanh^{-1}\left(\frac{|\mathrm{Im(coh)}|}{\sqrt{1 - \mathrm{Re(coh)}^2}}\right).$$

The resulting z-imaginary coherence spectrum at the single tetrode-pair level was then averaged across
all 29e-MEC tetrode pairs, yielding a region pair-level average z-imaginary coherence spectrum.

To assess significance of the average z-imaginary coherence spectra across all 29e-MEC tetrode pairs,
we used a hierarchical, cluster-based permutation test (*53*, *150*). For each tetrode pair, a null distribution
of CSD values was generated by randomly flipping the sign of only the imaginary component of the
CSDs from retained windows (which leaves the real part and magnitude unchanged), then recalculating
the z-imaginary coherence spectrum after averaging the (imaginary) sign-flipped CSDs. This
permutation was repeated 1,000 times per tetrode pair. To generate a group-level (region-pair) null, we
randomly sampled one null spectrum from each of the above-mentioned individual tetrode-pair null
distributions 5,000 times and, for each random sample, calculated the average z-imaginary coherence
spectrum. The observed region pair-level z-imaginary coherence at a frequency was considered
significant if it exceeded the 95[th] percentile of the permutation group-level null distribution. Clusters
were defined as contiguous frequencies that are significant with no gap. To control the familywise error
rate of the comparisons across the whole frequency spectrum while respecting spectral contiguity, the
cluster size (sum of coherence values within the cluster) was compared to the distribution of maximal
cluster sizes from the group-level permutations. Clusters with corrected $p < 0.05$ were considered
significant.

Coherence change from initial to final epochs

For each tetrode pair, we computed the z-imaginary coherence spectrum in the initial (G = 1) and final
(G = G_{final}) experimental epochs after averaging their bivariate CSD matrices across accepted 1-second,
non-overlapping windows from that epoch. For each frequency band—theta (5–10 Hz) and gamma (50–
120 Hz)— and for each epoch (initial and final), we took the maximum z-imaginary coherence across
frequency bins within the band and used this value as the band-specific coherence for that epoch. The
normalized change in coherence for each band was then calculated as ΔCoherence = (final –
initial)/(final + initial).

These within-session changes from the initial to final epoch were analyzed using linear mixed-effects
models to account for repeated nature of theta- and gamma-band measurements from the same 29e–
MEC tetrode pair and for the non-independence of measurements among multiple tetrode pairs recorded
from the same animal.

Fixed effects included Session Type (*LM-control* vs. *LM-failure*), Band (theta vs. gamma), and their
interaction (Session Type × Band). Because each 29e-MEC tetrode pair contributed a pair of
observations (one for theta, one for gamma), and multiple tetrode pairs were recorded from the same rat,
observations were not independent. We modeled this nested dependence structure using three random
effects:

(i)   a rat-specific random intercept (1 | RatID), allowing variability of ΔCoherence across rats to
capture correlation among observations from the same animal (across all tetrode pairs and both
bands within each rat).

(ii) a rat-specific random band effect (Band | RatID), allowing variability of theta-gamma difference
2031   across rats to capture covariance between theta and gamma-band observations from the same
2032   animal (across different tetrode pairs within each rat).

2033   (iii) a rat-specific random intercept for tetrode pairs (1 | RatID:TetrodePairID), allowing repeated
2034   measurements from each tetrode pair (within a rat) to have its own baseline level of ΔCoherence,
2035   thereby capturing the correlation between theta and gamma measurements from the same tetrode
2036   pair.

The model was formulated as:

ΔCoherence ~ Session Type × Band + (1 | RatID) + (Band | RatID) + (1 | RatID:TetrodePairID).

We fit the model by restricted maximum likelihood (using Matlab's *fitlme* function) and evaluated
statistical significance of fixed effects—including both omnibus tests on main and interaction effects as
well as post-hoc comparisons—using Satterthwaite-approximated denominator degrees of freedom.
Omnibus effects were assessed with F-tests using MATLAB's *anova* function based on the fitted model.
When the Session Type × Band interaction was significant, post-hoc pairwise comparisons of simple
effects (Control vs. Failure within each band; Theta vs. Gamma within each session type) were
performed using t-tests on linear contrasts of the fixed effects from the fitted mixed-effects model
(implemented via MATLAB's *coefTest* function). Each condition mean (e.g., Control–theta vs. 0) was
tested against zero using analogous linear contrasts on the same fitted model with the *coefTest* function.
For each family of tests (four post-hoc pairwise comparisons and four vs-zero comparisons), *p*-values
were adjusted using Holm–Bonferroni correction.

Temporal relationship between ongoing changes in landmark-control strength and changes in LFP
coherence

The speed-filtered windows included in the spectral computation of LFP–LFP coherence were
distributed unevenly relative to the animal's unwrapped position (track angle) in the landmark frame of
reference, because windows were selected over the time domain, while both running speed and
landmark rotation varied over time. Consequently, consecutive accepted 1-second windows were
typically separated by 30–90° (median 45°) yet collectively covered the full range of unwrapped
landmark-frame positions within each session. To investigate the temporal relationship between ongoing
changes in landmark control strength and changes in LFP–LFP coherence for every tetrode pair during a
session, we resampled the CSD measurements from these unevenly distributed windows onto a uniform
spatial grid (90° spacing) in the landmark frame of reference, using local linear kernel regression with a
Gaussian kernel (135° standard deviation). The resulting spatially uniform, resampled CSDs were used
to quantify the position-resolved z-imaginary coherence spectrum between each tetrode pair.

Landmark control-strength score was computed at the same spatial resolution as follows. For each 90°
spatial bin, we calculated the mean firing rates of all spatially correlated neurons (excluding those from
29e) and constructed a population vector (PV). The full-lap PV correlation at each bin was calculated as
the average Pearson correlation between the PV at that position and those from the same location one
and two laps earlier (offsets of 360° and 720°). The half-lap PV correlation was calculated as the
average correlation between the PV at that position and PVs from bins 180° and 540° earlier. At each
spatial bin, the landmark-control strength was defined as (full-lap PV correlation – half-lap PV
correlation)/2, where the division by 2 rescales the difference to a correlation-like, normalized range of –
1 to +1. The resulting, position-resolved, normalized landmark-control strength serves as a measure of
landmark-anchored neuronal firing in non-29e populations throughout each session.

Prior to cross-correlation analysis, both the position-resolved landmark control-strength scores as well as
z-imaginary coherence values were further smoothed with a Gaussian filter (240° standard deviation). At
each frequency within theta and gamma frequency bands, we computed the Pearson correlation
coefficient between the smoothed trajectories of landmark-control strength and z-imaginary coherence
across multiple spatial lags ranging from –18 to 18 bins (± 4.5 laps). For each frequency band (theta and
gamma), we identified the lag that maximized this correlation and recorded its corresponding correlation
coefficient. We assessed the statistical significance of these band-specific correlation coefficients
relative to null distributions obtained by circularly shifting the landmark control-strength trajectory by
offsets ranging from 32 [8 laps] to N–32 bins, where N is the total number of spatial bins (typically
>200). For each shifted trajectory, we repeated the cross-correlation analysis and recorded the maximum
correlation values within both theta- and gamma-bands for each LFP–LFP coupling measure. This
procedure yielded a null distribution for each band and each coupling measure. The observed correlation
for each band and coupling measure was considered statistically significant if it exceeded the 95[th]
percentile of its respective null distribution.

For tetrode pairs showing significant correlations in each frequency band, we collected the lags at which
their maximum correlations occurred and tested whether these lags differed significantly from zero
within each frequency band (Wilcoxon signed-rank test) and whether lags differed between frequency
bands or between LFP–LFP coupling measures (i.e., coherence vs. Granger causality [see below];
Wilcoxon rank-sum test).

Statistical association between ongoing changes in landmark-control strength and changes in LFP
coherence

To further test the statistical significance of positive associations between ongoing changes in landmark-
control strength and changes in LFP–LFP coherence while accounting for non-independence among
observations from each animal, we performed linear mixed-effects modeling. We fit separate models for
each frequency band (theta and gamma). Each model used the position-resolved landmark-control
strength as the dependent variable (y) and the z-imaginary coherence at the maximally correlated
frequency within that band as the predictor (x).

For each tetrode pair and frequency band, position-resolved coherence values were mean-centered to
remove baseline differences across tetrode pairs, ensuring that the analysis captured the average within-
tetrode-pair association rather than being dominated by tetrode pairs with higher or lower overall
coherence. After mean-centering, data from all tetrode pairs were concatenated to form the predictor (x)
and dependent variable (y) vectors for the model.

Because multiple 29e–MEC tetrode pairs were recorded from the same rat, and coherence (x) and
landmark-control strength (y) values across spatial bins represented repeated measurements from each
tetrode pair, observations were not independent. We modeled this nested dependence structure using
four random effects: random intercepts and slopes for rats, and random intercepts and slopes for tetrode
pairs nested within rats:

$$y \sim x + (1 + \text{x} \mid \text{RatID}) + (1 + \text{x} \mid \text{RatID:TetrodePairID}).$$

This structure accounts for: (i) rat-level variation in baseline landmark-control strength and in the
strength of the x–y relationship, and (ii) tetrode-pair-level variation in baseline landmark-control
strength and in the strength of the x–$\underline{y}$ relationship within each rat. By allowing these parameters to vary
across rats and tetrode pairs, the model captures correlations among observations from the same rat and
among repeated measurements from the same tetrode pair, providing a conservative test of the
population-level fixed effect (i.e., the population-average slope of the association between coherence [x]
and landmark-control strength [y]).

Models were fit using restricted maximum likelihood (using MATLAB's *fitlme* function). The fixed
effect for each model was evaluated using t-test with Satterthwaite-approximated denominator degrees
of freedom, as described above. To map the fixed effect strength to a normalized range, the fixed-effect
slope was converted to a standardized coefficient (β) by multiplying the raw slope by the ratio of the
global standard deviations of *x* and *y*.

LFP–LFP Granger causality and directed asymmetry index (DAI)

For Granger causality analysis, LFP signals were segmented into candidate windows of 3 seconds with
98% overlap (0.05 s step size). Due to the larger window size used in this analysis compared to LFP–
LFP coherence analysis, we adjusted the speed-filtering criterion: velocity was required to exceed 4
cm/s for more than 50% of the window, rather than the 75% threshold used previously. If a candidate
window did not satisfy this adjusted criterion, it was discarded.

Within each accepted 3-second window, LFP signals were detrended and transformed to the frequency
domain using a multi-taper Fourier transform with Slepian tapers (0.4 Hz resolution) via FieldTrip (*136*).
For low frequencies (<20 Hz), the entire 3-second window was used. For higher frequencies (≥20 Hz),
each 3-second window was divided into two equal, non-overlapping 1.5-second segments; spectral
analysis was performed separately for each segment to improve robustness against nonstationarity of
LFP signals in the higher frequencies, and the results were subsequently averaged. The combined
spectra from both frequency ranges yielded a full-frequency spectral representation for each window.

Using these full-frequency spectral representations, we computed conditional (partialized) Granger
causality (*56*, *151*) for each 29e-MEC tetrode pair and each accepted 3-second window in both
directions (i.e., $GC_{29e \to MEC}$ and $GC_{MEC \to 29e}$), conditioning on all other simultaneously recorded tetrodes
in 29e, MEC, and other parahippocampal regions via FieldTrip's *ft_connectivityanalysis* function
(configuration: method = 'granger', conditional = 'yes'). As with our use of imaginary coherency to guard
against spurious synchronization from common input, this multi-channel conditioning guards against
spurious directionality by removing indirect and common-source effects, to the extent such influences
are represented in the conditioning signals. However, whereas imaginary coherency is robust only
against instantaneous (zero-lag) common input, conditional Granger causality can mitigate common-
source confounds even when shared input reaches 29e and MEC with different delays (*57*, *152–154*). To
smooth the resulting Granger causality spectra, a moving average filter spanning 2 frequency bins (i.e.,
half-width 1 Hz) was applied in the frequency domain.

Baseline DAI levels

2149 For each 29e–MEC tetrode pair, the smoothed Granger causality spectra from all accepted windows in
2150 the initial stationary landmark epoch (G = 1) were averaged, yielding two mean spectra: one for the
2151 29e→MEC direction ($GC_{29e \to MEC}$), and one for the reciprocal MEC→29e direction ($GC_{MEC \to 29e}$). The
2152 directed asymmetry index (DAI; *58*) was then calculated for each pair as the normalized difference
2153 between the two:

$$\text{DAI} = \frac{GC_{29e \to MEC} - GC_{MEC \to 29e}}{GC_{29e \to MEC} + GC_{MEC \to 29e}}.$$

To assess the statistical significance of DAI values across tetrode pairs, we used a cluster-based
permutation approach. At each frequency, a one-sample t-test determined whether DAI values across
tetrode pairs for a region pair significantly deviated from zero. To control the familywise error rate, null
distributions of maximal cluster size were generated by randomly flipping the region labels of tetrode
pairs and repeating the t-test procedure for 1000 permutations. Clusters were defined as contiguous
frequencies with a significant deviation from zero ($p < 0.05$, two-sided). For each observed cluster, the

sum of t-statistics within the cluster was compared to the permutation distribution to generate cluster-
level, permutation-corrected *p*-values. Clusters with corrected *p* < 0.05 were considered significant.

Granger Causality and DAI changes from initial to final epoch

For each tetrode pair, we calculated the average Granger Causality spectra for both directions (i.e., 29e→MEC and MEC→29e) during the initial (G = 1) and final (G = G_{final}) experimental epochs by averaging the Granger causality spectra from all accepted windows in each of these epochs. To extract theta- and gamma-band-specific Granger causality values for each direction and each epoch, we identified the frequency bin with the maximum Granger causality within each band and averaged over ±1 Hz around that peak. For each band and direction, the initial epoch-to-final epoch change in Granger causality was calculated as $(GC_{final} - GC_{initial})/(GC_{final} + GC_{initial})$. To calculate the change in DAI, we first computed theta- and gamma-band DAI in each epoch from corresponding Granger causalities in both directions as above, and the change was calculated as $\Delta DAI = DAI_{final} - DAI_{initial}$.

Analysis of ΔDAI values across session types (*LM-control vs. LM-failure)* and across frequency bands (theta vs. gamma) was performed using a linear-mixed effects model with the same random effect structure as in ΔCoherence analysis (see above).

Temporal relationship between ongoing changes in landmark-control strength and changes in DAI

Position-resolved, uniformly sampled DAI values were computed using the same resampling procedure described for z-imaginary coherence (see "*Temporal relationship between ongoing changes in landmark-control strength and changes in LFP coherence*"). Briefly, Granger causality spectra from unevenly distributed windows were resampled onto a uniform spatial grid (90° spacing) using local linear kernel regression with a Gaussian kernel (135° standard deviation). DAI was then computed at each spatial bin from the resampled Granger causality spectra in both directions. Both position-resolved DAI and landmark-control strength trajectories were smoothed with a Gaussian filter (240° standard deviation) prior to cross-correlation analysis. Cross-correlation and statistical significance testing followed the same procedures as for z-imaginary coherence.

Statistical association between ongoing changes in landmark-control strength and changes in LFP coherence

To test the significance of associations between ongoing changes in landmark-control strength and changes in DAI while accounting for non-independence among observations, we applied linear mixed-effects modeling using the same approach as for z-imaginary coherence (see "*Statistical association between ongoing changes in landmark-control strength and changes in LFP coherence*"). Briefly, for each frequency band (i.e., theta or gamma), position-resolved DAI values were mean-centered within each tetrode pair, and data from all tetrode pairs were concatenated to form the predictor (*x*) and dependent variable (*y*) vectors. The model used the same random-effects structure described above:

$$y \sim x + (1 + x \mid \text{RatID}) + (1 + x \mid \text{RatID:TetrodePairID}),$$

where *x* denotes the position-resolved DAI and *y* denotes the position-resolved landmark-control strength. Models were fit using restricted maximum likelihood (using MATLAB's *fitlme* function), and fixed effects were evaluated using t-tests with Satterthwaite-approximated denominator degrees of freedom. Fixed-effect slopes were converted to standardized coefficients (β) as described above.

We also assessed the overall relationship between position-resolved trajectories of landmark control-strength score and band-specific LFP–LFP coupling measures across all sessions, animals, and tetrode pairs by fitting linear mixed-effects (LME) models. Separate models were fit for each coupling measure (z-imaginary coherence and DAI) and each frequency band (theta and gamma). In each model, the

position-resolved landmark control-strength score served as the predictor variable (fixed effect), and the
coupling measure at the frequency yielding the maximal correlation within the corresponding frequency
band (theta or gamma) served as the dependent variable. The models included random intercepts and
slopes for each animal and random intercepts for each tetrode pair nested within animal to account for
the repeated-measures structure of the data. Model parameters were estimated using restricted maximum
likelihood (using MATLAB's *fitlme* function), which also provided statistical significance of the fixed-
effect slope based on the t-statistic and associated *p*-value.

Cross-regional spike–LFP coupling

To quantify cross-regional spike–LFP coupling between 29e and MEC, we computed pairwise phase
consistency (PPC) spectra for each neuron relative to LFP channels in the other region. The LFP phase
at each frequency was estimated by convolving the continuous LFP trace with complex wavelet-like
filters constructed from DPSS tapers using the multitaper convolution method of the FieldTrip toolbox
(*132–134, 136*; frequency range: 1–140 Hz; spectral smoothing: 0.25 × center frequency, corresponding
to ~3 tapers; time window: 7 cycles per frequency). For each frequency, the phases at all spike times
within the analysis window (e.g., epoch 1) were extracted to form a spike-phase distribution, from which
we calculated PPC to obtain an unbiased estimate of spike–LFP coupling strength. We interpolated the
obtained PPC spectra to a uniform grid between 1 to 140 Hz with 1 Hz spacing.

Baseline coupling levels

We restricted the analysis windows for PPC computation to spikes from the initial epoch with stationary
landmarks (i.e., epoch 1, $G = 1$) and included only neurons with at least 40 spikes in that epoch. Because
neurons may show large variability in their coupling to LFP across laminar positions in the other region,
we constructed a representative spectrum for each neuron by taking, at each frequency, the maximum
PPC across all available LFP channels, capturing the strongest cross-regional coupling for between-area
comparisons.

Spike–LFP coupling change from initial to final epoch

To examine how cross-regional spike–LFP coupling changed over the course of a session, we compared
PPC values between the initial epoch ($G = 1$) and the final epoch ($G = G_{final}$), calculated for each neuron
in each region relative to the other region's LFP (29e relative to MEC and MEC relative to 29e). Only
neurons that fired at least 40 spikes in each of these epochs were included.

When multiple LFP channels were available, we used, for each neuron, its median PPC across channels
(instead of the maximum used for between-area comparisons in the baseline) to prevent within-neuron
initial-to-final epoch comparisons from being confounded by shifts in the optimal channel. From the
median PPC spectra for each neuron in each epoch, we extracted the maximum PPC value within the
theta and gamma bands, and quantified the initial-to-final change for each band as (PPC_{epoch3} −
PPC_{epoch1}) / (PPC_{epoch3} + PPC_{epoch1}).

Temporal relationship between ongoing changes in spike–LFP and landmark-control strength

To examine the temporal relationship between ongoing changes in cross-regional spike–LFP coupling
and landmark control strength, we performed a cross-correlation analysis. Only neurons with at least 8
spikes per lap in at least 75% of laps were included.

For each neuron, we computed the PPC spectrum relative to the LFP from the other region separately for
each lap (360° segment) in the reference frame of the rotating landmarks. When multiple LFP channels
from the other region were available, we computed the median PPC spectrum across channels for each
lap. Both the lap-wise PPC spectra and landmark control-strength scores were smoothed with a Gaussian

filter (320° standard deviation) and resampled onto a common grid of 90° spatial bins in the reference
frame of rotating landmarks.

We then performed a cross-correlation analysis between the position-resolved trajectories of landmark
control strength and PPC at each frequency, computing the Pearson correlation coefficient across spatial
lags ranging from −18 to +18 bins (±4.5 laps) and identifying the lag that maximized this correlation.
The maximal correlation coefficient, its corresponding lag and frequency within the theta and gamma
bands, and the corresponding PPC values at those frequencies were recorded for each neuron. (These
theta and gamma band PPC trajectories were also used in the subsequent section.)

Statistical significance of each neuron's maximal correlation coefficient within each frequency band
(i.e., theta or gamma) was assessed by comparison with a null distribution. The null was generated by
circularly shifting the landmark control-strength trajectory by offsets ranging from 32 bins (8 laps) to
N−32 bins, and repeating the identical cross-correlation procedure for each shift—i.e., computing
correlations across all frequencies and lags, and recording the maximum within each frequency band. A
neuron's observed correlation in a frequency band was considered statistically significant if it exceeded
the 95th percentile of its respective null distribution.

For neurons showing significant correlations in each region, we collected the lags at which their
maximum correlations occurred. We then tested whether these lags differed significantly from zero
within each region (Wilcoxon sign-rank test) and whether lags differed between 29e and MEC
(Wilcoxon rank-sum test).

Statistical association between ongoing changes in landmark-control strength and changes in spike–
LFP coupling

To test the significance of associations between ongoing changes in landmark-control strength and
changes in spike–LFP coupling while accounting for non-independence among observations, we applied
linear mixed-effects modeling using a similar approach as for LFP–LFP coherence (see "*Statistical*
association between ongoing changes in landmark-control strength and changes in LFP coherence").
Separate models were fitted for neurons in each region (29e neurons to MEC gamma and MEC neurons
to 29e theta), yielding two models in total.

For each neuron, we used the position-resolved, cross-regional PPC values identified in the previous
section, corresponding to the theta- and gamma-band frequencies with the maximum correlation
coefficient. These PPC values were mean-centered within each neuron to remove baseline differences,
ensuring that the analysis captured within-neuron variation rather than being dominated by neurons with
higher or lower overall coupling strength. After mean-centering, data from all neurons within a region
were concatenated to form the predictor (*x*: landmark-control strength) and dependent variable (*y*: PPC)
vectors for the model of each regions' neurons.

Because multiple neurons were recorded from the same rat, and values of position-resolved PPC and
landmark-control strength represented repeated measurements across spatial bins from all neurons
recorded in a session, we used random effects (both intercepts and slopes) at both the rat and neuron
levels. In mixed-effects notation, each model took the form:

$$y \sim x + (1 + x \mid \text{RatID}) + (1 + x \mid \text{RatID:CellID}),$$

This random-effects structure allowed the relationship between landmark-control strength (*x*) and PPC
(*y*) to vary across rats and neurons, capturing correlations among observations from the same rat and
among repeated measurements from the same neuron, and thereby providing a conservative test of the
population-level fixed effect. As in LFP–LFP analyses, model parameters were estimated using
restricted maximum likelihood (using MATLAB's *fitlme* function), and the fixed effect (i.e., the

population-average slope) was evaluated using t-tests with Satterthwaite-approximated denominator
degrees of freedom.

Identification of monosynaptic excitatory connections

2294 We ran the GLMCC algorithm (*155*) in likelihood-ratio mode (*156*), which returned a directed
2295 connectivity matrix among all simultaneously recorded neurons in each session. A neuron was classified
2296 as excitatory if $\geq$ 80% of its outgoing weights were positive. Monosynaptic connections of excitatory
2297 neurons were then defined as outgoing edges with weights exceeding a threshold of 1, as suggested in
2298 (*155*).

2299

2300 **Supplementary Figures**

29e: Area 29e, **MEC:** Medial entorhinal cortex, **PA:** Parasubiculum, **PO:** Postrhinal cortex, **PR:** Dorsal presubiculum

Fig. S1. Representative Nissl-stained sagittal sections showing tetrode recording sites across parahippocampal regions in Rat 1061. Four sections are arranged from lateral (top left) to medial (bottom right). As in Fig. 1, colored arrowheads indicate the superficial borders of recorded parahippocampal regions namely, area 29e (black), MEC (orange), PA (blue), PO (pink), and PR (green). Dots mark tetrode recording locations, with dot color corresponding to the recorded region.

A

994, section 52　　994, section 59　　995, section 48

995, section 59　　1062, section 58　　1062, section 64

B

1041, section 60　　1062, section 70

29e: Area 29e, **MEC:** Medial entorhinal cortex,
PA: Parasubiculum, **PO:** Postrhinal cortex,
PR: Dorsal presubiculum

C

Single-neuron
physio features

Neuron 1
Neuron M

Single-neuron
model

XGBoost

CV (training)
• Features
  selection
• Parameter
  optimization
• Probability
  calibration

Stacked-generalization

Permutation-
invariant
transform

Logistic
regression
model

CV (training)
• Synthetic multi-
  neuron training
  data generation
• Parameter
  optimization
• Probability
  calibration

Threshold application

Negative
class

Reject

Positive
class

CV (training)
• Synthetic multi-
  neuron training
  data generation
• Threshold
  optimization

D

29e vs. Rest

		Simultaneously recorded neuron count			
		N = 1	N = 3	N = 5	N = 7
Rejection rate	Positive class	0.44	0.24	0.13	0.12
	Negative class	0.36	0.29	0.23	0.16
Performance after reject	Sensitivity	0.56	0.96	0.99	1
	Specificity	0.88	0.94	0.95	0.96

PR vs. MEC/PA/PO

		Simultaneously recorded neuron count			
		N = 1	N = 3	N = 5	N = 7
Rejection rate	Positive class	0.45	0.42	0.38	0.35
	Negative class	0.44	0.43	0.4	0.36
Performance after reject	Sensitivity	0.6	0.67	0.8	0.85
	Specificity	0.77	0.83	0.86	0.86

2306

Fig. S2. Histologically identified recording sites in area 29e and machine-learning pipeline using single-neuron physiological properties applied to resolve ambiguous histological cases (see Methods).

(A) Example histological sections illustrating recording sites in area 29e from three rats (994, 995, and 1062). See fig. S1 for an additional histologically identified 29e site (rat 1061). Colored arrowheads indicate boundaries between areas following the same color scheme as Fig. 1A and fig. S1: area 29e (black), MEC (orange), PA (blue), PO (pink) and PR (green).

(B) Example histological sections illustrating recording sites with ambiguous localization between area 29e and neighboring regions. Left: Rat 1041, section 60 shows a recording location at the 29e/PA border (indicated by the red star). Application of the machine learning model (see panels C, D and the methods) classified this location as area 29e. Right: Rat 1062, section 70 shows a recording location at the 29e/PR border (red star) that was likewise classified as area 29e by the machine learning model.

(C) Machine-learning model architecture. When histological identification of a tetrode's recording region was ambiguous (as in panel B), we employed a machine-learning classification approach using physiological features of single neurons recorded on the tetrode to identify the region the tetrode was located in. Because ambiguous tetrode tracks were most often along the borders of area 29e or PR with neighboring regions, this approach relied on two separate one-vs.-rest classification pipelines (29e vs. rest and PR vs. rest) to distinguish recording sites in 29e from other regions and those in PR from other regions, respectively. Each pipeline was trained exclusively on neurons from tetrodes with unambiguous histological region assignments (ground-truth data) and proceeded through three steps: *Step 1 (Single-neuron classifier):* An XGBoost classifier used each neuron's physiological features to estimate the probability that it belonged to the target region. Cross-validation (CV) on the ground-truth training data was used for feature selection, hyperparameter optimization, and probability calibration. *Step 2 (Stacked-generalization as multi-neuron evidence accumulator)*: Probability outputs from the single-neuron classifiers were aggregated across all simultaneously recorded neurons on the same tetrode into a single tetrode-level probability estimate by applying a permutation-invariant transformation followed by a logistic-regression "meta-model" used for stacked-generalization. For hyperparameter optimization and probability calibration of this meta model, CV was performed on synthetic multi-neuron samples generated by combining and subsampling ground-truth neurons. *Step 3 (Reject-option thresholding):* Class-specific confidence thresholds were applied to the tetrode-level probabilities to assign either the positive (e.g., 29e) or negative (e.g., not 29e) label only when the corresponding probability exceeded its threshold; otherwise, we made no assignment (reject). Threshold values were optimized via CV on the synthetic multi-neuron samples (as in step 2) to balance coverage and accuracy. See Methods for further details.

(D) Machine-learning model validation and application criteria. To validate model performance, we performed tetrode-stratified fivefold cross-validation (CV), ensuring neurons from the same tetrode never appeared in both training and test sets. The table reports performance metrics evaluated on held-out test data as a function of the number of simultaneously recorded neurons per tetrode (N = 1, 3, 5, or 7). Because individual held-out tetrodes had variable neuron counts, synthetic test cases were generated by pooling neurons across held-out tetrodes within each class and randomly sampling N neurons to simulate tetrodes with a specific neuron count. For each pipeline (29e vs. rest, top; PR vs. rest, bottom), metrics include: (i) the rejection rate: the fraction of test cases where the pipeline declined to assign a label because neither the positive-class nor negative-class probability exceeded its respective confidence threshold. Rows show separate rejection rates for tetrodes truly belonging to the positive class (e.g., 29e/PR) versus the negative class (e.g., not 29e/not PR). Lower rejection rates indicate the pipeline more frequently made confident assignments. (ii) performance after reject: Among the test cases that were not rejected (i.e., received a confident assignment), sensitivity is the proportion of true positive-class tetrodes correctly identified as positive, and specificity is the proportion of true negative-class tetrodes correctly identified as negative. As the number of simultaneously recorded cells on a tetrode increased, rejection rate decreased, and sensitivity/specificity increased for both models. To ensure reliable assignments (both sensitivity and specificity ≥ 0.85), we applied the 29e vs. rest pipeline only to tetrodes with at least 5 simultaneously recorded neurons and the PR vs. rest pipeline only to tetrodes with at least 7 simultaneously recorded neurons. Final models trained on all ground-truth data (i.e., data from tetrode recording locations with histologically unambiguous region assignments) were then applied to classify histologically ambiguous tetrode recordings meeting these criteria (see Methods).

2353

Fig. S3. Examples of landmark anchoring in 29e relative to other regions. Figure organization and panels are similar to Fig.
1E-F.
(A) Example *LM-control* session in which spatially correlated neurons remain anchored to the rotating landmarks. Top panel
shows the experimental gain protocol (gain = 1 at baseline, then ramped to a higher target gain and held there for 10-30 laps).
Middle panel shows the spatial firing rates of simultaneously recorded neurons across laps in the landmark reference frame.
Neurons fired at fixed angles relative to landmarks throughout the session, producing stationary, horizontal bands of firing
rates in the landmark frame. Bottom shows the spatial autocorrelograms of the last 12 laps for example neurons. Clear peaks
indicate strong 1-lap periodicity; the accompanying score quantifies this periodicity (see "landmark-locking score" in
Methods).
(B) Example *LM-failure* session in which most spatially correlated neurons from regions other than 29e lose landmark
anchoring. Conventions as in (A). Around laps 65-75, firing fields of neurons recorded outside 29e started to drift in the
landmark frame, yielding autocorrelograms that were not lap-to-lap periodic and that exhibited low landmark-locking scores.
By contrast, a subset of 29e neurons (like the example in black) remained locked to landmarks, firing stably at fixed angular
locations in the landmark frame. In this example, the 29e cell had 2 firing fields: one at ~90° and one at ~180°. Early in the
session, the field at ~90° fired variably while the field at ~180° was more consistent. Shortly after the landmark-control failure
outside 29e, the field at ~90° strengthened, while the field at ~180° weakened. This reversed pattern persisted for about 10
laps, until lap 90. After this point, fields returned to their original relative firing levels, with the field at ~180° firing more
consistently than the field at ~90°. Despite these fluctuations in relative field strength, the locations of fields remained
anchored to landmarks throughout. This landmark-anchoring of 29e neurons was reflected in their spatial autocorrelograms
as a clear 1-lap periodic pattern (black trace in the bottom inset).

2374

2375

2376
2377

Fig. S4. Greater landmark anchoring in neurons of 29e compared to other regions during open-loop manipulation of visual landmarks. **(A)** Example rat and landmark trajectories in closed- and open-loop experiments. Top: Closed-loop (G > 1) session in which landmark rotation was coupled to the rat's movement. Bottom: Open-loop session in which landmark rotation was yoked to a rat's trajectory from a different session; note multi-second bout of immobility (starting at ~15 s) where the rat was stationary while landmarks rotated past it.

(B) Landmark-frame spatial firing rate map plot across laps in the landmark frame of reference (format same as in Fig. 1E,F) for an example 29e neuron (middle; black) and a PR neuron (bottom; green). The open-loop experiment gain is shown on top. The example PR neuron loses landmark anchoring towards the end of the gain ramp epoch, whereas the 29e neuron remained anchored to landmarks throughout. Notably, the 29e neuron maintains a lap-to-lap periodicity of 1 even when landmarks are moved in open-loop fashion, as evident in the autocorrelation trace, in contrast to the PR neuron, which is no longer controlled by landmarks.

(C) Distributions of landmark-locking scores in open-loop experiments for 29e (n = 29 neurons) versus non-29e (n = 39 neurons) regions. Neurons outside area 29e were pooled into a single non-29e group due to lower sample sizes with unambiguous histological assignments and similar closed-loop response profiles across regions. Scores were computed with velocity filtering in the laboratory frame (top) and in the landmark frame (bottom). 29e neurons showed higher landmark-locking scores in both cases (Lab frame: median [IQR] 29e = 0.431 [0.193, 0.741], Non-29e = 0.231 [0.107, 0.536], Wilcoxon rank-sum test: $Z = 2.28$, $p = 0.023$; Landmark frame: median [IQR] 29e = 0.484 [0.231, 0.725], Non-29e = 0.275 [0.139, 0.474], Wilcoxon rank-sum test: $Z = 2.08$, $p = 0.037$).

(D) Comparison of circular spatial firing rate maps across track angles in the landmark reference frame. Representative examples are shown for a landmark-anchored 29e neuron (top) and a landmark-anchored PA neuron (bottom) under open-loop conditions when the rat was *mobile* (left) and *stationary* (right; see Methods for details). Note that the spatial firing rate map for the 29e neuron shows greater similarity between the *mobile* and *stationary* conditions compared to the PA neuron.

(E) Cumulative distributions of landmark frame rate-map correlations between stationary and mobile periods (*stationary*: the animal's speed in laboratory frame < 4 cm/s, *mobile*: otherwise; see Methods) for landmark-anchored 29e (n = 14) and non-29e (n = 11) neurons. 29e landmark-anchored neurons showed higher stationary-vs-moving rate map correlations than non-29e landmark-anchored neurons (median [IQR] 29e = 0.856 [0.746, 0.924], Non-29e = 0.306 [-0.042, 0.425], Wilcoxon rank-sum test: $Z = 3.75$, $p < 0.001$).

(F) Proportion of landmark-anchored neurons with significant stationary-vs.-moving rate-map similarity based on a shuffle test (10,000 circular-shifted shuffles of the stationary map relative to the moving map; α=0.05). Most 29e neurons remain significantly landmark-anchored when the animal was stationary, unlike non-29e neurons (Fisher's exact test: $p < 0.001$).

Fig. S5. Regional firing-rate responses to landmark removal and relationship between contrast tuning and landmark locking. **(A)** Average baseline-normalized firing-rate deviation around landmark removal (time 0) across all tetrodes in each region. Because landmarks were removed only once at a single timepoint in each session, most neurons with spatially localized firing fields would not have been active at that moment (i.e., rat's location at the time of removal would not necessarily be within the spatial receptive field boundaries), making landmark removal effects difficult to quantify reliably at the single-neuron level. We therefore used tetrode-level multi-unit activity. For each tetrode, we computed an unsmoothed peri-event firing-rate trace, standardized it using that tetrode's own pre-removal baseline mean and variability (z-scoring relative to the pre-removal window), and took the absolute value to capture upward and downward deviations from the baseline mean symmetrically. The resulting rectified z-scored trace of each tetrode was then divided by its mean value during the pre-removal, baseline period. and traces were averaged across tetrodes within each region to compute the regional mean. Although the rescaled trace may show large fluctuations around 1 for a single tetrode, the regional means are expected to fluctuate closer to 1 during the pre-removal baseline period as a result of averaging across many tetrodes, as seen in the figure for all regions when t < 0. For the post-removal period, however, firing rates of tetrodes in 29e and PO consistently showed large deviations (|z| > 1) from their average baseline levels, , indicating a region-wide response to the removal of landmarks. **(B)** Absence of significant correlation between landmark-locking scores and contrast tuning strength in MEC and PR neurons. Each dot represents a single neuron's landmark-locking score (*x*-axis) versus its partial R² for landmark contrast tuning (*y*-axis) in MEC (left) and PR (right). Pearson's correlation coefficients (*r*) and corresponding *p*-values (computed using Fisher z-transformed landmark-locking scores) are shown above each plot. An ANCOVA with landmark-locking score as a continuous covariate and region as a categorical factor revealed a significant positive relationship between landmark-locking and contrast tuning across regions ($F(1, 67) = 5.68$, $p = 0.020$). There was also a significant main effect of region ($F(2, 67) = 3.71$, $p = 0.030$), likely reflecting higher contrast tuning in 29e (Fig. 2), but no significant interaction ($F(2, 67) = 1.92$, $p = 0.155$).

Fig. S6. Two-dimensional allocentric spatial tuning in open arena across parahippocampal regions.
(A) Schematic of the square open arena apparatus used to identify functional parahippocampal cell types. The open arena was placed on the floor in a room with multiple distal visual cues (not shown), providing stable landmarks.
(B) Tuning to allocentric position and head direction across regions. Five representative neurons are displayed per region. Same format as Fig. 3A.

2450

Fig. S7. Egocentric tuning of example neurons across different regions. Same format as Fig. 3C.
(A) Example HD neurons that show classic, allocentric tuning. The preferred HD vectors are roughly parallel at all locations
across the open arena, as expected from their allocentric nature. Some cells showed a modulation by both place and HD. Note
that it is impossible (and meaningless) to distinguish an allocentric HD cell from an egocentric HD cell with an anchor point
infinitely far away. Rat ID and tetrode number for each neuron are indicated along with the region identity above the
respective panels.
(B) Example egocentric HD neurons from 29E.
(C) Example egocentric HD neurons from MEC, PA, PO, and PR.

2459

Fig. S8. Workflow for computing the anchor-free egocentric-vector-field (EVF) score, illustrated with four example head-
direction (HD) cells.
A key question for HD-tuned neurons is whether their preferred direction is allocentric or egocentric. Standard approaches to
quantifying egocentric tuning rely on an exhaustive search for an optimal anchor point in the environment that serves as a
reference point to calculate the bearing angle between the rat's current allocentric HD at a particular location in the
environment and an anchor point (as illustrated by the rat cartoons on the left). The sharpest tuning curve among multiple
candidates (each corresponding to a different anchor point) is considered the "true" egocentric tuning curve, against which a
classic allocentric tuning curve is compared. This procedure creates a statistical bias in favor of egocentric tuning (the best
curve selected from multiple candidates) over allocentric tuning (a single curve). To enable unbiased comparison, we
developed an anchor-free Egocentric Vector Field (EVF) score that quantifies the spatial consistency of local head-direction
preferences across the arena without specifying any anchor point. See Methods for further details.
Figure organization: Each row corresponds to one example neuron. The leftmost column is for illustration purposes and
approximately depicts the neuron's preferred egocentric bearing, defined as the angle (red arc) between the allocentric HD
vector (black arrow) located at the rat's head and the head-to-anchor point vector (blue dashed line). The remaining columns
illustrate the three-step workflow for computing the EVF score to quantify such egocentric tuning of single neurons.
Input: HD vector field. The HD tuning curve within each spatial bin (x, y) is reduced to a single vector corresponding to the
first Fourier component of the circular tuning curve (i.e., the single sinusoid that, in a least-squares sense, best fits the full
circular tuning curve). The phase of this Fourier component is mathematically equivalent to the preferred HD (i.e., the mean
angular direction) and its amplitude equals the summed firing rate of the HD tuning curve × the mean resultant length of the
HD tuning curve. Collectively, these HD tuning vectors define a two-dimensional vector field $v_{hd}(x, y) = [P(x, y); Q(x, y)]$,
where P and Q are the horizontal and vertical components of each vector, respectively.
Step 1: Curl- and divergence-maps. For every spatial bin, we compute a curl score as $curl(x, y) = \frac{\partial Q(x,y)}{\partial x} - \frac{\partial P(x,y)}{\partial y}$ and a
divergence score as $div(x, y) = \frac{\partial P(x,y)}{\partial x} + \frac{\partial Q(x,y)}{\partial y}$. The resulting scalar values are displayed as heat maps (curl on top,
divergence on bottom), where lighter colors correspond to values close to 0 and darker colors correspond to progressively
larger absolute values. Therefore, the darkest color can represent either a large positive (e.g., CW, centrifugal) or a large
negative (e.g., CCW, centripetal) value, depending on the neuron. The color scale is normalized separately for each neuron.
Step 2: Curl/divergence vectors. For each spatial bin, the ordered pair (curl, div) is treated as a new 2D vector (horizontal =
curl, vertical = div), forming the curl-divergence field shown next. Curl captures rotational patterns (preferred egocentric
bearing to the left or right; top two rows). Divergence captures radial patterns (preferred egocentric bearing to the front or
back; bottom two rows).
Step 3: Calculation of the EVF score. All (curl, div) vectors are replotted in polar coordinates, with the black thick arrow
showing their mean resultant vector. The circular statistics of this (curl, div) vector collection define the preferred bearing
angle of an egocentric neuron and its EVF score as follows:

•   The circular mean indicates the neuron's preferred egocentric bearing angle underlying the orientation of its preferred
HD vectors across the arena as shown in the HD vector field in the input column. The examples from top to bottom
illustrate this relationship for bearing angles associated with the four cardinal egocentric directions as depicted by the rat
cartoons: rightward-pointing (curl,div) (i.e., pure positive curl) for right-anchoring, leftward-pointing (curl,div) (i.e., pure
negative curl) for left-anchoring, upward-pointing (curl,div) (i.e., pure positive divergence) for front-anchoring, and
downward-pointing (curl,div) (i.e., pure negative divergence) for back-anchoring.
•   Measuring the circular dispersion of the vector collection (i.e., its disorganization across the arena) with the mean
resultant length gives the anchor-free EVF score (0 = randomly distributed vectors, 1 = perfectly aligned vectors).
2501

Fig. S9. Functional properties of egocentric- versus allocentric-HD tuned neurons across parahippocampal regions.
(A) Relationship between neurons' landmark-locking scores (from *LM-failure* sessions) and their preferred frame of reference (egocentric vs. allocentric) in MEC, PA, PO, and PR. Unlike area 29e, where neurons are approximately equally split between allocentric and egocentric reference frames (Fig. 3D) and landmark-locking scores show considerable heterogeneity during LM-failure sessions (Fig. 2H, right), other regions showed imbalanced egocentric-allocentric splits (predominantly allocentric) with relatively homogeneous (and lower) landmark-locking scores. Given the small number of egocentric neurons and limited variability in landmark-locking scores, any observed differences within non-29e regions could reflect session-to-session or animal-to-animal variability rather than true differences in landmark anchoring between egocentric and allocentric neurons. We therefore did not perform a direct statistical comparison between 29e and other regions. Nevertheless, data from other regions are shown here for completeness, with *p*-values from Wilcoxon's rank-sum tests (egocentric vs. allocentric) displayed above each plot for reference.
(B) Relationship between neurons' partial R^2 values for landmark-contrast tuning (from luminance-manipulation sessions) and their preferred reference frame (egocentric vs. allocentric) in area 29e (left), MEC (middle), and PR (right). As with landmark-locking scores (panel A), contrast tuning was stronger and more heterogeneous in 29e than in other regions (Fig. 2), making this comparison primarily relevant for 29e. For similar reasons as described in panel A, combined with lower neuron counts in luminance-manipulation sessions overall, we did not perform a direct statistical comparison across areas. Instead, we provide within-region comparisons (p-values from two-sample *t*-tests shown above each plot) for reference. Conventions as in panel A. The panels shows that even though no region shows statistical significance, all regions show a trend in the same direction.
(C) Polar histograms showing the distribution of preferred bearing angles for egocentric HD cells in MEC, PA, PO, and PR. Same format as Fig. 3G. All regions showed a qualitative leftward bias in preferred bearing angles, contralateral to the recording hemisphere. This lateralization was statistically significant in MEC (left: n = 30, right: n = 14; binomial test, *p* = 0.023) and PA (left: n = 37, right: n = 17; *p* = 0.009), but not in PO (left: n = 11, right: n = 7; *p* = 0.481) or PR (left: n = 13, right: n = 8; *p* = 0.383).

2529

Fig. S10. Reduced theta power and firing rates but comparable gamma power in area 29e compared to other areas.

(A) Cumulative distribution of 1/f-corrected theta-band power across regions, computed at unique tetrode locations (29e: n =

20, MEC: n = 104, PA: n = 64, PO: n =22, PR: n = 24). A linear-mixed effects model including region identity as the fixed

effect and two random intercepts (rat identity and region identity within each rat) showed a significant main effect of the

region ($F(4, 9.7) = 4.6$, $p = 0.024$). Post-hoc testing of pairwise contrasts comparing each region to 29e (Holm-Bonferroni

corrected) based on the fitted model determined that theta-band power in 29e was significantly lower than MEC ($t(10.1) = -$

3.07, p = 0.0345 and PO ($t(14.4) = -3.37$, $p = 0.018$) but not PA ($t(10.9) = -1.86$, $p = 0.179$) or PR ($t(12.2) = -0.154$, $p =$

0.880).

(B) Region-wise cumulative distribution of 1/f-corrected gamma-band power computed at unique tetrode locations. (sample

size same as A). A linear mixed-effects model with the same fixed- and random-effects structure as the one in panel A

identified main effect of the regional identity as statistically significant ($F(4, 220.4) = 0.038$). When post-hoc pairwise

contrasts from the fitted model compared each region against 29e, however, we found that gamma-band power in 29e was

comparable to MEC ($t(204.2) = 0.32$, $p = 0.992$), PA ($t(203.6) = -1.25$, $p = 0.793$), PO ($t(217.3) = -1.29$, $p = 0.793$), and PR

($t(225.3) = -0.68$, $p = 0.992$). Thus, although there was a statistically significant difference among recorded parahippocampal

regions, it was not driven by a difference between 29e and another region.

(C) Cumulative distribution of single neuron firing rates across regions. 29e neurons exhibited significantly lower mean

firing rates (median [IQR] = 2.34 [0.90, 4.81]) compared to other regions except PO (MEC = 3.03 [1.45, 8.11], PA = 3.86

[1.61, 9.75], PR = 2.90 [1.42, 7.79], PO = 3.07 [0.97, 7.31]; Kruskal-Wallis test $\chi^2(4) = 19.4$, $p < 0.001$; post-hoc Wilcoxon

rank-sum tests with Holm-Bonferroni correction 29e vs. MEC: $Z = -2.97$, $p = 0.008$; 29e vs. PA: $Z = -4.24$, $p < 0.001$; 29e

vs. PO: $Z = -1.61$, $p = 0.105$; 29e vs. PR: $Z = -2.28$, $p = 0.044$).

2550

2551

Fig. S11. Relationship between single neurons' spike–LFP theta phase coupling and their functional properties across non-29e regions.

(A) Theta PPC in neurons with low (≤ median) versus high (> median) landmark-locking scores in non-29e regions during *LM-failure* sessions in the Dome VR. Dots indicate single neurons; bars show mean ± s.e.m.; p-values from Wilcoxon's rank-sum tests on arcsinh-transformed PPC values are shown above each plot. Unlike area 29e, where landmark-locking scores show considerable heterogeneity (Fig. 2H, right), other regions showed relatively homogeneous (and lower) landmark-locking scores, making their median-split groups sensitive to session-to-session and animal-to-animal variability rather than truly distinguishing neurons based on their landmark anchoring. Additionally, baseline theta PPC differed substantially across regions (Fig. 4F), complicating any cross-regional comparison of the effect. We therefore did not perform a direct statistical comparison between 29e and other regions. Nevertheless, data from other regions are shown here for completeness, with p-values displayed above each plot for reference.

(B) Same data as (A), shown as scatter plots to illustrate the continuous relationship between theta PPC (*x*-axis) and landmark-locking score (*y*-axis) across single neurons from each region recorded in *LM-failure* sessions. Note that, unlike panel A, this panel includes, because the main text and the accompanying main figure (Fig. 4) showed 29e data only in the median-split format as in panel A of this figure. Above each region's plot, the Spearman's correlation coefficient (r_s) and the corresponding p-value are given. Only 29e showed a significant correlation, which was negative.

(C) Same as (B) but with arcsinh-transformed theta PPC on the x-axis to account for the skewed distribution of PPC values, which can take zero or small negative values. The transformation spreads the data more evenly along the *x*-axis by reducing compression at low PPC values, allowing use of Pearson correlation (r_p) instead of Spearman's (r_s). The negative relationship in 29e remains significant, appearing more uniformly linear across the full range of transformed values, whereas it remains non-significant in other regions.

(D) Theta PPC in neurons with low (≤ median) versus high (> median) partial R^2 for contrast tuning in MEC and PR during luminance manipulation sessions in the Dome VR. Conventions as in panel A. As with landmark-locking scores (panel A), contrast tuning was stronger and more heterogeneous in 29e than in other regions, making median-split groups in non-29e regions more sensitive to session-to-session and animal-to-animal variability rather than truly distinguishing neurons based on their contrast tuning. Additionally, baseline theta PPC differed substantially across regions (Fig. 4F), and neuron counts were lower in luminance-manipulation sessions overall. We therefore did not perform a direct statistical comparison between 29e and other regions. Nevertheless, data from other regions are shown here for completeness.

(E) Same data as (A), shown as scatter plots to illustrate the continuous relationship between theta PPC (x-axis) and partial R^2 for contrast tuning across single neurons from each region recorded in luminance manipulation sessions. As with panel B, we included Area 29e in this plot because Fig. 4 of the main text showed 29e data only in the median-split format as in panel D.

(F) Same as (E) but with arcsinh-transformed theta PPC to reduce skewness, which is more suitable for Pearson's correlation (r_p). The significant negative relationship in 29e persisted after transformation, whereas other regions remained non-significant.

(G) Theta PPC in egocentric versus allocentric neurons across non-29e regions. Conventions as in panel A. As explained in fig. S9A, non-29e regions showed imbalanced egocentric–allocentric splits (predominantly allocentric), unlike area 29e, where neurons are approximately equally split between egocentric and allocentric reference frames (Fig. 3D). For similar reasons as described there, we did not perform a direct statistical comparison between 29e and other regions. Nevertheless, data from other regions are shown here for completeness, with within-region p-values displayed above each plot for reference. Although some regions did not reach statistical significance, all showed a trend in the same direction, with allocentric neurons exhibiting higher theta PPC. than egocentric neurons, suggesting this as a general organizing principle of spatial processing across the parahippocampal cortex.

2595

Fig. S12. Changes in directed LFP interactions of 29e and MEC tetrode pairs during landmark rotations and the relationship
between their cross-regional spike–LFP coupling and landmark anchoring.
(A) Normalized changes in theta- and gamma-band Granger causality between 29e and MEC LFPs from the initial epoch to
the final epoch across *LM-control* and *LM-failure* sessions. Theta band GC showed significant changes in the MEC→29e
direction during both *LM-control* (mean ± s.e.m. = –0.05 ± 0.01, Holm-Bonferroni corrected one-sample t-test $t(15) = -3.2$, p
= 0.022) and *LM-failure* sessions (0.06 ± 0.01, $t(17) = -3.8$, $p = 0.005$); theta-band GC changes in the reciprocal direction
(29e→MEC) were not significant during both *LM-control* (–0.018 ± 0.008, $t(15) = p = 0.057$) and *LM-failure* sessions (0.003
± 0.009, $t(17) = 0.35$ $p = 0.726$). For gamma-band GCs, changes were significant in the 29e→MEC direction during *LM-*
failure sessions (–0.047 ± 0.014, $t(17) = -3.3$ $p = 0.013$) and marginally significant during *LM-control* sessions (0.041 ±
0.017, $t(15) = 2.4$ $p = 0.074$). Three-way repeated measures ANOVA (factors: session type, frequency band, and direction)
identified significant session type × frequency band ($F(1,32) = 16.2$, $p < 0.001$) and significant session type × direction
($F(1,32) = 25.9$, $p < 0.001$) interactions, in addition to a significant main effect of direction ($F(1,32) = 5.0$, $p = 0.031$). Holm-
Bonferroni corrected post-hoc pairwise t-tests revealed that, within the theta band, changes in MEC→29e GC during *LM-*
failure sessions were significantly higher compared to the changes in [i] MEC→29e GC during *LM-control* sessions ($t(32) =$
4.9, $p < 0.001$), [ii] 29e→MEC GC during *LM-control* sessions ($t(32) = 4.5$, $p < 0.001$), and [iii] 29e→MEC GC during *LM-*
failure sessions ($t(17) = 2.9$, $p = 0.032$). Within the gamma band, directionally opposite but statistically similar differences
were observed: changes in 29e→MEC gamma-band GC during *LM-failure* sessions were significantly lower compared to the
changes in [i] 29e→MEC gamma-band GC during *LM-control* sessions ($t(32) = -3.9$, $p = 0.002$), [ii] MEC→29e gamma-

band GC during *LM-control* sessions ($t(32) = -3.4$ $p = 0.008$), and [iii] MEC→29e gamma-band GC during *LM-failure* sessions ($t(17) = -3.1$, $p = 0.027$).

(B-C) Permutation-based significance testing of the frequency-selective landmark locking–LFP phase coupling pattern for 29e **(B)** and MEC **(C)** neurons. At each permutation, landmark-locking scores from LM-failure sessions were randomly shuffled across 29e or MEC neurons. For each frequency bin (as in Fig. 6C), the correlation across neurons between their baseline PPC values (from the initial stationary-landmark epoch) and the shuffled landmark-locking scores was recalculated, generating a new correlation spectrum for each permutation. From this spectrum, the difference between the maximum correlation above 50 Hz and the minimum correlation below 20 Hz was computed. Repeating this procedure for 1000 permutations yielded null distributions of frequency-based correlation-coefficient differences. The observed difference (solid line) significantly exceeded the 95th percentile of the permutation distribution (dashed line) in 29e (B; $p = 0.006$) but not in MEC (C; $p = 0.078$).

(D-E) Relationship between landmark locking of 29e and MEC neurons and loss of their cross-regional spike–LFP coupling during LM-failure sessions. (D) Median difference in landmark-locking scores between neurons that lost coupling ("PPC loss") versus those that gained coupling ("PPC gain") for 29e (black) and MEC (orange) populations across frequencies. At each frequency, neurons were divided into PPC loss or PPC gain groups, and the median difference in landmark-locking scores between groups was calculated across the population. The statistical significance of the difference between medians at each frequency was assessed with a Wilcoxon signed-rank test. To control for multiple comparisons, a cluster-based permutation test was used in conjunction with this rank test: contiguous frequency bins with uncorrected $p < 0.05$ were defined as clusters, and cluster-level statistics were compared to a null distribution generated by 1000 random permutations of group labels. Asterisks indicate frequency ranges with significant differences after cluster-level, permutation-based correction ($p < 0.05$). Among 29e neurons, those that lost coupling to MEC gamma had significantly higher landmark-locking scores than those that gained coupling, whereas those that gained coupling to MEC theta had significantly lower landmark-locking scores. No such pattern was observed for MEC neurons.

(E) Logistic regression analysis testing whether landmark-locking score of a neuron predicts its "PPC loss" or "PPC gain" status, after controlling for baseline coupling levels. Because the frequency-dependent pattern among 29e neurons in panel D closely resembled the relationship between their landmark-locking score and their baseline PPC values (Fig. 6C), baseline coupling level may have confounded the observed differences in landmark-locking score between "PPC loss" and "PPC gain" groups by affecting both whether a neuron loses or gains coupling toward a mean level and its degree of landmark anchoring. To control for this potential confound, we performed a logistic regression analysis, using baseline PPC values and landmark-locking scores of each neuron during *LM-failure* sessions as predictors, and the direction of PPC change (loss vs. gain) as the outcome variable. The log odds ratio (β coefficient from logistic regression; *y*-axis) for landmark-locking score as a predictor of PPC loss (ΔPPC < 0) versus gain (ΔPPC > 0) is plotted as a function of frequency (*x*-axis) for 29e neurons' coupling to MEC LFP (black) and MEC neurons' coupling to 29e LFP (orange). A log odds ratio of 0 indicates no change in the odds of PPC loss per one standard-deviation increase in landmark-locking score, while a log odds ratio of 1.5 indicates an increase in odds by a factor of $\exp(1.5) \approx 4.5$ per the same amount of change in landmark-locking score. Asterisks and grey, thickened segments indicate the significant frequency ranges, determined by a cluster-based permutation test as before. Inset shows landmark-locking scores for 29e neurons grouped by whether they lost (ΔPPC < 0) or gained (ΔPPC > 0) coupling at the frequency with largest odds ratio. Overall, the logistic regression revealed that, after accounting for baseline PPC levels, the landmark-locking score remained a significant positive predictor of 29e neuron's PPC loss to MEC gamma (particularly 100–120 Hz; Fig. 6D). No significant association remained relative to lower-frequency oscillations of MEC LFP (e.g., theta), nor was it observed for MEC neurons' coupling to 29e LFP across the entire frequency spectrum. Collectively, the results in panels D and E indicate that landmark-anchored 29e neurons lost their coupling to MEC gamma, independent of their initial coupling levels, as landmark control was lost over parahippocampal populations during *LM-failure* sessions.

2666

2667

Fig. S13. Distributions of landmark-locking score differences between putative pre- and post-synaptic neuron pairs across
regions. Putative monosynaptic connections were identified from spike cross-correlograms between simultaneously recorded
neurons. Interregional connections were too scarce for a meaningful analysis, so we examined intraregional connections
within each region. Each histogram shows the distribution of landmark-locking score differences (presynaptic – postsynaptic)
for putative monosynaptic pairs recorded in LM-failure sessions. For each region, we tested whether landmark-locking scores
differed between pre- and postsynaptic neurons of putative monosynaptic pairs recorded in LM-failure sessions. Presynaptic
neurons exhibited significantly larger landmark-locking scores than their postsynaptic targets only in area 29e (mean ± s.e.m.
= 0.19 ± 0.06; Holm-Bonferroni corrected one-sample t-test: $t(31) = 3.04$, $p = 0.02$). In other regions (MEC, PA, PO, and
PR), however, landmark-locking scores did not differ significantly between pre- and post-synaptic neurons of putative
monosynaptic connections (MEC: mean ± s.e.m. = –0.04 ± 0.03, $t(73) = -1.34$, $p = 0.724$; PA: 0.02 ± 0.04, $t(85) = 0.57$, $p =$
1.000; PO: –0.04 ± 0.04, $t(48) = 1.00$, $p = 0.957$; PR: –0.06 ± 0.12, $t(30) = 0.48$, $p = 1.000$). One-way ANOVA comparing
these landmark-locking score differences across regions showed a significant effect of region ($F(4,267) = 2.59$, $p = 0.038$).
Post-hoc t-tests with Holm-Bonferroni correction comparing 29e against each other region showed that 29e differed
significantly from MEC ($t(104) = 3.97$, $p < 0.001$) and PO ($t(79) = 3.18$, $p = 0.006$), but not PA ($t(116) = 1.93$, $p = 0.110$) or
PR ($t(61) = 1.96$, $p = 0.110$).

2683

2684

2685

2686

2687

2688

2689

2690

2691

2692

2693

2694

2695

2696

2697

2698

2699

2700

2701

2702

2703

2704

Supplementary Tables

2705

2706
2707
2708 **Table S1.** – Number of sessions and neurons recorded with 3 versus 4 visual landmarks during closed-loop gain
2709 manipulation experiments in the Dome. M = male; F = female. Results were qualitatively similar regardless of
2710 whether 3 or 4 landmarks were used.

		Number of sessions	Number of neurons
Rat 994 (M)	3 landmarks	4	58
	4 landmarks	3	62
Rat 995 (M)	3 landmarks	3	30
	4 landmarks	7	93
Rat 1041 (M)	3 landmarks	6	238
	4 landmarks	0	0
Rat 1061 (F)	3 landmarks	7	114
	4 landmarks	0	0
Rat 1062 (F)	3 landmarks	1	21
	4 landmarks	1	10

2711

2712

2713

2714

2715

2716

2717

2718

2719

2720

2721

2722

2723

Table S2. – Median values of Dome landmark selectivity metrics and landmark-locking scores in *LM-failure* sessions.

		Z index	Local similarity index		Landmark locking score
Rat 994 (M)	*29e* (*n*=2)	0.101	–0.169	*29e* (*n*=2)	0.425
	MEC (*n*=37)	0.356	–0.213	*MEC* (*n*=32)	0.062
	PA (*n*=44)	0.372	–0.187	*PA* (*n*=25)	0.063
	PO (*n*=35)	0.382	–0.221	*PO* (*n*=22)	0.055
	PR (*n*=3)	0.068	–0.201	*PR* (*n*=2)	0.355
Rat 995 (M)	*29e* (*n*=10)	1.319	0.412	*29e* (*n*=9)	0.767
	MEC (*n*=93)	0.345	–0.186	*MEC* (*n*=54)	0.280
	PA (*n*=12)	0.297	–0.238	*PA* (*n*=9)	0.204
	PO (*n*=3)	0.188	–0.240	*PO* (*n*=2)	0.297
	PR (*n*=6)	0.283	–0.149	*PR* (*n*=6)	0.148
Rat 1041 (M)	*29e* (*n*=19)	0.625	–0.025	*29e* (*n*=19)	0.610
	MEC (*n*=60)	0.441	–0.032	*MEC* (*n*=57)	0.296
	PA (*n*=110)	0.512	–0.235	*PA* (*n*=97)	0.253
	PO (*n*=49)	0.466	–0.232	*PO* (*n*=41)	0.197
	PR (*n*=0)	–	–	*PR* (*n*=0)	–
Rat 1061 (F)	*29e* (*n*=17)	0.443	–0.133	*29e* (*n*=3)	0.542
	MEC (*n*=67)	0.295	–0.284	*MEC* (*n*=33)	0.156
	PA (*n*=12)	0.280	–0.324	*PA* (*n*=7)	0.395
	PO (*n*=2)	0.831	–0.279	*PO* (*n*=1)	0.819
	PR (*n*=17)	0.329	–0.291	*PR* (*n*=10)	0.157
Rat 1062 (F)	*29e* (*n*=24)	0.613	0.047	*29e* (*n*=24)	0.292
	MEC (*n*=0)	–	–	*MEC* (*n*=0)	–
	PA (*n*=0)	–	–	*PA* (*n*=0)	–
	PO (*n*=0)	–	–	*PO* (*n*=0)	–
	PR (*n*=7)	0.496	–0.323	*PR* (*n*=7)	0.103

Table S3. – Median values of open-arena spatial tuning scores per animal.

		Skagg's SI score	Rayleigh score	EVF score
Rat 994 (M)	*29e* (*n*=7)	0.016	0.179	0.343
	MEC (*n*=148)	0.031	0.084	0.231
	PA (*n*=122)	0.047	0.121	0.198
	PO (*n*=45)	0.029	0.087	0.208
	PR (*n*=15)	0.043	0.363	0.160
Rat 995 (M)	*29e* (*n*=19)	0.029	0.058	0.435
	MEC (*n*=137)	0.036	0.051	0.185
	PA (*n*=16)	0.029	0.182	0.298
	PO (*n*=4)	0.045	0.041	0.378
	PR (*n*=14)	0.035	0.050	0.218
Rat 1041 (M)	*29e* (*n*=20)	0.035	0.491	0.516
	MEC (*n*=79)	0.045	0.041	0.136
	PA (*n*=133)	0.035	0.063	0.176
	PO (*n*=40)	0.024	0.112	0.298
	PR (*n*=0)	–	–	–
Rat 1061 (F)	*29e* (*n*=28)	0.012	0.117	0.238
	MEC (*n*=66)	0.023	0.107	0.265
	PA (*n*=23)	0.014	0.351	0.276
	PO (*n*=3)	0.028	0.303	0.381
	PR (*n*=52)	0.012	0.069	0.253
Rat 1062 (F)	*29e* (*n*=49)	0.017	0.225	0.324
	MEC (*n*=0)	–	–	–
	PA (*n*=0)	–	–	–
	PO (*n*=0)	–	–	–
	PR (*n*=10)	0.021	0.078	0.315

2747

Table S4. – Median values of single-cell physiological features.

		Theta rhythmicity index	Gamma rhythmicity index	Theta PPC
Rat 994 (M)	*29e* (*n*=8)	1.958	1.561	0.002
	MEC (*n*=190)	8.730	0.674	0.050
	PA (*n*=161)	5.983	0.529	0.104
	PO (*n*=58)	3.892	0.926	0.038
	PR (*n*=27)	8.783	0.322	0.022
Rat 995 (M)	*29e* (*n*=12)	1.168	0.840	0.0004
	MEC (*n*=166)	5.722	0.755	0.057
	PA (*n*=27)	5.259	0.529	0.071
	PO (*n*=5)	2.670	1.760	0.0006
	PR (*n*=18)	4.481	0.627	0.064
Rat 1041 (M)	*29e* (*n*=21)	2.649	0.741	0.0007
	MEC (*n*=89)	2.922	0.671	0.046
	PA (*n*=195)	9.755	0.418	0.146
	PO (*n*=67)	6.281	0.730	0.097
	PR (*n*=0)	–	–	–
Rat 1061 (F)	*29e* (*n*=22)	1.509	1.241	0.005
	MEC (*n*=121)	2.279	1.099	0.014
	PA (*n*=31)	2.309	0.555	0.023
	PO (*n*=3)	3.028	0.892	0.002
	PR (*n*=65)	1.897	1.048	0.006
Rat 1062 (F)	*29e* (*n*=60)	2.133	0.943	0.0015
	MEC (*n*=0)	–	–	–
	PA (*n*=0)	–	–	–
	PO (*n*=0)	–	–	–
	PR (*n*=14)	7.411	0.280	0.359

2748