Tick Genome Assemblies: Overcoming biological limitations through advances in

sequencing and assembly

Katie C. Dillon[*1], Julia C. Frederick[2], Hein Sprong[3], Travis C. Glenn[*1,2] and Isobel

Ronai[4-6]

[1]Institute of Bioinformatics, University of Georgia, Athens, GA, USA

[2]Department of Environmental Health Science, University of Georgia, Athens, GA, USA

[3]Centre for Infectious Diseases, National Institute of Public Health and Environment,

Bilthoven, The Netherlands

[4]University of Sydney, Sydney, NSW, Australia

[5]Department of Organismic and Evolutionary Biology, Harvard University, Cambridge,

MA, USA

[6]Howard Hughes Medical Institute, Chevy Chase, Maryland 20815, USA

[*]Corresponding authors

Further author information

Katie C. Dillon   katie.dillon@uga.edu / 0009-0005-7125-5793

Julia Frederick   julia.frederick@uga.edu / 0000-0003-4499-8688

Hein Sprong   hein.sprong@rivm.nl / 0000-0002-0218-4320

Travis Glenn   travisg@uga.edu / 0000-0001-7725-3637

Isobel Ronai   ironai@fas.harvard.edu / 0000-0002-1658-857X

RUNNING TITLE: Tick Genomes

ABSTRACT

Ticks are blood-feeding arthropods with approximately 1,000 species, however, only 24 species currently have a genome assembly. These genome assemblies are important resources to advance tick biology and control of tick-associated diseases. Generating tick genome assemblies is challenging due to their small body size (low DNA input), large genome size (approximately 2.4 Gbp for hard ticks), DNA contamination (from microbiota and host bloodmeals), and abundant transposable elements. Advances in sequencing technologies and assembly software have facilitated an increasing number of tick assemblies from 2011 to 2025. We characterize and assess the 54 tick genome assemblies within public genome databases using QUAST-LG and BUSCO compleasm. Then we evaluate the impact on these tick genome assemblies of biological source material, sequencing platform, and gene and repetitive element annotation. We identify 34 high-quality assemblies from 21 species that are well-suited for a variety of downstream analyses. Future tick genome assemblies should use long-read sequencing and Hi-C scaffolding to improve genomic resources for these unique blood-feeding parasites.

KEY WORDS: Ixodida, Ixodidae, Argasidae, vector biology, genetics, RNA-seq

THE IMPORTANCE OF TICKS

Ticks are hematophagous mites with a global health impact on humans, domesticated animals including livestock, and wildlife. As blood-feeders, ticks are disease vectors, transmitting a diverse range of disease agents, including viruses (for example, tick-borne *Flaviviridae*, *Dabie bandavirus*, and *Orthonairovirus*), bacteria (for example, *Borrelia* spp*., Anaplasma* spp. and Spotted Fever Group *Rickettsiae*), protozoa (for example, *Babesia* spp., *Theileria* spp.), and filarial nematodes[1-5]. The occurrence and distribution of these tick-associated diseases are increasing due to urbanization and climate changes[6,7]. Furthermore, ticks are developing resistance to current control strategies[8-11].

Ticks (order Ixodida) are comprised of three extant families: hard ticks (Ixodidae), soft ticks (Argasidae), and Nuttalliellidae. While there are approximately 1,000 valid tick species[12,13], few have genome assemblies available on databases such as the National Center for Biotechnology Information (NCBI). This lack of whole genome sequencing, which determines the complete DNA sequence for all chromosomes, is in part due to the unique biology of ticks.

Tick Biology Considerations for Genome Sequencing

The number of autosomes varies across tick species from 12 to 28 autosomes (diploid)[14-20] and the sex chromosome system is typically XY or XO[18]. Key aspects of tick biology should be taken into consideration prior to whole-genome sequencing and assembly.

Ticks have large genomes that are more similar in size to mammalian genomes than other vector species. We identified 17 genome assemblies of human disease vector species, spanning six taxonomic orders, from the World Health Organization's list of major human vector-borne diseases globally (https://www.who.int/news-room/fact-sheets/detail/vector-borne-diseases). For these 17 vector assemblies, the five largest all belong to ticks (Fig. 1). These tick assemblies have an average size that is up to 19 times larger than the other vector species, such as the human body louse (Fig. 1). The large genome-size of ticks has hindered the generation of genome assemblies, as this requires large amounts of sequencing and bioinformatic resources to achieve high

quality genomes with sufficient genome coverage (the number of times a single

nucleotide has been sequenced).

Tick genomes contain numerous highly repetitive regions, with transposable

elements (mobile DNA sequences that move around the genomes[21-23]) making up

almost 70% of some genomes[23]. This proportion is amongst the highest observed in

animals[12,24,25]. The high proportion of transposable elements in ticks increases

genome size and complexity, greatly complicating genome assembly.

The body size of a tick is relatively small for an animal; the largest life stage

(adult) and larger sex (female) is typically only a few millimeters when unengorged[26].

This small amount of tissue leads to a low total DNA yield (typically less than 6,000 ng)

from a single individual[23,27-29]. As tick genomes are large and highly repetitive,

relatively large amounts of high molecular weight DNA are required to meet optimal

sample (library) preparation minimums for genome sequencing that were originally

established for the generation of vertebrate genomes. Tick assemblies from individuals

initially needed to rely on amplification-based sample preparation[29]. In 2025, the

minimum amount of high molecular weight genomic DNA needed for non-amplification-

based sample preparation can be met by single-tick DNA extractions, such as Oxford

Nanopore 1,000 ng (https://store.nanoporetech.com/us/ligation-sequencing-kit-

v14.html), and PacBio Standard HiFi Sequencing 500 ng (https://www.pacb.com/revio).

Therefore, whole-genome sequencing of individual ticks has only become technically

feasible within the last few years[23].

DNA extractions from ticks are particularly high in contaminants, as they contain

DNA from their hosts and microbes[30]. Most tick species consume three blood meals

throughout their lifecycle, often from multiple host species. In addition, these

bloodmeals initiate the transfer of microbes from host vertebrate to tick[31].

Furthermore, tick species generally possess facultative and sometimes also obligatory

endosymbionts[32]. This contaminating DNA from other organisms, as well as

environmental contaminants means that genome sequencing of ticks generates many

reads of non-tick origin[23,33] (for example, ~14% total reads for an *I. ricinus*

assembly[23]). In summary, generating tick genome assemblies is challenging due to

their unique biology.

Survey of tick genome assemblies and quality assessment

We conducted a survey for tick whole-genome assemblies (keyword Ixodida) in the publicly available sequence databases: United States National Center for Biotechnology Information (NCBI); European Nucleotide Archive (ENA); and China National Center for Bioinformation (CNCB). A total of 70 tick genome assemblies: found on NCBI (65 assemblies) or ENA (one assembly), as of April 30, 2025; found on CNCB (one assembly), as of February 28, 2025; or came from in-house (three assemblies), which were not yet publicly available. We identify 16 assemblies (Table S1) on NCBI that have been mistakenly classified as tick genomes; they are endosymbiont genomes[34,35] or transcriptomes[36]. Therefore, in this study, we examine 54 genome assemblies, representing 24 species of tick (Table S2). Species designations were obtained from tick genome assembly publications, NCBI, and personal communications. We assessed quality metrics for these tick genome assemblies using QUAST-LG v.5.2.0[37] and BUSCO (benchmarking universal single-copy orthologs) lineage arthropoda_odb10, using compleasm v.0.2.5[38]. Here, we evaluated how biological source material, sequencing platforms, bioinformatic assembly methods, and gene and repetitive element annotation impact the quality of these tick genome assemblies.

BIOLOGICAL SOURCE MATERIAL FOR GENOME SEQUENCING OF TICKS

We compiled the metadata of the biological material for the 54 tick genome assemblies. For the zoogeographic origin of the tick material[39], the highest number of assemblies (~35%) are generated from ticks collected in the Nearctic zoogeographic region (North America), followed by ~31% Palearctic (Europe and China) and ~26% Afrotropical (Africa) (Fig. 2A, Table S3). We also note that some tick species are invasive and have been collected outside their native zoogeographic region and a genome assembly generated from this material, such as *Haemaphysalis longicornis* (see [40,41]). Currently, there are no genome assemblies from ticks collected in the Oriental region (South Asia and Southeast Asia) (Fig. 2A, Table S3), which has the second highest number of hard tick species, nor the Neotropical region (Central America and South America) (Fig. 2A, Table S3) which has the fourth highest number of hard tick

species[39]. We recommend the Neotropical and Oriental zoogeographic regions be
prioritized in future tick genome projects.

For the collection source of the tick material, the highest number of assemblies
are generated from tick colonies (Fig. 2B, Table S3). These ticks from colonies were
sourced ~24% from the Deutch colonies[42] followed by ~14% Oklahoma State
University colonies[29,43,44] (Fig. 2B, Table S3). The second highest number of
assemblies are from ticks caught in the field (Fig. 2B, Table S3). The third highest
number of tick assemblies are from laboratory reared ticks that were caught in the field
and maintained in the laboratory until a desired life stage had been obtained, and the
smallest number of assemblies are from tick cell lines (Fig. 2B, Table S3). These cell
lines are typically created from heterogenous embryonic cells and maintained
indefinitely in a laboratory, which leads to large-scale genome duplications,
rearrangements, and deletions[45]. We recommend the biological source material for
assemblies be obtained from laboratory colonies, laboratory reared, or field caught.

Tick material has varied in sample size for assemblies, with ~56% individual ticks
and ~44% of pooled ticks (Fig 2C, Table S3). For assemblies generated from
individuals, all are adult ticks, apart from three nymphs (Fig. 2D, Table S3). For
assemblies generated from pooled tick material, the highest number of assemblies
came from eggs (~46%) (Fig. 2D, Table S3). For the assemblies generated from
individuals, ~78% were from females (Fig. 2E, Table S3), most likely because they
generally contain larger amounts of DNA than males[46]. Until 2020, pooling DNA from
multiple individuals (Table S2-S3) was needed exclusively to achieve the minimum
amount of DNA required for sequencing. Many of the most recent assemblies use DNA
from a single individual to generate the data used for primary (contig) assembly, then
use pooled DNA to generate sequence data for high-throughput chromosome
conformation capture (Hi-C) assembly scaffolding[29]. Pooling individual ticks or eggs
provides sufficient DNA yield for a variety of sample preparation and sequencing
approaches, however, the high amount of heterozygosity among individuals can lead to
significant difficulty in producing highly contiguous assemblies[21]. We recommend
prioritizing tick assemblies generated from single, individual ticks, which can achieve a
high-quality haploid assembly.

TICK WGS AND ASSEMBLY DEVELOPMENTS FROM 2011 – 2025

We traced the history of the 54 tick genome assemblies. The first tick assembly, JCVI_ISG_i3_1.0 for *I. scapularis*, was released in 2011 (Fig. 3, Table S2). This assembly project[33] resulted from a 2005 community white paper, with this species being prioritized due to its medical importance in the United States as the key species associated with Lyme disease[47]. The JCVI_ISG_i3_1.0 assembly is the only tick assembly to use Sanger dideoxy chain termination on capillary instruments (first-generation sequencing)[48,49]. This method costs millions of dollars in reagents per tick genome, due to the large genome size[48]. Use of pooled individuals from an inbred colony, combined with modest sequence read lengths[50] was unable to produce a highly contiguous assembly due to the highly repetitive and heterozygous tick genomes.

The next tick genome assemblies used 454-pyrosequencing (second generation sequencing), CCG_Rmi1.0[50] and Rmi2.0[51] *Rhipicephalus microplus*. This sequencing platform obtains millions of ~500 nt reads at a time, thus reducing costs, but has similar limitations in handling repetitive and heterozygous assemblies as Sanger sequencing[52]. By 2016, published tick assemblies were using the Illumina (second-generation) short-read sequencing platform (Fig. 3, Table S2). Illumina reads are 50-600 nt in total, most commonly paired-end 100 or 150 nt reads (i.e., 200 or 300 nt total) and sequencing is massively paralleled for high throughput, providing higher genome coverage[52]. As Illumina uses short reads, it cannot be used to reliably map or assemble long repetitive regions present in tick genomes.

After 2017, tick genomes were assembled using the data from the Pacific Biosciences (PacBio) (third-generation sequencing) sequencing platform (Fig. 3, Table S2). PacBio uses specialized library preparation (double-stranded DNA is circularized via SMRTbell adapters) and single-molecule sequencing to create long-read reads (up to ~100 kb for single-pass CLR reads or modal length of ~20 kb when sequenced repeatedly for HiFi reads) to handle the high proportion of repetitive elements in ticks. Initially, PacBio CLR reads were used to create tick assemblies with much better contiguity[53] and Illumina reads were often used to correct uncalled or mis-called bases (Fig. 3, Table S2). Now the standard output for PacBio is HiFi reads, which have

greatly improved base-calling accuracy (trading-off reads >25kb) for tick

assemblies[29,54].

From 2020 to 2023, 22 tick assemblies were released and more than half used

the PacBio sequencing platform (Fig. 3, Table S2). In 2023, the Oxford Nanopore

(another third-generation sequencing) sequencing platform was first used for tick

genome sequencing (Fig. 3, Table S2). This platform uses long-reads (reads 50 bp to >

4 Mbp) generated in real time via flow cells with nanopores[55]. While Oxford Nanopore

base-calling accuracy has greatly improved, ~33% of Nanopore tick assemblies have

been generated in combination with Illumina short-reads to correct base-calling errors

(Fig. 3, Table S2). In 2024 the most genome assemblies were released (20 assemblies)

with ~40% using Nanopore and the remaining using combinations of Nanopore or

PacBio, and Illumina (Fig. 3, Table S2). By April 2025, six assemblies were released,

with four using Nanopore (Fig. 3). The accuracy of Oxford Nanopore reads has now

improved sufficiently that Illumina reads are often no longer necessary for error

correction[56] or are only needed in scenarios where near perfect sequence accuracy is

required[57].

The development of new generations of sequencing platforms has been critical

for overcoming the limitations of tick biology (large genome size and high proportion of

repeats) leading to an increasing number of tick genome assemblies. Standalone first-

generation sequencing generated ~4% of tick assemblies, second-generation

sequencing ~20%, and third-generation sequencing ~39% (Fig. 3, Table S2). By 2018,

first generation sequencing was no longer being used to generate tick assemblies, while

third-generation sequencing was first used for ticks in 2017 (Fig. 3, Table S2). In

addition, ~37% of tick genome assemblies have been generated using combinations of

different generation sequencing (Fig. 3, Table S2).

For the 54 tick assemblies, ~37% used one or more integrations (Hi-C or RNA-

sequencing) to the sequencing platform. Hi-C is a molecular approach[58,59], which

improves contiguity of an assembly and is useful for ticks with their large, repetitive,

heterozygous genomes. Alternatively, RNA-seq provides direct evidence of

transcription, as well as the sequence of genes and isoforms[60]. Of the 20 assemblies

using integrations, 55% used a combination of Hi-C and RNA-sequencing, 30% RNA-

sequencing alone, and 15% Hi-C alone (Fig. 3, Table S2). Hi-C was first used to

generate a tick assembly in mid-2020, since then, ~26% of assemblies have integrated

Hi-C (Fig. 3, Table S2).

MAJOR CHALLENGES IN STANDARDIZATION OF ANNOTATION FOR TICK
236

GENOME ASSEMBLIES

Annotation of genes and repetitive elements (in particular, the transposable elements) in

tick genome assemblies has not been standardized. These non-standardized

annotations currently limit our ability to compare the genes and repetitive elements

between tick assemblies, even for the same species.

While genome assemblies commonly include gene annotation, tick genome

assemblies have not had a standardized approach. In particular, some assemblies

report the number of protein-coding genes and others the total number of genes (Table

S2). Gene annotation for tick genome assemblies has four common approaches. First,

custom gene annotation pipelines individually created for each assembly. Second,

taking a custom pipeline annotation and aligning RNA-seq data (either pre-existing or

generated alongside the genome assembly) using gene modeling programs (such as

AUGUSTUS[61] and BRAKER[61,62]). Third, manual pipeline including aligning RNA-

seq and proteome data. Fourth, requesting NCBI to perform a RefSeq annotation,

running the NCBI Eukaryotic Genome Annotation Pipeline[63], which incorporates

multiple data types (including RNA-seq and proteome) from within NCBI.

Gene annotation has been performed for 30 tick assemblies, representing 18

species (Figure S1, Table S2). A custom gene annotation pipeline was used for seven

tick assemblies, with a median of ~27,100 total genes and ~26,400 protein-coding

genes (Figure S1, Table S2). Also, a custom gene annotation pipeline with RNAseq

data has been used for 17 tick assemblies with a median of ~28,500 total genes and

~27,000 protein-coding genes (Figure S1, Table S2). A manual pipeline was used for

four tick assemblies, with a median of ~23,700 protein-coding genes (Figure S1, Table

S2). The NCBI Pipeline has been used for seven tick assemblies with a median of

~27,000 total genes and ~26,900 protein-coding genes (Figure S1, Table S2). While

four alternative approaches have been used to annotate the genes in tick assemblies

and resulted in variable gene number estimates, the median number for total genes (15

assemblies) is between ~27,000 to ~28,500 and protein-coding genes (24 assemblies)

is between ~23,700 to ~27,000. We recommend future studies that generate tick

genome assemblies achieve high quality gene annotation by including a transcriptome,

ideally via PacBio Iso-Seq, and submitting their assembly to NCBI for annotation.

The highly repetitive genomes of ticks mean that these repetitive elements

should also be annotated. The common approach to annotating repetitive elements is

using an automated pipeline, such as one or a combination of RepeatModeler[64] and

RepeatMasker[65]. Repetitive elements have been annotated in 25 tick genome

assemblies, representing 15 species (Figure S2, Table S2). An automated pipeline for

repetitive elements has been performed on 23 tick assemblies with a median of ~62%

repetitive DNA, composed of ~1.5% tandem repeats and ~59% transposable elements

(Figure S2A, Table S2). When these transposable elements have been further

classified, the median is ~24% class I, ~6% class II, and ~2.4% unclassified (Figure

S2B, Table S2). A manual pipeline for repetitive elements has been performed on three

Ixodes assemblies from one study[23] with a median of 73% repetitive DNA, composed

of ~4.3% tandem repeats and ~68% transposable elements (Figure S2A, Table S2).

Further classification gave a median ~32% class I and ~36% class II (Figure S2B, Table

S2). We recommend future tick genome assemblies perform manual curation of the

transposable elements until a standardized, automated pipeline is established.

QUALITY ASSESSMENT OF TICK GENOME ASSEMBLIES

We comprehensively assessed the impact of sequencing platform and Hi-C scaffolding

on tick assembly quality, using metrics for contiguity (auN, N50, L50, and L90) and

BUSCO complete scores (a higher BUSCO score indicates higher assembly

completeness), which are common assessments of genome assembly quality[66]. The

complete BUSCO score is the sum of the single copy and duplicated genes that can be

completely aligned to the assembly with one or more than one gene copy present,

respectively. Sanger and 454-FLX sequencers were used for the first two tick

assemblies (Table S2), generating highly fragmented assemblies (Fig. 4A-D, Table S4).

While the complete BUSCO score for the Sanger assembly was close to 90%, the 454-
FLX partial assembly was close to 0% (Fig. 4A-D, Table S5).

Illumina sequencing has been used for 12 tick assemblies (Table S2). Only one
Illumina assembly integrated Hi-C scaffolding, *I. ricinus* (IXRI_v2)[27], which generated
the highest quality Illumina tick assembly with a ~95% complete BUSCO score (Fig. 4A-
D, Table S4-S5). The other 11 Illumina assemblies are more fragmented and seven did
not produce sufficient reads to cover the genome, resulting in lower complete BUSCO
scores of 0 to ~65% (Fig. 4A-D, Table S4-S5). Therefore, standalone short read
platforms that are of insufficient depth and do not integrate Hi-C scaffolding are not
sufficient to handle the genome complexity of ticks.

PacBio sequencing has been used for eight tick assemblies, two of which
implemented Hi-C scaffolding (Table S2). Interestingly, one of the PacBio assemblies
with Hi-C had a similar quality to the Illumina assembly that integrated Hi-C (Fig. 4A-D,
Table S4-S5). The remaining six PacBio assemblies without Hi-C varied in quality (Fig.
4A-D, Table S4-S5).

A combination of PacBio sequencing error-corrected with Illumina short-reads
has been used for 13 tick assemblies (Table S2). Twelve of these assemblies have
complete BUSCO scores over 92%, indicating sufficient sequencing coverage of the
genome (Fig. 4A-D, Table S4-S5). Ten of these 13 assemblies implemented Hi-C
scaffolding and the three without Hi-C were lower in quality regarding contiguity and
coverage (Fig. 4A-D, Table S4-S5). Our findings indicate that Hi-C positively impacts
assembly contiguity and quality for PacBio tick assemblies with or without additional
Illumina reads.

Nanopore sequencing has been used for 12 tick assemblies (Table S2). These
assemblies overlap in quality with the Illumina, PacBio, and PacBio+Illumina tick
assemblies (Fig. 4A-D, Table S4-S5). None of the standalone Nanopore assemblies
implemented Hi-C (Fig. 4A-D). The quality and contiguity of Nanopore assemblies is
moderate when compared to other sequencing platforms.

A combination of Nanopore error-corrected with Illumina short-reads has been
used for six tick assemblies (Table S2). For all Nanopore+Illumina assemblies, the
complete BUSCO values were higher than 83%, which is unexpectedly lower than the

standalone Nanopore assemblies, which were ~94% (Fig. 4A-D, Table S4-S5). One of

these six assemblies integrated Hi-C scaffolding, which had higher quality and

contiguity compared to Nanopore+Illumina tick assemblies without Hi-C (Fig. 4A-D,

Table S4-S5). The quality of the remaining five Nanopore+Illumina assemblies without

Hi-C were similar in contiguity, but lower in quality in regard to BUSCO complete scores

compared to the standalone Nanopore tick assemblies (Fig. 4A-D, Table S4-S5).

Therefore, the highest quality Nanopore assemblies were generated without error-

correction using Illumina reads, with exception of the Nanopore+Illumina assembly that

integrated Hi-C.

Across all the sequencing platforms, on average, assemblies integrating Hi-C

data improved quality metrics and complete BUSCO scores, resulting in assemblies

with higher contiguity, quality, and completeness (Fig. 4A-D, Table S4-S5). Additionally,

assemblies generated by a combination of long-read platforms (PacBio and Nanopore)

then error-corrected with Illumina reads had, on average, better contiguity but lower

genome completeness compared to those generated by long-read platforms without

error-correction, suggesting a trade-off in contiguity versus errors (Fig. 4A-D, Table S4-

S5). In summary, we recommend future tick genome assembly projects use long-read

sequencing platforms with Hi-C integration to produce high-quality assemblies.

CHARACTERIZATION OF HIGH-QUALITY TICK ASSEMBLIES

We assessed the quality of the tick genome assemblies using number of contigs,

contig N50, and BUSCO complete score to identify 34 high quality tick genome

assemblies that are suitable for downstream analyses (Fig. 5). These high-quality

assemblies have fewer than 100,000 contigs, a contig N50 greater than 45 Kbp, and a

complete BUSCO score higher than 83% (Fig. 5, Table S4-S6). These 34 high-quality

assemblies were all generated since 2020 (Fig. 3) and used ~15% Illumina, ~18%

Nanopore, ~18% Nanopore+Illumina, ~21% PacBio, and ~29% PacBio+Illumina (Fig. 5,

Table S4-S6). Approximately 41% of the high-quality assemblies integrated Hi-C (Fig. 5,

Table S4-S6).

We also identify a cluster of 14 highest quality tick genome assemblies which

have fewer than 7,500 contigs, a contig N50 greater than 84 Mbp, and a complete

BUSCO score greater than 93% (Fig. 5, Table S4-S6). These 14 highest-quality

assemblies were all generated since 2020 (Fig. 3) and ~7% used Illumina, ~7%

Nanopore+Illumina, ~21% PacBio, and ~64% PacBio+Illumina (Fig. 5, Table S4-S6). All

14 assemblies integrated Hi-C, except for *Rhipicephalus microplus* (USDA_Rmic)[15]

(Fig. 5). Our use of comprehensive metrics for the highest-quality tick assemblies

means future tick assembly projects now have metrics to benchmark against.

TICK SPECIES WITH HIGH QUALITY GENOME ASSEMBLIES

The 34 high-quality genome assemblies cover 21 species of tick (Fig. 6). As some

species have multiple high-quality assemblies, we identify the best quality genome

assembly for each of the 21 tick species (Fig. 6, Table S6).

For the 21 best quality assemblies, we find the hard ticks have an average

assembly length of ~2,390 Mbp and soft ticks ~1,176 Mbp (Fig. 6, Table S4). The hard

tick *Ixodes* spp. (Prostriata) have an average assembly length of ~2,292 Mbp and the

non-*Ixodes* spp. (Metastriata) ~2,425 Mbp (Fig. 6, Table S4). Across all tick species

Rhipicephalus spp. have the largest assembly length, with an average of ~2,689 Mbp,

whereas the two *Hyalomma* spp. have the smallest assembly lengths for the hard ticks

(average ~1,840 Mbp) (Fig. 6, Table S4). The two soft tick *Ornithodoros* spp. have the

smallest assembly lengths for all ticks with an average of ~1,176 Mbp (Fig. 6, Table S4).

Assembly size (length of gapped DNA sequence) is not equivalent to genome size. The

average genome size for soft ticks is 1.3 Gbp and for hard ticks is 2.7 Gbp using flow

cytometry[67]. The average percent difference of the estimated genome size to both the

NCBI assembly size and assembly total length is 95.5% (Table S7). We hypothesize

that hard ticks have larger genomes compared to soft ticks due to larger proportions of

repetitive elements.

The 21 species with high-quality assemblies are two soft ticks (Argasidae) and

19 hard ticks (Ixodidae), or five Ixodes and 14 non-Ixodes species (Table S7). This

makes up 0.9% (2/216) of the known soft tick species and 2.4% (19/786) of the known

hard tick species[12,13]. We recommend that generating genome assemblies for

additional soft ticks should be a priority. So far, tick species sequenced are those of

medical and veterinary importance[47]. Therefore, we also recommend future tick

genome projects prioritize capturing the diversity of tick species, particularly those of

medical, veterinary, and phylogenetic importance[68,69].

CONCLUDING REMARKS

We have assessed 54 genome assemblies for ticks that have been generated,

identifying 34 high-quality assemblies for 21 tick species. Our assessment of the tick

genome assemblies has provided an opportunity to make recommendations for future

tick genome assembly projects. We recommend future assemblies better represent the

biodiversity of tick species, particularly the soft ticks and from unrepresented

zoogeographic regions. In addition, to obtain the highest quality tick genome

assemblies, future projects should prioritize obtaining individual, adult ticks for DNA

extractions and for sequencing platforms use a combination of PacBio HiFi with Hi-C

Illumina scaffolding (or Oxford Nanopore with Hi-C scaffolding) with transcriptomes.

High-quality annotations should include transcriptomes from as many life-stages as

possible, plus TE manual curation. With these recommendations, future tick genome

assemblies will improve in quality and representation of tick species.

With the availability of high-quality tick genome assemblies, there are

opportunities to move beyond sequencing a single individual per species to better

understand tick biology. Genomic resources that are well established in other non-

model arthropods but are needed in ticks include a chromosome-anchored reference

genome using fluorescent *in situ* hybridization (FISH)[70] , sequencing multiple

individuals per species to increase phylogeographic representation[71], and cross-

species whole genome alignments[71]. New approaches to make even more advances

include new sequencing technologies, surveys for transposase-accessible chromatin

sequencing (ATAC-seq)[72] to characterize regulatory elements in the genome, and

telomere-to-telomere (T2T) gapless sequencing[73] to understand tick chromosome

structures. These additional approaches will greatly improve our understanding of tick

genomes and tick biology.

Over the past decade, major developments in sequencing technologies have

brought about a renaissance for tick genomics. These genome assemblies are critical

for advancing tick research from vectorial capacity[74] to anti-tick vaccines[75]. The tick

genome assemblies have so far been referred to or used in more than 130 follow up

studies (Table S8) since 2011. These follow up studies have four key themes:

identifying gene targets for anti-tick vaccination studies, tick control, and acaracide

resistance; annotation of repetitive elements; intra-species and inter-species

comparative genomics studies; transcriptomics and proteomics studies. Therefore, the

tick genome assemblies have enabled major advances in our understanding of tick

biology.

We look forward to new tick genome assemblies providing more insights into the

fascinating biology of these medically and veterinary important blood-feeding parasites.

Ethics approval

None.

Availability of data and materials

All supporting datasets for this study are available in the supplementary material.

Funding

KCD and TCG have been supported in part with federal funds from the Centers of

Disease Control and Prevention Pathogen Genomics Center of Excellence (Contract

No. NU50CK000626). JCF was partially supported by Grant No. DGE-1545433 from the

National Science Foundation and the IDEAS programme at the University of Georgia.

HS was financially supported by the Dutch Ministry of Health, Welfare and Sport (VWS).

IR is a Howard Hughes Medical Institute Awardee of the Life Sciences Research

Foundation.

Acknowledgements

We thank all the researchers who have pushed the boundaries of tick genomics and

shared with us additional information. In particular, we thank the following researchers

for their willingness and transparency to provide information on unpublished tick

genome assemblies that they have released on genome databases, in alphabetical

order: Joshua Benoit, Seemay Chou, Christopher Faulk, Jan Forth, Melissa Klein,

Michel Labuschagne, Susan Noh, Pia Olafson, Perot Saelao, Ryan O. M. Rego, Claude

Rispe, Hailey Schau, and Pavlina Vechtova. We also acknowledge Roslyn Hickson for

facilitating the resources that made the *O. gurneyi* genome assembly possible. The use

of the unpublished genome assemblies in this paper does not constitute their

announcement, as details related to preparation, assembly and gene annotation

processes have yet to be described.

Author contributions

Conceptualization: KCD, IR, TCG, JF

Validation: KCD

Formal analysis: KCD

Investigation: JF, KCD

Resources: HS

Data Curation: KCD

Writing - Original Draft: KCD, IR

Writing - Review & Editing: KCD, IR, TCG, HS, JF

Visualization: KCD, IR

Supervision: IR, TCG

Project administration: IR, TCG

Funding acquisition: HS, IR, TCG

Declaration of interests

470 None declared.

FIGURES

Figure 1. Ticks (Ixodida) have the largest genome assembly sizes when compared with other major human disease vectors. Genome assembly size for 17 disease vectors, spanning six taxonomic orders, from the major human vector-borne diseases global list (https://www.who.int/news-room/fact-sheets/detail/vector-borne-diseases). For human vector-borne diseases associated with multiple vector species, the species associated with the highest number of disease cases and has an available assembly is shown. When a vector species has multiple assemblies, the current reference assembly from NCBI is used. For each assembly, the NCBI assembly size is reported.

Figure 2. Biological source material for 54 tick genome assemblies. A) Geographic

origin of the tick material. Two assemblies were generated using ticks from three

locations. **B)** Biological sources of the tick material. A laboratory colony is defined as

ticks being maintained across generations in a lab environment. We define laboratory

reared ticks as those collected from the field and maintained in the laboratory until a

desired life stage has been obtained, not across generations. Field caught ticks are

collected from either the environment or hosts. Tick cell lines are preserved cells,

typically from embryos. **C)** Sample size of the tick material, either is an individual

specimen or pooled specimens. **D)** Life stage of the tick in individual versus pooled

material. **E)** The sex of the adult ticks in individual versus pooled material. See Table S3

for further details.

Figure 3. Timeline of assembly release for 54 tick genome assemblies, first

release was in 2011. Each point is the release date on a genome database of a

genome project; a project has included up to five tick assemblies. Sequencing platforms

are in chronological order, with first generation (Sanger or 454-FLX), second generation

(Illumina) and third generation (PacBio or Nanopore). Followed by combinations of

sequencing platforms. Integrations to the sequencing platforms were none, RNA-seq

alone, Hi-C alone or Hi-C with RNA-seq. See Table S2 for further details.

Figure 4. Quality metrics for 54 tick genome assemblies across sequencing platforms. A) auN (comprehensive metric for contiguity), as auN increases so does assembly quality. **B)** Contig N50 (the length of the shortest contig that must be included

to cover half an assembly), as N50 increases, fewer longer contigs are used to cover half of the assembly, leading to a higher quality assembly. **C)** L50 (minimal number of contigs that cover half of the assembly), as L50 decreases assembly quality can increase, however, the BUSCO score needs to be included in the assessment. **D)** L90 (minimal number of contigs that cover 90% of the assembly), as L90 decreases assembly quality can increase, however the BUSCO score needs to be included in the assessment. Each point is a genome assembly. Sequencing platforms are ordered by first generation (Sanger and 454-FLX), second generation (Illumina) and third generation (PacBio and Nanopore). Followed by sequencing platform combinations. Scaffolding for the sequencing platforms were none or Hi-C. See Table S2, S4, and S5 for further details.

Figure 5. Quality assessment of 54 tick genome assemblies. Identified 34 high-

quality assemblies (lower contig number, higher N50, and higher BUSCO score),

including 21 highest quality assemblies. Each dot is a genome assembly. Contig N50 is

the length of the shortest contig that must be included to cover half an assembly, a

higher N50 means less small contigs which is a more complete, less fragmented

assembly. For the number of contigs, fewer contigs typically result in a more contiguous

assembly as there is less fragmentation. However, few contigs associated with a low

N50 and small assembly length can lead to poor assembly quality. As not all assemblies

were obtained from NCBI, we used a metric that could be calculated for all 54

assemblies: the total length of the assembly (generated by QUAST) which includes

uncalled bases. However, QUAST total length more closely represents the assembly

size, not genome size. BUSCO complete score, a higher BUSCO score indicates higher

assembly completeness and a higher quality assembly. Estimated haploid assembly

size from NCBI. See Table S4-S6 for further details.

Figure 6. Assembly total length of the best quality genome assembly for 21 tick species. The best assembly for each tick species was identified from the high and highest quality assemblies. Assembly length (Mbp) of the best quality assembly for each species. Tick species are ordered by assembly total length, from smallest to largest. See Table S6 for further details.

References Cited

1. Madison-Antenucci, S. *et al.* (2020) Emerging Tick-Borne Diseases. *Clinical Microbiology Reviews* 33, 10.1128/cmr.00083–00018. https://doi.org/10.1128/cmr.00083-18

2. Parola, P. *et al.* (2013) Update on tick-borne rickettsioses around the world: a geographic approach. *Clinical microbiology reviews* 26, 657–702. https://doi.org/10.1128/cmr.00032-13

3. Kim, E.-H. and Park, S.-J. (2023) Emerging tick-borne *Dabie bandavirus*: virology, epidemiology, and prevention. *Microorganisms* 11, 2309. https://doi.org/10.3390/microorganisms11092309

4. Bishop, R. *et al.* (2004) *Theileria*: intracellular protozoan parasites of wild and domestic ruminants transmitted by ixodid ticks. *Parasitology* 129, S271–S283. https://doi.org/10.1017/S0031182003004748

5. Ajileye, O.D. *et al.* (2025) A review of filarial nematodes parasitizing tick vectors: unraveling global patterns in species diversity, host associations, and interactions with tick-borne pathogens. *Parasites & Vectors* 18, 50. https://doi.org/10.1186/s13071-025-06690-6

6. Kilpatrick, A.M. and Randolph, S.E. (2012) Drivers, dynamics, and control of emerging vector-borne zoonotic diseases. *The Lancet* 380, 1946–1955. https://doi.org/10.1016/S0140-6736(12)61151-9

7. Saker, L. *et al.* (2004) Globalization and infectious diseases, a review of the linkages. *UNDP/World Bank/WHO Special Programme on Tropical Diseases Research, Geneva.* https://researchonline.lshtm.ac.uk/id/eprint/15642

8. Burtis, J.C. *et al.* (2021) Susceptibility of *Ixodes scapularis* (Acari: Ixodidae) to Permethrin Under a Long-Term 4-Poster Deer Treatment Area on Shelter Island, NY. *Journal of Medical Entomology* 58, 1966–1969. https://doi.org/10.1093/jme/tjab054

9. Vudriko, P. *et al.* (2016) Emergence of multi-acaricide resistant *Rhipicephalus* ticks and its implication on chemical tick control in Uganda. *Parasites & Vectors* 9, 4. https://doi.org/10.1186/s13071-015-1278-3

10. Ziapour, S. *et al.* (2016) Susceptibility status of field populations of *Rhipicephalus bursa* (Acari: Ixodidae) to pyrethroid insecticides. *Tropical Biomedicine* 33, 446–461

11. Lempereur, L. *et al.* (2025) Research-led innovation for integrated tick management and acaricide resistance mitigation in livestock. *Trends in Parasitology* 0. https://doi.org/10.1016/j.pt.2025.09.002

12. Mans, B.J. *et al.* (2019) Argasid and ixodid systematics: implications for soft tick evolution and systematics, with a new argasid species list. *Ticks and Tick-borne Diseases* 10, 219–240. https://doi.org/10.1016/j.ttbdis.2018.09.010

13. Robbins, R.G. *et al.* (2025) Type specimens of the world's hard tick species (Acari: Ixodida: Ixodidae): collection data and depositories for all valid names and the current status of invalid names. *Zootaxa* 5606, 1–287. https://doi.org/10.11646/zootaxa.5606.1.1

14. Wysoki, M. and Bolland, H.R. (1978) Spermatogenesis, chromosomes and sex determination of four *Rhipicephalus* species (Acari: Ixodidae) from East Africa. *Genetica* 48, 233–238. https://doi.org/10.1007/BF00155576

15. Tidwell, J.P. *et al.* (2024) Identifying the sex chromosome and sex determination genes in the cattle tick, *Rhipicephalus (Boophilus) microplus. G3 Genes|Genomes|Genetics* 14. https://doi.org/10.1093/g3journal/jkae234

16. Gunn, S.J. and Hilburn, L.R. (1990) Cytosystematics of Five North American *Dermacentor* (Acari: Ixodidae) Species. *Journal of Medical Entomology* 27, 620–627. https://doi.org/10.1093/jmedent/27.4.620

17. Yilmazer, N. (2018) The genome size of some species of *Hyalomma* ticks from Turkish Thrace. *Systematic and Applied Acarology* 23, 2122–2128. https://doi.org/10.11158/saa.23.11.5

18. Oliver, J.H. (1977) Cytogenetics of Mites and Ticks. *Annual Review of Entomology* 22, 407–429. https://doi.org/10.1146/annurev.en.22.010177.002203

19. Kahn, J. (1964) Cytotaxonomy of ticks. *Journal of Cell Science* 3, 123–137. https://doi.org/10.1242/jcs.s3-105.69.123

20. Gunn, S.J. and Hilburn, L.R. (1989) Differential Staining of Tick Chromosomes: Techniques for C-Banding and Silver-Staining and Karyology of *Rhipicephalus sanguineus* (Latreille). *The Journal of Parasitology* 75, 239–245. https://doi.org/10.2307/3282772

21. Petersen, M. *et al.* (2019) Diversity and evolution of the transposable element repertoire in arthropods with particular reference to insects. *BMC Ecology and Evolution* 19, 11. https://doi.org/10.1186/s12862-018-1324-9

22. Wu, C. and Lu, J. (2019) Diversification of transposable elements in arthropods and its impact on genome evolution. *Genes* 10, 338. https://doi.org/10.3390/genes10050338

23. Ronai, I. *et al.* (2024) The repetitive genome of the *Ixodes ricinus* tick reveals transposable elements have driven genome evolution in ticks. *bioRxiv*, 2024.2003.2013.584159. https://doi.org/10.1101/2024.03.13.584159

24. Osmanski, A.B. *et al.* (2023) Insights into mammalian TE diversity through the curation of 248 genome assemblies. *Science* 380, eabn1430. https://doi.org/10.1126/science.abn1430

25. Shao, C. *et al.* (2023) The enormous repetitive Antarctic krill genome reveals environmental adaptations and population insights. *Cell* 186, 1279–1294. e1219. https://doi.org/10.1016/j.cell.2023.02.005

26. Keirans, J.E. and Durden, L.A. (2005) Tick systematics and identification. *Tick-borne Diseases of Humans*, 123–140. https://doi.org/10.1128/9781555816490.ch7

27. Cerqueira de Araujo, A. *et al.* (2025) Genome sequences of four *Ixodes* species expands understanding of tick evolution. *BMC biology* 23, 17. https://doi.org/10.1186/s12915-025-02121-1

28. Ribeiro, J.M. *et al.* (2023) A draft of the genome of the Gulf Coast tick, *Amblyomma maculatum. Ticks and tick-borne diseases* 14, 102090. https://doi.org/10.1016/j.ttbdis.2022.102090

29. De, S. *et al.* (2023) A high-quality *Ixodes scapularis* genome advances tick science. *Nature Genetics* 55, 301–311. https://doi.org/10.1038/s41588-022-01275-w

30. Díaz-Sánchez, S. *et al.* (2019) Evolutionary insights into the tick hologenome. *Trends in Parasitology* 35, 725–737. https://doi.org/10.1016/j.pt.2019.06.014

31. Anderson, J.F. and Magnarelli, L.A. (2008) Biology of ticks. *Infectious disease clinics of North America* 22, 195–215. https://doi.org/10.1016/j.idc.2007.12.006

32. Kolo, A.O. and Raghavan, R. (2023) Impact of endosymbionts on tick physiology and fitness. *Parasitology* 150, 859–865. https://doi.org/10.1017/S0031182023000793

33. Gulia-Nuss, M. *et al.* (2016) Genomic insights into the *Ixodes scapularis* tick vector of Lyme disease. *Nature Communications* 7. https://doi.org/10.1038/ncomms10507

34. Buysse, M. *et al.* (2021) A dual endosymbiosis supports nutritional adaptation to hematophagy in the invasive tick Hyalomma marginatum. *elife* 10, e72747. https://doi.org/10.7554/eLife.72747

35. Floriano, A.M. *et al.* (2025) Comparative genomics of *Rickettsiella* bacteria reveal variable metabolic pathways potentially involved in symbiotic interactions with arthropods. *Peer Community Journal* 5. https://doi.org/10.24072/pcjournal.633

36. Forth, J.H. *et al.* (2020) Identification of African swine fever virus-like elements in the soft tick genome provides insights into the virus' evolution. *BMC biology* 18, 136. https://doi.org/10.1186/s12915-020-00865-6

37. Mikheenko, A. *et al.* (2018) Versatile genome assembly evaluation with QUAST-LG. *Bioinformatics* 34, i142–i150. https://doi.org/10.1093/bioinformatics/bty266

38. Huang, N. and Li, H. (2023) compleasm: a faster and more accurate reimplementation of BUSCO. *Bioinformatics* 39, btad595. https://doi.org/10.1093/bioinformatics/btad595

39. Guglielmone, A.A. *et al.* (2023) Geographic distribution of the hard ticks (Acari: Ixodida: Ixodidae) of the world by countries and territories. *Zootaxa* 5251, 1–274. https://doi.org/10.11646/zootaxa.5251.1.1

40. Guerrero, F.D. *et al.* (2019) The Pacific Biosciences de novo assembled genome dataset from a parthenogenetic New Zealand wild population of the longhorned tick, *Haemaphysalis longicornis* Neumann, 1901. *Data in Brief* 27, 104602. https://doi.org/10.1016/j.dib.2019.104602

41. Moustafa, M.A.M. *et al.* (2025) Genome of the invasive North American *Haemaphysalis longicornis* tick as a template for bovine anti-tick vaccine discovery. *BMC Genomics* 26, 307. https://doi.org/10.1186/s12864-025-11477-1

42. Davey, R. *et al.* (1980) Ovipositional biology of the cattle tick, *Boophilus annulatus* (Acari: Ixodidae), in the laboratory. *Journal of Medical Entomology* 17, 287–289. https://doi.org/10.1093/jmedent/17.3.287

43. Nuss, A.B. *et al.* (2023) The highly improved genome of *Ixodes scapularis* with X and Y pseudochromosomes. *Life science alliance* 6. https://doi.org/10.26508/lsa.202302109

44. Chou, S. *et al.* (2023) De novo assembly of a long-read *Amblyomma americanum* tick genome. *Arcadia Science*. https://dx.doi.org/10.57844/arcadia-9b6j-q683

678	45.	Kotsarenko, K. *et al.* (2020) Karyotype changes in long-term cultured tick cell
679		lines. *Scientific reports* 10, 13443. https://doi.org/10.1038/s41598-020-70330-5
680	46.	Ammazzalorso, A.D. *et al.* (2015) To beat or not to beat a tick: comparison of
681		DNA extraction methods for ticks (*Ixodes scapularis*). *PeerJ* 3, e1147.
682		https://doi.org/10.7717/peerj.1147
683	47.	Hill, C.A. and Wikel, S.K. (2005) The *Ixodes scapularis* Genome Project: an
684		opportunity for advancing tick research. *Trends in parasitology* 21, 151–153.
685		https://doi.org/10.1016/j.pt.2005.02.004
686	48.	Kircher, M. and Kelso, J. (2010) High-throughput DNA sequencing–concepts and
687		limitations. *Bioessays* 32, 524–536. https://doi.org/10.1002/bies.200900181
688	49.	Sanger, F. *et al.* (1977) DNA sequencing with chain-terminating inhibitors.
689		*Proceedings of the national academy of sciences* 74, 5463–5467.
690		https://doi.org/10.1073/pnas.74.12.5463
691	50.	Guerrero, F.D. *et al.* (2010) Reassociation kinetics-based approach for partial
692		genome sequencing of the cattle tick, *Rhipicephalus (Boophilus) microplus*. *BMC*
693		*genomics* 11, 374. https://doi.org/10.1186/1471-2164-11-374
694	51.	Barrero, R.A. *et al.* (2017) Gene-enriched draft genome of the cattle tick
695		*Rhipicephalus microplus*: assembly by the hybrid Pacific Biosciences/Illumina
696		approach enabled analysis of the highly repetitive genome. *International journal*
697		*for parasitology* 47, 569–583. https://doi.org/10.1016/j.ijpara.2017.03.007
698	52.	Satam, H. *et al.* (2023) Next-generation sequencing technology: current trends
699		and advancements. *Biology* 12, 997. https://doi.org/10.3390/biology12070997
700	53.	Miller, J.R. *et al.* (2018) A draft genome sequence for the *Ixodes scapularis* cell
701		line, ISE6. *F1000Research* 7, 297.
702		https://doi.org/10.12688/f1000research.13635.1
703	54.	Benoit, J. (in prep.),
704	55.	Branton, D. *et al.* (2008) The potential and challenges of nanopore sequencing.
705		*Nature Biotechnology* 26, 1146–1153. https://doi.org/10.1038/nbt.1495
706	56.	Mostafa, H.H. (2024) An evolution of Nanopore next-generation sequencing
707		technology: implications for medical microbiology and public health. *Journal of*
708		*clinical microbiology* 62, e00246–00224. https://doi.org/10.1128/jcm.00246-24
709	57.	Luan, T. *et al.* (2024) Benchmarking short and long read polishing tools for
710		nanopore assemblies: achieving near-perfect genomes for outbreak isolates.
711		*BMC genomics* 25, 679. https://doi.org/10.1186/s12864-024-10582-x
712	58.	Dekker, J. *et al.* (2002) Capturing Chromosome Conformation. *Science* 295,
713		1306–1311. https://doi.org/10.1126/science.1067799
714	59.	Lieberman-Aiden, E. *et al.* (2009) Comprehensive Mapping of Long-Range
715		Interactions Reveals Folding Principles of the Human Genome. *Science* 326,
716		289–293. https://doi.org/10.1126/science.1181369
717	60.	Guizzo, M.G. *et al.* (2023) A comparison of Illumina and PacBio methods to build
718		tick salivary gland transcriptomes confirms large expression of lipocalins and
719		other salivary protein families that are not represented in available tick genomes.
720		*Ticks and tick-borne diseases* 14, 102209.
721		https://doi.org/10.1016/j.ttbdis.2023.102209

61.   Stanke, M. *et al.* (2008) Using native and syntenically mapped cDNA alignments
to improve de novo gene finding. *Bioinformatics* 24, 637–644.
https://doi.org/10.1093/bioinformatics/btn013
62.   Stanke, M. *et al.* (2006) Gene prediction in eukaryotes with a generalized hidden
Markov model that uses hints from external sources. *BMC bioinformatics* 7, 62.
https://doi.org/10.1186/1471-2105-7-62
63.   Thibaud-Nissen, F. *et al.* (2013) Eukaryotic genome annotation pipeline. *The*
NCBI Handbook 2,
64.   Flynn, J.M. *et al.* (2020) RepeatModeler2 for automated genomic discovery of
transposable element families. *Proceedings of the National Academy of Sciences*
117, 9451–9457. https://doi.org/10.1073/pnas.1921046117
65.   Smit, A. *et al.* (2015). RepeatMasker Open-4.0. 2013–2015. Seattle, USA
66.   Feron, R. and Waterhouse, R.M. (2022) Assessing species coverage and
assembly quality of rapidly accumulating sequenced genomes. *GigaScience* 11,
giac006. https://doi.org/10.1093/gigascience/giac006
67.   Geraci, N.S. *et al.* (2007) Variation in genome size of argasid and ixodid ticks.
Insect Biochemistry and Molecular Biology 37, 399–408.
https://doi.org/10.1016/j.ibmb.2006.12.007
68.   Van Zee, J.P. *et al.* (2007) Tick genomics: the *Ixodes* genome project and
beyond. *International journal for parasitology* 37, 1297–1305.
https://doi.org/10.1016/j.ijpara.2007.05.011
69.   Murgia, M.V. *et al.* (2019) Meeting the challenge of tick-borne disease control: a
proposal for 1000 *Ixodes* genomes. *Ticks and Tick-borne Diseases* 10, 213–218.
https://doi.org/10.1016/j.ttbdis.2018.08.009
70.   Sharakhova, M.V. *et al.* (2018) Physical genome mapping using fluorescence in
situ hybridization with mosquito chromosomes. In *Insect Genomics: Methods and*
Protocols, pp. 177–194, Springer
71.   Kim, B.Y. *et al.* (2024) Single-fly genome assemblies fill major phylogenomic
gaps across the Drosophilidae Tree of Life. *PLoS Biology* 22, e3002697.
https://doi.org/10.1371/journal.pbio.3002697
72.   Buenrostro, J.D. *et al.* (2013) Transposition of native chromatin for fast and
sensitive epigenomic profiling of open chromatin, DNA-binding proteins and
nucleosome position. *Nature methods* 10, 1213–1218.
https://doi.org/10.1038/nmeth.2688
73.   Nurk, S. *et al.* (2022) The complete sequence of a human genome. *Science* 376,
44–53. https://doi.org/10.1126/science.abj6987
74.   Frederick, J. *et al.* (2022) Phylogeography of the blacklegged tick (*Ixodes*
scapularis) throughout the USA identifies candidate loci for differences in
vectorial capacity. *Molecular Ecology* 32. https://doi.org/10.1111/mec.16921
75.   Willadsen, P. (2004) Anti-tick vaccines. *Parasitology* 129, S367–S387.
https://doi.org/10.1017/S0031182003004657