Analysis of SIV Spatiotemporal Dissemination Patterns in Rhesus Macaques During

Early Rectal Transmission Demonstrates Systemic Infection Does Not Require

Viral Local Amplification at the Entry Portal

Yilun Cheng[1], Jackson Chen,[1,2], Miaoyun Zhao[1,2], Subhra Mandal[1], Mark Lewis[3], Ma Luo[4,5],

Michael Gale, Jr.[6,7] Qingsheng Li[1,2*]

[1]Nebraska Center for Virology and School of Biological Sciences, University of Nebraska-

Lincoln, Lincoln, NE, USA.

[2] The Wistar Institute, HIV-1 Cure and Viral Diseases Center, 3601 Spruce Street Philadelphia,

PA, USA.

[3] BIOQUAL, Inc., Rockville, MD, USA.

[4]Department of Medical Microbiology and Infectious Diseases, University of Manitoba,

Winnipeg, Manitoba, Canada. [5]National Microbiology Laboratory, Public Health Agency of

Canada, Winnipeg, Manitoba, Canada.

[6]Department Department of Microbiology and Immunology, University of Minnesota, [7]Institute on

Infectious Diseases, University of Minnesota, MN, USA.

* Corresponding author:

 Qingsheng Li, qli@wistar.org

Abstract:

Receptive anal sex is a dominant mode of HIV-1 (HIV) transmission, especially among men who have sex with men (MSM). However, the early events during HIV rectal transmission are not well understood and many questions remain, such as, does virus local amplification at the entry portal require for viral distal seeding? What is the spatiotemporal dissemination pattern across whole body? How do three viral forms, e.g., cell-free virus (V_{CF}), cell-associated virus (V_{CA}), and follicular dendritic cell-trapped virus (V_{FDC}), evolve during early infections? To close this knowledge gap, we comprehensively examined rectum, draining and distant lymph nodes, as well as non-lymphatic organs of brain, lungs, and liver of 39 Indian rhesus macaques at the early time points following intrarectal SIV inoculation (1, 2, 3, 4, 6, 10 14, 28 day post inoculation, dpi). Our findings unambiguously demonstrated SIV rapidly disseminates to distant tissues and organs (<=1 dpi), virus local replication and amplification in rectum and draining lymph nodes is not required for viral distal seeding for the establishment of systemic infection, viral forms shift from V_{CF} and V_{CA} to V_{FDC} accompanying from T cell zones into B cell follicles. These findings indicates that an effective HIV vaccine needs to elicit comprehensive immune protection at both local sites of viral entry and the entire level.

Keyword: HIV/SIV early rectal transmission, local viral amplification, systemic infection, HIV vaccine

INTRODUCTION

Viral infection often begins with virus crossing the mucosal barrier and disseminating to distal sites through blood circulation, lymphatic system, neurological pathways along or in combination. A better understand of virus dissemination patterns and the role of virus local amplification at the entry portal after exposure is critical for developing vaccines and pre- and post-exposure prophylaxis. Different viruses have distinct dissemination patterns after the exposure. For example, rabies virus dissemination follows a multi-stage process where viral local replication and amplification in muscle tissues after animal bite and scratch is critical for virus dissemination and widespread infection in the central nervous system (1, 2) and Poliovirus local replication and amplification in the pharynx and the gastrointestinal tract has been demonstrated to be required for spreading to the central nervous system (3). However, the role of HIV local replication and amplification at the portal of entry for systemic infection is unknown.

HIV infection remains a global health threat. About 39 million people are living with HIV worldwide. Despite the presence of a combined antiretroviral therapy (ART) and several preventive measures, approximately 1.3 million HIV new infections were still reported in 2022 (https://unaids.org/en). Receptive anal sex is a major mode of HIV transmission, especially in men who have sex with men (MSM) (4-7), which accounted for 66% of all new HIV diagnoses in males in the United States (CDC, HIV surveillance report 2024, https://www.cdc.gov/hiv-data/nhss/hiv-diagnoses-deaths-and-prevalence-2025.html). Moreover, HIV vaccine development over the past 4 decades has made some progress, however, vaccines that prevent infection remain elusive. Thus, a better understanding of very early events of HIV mucosal transmission and viral spatiotemporal distribution patterns crossing viral entry site, draining and distal lymphatic organs and non-lymphoid organs of whole body, especially the role of viral local amplification at the portal of entry in the establishment of systemic infection, is needed for guiding HIV vaccine development (7-11).

For the convenience and ready availability of peripheral blood in clinical research, most

investigations on HIV transmission have primarily depended on the continuous monitoring of

peripheral blood samples (12-14). However, HIV target cells are mainly localized in mucosal

and the secondary lymphatic tissues (LTs) and there is an eclipse phase between the point of

initial infection to the time of viremia detection (15, 16). Due to ethical constraints in humans, in

vivo data regarding viral entry and early dissemination patterns can only be studied using

rhesus macaques (RMs) and Simian Immunodeficiency virus (SIV) model(17, 18). Nevertheless,

previously published RM SIV rectal transmission studies mainly focused on infection at the entry

site and viral-host mucosal interactions (19-21). Patricia and colleagues found that SIV viral

DNA was detected as early as 4 hours pi in colic lymph nodes, but not in axillary lymph nodes

until 2 dpi (5), Sui and colleagues reported only 15% RMs had detectable viral RNA (vRNA) in

the plasma as early as 4 day pi (6). Mair and colleagues reported that at 48 hours pi, infected

cell foci consisting of only a few cells scattered throughout the anal and rectal tissues. (22). We

previously reported that SIV vRNA was only detectable in the rectum at 3 dpi but not 1 and 2 dpi

using S^{35} labeled SIV riboprobe in situ hybridization (ISH), which was much less sensitive than

RNAscope ISH(19). In short, there was a significant knowledge gap in the very early events of

HIV rectal transmission before the current study.

In this research, we comprehensively evaluated the viral spatiotemporal distribution patterns

across whole body, the dynamic change of viral forms, and the role of virus local amplification

for the establishment of systemic infection using a highly sensitive RNAscope ISH and a

combination of CODEX (CO-Detection by inDEXing) immunostaining with RNAscope ISH

(Comb-CODEX-RNAscope). We found that virus local amplification at the portal of entry and

draining lymph nodes (dLN) are not required for viral systemic infection; at the 1 dpi, vRNA were

readily detected not only in the rectal and its dLN tissues, but also in the distal LN (disLN)

tissues as well as non-lymphatic organs of liver, lungs and brain. These findings support a new

model of SIV rectal transmission where SIV rapidly seeds to distal sites and viral local

amplification is not required for the establishment of systemic infection. These findings suggest

that an effective HIV vaccine needs to elicit robust immune protections at both the portal of viral

entry and at the systemic level.

MATERIALS AND METHODS

Animals and Ethics Statement

This report studied the samples from two cohorts of adult male rhesus macaques (*Macaca*

mulatta) of Indian origin. All the animals were specific pathogen free (SPF, negative for HIV-2,

SIV, type D retrovirus, and simian type D retrovirus 1) and without the protective major

histocompatibility complex class I alleles Mamu A01, B01, and B17. The first macaque cohort

study was approved by the Institutional Animal Care and Use Committee at the University of

Nebraska-Lincoln and BIOQUAL, INC. The rhesus macaque animal work was conducted at

BIOQUAL, Inc. (Rockville, Maryland). Twenty macaques were intrarectally inoculated with

SIVmac251 at a dose of 3.1×10^4 TCID50 and were euthanized at various time points of post

inoculation (3 animals at each time points of 3, 10, 14 and 28 dpi and 4 animals at 4 and 6 dpi),

while three rhesus macaques without virus challenge were served as uninfected controls (n=3).

The second cohort macaque study was approved by the University of Washington

Environmental Health and Safety Committee, the Occupational Health Administration, the

Primate Center Research Review Committee, and the Institutional Animal Care and Use

Committee. The rhesus macaque animal work was conducted at the Washington National

Primate Research Center. The macaques were intrarectally inoculated with SIVmac251 at a

dose of 6000 $TCID_{50}$ and were euthanized at various time points (n=4 at 1, 2, 3, 6 dpi). All the

tissues during necropsy were fixed with SafeFix II (Cat# 23-042600, Fisher Scientific), 4%

paraformaldehyde or neutral-buffered formalin and embedded with paraffin as we previously

reported (20, 23).

RNAscope In Situ Hybridization (ISH)

RNAscope ISH (RNAscope) from the Advanced Cell Diagnostics (ACD) was performed according to our previously published method(24, 25). RNAscope® Probe-SIVmac239 (anti-sense, Cat# 312811, ACD) was used and the signals were amplified and detected with RNAscope® 2.5 HD assay-Red kit (Cat# 322360, ACD). The RNAscope® negative control probe-DapB (Cat# 310043, ACD) was used as negative probe control and uninfected macaque tissues were used as negative tissue control. Tissue sections on slides were digitized using our Aperio ScanScope into our image server and perform image quantification analysis using our Aperio Spectrum system as previously reported(25).

Antibody-oligonucleotide conjugation:

To find a rhesus macaque reactive CD3 antibody for CODEX, an anti-human CD3 antibody in a carrier-free PBS solution (Clone#: SP162, Cat#: ab245731, Abcam) was conjugated with the barcode-oligonucleotide (Cat#: 5350002, Akoya) using CODEX conjugation kit (Cat#: 7000009, Akoya) following CODEX conjugation manual as we previously reported(26).

Combination of CODEX immunostaining with RNAscope ISH

CODEX (Co-detection by indexing) is a cutting-edge multiplexed tissue imaging technique that enables simultaneous visualization of dozens of protein markers while preserving tissue architecture. The Combination of CODEX immunostaining with RNAscope ISH (Comb-CODEX-RNAscope) enables detection of viral RNA via RNAscope and dozens of proteins. Comb-CODEX-RNAscope was performed according to our previously published method (26). CODEX and subsequent RNAscope ISH approach and the combination of pretreatments of CODEX and RNAscope together at the beginning of Comb-CODEX-RNAscope were used(26). Briefly, tissue sections of 6-µm in thickness were mounted on poly-lysine-coated coverslips. The

deparaffinization process began by heating samples in a 60°C incubator for 2 hours, followed by

two 5-minute xylene washes. Samples were then rehydrated through a series of decreasing

ethanol concentrations and DEPC water. Prior to initiating the CODEX cycling, tissues

underwent pretreatment with 3% hydrogen peroxide and antigen retrieval in pH 6 citrate buffer

(Sigma 21545).

The retrieved tissues were then incubated with a DNA-barcoded antibody mixture for 3 hours at

room temperature. Unbound antibodies were removed through three consecutive 2-minute PBS

washes. Post-fixation occurred in two stages: first with 1.6% paraformaldehyde for 10 minutes

at room temperature, followed by washing and a second fixation step using ice-cold methanol

for 5 minutes. After additional washing, samples were stored in buffer at 4°C for up to five days

or immediately processed in the CODEX instrument. Fluorescent reporters consisting of labeled

oligonucleotide probes matching the DNA-barcoded antibodies were prepared in a 96-well plate

according to manufacturer specifications. The CODEX fluidics system and Keyence microscope

operated under control of the CODEX instrument manager and Keyence software following

standard protocols. Upon completion of CODEX cycling, tissue-coverslips were transferred to

the RNAscope workflow beginning at the protease plus treatment step. Following RNAscope

completion, samples were counterstained with DAPI (1:2000 dilution) for 5 minutes at room

temperature, washed, and returned to the CODEX instrument for RNAscope image acquisition.

Integration of RNAscope data with CODEX images requires adding a blank cycle with Cy5-

channel assignment for RNAscope detection. Final image data was processed using CODEX

processor software after transferring to the appropriate file location.

Data Analysis for the Comb-CODEX-RNAscope

After the processing of Comb-CODEX-RNAscope raw data with CODEX processor software,

FlowJo v10 with CodexMAV extension was used to analyze data. Flow cytometry like gating

strategy was used to determine cell populations. The statistical analysis and figures of cell

populations and viral load dissemination change were made with Graphpad Prism 9.5.

Spatial Cluster Analysis of vRNA Distribution

Spatial analysis of vRNA distribution (Figure 4) was performed using QuPath software

(https://qupath.github.io/). Prior to analysis, the metadata for all whole tissue section images

was manually verified to ensure correct pixel size calibration. A pixel classifier was first trained

to segment tissue regions by distinguishing tissue from the white background. To separate the

stains, color deconvolution was performed. Initial stain vectors for nuclear stains with

Hematoxylin and vRNA signal of Fast Red were identified using the 'Estimate Stain Vectors'

command, followed by manual adjustment to ensure optimal stain separation. Following

segmentation, vRNA-positive signals were identified using the 'Positive Cell Detection'

command. The analysis was run at a requested pixel size of 0.5 µm/pixel. Key parameters were

set as follows: the background radius was 5 µm, the minimum and maximum detection areas

were set to 0.5 µm² and 5.0 µm², respectively, and cell expansion was set to 0 µm.

To identify spatial clusters of vRNA-positive cells, the Density-Based Spatial Clustering of

Applications with Noise (DBSCAN) algorithm was subsequently applied(27). The algorithm's

parameters were set with an epsilon (maximum search radius) of 50 µm and a minimum of 3

points (minPts) required to form a dense cluster.

RESULTS

SIV fast disseminates to draining and distal lymphatic tissues as well as non-lymphoid

organs of lungs, liver and brain during very early rectal transmission.

To assess SIV spatiotemporal dissemination after rectal transmission, adult male *Indian* rhesus

macaques (RMs) were intrarectally inoculated with a swarm of SIVmac251. The RMs were

euthanized at early time points of 1 (24 hrs)-, 2 (48 hrs)-, 3-, 4-, 6-, 10-, 14-, and 28-days post

inoculation (dpi, Fig. 1). Additionally, three RMs without SIV inoculation were euthanized to

serve as uninfected controls (Fig 1). A comprehensive necropsy was performed on each animal

to collect samples, including rectum, draining lymph nodes (dLN, mesenteric) of rectum, distal

lymph nodes (disLN, axillary and mandibular), spleen, and non-lymphoid organs of liver, lung,

and brain tissues (Fig.1). Tissue samples were divided into three portions for Safefix II fixation, 4%

PFA fixation, and snap-freezing, respectively. Peripheral blood mononuclear cells (PBMC) were

isolated before and after virus inoculation, and plasma was stored at -80C for SIV vRNA

quantification using qRT-PCR.

To detect SIV viral RNA (vRNA) in tissues that were collected during necropsies as described

above, RNAscope in situ hybridization (RNAscope) using SIVmac239 anti-sense probe (Cat#

312811, ACD) was performed, of which the detection sensitivity is a single copy of vRNA. The

RNAscope® negative control probe-DapB (Cat# 310043, Advanced Cell Diagnostics) was used

as negative control and vRNA signals were negative in the RMs of the uninfected control that

received sham inoculation by RNAscope with SIVmax239antisense-probe. on 1 dpi (24 hrs pi),

vRNA was readily detected in the rectal tissues and mesenteric dLN tissues of all 4 animals (Fig.

2). Importantly, vRNA was also detected in the disLN tissues of axillary and mandibular lymph

nodes (Fig. 2) and non-lymphoid organs of lungs, liver and frontal cortex and basal ganglion of

brain (Fig. 3, sTable 1), demonstrating that at 1 dpi, SIV already disseminated from the portal of

rectal entry to dLN, disLN, non-lymphatic organs of lungs, liver and brain. Of note, the

frequency and abundance of detected vRNA signals were extremely low, manifesting as a single

dot, which morphologically resembles a single virion derived signal. At this point of time of 1 dpi,

vRNA in peripheral plasma was not detectable using qRT-PCR. At 2-, 3- (Fig 4 and sTable 1)

and 4-dpi and thereafter, vRNA signals were readily detected in all the tissues that we examined,

including rectal, dLN, disLN, and nonlymphoid organs of lungs, brain and liver in all the animals.

To our knowledge, this is the first study that comprehensively investigated the spatiotemporal

viral dissemination across tissues and organs in such an early timeframe using the highly

sensitive RNAscope.

Virus Local Amplification at the Portal Entry Is Not Required for Establishing Systemic

infection.

To determine whether viral distal seeding is the consequence of viral local amplification at the

portal of entry and dLN, we ran a spatial cluster program to quantify virus replication and

amplification clustering. A viral cluster is defined as there are 3 or more distinct virions derived

from vRNA signals in a closely adjacent area of smaller than 50um in radius. As the diameter of

CD4 T cells is generally 5–10 µm (28) and three or more of virion in such as small area indicate

viral local amplification. On 1 dpi when vRNA was detected in disLN, spleen and non-lymphoid

organs of liver, lungs and brains, we did not find any vial clusters in rectal, dLN, disLN in all four

animals (Fig. 2), indicating virus local replication and amplification at portal rectal entry and dLN

is not essential for virus distal seeding. We detected viral clusters in both dLN and disLN in 2, 3,

and 4 dpi (Fig.4) and the time point thereafter, when SIV already widely disseminated to distal

tissues and organs, indicating vRNA replication and amplification across the body after

establishing the systemic infection. To our knowledge, this is the first study showed evidence

that SIV local amplification at the portal of entry is not required for the establishment of systemic

infection.

The Spatiotemporal Distribution Patterns of Three Different Vial Forms and Virally

Infected CD4 and Macrophage during Early Rectal Transmission

Three viral forms, e.g. cell-free (Vcf), cell-associated (Vca), and follicular dendritic cells (FDC)-

trapped (V_{FDC}), coexist in vivo in the secondary lymphatic tissues (LTs) during HIV infection.

Different viral forms play distinct roles in the establishment and maintenance of infection and

pathogenesis. FDC localizes in the light zones of germinal center in B cell follicles of the LTs,

which can trap HIV virions for an extremely long-period of time via immune complex. V_{FDC} has

been demonstrated to be infectious to migrating CD4 T follicular help cells (Tfh) and play an

important role in the HIV pathogenesis(29). In the very early infection (≤6dpi), all the SIV vRNA

we detected are Vca and Vcf (single dot outside cells) (Fig 5A). V_{FDC} was first detected on 10

dpi and became dominant viral form in LTs at 14 and 28 dpi (Fig.5).

To quantitatively examine the spatial distribution patterns of virally infected CD4 and

macrophage during early rectal infection in LTs, we used the Comb-CODEX-RNAscope, a

multiplexed proteins and SIV vRNA co-detection method(26) to visualize and quantify vRNA and

multiple immune cell markers simultaneously. A cocktail of antibodies to CD3, CD4, CD68,

CD20, CD21, CD31, Ki67, HLA-Dr, and IDO were used to identify immune cell types and

immune activation state in combination with vRNA detection. Figure 6 (Rh5429, 28dpi) shows

representative images for co-detection of vRNA and various immune markers. Using the Comb-

CODEX-RNAscope, we found that CD4 T cells are the primary cell type supporting productive

viral replication (vRNA+), making up greater than 90% of infected cells, with macrophages

constituting a minor fraction below 10% (data not show). To examine the spatial distribution

patterns of virally infected cells in LTs during early infection, we quantified v*RNA*+ CD4 T cells

and macrophages in T cell zones and B cell follicles of LN tissues at 10. 14, 28 dpi (n=3/each).

We first identified virally infected CD4 (CD4+ & vRNA+) or macrophage (CD68+ & vRNA+) by

combining two channels of signals, then used B-cell follicle marker CD20 as a gate to

differentiate virally infected cells in B-cell follicles vs. T- cell zones. We found that the population

of virally infected CD4 T cells distribution patterns shifted from mainly in T cell zones at 10 dpi to

mainly in B-cell follicles at 14 and 28 dpi (Fig. 7A). This shift is also true for virally infected

macrophage (Fig. 7B), which parallel with the shift of vRNA loads mainly in T-cell zones at 10

dpi to the B-cell follicles at 14 and 28 dpi in the viral form of V_{FDC} (Fig. 7C).

We also quantified vRNA in T-cell zones versus B-cell follicles, CD4 counts and immune

activation markers in the LN tissues of macaques at 0 (uninfected control), 1, 2, 3, 6, 10 and 28

dpi (n=4 at 1, 2, 3 dpi, and n=3 at 0, 6, 10, 14, 28 dpi). SIV vRNA in vivo growth curve in T-cell

zones peaked at 10 dpi and then declined (Fig. 7D), while vRNA in B-cell follicles was

significantly increased at 10 dpi and maintained at high level till 28 dpi (Fig. 7E).   Moreover,

there was a significant CD4 T cells decline and a significantly increased expression of immune

activation protein markers of ki67, HLA-Dr (Supplemental Fig. 3) as well as immune

suppression protein marker of Indoleamine 2,3-dioxygenase 1 (IDO1) that may counteract the

overall immune activation induced by SIV infection (Supplemental Fig. 3).

Discussion

The period of time between HIV initial exposure at the mucosal site and the virus spreads

throughout the body represents a critical window of time during which interventions could be

most effective (30, 31). Historically using less sensitive viral detection methods, it was believed

that a prolonged local viral amplification phase for small founder populations of infected cells

near the portal of entry is essential for the establishment of systemic infection (10, 32-37).

Deleage, C and colleagues, using barcoded-virus experiments and high-resolution mapping,

indicate that multiple microfoci within the female genital tract appear very early (hours–days)

preceded lymphatic dissemination, also before expansion and spread to distal sites (35).

Historically, tissue-level studies using radiolabeled ISH and immunohistology showed high LN

viral loads early after a few days of infection (24), and more modern RNAscope/IF-FISH

approaches have increased sensitivity and allowed cell-type colocalization (26, 38-40). A

convergence of data from rigorous necropsy studies in non-human primate rectal inoculation

models now reveals a dramatically accelerated timeline of systemic infection (41, 42). Results

from Ribeiro Dos Santos P and colleagues showed viral DNA or infectious virus in

draining/mesenteric lymph nodes and local tissues as early as 4 hours after rectal or mucosal

exposure, consistent with very rapid trans-epithelial transport or virus-bearing cell trafficking.

Within 2-3 days, robust replication was evident in systemic lymph nodes and the gut-associated

lymphoid tissue (GALT), which rapidly becomes a major site of viral amplification (42). When

talking about mechanisms of rapid viral spread during acute infection stage, some papers

demonstrate that cell-associated inoculation can seed tissues faster because infected cells

migrating from the mucosa into lymphatic channels appear to be one vehicle for early

dissemination (35, 43). However, the whole-body analysis of viral spatiotemporal distribution

patterns following early rectal transmission is unknow until this study.

In this study, we unambiguously demonstrated that SIV rapidly disseminated to rectal, draining

lymph nodes (dLN), distal lymph nodes (disLN), spleen, liver, lungs and brain at 1 day post

inoculation in all 4 macaques we studied using the ultra-sensitive RNAscope ISH that enables

us to detect single SIV virion. We detected both Vca and Vcf in multiple tissues of dLN, disLN

and non-lymphatic lung, brain and liver at 1dpi, indicating that both blood and lymphatic

circulation are likely involved in this early dissemination process (Fig. A). Our findings align with

the research conducted by Whitney and colleagues (14), who examined the effects of initiating

antiretroviral therapy (ART) at 6 hours and 1 day following intrarectal SIVmac251 infection.

Following cessation of the 6-month ART regimen, viral rebound rates were 0% in animals that

began treatment at 6 hours post-infection and 20% in those that started treatment at 1 dpi.

To determine the role of viral local amplification at the portal of rectal entry and dLN for a distal

seeding for the establishment of systemic infection, we comprehensively analyzed vRNA spatial

distribution in rectal, dLN and disLN in macaques on 1, 2, 3, 4 dpi. We hypothesized that if a

local viral amplification plays a crucial role in establishing distant viral seeding during initial

systemic dissemination (1 dpi), then the evidence of viral local replication and expansion should

be detectable within rectal and dLN tissues. We tested this hypothesis by using ultra-sensitive

RNAscope ISH and extensively examining vRNA in multiple sections of rectum, dLN, however

we did not detect any viral cluster, which is defined as at least 3 viruses within 50 micron radium

(about 5-8 lymphocyte distance), indicating that viral local replication is not essential for virus

distal seeding and establishment systemic infection. These findings suggest that an effective

HIV vaccine needs to protect not only the portal of viral entry but also at systemic level.

Understanding where virally infected-cells reside in the secondary lymphoid tissues (LTs) and

when virus replication and spread during the earliest stages of infection is critical to defining

founder populations, understanding early virus-host interaction and its immunopathogenesis,

and designing interventions to block systemic spread. In this study, we quantitatively analyze

the spatial distribution patterns of viruses and the temporal changes in percentage and

distribution of infected CD4$^+$ T cells and macrophages during very early rectal infection in LTs.

We employed Comb-CODEX-RNAscope to visualize and quantify vRNA and immune cell

markers concurrently and found that CD4$^+$ T cells dominate early infected-cell populations while

macrophages are secondary. Previous researches demonstrate that Tfh cells and effector-

memory CD4$^+$ subsets in visceral lymphoid tissues are important early and persistent reservoirs

(44), and macrophage-associated vRNA can localize to subcapsular sinus macrophages or

interfollicular macrophage populations depending on the LN microanatomy (45, 46). Here, using

Comb-CODEX-RNAscope and spatial analysis, we found that Vca of CD4$^+$ T cells and

macrophages as well as Vcf are mainly localized in T cell zones at and before 10 dpi, in contrast,

at 14 and 28 dpi , vRNA loads mainly shift to B-cell follicles that are associated with FDC

networks in the viral form of V_{FDC} and to a less degree as the viral form of Vca in CD4 and

Macrophages.   The B-cell follicles in LTs are unique anatomically and functionally, as there is a

low frequency of CD8 T cells and a high abundances of infectious V_{FDC}, thus B follicles may be

more favorable niche for viruses to survival by avoiding host's immune attack and uninfected

CD4 T cell and macrophages get infected by V_{FDC}.

In summary, using the RMs-SIV rectal transmission model, highly sensitive RNAscope and

Comb-CODEX-RNAscope, and by comprehensively examine rectum, dLN, DisLN and non-

lymphatic organs of liver, lung and brains, we unambiguously demonstrated that SIV rectal

transmission is not a staged disseminate model and viral local replication and amplification is

not a precondition for the establishment of systemic infection, and an effective HIV vaccine

needs to provide robust protection not only at the portal of virus entry but at the systemic level.

Reference:

1.   Davis BM, Rall GF, Schnell MJ. 2015. Everything You Always Wanted to Know About Rabies Virus
(But Were Afraid to Ask). Annu Rev Virol 2:451-71.
2.   Murphy FA, Bauer SP, Harrison AK, Winn WC, Jr. 1973. Comparative pathogenesis of rabies and
rabies-like viruses. Viral infection and transit from inoculation site to the central nervous system.
Lab Invest 28:361-76.
3.   Racaniello VR. 2006. One hundred years of poliovirus pathogenesis. Virology 344:9-16.
4.   Beyrer C, Baral SD, Van Griensven F, Goodreau SM, Chariyalertsak S, Wirtz AL, Brookmeyer R.
2012. Global epidemiology of HIV infection in men who have sex with men. The Lancet 380:367-
377.
5.   Ribeiro Dos Santos P, Rancez M, Pretet JL, Michel-Salzat A, Messent V, Bogdanova A, Couedel-
Courteille A, Souil E, Cheynier R, Butor C. 2011. Rapid dissemination of SIV follows multisite
entry after rectal inoculation. PLoS One 6:e19493.
6.   Yongjun Sui EML, Yichuan Wang, Alison Hogg, Blake Frey, David Venzon, Ranajit Pal, and Jay A.
Berzofsky. 2016. Early SIV Dissemination After Intrarectal SIVmac251 Challenge Was Associated
With Proliferating Virus-Susceptible Cells in the Colorectum. J Acquir Immune Defic Syndr
71:353–358.
7.   Kelley CF, Kraft CS, de Man TJ, Duphare C, Lee HW, Yang J, Easley KA, Tharp GK, Mulligan MJ,
Sullivan PS, Bosinger SE, Amara RR. 2017. The rectal mucosa and condomless receptive anal
intercourse in HIV-negative MSM: implications for HIV transmission and prevention. Mucosal
Immunol 10:996-1007.
8.   Deleage C, Immonen TT, Fennessey CM, Reynaldi A, Reid C, Newman L, Lipkey L, Schlub TE,
Camus C, O'Brien S, Smedley J, Conway JM, Del Prete GQ, Davenport MP, Lifson JD, Estes JD,
Keele BF. 2019. Defining early SIV replication and dissemination dynamics following vaginal
transmission. Sci Adv 5:eaav7116.
9.   Deleage C, Wietgrefe SW, Del Prete G, Morcock DR, Hao XP, Piatak M, Jr., Bess J, Anderson JL,
Perkey KE, Reilly C, McCune JM, Haase AT, Lifson JD, Schacker TW, Estes JD. 2016. Defining HIV
and SIV Reservoirs in Lymphoid Tissues. Pathog Immun 1:68-106.

10.   Haase AT. 2011. Early events in sexual transmission of HIV and SIV and opportunities for
402   interventions. Annu Rev Med 62:127-39.

403   11.   Martins MA, Gonzalez-Nieto L, Ricciardi MJ, Bailey VK, Dang CM, Bischof GF, Pedreño-Lopez N,
404   Pauthner MG, Burton DR, Parks CL, Earl P, Moss B, Rakasz EG, Lifson JD, Desrosiers RC, Watkins DI.
405   2020. Rectal Acquisition of Simian Immunodeficiency Virus (SIV) SIVmac239 Infection despite
406   Vaccine-Induced Immune Responses against the Entire SIV Proteome. J Virol 94.

407   12.   Brandon F. Keele FVH, Yingying Li, Elizabeth Bailes, Jun Takehisa, Mario L. Santiago, Frederic
408   Bibollet-Ruche, Yalu Chen, Louise V. Wain, Severin Loul, Eitel Mpoudi Ngole, Yanga Bienvenue,
409   Eric Delaporte, John F. Y. Brookfield, Paul M. Sharp, George M. Shaw, Martine Peeters, Beatrice H.
410   Hahn. 2006. Chimpanzee reservoirs of pandemic and nonpandemic HIV-1. Science.

411   13.   Smyth RP, Davenport MP, Mak J. 2012. The origin of genetic diversity in HIV-1. Virus Res 169:415-
412   29.

413   14.   Whitney JB, Lim SY, Osuna CE, Kublin JL, Chen E, Yoon G, Liu PT, Abbink P, Borducci EN, Hill A,
414   Lewis MG, Geleziunas R, Robb ML, Michael NL, Barouch DH. 2018. Prevention of SIVmac251
415   reservoir seeding in rhesus monkeys by early antiretroviral therapy. Nat Commun 9:5429.

416   15.   Shaw GM, Hunter E. 2012. HIV transmission. Cold Spring Harb Perspect Med 2.

417   16.   Yu M, Vajdy M. 2010. Mucosal HIV transmission and vaccination strategies through oral
418   compared with vaginal and rectal routes. Expert Opin Biol Ther 10:1181-95.

419   17.   Evans DT, Silvestri G. 2013. Nonhuman primate models in AIDS research. Curr Opin HIV AIDS
420   8:255-61.

421   18.   García-Lerma JG, Heneine W. 2012. Animal models of antiretroviral prophylaxis for HIV
422   prevention. Current Opinion in HIV and AIDS 7:505-513.

423   19.   Lu W, Ma F, Churbanov A, Wan Y, Li Y, Kang G, Yuan Z, Wang D, Zhang C, Xu J, Lewis M, Li Q. 2014.
424   Virus-host mucosal interactions during early SIV rectal transmission. Virology 464-465:406-414.

425   20.   Demers A, Kang G, Ma F, Lu W, Yuan Z, Li Y, Lewis M, Kraiselburd EN, Montaner L, Li Q. 2014. The
426   mucosal expression pattern of interferon-ε in rhesus macaques. J Leukoc Biol 96:1101-7.

427   21.   Anne Coue¨del-Courteille, Nathalie Barget SbJ, a Jean-Ge´rard Guillet,, Alain Venet c, and Ce´cile
428   Butor. 2003. Delayed viral replication and CD4+ T cell depletion in the rectosigmoid mucosa of
429   macaques during primary rectal SIV infection. Virology 316:290-301.

430   22.   Maric D, Corbin L, Greco N, Lorenzo-Redondo R, McRaven MD, Veazey RS, Hope TJ. 2023.
431   Temporal and spatial characterization of HIV/SIV infection at anorectal mucosa using rhesus
432   macaque rectal challenge model. bioRxiv doi:10.1101/2023.02.22.529624.

433   23.   Lu W, Demers AJ, Ma F, Kang G, Yuan Z, Wan Y, Li Y, Xu J, Lewis M, Li Q. 2015. Next-Generation
434   mRNA Sequencing Reveals Pyroptosis-Induced CD4+ T Cell Death in Early Simian
435   Immunodeficiency Virus-Infected Lymphoid Tissues. J Virol 90:1080-7.

436   24.   Yuan Z, Wang N, Kang G, Niu W, Li Q, Guo J. 2017. Controlling Multicycle Replication of Live-
437   Attenuated HIV-1 Using an Unnatural Genetic Switch. ACS Synth Biol 6:721-731.

438   25.   Ko A, Kang G, Hattler JB, Galadima HI, Zhang J, Li Q, Kim WK. 2019. Macrophages but not
439   Astrocytes Harbor HIV DNA in the Brains of HIV-1-Infected Aviremic Individuals on Suppressive
440   Antiretroviral Therapy. J Neuroimmune Pharmacol 14:110-119.

441   26.   Cheng Y, Burrack RK, Li Q. 2022. Spatially Resolved and Highly Multiplexed Protein and RNA In
442   Situ Detection by Combining CODEX With RNAscope In Situ Hybridization. J Histochem Cytochem
443   70:571-581.

444   27.   Rangaprakash D, Odemuyiwa T, Narayana Dutt D, Deshpande G. 2020. Density-based clustering
445   of static and dynamic functional MRI connectivity features obtained from subjects with cognitive
446   impairment. Brain Inform 7:19.

447   28.   Jiang X, Dudzinski S, Beckermann KE, Young K, McKinley E, McIntyre JO, Rathmell JC, Xu J, Gore JC.
448   2020. MRI of tumor T cell infiltration in response to checkpoint inhibitor therapy. Journal for
449   ImmunoTherapy of Cancer 8:e000328.

450   29.   Heesters BA, Lindqvist M, Vagefi PA, Scully EP, Schildberg FA, Altfeld M, Walker BD, Kaufmann DE,
451   Carroll MC. 2015. Follicular Dendritic Cells Retain Infectious HIV in Cycling Endosomes. PLoS
452   Pathog 11:e1005285.

453   30.   Wang X, Vincent E, Siddiqui S, Turnbull K, Lu H, Blair R, Wu X, Watkins M, Ziani W, Shao J, Doyle-
454   Meyers LA, Russell-Lodrigue KE, Bohm RP, Veazey RS, Xu H. 2022. Early treatment regimens
455   achieve sustained virologic remission in infant macaques infected with SIV at birth. Nature
456   Communications 13:4823.

457   31.   Igarashi T, Endo Y, Nishimura Y, Buckler C, Sadjadpour R, Donau OK, Dumaurier MJ, Plishka RJ,
458   Buckler-White A, Martin MA. 2003. Early control of highly pathogenic simian immunodeficiency
459   virus/human immunodeficiency virus chimeric virus infections in rhesus monkeys usually results
460   in long-lasting asymptomatic clinical outcomes. J Virol 77:10829-40.

461   32.   Haase AT. 2010. Targeting early infection to prevent HIV-1 mucosal transmission. Nature
462   464:217-23.

463   33.   Staprans SI, Dailey PJ, Rosenthal A, Horton C, Grant RM, Lerche N, Feinberg MB. 1999. Simian
464   immunodeficiency virus disease course is predicted by the extent of virus replication during
465   primary infection. J Virol 73:4829-39.

466   34.   Veazey RS, Gauduin M-C, Mansfield KG, Tham IC, Altman JD, Lifson JD, Lackner AA, Johnson RP.
467   2001. Emergence and Kinetics of Simian Immunodeficiency Virus-Specific CD8$^+$ T Cells in the
468   Intestines of Macaques during Primary Infection. 75:10515-10519.

469   35.   Deleage C, Immonen TT, Fennessey CM, Reynaldi A, Reid C, Newman L, Lipkey L, Schlub TE,
470   Camus C, O'Brien S, Smedley J, Conway JM, Del Prete GQ, Davenport MP, Lifson JD, Estes JD,
471   Keele BF. 2019. Defining early SIV replication and dissemination dynamics following vaginal
472   transmission. 5:eaav7116.

473   36.   Abel K, Pahar B, Van Rompay KK, Fritts L, Sin C, Schmidt K, Colón R, McChesney M, Marthas ML.
474   2006. Rapid virus dissemination in infant macaques after oral simian immunodeficiency virus
475   exposure in the presence of local innate immune responses. J Virol 80:6357-67.

476   37.   Milush JM, Chen H-L, Atteberry G, Sodora D. 2013. Early Detection of Simian Immunodeficiency
477   Virus in the Central Nervous System Following Oral Administration to Rhesus Macaques. Volume
478   4 - 2013.

479   38.   Deleage C, Chan CN, Busman-Sahay K, Estes JD. 2018. Next-generation in situ hybridization
480   approaches to define and quantify HIV and SIV reservoirs in tissue microenvironments.
481   Retrovirology 15:4.

482   39.   Jiang S, Chan CN, Rovira-Clavé X, Chen H, Bai Y, Zhu B, McCaffrey E, Greenwald NF, Liu C, Barlow
483   GL, Weirather JL, Oliveria JP, Nakayama T, Lee IT, Matter MS, Carlisle AE, Philips D, Vazquez G,
484   Mukherjee N, Busman-Sahay K, Nekorchuk M, Terry M, Younger S, Bosse M, Demeter J, Rodig SJ,
485   Tzankov A, Goltsev Y, McIlwain DR, Angelo M, Estes JD, Nolan GP. 2022. Combined protein and
486   nucleic acid imaging reveals virus-dependent B cell and macrophage immunosuppression of
487   tissue microenvironments. Immunity 55:1118-1134.e8.

488   40.   Moysi E, Del Rio Estrada PM, Torres-Ruiz F, Reyes-Terán G, Koup RA, Petrovas C. 2021. In Situ
489   Characterization of Human Lymphoid Tissue Immune Cells by Multispectral Confocal Imaging
490   and Quantitative Image Analysis; Implications for HIV Reservoir Characterization. Volume 12 -
491   2021.

492   41.   Raehtz KD, Xu C, Deleage C, Ma D, Policicchio BB, Brocca-Cofano E, Piccolo D, Weaver K, Keele BF,
493   Estes JD, Apetrei C, Pandrea I. 2024. Rapid systemic spread and minimal immune responses
494   following SIVsab intrarectal transmission in African green monkeys. JCI Insight 9.

495   42.   Ribeiro dos Santos P, Rancez M, Prétet J-L, Michel-Salzat A, Messent V, Bogdanova A, Couëdel-
496   Courteille A, Souil E, Cheynier R, Butor C. 2011. Rapid Dissemination of SIV Follows Multisite
497   Entry after Rectal Inoculation. PLOS ONE 6:e19493.

498   43.   Allers K, Kunkel D, Hofmann J, Stahl-Hennig C, Moos V, Schneider T. 2018. Cell-Associated Simian
499   Immunodeficiency Virus Accelerates Initial Virus Spread and CD4+ T-Cell Depletion in the
500   Intestinal Mucosa. The Journal of Infectious Diseases 217:1421-1425.

501   44.   Rabezanahary H, Moukambi F, Palesch D, Clain J, Racine G, Andreani G, Benmadid-Laktout G,
502   Zghidi-Abouzid O, Soundaramourty C, Tremblay C, Silvestri G, Estaquier J. 2020. Despite early
503   antiretroviral therapy effector memory and follicular helper CD4 T cells are major reservoirs in
504   visceral lymphoid tissues of SIV-infected macaques. Mucosal Immunology 13:149-160.

505   45.   Li Y, Kang G, Duan L, Lu W, Katze MG, Lewis MG, Haase AT, Li Q. 2015. SIV Infection of Lung
506   Macrophages. PLOS ONE 10:e0125500.

507   46.   Ponte R, Rancez M, Figueiredo-Morgado S, Dutrieux J, Fabre-Mersseman V, Charmeteau-de-
508   Muylder B, Guilbert T, Routy JP, Cheynier R, Couëdel-Courteille A. 2017. Acute Simian
509   Immunodeficiency Virus Infection Triggers Early and Transient Interleukin-7 Production in the
510   Gut, Leading to Enhanced Local Chemokine Expression and Intestinal Immune Cell Homing. Front
511   Immunol 8:588.

Figure 1. Experimental Design and Euthanasia Timeline of Rhesus Macaques Following Rectal SIVmac251 Inoculation. Upper panel shows the timeline for animal euthanasia following SIV rectal inoculation with animal numbers per time point. Lower panel shows the sampling strategy and assays utilized to track viral spatiotemporal distribution patterns across macaque organ systems.

1 dpi   Rectum   **Draining LN** (Mesenteric)   **Distal LN** (Submandibular)

Figure 2. SIV viral RNA (vRNA) was detected in rectum draining lymph node (dLN) and distal lymph node (disLN) tissues obtained from four macaques on day 1 post SIV inoculation. Representative micrographs showing SIV vRNA in rectal tissue, dLN (mesenteric), and disLN (mandibular) from all four rhesus macaques at 1 dpi. The vRNA was visualized as distinct red punctate signals with SIVmac239 antisense probes and Fast Red chromogen and cell nuclei were counterstained blue using hematoxylin. Rows designated as A1-4, each represents an individual macaque, and columns show the rectum, dLN and disLN tissues, respectively. Red arrow and green circle highlight vRNA signals. Higher magnification insets from the boxed regions of each tissue section are displayed.

Figure 3. SIV vRNA was detectable in non-lymphoid organs of lungs, liver and brain tissues obtained from four macaques on day 1 post SIV inoculation.

Representative micrographs from a small boxed area of insets showing SIV vRNA in lung (**A**), liver (**B**), and brain basal ganglia (**C**) of a macaque A1 at 1 dpi. The vRNA was visualized as distinct red punctate signals with SIVmac239 antisense probes and Fast Red chromogen and cell nuclei were counterstained blue using hematoxylin. Red arrows and green circles highlight vRNA signals.

Figure 4. SIV vRNA clusters present in lymph node tissues on 2, 3, and 4 dpi, but not 1 dpi.

Representative micrographs from a small boxed area of insets show SIV vRNA clusters within draining lymph node (dLN) tissues collected from macaques at 2, 3, and 4 dpi. vRNA clusters were identified through implementation of the Density-Based Spatial Clustering of Applications with Noise (DBSCAN) algorithm within QuPath image analysis software. A vRNA cluster is defined as ≥3 individual vRNA signals within 50 µm diameter regions. vRNA was detected using RNAscope. (A) shows 2 dpi, (B) 3 dpi, and (C) 4 dpi. No viral clustering was observed at 1 day post-infection across rectal tissue, draining lymph nodes, or distal lymph node specimens. Higher magnification insets from the boxed regions of each tissue section are displayed. Red arrows and green circles highlight vRNA signals.

Figure 5. Temporal and spatial dynamics of three distinct viral forms in lymph node tissues during acute SIV rectal infection

Representative micrographs show three distinct viral forms, e.g., cell-free viruses (Vcf), cell-associated viruses (Vca), and follicular dendritic cell-trapped viruses (VFDC) in LN tissues during early rectal infection. At 6 dpi (panels A and B), vRNA was exclusively detected as Vca and Vcf populations, with Vcf appearing as discrete signals external to cellular boundaries. By 10 dpi (panels C and D), VFDC first became detectable, characterized by diffuse vRNA signal patterns in the B follicles. The VFDC population emerged as the predominant viral form at both 14 dpi (panels E and F) and 28 dpi (panels G and H). Higher magnification images on the right side of each panel provide detailed visualization of the boxed regions from the corresponding left-side overview images.

Figure 6. Representative Comb-CODEX-RNAscope imaging of lymph node tissues.

The upper-left image displays the complete lymph node cross-section (28 dpi, Rh5429) with overlay of some fluorescent signal channels, with respective markers and corresponding color assignments as indicated. **The lower smaller panels** display magnified views of the designated boxed regions from the overview image, with each image showing individual detection channels. The **upper-right image shows a**n additional magnified view of the designated boxed regions from the overview image.

Fig 7　The spatiotemporal distribution patterns of virally infected cells in lymphatic tissues during early rectal transmission

SIV vRNA was detected using Comb-CODEX-RNAscope and the spatial distribution patterns of vRNA+ cells in T-cell zones versus B-cell follicles of mesenteric LN tissues over time were quantified CODEX processor software. The percentage of SIV vRNA+ CD4 T cells (**A**), vRNA+ macrophages (**B**) in T-cell zones versus B-cell follicles as well as the percentage of follicular

dendritic cell–associated virions (VFDC) in total vRNA signals over time of 10, 14 and 28 dpi

(n=3/each time) . (**C**) SIV vRNA in vivo viral growth curves in T-cell zones (**D**) and in B-cell

follicles, include Vca and VFDC (**E**) throughout early rectal transmission (n=4 at 1, 2, 3 dpi and

n=3 at 6, 10, 14, 28 dpi).

Supplemental Table 1

SIV vRNA Detection and Quantification in Brain, Liver and Lung Tissues During Early Rectal Transmission Using RNAscope In Situ Hybricyzation

ID	vRNA	Frontal Cortex				Basal Ganglia				Liver				Lung			
		vRNA	Area	Pos #	#/mm2	vRNA	Area	Pos #	#/mm2	vRNA	Area	Pos #	#/mm2	vRNA	Area	Pos #	#/mm2
A1	1	+	2.7	1	0.37	+	21.5	21	0.98	+	37.2	2	0.05	+	45.7	88	1.93
A2	1	+	28.9	38	1.32	+	14.8	8	0.53	+	41.7	2	0.05	+	26.9	43	1.6
A3	1	+	49.6	67	1.35	+	29.7	30	1.01	+	24.5	2	0.08	+	35.0	40	1.14
A4	1	+	12.9	4	0.31	+	30.8	23	0.75	+	15.8	1	0.06	+	14.4	46	3.21
B1	2	+	9.0	12	1.34	+	13.8	20	1.45	+	17.6	12	0.68	+	18.3	14	0.77
B2	2	+	36.9	7	0.19	+	7.7	27	3.53	+	56.3	9	0.16	+	15.6	33	2.12
B3	2	+	53.3	34	0.64	+	93.1	45	0.48	+	37.7	15	0.4	+	30.1	49	1.63
B4	2	+	33.8	5	0.15	+	73.1	50	0.68	+	47.8	4	0.08	+	19.9	38	1.91
C1	3	+	18.8	26	1.38	+	33.6	111	3.31	+	62.6	13	0.21	+	33.3	29	0.87
C2	3	+	52.6	1	0.02	+	62.1	62	1.00	+	52.1	32	0.61	+	14.0	27	1.94
C3	3	+	13.9	5	0.36	+	51.1	53	1.04	+	47.3	8	0.17	+	39.0	23	0.59
C4	3	+	10.8	191	17.7	+	19.3	27	1.40	+	43.3	26	0.6	+	13.8	38	2.76
Rh4971	3	+	103.8	6	0.06	+	119.0	8	0.07	+	97.5	113	1.16	+	101.2	264	2.61
Rh4972	3	+	76.37	13	0.17	+	70.5	1	0.01	+	160.9	3	0.02	+	202.2	108	0.53
Rh4973	3	+	110.7	35	0.32	+	112.5	4	0.04	+	110.2	7	0.06	+	179.8	131	0.73
RhL983	4	+	149.7	10	0.07	+	78.3	1	0.01	+	37.2	108	2.90	+	73.7	128	1.74
Rh5043	4	+	53.6	15	0.28	-	63.9	0	0.00	+	74.6	48	0.64	+	46.3	9	0.19
Rh5044	4	+	35.9	3	0.08	+	56.5	1	0.02	+	36.9	26	0.70	+	37.3	3	0.08
Rh5045	4	+	81.1	18	0.22	+	72.4	10	0.14	+	31.3	30	0.96	+	93.6	28	0.3
RhL864	6	+	32.5	S6	1.72	+	97.3	11	0.11	+	122.5	8	0.07	+	36.5	18	0.49
Rh4809	6	+	97.1	89	0.92	-	128.5	0	0.00	+	31.9	37	1.16	+	S5.2	8	0.15
Rh5052	6	+	89.7	22	0.25	+	39.3	6	0.15	+	43.3	11	0.25	+	45.4	11	0.24
Rh5053	6	+	70.6	9	0.13	+	27.4	3	0.11	+	85.3	35	0.41	+	53.7	71	1.32
D1	6	+	61.3	3	0.05	+	124.3	37	0.30	+	42.3	75	1.77	+	63.5	21	0.33
D2	6	+	12.3	6	0.49	+	43.9	16	0.36	+	65.3	2	0.03	+	14.8	5	0.34
D3	6	+	12.37	1	0.08	+	17.2	10	0.58	+	23.7	40	1.69	+	26.1	7	0.27
D4	6	+	3.29	18	5.47	+	27.7	12	0.43	+	36.0	15	0.42	+	12.3	7	0.57

Supplemental Figure1. SIV viral RNA (vRNA) in the draining lymph node (dLN) and distal lymph node (disLN) tissues on 2 and 3 dpi. Representative micrographs showing SIV vRNA in dLN (mesenteric), and disLN (mandibular) from all 2 rhesus macaques at 2 dpi (B1 & B4) and 3 dpi (C1 &C2). The vRNA was visualized as distinct red punctate signals with SIVmac239 antisense probes and Fast Red chromogen and cell nuclei were counterstained blue using hematoxylin. Green circles highlight vRNA signals. Higher magnification images from the boxed regions of each corresponding tissue section are displayed. Scale bar equals 100 microns.

Supplementary Figure 2. Immune activation and CD4 T cell decline in the LN tissues during

early SIV rectal infection. Immune cell markers were detected using Comb-CODEX-RNAscope.

CD4 T cells and immune activation markers in the LN tissues of macaques without SIV infection,

6 10 and 28 dpi (n=3/time point) were quantified. There was a significant CD4 T cells decline

(CD4/Cd3 ratio), an increased expression of immune activation protein markers of ki67, HLA-Dr

as well as immune suppression protein marker of Iindoleamine 2,3-dioxygenase 1 (IDO1).