Phase Behavior of TDP-43 and hnRNPH1: From Soluble and Aggregated States to Liquid-Like Droplets

Travis R Wang, Jiwu Wang

Allele Biotechnology and Pharmaceuticals, Inc., 6404 Nancy Ridge Drive, San Diego, CA 92121
The Scintillon Institute, 6868 Nancy Ridge Drive, San Diego, CA 92121

Corresponding Author:

Jiwu Wang
jiwuwang@allelebiotech.com

jiwuwang@scintillon.org

Abstract (200 words)

Cytotoxic inclusions of TAR DNA-binding protein 43 (TDP-43) have been identified in various neurodegenerative diseases. Mediated by its low-complexity C-terminal domain (CTD), TDP-43 can undergo liquid-liquid phase separation (LLPS) and form aggregates and fibrils. These phase transitions are associated with pathological TDP-43 mislocalization and cytoplasmic aggregation. What determines whether and when TDP-43 adopts a liquid droplet state or forms solid aggregates, transiently or irreversibly? A full understanding of the underlying biophysical mechanisms will help explain the cellular toxicity of aberrant forms of TDP-43 and similar proteins, such as some of the large family of RNA-binding proteins. In this study, we generated recombinant full-length TDP-43 and maintained soluble-protein working solutions without the need of a solubility tag. By focusing on conditions that tune the electrostatic characteristics, we induced phase transitions among states of soluble TDP-43, solid aggregates, and liquid droplet phases in an expedient and controlled manner. We also obtained soluble working solutions of TDP-CTD and another crucial splicing factor, hnRNPH1, and investigated their phase behaviors as individual proteins and their interactions between each other during phase transitions. Using this system, we discovered a previously unreported aggregate-to-droplet transition and found evidence that phase separation of low-complexity domain (LCD)-containing proteins likely involves domain matching when they coaggregate out of solution.

Introduction

Protein aggregation is a defining feature of many neurodegenerative disorders, including the accumulation of amyloid-β and Tau in Alzheimer's disease, α-synuclein in Parkinson's disease, and TDP-43 in amyotrophic lateral sclerosis (ALS). A central question in ALS study is what stimuli trigger TDP-43 aggregation and which elements and properties of TDP-43 respond to the triggering stimuli. TDP-43 is a ubiquitous RNA-binding protein (RBP) that regulates RNA splicing[1], with roles in regulatory control on mRNA homeostasis, translation, transcription, and transport[2].

Though predominantly residing in the nucleus with an N-terminal nuclear localization signal, TDP-43 shuttles between the nucleus and cytoplasm, a process found to be influenced by cellular conditions such as stress[3,4]. The N-terminus region promotes self-association into dimers and ordered oligomers[5], with evidence showing oligomerization into higher-order structures prerequisite to elements of protein function[6,7]. Congruent to its role as an RBP, TDP-43 contains two conserved RNA recognition motifs (RRMs) which mediate binding to RNA and DNA and are central for TDP-43's biological roles. The C-terminus includes a low-complexity domain (LCD) which is glycine-rich and aggregation-prone that regulates protein-protein interactions[8], splicing, and solubility during physiological function[4,7,9].

Pathologically, the accumulation of TDP-43 aggregates is observed in 90% of cases of ALS, 50% of frontotemporal lobar degeneration cases, and up to 75% in severe cases of Alzheimer's disease, according to some estimates[10]. However, even in physiological functions, TDP-43 does not always exist as a free soluble protein; rather, it transitions between solution and reversible liquid-like droplets (LLDs) with limited mobility[3]. These droplets can change in size and shape in response to cellular conditions. It appears that individual TDP-43 molecules and the oligomer complexes they form exist in a constant state of flux; protein-protein interaction and aggregate formation are always in play without necessarily leading to a pathological endpoint.

Under stress conditions, TDP-43 relocates to the cytoplasm to scaffold with other RBPs and RNA into dynamic stress granules, which lock down translation and sequester specific molecules as a coordinated cellular response to stress[11,12]. Extensive studies into stress granule composition have illustrated that increased concentrations of TDP-43 lead to liquid phase separation (LLPS; demixing) and depletion of native protein[13]. When forming different forms of aggregates, the protein concentration and interaction strength among protein molecules are key factors that

modulate LLPS from a proteome-wide perspective[14]. One proposed pathway for how soluble

TDP-43 congregates and misfolds into toxic aggregates is through prolonged or repeated

homotypic-interactions at elevated concentrations within these granules, with other triggers

playing a role such as oxidative stress, amino acid mutation and modification, and the presence

of RNA[15]. However, studies using tracking and localization microscopy uncovered not only distinct

substructures of TDP-43 within stress granules, but also numerous regions of slow-moving TDP-

43 species dispersed throughout the cytoplasm, indicating TDP-43 oligomerization and

aggregation can also occur outside of stress granules[16,17], where protein concentrations are not

constrained by granule confinement. The precise mechanisms governing the dynamic balance

among soluble proteins, oligomer formation, and subsequent transitions into droplets and fibrillar

assemblies are subjects of intense study, yet remain poorly understood even with significant

progress made over a decade by many groups of researchers.

The term "aggregate" relevant to low-complexity domain (LCD)-containing RNA-binding proteins

commonly include oligomers that may be in the form of liquid-like droplets, hydrogel-like "glassy"

condensates, or solid formations of various shapes, often progressing in that order [16-18]. Stress

granules and similar membraneless organelles, in which most of this process has been studied,

dynamically organize protein–protein and protein–RNA interactions in the cytoplasm and through

LLPS segregate into a concentrated phase from a dilute phase. LLPS in proteins is often driven

by LCDs, which mediate multivalent interactions. Both full-length TDP-43 and its C-terminal

domain (CTD) are known to undergo phase separation in vitro[14,15].

Insoluble inclusions of TDP-43 proteins containing fragments of its C-terminus are a hallmark in

numerous neurodegenerative diseases. It is quite remarkable that nearly all pathology-relevant

mutations of TDP-43 reside in its CTD[8], indicating the sensitivity of CTD's intrinsic functions to

any sequence perturbation despite its disordered structural morphology. This may suggest that,

prior to aggregation, factors beyond structural elements, such as the charges of key amino acids,

which can be influenced by extensive phosphorylation, acetylation, and other posttranslational

modifications concentrated within the region[8,19]. Additionally, charge distribution along the

sequence and neighboring effects, including stacking and hydrophobic interactions with specific

charged amino acids, may be essential for their behavior during phase transition[20]. In one report,

extensive mutagenesis was conducted to identify amino acid sequence as the "molecular

grammar" that dictates phase behavior in the Fused in Sarcoma (FUS) family[21]. In another study,

adjusting the placement of charged amino acids within an otherwise hydrophobic sequence

resulted in a twofold change in the tendency to phase-separate, without altering the net charge of

the polypeptide[22]. To manipulate the phase behaviors of TDP-43 through electrostatic alteration,

which shares many similarities with FUS, we created conditions that strongly altered the surface

charge of the protein and observed that it drove phase transitions into liquid droplets either directly

from soluble protein or, unexpectedly, through a novel reversal pathway from aggregates.

Phase separation is typically a thermodynamic process that occurs when protein-protein

interactions are more favourable than protein-solution interactions, separating proteins and

solutes into a dense phase[23]. Observed TDP-43-rich droplets in cells often contain internal

vacuoles filled with nucleoplasm materials, and these coexisting yet immiscible multiphase

occurrences correlate with ALS-causative mutations in CTD, indicating a higher-level complexity

of phase separation exists and is pathologically relevant[24]. Intriguingly, a reversible double-

emulsion structure (droplets-in-droplets) was created in vitro using Poly-rA RNA and PEG and

found to be due to kinetic trapping as opposed to thermodynamic equilibrium[25], but multi-phase

droplets have yet to be recreated with TDP-43 or other similar RNA-binding proteins in vitro in a

similarly controlled manner. In this report, we present our system as being able to show multi-

emulsion droplet formation with simple triggers and a consistent time course.

The CTD of TDP-43 has been found to form LLDs[26,27]. NMR and in vitro assays identified residue

321–335 of TDP-43 in the CTD region to be part of a helical core that tunes LLPS[14] whereas

proteolytic and cryo-EM studies identified residue 279-360 as the fibrillar core of TDP-43

filaments[28], encompassing the LLPS core. As for full-length TDP-43, phase separation has been

reported under unique poly(ADP-ribose) (PAR) binding conditions with concentrations of TDP-43

at 10μM[29], though in vitro LLPS of the full-length protein is still difficult to induce or control even

with favorable conditions such as using crowding agent PEG or aggregation-inhibiting molecules

dopamine and polyphenol EGCG[30,31]. Here, we present a novel system of manipulating liquid

phase separation of CTD and the full-length TDP-43, and for comparison, hnRNPH1. The current

paradigm states that TDP-43 LLPS droplets age into hydrogels, inclusions, and amyloid fibrils.

LLPS acts as a scaffold accelerating solid aggregate formation, often leading to direct

amyloidosis[30]. Our hypothesis is that the phasing-out of the soluble form starts with a "seed" area

in a protein-specific manner, which leads to early-stage substructural fibrils, forming solid

aggregates or LLDs depending on the predicted charge of this region and subsequent

electrostatic interactions. We further show the protein-specific seeding results by using switchable

combinations of fluorescent labelling schemes.

Our findings represent an unusual reversal of the conventional uni-directional liquid-to-solid

maturation of protein condensates, showing a more dynamic nature of the three macroscopic

phases of LCD-containing RNA-binding proteins, and reveal the likely role of electrostatic charge

in regulating LLPS.

Results and Discussion

Multidirectional phase transitions of TDP-43 are reversible and tunable

Liquid–liquid phase separation of full-length TDP-43 has been viewed as difficult to recreate in

vitro using recombinant proteins[32]. Several reported cases relied on an enzymatic cleavage of a

solubility tag to initiate aggregation, which is typically monitored by increased turbidity with the

enzymatic reaction difficult to control[7,33-35]. In other systems, a solution jump triggers phase

separation either almost instantly or very slowly—within minutes after a solution jump[29], or over 5

days[28]. In all cases, it has been challenging to fine-tune the process and capture time points of

phase transitions. To better control the transition process to gain clearer insight into the phase

behaviors of these proteins, we sought to generate recombinant full-length TDP-43 of sufficiently

high soluble concentration and stability without a large solubility tag.

The first step was to choose a method to keep TDP-43 soluble through purification. Most previous

attempts to produce full-length TDP-43 have relied on large solubility tags, such as GST[36], MBP[37],

and SUMO[15,28,31], which keep the fusion protein soluble until aggregation is triggered by enzymatic

tag cleavage or solution jump. Alternatively, like all proteins, recombinant TDP-43 can be readily

solubilized using strong denaturants such as 6-8 M urea or guanidine, with or without a solubilizing

tag, and this method has been employed by different investigators[29,36]. However, as complete

removal of these denaturants is often problematic because the protein tends to precipitate upon

denaturant elimination, a residual low concentration (e.g., ~0.6 M urea or guanidine) is typically

retained in storage buffer after purification, which may interfere with subtle aspects of phase

behavior. For our in vitro phase analysis system, we aimed to establish a baseline using full-

length TDP-43 without interference from large solubility tags or residual denaturants. We chose

to produce full-length TDP-43 with only a small His6-tag for purification and use urea to keep it

soluble during production, then refold to urea-free storage buffer. We solubilized full-length TDP-

43 expressed in E. coli using 6M urea that completely denatured its secondary structures without

affecting affinity purification via the His-tag. After washing and eluting with imidazole from cobalt

beads in 6 M urea-containing buffers, urea was gradually removed by stepwise dialysis against

buffer at 100-fold the sample volume. The urea concentration was reduced by half every 1–2

hours over 5-6 steps, concluding with a final exchange into urea-free buffer.

As the second step, we formulated the protein storage buffer by first testing whether high or low

salt conditions promote solubility. In the field of membraneless organelles and coacervates,

critical salt concentration is determined by the intrinsic properties such as charge-to-neutral ratio

among monomers of polymeric molecules such as DNA, RNA or protein, and charge distribution

and pattern[22,38]. Reducing the salt concentration below the critical concentration leads to energy-

favorable segregation of many types of proteins. This would suggest that keeping salt

concentration high should avoid forming aggregates during purification steps and in storage.

Many previous studies indeed used storage buffers containing high salt, e.g. 0.5 M NaCl or

KCl[28,31,36,37]. However, we found that TDP-43 tends to aggregate more under high-salt conditions

(e.g., 0.5 M NaCl) than under low-salt conditions (Supplementary Fig. 1a). Recent studies have

also reported that high salt promotes phase separation of soluble TDP-43[29,37].

The observation that high salt destabilizes the soluble state of TDP-43 suggests that charge

screening enhances hydrophobic and other non-ionic interactions, thereby promoting phase

separation. We hypothesized that similar effects could be achieved by adjusting the buffer pH

toward the protein's isoelectric point (pI). To test this, we prepared TDP-43 (5 μM) in a low-

capacity pH buffer (20 mM Tris-HCl, pH 8.0) and low salt (100 mM NaCl) and titrated the pH by

adding 1 M Tris-HCl at the target pH, maintaining 100 mM NaCl. Approximately 10% of the final

volume was required to reach the desired pH. Under these conditions, TDP-43 remained soluble

at pH 8.0, but the solution became turbid with diffuse assemblies at pH 7.0. At pH 6.0—near the

pI of full-length TDP-43—distinct solid aggregates became evident (Supplementary Fig.1b). This

observation is consistent with findings from the Chiti group[29], which implied that electrostatic

repulsion normally inhibits phase separation. Based on these results, we set the storage buffer

for recombinant TDP-43 as 20 mM Tris-HCl, pH 8.0, 100 mM NaCl.

By including urea during purification with a solubility-friendly storage buffer, we obtained working

solutions of TDP-43 suitable for phase transition analysis. A few points are noteworthy: a stepwise

removal of urea in our protocol enables the protein to refold through a more controlled process,

rather than by dropwise dilution or leaving residual denaturant[29,36]. Complete denaturation with

urea also eliminates the need for extensive RNA digestion followed by ion-exchange and gel-

filtration chromatography to remove nucleotides, which, if retained, can interfere with phase

separation[35].

To initiate analysis of the solution-to-aggregation transition, we altered salt concentration and pH

by adding 5 M NaCl and 1 M Tris-HCl (pH 6.0) to aliquots of TDP-43 (5 µM) in storage solution.

Once the buffer composition changes to high salt (0.5 M NaCl) at pH 6.0, TDP-43 almost

immediately aggregated out of solution and formed solid globules (Fig 1A). Over several hours,

these globules gradually assembled into larger structures composed of subunits that merged into

similarly disordered arrangements, reminiscent of fractal-like architectures (Fig. 1a). After 3 hours,

these aggregates ranged from clusters of only a few smaller congregates to assemblies spanning

hundreds of microns. The appearance of these structures closely resembles those described by

Lashuel and colleagues, which were composed of full-length TDP-43 oligomers released from

immobilized metal affinity chromatography only in high-imidazole (500 mM) fractions[28]. Unlike our

process, their assemblies required approximately five days to form. To assess whether a

commonly used solubility tag would alter TDP-43 phase behavior, we tested SUMO, a 101-

residue natural fragment with a pI of 5.2. When fused to TDP-43, the resulting construct has a pI

of 5.6, close to untagged TDP-43, leading us to predict that both versions would exhibit

comparable phase properties under our pH- and salt-based phase transition conditions. We

prepared a SUMO–TDP-43 fusion using the same 6 M urea denaturation and renaturation

protocol and observed very similar phase behavior to untagged TDP-43 (Supplementary Fig. 2a).

These results suggest that SUMO tagging preserves TDP-43 phase characteristics and may be

suitable to study phase transitions of other proteins, although pI matching should be considered.

Our findings that a change in TDP-43 net charge can modulate aggregate formation prompted us

to investigate whether alternative strategies for manipulating charge could similarly influence

TDP-43 phase behavior. The strong anionic detergent sodium dodecyl sulfate (SDS) micelles

facilitate denaturation of proteins and destabilization of protein complexes. The extent of

destabilization depends on the SDS concentration and the protein−SDS stoichiometry and the

resistance to 0.1-2% SDS has been used as a test of TDP-43 fibrillation strength[31,39]. Because

SDS introduces negative charges along the entire polypeptide, we therefore employed SDS at a

very low concentration, 0.03%, as a means of mildly modulating the overall protein surface

charge. Upon addition to TDP-43 aggregates, strikingly, the solid aggregates dispersed in the

field of view began to 'melt' into glassy material, but instead of dissolving and disappearing,

progressively transforming in shape and eventually forming round condensates, in about 6–

7 hours (Fig 1b). Formed droplets were quantified in circularity, roundness and solidity using

ImageJ quantification functions (Fig 1c). Coating aggregated TDP-43 with negative charge via

SDS did not dissolve the aggregates into a soluble state but instead transformed them into

droplets. We next asked whether introducing net negative charge to soluble TDP-43 would directly

induce LLPS. As shown in Fig. 1d, adding SDS to soluble TDP-43 indeed triggered droplet

formation, although the droplets were smaller and less dense than liquid-like formations formed

from aggregates. These early-phase droplets could be further enlarged and increased in

abundance by adding PEG8000 (Supplementary Fig. 3), a crowding agent commonly used to

promote LLPS[29,36]. The transitions from soluble state to solid aggregates induced by pH and salt,

and from soluble state or, as a novel observation, from solid aggregates to liquid-like droplets

upon SDS addition are schematically illustrated in Fig 1e.

The current paradigm for phase separation of LCD-containing RNA-binding proteins describes a

progression from soluble molecules to liquid-like condensates, then to glassy solids, and

ultimately to amyloid fibers[18]. It is interesting to compare the SDS-induced LLDs with LLDs formed

by means of adding negative net charges other than SDS. For instance, phosphomimetic

mutations reduced TDP-43 aggregates and increased the appearance of larger, more liquid-like

droplets[33]. Inside cells, phosphorylation is expected to add negative charges and there are many

other factors that can alter the net charge and charge distribution of proteins. McGurk *et al.*

reported that non-covalent binding of TDP-43 to poly(ADP-ribose) (PAR), a negatively charged

biopolymer and cellular regulator of protein localization and liquid demixing, promotes TDP-43

LLPS in vitro and is essential for its accumulation in stress granules in neurons[15]. Non-covalent

binding to PAR resulted in droplets comparable to those in our system using SDS, likely from

similar effects of adding net negative charge to TDP-43. As a matter of fact, binding to RNA would

add net negative charge to TDP-43 as well in normal cellular environment, which could enhance

LLPS according to our model along with suggestions by others[40]. Although SDS has not, to our

knowledge, been used as a trigger to induce LLPS in TDP-43 or other LCD-containing RNA-

binding proteins, it has been reported to promote LLPS of amyloid-β oligomers at

supraphysiological concentrations[41], suggesting its broader potential as a tool for modulating

phase behavior across diverse proteins. Demonstrating reversibility of solid formations with strong

anionic modulators could shed light on understanding TDP-43 pathology and lead to the

development of new therapeutic strategies.

SDS-induced droplets possess typical liquid-like properties

Three lines of evidence indicated that droplets derived from solid aggregates shown above exhibit

the same liquid properties as those formed by LLPS. First, these droplets fuse upon close contact

(Fig. 2a). Second, they dissolve in 1,6-hexanediol, similar to typical LLPS droplets (Fig. 2b).

Recent work has shown that glycine and other amino acids can bind phase-separating proteins

and modulate condensate stability and liquid-like dynamic[42], therefore, as a third test, we added

glycine to the droplets at 50 mM and observed pronounced changes within minutes: namely, a

dynamic formation of droplets-in-droplets, subsequent growth, and eventual remerging over time

(Fig. 2c). These dynamic changes were quantified by tracking the number of internal droplets

relative to the size of the encompassing outer droplets (Fig. 2d), as well as the average count of

droplets over a 12-minute period (Fig. 2e). Upon glycine addition, there was a rapid increase in

internal droplet formation, coordinated with the growth of the encompassing droplets. This was

followed by a consolidation phase, during which the average number of internal droplets

decreased. Altogether these corroborating assays provide strong evidence that the dynamic

liquid-like behavior of these solid-derived droplets.

When Cy5-labeled mRNA of a fluorescent protein mWasabi as biologically irrelevant mRNA was

allowed to bind TDP-43 prior to phase separation, the Cy5 imaging revealed that the mRNA

colocalized with the protein within the dense phase of the multi-emulsion droplets (Supplementary

Fig. 4), consistent with previous findings that full-length TDP-43 oligomers retain RNA-binding

capability[31]. These observations further indicate that the contents of these droplets exhibit liquid-

like properties similar to typical LLPS condensates, despite originating from solid assemblies

transformed by forced charge alteration.

Phase transition behavior of TDP-43 can be generalized to other RNA-binding proteins.

To see if any of these phase behaviors are specific to full-length TDP-43, we wanted to compare

TDP-43 to hnRNPH1, another nuclear RNA-binding protein that regulates alternative splicing with

implications in ALS and other diseases[43]. Similar to TDP-43, hnRNPH1 also has an LCD at the

C-terminus of the polypeptide composed of 449 residues, close to 414 in TDP-43; its pI is the

same at pH 6, allowing us to apply the same pH and salt conditions to manipulate the electrostatic

and ionic environment for phase transitions as TDP-43. We also wanted to compare the whole

TDP-43 protein to its aggregation-prone domain. In vivo, neuronal inclusions contain ubiquitinated

and phosphorylated aggregates of TDP-43 C-terminal (TDP-CTD) fragments in various sizes

(e.g., TDP-25 and TDP-35), created by aberrant cleavage events that contribute to ALS[44,45]. TDP-

CTD was initially identified as the fragment encoding amino acids 274-414 by pioneering

studies[14,26]. Later, 266-414[46], 267-414[27], and longer CTD 193-414 were also studied[47], and all

CTD segments have been found to play a major role in phase separation as well as aggregations

of TDP-43. The 274-414 fragment CTD has a pI close to pH 10. It is previously reported to have

a different phase separation response to pH gradients from the full-length[29], likely related to

different net charge at different pH compared to the full-length TDP-43 protein. We therefore opted

to study an intermediate-sized version of TDP-CTD, spanning 208 to 414 amino acids, which has

a predicted pI at pH 6. This allowed us to compare it to the full-length TDP-43 in parallel and

during protein mixing experiments.

As with full-length TDP-43, both TDP-CTD (208-414, hereby referred to as TDP-CTD in this

report) and hnRNPH1 formed solid assemblies when their solutions were adjusted to pH 6.0 and

0.5 M NaCl. Upon addition of 0.03% SDS, these assemblies transitioned into droplets, exhibiting

behavior broadly similar to that of full-length TDP-43 (Fig. 3a). However, the dynamics and

morphological details of the resulting aggregates and droplets varied among the proteins. For

example, TDP-CTD consistently formed larger and more compact aggregates compared to TDP-

43 and hnRNPH1. Whilst the primary focus of this report is to demonstrate macroscale phase

transition processes that can be modulated by solution conditions, further insights into the fine

structural differences underlying the various assembly forms of these proteins could be gained by

high-resolution imaging in future studies. This result nevertheless supports that the pI and high

salt promotion of solid aggregation and their SDS-induced conversion into droplets are not

specific to full-length TDP-43. This is consistent with the notion that phase behavior can be

dictated by a general mechanism based on electrostatic and hydrophobic interactions.

TDP-CTD droplets and aggregates are known to contain amyloid filaments[27,48,49]. To examine the

amyloid nature of our TDP-43 and TDP-CTD solid aggregates, we stained them with classical

amyloid-binding dyes, including thioflavin-T (ThT) and Congo Red (CR), both of which strongly

fluoresce upon binding to β-sheet-rich amyloid structures. They exhibited clear and nearly uniform

positive staining, suggesting that amyloid structures are present throughout the formations and

likely organized at the substructural level rather than being confined to larger, assembled

structures (Fig 3b). Next, we stained the droplets formed from both TDP-43 and TDP-CTD solid

aggregates with SDS addition and discovered that these droplets too were remarkably ThT and

CR positive (Fig 3c). Extensive studies by many groups on LLPS of TDP-CTD have shown that

amyloid fibrillation can occur independently of LLPS, as observed in our solid aggregates;

whereas ThT-positive amyloid fibrils have also been detected within liquid droplets, consistent

with our findings in solid-derived droplets[27,50]. Although dye-staining with ThT and CR by itself is

not unique to amyloids, the strong and consistent signals here nevertheless provide clear

likelihood that these assemblies contain amyloid elements and justify further investigations using

high-resolution imaging techniques when available for analysis of the assembly dynamics.

Namely, how the helical structures present in both full-length TDP-43 and its CTD during

assembly transitions can potentially give rise to β-sheet-rich structures[26,39,46].

The formation of multi-emulsion or multiphase droplets is a characteristic feature of complex

coacervate (polyelectrolyte material) mixtures, which are widely viewed as models for studying

charge-mediated LLPS[38]. By patterning charged monomers along a polymer backbone such as

polypeptides, one can precisely adjust local charge organization and the resulting electrostatic

interactions between coacervate-forming chains, creating multi-phase or multi-emulsion

droplets[22]. In TDP-43, charged amino acids are naturally distributed along the polypeptide, with

additional charges that can be introduced in vivo through phosphorylation and other post-

translational modifications, particularly concentrated in the CTD. In our in vitro phase-separation

system, a droplets-in-droplets morphology can be induced simply by adding glycine to SDS-

induced droplets (Fig 2c). As previously reported, glycine interacts weakly with amide groups in

the protein backbone and aromatic side chains, reducing backbone–backbone contacts while

enhancing interactions among aromatic residues[42]. The biological relevance lies in the surprising

observation that free amino acids account for more than 25% of the total volume and 6% of the

total dry mass of a mammalian cell[51], as well as their natural role in stabilizing proteins and

colloidal dispersions[52]. We set out to determine the nature of multi-emulsion droplets formed

following glycine addition to SDS-induced single droplets. To do this, we repeated the experiment

shown in Fig. 2c but in the presence of ThT, and used confocal imaging to visualize phase

behavior and amyloid properties. Z-stacks of representative complex droplets revealed alternating

dense (coacervate) and dilute (supernatant) phases within larger droplets (Fig. 3d). The 'bubbles'

were ThT-negative, whereas the coacervates were ThT-positive, confirming the presence of

amyloid substances in protein-rich coacervates at the 3D level. This result further highlights the

liquid-like dynamics of SDS-induced droplets and the secondary phase separation that can be

triggered by electrostatic effectors such as glycine.

Amyloid formation starts upon phase separation

Compared to the CTD of TDP-43, findings reported in the literature on the phase separation

behavior of full-length TDP-43 and the role of amyloid fibrillation are less consistent, presumably

due to the challenges of producing recombinant full-length TDP-43 without large solubility tags or

denaturing agents such as urea, even at low concentrations[32]. Among the few studies on full-

length TDP-43, some reported the presence of aggregates composed of proteins with disordered

secondary structure that failed to bind ThT[19,53,54]. Other studies observed ThT-positive segregated

structures, but the process required extended time, approximately 10 hours in one case[55] and 9

days with agitation in another[56]. Understanding this time course may provide valuable insights

into the fibrillation pathway, including the involvement of LLPS and solid aggregates, and is

essential for clarifying how disease-associated amyloid fibrils form. Leveraging the controllability

of aggregate and droplet formation using our solubility tag-free recombinant proteins and their

strong ThT staining once high order structures form, we performed fluorescence video recording

to capture the dynamic process of TDP-43 solutions transitioning into droplets with SDS or

forming solid aggregates under conditions of pH near the pI and high salt. Time-lapse images

extracted from videos recorded at fixed fields of view from two independent repeats show that,

upon addition of SDS to 0.03% at time zero, small droplets began to emerge and progressively

increased in size, through joining from soluble proteins or by merging with other condensates (Fig.

4a). Note that ThT was included in the protein solution prior to SDS addition and showed positive

staining in both small and large droplets at the first captured time point. While fluorescence

intensity increased over time, the rate of increase appeared generally in line with the growth in

droplet size, with a slight delay that could reflect amyloid fibrillation progressed in addition to

droplet growth; the total droplet counts increased during the first 4 hours before decreasing

through merging (Fig. 4b). A low concentration of glycine at 5mM, 10% of what was used in Fig

2c, was also included, and by hour 7, double-emulsion droplets shown as dark circles began to

appear, consistent with the results shown in Fig. 2c but at a slower pace. We then applied the

same video recording approach to monitor solid aggregate formation from soluble solutions.

Similar to SDS-induced droplet formation, aggregates triggered by pH near the isoelectric point

and high salt showed ThT staining shortly after the induction. These amorphous, fractal-like solids

continued to grow, originating entirely from recruiting soluble proteins as no merging events were

observed (Fig. 4c). Analysis of the area and intensity of the solid formations supports our

interpretation that aggregates rapidly grew from unstained protein molecules and began to

plateau at around 3 hours (Fig. 4d). These observations were consistent across multiple trials and

variations using different imaging systems, all showing a smooth transition from unstained

solution to clear and immediate amyloid-positive staining. This behavior contrasts with some of

the previous reports on full-length TDP-43, as discussed above. The discrepancy may stem from

our approach: our TDP-43 protein was fully denatured and then slowly renatured during

recombinant protein preparation, likely maintaining a more native-like conformation ready for

orderly assembly. Additionally, our induction was carried out by adding the triggering solution at

less than 10% of the final volume, creating lower perturbation or stress to the system than other

protocols by dropping high concentration protein solutions into low concentration buffers, perhaps

better reflecting biological events in vivo.

Switchable fluorescent protein tags are a valuable tool for studying protein phase behavior

We were also interested in determining whether different proteins, sharing the same or similar but

distinct LCDs, would co-aggregate or remain segregated upon triggering phase separation. To

address this, we designed a system to track multiple proteins using distinguishable fluorescent

labels. Because phase separation in our system depends on charges that influence electrostatic

and hydrophobic interactions, we avoided commonly used dyes that bind to amino acids and may

alter charge. Structural dyes such as ThT and CR are unsuitable for aggregate combination

studies because they lack protein-specific tracking capability. Instead, we adapted a switchable

tagging system to label TDP-43, TDP-CTD, and hnRNPH1 with spectrally distinct fluorescent

proteins (FPs). A 13-amino-acid SpyTag was added to the C-terminus of each protein via a 7-

amino-acid linker, and the 116-amino-acid SpyCatcher was fused to the N-terminus of

mNeonGreen[57], mScarlet3[58], and mTurquoise2[59], all of which represent FPs with optimal

brightness, physical and photo stability, and more stringent monomeric properties within their

respective spectrum to maximize fluorescence signals while minimizing aggregation artifacts (Fig.

5a). By briefly mixing a pair of SpyTag- and SpyCatcher-fused proteins, a fluorescent label is

covalently attached to the target protein via SpyTag–SpyCatcher bonding. The protein ratios

within each pair were optimized, with a slight excess of SpyCatcher-FP to ensure that all SpyTags

on target proteins were occupied (Supplementary Fig. 5).

The pI of all three SpyCatcher-FPs ranges between 5.0 and 5.5. When fused with TDP-43, TDP-

CTD, and hnRNPH1, the resulting fusion proteins have pIs between 5.5 and 6.0, predicting low

net charges like the untagged proteins under our aggregation induction conditions. As expected,

when the buffer was adjusted to pH 6.0 with 0.5M NaCl, the fused proteins formed solid

aggregates, exhibiting strong and uniform fluorescence (Fig. 5b). Each protein can be fused to

any FP without apparent differences in behavior when the FP was switched among the three

monomeric variants (Supplementary Fig. 6). We concluded that adding an FP to the C-terminus

of these proteins does not significantly alter their phase behavior, while it affords a convenient

and switchable end labeling system to study these proteins. Older versions of green and red FPs

have been previously used in fusions to study TDP-43 both in vivo and in vitro[37,53,60]. Our data is

consistent with these findings, indicating that FP tags offer reliable information about native

proteins. Most commonly used FPs have a slightly acidic pI, making them unlikely to interfere with

pH-dependent effects on net charge or electrostatic interactions of proteins with similar pIs as that

of TDP-43. However, caution may be warranted about the tendency of conventional avGFP-

derived GFP, YFP, and their variations to form weak dimers[61], particularly when they are used to

study oligomerization and seeding for aggregation.

With fluorescence signals appended to each protein, we performed time-lapse imaging to track

the transition of individual aggregates of TDP-43 and TDP-CTD that were fused to mScarlet3 and

mNeonGreen, respectively. Spread-out aggregates of each protein began condensing within 10

minutes of SDS addition and self-merged into spheroid structures within 1 hour (Fig. 5C), in

agreement with results shown with non-fusion TDP-43 (Fig. 1b).

Electrostatic-driven phase separation is differential to aggregation seeding

Having observed that TDP-43, its CTD, and hnRNPH1—the latter a distinct protein with

comparable features to TDP-43—all exhibited parallel behaviors, we explored whether phase

separation is directed by protein-specific features, or if it is driven by a common physical property-

based process, such as hydrophobic inclusion. We pairwise-mixed all three proteins, each tagged

with a different FP, and examined aggregates for the presence of two different proteins.

Interestingly, coaggregates were readily observed between full-length TDP-43 and TDP-CTD (Fig.

6). However, in all view areas obtained from repeated attempts, no coaggregates were detected

between hnRNPH1 and either TDP-43 or TDP-CTD. Because TDP-CTD is a segment of TDP-43

and known to be required for phase separation[24,26], we hypothesized that the cocomplex between

TDP-43 and TDP-CTD forms through the shared helical seed within the LLPS core previously

identified by biochemical and structural studies[14,28]. The LCD core sequence of hnRNPH1 differs

from that of TDP-43's while it is also known to be involved in phase separation[43]. These results

suggest that protein assembly is not driven solely by hydrophobic or other nonspecific physical

forces, as the process appears protein-specific. Instead, it is plausible that under conditions

controlling electrostatic interactions combined with hydrophobic segregation from the aqueous

phase, the phase separation appears mediated, or even initiated, at their LCD cores, similar to

the concept of "steric zipper"[62]. These flexible regions may allow transient 'kiss-and-run'

interactions when no matching partner is found, or bundle to form fibrils and develop amyloids

when matching occurs. Recently, non-canonical mechanism of protein recognition, wherein

transient interactions surrounding the TDP-43 conserved helical region drive the phase separation

of CTD through direct enhancement of site-specific, helix-mediated interactions, was illustrate as

an example of how the amino acid sequence can be used as a code for segregation[63]. It was

previously reported that the mutations within the core of 279-360 (as defined by Kumar *et al.*[28])

rendered TDP-43 less liquid-like[24], and led to defects in stress granule formation[64]. The sequence

specificity in multiple protein assembly formation revealed by our results could be useful for

understanding the effects of these mutations.

Toward modulation of phase transition

Based on the experimental data presented in this study, we propose a unifying perspective about phase behavior of proteins such as TDP-43. Electrostatics related to charge, whether global or localized, arising from natural amino acid composition and post-translational modifications is a major determinant. In vitro attunement to pH, salt screen, zwitterionic glycine (and other amino acids) or charge decoration by ionic detergent SDS at very low concentration, can modulate how proteins like TDP-43 transition among various macroscopic phases. These manipulations can induce phase transitions by shifting some microscopic substructures such as dimers, oligomers, nonamyloid fibrils and amyloids to reach the lowest-energy steady state. These substructures all retain a degree of fluidity and reversibility, allowing the system to respond dynamically to environmental changes.

Protein assemblies can be visualized at the micron scale using optical microscopy, as shown in this study, or at the nanometer scale with electron microscopy and other high-resolution techniques like NMR, revealing details such as filament-containing helical amyloid core[28]. Large assemblies likely originate from smaller subunits that act as building blocks, beginning as oligomers and progressing through stepwise assembly. The process typically takes time, occurring over hours or days[31], though notably faster in our system. These substructures may share physical and chemical characteristics with the larger assemblies they form, either through fractal growth into amorphous solids or by coexisting in a milieu that phase-separates into liquid-like droplets. In our system, strong ThT and CR staining was observed in both solids and droplets, during and after solid-to-droplet transitions with no delays, suggesting that different forms might share related substructures. These staining events may originate from a seed region containing an amphipathic helix shared among aggregating molecules[29], which undergo conformational transitions toward increased β-sheet stacking during maturation, under stress, or with mutations,

as previously proposed[30,49,65]. Such seeding could allow the initial helical core and adjacent

sequences to act as a sorting mechanism, determining interaction partners and conditions. It is

important to note that RNA-binding proteins can also interact between each other as soluble

molecules via LCDs, such as glycine-rich regions, to perform biological functions without

undergoing phase transitions, as shown in our earlier in vitro protein-protein interaction

studies[66,67].

TDP-43 is known to co-segregate under physiological conditions with numerous other proteins,

including hnRNPA1, hnRNPA2, and FUS[3,68,69], through domain-specific physical contact, via RNA

binding, or perhaps coaggregation based on matching helix or β-sheet stacking at seed region.

The phase transitions observed here with TDP-43 and hnRNPH1 may extend beyond RNA-

binding proteins involved in splicing regulation. A notable early example is Sear's 2007 prediction

that the Dvl protein would undergo LLPS[70], an all-or-none switch that has since been shown to

play a critical role in the Wnt signaling pathway[71].

The discovery that solid aggregates can be driven into a more liquid-like state may open new

avenues for identifying therapeutic intervention points by preserving normal, reversible LLPS that

does not progress to irreversible fibril amyloid formation[72]. Potential strategies include using short

RNAs to provide specificity by targeting RNA-binding proteins, and amino acids or oligopeptides

to achieve charge-modifying effects.

Materials and Methods

Protein production

Plasmids encoding TDP-43, TDP-CTD, hnRNPH1, and SUMO–TDP-43 were cloned into the

pNCS vector[57] and transformed into NEB SHuffle E. coli competent cells. Selected colonies were

inoculated into culture tubes containing 5mL 0.1 mg/mL carbenicillin 2xYT broth and incubated at

30°C overnight while shaking, the next day transferred to 500mL 2xYT broth and again incubated

overnight while shaking. For expression, small cultures of transformed bacteria were inoculated

in 500 mL 2×YT broth supplemented with 0.1 mg/mL carbenicillin at 30°C while shaking overnight.

Cells were harvested by centrifugation at 8,000 × g, and the pellets were resuspended in lysis

buffer containing 20 mM Tris-HCl (pH 8.0), 6 M urea, 500 mM NaCl, and 10 mM imidazole.

Cells were lysed by sonication on ice (30% amplitude, 5 min total, 10 s on/10 s off) and centrifuged

at 8,000 × g to remove debris. The supernatant was incubated at 4 °C for 1 h to overnight with

nickel beads to capture His-tagged proteins. Bead–protein complexes were transferred to

purification columns and washed 3–4 times with wash buffer (20 mM Tris-HCl, pH 8.0; 500 mM

NaCl; 10 mM imidazole; 6 M urea). Proteins were eluted in a single step using elution buffer (20

mM Tris-HCl, pH 8.0; 500 mM NaCl; 200 mM imidazole; 4.5 M urea) and filtered through 0.22 µm

PES syringe filters.

Protein purity was assessed by SDS-PAGE, and concentrations were measured using NanoDrop.

Eluted proteins were standardized to 5 µM, or about 0.2 mg/ml, after removal of urea in gradual

5-6 step dialysis using Spectra/Por dialysis tubing (MW cutoff 7,000) against storage buffer of 20

mM Tris-HCl, pH 8.0 100 mM NaCl with urea concentration decreasing ~50% per step until all

urea was removed. Soluble proteins were collected and stored at −20 °C.

Formation of solid aggregates

Phase separation reactions for TDP-43, TDP-CTD, and hnRNPH1 were initiated from soluble

protein fractions by adjusting the storage buffer from pH 8 to pH 6 using small-volume additions

of 1 M Tris-HCl pH buffer. Additional modifications, such as changes in salt concentration, were

made by adding small volumes of high-molar salt solutions. The total reaction volume was

approximately 40 µL. All added reagents to modify experimental conditions resulted in <10%

increase to the total reaction volume. Aggregates from all proteins were monitored at designated

time points, with complete formation typically occurring within 5 hours.

Formation of liquid-like condensates

Liquid-like condensates were generated from either soluble proteins or pre-formed solid

aggregates by adding 1% SDS directly to the reaction mixture. The transition typically completed

within 8 hours. The final SDS concentration was 0.03%, consistent across all stated methods.

In experiments where glycine was added to probe the effects of addition, the final concentration

of glycine was 50 mM for experiments shown in Fig. 2c, and 5mM in Fig. 4a. Concentrated glycine

was diluted directly to the experimental conditions.

To quantify droplet shape characteristics, three geometric descriptors were calculated from binary

images: circularity, roundness, and solidity.

Circularity was calculated using the formula:

Circularity = $(4 \times \pi \times \text{Area}) / (\text{Perimeter}^2)$

This metric ranges from 0 to 1, where a value of 1 indicates a perfect circle. Lower values suggest

more elongated or irregular shapes.

Roundness was defined as:

Roundness = $(4 \times \text{Area}) / (\pi \times (\text{Major Axis Length})^2)$

Roundness also ranges from 0 to 1, with higher values indicating shapes closer to a perfect circle,

but it is more sensitive to elongation than circularity.

Solidity was computed as:

583 Solidity = Area / Convex Hull Area

584 Solidity measures the extent to which a shape is convex. A value of 1 indicates a completely

585 convex shape, while lower values suggest the presence of concavities or irregular boundaries.

586 All measurements were performed using image analysis software capable of extracting shape

587 descriptors from segmented droplet images.

589 **Imaging**

590 Microscopy imaging included bright field, phase contrast, widefield fluorescence, and confocal

591 fluorescence modalities. The primary instrument was the Keyence BZ-X800 fluorescence

592 microscope for both light and fluorescence imaging. Additional bright-field and widefield

593 fluorescence images were captured using the EVOS FL Fluorescence Digital Microscope, while

594 confocal fluorescence imaging was performed with the BioTek Cytation C10 Confocal Imaging

595 Reader. Images taken by Cytation C10 were at 40x magnification. All other images were taken at

596 20x magnification.

598 **Image analysis and quantifications**

599 Image contrast and brightness were adjusted uniformly for clarity using ImageJ. Quantitative

600 measurements were also performed using ImageJ's built-in analysis functions.

602 For quantifications of fluorescence images, ImageJ's color threshold function was used to

603 delineate positive signals from background signal. The analysis function provided a count of

604 detected structure, the area covered by structures of the total visible area Area%, as well as

605 fluorescence intensity IntDen.

607 IntDen or Integrated Density is the product of mean gray value (a stand-in for intensity) and area,

608 representing the amount of dye bound and fluorescing. The units are (relative fluorescence units)

* (μm^2). Corrected Total View Fluorescence is used to represent the intensity of fluorescence in

our measurements, which takes the IntDen value from each measured object and corrects for

background fluorescence. The equation is as follows:

CTFV = IntDen – (Area of Droplet * Mean fluorescence of background)

The mean fluorescence of background was measured by averaging 3 separate readings of the

background for each time point and view of the well.

Dye staining

Protein assemblies were stained by incubating at room temperature for 2–3 hours with Thioflavin

T (ThT) or Congo Red (CR) from Sigma. Final concentrations in wells were 40 μM ThT or 0.5%

CR. For time course experiments shown in Fig. 4, ThT was added to the solution before the phase

separation was induced.

SpyTag/SpyCatcher Conjugation

Protein constructs were conjugated by incubating together at room temperature for 1 hour. FP–

SpyCatcher was added in slight excess (10%) over TDP/CTD/hnRNPH1–SpyTag to achieve a

1.1:1 molar ratio. For quenching, excess SpyTag-tagged nanobodies (at least 4× the molarity of

SpyCatcher–FP) were added to bind any remaining free SpyCatcher–FP, followed by an

additional 1-hour incubation at room temperature.

Acknowledgment

We thank Alex Ward for some of the initial DNA constructions and initial attempts at protein

production, Andrew Chammas and Nathan Shaner for help with designing of SpyTag and

SpyCatcher constructs, Aryan Mudliar for lab assistance and discussions.

Author Contributions

JW conceptually initiated the project; TRW designed and carried out the experiments; TRW and

JW interpreted data and drafted the manuscript.

Competing Interest

JW is a current employee of Allele Biotech, a biotech company in the US.

Data Availability

Additional, related data of the present report is available from the corresponding author upon

reasonable request.

Supplementary Material

Supplementary material.doc: Supplementary figures, materials and methods

References

Polymenidou, M. *et al.* Long pre-mRNA depletion and RNA missplicing contribute to neuronal vulnerability from loss of TDP-43. *Nat Neurosci* **14**, 459-468 (2011). https://doi.org/10.1038/nn.2779

Ishiguro, T. *et al.* Regulatory Role of RNA Chaperone TDP-43 for RNA Misfolding and Repeat-Associated Translation in SCA31. *Neuron* **94**, 108-124 e107 (2017). https://doi.org/10.1016/j.neuron.2017.02.046

Gasset-Rosa, F. *et al.* Cytoplasmic TDP-43 De-mixing Independent of Stress Granules Drives Inhibition of Nuclear Import, Loss of Nuclear TDP-43, and Cell Death. *Neuron* **102**, 339-357 e337 (2019). https://doi.org/10.1016/j.neuron.2019.02.038

Ayala, Y. M. *et al.* Structural determinants of the cellular localization and shuttling of TDP-43. *J Cell Sci* **121**, 3778-3785 (2008). https://doi.org/10.1242/jcs.038950

Shiina, Y., Arima, K., Tabunoki, H. & Satoh, J. TDP-43 dimerizes in human cells in culture. *Cell Mol Neurobiol* **30**, 641-652 (2010). https://doi.org/10.1007/s10571-009-9489-9

Perez-Berlanga, M. *et al.* Loss of TDP-43 oligomerization or RNA binding elicits distinct aggregation patterns. *EMBO J* **42**, e111719 (2023). https://doi.org/10.15252/embj.2022111719

7   Wang, A. *et al.* A single N-terminal phosphomimic disrupts TDP-43 polymerization, phase
separation, and RNA splicing. *EMBO J* **37** (2018).
https://doi.org/10.15252/embj.201797452

8   Santamaria, N., Alhothali, M., Alfonso, M. H., Breydo, L. & Uversky, V. N. Intrinsic disorder
in proteins involved in amyotrophic lateral sclerosis. *Cell Mol Life Sci* **74**, 1297-1318
(2017). https://doi.org/10.1007/s00018-016-2416-6

9   Buratti, E. *et al.* TDP-43 binds heterogeneous nuclear ribonucleoprotein A/B through its
C-terminal tail: an important region for the inhibition of cystic fibrosis transmembrane
conductance regulator exon 9 splicing. *J Biol Chem* **280**, 37572-37584 (2005).
https://doi.org/10.1074/jbc.M505557200

10   Jo, M. *et al.* The role of TDP-43 propagation in neurodegenerative diseases: integrating
insights from clinical and experimental studies. *Exp Mol Med* **52**, 1652-1662 (2020).
https://doi.org/10.1038/s12276-020-00513-7

11   Lang, R., Hodgson, R. E. & Shelkovnikova, T. A. TDP-43 in nuclear condensates: where,
how, and why. *Biochem Soc Trans* **52**, 1809-1825 (2024).
https://doi.org/10.1042/BST20231447

12   Lee, Y. B. *et al.* Cytoplasmic TDP-43 is involved in cell fate during stress recovery. *Hum Mol*
Genet **31**, 166-175 (2021). https://doi.org/10.1093/hmg/ddab227

13   Chen, Y. & Cohen, T. J. Aggregation of the nucleic acid-binding protein TDP-43 occurs via
distinct routes that are coordinated with stress granule formation. *J Biol Chem* **294**, 3696-
3706 (2019). https://doi.org/10.1074/jbc.RA118.006351

14   Conicella, A. E. *et al.* TDP-43 alpha-helical structure tunes liquid-liquid phase separation
and function. *Proc Natl Acad Sci U S A* **117**, 5883-5894 (2020).
https://doi.org/10.1073/pnas.1912055117

15   McGurk, L. *et al.* Poly(ADP-Ribose) Prevents Pathological Phase Separation of TDP-43 by
Promoting Liquid Demixing and Stress Granule Localization. *Mol Cell* **71**, 703-717 e709
(2018). https://doi.org/10.1016/j.molcel.2018.07.002

16   Fonda, B. D., Jami, K. M., Boulos, N. R. & Murray, D. T. Identification of the Rigid Core for
Aged Liquid Droplets of an RNA-Binding Protein Low Complexity Domain. *Journal of the*
American Chemical Society **143**, 6657-6668 (2021). https://doi.org/10.1021/jacs.1c02424

17   Lin, Y., Protter, D. S., Rosen, M. K. & Parker, R. Formation and Maturation of Phase-
Separated Liquid Droplets by RNA-Binding Proteins. *Mol Cell* **60**, 208-219 (2015).
https://doi.org/10.1016/j.molcel.2015.08.018

18   Shin, Y. & Brangwynne, C. P. Liquid phase condensation in cell physiology and disease.
Science **357** (2017). https://doi.org/10.1126/science.aaf4382

19   Johnson, B. S. *et al.* TDP-43 is intrinsically aggregation-prone, and amyotrophic lateral
sclerosis-linked mutations accelerate aggregation and increase toxicity. *J Biol Chem* **284**,
20329-20339 (2009). https://doi.org/10.1074/jbc.M109.010264

20   François-Moutal, L. *et al.* Structural Insights Into TDP-43 and Effects of Post-translational
Modifications. *Frontiers in Molecular Neuroscience* **12** (2019).
https://doi.org/10.3389/fnmol.2019.00301

21   Wang, J. *et al.* A Molecular Grammar Governing the Driving Forces for Phase Separation
of Prion-like RNA Binding Proteins. *Cell* **174**, 688-699 e616 (2018).
https://doi.org/10.1016/j.cell.2018.06.006

22   Chang, L. W. *et al.* Sequence and entropy-based control of complex coacervates. *Nat*
Commun **8**, 1273 (2017). https://doi.org/10.1038/s41467-017-01249-1

23   Millar, S. R. *et al.* A New Phase of Networking: The Molecular Composition and Regulatory
Dynamics of Mammalian Stress Granules. *Chem Rev* **123**, 9036-9064 (2023).
https://doi.org/10.1021/acs.chemrev.2c00608

24   Schmidt, H. B. & Rohatgi, R. In Vivo Formation of Vacuolated Multi-phase Compartments
Lacking Membranes. *Cell Rep* **16**, 1228-1236 (2016).
https://doi.org/10.1016/j.celrep.2016.06.088

25   Erkamp, N. A. *et al.* Spatially non-uniform condensates emerge from dynamically arrested
phase separation. *Nature Communications* **14** (2023). https://doi.org/10.1038/s41467-
023-36059-1

26   Conicella, A. E., Zerze, G. H., Mittal, J. & Fawzi, N. L. ALS Mutations Disrupt Phase
Separation Mediated by alpha-Helical Structure in the TDP-43 Low-Complexity C-Terminal
Domain. *Structure* **24**, 1537-1549 (2016). https://doi.org/10.1016/j.str.2016.07.007

27   Babinchak, W. M. *et al.* The role of liquid-liquid phase separation in aggregation of the
TDP-43 low-complexity domain. *J Biol Chem* **294**, 6306-6317 (2019).
https://doi.org/10.1074/jbc.RA118.007222

28   Kumar, S. T. *et al.* Seeding the aggregation of TDP-43 requires post-fibrillization
proteolytic cleavage. *Nature Neuroscience* **26**, 983-996 (2023).
https://doi.org/10.1038/s41593-023-01341-4

29   Staderini, T., Bigi, A., Mongiello, D., Cecchi, C. & Chiti, F. Biophysical characterization of
full-length TAR DNA-binding protein (TDP-43) phase separation—solution-jump. *Protein*
Sci **31**, e4509 (2022). https://doi.org/10.1002/pro.4509

30   Liu, Y. *et al.* The Regulation of TDP-43 Structure and Phase Transitions: A Review. *Protein*
J **44**, 113-132 (2025). https://doi.org/10.1007/s10930-025-10261-0

31   Yang, L., Jasiqi, Y. & Lashuel, H. Recombinant Full-Length TDP-43 Oligomers Retain Their
Ability to Bind RNAs, Are Not Toxic, and Do Not Seed TDP-43 Aggregation in Vitro. *ACS*
Chem Neurosci **15**, 193-204 (2024). https://doi.org/10.1021/acschemneuro.3c00691

32   Staderini, T. *et al.* Biophysical characterization of the phase separation of TDP-43 devoid
of the C-terminal domain. *Cell Mol Biol Lett* **29**, 104 (2024).
https://doi.org/10.1186/s11658-024-00615-4

33   Gruijs Da Silva, L. A. *et al.* Disease‐linked TDP‐43 hyperphosphorylation suppresses
TDP‐43 condensation and aggregation. *The EMBO Journal* **41** (2022).
https://doi.org/10.15252/embj.2021108443

34   Mann, J. R. *et al.* RNA Binding Antagonizes Neurotoxic Phase Transitions of TDP-43.
Neuron **102**, 321-338 e328 (2019). https://doi.org/10.1016/j.neuron.2019.01.048

35   Molliex, A. *et al.* Phase separation by low complexity domains promotes stress granule
assembly and drives pathological fibrillization. *Cell* **163**, 123-133 (2015).
https://doi.org/10.1016/j.cell.2015.09.015

36   Vivoli Vega, M. *et al.* Isolation and characterization of soluble human full-length TDP-43
associated with neurodegeneration. *FASEB J* **33**, 10780-10793 (2019).
https://doi.org/10.1096/fj.201900474R

37 Krainer, G. *et al.* Reentrant liquid condensate phase of proteins is stabilized by
hydrophobic and non-ionic interactions. *Nature Communications* **12** (2021).
https://doi.org/10.1038/s41467-021-21181-9

38 Lu, T. & Spruijt, E. Multiphase Complex Coacervate Droplets. *J Am Chem Soc* **142**, 2905-
2914 (2020). https://doi.org/10.1021/jacs.9b11468

39 Garg, D. K. & Bhat, R. Modulation of assembly of TDP-43 low-complexity domain by
heparin: From droplets to amyloid fibrils. *Biophysical Journal* **121**, 2568-2582 (2022).
https://doi.org/10.1016/j.bpj.2022.05.042

40 Wang, C. *et al.* Stress Induces Dynamic, Cytotoxicity-Antagonizing TDP-43 Nuclear Bodies
via Paraspeckle LncRNA NEAT1-Mediated Liquid-Liquid Phase Separation. *Mol Cell* **79**,
443-458 e447 (2020). https://doi.org/10.1016/j.molcel.2020.06.019

41 Gui, X. *et al.* Liquid-liquid phase separation of amyloid-beta oligomers modulates amyloid
fibrils formation. *J Biol Chem* **299**, 102926 (2023).
https://doi.org/10.1016/j.jbc.2023.102926

42 Xu, X. *et al.* Amino acids bind to phase-separating proteins and modulate biomolecular
condensate stability and dynamics. *Nature Communications* **16** (2025).
https://doi.org/10.1038/s41467-025-63755-x

43 Kim, G. H. & Kwon, I. Distinct roles of hnRNPH1 low-complexity domains in splicing and
transcription. *Proc Natl Acad Sci U S A* **118** (2021).
https://doi.org/10.1073/pnas.2109668118

44 Neumann, M. *et al.* Ubiquitinated TDP-43 in frontotemporal lobar degeneration and
amyotrophic lateral sclerosis. *Science* **314**, 130-133 (2006).
https://doi.org/10.1126/science.1134108

45 Zhang, Y. J. *et al.* Aberrant cleavage of TDP-43 enhances aggregation and cellular toxicity.
Proc Natl Acad Sci U S A **106**, 7607-7612 (2009).
https://doi.org/10.1073/pnas.0900688106

46 Li, H. R. *et al.* The physical forces mediating self-association and phase-separation in the
C-terminal domain of TDP-43. *Biochim Biophys Acta Proteins Proteom* **1866**, 214-223
(2018). https://doi.org/10.1016/j.bbapap.2017.10.001

47 Prasad, A., Sivalingam, V., Bharathi, V., Girdhar, A. & Patel, B. K. The amyloidogenicity of
a C-terminal region of TDP-43 implicated in Amyotrophic Lateral Sclerosis can be affected
by anions, acetylation and homodimerization. *Biochimie* **150**, 76-87 (2018).
https://doi.org/10.1016/j.biochi.2018.05.003

48 Guo, W. *et al.* An ALS-associated mutation affecting TDP-43 enhances protein
aggregation, fibril formation and neurotoxicity. *Nat Struct Mol Biol* **18**, 822-830 (2011).
https://doi.org/10.1038/nsmb.2053

49 Jiang, L.-L. *et al.* Structural Transformation of the Amyloidogenic Core Region of TDP-43
Protein Initiates Its Aggregation and Cytoplasmic Inclusion. *Journal of Biological Chemistry*
288, 19614-19624 (2013). https://doi.org/10.1074/jbc.m113.463828

50 Pakravan, D. *et al.* Liquid-Liquid Phase Separation Enhances TDP-43 LCD Aggregation but
Delays Seeded Aggregation. *Biomolecules* **11** (2021).
https://doi.org/10.3390/biom11040548

51 Rollin, R., Joanny, J. F. & Sens, P. Physical basis of the cell size scaling laws. *Elife* **12** (2023).
https://doi.org/10.7554/eLife.82490

52   Mao, T. *et al.* Stabilizing effect of amino acids on protein and colloidal dispersions. *Nature*
645, 915-921 (2025). https://doi.org/10.1038/s41586-025-09506-w
53   Sun, Y. *et al.* Physiologically Important Electrolytes as Regulators of TDP-43 Aggregation
and Droplet-Phase Behavior. *Biochemistry* **58**, 590-607 (2019).
https://doi.org/10.1021/acs.biochem.8b00842
54   Capitini, C. *et al.* Full-length TDP-43 and its C-terminal domain form filaments in vitro
having non-amyloid properties. *Amyloid* **28**, 56-65 (2021).
https://doi.org/10.1080/13506129.2020.1826425
55   Lin, P. H. *et al.* TDP-43 Amyloid Fibril Formation via Phase Separation-Related and -
Unrelated Pathways. *ACS Chem Neurosci* **15**, 3767-3775 (2024).
https://doi.org/10.1021/acschemneuro.4c00503
56   Esposto, J. C. & Martic, S. Phosphorylated TAR DNA-binding protein-43: Aggregation and
antibody-based inhibition. *Biochim Biophys Acta Mol Basis Dis* **1867**, 166234 (2021).
https://doi.org/10.1016/j.bbadis.2021.166234
57   Shaner, N. C. *et al.* A bright monomeric green fluorescent protein derived from
Branchiostoma lanceolatum. *Nature Methods* **10**, 407-409 (2013).
58   Gadella, T. W. J., Jr. *et al.* mScarlet3: a brilliant and fast-maturing red fluorescent protein.
Nat Methods **20**, 541-545 (2023). https://doi.org/10.1038/s41592-023-01809-y
59   Goedhart, J. *et al.* Structure-guided evolution of cyan fluorescent proteins towards a
quantum yield of 93%. *Nat Commun* **3**, 751 (2012). https://doi.org/10.1038/ncomms1738
60   Schmidt, H. B., Barreau, A. & Rohatgi, R. Phase separation-deficient TDP43 remains
functional in splicing. *Nat Commun* **10**, 4890 (2019). https://doi.org/10.1038/s41467-019-
12740-2
61   Nakagawa, C., Nishimura, S., Senda-Murata, K. & Sugimoto, K. A rapid and simple method
of evaluating the dimeric tendency of fluorescent proteins in living cells using a truncated
protein of importin alpha as fusion tag. *Biosci Biotechnol Biochem* **76**, 388-390 (2012).
https://doi.org/10.1271/bbb.110677
62   Sawaya, M. R., Hughes, M. P., Rodriguez, J. A., Riek, R. & Eisenberg, D. S. The expanding
amyloid family: Structure, stability, function, and pathogenesis. *Cell* **184**, 4857-4873
(2021). https://doi.org/10.1016/j.cell.2021.08.013
63   Mohanty, P. *et al.* A synergy between site-specific and transient interactions drives the
phase separation of a disordered, low-complexity domain. *Proc Natl Acad Sci U S A* **120**,
e2305625120 (2023). https://doi.org/10.1073/pnas.2305625120
64   Dubinski, A. *et al.* Stress granule assembly<i>in vivo</i>is deficient in the CNS of mutant
TDP-43 ALS mice. *Human Molecular Genetics* **32**, 319-332 (2023).
https://doi.org/10.1093/hmg/ddac206
65   Yan, X. *et al.* Intra-condensate demixing of TDP-43 inside stress granules generates
pathological aggregates. *Cell* **188**, 4123-4140 e4118 (2025).
https://doi.org/10.1016/j.cell.2025.04.039
66   Wang, J. & Bell, L. R. The Sex-lethal amino terminus mediates cooperative interactions in
RNA binding and is essential for splicing regulation. *Genes Dev* **8**, 2072-2085 (1994).
https://doi.org/10.1101/gad.8.17.2072

67   Wang, J., Dong, Z. & Bell, L. R. Sex-lethal interactions with protein and RNA. Roles of
839   glycine-rich and RNA binding domains. *J Biol Chem* **272**, 22227-22235 (1997).
840   https://doi.org/10.1074/jbc.272.35.22227
841   68   Ryan, V. H. *et al.* Mechanistic View of hnRNPA2 Low-Complexity Domain Structure,
842   Interactions, and Phase Separation Altered by Mutation and Arginine Methylation. *Mol*
843   *Cell* **69**, 465-479 e467 (2018). https://doi.org/10.1016/j.molcel.2017.12.022
844   69   Freibaum, B. D., Chitta, R. K., High, A. A. & Taylor, J. P. Global analysis of TDP-43 interacting
845   proteins reveals strong association with RNA splicing and translation machinery. *J*
846   *Proteome Res* **9**, 1104-1120 (2010). https://doi.org/10.1021/pr901076y
847   70   Sear, R. P. Dishevelled: a protein that functions in living cells by phase separating. *Soft*
848   *Matter* **3**, 680-684 (2007). https://doi.org/10.1039/b618126k
849   71   Kang, K., Shi, Q., Wang, X. & Chen, Y.-G. Dishevelled phase separation promotes Wnt
850   signalosome assembly and destruction complex disassembly. *Journal of Cell Biology* **221**
851   (2022). https://doi.org/10.1083/jcb.202205069
852   72   Wang, F. & Zhang, Y. Physiology and pharmacological targeting of phase separation. *J*
853   *Biomed Sci* **31**, 11 (2024). https://doi.org/10.1186/s12929-024-00993-z
854

Figures

t = 5 min   t = 30 min   t = 60 min   t = 240 min   t = 360 min

a.

t = 0 min   t = 20 min   t = 60 min

b.

c.

t = 120 min   t = 240 min   t = 410 min

t = 0 min   t = 360 min   t = 390 min

d.

t = 420 min   t = 450 min   t = 480 min

e.

-SDS

Aggregation
Favorable pH
Salt

+SDS

+SDS

Figure 1. Three-phase transition of recombinant full-length TDP-43 induced by charge

modulation

a Gradual formation of solid aggregates from soluble TDP-43 at pH 6 and 0.5 M NaCl. Scale bar

is 40μm. **b** Solid aggregates converted into a hydrogel upon addition of SDS (final concentration

0.03%), then progressively transformed into spherical droplets over 7 hours. Time notations begin

at the addition of SDS. Scale bar is 40μm. **c** The diagram illustrates the time-dependent

morphological changes measured by circularity, roundness, and solidity (see Materials and

Methods). Error bars are Standard Errors of the Mean. **d** Droplets formed directly from solution

following SDS addition, with the most pronounced transition occurring between the 6th and 8th

hour. Scale bar is 20μm. **e** Schematic representation showing how TDP-43 can transition among

all three phases through controlled induction.

Figure 2. Droplets induced from solid aggregates exhibit liquid-like properties

a Time-lapse images showing a small droplet (arrow) gradually merging with a larger droplet. Scale bar is 10µm. **b** Droplets derived from aggregates dissolve upon treatment with 1,6-hexanediol, consistent with typical LLPS behavior. Scale bar is 40µm. **c** Formation of multi-emulsion droplets upon addition of glycine (50 mM) to initial aggregate-derived droplets. Time indications begin at addition of glycine. Scale bar is 20µm. The transition occurs on the scale of minutes, progressing from small droplets (left) to multiple inner droplets encapsulated within larger outer droplets (middle), and finally to fused, larger droplets (right). **d** Graph showing the number of internal droplets (Y-axis) versus the size of the encompassing droplet (X-axis) and time in seconds (color legend). **e** Graph displaying changes in average number of internal droplets per encompassing droplet (Y-axis) over time in seconds (X-axis). Error bars are Standard Errors of the Mean.

Figure 3. TDP-43, TDP-CTD, and hnRNPH1 form amyloid-positive aggregates and droplets

a Side-by-side comparison of solid aggregates formed by inducing soluble TDP-43, TDP-CTD, and hnRNPH1 at pH 6.0 and 0.5 M NaCl (top), and droplets formed after further addition of SDS to a final concentration of 0.03% (bottom). Images were taken 8 hours after induction. Scale bar

is 20µm. **b-d** Aggregates formed by TDP-43 (**b**) and TDP-CTD (**c**) stain positive with ThT and

Congo Red (top: bright field; bottom: fluorescence images). **d** hnRNPH1 also stains positive for

ThT. Scale bar is 40µm. **e** Representative data showing droplets formed directly from soluble

TDP-43 and TDP-CTD solutions exhibit ThT staining. Scale bar is 40µm. **f** Multi-emulsion droplet

formed upon addition of glycine (50 mM), with ThT included; shown are confocal fluorescence

images from a representative Z-stack acquired using Cytation10 (40×). The z-distance between

the two slices is 20µm. Scale bar is 100µm.

Figure 4. Time course of amyloid-containing droplet and solid aggregate formation

a Still photos taken at indicated time points from a time-lapse series of fluorescence image capturing droplets formed from soluble TDP-43. ThT was added to protein solution before addition of SDS (0.03% final concentration) at time 0. Arrows point to example droplets in the merging process. Scale bar is 100μm. **b** Quantitative analysis of count, percentage of view covered, and Corrected Total View Fluorescence (CTVF) of the droplets formed from 3 separate, non-overlapping views of the same time lapse course. Droplet count and area covered are averaged from the 3 views while CTVF is presented from each time point and view. **c** Still photos taken at indicated time points from a time-lapse series of fluorescence image capturing solid aggregates from soluble solution of TDP-43. ThT was added to protein solution before addition of 1M Tris-HCl pH6 and 5M NaCl at 10% final volume at time 0. Scale bars are 50μm. All reactions were carried out at room temperature, ~25°C. **d** Quantitative analysis of view covered and fluorescence intensity of the solid aggregates formed. Data came from 1 representative view of the time lapse.

Figure 5. Labeling phase-transition proteins with switchable fluorescent tags

a Schematic representation of protein domains fused with SpyTag/SpyCatcher modules. **b** Solid aggregates formed by TDP-43–mScarlet3 (left), TDP-CTD–mNeonGreen (middle), and hnRNPH1–mTurquoise2 (right) under SDS induction conditions. Scale bar is 40μm. **c, d** Time-lapse images showing TDP-43–mScarlet3 (**c**) and TDP-CTD–mNeonGreen (**d**) transitioning from solid aggregates into droplets; the corresponding bright-field image for the final fluorescence frame is shown on the right. Time notations begin after addition of SDS. Scale bars are 10μm.

TDP-CTD-mNG **TDP-43-mSCI3** **TDP-CTD and TDP-43 Overlay**

992 **TDP-CTD-mNG** **hnRNPH1-mTq2** **TDP-CTD and hnRNPH1 Overlay**

997 **TDP-43-mSCI3** **hnRNPH1-mTq2** **TDP-43 and hnRNPH1 Overlay**

1000

1001

1002

1003 **Figure 6. Seeding of phase transition depends on protein compatibility**

1004 Pairwise mixing of TDP-CTD–mNeonGreen (mNG), TDP-43–mScarlet3 (mSCI3), and

1005 hnRNPH1–mTurquoise2 (mTq2) in solution prior to induction to form aggregates. Coaggregates

1006 between TDP-CTD–mNeonGreen and TDP-43–mScarlet3 appear as yellow-orange due to

1007 fluorescence overlap. Scale bar is 20µm.

1008