Multiparametric MRI and imaging transcriptomics reveal molecular and cellular correlates of neurodegeneration in experimental parkinsonism

Eugene Kim[1], Diana Cash[1], Daniel Martins[1,2,3], Camilla Simmons[1,4], Antonio Heras-Garvin[5], Elena Klippel[5], Florian Krismer[5], Laura Mantoan Ritter[6,7]*, Nadia Stefanova[5]*#

1. The Brain Centre, Department of Neuroimaging; Institute of Psychiatry, Psychology, and Neuroscience; King's College London, London, United Kingdom
2. Department of Clinical Neurosciences and Mental Health, Faculty of Medicine, University of Porto, Porto, Portugal
3. RISE-Health (Neurosciences), Faculty of Medicine, University of Porto, Porto, Portugal
4. Sygnature Discovery, Nottingham, United Kingdom
5. Division of Neurobiology, Department of Neurology, Medical University Innsbruck, Innsbruck, Austria
6. Maurice Wohl Clinical Neuroscience Institute, Institute of Psychiatry, Psychology and Neuroscience; King's College London, London, United Kingdom
7. Clinical Neurosciences Department, King's College Hospital, London, United Kingdom

*Contributed equally

#corresponding author: Nadia Stefanova, Division of Neurobiology, Department of Neurology, Medical University of Innsbruck, Innrain 66/G3, 6020 Innsbruck, Austria, E-mail: nadia.stefanova@i-med.ac.at; ORCID: 0000-0001-8188-639X

ABSTRACT

Multiple system atrophy (MSA) is an atypical Parkinsonian disorder marked by oligodendroglial α-synucleinopathy and selective neurodegeneration. Although MRI can capture regional atrophy and microstructural alterations in MSA brain, the molecular substrates underlying these phenotypes remain poorly defined. Imaging transcriptomics provides a computational framework to relate spatial imaging patterns to brain-wide gene expression. While this approach has been applied to human MSA, interpretation is constrained by limited experimental control and lack of disease-matched molecular validation. Here, we apply imaging transcriptomics in a controlled preclinical setting by integrating high-resolution ex vivo multimodal MRI with transcriptomic mapping in the PLP-αSyn mouse model of MSA. Structural and diffusion MRI revealed distinct patterns of regional atrophy and microstructural abnormalities. Atlas-based analyses associated imaging phenotypes to gene programs related to oligodendrocyte biology, energy metabolism, and neuroinflammation, with modality-specific signatures. These associations were supported by independent RNA-sequencing and showed convergence with human MSA findings. Our work benchmarks MRI–transcriptomic relationships in MSA and provides a translational framework for interpreting imaging biomarkers in synucleinopathies.

Keywords

INTRODUCTION

Multiple system atrophy (MSA) is a rapidly progressive neurodegenerative disorder clinically characterized by autonomic failure, parkinsonism, and cerebellar ataxia. Its etiology remains unclear, diagnosis is challenging, and no effective disease-modifying therapy is currently available [1]. MSA is classified as an α-synucleinopathy due to the hallmark accumulation of α-synuclein (α-syn) fibrils within oligodendrocytes, forming glial cytoplasmic inclusions (GCIs) [2]. At the systems level, MSA is characterized by regionally selective vulnerability that likely reflects complex interactions across molecular-, cellular-, and network-scale processes. Magnetic resonance imaging (MRI) provides a powerful, non-invasive window into these processes by capturing macroscopic and microstructural brain alterations in vivo (**Table 1**). However, while MRI phenotypes such as atrophy or diffusion abnormalities are sensitive markers of disease progression [3, 4], they lack specificity and the molecular and cellular substrates underlying these imaging changes remain incompletely characterized. To address this limitation, imaging transcriptomics has emerged as a framework for linking spatial patterns of neuroimaging phenotypes to normative brain-wide gene expression maps derived from resources such as the Allen Brain Atlas. Rather than measuring disease-state transcription directly, imaging transcriptomics exploits the correspondence between the spatial organization of normative gene expression and the regional distribution of disease-related imaging alterations to identify molecular and cellular programs associated with regional vulnerability across the brain. This approach has been applied extensively in humans to interpret MRI-derived cortical thinning, volumetric atrophy, and functional alterations in neurodegenerative disorders, including Alzheimer's disease and Parkinsonian syndromes, where imaging-derived effect maps are correlated with transcriptomic gradients to infer biological pathways and cell-type contributions underlying disease-specific patterns of neurodegeneration [5-7]. More recently, imaging transcriptomics has been applied to MSA in humans. In a landmark study, Chougar and colleagues [8] correlated regional atrophy maps derived from structural MRI in patients with MSA with gene expression data from the Allen Human Brain Atlas. Using multivariate statistical approaches, the authors identified molecular

signatures enriched for oligodendrocyte-related and mitochondrial pathways that spatially

aligned with MSA-specific atrophy patterns, providing new insights into the biological

programs associated with selective regional vulnerability in this disorder. Importantly, this

work demonstrated the feasibility and interpretive power of imaging transcriptomics in a rare

and rapidly progressive neurodegenerative disease. At the same time, it is necessary to

recognize the intrinsic limitations of imaging transcriptomics studies in clinical populations.

The reliance on postmortem normative transcriptomic atlases precludes direct assessment of

disease-state gene expression, and inter-individual variability, disease heterogeneity, and

limited experimental control restrict mechanistic validation. In the context of MSA, these

constraints make it difficult to disentangle whether imaging transcriptomic associations reflect

primary drivers of neurodegeneration, downstream consequences, or compensatory

responses. Moreover, validation of transcriptomic inferences at the tissue or cellular level is

typically not feasible in human cohorts.

Experimental animal models offer a unique opportunity to address these limitations. The

proteolipid protein (PLP)-αSyn transgenic mouse, which overexpresses human α-syn in

oligodendrocytes [9], recapitulates key pathological features of MSA, including early

oligodendroglial dysfunction and neurotrophic deficits, progressive α-syn aggregation, myelin

dysfunction, gliosis, and selective neurodegeneration associated with progressive motor and

non-motor deficits [10, 11]. Crucially, this model enables controlled, whole-brain MRI

phenotyping, direct access to tissue for molecular validation, and integration of imaging

findings with independently acquired transcriptomic data from the same disease context.

Applying imaging transcriptomics in the PLP-αSyn mouse therefore serves several

complementary purposes. First, it provides a preclinical testbed for evaluating whether

imaging transcriptomic associations observed in human MSA are conserved across species

and disease contexts. Second, it allows assessment of how different MRI modalities—such

as volumetric and diffusion-based measures—map onto partially distinct molecular and

cellular programs within a single, well-characterized model. Third, the availability of model-

specific RNAseq data enables empirical evaluation of whether atlas-derived transcriptomic

associations align with disease-relevant gene expression changes, thereby strengthening

biological interpretability beyond what is possible in human studies alone. Importantly, the

goal of imaging transcriptomics in the animal model is not to assert direct causality between

gene expression and MRI phenotypes, but rather to identify molecular correlates of regional

imaging vulnerability that can guide mechanistic hypotheses and targeted validation.

Establishing these relationships at advanced stages of disease can provide a critical

benchmark for interpreting earlier and subtler imaging changes, and will lay the groundwork

for future studies focused on prodromal biomarkers and longitudinal progression.

In this study, we combine high-resolution ex-vivo structural and diffusion MRI with imaging-

transcriptomics analysis in the PLP-αSyn mouse model of MSA. By integrating multimodal

MRI phenotypes with more than 4,000 normative gene expression maps from the Allen Brain

Atlas, followed by gene enrichment analyses to determine the associated biological

processes and cell types [7, 12, 13] and validating key findings using model-specific RNAseq

data, we aim to delineate the molecular and cellular programs associated with MRI-detected

neurodegeneration and microstructural alterations in the MSA mouse model. This work

extends imaging transcriptomics into a controlled experimental setting, complements recent

human studies, and establishes a translatable framework for mechanistically informed

interpretation of MRI biomarkers in synucleinopathies.

RESULTS

Multiparametric MRI identifies atrophy and microstructural alterations in the MSA

mouse brains

Ex vivo deformation-based morphometry revealed widespread brain atrophy in PLP-αSyn

mice compared with age-matched WT controls (**Figure 1a; Suppl. Fig. S2a**). Significant age

× genotype interactions were detected in some regions; therefore, post hoc comparisons

were performed. At 10 months, PLP-αSyn mice exhibited pronounced volume loss in the

midbrain, hindbrain, and cerebellum, accompanied by ventricular enlargement, consistent

with neurodegeneration. Conversely, localized volume increases were observed in regions of

the cortex, amygdala, and striatum (**Suppl. Fig. S2a**). By 15 months, atrophy extended to

additional brain areas, and no regions displayed volume gain (**Suppl. Fig. S2b**). This pattern

was at least partly associated with an age-related volume increase in WT mice (**Suppl. Fig.**

S2c).

Given the limited group sizes and the absence of age or age × genotype effects in the

diffusion metrics (FA and MD), subsequent analyses focused on the main genotype effect.

Significant atrophy was identified in multiple motor-related structures, including the pons,

extrapyramidal fiber tracts, medulla, pedunculopontine tegmental nucleus, and cerebellum

(**Table 2; Figure 2a**). Non-motor regions implicated in sleep regulation, autonomic control,

anxiety, and pain processing—such as the epithalamus, ventral tegmental area, raphe nuclei,

insular cortex, reticular nucleus, and intralaminar thalamic nuclei—also showed reduced

volume (**Suppl. Fig. S3a**).

FA was reduced across widespread white-matter tracts (**Table 2; Suppl. Fig. S3b**), including

interhemispheric pathways (anterior commissure, corpus callosum), descending motor tracts

(internal capsule, cerebellar peduncles), and fibers associated with non-motor systems (optic

nerve, olfactory bulb, epithalamus, stria terminalis) (**Table 2; Figure 2b**). In contrast, MD was

globally increased, with the most significant elevations in the brainstem and cerebellum

(**Table 2; Figure 2c; Suppl. Fig. S3c**).

Imaging transcriptomics highlights associations between MRI phenotypes and gene

expression signatures of myelination, gliogenesis, neuroinflammation, and neural

network regulation

To relate MRI changes to underlying molecular mechanisms, we performed imaging

transcriptomic analyses, examining spatial correlations between MRI-derived t-statistic maps

(volume, FA, MD) and normative gene expression patterns from the Allen Brain Atlas. We

identified 3,214 genes significantly correlated with regional volume changes ($p_{adj} < 0.05$),

1,850 with FA, and 2,334 with MD (**Suppl. Tables S1–S3**). All correlations exceeded those

expected by chance (p = 0 vs. 5,000 BrainSMASH surrogates). The top 30 volume- and top

55 FA-correlated genes were predominantly negatively associated with the corresponding

MRI features, while among the top 60 MD-correlated genes, 20 showed positive associations

(**Suppl. Tables S1–S3**).

Genes most strongly correlated with atrophy included *Arhgef10* (axon guidance, nerve

conduction), *Cldn11* (axon ensheathment), *Tspan2* (oligodendrocyte differentiation), *Aspa* (N-

acetylaspartate metabolism relevant to myelin synthesis), *Cnp* (myelinating cell RNA

regulation), *Ugt8a* (galactocerebroside synthesis), and *Fa2h* (myelin sphingolipid

biosynthesis). Top FA-correlated genes overlapped substantially, featuring *Cldn11*, *Tspan2*,

Cnp, and *Fa2h*, alongside immune and proteostasis regulators (*Lpar1*, *CD9*, *App*, *Adamts4*,

Cryab). MD-correlated genes included neuronal and metabolic markers such as *Pvalb*

(inhibitory interneurons), *Scn1a* (neuronal excitability), *Uqcrh* (mitochondrial respiration), and

Abcd2 (energy metabolism).

To examine associations between PLP-driven human α-syn overexpression and MRI

phenotypes, we correlated MRI t-statistic maps with regional *Plp1* expression (**Figure 3**).

Regional volume loss and FA reductions showed negative correlations with *Plp1* expression,

highlighting a spatial association between these MRI measures and oligodendrocyte-related

transcriptional patterns. In contrast, MD showed no association with *Plp1* expression,

consistent with its enrichment for neuronal and mitochondrial gene signatures and

suggesting that diffusivity changes may reflect processes distinct from primary

oligodendroglial pathology.

Gene Ontology (GO) enrichment analysis of FA-correlated genes revealed significant (FDR <

0.05) overrepresentation of processes related to myelination, gliogenesis, and astrocyte and

oligodendrocyte differentiation, all prominently featuring *Plp1* (**Figure 4; Suppl. Tables S4–**

S5). By contrast, genes associated with MD changes were enriched for dendritic spine

development, reflecting neuronal programs without implying mechanistic causality (**Suppl.**

Table S6). The modality-specific associations of MRI maps with oligodendrocyte-related

transcriptional programs argue against a purely circular effect and indicate that different MRI

measures reflect multiple, complementary biological processes.

Cell-type enrichment implicates oligodendrocytes, astrocytes, and specific neuronal

subpopulations

Cell-type–specific enrichment analysis demonstrated that genes spatially correlated with

regional volume loss were significantly enriched (FDR < 0.05) for oligodendrocyte markers in

the cortex and cerebellum, and for cholinergic neurons of the spinal cord and brainstem

(**Figure 5a; Suppl. Table S10**). Subthreshold enrichment ($0.05 < p < 0.1$) was observed for

brainstem serotonergic neurons.

Similarly, genes correlated with FA reduction were enriched for cortical and cerebellar

oligodendrocytes, cerebellar stellate and basket cells, and Purkinje cells (pSI < 0.001), as

well as cortical and cerebellar astrocytes and oligodendrocyte progenitors (pSI < 0.05)

(**Figure 5b; Suppl. Table S11**). Subthreshold enrichment was observed for Bergmann glia

and cerebellar granule cells. No cell type reached significance for genes associated with

increased MD (**Figure 5c; Suppl. Table S12**), although mild enrichment was noted for

cortical layer 6 pyramidal cells and prepronociceptin (PNOC)-expressing neurons.

RNAseq supports the biological relevance of molecular signatures of

neurodegeneration identified by imaging transcriptomics

To test if the genes we have identified indirectly to be associated with disorder-related

imaging phenotypes are indeed dysregulated in the post-mortem brain tissue of the same

model, we performed RNAseq and DESeq2 analysis on cerebellar tissue from PLP-αSyn

mice [14], which we found significantly affected in MRI modalities (**Figure 1, 2**), and cross-

referenced the 2613 differentially expressed genes (DEGs, FDR<0.05) with those correlated

to MRI-derived maps. We identified 664 DEGs overlapping with volume-correlated genes,

441 with FA-correlated genes, and 510 with MD-correlated genes (**Suppl. Tables S7-S9**).

The log2(FC) values of cerebellar DEGs were weakly but significantly correlated with FA-

correlated genes (ρ =-0.106, p=0.026) and MD-correlated genes (ρ=-0.1895, p<0.0001)

(**Figure 6**). Further enrichment analysis showed that the DEGs were significantly

overrepresented (p=0.0376) among the genes whose normative spatial expression patterns

correlated most strongly with volume changes in the PLP-αSyn mice (**Table 3**). The overlap

between DEGs and MRI-correlated genes is consistent with the biological relevance of the

imaging transcriptomics findings and highlights an association between MRI-detected

alterations and molecular signatures of neurodegeneration in MSA.

DISCUSSION

Previous work has established the PLP-αSyn mouse as a valuable MSA model for

investigating α-synuclein–driven mechanisms of neurodegeneration and testing disease-

modifying therapies that mitigate α-syn aggregation, neuroinflammation, and neuronal loss

[15-21]. Here, we provide the first comprehensive, whole-brain MRI characterization of this

model compared with wild-type controls. High-resolution ex vivo structural MRI revealed a

robust pattern of regional atrophy affecting white matter tracts, brainstem nuclei (medulla,

pons, midbrain), cerebellum, thalamus, hypothalamus, and subiculum. The pattern of volume

loss resembled that reported in MSA patients, with involvement of the hippocampus,

pons/medulla, cerebellum, and pyramidal tracts (**Table 1**), supporting the translational

relevance of MRI-detected neurodegeneration in the PLP-αSyn model. DTI revealed

widespread microstructural abnormalities: FA was reduced across major white-matter tracts

(e.g., corpus callosum, fornix, anterior commissure, striatothalamic fibres, and cerebellar

peduncles), consistent with clinical MSA and suggestive of altered myelination and axon–glia

interactions rather than overt demyelination. Focal FA increases in the hippocampus may

reflect gliosis-related reorganisation or loss of crossing fibres. In contrast, MD was

predominantly elevated in brainstem and cerebellum, consistent with tissue rarefaction and

extracellular water increase. The modest overlap between FA and MD effects indicates that

these metrics capture complementary aspects of pathology.

Applying an imaging transcriptomics framework to these multimodal MRI phenotypes, we

observed that atrophy and microstructural alterations were systematically organized and

aligned with transcriptional programs related to oligodendrocyte biology, energy metabolism,

and neuroinflammatory processes. These findings broadly converge with human MSA

imaging transcriptomics [8], despite differences in species, disease duration, imaging

resolution, and experimental context, supporting the view that imaging–transcriptomic

associations capture core molecular correlates of regional vulnerability.

A key extension beyond prior human studies is the integration of volumetric and diffusion-

derived MRI measures. While structural atrophy reflects cumulative tissue loss, diffusion

metrics such as FA and MD capture complementary microstructural features. FA and volume,

but not MD, correlated with *Plp1* expression, linking these signatures to the oligodendroglial

α-synucleinopathy, whereas MD aligned with neuronal and mitochondrial programs. Gene

Ontology and cell-type enrichment analyses reinforce these modality-specific interpretations:

oligodendrocytes and astrocytes were associated with FA and volume changes, whereas

neuronal subtypes correlated with MD-related alterations. Importantly, different MRI

phenotypes mapped onto partially distinct oligodendrocyte-associated transcriptional

signatures, suggesting that regional vulnerability within the oligodendrocyte lineage—

including myelination, metabolic support, and glial reactivity—may influence circuit-level

susceptibility and pathological manifestation. In the controlled PLP-αSyn model, modality-

specific signatures—previously unresolvable in clinical MSA—position diffusion MRI as a

biologically informative complement to volumetric imaging for translational studies.

Imaging transcriptomics relies on normative gene expression atlases, which do not capture

dynamic or region-specific transcriptional changes during neurodegeneration [5-7, 12]. In the

PLP-αSyn model, we partially addressed this by comparing atlas-derived signatures with

independent RNAseq data [14], revealing concordance that supports the biological relevance

of the associations. While molecular validation was restricted to the cerebellum—a core site

of pathology in both human MSA and the mouse model [1, 14]—the spatial alignment of MRI-

detected alterations with known neurodegeneration sites reinforces the relevance of these

imaging markers [10, 22-25].

We should point out that an immediate systematic histological confirmation was beyond the

scope of this study but the model has been previously extensively histologically

characterized [10, 22-27]. Among the limitations of this study is the reliance on ex vivo

imaging in a modest cohort of female mice, limiting longitudinal assessment and raising the

possibility of fixation-related bias. Future studies should incorporate longitudinal in vivo MRI

with time-resolved transcriptomic profiling, and experimental manipulations such as

conditional gene targeting or pharmacological interventions to establish causal links between

cellular dysfunction and MRI phenotypes.

The primary value of applying imaging transcriptomics in the PLP-αSyn model lies in

benchmarking and interpretability. By establishing how MRI phenotypes align with

transcriptional programs in a controlled system, this work provides a reference framework for

interpreting imaging findings in preclinical and clinical studies and enables direct comparison

with human MSA data, strengthening confidence in shared disease mechanisms. Rather

than replacing molecular or histological analyses, imaging transcriptomics adds value by

integrating systems-level imaging with cellular biology, offering a scalable approach to

contextualize MRI biomarkers and bridge the gap between macroscopic imaging readouts

and cellular pathology in synucleinopathies.

In summary, our study is the first to apply whole-brain imaging to an MSA relevant mouse

model and to show that its imaging phenotype parallels key features reported in human MSA.

We also extend recent human work by applying imaging transcriptomics and we highlight

both the strengths and limitations of this integrative approach. By combining multimodal MRI

with atlas-based transcriptomic mapping and model-specific RNAseq, we provide a

multiscale perspective on brain vulnerability in MSA and establish a foundation for future

longitudinal and interventional studies to identify early imaging biomarkers and test

mechanistic hypotheses.

METHODS

Animals

This study included nine female PLP-αSyn mice (four 10-month-old and five 15-month-old)

and nine age-matched female wild-type (WT) mice (six 11-month-old and three 15-month-

old). The animals were bred and aged in the Animal Facility of the Medical University

Innsbruck under temperature-controlled pathogen-free conditions with a light/dark 12-hour

cycle and free access to food and water. Mice were anesthetized with an overdose of

thiopental and intracardially perfused with phosphate buffered saline (PBS pH 7.4, Sigma-

Aldrich, Vienna, Austria) followed by 4% paraformaldehyde (PFA, pH 7.4, Sigma-Aldrich,

Vienna, Austria). The heads were removed, postfixed in 4% paraformaldehyde for 24h at 4°C

and stored in PBS with 0.05% sodium azide until scanning. All experiments were performed

according to the ethical guidelines with the permission of the Austrian Federal Ministry of

Science and Research (BMFWF-2022-0.899.398).

The sample size (n = 9 per group) was determined using the resource equation method,

which ensures an efficient balance between statistical validity and ethical reduction principles

[28]. In this design, the calculated error degrees of freedom (E = total number of animals –

total number of groups = 18 – 2 = 16) fall within the acceptable range of $10 \leq E \leq 20$,

indicating adequate power for detecting biologically meaningful effects while minimizing

unnecessary animal use. The chosen group size is consistent with previous preclinical MRI

studies in transgenic models of neurodegeneration. Only female mice were included to

reduce variability associated with sex differences in brain structure.

Magnetic resonance imaging

Image acquisition

Skulls were placed four at a time in a 50-ml Falcon tube, separated and secured by a custom

3D-printed holder, and immersed in perfluoropolyether (Galden®, Solvay). These were

scanned in a 9.4T Bruker BioSpec 94/20 preclinical MRI scanner with a 39-mm radio-

frequency coil to acquire T2-weighted (T2w) structural images and diffusion tensor imaging

(DTI) data as previously described [29]. In brief, T2w images with 100-micron isotropic voxel

size were acquired with a 3D fast spin echo sequence, and DTI images with 200-micron

isotropic voxel size were acquired with a 2D multi-slice pulsed gradient spin echo sequence

(4 b0 images, 30 diffusion directions, b=1500 s/mm2).

Image processing and analysis

Diffusion tensor fitting was performed using the dtifit command in FMRIB Software Library

(FSL version 5.0.10). Multi-modal study-specific templates were created from the resulting

fractional anisotropy (FA) and mean diffusivity (MD) maps and the T2w images using the

antsMultivariateTemplateConstruction2.sh script included in Advanced Normalization Tools

(ANTs version 2.2.0). Template and individual subject brain masks were created by

multiplying the reciprocal of the MD map and the T2w image (fslmaths) and passing this

image product to the Rapid Automatic Tissue Segmentation (RATS) tool for skull-stripping of

rodent brain MRI [30, 31]. Masked subject images were registered to the masked templates

using antsRegistration to perform sequential rigid-body, affine, and diffeomorphic SyN multi-

modal registrations with equal weight given to the FA, MD, and T2w images. To quantify

inter-subject voxel-level variations in local brain volume, maps of the Jacobian determinants

of the composite transformations were computed using the ANTs

CreateJacobianDeterminantImage command. The registered FA and MD maps were

smoothed with a full-width half-maximum of 0.2 mm using the 3dBlurInMask command in the

Analysis of Functional NeuroImages (AFNI) software suite (version 17.2.17). Univariate, 2×2

ANOVAs were performed on the log-transformed Jacobian determinant maps and the

smoothed DTI maps to test for the effects of age (old vs. middle age) and genotype (PLP-

αSyn vs. WT) on volume, FA, and MD. This was done using FSL's randomise tool for

nonparametric permutation inference to perform 10,000 permutations, threshold-free cluster

enhancement, and family-wise error correction.

For region-of-interest (ROI)-based analysis, the masked T2w study template was registered

to the Allen Mouse Brain Common Coordinate Framework (CCFv3) using antsRegistration as

described above. The CCFv3 atlas labels were merged into 87 ROIs and warped to study

template space. ROI volumes were computed by taking the sum of all Jacobian determinant

values within each ROI and multiplying by the volume of one voxel. Mann-Whitney U (MWU)

tests were performed to compare ROI volumes and mean FA and MD values from the PLP-

αSyn and WT groups. The Benjamini-Hochberg procedure was used to correct for multiple

comparisons (87 ROIs) for each MRI parameter.

Spatial gene enrichment analysis

Given the limited sample size, all mice (10-15 months of age) were pooled and only the main

effects of genotype were considered for spatial gene enrichment analysis. Only the coronal

dataset of the Allen *in situ* hybridization data was used because, although it contains far

fewer genes than the sagittal dataset, it has superior brain coverage and quality. After

excluding gene expression energy maps that had missing data in greater than 20% of voxels

within the brain, the final number of maps included in the analysis was 4071 (some genes

have duplicates, which were treated separately and all kept for correlation analysis).

Imaging-transcriptomic correlation

The volume, FA, and MD t-statistic maps generated by randomise were warped to CCFv3

space and down-sampled to 200-micron isotropic voxel size to match the gene expression

energy maps. The Spearman correlation coefficient (ρ) was calculated between each MRI t-

statistic map and each gene expression energy map.

To calculate non-parametric p-values for the ρ values, the BrainSMASH Python package [32]

was used to generate 5000 surrogate maps that match the spatial autocorrelation of each of

the volume, FA, and MD t-statistic maps. The Spearman correlation coefficient was

computed between each surrogate map and each gene expression energy map to build null

distributions from which p-values were computed (**Suppl. Fig. S1**).

Gene set enrichment analysis

Two separate gene set enrichment analyses were performed using two categories of gene

sets: gene sets associated with the Gene Ontology (GO) biological process terms, and cell-

type specific expression (CSE) gene sets curated by Dougherty et al. [33].

For each of 27 different cell types, the authors identified genes that are enriched in that cell

population. Further, they truncated these gene lists at four Specificity Index thresholds (pSI)

(a smaller pSI corresponds to a more exclusive list of genes that are more specific to a

particular cell type) (http://doughertytools.wustl.edu/CSEAtool.html). Two cell types—cone

and rod cells—were deemed irrelevant to this study and excluded.

For each type of analysis, only genes found in both the filtered Allen coronal dataset and the

GO or CSE gene sets were included, resulting in 3606 and 3646 expression maps of 3385

and 3422 unique genes, respectively. The GO biological process terms were filtered by

removing those with fewer than 10 or more than 200 remaining genes, à la Ritchie [34].

Furthermore, terms with identical gene sets (exactly equal) to another term were discarded,

leaving a final count of 2393 GO biological process terms. The curation of the GO and CSE

gene sets were performed with custom Python scripts.

For each MRI parameter, genes were ranked by FDR-corrected p-value and then by the

absolute value of ρ. For genes with duplicates, only the highest ranked duplicate was kept for

gene set enrichment analysis. The minimum hypergeometric (mHG) score [35] was used to

quantify the enrichment of GO and CSE gene sets at the top of these ranked lists. To account

for propagating effects of spatial autocorrelation into the enrichment analysis, a null

distribution of mHG scores for each MRI parameter and gene set pair was calculated from

the BrainSMASH-derived surrogate maps. For each observed mHG score s, a p-value was

calculated as the fraction of mHG scores in the corresponding null distribution $s' < s$. To

control the false discovery rate in each MRI parameter, the Benjamini-Hochberg procedure

was performed across: 1) all 2393 GO gene sets, 2) the 42 GO gene sets that contain Plp1,

and 3) the 25 CSE gene sets for each pSI level.

Imaging transcriptomics and RNAseq correlation

To consolidate the relevance of the disease-associated gene maps defined by imaging

transcriptomics, we identified the significantly correlated genes to volume, FA and MD

(p<0.05), which are found significantly dysregulated in the PLP-αSyn cerebellum by RNAseq

and DESeq2 analysis (p_{adj}>0.05,[14]). We calculated the Spearman correlation of the

Spearman ρ values from the imaging transcriptomics and the Log2(FC) values from the

DESeq2 analysis applying GraphPad Prism 10.5.0 software. In addition, we performed the

enrichment analysis described above using the significantly dysregulated genes as a gene

set. Of the 3494 genes found in both the Allen Atlas and RNAseq datasets, 758 genes were

differentially expressed and considered in this enrichment analysis.

Figure legends

Figure 1. Multiparametric MRI phenotype of PLP-⬚Syn brains compared to wild-type

mouse brains. Voxel-wise differences in a) volume, b) fractional anisotropy (FA), and c)

mean diffusivity (MD) overlaid on the Allen Mouse Brain Common Coordinate Framework

(CCFv3) template. The difference in group means (PLP-⬚Syn minus wild-type) is mapped to

the color of the overlays, while the family-wise error rate (FWER) is mapped to the

transparency of the overlays. The overlays are completely transparent where the FWER >

0.5, while areas where the FWER < 0.05 are contoured by black lines.

Figure 2. Most affected brain regions in PLP-⬚Syn mice according to atlas-based

statistics of the MRI phenotype. a) Dot and box plots of regional volume, mean fractional

anisotropy (FA), and mean mean diffusivity (MD) for select regions-of-interest (ROI). Two-

tailed Mann-Whitney U tests, FDR-corrected p < 0.1*, 0.01**, 0.001***. b) 3-dimensional

representation of the selected ROIs.

Figure 3. Spatial correlation between the PLP-⬚Syn MRI phenotype and Allen gene

expression maps. Left: coronal slices of the Plp1 expression energy map from the Allen

Mouse Brain Atlas. Middle: corresponding slices of t-statistic maps from voxel-wise

comparisons of volume, fractional anisotropy (FA), and mean diffusivity (MD) in PLP-⬚Syn

mice vs wild-types. Right: corresponding slices of gene expression energy maps that

correlated most strongly with the t-statistic maps. The Spearman correlation coefficients (ρ)

between the t-statistic and gene expression maps are shown in the double arrows.

Figure 4. Enriched Gene Ontology terms among genes most spatially correlated with

the PLP-⬚Syn MRI phenotype. Directed acyclic graphs of Gene Ontology (GO) biological

process terms containing Plp1 that are significantly enriched (Benjamini-Hochberg-corrected

p < 0.05) in the top of ranked lists of genes most correlated with PLP-⬚Syn changes in a)

volume and b) fractional anisotropy (FA). For readability purposes, parent terms were

included in the graph only if they are in the shortest path between a significantly enriched

term and the root "biological process" term. The GO term nodes are colored if corrected p <

0.1, with the node outline color indicating the enrichment score. For a given GO term,

enrichment = (b/n) / (B/N), where N is the total number of genes in the ranked list; B is the

number of genes associated with the GO term; n is the size of the target set, i.e., the number

of genes in the top of the ranked list that gives the minimum hypergeometric (mHG) score;

and b is the number of genes in the target set that are associated with the GO term.

Figure 5. Enriched cell types in genes most spatially correlated with the PLP-⍺Syn MRI

phenotype. Each cell type is represented by four concentric hexagons, the areas of which

are proportional to the number of gene that are enriched in that cell type at four different

Specificity Index (pSI) thresholds. The smallest, central hexagon represents the list of genes

that are most specifically enriched in a particular cell type. The hexagons are grouped by

brain region, indicated by the background color. Hexagons are colored if there is an

enrichment (Benjamini-Hochberg-corrected p < 0.1) of genes associated with the

corresponding cell-type in the ranked lists of genes most correlated with PLP-⍺Syn changes

in a) volume, b) fractional anisotropy (FA), and c) mean diffusivity (MD).

Figure 6. Spearman correlation analysis of DEGs in the cerebellum and MRI–linked

genes identified by imaging transcriptomics. The Spearman correlation coefficients

between genes' normative spatial expression patterns (Allen Mouse Brain Atlas) and MRI-

measured volume, fractional anisotropy (FA), and mean diffusivity (MD) changes in the PLP-

αSyn brain are plotted against the genes' log2(FC) from DESeq2 analysis of PLP-αSyn

cerebellar tissue.

Tables

Table 1: Regions of interest with significantly altered MRI volumes and diffusivity in human

MSA compared to healthy controls.

Brain region	MRI			
	volume	**FA**	**MD**	*Ref.*
Cerebral white matter	↑	↓	↑	*[36-41]*
Amygdala	-	↓	-	*[42]*
Pyramidal tract	↓	↓	↑	*[37, 43-49]*
Capsula interna	-	↓	↑	*[36-38, 43, 45, 50, 51]*
Putamen/Striatum	↓	↓	↑	*[36, 42, 44, 52-62]*
Globus pallidus	-	↑	-	*[42, 60]*
Midbrain	↓	↓	↑	*[38, 58, 60, 63, 64]*
Cerebral peduncle		↓	↑	*[38, 51]*
Substantia nigra	↓	↓	-	*[42, 56, 58, 65, 66]*
Red nucleus	-	-	↑	*[42]*
Subthalamic nucleus	-	-	↑	*[42]*
Brainstem	-	-	↑	*[67]*
Medulla	↓	-	-	*[56]*
Pons	↓	↓	↑	*[37, 38, 44, 45, 48, 50, 51, 56, 58-61, 63, 65, 68-72]*
Cerebellum	gray matter: ↓ white matter: ↑/↓ total: ↓	↓	↑	*[38-42, 44-46, 48, 52, 53, 56, 58-61, 67, 70, 71, 73, 74]*
Middle cerebellar peduncle (MCP)	↓	↓	↑	*[36-38, 42, 44-46, 48-50, 56, 58-61, 68, 70, 75-79]*
Superior cerebellar peduncle (SCP)	-	↓	↑	*[36-38, 60, 78]*

476   ↑ = *significantly increased*
477   ↓ = *significantly decreased*
478   - = *no data available*
479   *FA = fractional anisotropy*
480   *MD = mean diffusivity*
481

482   **Table 2.** Most affected brain regions according to atlas-based statistics comparing PLP-αSyn

483   and wild-type mice.

volume			FA			MD		
ROI	% difference in group medians	FDR-corrected p-value	ROI	% difference in group medians	FDR-corrected p-value	ROI	% difference in group medians	FDR-corrected p-value
epithalamus related	-8.89	0.004	anterior commissure, olfactory limb	-12.32	5.10E-04	Pons, behavioral state related	4.86	0.05
Perithalamus	-14.15	0.004	optic nerve	-12.59	5.10E-04	Cerebellar nuclei	4.16	0.054
Ventral tegmental area	-19.98	0.004	thalamus related	-13.64	5.10E-04	Red nucleus	7.08	0.054
Raphe nuclei	-14.7	0.004	fornix system	-13.59	5.10E-04	(Para)flocculus	7.96	0.054
Pons, sensory related	-11.73	0.004	epithalamus related	-13.33	5.10E-04	Cerebellar vermis	8.84	0.064
Pons, motor related	-10.33	0.004	lateral ventricle	-19.32	5.10E-04	Cerebellar hemispheres	15.12	0.115
medial lemniscus	-12.7	0.004	third ventricle	-14.36	5.10E-04	Inferior colliculus	9.6	0.18
extrapyramidal fiber systems	-11.97	0.004	corpus callosum	-13.08	7.20E-04	third ventricle	19.3	0.18
Insular cortex	7.82	0.005	fourth ventricle	-9.03	7.20E-04	Pedunculotegmental nucleus	5.36	0.18
Reticular nucleus	-11.5	0.005	choroid plexus	-18.24	7.20E-04	arbor vitae	3.87	0.18
Medulla, motor related	-10.04	0.005	stria terminalis	-13.1	0.001	Pons, motor related	4.07	0.217
Intralaminar thalamic nuclei	-10.82	0.006	Olfactory bulb	-6.93	0.003	Ventral thalamus	5.39	0.228
Pedunculotegmental nucleus	-9.34	0.006	Geniculate nucleus	-9.51	0.003	facial nerve	4.08	0.241
cerebellar peduncles	-14.15	0.006	Perithalamus	-9.83	0.003	Raphe nuclei	5.49	0.288
internal capsule	-17.15	0.007	Pretectal region	-13.58	0.003	lateral ventricle	36.01	0.292
Pons, behavioral state related	-9.64	0.008	Superior colliculus	-9.94	0.003	oculomotor nerve	5.18	0.319
Cerebellar nuclei	-15.68	0.008	arbor vitae	-9.3	0.003	Medulla, sensory related	2.72	0.319
vestibulocochlear nerve	-13.28	0.008	internal capsule	-9.75	0.004	corticospinal tract	-2.03	0.319
stria terminalis	-9.81	0.013	Cerebellar hemispheres	-10.74	0.005	Primary somatosensory cortex	6.51	0.319
Cuneiform nucleus	-6.82	0.016	Cerebellar nuclei	-4.42	0.007	Habenula	8.99	0.319

486   **Table 3.** Cerebellar DEG enrichment analysis.

MRI parameter	N	B	b	n	enrichment	smHG	p-value
volume	3494	758	146	504	1.33529128	1.96E-05	0.0376
FA	3494	758	9	18	2.30474934	0.00735212	0.3692
MD	3494	758	5	6	3.8412489	0.00234034	0.2398
DEG: differentially expressed gene, N: total number of genes found in both the Allen Atlas and RNAseq datasets, B: number of DEGs in the N total genes, b: number of DEGs in the top n ranked genes, enrichment = (b/n)/(B/N), smHG: minimum hypergeometric test statistic, FA: fractional anisotropy, MD: mean diffusivity							

Supplementary material

Additional file 1

Title: Supplementary Figures: Fig S1 to S3.

Format: .pdf

Additional file 2

Title: Supplementary Tables: Tables S1 to S12.

Format: .xls

Captions:

Suppl Table S1. Genes significantly correlated with regional volume changes.

Suppl Table S2. Genes significantly correlated with regional FA changes.

Suppl Table S3. Genes significantly correlated with regional MD changes.

Suppl Table S4. GO enrichment analysis of genes correlated with volume changes.

Suppl Table S5. GO enrichment analysis of genes correlated with FA changes.

Suppl Table S6. GO enrichment analysis of genes correlated with MD changes.

Suppl Table S7. DEGs in RNAseq of cerebellum (p<0.05, Log2(FC)) and significantly

correlated with volume changes (p<0.05, Spearman r).

Suppl Table S8. DEGs in RNAseq of cerebellum (p<0.05, Log2(FC)) and significantly

correlated with FA changes (p<0.05, Spearman r).

Suppl Table S9. DEGs in RNAseq of cerebellum (p<0.05, Log2(FC)) and significantly

correlated with MD changes (p<0.05, Spearman r).

Suppl Table S10. Cell-type–specific enrichment analysis of genes correlated with volume

changes

Suppl Table S11. Cell-type–specific enrichment analysis of genes correlated with FA

changes

Suppl Table S12. Cell-type–specific enrichment analysis of genes correlated with MD

changes

Availability of data and materials

All data generated or analysed during this study are included in this published article and its

supplementary information files.

Competing interests

EK, declares no competing interests

DC, declares no competing interests

DM, declares no competing interests

CS, declares no competing interests

EK, declares no competing interests

AHG, received grant support from the Austrian Science Fund (FWF) and the Tyrolian Science

Fund (TWF) outside of the submitted work.

FK, received personal fees from AbbVie, Bial, Ionis Pharmaceuticals, Koneksa, Merz,

Österreichische Apotheker-Verlagsgesellschaft, Sanofi, Takeda Pharmaceuticals, and Teva

and his institution has ongoing grant support from the Austrian Science Fund (FWF), the

National Institutes of Health and the Michael J. Fox Foundation, outside of the submitted

work.

LM, declares no competing interests

NS, received grant support from the Medical University of Innsbruck and The Michael J. Fox

Foundation, outside of the submitted work; Director-at-large, Board of Directors and Chair of

the Research Steering Council of Mission MSA, USA; Advisory board, Karl Golser

Foundation, Italy; Advisory services for IONIS, Mitsubishi Pharma

Funding
542
543 DM is supported by NIHR Maudsley Biomedical Research Centre, South London and

544 Maudsley NHS Trust.

Authors' contributions

EK has contributed to the methodology, data analysis of the study and drafting parts of the

manuscript

DC has contributed to the conceptualization of the study, data interpretation, and drafting

parts of the manuscript

DM has contributed to the methodology of the study

CS has contributed to the methodology of the study

EK has contributed by summarizing clinical MRI/DTI studies in MSA

AHG has contributed to data analysis and tissue preparation

FK has contributed to the interpretation of the clinical relevance of the data

LM has contributed to the conceptualization of the study

NS has contributed to the conceptualization of the study, analysis of the data, data

interpretation, and drafting parts of the manuscript

All authors have reviewed and approved the final version of the manuscript.

References

1.   Poewe W, Stankovic I, Halliday G, Meissner WG, Wenning GK, Pellecchia MT, et al. Multiple

system atrophy. Nat Rev Dis Primers. 2022;8:56.

2.   Stefanova N, Wenning GK. Multiple system atrophy: at the crossroads of cellular, molecular

and genetic mechanisms. Nat Rev Neurosci. 2023;24:334-346.

3.   Fabbri M, Del Campo N, Meissner WG, Rousseau V, Sommet A, Payoux P, et al. Disease

Progression in Multiple System Atrophy: The ASPIRE Multi-Modal Biomarker Study. Annals of

neurology. 2025;10.1002/ana.70028.

4.   Krismer F, Seppi K, Poewe W. The role of neuro-imaging in multiple system atrophy. J Neural

Transm (Vienna). 2025;10.1007/s00702-025-02964-6.

5.   Adewale Q, Khan AF, Carbonell F, Iturria-Medina Y, Alzheimer's Disease Neuroimaging I.

Integrated transcriptomic and neuroimaging brain model decodes biological mechanisms in aging and

Alzheimer's disease. Elife. 2021;10.

6.   Mroczek M, Desouky A, Sirry W. Imaging Transcriptomics in Neurodegenerative Diseases. J

Neuroimaging. 2021;31:244-250.

7.   Arnatkeviciute A, Fulcher BD, Bellgrove MA, Fornito A. Imaging Transcriptomics of Brain

Disorders. Biol Psychiatry Glob Open Sci. 2022;2:319-331.

8.   Chougar L, Tremblay C, Delva A, Filiatrault M, Vo A, Hansen JY, et al. MRI-derived atrophy in

multiple system atrophy aligns with mitochondrial and glial gene expression patterns. NPJ Parkinson's

disease. 2025;12:16.

9.   Kahle PJ, Neumann M, Ozmen L, Muller V, Jacobsen H, Spooren W, et al.

Hyperphosphorylation and insolubility of alpha-synuclein in transgenic mouse oligodendrocytes.

EMBO Rep. 2002;3:583-588.

10.   Refolo V, Bez F, Polissidis A, Kuzdas-Wood D, Sturm E, Kamaratou M, et al. Progressive

striatonigral degeneration in a transgenic mouse model of multiple system atrophy: translational

implications for interventional therapies. Acta Neuropathol Commun. 2018;6:2.

11.   Stefanova N. A Mouse Model of Multiple System Atrophy: Bench to Bedside.

Neurotherapeutics. 2023;20:117-126.

12.   Martins D, Giacomel A, Williams SCR, Turkheimer F, Dipasquale O, Veronese M, et al.

Imaging transcriptomics: Convergent cellular, transcriptomic, and molecular neuroimaging signatures

in the healthy adult human brain. Cell Rep. 2021;37:110173.

13.   Edwards LJ, McColgan P, Helbling S, Zarkali A, Vaculciakova L, Pine KJ, et al. Quantitative

MRI maps of human neocortex explored using cell type-specific gene expression analysis. Cereb

Cortex. 2023;33:5704-5716.

14.   Heras-Garvin A, Fellner L, Granata R, Wenning GK, Stefanova N. Transcriptional

dysregulation in the cerebellum triggered by oligodendroglial alpha-synucleinopathy: insights from a

transgenic mouse into the early disease mechanisms of MSA. J Neural Transm (Vienna).

2025;10.1007/s00702-025-02892-5.

15.   Heras-Garvin A, Refolo V, Schmidt C, Malfertheiner K, Wenning GK, Bradbury M, et al.

ATH434 Reduces alpha-Synuclein-Related Neurodegeneration in a Murine Model of Multiple System

Atrophy. Mov Disord. 2021;36:2605-2614.

16.   Luque M, Matic M, Heras-Garvin A, Amo-Aparicio J, Luk KC, Haindl MT, et al. Pharmacologic

NLRP3 Inhibition Modulates Parkinson's Disease-Associated Microglial Transcriptomic Signatures and

Mitigates α-Synuclein–Triggered Neurodegeneration. bioRxiv.

2025;10.1101/2025.10.22.683837:2025.2010.2022.683837.

17.   Cheng A, Kawahata I, Wang Y, Jia W, Wang H, Sekimori T, et al. Epsin2, a novel target for

multiple system atrophy therapy via alpha-synuclein/FABP7 propagation. Brain. 2023;146:3172-3180.

18.   Heras-Garvin A, Weckbecker D, Ryazanov S, Leonov A, Griesinger C, Giese A, et al.

Anle138b modulates α-synuclein oligomerization and prevents motor decline and neurodegeneration

in a mouse model of multiple system atrophy. Mov Disord. 2019;34:255-263.

19.   Lemos M, Venezia S, Refolo V, Heras-Garvin A, Schmidhuber S, Giese A, et al. Targeting α-

synuclein by PD03 AFFITOPE® and Anle138b rescues neurodegenerative pathology in a model of

multiple system atrophy: clinical relevance. Translational neurodegeneration. 2020;9:38.

20.   Venezia S, Kaufmann WA, Wenning GK, Stefanova N. Toll-like receptor 4 deficiency facilitates

α-synuclein propagation and neurodegeneration in a mouse model of prodromal Parkinson's disease.

Parkinsonism Relat Disord. 2021;91:59-65.

21.   Venezia S, Refolo V, Polissidis A, Stefanis L, Wenning GK, Stefanova N. Toll-like receptor 4

stimulation with monophosphoryl lipid A ameliorates motor deficits and nigral neurodegeneration

triggered by extraneuronal α-synucleinopathy. Molecular neurodegeneration. 2017;12:52.

22.   Kuzdas D, Stemberger S, Gaburro S, Stefanova N, Singewald N, Wenning GK.

Oligodendroglial alpha-synucleinopathy and MSA-like cardiovascular autonomic failure: experimental

evidence. Exp Neurol. 2013;247:531-536.

23.   Stemberger S, Poewe W, Wenning GK, Stefanova N. Targeted overexpression of human

alpha-synuclein in oligodendroglia induces lesions linked to MSA-like progressive autonomic failure.

Exp Neurol. 2010;224:459-464.

24.   Boudes M, Uvin P, Pinto S, Voets T, Fowler CJ, Wenning GK, et al. Bladder dysfunction in a

transgenic mouse model of multiple system atrophy. Mov Disord. 2013;28:347-355.

25.   Stefanova N, Reindl M, Neumann M, Haass C, Poewe W, Kahle PJ, et al. Oxidative stress in

transgenic mice with oligodendroglial alpha-synuclein overexpression replicates the characteristic

neuropathology of multiple system atrophy. Am J Pathol. 2005;166:869-876.

26.   Krismer F, Wenning GK, Li Y, Poewe W, Stefanova N. Intact olfaction in a mouse model of

multiple system atrophy. PLoS One. 2013;8:e64625.

27.   Stefanova N, Kaufmann WA, Humpel C, Poewe W, Wenning GK. Systemic proteasome

inhibition triggers neurodegeneration in a transgenic mouse model expressing human alpha-synuclein

under oligodendrocyte promoter: implications for multiple system atrophy. Acta Neuropathol.

2012;124:51-65.

28.   Charan J, Kantharia ND. How to calculate sample size in animal studies? Journal of

pharmacology & pharmacotherapeutics. 2013;4:303-306.

29.   White MA, Lin Z, Kim E, Henstridge CM, Pena Altamira E, Hunt CK, et al. Sarm1 deletion

suppresses TDP-43-linked motor neuron degeneration and cortical spine loss. Acta Neuropathol

Commun. 2019;7:166.

30.   Oguz I, Zhang H, Rumple A, Sonka M. RATS: Rapid Automatic Tissue Segmentation in rodent

brain MRI. J Neurosci Methods. 2014;221:175-182.

31.   Yin Y, Zhang X, Williams R, Wu X, Anderson DD, Sonka M. LOGISMOS--layered optimal

graph image segmentation of multiple objects and surfaces: cartilage segmentation in the knee joint.

IEEE Trans Med Imaging. 2010;29:2023-2037.

32.   Burt JB, Helmer M, Shinn M, Anticevic A, Murray JD. Generative modeling of brain maps with

spatial autocorrelation. Neuroimage. 2020;220:117038.

33.   Dougherty JD, Schmidt EF, Nakajima M, Heintz N. Analytical approaches to RNA profiling data

for the identification of genes enriched in specific cells. Nucleic Acids Res. 2010;38:4218-4230.

34.   Ritchie J, Pantazatos SP, French L. Transcriptomic characterization of MRI contrast with focus

on the T1-w/T2-w ratio in the cerebral cortex. Neuroimage. 2018;174:504-517.

35.   Eden E, Lipson D, Yogev S, Yakhini Z. Discovering motifs in ranked lists of DNA sequences.

PLoS Comput Biol. 2007;3:e39.

36.   Rulseh AM, Keller J, Rusz J, Syka M, Brozova H, Rusina R, et al. Diffusion tensor imaging in

the characterization of multiple system atrophy. Neuropsychiatr Dis Treat. 2016;12:2181-2187.

37.   Gorges M, Maier MN, Rosskopf J, Vintonyak O, Pinkhardt EH, Ludolph AC, et al. Regional

microstructural damage and patterns of eye movement impairment: a DTI and video-oculography

study in neurodegenerative parkinsonian syndromes. J Neurol. 2017;264:1919-1928.

38.   Dash SK, Stezin A, Takalkar T, George L, Kamble NL, Netravathi M, et al. Abnormalities of

white and grey matter in early multiple system atrophy: comparison of parkinsonian and cerebellar

variants. Eur Radiol. 2019;29:716-724.

39.   Jao CW, Soong BW, Huang CW, Duan CA, Wu CC, Wu YT, et al. Diffusion Tensor Magnetic

Resonance Imaging for Differentiating Multiple System Atrophy Cerebellar Type and Spinocerebellar

Ataxia Type 3. Brain Sci. 2019;9.

40.   Ge Y, Zheng W, Li Y, Dou W, Ren S, Chen Z, et al. Altered Brain Volume, Microstructure

Metrics and Functional Connectivity Features in Multiple System Atrophy. Front Aging Neurosci.

2022;14:799251.

41.   Raghavan S, Lesnick TG, Castillo AM, Reid RI, Fought AJ, Thostenson KB, et al. White Matter

Abnormalities Track Disease Progression in Multiple System Atrophy. Movement disorders clinical

practice. 2024;11:1085-1094.

42.   Du G, Lewis MM, Kanekar S, Sterling NW, He L, Kong L, et al. Combined Diffusion Tensor

Imaging and Apparent Transverse Relaxation Rate Differentiate Parkinson Disease and Atypical

Parkinsonism. AJNR Am J Neuroradiol. 2017;38:966-972.

43.   Ito M, Watanabe H, Atsuta N, Senda J, Kawai Y, Tanaka F, et al. Fractional anisotropy values

detect pyramidal tract involvement in multiple system atrophy. J Neurol Sci. 2008;271:40-46.

44.   Ito M, Watanabe H, Kawai Y, Atsuta N, Tanaka F, Naganawa S, et al. Usefulness of combined

fractional anisotropy and apparent diffusion coefficient values for detection of involvement in multiple

system atrophy. J Neurol Neurosurg Psychiatry. 2007;78:722-728.

45.   Prakash N, Hageman N, Hua X, Toga AW, Perlman SL, Salamon N. Patterns of fractional

anisotropy changes in white matter of cerebellar peduncles distinguish spinocerebellar ataxia-1 from

multiple system atrophy and other ataxia syndromes. Neuroimage. 2009;47 Suppl 2:T72-81.

46.   Wang PS, Wu HM, Lin CP, Soong BW. Use of diffusion tensor imaging to identify similarities

and differences between cerebellar and Parkinsonism forms of multiple system atrophy.

Neuroradiology. 2011;53:471-481.

47.   Ji L, Zhu D, Xiao C, Shi J. Tract based spatial statistics in multiple system atrophy: a

comparison between clinical subtypes. Parkinsonism Relat Disord. 2014;20:1050-1055.

48.   Faber J, Giordano I, Jiang X, Kindler C, Spottke A, Acosta-Cabronero J, et al. Prominent

White Matter Involvement in Multiple System Atrophy of Cerebellar Type. Mov Disord. 2020;35:816-

824.

49.   Pang H, Yu Z, Yu H, Li X, Bu S, Liu Y, et al. Advanced Cognitive Patterns in Multiple System

Atrophy Compared to Parkinson's Disease: An Individual Diffusion Tensor Imaging Study. Acad Radiol.

2024;10.1016/j.acra.2024.01.006.

50.   Shiga K, Yamada K, Yoshikawa K, Mizuno T, Nishimura T, Nakagawa M. Local tissue

anisotropy decreases in cerebellopetal fibers and pyramidal tract in multiple system atrophy. J Neurol.

2005;252:589-596.

51.   Yang H, Wang X, Liao W, Zhou G, Li L, Ouyang L. Application of diffusion tensor imaging in

multiple system atrophy: the involvement of pontine transverse and longitudinal fibers. Int J Neurosci.

2015;125:18-24.

52.   Tir M, Delmaire C, le Thuc V, Duhamel A, Destée A, Pruvo JP, et al. Motor-related circuit

dysfunction in MSA-P: Usefulness of combined whole-brain imaging analysis. Mov Disord.

2009;24:863-870.

53.   Novak P, Williams A, Ravin P, Zurkiya O, Abduljalil A, Novak V. Treatment of multiple system

atrophy using intravenous immunoglobulin. BMC Neurol. 2012;12:131.

54.   Baudrexel S, Seifried C, Penndorf B, Klein JC, Middendorp M, Steinmetz H, et al. The value of

putaminal diffusion imaging versus 18-fluorodeoxyglucose positron emission tomography for the

differential diagnosis of the Parkinson variant of multiple system atrophy. Mov Disord. 2014;29:380-

387.

55.   Meijer FJ, van Rumund A, Tuladhar AM, Aerts MB, Titulaer I, Esselink RA, et al. Conventional

3T brain MRI and diffusion tensor imaging in the diagnostic workup of early stage parkinsonism.

Neuroradiology. 2015;57:655-669.

56. Huppertz HJ, Möller L, Südmeyer M, Hilker R, Hattingen E, Egger K, et al. Differentiation of

neurodegenerative parkinsonian syndromes by volumetric magnetic resonance imaging analysis and

support vector machine classification. Mov Disord. 2016;31:1506-1517.

57. Planetta PJ, Ofori E, Pasternak O, Burciu RG, Shukla P, DeSimone JC, et al. Free-water

imaging in Parkinson's disease and atypical parkinsonism. Brain. 2016;139:495-508.

58. Krismer F, Seppi K, Göbel G, Steiger R, Zucal I, Boesch S, et al. Morphometric MRI profiles of

multiple system atrophy variants and implications for differential diagnosis. Mov Disord. 2019;34:1041-

1048.

59. Seki M, Seppi K, Mueller C, Potrusil T, Goebel G, Reiter E, et al. Diagnostic Potential of

Multimodal MRI Markers in Atypical Parkinsonian Disorders. J Parkinsons Dis. 2019;9:681-691.

60. Krismer F, Beliveau V, Seppi K, Mueller C, Goebel G, Gizewski ER, et al. Automated Analysis

of Diffusion-Weighted Magnetic Resonance Imaging for the Differential Diagnosis of Multiple System

Atrophy from Parkinson's Disease. Mov Disord. 2021;36:241-245.

61. Wenning GK, Stankovic I, Vignatelli L, Fanciulli A, Calandra-Buonaura G, Seppi K, et al. The

Movement Disorder Society Criteria for the Diagnosis of Multiple System Atrophy. Mov Disord.

2022;37:1131-1148.

62. Kübler D, Kobylecki C, McDonald KR, Anton-Rodriguez JM, Herholz K, Carter SF, et al.

Structural and metabolic correlates of neuropsychological profiles in multiple system atrophy and

Parkinson's disease. Parkinsonism Relat Disord. 2023;107:105277.

63. Oba H, Yagishita A, Terada H, Barkovich AJ, Kutomi K, Yamauchi T, et al. New and reliable

MRI diagnosis for progressive supranuclear palsy. Neurology. 2005;64:2050-2055.

64. Cosottini M, Ceravolo R, Faggioni L, Lazzarotti G, Michelassi MC, Bonuccelli U, et al.

Assessment of midbrain atrophy in patients with progressive supranuclear palsy with routine magnetic

resonance imaging. Acta Neurol Scand. 2007;116:37-42.

65. Mangesius S, Krismer F, Gizewski ER, Müller C, Hussl A, Schocke M, et al. 1.5 Versus 3 tesla

magnetic resonance planimetry in neurodegenerative parkinsonism. Mov Disord. 2016;31:1925-1927.

66. Andersen AM, Kaalund SS, Marner L, Salvesen L, Pakkenberg B, Olesen MV. Quantitative

cellular changes in multiple system atrophy brains. Neuropathol Appl Neurobiol. 2023;49:e12941.

67. Nicoletti G, Rizzo G, Barbagallo G, Tonon C, Condino F, Manners D, et al. Diffusivity of

cerebellar hemispheres enables discrimination of cerebellar or parkinsonian multiple system atrophy

from progressive supranuclear palsy-Richardson syndrome and Parkinson disease. Radiology.

2013;267:843-850.

68.   Blain CR, Barker GJ, Jarosz JM, Coyle NA, Landau S, Brown RG, et al. Measuring brain stem

and cerebellar damage in parkinsonian syndromes using diffusion tensor MRI. Neurology.

2006;67:2199-2205.

69.   Makino T, Ito S, Kuwabara S. Involvement of pontine transverse and longitudinal fibers in

multiple system atrophy: a tractography-based study. J Neurol Sci. 2011;303:61-66.

70.   Ito K, Ohtsuka C, Yoshioka K, Maeda T, Yokosawa S, Mori F, et al. Differentiation Between

Multiple System Atrophy and Other Spinocerebellar Degenerations Using Diffusion Kurtosis Imaging.

Acad Radiol. 2019;26:e333-e339.

71.   Krismer F, Seppi K, Wenning GK, Papapetropoulos S, Abler V, Goebel G, et al. Abnormalities

on structural MRI associate with faster disease progression in multiple system atrophy. Parkinsonism

Relat Disord. 2019;58:23-27.

72.   Chung SJ, Cho KH, Lee YH, Yoo HS, Baik K, Jung JH, et al. Diffusion tensor imaging-based

pontine damage as a degeneration marker in synucleinopathy. J Neurosci Res. 2021;99:2922-2931.

73.   Abos A, Baggio HC, Segura B, Campabadal A, Uribe C, Giraldo DM, et al. Differentiation of

multiple system atrophy from Parkinson's disease by structural connectivity derived from probabilistic

tractography. Scientific reports. 2019;9:16488.

74.   Vemuri P, Castillo AM, Thostenson KB, Ward CP, Raghavan S, Reid RI, et al. Imaging

biomarkers for early multiple system atrophy. Parkinsonism Relat Disord. 2022;103:60-68.

75.   Nilsson C, Markenroth Bloch K, Brockstedt S, Lätt J, Widner H, Larsson EM. Tracking the

neurodegeneration of parkinsonian disorders--a pilot study. Neuroradiology. 2007;49:111-119.

76.   Chen B, Fan G, Sun W, Shang X, Shi S, Wang S, et al. Usefulness of diffusion-tensor MRI in

the diagnosis of Parkinson variant of multiple system atrophy and Parkinson's disease: a valuable tool

to differentiate between them? Clin Radiol. 2017;72:610.e619-610.e615.

77.   Ito K, Ohtsuka C, Yoshioka K, Kameda H, Yokosawa S, Sato R, et al. Differential diagnosis of

parkinsonism by a combined use of diffusion kurtosis imaging and quantitative susceptibility mapping.

Neuroradiology. 2017;59:759-769.

78.   Zanigni S, Evangelisti S, Testa C, Manners DN, Calandra-Buonaura G, Guarino M, et al.

White matter and cortical changes in atypical parkinsonisms: A multimodal quantitative MR study.

Parkinsonism Relat Disord. 2017;39:44-51.

79.   Beliveau V, Krismer F, Skalla E, Schocke MM, Gizewski ER, Wenning GK, et al.

Characterization and diagnostic potential of diffusion tractography in multiple system atrophy.

Parkinsonism Relat Disord. 2021;85:30-36.

GO terms: Volume-correlated genes

Go terms: FA-correlated genes