Title:

**Early impairment in dentate pattern separation in a rodent model of temporal lobe epilepsy.**

Author list:

**Laura A. Ewell[1,2,3,4], Ario Ramezani[1], Gabrielle Suarez[1], Verónica C. Piatti[1], Vicky Lam[1], Tatyana**

**G. Mills[1], Anelah McGinness[1], Raafat Kuk[1], Mika Ukai[1], Anna-Lena Schlenner[1], Stefan**

**Leutgeb[1,5]*, and Jill K. Leutgeb[1,5,6]***

Affiliations:

[1]Neurobiology Department, School of Biological Sciences, University of California, San Diego,

USA

[2]Department of Anatomy and Neurobiology, University of California, Irvine, USA

[3]Department of Neurobiology and Behavior, University of California, Irvine, USA

[4]Center for the Neurobiology of Learning and Memory, University of California, Irvine, USA

[5]Kavli Institute for Brain and Mind, University of California, San Diego, USA

[6]Lead contact

*Correspondence: jleutgeb@ucsd.edu (J.K.L), sleutgeb@ucsd.edu (S.L.)

**SUMMARY**

The dentate gyrus of the hippocampal formation utilizes sparse neural coding to support

memory formation. In temporal lobe epilepsy (TLE), the dentate undergoes extensive

restructuring, including loss of local inhibition. It is unknown how circuit changes impact

dentate network computations in awake behaving animals. Here we perform high-density

tetrode recordings from the dentate in male rats treated with systemic kainic acid (KA) to

induce TLE. In KA-treated rats, dentate spatial representations were less precise at the single

neuron level, however the stability of the spatial code in representing repeated visits to the

same location was retained at the population level. Despite spared spatial representation, the

network computations for pattern separation were disrupted. The reorganized dentate

network in KA-treated rats generated less distinct population activity patterns for different

environments, specifically as a result of impaired rate-remapping. Changes in network

computations paralleled impairments in a dentate-dependent memory task, potentially

providing a link between altered pattern separation and behavioral discrimination. These

deficits were present in animals with confirmed dentate circuit reorganization that had not yet

developed spontaneous seizures – suggesting that the dentate network changes underlying

altered rate-remapping and impaired dentate-dependent memory occur early in

epileptogenesis.

**Keywords:** dentate gyrus, medial temporal lobe epilepsy, pattern separation, hippocampus,

memory comorbidity

**INTRODUCTION**

A critical feature of episodic memory is the ability of neural networks in the hippocampus to represent different events uniquely such that they can be stored and later remembered with minimal interference from memories that share common features. Accurate memory retrieval is thus dependent on encoding memories distinctly when they are formed. The neuronal mechanism proposed to mediate this process is pattern separation, in which the pattern separator receives overlapping input representing similar experiences, and transforms those inputs to non-overlapping, separated outputs such that a memory is represented by a distinct pattern of activity [1-3]. Experimental evidence converges to suggest that the first processing station of the hippocampal trisynaptic circuit, the dentate gyrus, is an optimal pattern separator, confirmed both in rodent models [2,4-7] and in humans [5,8,9]. The dentate gyrus is well positioned to perform this operation based on unique network cytoarchitecture, plasticity, and neuronal activity [10,11]. The dentate gyrus receives dense cortical input from higher cortical areas and sends output (mossy fibers) to the downstream CA3, the auto-associative recurrent network at the core of the hippocampal formation which operates most efficiently for memory encoding when incoming patterns are maximally separated by the dentate gyrus [1,12].

At the heart of the capacity for the dentate gyrus neural network to perform pattern separation is its unique sparsity unlike other neural networks in memory circuits of the brain. An extremely sparse neuronal population code is utilized to separate experiences [13,14,15]. Such sparse activation selection is achieved in part by the large network expansion between the cortex (relatively low numbers of neurons in layer 2 of entorhinal cortex) and the much larger

dentate gyrus network [16] . Another essential component is strong feedforward [17] and feedback

inhibition [18] that are present within dentate gyrus microcircuits, which effectively silence the

majority of neurons, enabling a highly competitive winner take all coding scheme [19] and

support sub-second toggling between distinct co-active ensembles [20].

Further capacity to generate unique population activity is achieved by combining a

distributed cell activation code with a distributed rate code for representing distinct

experiences. The active subset of dentate neurons vary their firing rates (rate-remapping) or

spatial receptive fields (spatial remapping) to represent different features of similar experiences

[2]. These collective neuronal changes magnify memory capacity by expanding the possible

unique combinations of neuronal activity at the population level [21,22]. These network dynamics

were described in *in vivo* electrophysiological recordings in the dentate gyrus examining the

neural mechanism of pattern separation in rodents [2,4,6], and can serve as important windows

into memory processes. With this mechanistic understanding, the neuronal mechanisms for

pattern separation can be assessed as a readout of proper memory encoding and storage of

precise spatial memory.

There are several lines of evidence indicating that reduced sparsity in the dentate gyrus

weakens pattern separation and leads to memory deficits [23-26]. Perhaps the most significant line

of evidence comes from temporal lobe epilepsy (TLE), a disease defined by increased neuronal

activity and synchrony, and thus reduced sparsity. Patients with TLE are unable to pattern

separate similar visual scenes [27] and mouse models of TLE exhibit reduced *in vitro* dentate

pattern separation [28]. Several changes occur in the dentate gyrus network in epilepsy that could

account for dentate-dependent memory deficits. Granule cells sprout de novo axons which

create aberrant connections between granule cells [29-31]. Such increased recurrence has been

hypothesized to promote hypersynchronous activation of granule cells, promoting pathological

high frequency events [32]. Other changes include death of inhibitory interneurons [33,34], which

lead to reduced inhibition onto granule cells [35]. Finally, granule cells exhibit changes in dendritic

and synaptic properties which alter the plasticity rules of the few active neurons [36,37]. Together,

these changes can facilitate overactivation of dentate granule cells [38], which leads to runaway

excitation and seizure propagation [39]. It is currently unknown how changes in (1) network

recurrence, (2) network inhibition, and (3) seizure burden, impact the neural-network

computations in the dentate gyrus that underlie pattern separation and ultimately memory. To

address these questions, we first assessed circuit level structural changes and dentate-

dependent memory [40,41] in rats with ongoing epileptogenesis prior to seizure generation as well

as in rats with chronic epilepsy. At the same time-point in disease progression, we performed

high-density single unit recordings from the dentate gyrus in a task where the network

mechanisms of pattern separation by place field rate-remapping have been extensively

characterized. We report that even prior to developing seizures, rats had impaired dentate-

dependent memory. This behavioral deficit emerged at a timepoint when we show that the

dentate network experienced loss of inhibitory neurons, altered spatial representations and

reduced specificity in the population code via reduced rate-remapping between similar

environments.

**RESULTS**

**Kainic acid induced status epilepticus results in structural changes of the dentate gyrus and**

**impaired behavioral discrimination**

To test how circuit reorganization in the dentate gyrus network as a result of

epileptogenesis and eventually epilepsy impacts dentate-dependent behavior, we induced

*status epilepticus* in male rats utilizing the low-dose systemic kainic acid (KA) model while

littermates served as controls. Two months after rats were treated with KA, they were

continuously video monitored (24 hours a day) for one month to determine disease progression

(those with >2 spontaneous behavioral seizures were classified as epileptic) and then were

sacrificed for histological analysis. At this timepoint, roughly half of the treated rats were

epileptic, which allowed us to track which changes in the dentate gyrus were associated with

seizures (epilepsy) versus associated with epileptogenesis. In both control and KA-induced rat

populations, we measured two hallmarks of epilepsy that could have direct impacts on dentate

network sparsity; mossy fiber sprouting and hilar somatostatin positive (SST+) interneuron cell

loss (Figure 1A-D). Mossy fibers are the excitatory output axons of dentate granule cells and

SST+ neurons are interneurons that provide feedback inhibition to granule cells. Mossy fiber

sprouting was elevated in KA-treated rats that suffered behavioral seizures, but not in those

that were seizure free (Figure 1A,C; Figure S1) (ctrl: n = 28 rats, mean ± S.E.M., 0.7 ± 0.1; KA-no

seizures: n = 15 rats, 1.1 ± 0.2; KA-seizures: n = 13 rats, 2.1 ± 0.2; one-way ANOVA, df = 2, f =

18.6, p = $7.5 * 10^{-7}$, multiple comparisons: ctrl vs KA-no seizures, n.s., ctrl vs KA-seizures, p = 3.9

$* 10^{-7}$, seizures vs no seizures, p = 0.0006). We processed a subset of the tissue to stain for SST+

neurons. The number of SST+ neurons in the dorsal hilus were reduced in both groups of KA-

treated rats, regardless of whether they had yet developed spontaneous behavioral seizures

(Figure 1 B,D; ctrl: n = 10 rats, mean ± S.E.M., 2525 ± 271 SST+; KA-no seizures: n = 6 rats, 1183

± 268 SST+; KA-seizures: n = 7 rats, 1371 ± 110 SST+, one-way ANOVA, df = 2, f = 9.52, p = 0.001,

multiple comparisons: ctrl vs KA-no seizures, p = 0.005; ctrl vs KA-seizures, p = 0.0009, seizures

vs no seizures, n.s.).

A subset of the KA induced animals (n = 18) were tested on a dentate-dependent spatial

discrimination task during the period that they were being video monitored. In this reference

memory task, on each day of training animals choose between two adjacent open arms of an 8-

arm radial arm maze, one of which is consistently baited with a food reward (Figure 1E). To

confirm dentate dependence of this task, we added an additional group of animals in which we

lesioned the dentate gyrus [42]. Dorsal dentate granule cells were selectively lesioned with

bilateral infusions of colchicine in the dorsal region of the dentate gyrus of the hippocampus. In

the lesioned rats (n = 11), there was a 67.3 ± 4.9% reduction in the volume of the dorsal dentate

gyrus granule cell layer, as quantified using the Cavalieri method. We found that lesioning the

dentate was sufficient to disrupt learning (requiring more trials to reach criterion) and

strikingly, KA-treated rats behaved as poorly as the lesion group (Figure 1F) (ctrl: 70 ± 5 trials, n

= 30 rats; KA: 95 ± 10 trials, n = 18 rats; DG Lesion: 105 ± 15 trials, n = 11, one-way ANOVA, df =

2,f = 6.27, p = 0.003; multiple comparisons: ctrl vs KA, p = 0.03, ctrl vs lesion, p = 0.009, KA vs

lesion, p = 0.7). Within the KA group, there was no behavioral difference between KA animals

with or without seizures (seizures: 110 ± 20 trials, n = 7 rats; no seizures: 90 ± 10 trials, n = 11,

unpaired t-test, t = 1.19, df = 16, p = 0.25), suggesting that brain changes associated with

epileptogenesis were sufficient to disrupt dentate-dependent memory. Collectively, these data

imply that early changes to the dentate gyrus network during disease progression, i.e. before

the emergence of the spontaneous seizures that define epilepsy, are sufficient to alter the

network computations necessary to support behaviors that require fine spatial discrimination.

**Impaired spatial information and reduced spatial sparsity in the neural activity of KA-treated**

**rats**

To link how disease related circuit reorganization in the dentate gyrus resulted in

altered dentate-dependent memory we next focused on characterizing changes in dentate

network activity in KA treated rats. We first asked whether structural changes such as loss of

inhibition and increases in excitatory connectivity in epileptogenesis are accompanied by

physiological changes in dentate neurons in awake-behaving rats. We recorded activity from

dentate principal neurons while rats explored distinct spatial environments, a condition where

dentate neurons in the healthy brain were previously shown to represent different

environments with unique firing patterns [2,4,6]. Rats were trained to forage for food in open-field

environments that differed in shape (Ctrl: n = 39 neurons, n = 6 rats; KA: n = 60 neurons, n = 6

rats). We tested whether there were general differences in dentate cellular excitability between

control animals and those that had undergone KA induced *status epilepticus* 2 months prior.

Distributions of peak firing rates during foraging were similar between the two groups (Median,

IQR; Ctrl: 2.2 Hz, 0.3 - 5.6 Hz; KA: 2.0 Hz, 0.5 - 7.8 Hz). Neurons with peak firing rates >2 Hz were

considered active during foraging and comprised a similar proportion of total neurons in control

and KA-treated rats (Ctrl, n = 22 neurons or 56%; KA, n = 30 neurons or 50%; Figure 2B). Of the

active neurons (>2 Hz), KA-treated dentate neurons had similar average firing rates (M, IQR: 0.9

Hz, 0.6 - 1.7 Hz) compared to control neurons (M, IQR: 0.3 Hz, 0.2 - 0.6 Hz; p = 0.13, rank-sum;

z-val: -1.5; Figure 2C). Despite similar firing rate distributions, we observed differences in the

spatial receptive fields of active cells, measured by assessing the quality of their spatial

representations while rats explored the open field. In control rats, even neurons with low

spatial information scores (a measure of how well a neuron's firing rate predicts the rat's

position) exhibited well defined place fields, the difference to higher scores being that cells with

lower spatial information had multiple rather than single well defined firing fields within the

same environment (Figure 2D, left, bottom row). In contrast, in KA-treated rats, cells had spatial

information scores that were below the control range and the cells with the lowest scores were

characterized by distributed firing patterns such that the neurons fired across most spatial bins

in the open-field (Figure 2D, right, bottom row). In line with this observation, across the

population of active neurons, spatial information was lower in KA-treated rats (M, IQR: 0.8, 0.4

- 1.7) compared to control rats (M, IQR: 1.8, 1.2 - 2.2; p = 0.0056; rank-sum, z-val: 2.8; Figure 2E,

left). Furthermore, we observed differences in spatial non-selectivity (a measure of the

proportion of spatial bins within the environment that the neuron was active in). Across the

population, KA-treated dentate neurons exhibited higher spatial non-selectivity (KA-treated

neurons: M, IQR, 0.42, 0.19 - 0.60 compared to control neurons: M, IQR, 0.19, 0.15 - 0.29; p =

0.017, rank-sum, z-val: -2.4; Figure 2E, right). These results confirm that the spatial firing

patterns of dentate neurons following epileptogenesis are altered despite retained average

firing rates. Together, the reduced spatial information and increased spatial non-selectivity

indicate that in KA-treated rats, at each location within an open-field, a higher number of

dentate neurons are active. Therefore, during an experience at any given moment in time, the

dentate network is indeed less sparse in KA-treated rats in comparison to controls. We

observed behavioral seizures (Racine III-V) in four of the six KA-treated rats (4 hrs/day of video

monitoring, see methods for more details). We therefore asked whether the observed changes

in place field quality were driven by neurons recorded in animals with observed behavioral

seizures. When testing between neurons recorded from KA-treated rats with seizures (n = 17)

vs. without seizures (n = 13), we found no differences in average firing rate (with seizures: 0.6,

0.3  - 1.6; without seizures:  0.3, 0.2-1.2, rank-sum, p = 0.4, z-val:-0.8; Figure S2A), spatial

information (with seizures: 0.8, 0.4 – 1.7; without seizures:  0.8, 0.6-1.3, rank-sum, p = 0.77, z-

val: 0.29; Figure S2B), or spatial non-selectivity (with seizures: 0.5, 0.2-0.7, without seizures:

0.4,0.3-0.5, rank-sum, p = 0.62, z-val: -0.5; Figure S2C), suggesting changes in neuronal firing

patterns occurred prior to seizure generation. We therefore pooled all KA-treated animals for

further electrophysiological analysis.

One feature of neurons in the dentate gyrus is that individual place cells often have

many distinct spatial firing fields. Multiple place fields per cell endow the sparse network with

the capacity to create as many distinct population codes with fewer active cells as a much

larger network of active cells can generate from fewer place fields per cell [2]. We observed that

the number of fields per neuron and the size of fields were not statistically different between

control and KA-treated rats (Figure 2F). Note, however, that we observed a skewed distribution

for number of fields in dentate neurons recorded in KA-treated animals, with a quarter of cells

having >5 fields. KA-treated rats had significantly higher within field mean firing rates,

unmasking a hyperexcitability phenotype that was occluded when considering overall activity

(ctrl: n = 24 fields, 2.3 ± 0.3 Hz; KA: n = 55 fields, 3.1 ± 0.2 Hz, rank-sum, z-val = -2.4, p = 0.014;

Figure 2F, right). Collectively we observed several features of altered dentate neuronal activity

in KA-treated rats that suggests reduced network sparsity manifests early in disease progression

before seizures emerge. These results are in line with our observation that dentate-dependent

memory impairments were also present in animals treated with KA, but prior to the emergence

of behavioral seizures.

**Dentate network computations that encode and represent events distinctly are impaired in**

**KA-treated rats**

An important neural-network computation of the dentate gyrus is to distinguish

between similar input patterns, or events, by generating distinct population activity for each

input that can then be conveyed to downstream hippocampal subregions that are not

optimized for pattern separation [2,43]. Pattern separation is considered a critical network

computation for the encoding of memory that facilitates accurate memory retrieval. To

differentiate between environments, dentate neurons either change their spatial firing fields,

in-field firing rates, or turn on/off such that the neural representations of similar environments

are strongly decorrelated from one another at the population level [21,44].  To test whether

dentate network computations for pattern separation were altered in KA-treated animals, we

trained rats to explore an open field arena with flexible walls that could be in either a square or

circular configuration (Figure 3A). The arena remained in the same spatial location in the room,

with walls placed upon the same section of floor leaving the majority of distal and proximal

cues unchanged between trials except for the shape of the enclosure walls. Trials consisted of

four ten minute-sessions in which box shape varied between a square or circle randomly. This

design allowed us to test pattern separation by comparing dentate population activity in the

same location between visits when the same environmental cues were repeated (same shape)

and when the environmental cues differed (different shape) (Figure 3B).

To make these comparisons, we constructed population vectors that allow for testing of

dentate network decorrelation between environments. Population vectors include both the

spatial aspect (where in the environment the neurons are active) and the rate aspect (the

magnitude of the activity in each spatial bin) for the entire population of active cells in the

network [21] (Figure 3C). Place maps from each neuron were generated for each of the four

sessions (i.e., two for each of the two environment shapes). The maps of neurons that were

active in at least one of the shapes were stacked into a 3-dimensional matrix with x and y

corresponding to matching spatial bins and z corresponding to cell identity. Pairs of vectors for

each spatial bin(x,y), comprising the average rate for every neuron within that bin, were

correlated across either maps from the same shapes (square-square, circle-circle) or different

shapes (square-circle). First, we confirmed previous results [2] for dentate network computations

in control animals (Figure 3D). In control, same shape PV correlations were high, reflecting the

same population activity in repeated visits to the same locations (correlation = 0.79 ± 0.009). As

expected, different shape comparisons plunged to low correlation values reflecting different

patterns of activity associated with different environments (correlation = 0.31 ± 0.01, reduced

by 0.48 ± 0.01). In KA-treated rats, same shape PV correlations were also high (correlation =

0.82 ± 0.01), but were reduced to a lesser extent across different shapes (correlation 0.57 ±

0.01, reduced by 0.25 ± 0.01, p = 5 * $10^{-61}$, unpaired t-test; t stat: 17.05). Thus, the

discrimination of different environmental cues was diminished for dentate networks of KA-

treated rats compared to control rats.

Next, we further explored the nature of the network discrimination deficits in KA-

treated rats to determine whether changes in spatial representations or rate distributions were

differentially affected in the KA-treated dentate network. Spatial correlations between the

place maps generated from active dentate neurons across same-shape comparisons (circle-

circle and square-square) were high for control and KA-treated rats, indicating robust stability

of spatial representations for both groups (mean ± S.E.M.; Ctrl: n = 36 comparisons, 0.69 ± 0.04;

KA: n = 40 comparisons, 0.57 ± 0.05). Furthermore, a correlation of place cell maps between

different environments yielded significantly lower values than for the across-shape comparison

for both groups (one-way ANOVA, df = 3, f = 17.39, p = 3.2 * $10^{-10}$, p<0.001, post-hoc Tukey-

Cramer correction for group comparisons, mean ± S.E.M.; Ctrl: n = 76 comparisons, 0.31 ± 0.03,

p = 4.0 * $10^{-8}$; KA: n = 86 comparisons, 0.31 ± 0.04, p = 4.0 * $10^{-4}$; Figure 3E, left). In contrast to

the similar response of spatial representations in control versus KA-treated rats, we observed a

difference in rate discrimination by place fields (Figure 3E, right). Note that rate analyses were

done on individual place fields rather than the full spatial map of an environment to account for

the independent rate code of individual firing fields that have been shown to represent changes

in non-spatial cues across environments [2]. In control rats, the firing rates within place fields

were similar between visits to the same environment, but more distinct between visits to

different environments, as would be expected when changes in rate is used to discriminate

between different environmental cues (reduction of 0.3 ± 0.05, n = 24 fields, sign-rank, z-val =

4.01, p = 5.9 * $10^{-5}$). KA-treated rats also exhibited a higher rate difference between different

compared to same environments (reduction of 0.12 ± 0.04, n = 55 fields, sign-rank, z-val = 2.1, p

= 0.03). However, the magnitude of the rate change between same vs. different was

significantly larger in controls compared to KA-treated rats (rank-sum, z-val = 3.2, p = 0.0013).

Taken together, our results suggest that pattern separation supported by a redistribution of

firing rates, a mechanism for the generation of unique population codes for distinct events, is

selectively more sensitive to changes in the dentate network as a result of epileptogenesis.

**DISCUSSION**

The dentate gyrus undergoes massive restructuring in TLE, yet there have been few studies

examining how neural network computations of the dentate gyrus may or may not be

compromised. In line with previous work, we observed two histological changes in the dentate

gyrus with the induction of epileptogenesis – sprouting of recurrent granule cell axons and loss

of somatostatin interneurons – both of which would be predicted to lead to hyperexcitability of

dentate gyrus neurons and a loss of network sparsity. In parallel we observed a deficit in

dentate-dependent memory along with the degradation of neural-network computations

essential for the dentate gyrus contribution to memory encoding. Epileptogenesis induced

impairment in the spatial coding features of individual dentate neurons including less precise

spatial firing with higher within field firing rates. However, despite a less precise spatial

representation at the single neuron level, the stability of the spatial code for repeated visits to

the same environment was retained at the population level. The degraded pattern separation

in the dentate networks of KA-treated rats for representing distinctly different environments

was instead a result of reduced rate-remapping that impaired the capacity of the reorganized

dentate network to generate distinct population patterns needed to distinguish between

similar familiar environments, which may in turn preclude the cueing of accurate memory

retrieval in downstream neural networks.

In our experiments, KA-treated rats showed interneuron loss, behavioral deficits, and

impaired pattern separation in the dentate even if they had not yet developed the spontaneous

behavioral seizures that define epilepsy. These results suggest that dentate specific memory

deficits such as learning to discriminate adjacent spatial locations may be an early

manifestation of epileptogenic circuit changes that impair select dentate network computations

prior to the development of spontaneous seizures. The timeline in dentate may be different

than in other regions of the hippocampus. For example, in CA1, it has been reported that place

field spatial information reduced concurrently with the progressive increase in spontaneous

seizure frequency and that reductions in spatial stability emerged weeks later [45]. Earlier

impairments in dentate function may provide a biomarker for assessing the risk for developing

acquired epilepsy after brain injury or stroke. Indeed, epileptic patients clearly exhibit impaired

pattern separation [27,28], although to our knowledge no longitudinal study has been done to

assess whether performance on such tasks predict disease outcomes.

An early feature of epileptogenesis is interneuron loss in the hippocampus [46]. Specific to

the dentate gyrus, loss of interneurons in the hilus has been widely reported [35,47]. Even though

remaining interneurons sprout, they are unable to compensate, resulting in a net loss of

inhibition to the dentate network [48]. Confirming these reports, we also report a loss of

interneurons in the dentate hilus – specifically SST+ containing interneurons (Figure 1D). It is

interesting to consider how this early circuit change may be linked to the altered physiology we

observe. Lateral inhibition has been shown to exert a critical circuit level constraint on the

number of neurons active at a particular location [49,50], which may explain the reduced spatial

selectivity we observed in neurons recorded from KA-treated rats. SST+ interneurons are also

known to regulate spike bursting within place fields [51]. Consistent with this role, we observed

increased mean firing rate within fields (Figure 2F). The early influence of SST+ cell loss on

network rate-remapping could be a combination of both direct effects of the reduced inhibition

on select cortical inputs to dentate, as well as the broader effects of reduced inhibition on

dentate network sparsity.

Dentate network sparsity is hypothesized to play a critical role in setting the level of

pattern separation that can be achieved through unique combinations of population activity [14].

The fewer neurons capable of overcoming the strong inhibition characteristic of a competitive

network like the dentate gyrus [19], the more unique patterns of activity can be achieved from

the total number of dentate neurons. The possible number of unique population codes is

hypothesized to be directly related to the memory capacity of the hippocampus [1,12]. Moreover,

it has been shown experimentally that directly altering the inhibitory gain of the dentate

network through the loss of other inhibitory cells results in more neurons becoming active for a

given input representation (reduced sparsity) and impaired pattern separation [52,53]. However,

the loss of SST+ interneurons could have a more direct influence on rate-remapping, the

identified network computation selectively impaired early in epileptogenesis while spatial

coding is retained. Somatostatin interneurons inhibit the distal dendrites of dentate granule

neurons in the outer molecular layer [54]. The outer and inner molecular layers are defined by

select cortical inputs that terminate on distinct regions of dendrite. Input from the lateral

entorhinal cortex (LEC) reaches granule neurons via the lateral perforant path, terminating on

the distal dendrites of granule neurons where SST+ interneurons regulate activity [55]. Input from

the LEC has been shown to convey mainly nonspatial information to hippocampus [56,57] and to

be necessary for rate-remapping to occur in CA3 neural-networks under the same behavioral

conditions tested here [58]. These results suggest that a select disruption of LEC inputs to dentate

gyrus via a reduction in SST+ inhibition could also result in impaired rate-remapping with a

preservation of the spatial code from medial inputs. Although direct tests are needed to reveal

longitudinal timelines and include causal manipulations, our results suggest a direct link

between SST+ loss and an early impairment in dentate network computations for memory

rather than spatial coding, providing a therapeutic target for early intervention and the

potential rescue of memory comorbidities.

The second histological feature we considered as an early feature of epileptogenesis in

the hippocampus was mossy fiber sprouting. We only observed significant levels of sprouting in

animals in which we observed spontaneous seizures. The link between sprouting and

pathological population bursts in the dentate has been previously reported [29,59]. Our data

suggest that mossy fiber sprouting is better associated with seizure outcomes and is preceded

by changes to dentate network computations and dentate-dependent memory deficits. Our

findings do not preclude the possibility that sprouting and seizure activity would not further

exacerbate cognitive problems. In fact, seizure activity has been shown to impact hippocampal

neuronal activity and network computations in additional ways, included changes in spatial

information processing [60] [61] . These results highlight that the disease progression that leads to

dentate dependent memory problems versus seizure susceptibility are on separate trajectories

and may not share common mechanisms.

Together, these data highlight the importance of the dentate gyrus for pattern

separation and underscore its dysfunction in TLE cognitive comorbidities. Impairments in

dentate physiology emerge before spontaneous seizures, indicating that functional assessment

of dentate memory could be a powerful biomarker for determining disease outcomes. Testing

pattern separation in humans has been done at both the imaging and behavioral level [5,8,9],

further motivating the therapeutic importance for our finding that pattern separation

computations in the dentate gyrus are early outcomes of epileptogenesis and precede seizure

outcomes. These results also point to a fundamental role of inhibition in the dentate network

for supporting its function – indicating that interneuron transplant therapies [62], which have had

success in treating cognitive comorbidities after brain injury could be an exciting avenue for

dentate-dependent memory in TLE as well [63].

**Acknowledgements**: We thank M. Wong, Y. An, M. Appalaraju, and M. Andersson for technical

assistance. This work was supported by the Hellman Foundation, Epilepsy foundation grant

157927, NIMH grant R01MH119179, and the Walter F. Heiligenberg Professorship to J.K.

Leutgeb; NIH grants R01 NS102915, R21 MH100354, R01 NS084324, and R01 NS097772 to S.

Leutgeb. L.A. Ewell was supported by NIH training grant F32MH096526.

**Author Contributions**: L.A.E., S.L. and J.K.L. conceived experiments, designed study, and

interpreted data. S.L. and J.K.L. managed the project.  L.A.E., with assistance from L.M., T.G.,

and A-L.S, performed electrophysiological experiments. L.A.E., G.S., and A.M. scored seizure

monitoring, G.S. and A.M. performed immunohistochemistry and microscopy, V.P. provided

dentate lesion data, and A-L.S. quantified the lesion data. L.A.E., G.S., A.M., R.K., and M.K.

performed behavioral experiments. L.A.E. analyzed data and prepared figures. L.A.E. prepared

the manuscript with feedback and editing from J.K.L. and S.L.

**Ethics Declarations**: The authors declare no competing financial interests.

**METHODS**

**Subjects**

All experimental procedures were performed as approved by the Institutional Animal Care and

Use Committee at the University of California, San Diego and according to National Institutes of

Health and institutional guidelines. Epilepsy was induced as in previous studies [64] [61] using a

repeated low-dose kainate chronic model of temporal lobe epilepsy [65]. Male Long Evans rats (n

= 52, 40 days of age, Charles River, CA) were treated with kainic acid (KA) (5 mg/kg, i.p., Tocris)

each hour until the onset of status epilepticus, which was defined as >10 motor seizures per

hour of class IV or V on the Racine scale [66]. After the seizure induction protocol, the rats were

housed individually and maintained on a reverse 12 hr light/12 hr dark cycle. To confirm

whether chronic spontaneous seizures that define epilepsy had developed, rats were video-

monitored for 4 hours per day for electrophysiological studies or 24 hours per day for

behavioral and histological studies. Once two or more motor seizures of class III or greater on

the Racine scale were observed, the animal was considered epileptic. Twenty-eight KA-treated

rats (3.5 – 4.5 months old) were continuously video monitored for 1 month and paired with 28

control rats not injected with KA (4 months old), after which they were sacrificed for

histological analysis of circuit reorganization. Six KA-treated rats (6-12 months old) and six

control rats not injected with KA (6 -12 months old) were used for electrophysiological

experiments. Two of the control rats used for electrophysiological experiments were wildtype

GFAP-TK rats on a Long Evans background [67]. Eighteen KA-treated rats (6-8 months old) and 30

control rats not injected with KA (6-8 months old) were used for behavioral experiments.

Eleven wildtype GFAP-TK rats on a Long Evans background were injected with colchicine to

lesion the dentate gyrus and were used for behavioral experiments. The transgenic GFAP-TK rat

line was originally provided by Dr. Heather A. Cameron (National Institutes of Health) and

maintained by breeding with wildtype Long Evans rats obtained from Charles River Laboratory.

**Surgical procedures - chronic electrophysiological recording device**

Rats were anesthetized with isoflurane (2% to 2.5% in $O_2$) and an electrode assembly

(Hyperdrive, designed by the McNaughton laboratory [68]) comprising 14 independently movable

tetrodes was implanted above the right hippocampus (AP, 4.1 mm posterior to bregma; ML, 3.0

mm) and fixed to the skull using stainless steel screws and dental cement. Two screws were

used as animal ground and were implanted to touch the surface of cortex, anterior and lateral

to bregma. Tetrodes were prepared by twisting four insulated platinum wires together

(diameter = 0.017 mm, California Fine Wire Company). Leads were plated with platinum prior

to surgery to obtain stable impedances near 200 kOhm. Two tetrodes had all four leads shorted

with one of the two left in the cortex to record a differential reference signal, while the other

was advanced to stratum radiatum to record local field potentials (LFP). The other 12 tetrodes

were positioned in the dentate gyrus cell layer to record single units.

**Tetrode recording locations**

Histological sections through segments of the hippocampus with electrode tracks were

collected as described below (see Histology section). All tetrodes of the 14-tetrode bundle were

identified by matching tetrode positions in the bundle with electrode tracks (small lesions) in

postmortem tissue. Recordings from a tetrode were included in the data analysis if the

tetrode's deepest position could be clearly identified within or adjacent to (touching) the

dentate gyrus granule cell layer (see Figure 2A).

**Surgical procedures  - dentate gyrus lesions**

Dentate gyrus lesions were performed as described previously [69]. At the time of surgery, the rat

was anesthetized with isoflurane gas [2- 2.5% in $O_2$ (20 mL /1 L per minute)] and buprenorphine

(0.05 mg/kg, s.c.) was given as an analgesic. To create a lesion specific to granule cells of the

dorsal dentate gyrus, colchicine (2.5 mg/ml in 0.1M phosphate buffer, pH 7.2) was injected into

3 sites (0.04 µl per site at 0.012 µl per minute) in both hemispheres at the following coordinates

anterior-posterior, medio-lateral from bregma and ventral from dura (in mm): (1) -2.8, ± 0.9, -

3.7 (2) -4.0, ± 2.1, -3.6, (3) -5.2, ± 3.2, -3.7. Three animals received injections only at sites (1)

and (2). Infusions were made using a 1 µl Hamilton syringe and the needle was left in place for

8 min at each injection site to prevent spread. Colchicine has previously been shown to cause

selective cell death of dentate granule cells [70][71]. Using computer-based 3D volumetric analysis,

the volume of the dorsal dentate gyrus granule cell layer and dorsal CA1 and CA3 pyramidal cell

layers was quantified in cresyl violet stained, 40 µm thick brain sections. To assess the

completeness of the dorsal dentate gyrus lesions and to assess damage outside the dentate

gyrus, the Cavalieri estimator in the Stereo Investigator software (grid size 20 µm; 10x

objective) was used for volume quantification. The remaining granule cell volume was

normalized to a control volume of 1.54 * $10^9$ μm³ (n = 3 rats). The remaining granule cell

percent volume was low and comparable in the two dorsal hemispheres (left: 29.9 ± 4.7 %

remaining, right: 35.4 ± 5.7 % remaining). Only minor mechanical damage was observed in the

CA1 region above the dentate injection site. Similar damage was previously shown to not

significantly contribute to dentate-dependent memory impairment [69], or an impairment in CA1-

dependent memory [40].

**Recordings and data acquisition**

Single units and LFP were recorded during restful and running behavior using a data acquisition

system (Digital Lynx SX; Neuralynx) with 64 digitally programmable amplifiers. Unit activity was

amplified and band-pass filtered between 0.6 and 6 kHz. Spike waveforms above a trigger

threshold (40 μV) were time stamped and digitized at 32 kHz for 1 ms. The LFP was recorded

continuously in the 0.1 - 900 Hz band from one of the wires of each tetrode. All recording

sessions occurred during the animal's dark cycle (between 7 am - 7 pm). Rats were food

deprived to 85% of their baseline weight and trained to forage for randomly scattered food

reward (chocolate cereal crumbs) in open arenas. Open arenas were enclosed by a set of

flexible walls that could either be in a circular formation (diameter = 1.0 m, height = 0.5 m) or a

square formation (0.8 × 0.8 m, height = 0.5 m). Behavioral sessions consisted of four 10 min

foraging epochs preceded and followed by resting periods. The rats were also allowed to rest in

a holding box for 5 min between foraging epochs. For position tracking, light emitting diodes on

the head-mounted preamplifier were tracked at 30 Hz by processing video images.

**Cell clustering**

Single units were manually sorted using MClust (MClust 3.5, written by A. David Redish;

http://redishlab.neuroscience.umn.edu/MClust/MClust.html) adapted to aid in the tracking of

cell identity over extended periods of time [72]. Clusters that persisted in the same region of

parameter space throughout a behavioral session were accepted for analysis, such that the

action potentials of a given neuron included in the analysis did not change shape, were distinct

from noise signals, and remained inside the cluster boundary. The cluster was required to be

clearly defined in both the first and last rest period of the recording session, but was not

required to be present in individual foraging epochs, as many dentate gyrus single units are

silent within a particular environment.

**Neuronal firing rate map**

For each neuron, we constructed firing rate maps for each behavioral session by summing the

total number of spikes that occurred in a given location bin (5 × 5 cm), dividing by the total

amount of time that the rat occupied the bin, and smoothing with a 5 × 5 bin Gaussian filter

with a standard deviation of approximately one bin:

[0.0025 0.0125 0.0200 0.0125 0.0025;

0.0125 0.0625 0.1000 0.0625 0.0125;

0.0200 0.1000 0.1600 0.1000 0.0200;

0.0125 0.0625 0.1000 0.0625 0.0125;

0.0025 0.0125 0.0200 0.0125 0.0025].

Bins that were never within a distance of less than 2.5 cm from the tracked path or with total

occupancy of less than 150 ms were regarded as unvisited and were not included in the rate

map. Spatial firing patterns of individual cells were compared between sessions when that cell

was active (>2 Hz in at least one of the sessions). For each cell, the comparison was performed

by calculating the Pearson correlation coefficient for firing rates in corresponding spatial bins

across pairs of spatial maps.

**Active neurons**

The total number of single units was determined as the number that were tracked across

resting and foraging behavior. All included single units were active during the first and last

resting period (validating their stability), but not all single units had high firing rates during

foraging epochs. Thus, we determine the proportion of active neurons in behavior as the

number of neurons that had at least one neuronal firing rate map with a peak >2 Hz divided by

the total number of single units recorded (neurons that were clearly isolated and active in sleep

sessions before and after running sessions).

**Spatial information score**

Spatial information was calculated for each neuron for each foraging session in which the

neuron had a peak firing rate of at least 2 Hz in at least one spatial bin. We calculated the

spatial information per spike for each firing rate map as: $I = \sum_i P_i \frac{R_i}{R} log_2 \frac{R_i}{R}$ where *i* indexes

the spatial bins, $P_i$ is the probability of occupancy in each bin, $R_i$ is the mean firing rate in each

bin, and R is the mean firing rate across the spatial map [73].

**Place field boundaries**

Individual spatial firing fields of neurons were isolated as previously described [72]. A reference

map was calculated as the average of the common area of the four rate maps corresponding to

the four behavioral sessions. For each firing peak in the rate map (>0.8 Hz), contours were

calculated at 20 levels between zero and the peak rate. For overlapping contours between

fields, each shared contour was divided into segments at inflection points of the contour, and

each segment was assigned to the nearest field. After fields were determined, only fields with

peak rates >2 Hz and at least 50 mm$^2$ were included for further analysis.

**Population vector correlations**

For each behavioral session, rate vectors were constructed by arranging the spatial maps of all

cells recorded from that region from all animals in an x-y-z stack, where x and y represent the

two spatial dimensions and z represents the cell-identity of the population [44] [21]. To allow for

comparisons between the square and circular enclosure shapes, the analysis was restricted to

the 16 by 16 bins that were common to both shapes, yielding 256 x-y locations. Population

vector correlations were obtained by calculating, for each x-y location, the Pearson correlation

coefficient for firing rates along the z-dimension between pairs of sessions. Cells with firing

below 2 Hz in all x-y bins of the two sessions that were compared were excluded from the

population vectors before calculating the correlation coefficients. The correlation coefficients

from same shape comparisons (circle-circle and square-square) of all spatial bins were averaged

to estimate the population vector correlation for a pairwise comparison between same

sessions. The average same shape correlation was subtracted from each different shape

comparison correlation to quantify the extent of decorrelation in distinct environments.

**544    Dentate-dependent behavior**

We adopted a behavioral discrimination task that is dependent on the dentate gyrus. Rats

learned to discriminate between adjacent arms on an 8-Arm Radial Maze [42]. The maze was in a

room with spatial cues on the walls (not on the maze). The maze was comprised of 8 arms

arranged around an octagonal stem. Each arm was spaced 45 degrees apart from its adjacent

arms. The maze measured approximately 188 centimeters in diameter and was elevated

approximately 36 centimeters above the ground. Transparent Plexiglas walls were placed

around the central stem of the maze and attached with Velcro in order to block unused arms

and designate the two arms that were open per trial. For the adjacent arm discrimination task

open arms would either be arm 1 and 2 or arm 1 and 8. Uniform, chrome, silver bowls were

taped to the end of each arm. The bowl of the reward arm (arm 1) was the location of the food

reward (chocolate cereal) for all behavior testing performed during this experiment. Rats were

food restricted to 85% of their baseline weight and maintained at that weight for the duration

of habituation and testing. Habituation was conducted in three phases. The first phase involved

placement of a single food reward at the stem of the maze, midway down the three

experimental arms, just outside the silver bowls, and inside each bowl. The animal was given 10

minutes to explore during this phase of habituation. If all 10 rewards were eaten within the 10-

minute time frame, the rat moved to phase 2. For phase 2, reward was placed within the bowls

and just outside the bowls of the same three arms. Eating all rewards within 10 minutes

allowed the rat to pass to phase 3, in which reward was only in the three bowls. Rats had 5

minutes to eat all rewards to move onto testing. During testing, the rat ran 10 trials per day. For

each trial two adjacent arms were open (1 and 2) or (1 and 8). Each adjacent, non-rewarded

arm was used 5 times among the 10 trials in a pseudo-random order to ensure that the rat was

not relying on a right-left strategy when making its decision. Reward was always at arm 1. For

each day, the number of correct trials were counted. In between trials, the rat was enclosed

under an opaque bucket in the central stem. All arms were washed with water and dried

between trails. Testing continued until the rat reached a criterion of 90% correct trials for 2 of 3

consecutive days, or for a maximum of 17 days. The total number of trials to reach criterion

were summed. Initially, behavioral testing was done in rats that were not monitored 24 hours

for behavioral seizures. The behavioral performance of non-monitored KA-treated animals did

not differ from video-monitored KA-treated animals (non-monitored: 90 ± 10 trials, n = 8 rats;

monitored: 95 ± 10 trials, n = 18 rats; unpaired t-test, t = -0.45, df = 24, p = 0.66), therefore data

were pooled. Among monitored KA-treated animals, those that had seizures had similar

performance to those who did not (seizures: 110 ± 20 trials, n = 7 rats; no seizures: 90 ± 10

trials, n = 11; unpaired t-test, t = 1.19, df = 16, p = 0.25), therefore data were pooled. Control

animals from lesion experiments were 5 months old, while control animals from epilepsy

experiments were 6 months old. No statistical difference in behavioral performance was

observed between control groups (epilepsy control: 70 ± 5 trials, n = 21 rats; lesion control: 60 ±

5 trials, n = 9 rats; unpaired t-test, t = 1.18, df = 28, p = 0.25), therefore data were pooled.

**Histology**

The rat was deeply anesthetized with isoflurane and given a lethal injection of Pentobarbital

(0.8 mL - 2 mL). Rats were systemically perfused, first with 0.37% sulphide solution and then

with 4% paraformaldehyde (PFA). Brains were transferred to 30% sucrose in 1X PBS before

sectioning with a frozen microtome (40 μM sections, Leica).

*Somatostatin immunohistochemistry:* Every sixth, coronal section (40 μm) was processed for

staining. Sections were washed (3 x 5 min) in 1X PBS. The free-floating sections were then

blocked for 1 hour in blocking solution at room temperature (10% horse serum (HS)

and 0.3% Triton-X in PBS). After blocking, the sections were incubated in diluted 1:300

monoclonal mouse anti-somatostatin antibody (GeneTex, code GTX71935, clone SOM-018)

with 10% HS and 0.3% Triton-X in PBS over two nights at 4°C. After washing (3 x 5 min) in 1 x

PBS, the sections were then incubated for 1 hour at room temperature in diluted (1:200)

biotinylated horse anti-mouse IgG antibody (Vector BA-2000) with 0.3% Triton-X in PBS, and

then washed again. Endogenous peroxide activity was blocked with 0.3% $H_2O_2$ for 30 minutes at

room temperature. The sections were washed again and then incubated with the avidin-biotin-

peroxidase technique (Vectastain ABC Kit, Vector Labs, USA) for 30 minutes at room

temperature. The complex was developed using 0.05% of 3,3'-diaminobenxidine (DAB)

tetrahydrochloride, 0.3% $NiCl_2 \cdot 6H_2O$ stock, and 0.05% hydrogen peroxide as a substrate

for the peroxidase reaction. The sections were then mounted onto Fisherbrand Superfrost Plus

Microscope Slides. Stereology of the SST-immunostained tissue was performed using a Leica

CTR 6000 microscope and the optical fractionator method and the Cavalieri principle on Stereo

Investigator 10 (MBF Bioscience). The counting frame was 300 μm × 300 μm and the counting

grid was also 300 μm × 300 μm. After counting SST interneurons in the dorsal hilus of each rat,

the total SST interneuron population was estimated based on the thickness of the mounted

sections (40 μm).

*TIMM staining:* Sections were mounted onto Fisherbrand Superfrost Plus Microscope Slides

with a gelatin and PBS mounting solution and left to dry overnight. Slides were placed in

distilled water for 3 minutes and then incubated in developer solution in the dark. The

developer solution is: 120 mL of Gum Arabic (100 g/200 mL), 10 mL of filtered citrate buffer

(51g $C_6H_8O_7$/200 mL; 47g $C_6H_5Na_3$-$O_7$ - $2H_2O$/200 mL), 60 mL of 1.7% hydroquinone solution,

and 1 mL of 0.09% light-sensitive, silver nitrate solution. After 30 minutes of incubation, the

developer solution was mixed and the tissue was re-submerged for at least another 30 minutes.

If the stain was not dark enough, the tissue was left in solution until the stain was complete.

The slides were then dipped twice in warm distilled water (50°C) and placed in room

temperature distilled water for 15 to 30 minutes. Next, the slides were fixed in 5% sodium

thiosulfat pentahydrat for 30 seconds. Slides were dehydrated in an alcohol series and

coverslipped with permount. The tissue for a subset of animals used for electrophysiological

experiments were processed for TIMM staining and subsequently stained with cresyl violet to

facilitate tetrode tracking [74]. For TIMM scoring, the scorer was blinded to condition and scored

the extent of mossy fiber sprouting according to the criteria described in [75]. Every sixth section

through the dorsal dentate gyrus was given a TIMM score. Sections with no or only occasional

supragranular mossy fiber-like staining were scored 0. Sections with scattered mossy fiber-like

staining above all parts of the granular layer were given a score of 1. Sections that exhibited

either patches of heavy mossy fiber-like staining interspersed with regions of sparser staining or

a continuous band of staining intermediate in intensity between sections scored 1 and 3 were

given a score of 2. Finally, sections with a dense, continuous band of supragranular mossy fiber-

like staining were assigned a score of 3. For analysis, the worst score was selected to represent

the maximum extent of sprouting that was observed in a given rat.

**Statistics**

For each statistical comparison, normality was assessed with Lilliefors test and either

parametric or non-parametric tests were chosen accordingly. Multiple comparisons were made

with Tukey Kramer post-hoc tests. Throughout the text values are presented as mean ± S.E.M.

or median and inter-quartile interval.

 **FIGURE LEGENDS**

**Figure 1.  Dentate circuit reorganization and dentate-dependent memory impairment after**

**kainic acid treatment. A,** Coronal sections of hippocampi from two KA-treated rats with

different degrees of axonal sprouting. Timm stains label zinc concentration black, which is

selectively dense in the mossy fiber axons of dentate granule neurons. Magnified images to the

right are selected from the area defined by the black boxes centered in the hilar region of the

dentate gyrus in the images to the left. Top row, minor sprouting (Timm score 1). Bottom row,

intermediate sprouting (Timm score 2). Arrows indicate an example area of sprouting in the

inner molecular layer. See Figure S1 for more examples. Scale bars = 250 $\mu$m. DG: dentate

gyrus; GCL: granule cell layer. **B,** Representative coronal sections of the dentate hilar region

from a control rat (top), KA-treated rat without seizures (middle) and a KA-treated rat with

seizures (bottom). Somatostatin positive (SST+) neurons are visualized using cell type specific

immunohistochemistry. Magnified images to the right are selected from the area defined by

the black boxes. (Right) Examples of more abundant SST+ interneurons in the hilus (H) of

control rats compared to both KA-treated conditions. **C,** Timm scores indicating degree of

mossy fiber sprouting for all rats from each group. Each circle is the average score from one rat.

KA-treated rats with seizures have the most extensive mossy fiber sprouting as indicated by

significantly higher TIMM scores compared to control rats and KA-treated rats without seizures.

**D,** The number of SST+ interneurons in the dorsal hilus is significantly higher in control rats

compared to KA-treated rats with and without seizures. Each circle is the average across

sections from a single rat. **E**, (Left) Schematic of the dentate-dependent discrimination memory

task on an 8-arm radial maze. A single arm is baited with a reward, and two arms are available

to the animal, the baited arm and one of the two adjacent arms. The maze was in a small room

with distinct distal cues. **F,** The number of trials for rats to preferentially enter the rewarded

arm compared to adjacent non-rewarded arms (criterion: 2 of 3 days with ≥90% rewarded

trials) by groups (control, KA-treated, and dentate lesioned animals). KA-treated rats with and

without seizures are combined, as there were no significant differences between these groups.

Ctrl, control; KA, kainic acid treated; seiz, seizures.

**Figure S1. Range of TIMM scores observed in KA-treated rats.** (Left) Four images from four

different rats, each of which were scored with progressively higher TIMM scores (0-3, top left

corner), a measure of the degree of mossy fiber sprouting (see Methods for details on scoring

procedure). Sections with no or only occasional supragranular mossy fiber staining were scored

0. Sections with scattered mossy fiber staining above all parts of the granular layer were given a

score of 1. Sections that exhibited either patches of heavy mossy fiber staining interspersed

with regions of sparser staining or a continuous band of staining intermediate in intensity were

given a score of 2. Finally, sections with a dense, continuous band of supragranular mossy fiber

staining were assigned a score of 3. (Right) Magnified images from the area defined by white

boxes in the sections to the left. The magnified area spans the granule cell layer and inner

molecular layer (IML). A higher TIMM score reflects a more substantial reorganization of the

mossy fibers, including sprouting into regions where axons and terminals are typically not

found, such as the IML. For example, the bottom image with the highest TIMM score of 3 is

characterized by extensive black fiber tracks within the IML in comparison to control levels

(score of 0). GCL, granule cell layer.

**Figure 2. Spatial firing patterns of dentate neurons were less precise in KA-treated rats. A,**

Cresyl stained coronal section of the dentate gyrus, from a representative control rat with

chronic electrophysiological recordings during behavior. The end of a tetrode positioned within

the granule cell layer (GCL) is indicated by a circle (see Methods for anatomical tracking of

tetrode locations). Neurons are defined as putative granule neurons when the associated

tetrode is confirmed by histology to be in the GCL.  Scale bar = 250 μm. **B,** Cumulative

distribution functions of peak firing rates of recorded dentate neurons (from control and KA-

treated rats). Neurons with peak activity >2 Hz (indicated by black dashed line) were considered

active. **C,** The average rate of active neurons was similar between KA-treated and control rats.

**D,** Activity patterns of individual dentate neurons (10 per group) during 10 min random foraging

in a square environment. For each dentate neuron, the spatial information (blue number on top

of map) and peak firing rate (black number on bottom of map) were calculated. For each group

(control, left; KA-treated, right), the top row comprises rate maps from neurons with the

highest spatial information and the bottom row comprises rate maps from neurons with the

lowest spatial information. **E,** Firing patterns of dentate neurons carried significantly less spatial

information in KA-treated compared to control rats**.** Rate maps of KA-treated dentate neurons

contained significantly higher spatial non-selectivity compared to control rats – revealing that

spikes were more spatially dispersed after KA treatment.  **F,** Number of place fields of a single

dentate cell during one 10-min session in the open field and the average size of each place field

did not differ between groups. However, the within field average firing rate was higher in KA-

treated compared to control rats. Boxplots show median and quartiles, with whiskers extending

to minimum and maximum values. *P* values are reported for rank-sum tests. Ctrl, control; KA,

kainic acid treated.

**Figure S2. The spatial firing properties of dentate neurons from KA-treated rats without**

**seizures did not differ from rats that had developed epileptic seizures.** No statistical

differences were measured in (A) average firing rate, p = 0.4, (B) spatial information, p = 0.77,

or (C) spatial non-selectivity, p = 0.62, when comparing neurons from KA-treated rats with or

without observed seizure events (KA-treated no seizure: n = 2 rats, n = 13 active dentate

neurons; KA-treated seizure: n = 4 rats, n = 17 active dentate neurons). Active neurons are

defined as neurons with a peak rate >2 Hz. Statistics: rank-sum test. KA, kainic acid treated.

**Figure 3. Dentate network pattern separation is impaired in KA-treated rats through**

**reduction in the capacity of the network to perform rate remapping. A,** (Left) Behavioral

paradigm. Animals explored a highly familiar open-field arena in which the shape of the arena

could be changed from square to circular (four 10-min sessions; two of each shape presented in

random order each day). The arena remained in the same place in the room with common

visual cues and a shared floor. The activity of dentate neuron populations was recorded

throughout the entire behavioral paradigm. (Right) Dentate population activity was compared

across the behavioral sessions as indicated. **B,** Rate maps for all four sessions are shown for two

dentate neurons per group (left, control; right KA-treated). Only common areas between the

square and circle shape were compared (corresponding to the color-coded maps, which fill the

size of the square arena, but are inside the stippled line for the circular area). Rate maps are

reordered for visual comparison, regardless of their order during the experiment. All maps are

scaled to the peak rate across the entire recording sequence (indicated below the leftmost rate

map for each cell). Representative dentate neurons show similar activity patterns in repetitions

of the same shape environment, indicating that spatial representations are stable. Firing

patterns across shapes are more different as expected for pattern separation across distinct

environments. **C,** Schematic illustrating how population vector correlations are computed from

dentate population activity. Using separate stacks for recorded neurons from control and KA-

treated rats, population vectors were calculated for each spatial bin (x,y) in the environment

across all neurons (z). Example numeric population vectors (PV) corresponding to the bin

indicated by the pink line are shown below each stack. Bins that include firing fields that change

between environments (rate or location of firing) will have low correlation. **D,** (Left) Cumulative

distribution functions of population vector correlations between pairs of spatial bins

(comparisons indicated in A, right). The dentate population activity was well correlated

between repetitions in same-shaped environments in both control (light colored line) and KA-

treated rats (dark colored line). When comparing population activity between different shaped

environments (dashed lines), the control group exhibited strong decorrelation (light colored

dots), in comparison to the KA-treated group where a more moderate decorrelation was

observed (dark colored dots). (Right) Control neuron populations discriminated more between

different-shape comparisons versus same-shape comparisons than populations from KA-

treated rats (change in PV correlation was calculated by subtracting each different-shape-PV

from the median of the same-shape-PV), indicating stronger pattern separation in control

networks. **E,** (Left) Cumulative distribution functions of each neuron's spatial correlations

between pairs of firing rate maps (comparisons indicated in A).  Same-shape comparisons were

more correlated than different-shape comparisons in both groups (KA-treated, control). P

values are reported for post-hoc comparison for KA-treated or control between same and

different shapes. (Right) Cumulative distribution functions of the rate overlap between pairs of

firing fields (comparisons indicated in A). Only the control group demonstrated strong rate-

remapping across distinct environments.

## REFERENCES

1    O'Reilly, R. C. & McClelland, J. L. Hippocampal conjunctive encoding, storage, and recall:
avoiding a trade-off. *Hippocampus* **4**, 661-682 (1994). https://doi.org:10.1002/hipo.450040605
2    Leutgeb, J. K., Leutgeb, S., Moser, M. B. & Moser, E. I. Pattern separation in the dentate gyrus
and CA3 of the hippocampus. *Science* **315**, 961-966 (2007).
https://doi.org:10.1126/science.1135801
3    Josselyn, S. A., Kohler, S. & Frankland, P. W. Finding the engram. *Nat Rev Neurosci* **16**, 521-534
(2015). https://doi.org:10.1038/nrn4000
4    Neunuebel, J. P. & Knierim, J. J. CA3 retrieves coherent representations from degraded input:
direct evidence for CA3 pattern completion and dentate gyrus pattern separation. *Neuron* **81**,
416-427 (2014). https://doi.org:10.1016/j.neuron.2013.11.017
5    Berron, D. *et al.* Strong Evidence for Pattern Separation in Human Dentate Gyrus. *J Neurosci* **36**,
7569-7579 (2016). https://doi.org:10.1523/JNEUROSCI.0518-16.2016
6    GoodSmith, D., Lee, H., Neunuebel, J. P., Song, H. & Knierim, J. J. Dentate Gyrus Mossy Cells
Share a Role in Pattern Separation with Dentate Granule Cells and Proximal CA3 Pyramidal Cells.
*J Neurosci* **39**, 9570-9584 (2019). https://doi.org:10.1523/JNEUROSCI.0940-19.2019
7    Woods, N. I. *et al.* The Dentate Gyrus Classifies Cortical Representations of Learned Stimuli.
*Neuron* **107**, 173-184 e176 (2020). https://doi.org:10.1016/j.neuron.2020.04.002
8    Bakker, A., Kirwan, C. B., Miller, M. & Stark, C. E. Pattern separation in the human hippocampal
CA3 and dentate gyrus. *Science* **319**, 1640-1642 (2008).
https://doi.org:10.1126/science.1152882
9    Yassa, M. A. & Stark, C. E. Pattern separation in the hippocampus. *Trends Neurosci* **34**, 515-525
(2011). https://doi.org:10.1016/j.tins.2011.06.006
10   Amaral, D. G., Scharfman, H. E. & Lavenex, P. The dentate gyrus: fundamental neuroanatomical
organization (dentate gyrus for dummies). *Prog Brain Res* **163**, 3-22 (2007).
https://doi.org:10.1016/S0079-6123(07)63001-5
11   Kesner, R. P. An analysis of the dentate gyrus function. *Behav Brain Res* **254**, 1-7 (2013).
https://doi.org:10.1016/j.bbr.2013.01.012
12   Marr, D. Simple Memory - Theory for Archicortex. *Philos T Roy Soc B* **262**, 23-& (1971).
https://doi.org:DOI 10.1098/rstb.1971.0078
13   Chawla, M. K. *et al.* Sparse, environmentally selective expression of Arc RNA in the upper blade
of the rodent fascia dentata by brief spatial experience. *Hippocampus* **15**, 579-586 (2005).
https://doi.org:10.1002/hipo.20091

14  Piatti, V. C., Ewell, L. A. & Leutgeb, J. K. Neurogenesis in the dentate gyrus: carrying the message
or dictating the tone. *Front Neurosci* **7**, 50 (2013). https://doi.org:10.3389/fnins.2013.00050
15  Erwin, S. R. *et al.* A Sparse, Spatially Biased Subtype of Mature Granule Cell Dominates
Recruitment in Hippocampal-Associated Behaviors. *Cell Rep* **31**, 107551 (2020).
https://doi.org:10.1016/j.celrep.2020.107551
16  Patton, P. E. & McNaughton, B. Connection matrix of the hippocampal formation: I. The dentate
gyrus. *Hippocampus* **5**, 245-286 (1995). https://doi.org:10.1002/hipo.450050402
17  Ewell, L. A. & Jones, M. V. Frequency-tuned distribution of inhibition in the dentate gyrus. *J
Neurosci* **30**, 12597-12607 (2010). https://doi.org:10.1523/JNEUROSCI.1854-10.2010
18  Braganza, O., Mueller-Komorowska, D., Kelly, T. & Beck, H. Quantitative properties of a feedback
circuit predict frequency-dependent pattern separation. *Elife* **9** (2020).
https://doi.org:10.7554/eLife.53148
19  Rolls, E. T. Pattern separation, completion, and categorisation in the hippocampus and
neocortex. *Neurobiol Learn Mem* **129**, 4-28 (2016). https://doi.org:10.1016/j.nlm.2015.07.008
20  van Dijk, M. T. & Fenton, A. A. On How the Dentate Gyrus Contributes to Memory
Discrimination. *Neuron* **98**, 832-845 e835 (2018). https://doi.org:10.1016/j.neuron.2018.04.018
21  Leutgeb, S. *et al.* Independent codes for spatial and episodic memory in hippocampal neuronal
ensembles. *Science* **309**, 619-623 (2005). https://doi.org:10.1126/science.1114037
22  Colgin, L. L., Moser, E. I. & Moser, M. B. Understanding memory through hippocampal
remapping. *Trends Neurosci* **31**, 469-477 (2008). https://doi.org:10.1016/j.tins.2008.06.008
23  McAvoy, K., Besnard, A. & Sahay, A. Adult hippocampal neurogenesis and pattern separation in
DG: a role for feedback inhibition in modulating sparseness to govern population-based coding.
*Front Syst Neurosci* **9**, 120 (2015). https://doi.org:10.3389/fnsys.2015.00120
24  Yim, M. Y., Hanuschkin, A. & Wolfart, J. Intrinsic rescaling of granule cells restores pattern
separation ability of a dentate gyrus network model during epileptic hyperexcitability.
*Hippocampus* **25**, 297-308 (2015). https://doi.org:10.1002/hipo.22373
25  Kahn, J. B., Port, R. G., Yue, C., Takano, H. & Coulter, D. A. Circuit-based interventions in the
dentate gyrus rescue epilepsy-associated cognitive dysfunction. *Brain* **142**, 2705-2721 (2019).
https://doi.org:10.1093/brain/awz209
26  Madar, A. D., Ewell, L. A. & Jones, M. V. Pattern separation of spiketrains in hippocampal
neurons. *Sci Rep* **9**, 5282 (2019). https://doi.org:10.1038/s41598-019-41503-8
27  Lalani, S. J. *et al.* Impaired Behavioral Pattern Separation in Refractory Temporal Lobe Epilepsy
and Mild Cognitive Impairment. *J Int Neuropsychol Soc*, 1-13 (2021).
https://doi.org:10.1017/S1355617721000734
28  Madar, A. D. *et al.* Deficits in Behavioral and Neuronal Pattern Separation in Temporal Lobe
Epilepsy. *J Neurosci* **41**, 9669-9686 (2021). https://doi.org:10.1523/JNEUROSCI.2439-20.2021
29  Buckmaster, P. S. & Dudek, F. E. Network properties of the dentate gyrus in epileptic rats with
hilar neuron loss and granule cell axon reorganization. *J Neurophysiol* **77**, 2685-2696 (1997).
https://doi.org:10.1152/jn.1997.77.5.2685
30  Buckmaster, P. S. & Dudek, F. E. Neuron loss, granule cell axon reorganization, and functional
changes in the dentate gyrus of epileptic kainate-treated rats. *Journal of Comparative Neurology*
**385**, 385-404 (1997).
31  Buckmaster, P. S. & Dudek, F. E. In vivo intracellular analysis of granule cell axon reorganization
in epileptic rats. *J Neurophysiol* **81**, 712-721 (1999). https://doi.org:10.1152/jn.1999.81.2.712
32  Bragin, A., Benassi, S. K., Kheiri, F. & Engel, J., Jr. Further evidence that pathologic high-
frequency oscillations are bursts of population spikes derived from recordings of identified cells
in dentate gyrus. *Epilepsia* **52**, 45-52 (2011). https://doi.org:10.1111/j.1528-1167.2010.02896.x

33    Houser, C. R. & Esclapez, M. Vulnerability and plasticity of the GABA system in the pilocarpine
831             model of spontaneous recurrent seizures. *Epilepsy Res* **26**, 207-218 (1996). https://doi.org:Doi
832             10.1016/S0920-1211(96)00054-X

833    34    Zhang, W. *et al.* Surviving hilar somatostatin interneurons enlarge, sprout axons, and form new
834             synapses with granule cells in a mouse model of temporal lobe epilepsy. *J Neurosci* **29**, 14247-
835             14256 (2009). https://doi.org:10.1523/JNEUROSCI.3842-09.2009

836    35    Kobayashi, M. & Buckmaster, P. S. Reduced inhibition of dentate granule cells in a model of
837             temporal lobe epilepsy. *J Neurosci* **23**, 2440-2452 (2003).

838    36    Lynch, M., Sayin, U., Golarai, G. & Sutula, T. NMDA receptor-dependent plasticity of granule cell
839             spiking in the dentate gyrus of normal and epileptic rats. *J Neurophysiol* **84**, 2868-2879 (2000).
840             https://doi.org:10.1152/jn.2000.84.6.2868

841    37    Kelly, T. & Beck, H. Functional properties of granule cells with hilar basal dendrites in the
842             epileptic dentate gyrus. *Epilepsia* **58**, 160-171 (2017). https://doi.org:10.1111/epi.13605

843    38    Dengler, C. G., Yue, C., Takano, H. & Coulter, D. A. Massively augmented hippocampal dentate
844             granule cell activation accompanies epilepsy development. *Sci Rep* **7**, 42090 (2017).
845             https://doi.org:10.1038/srep42090

846    39    Krook-Magnuson, E. *et al.* In vivo evaluation of the dentate gate theory in epilepsy. *J Physiol* **593**,
847             2379-2388 (2015). https://doi.org:10.1113/JP270056

848    40    Gilbert, P. E., Kesner, R. P. & Lee, I. Dissociating hippocampal subregions: double dissociation
849             between dentate gyrus and CA1. *Hippocampus* **11**, 626-636 (2001).
850             https://doi.org:10.1002/hipo.1077

851    41    Lee, C. H. & Lee, I. Impairment of Pattern Separation of Ambiguous Scenes by Single Units in the
852             CA3 in the Absence of the Dentate Gyrus. *J Neurosci* **40**, 3576-3590 (2020).
853             https://doi.org:10.1523/JNEUROSCI.2596-19.2020

854    42    Morris, A. M., Churchwell, J. C., Kesner, R. P. & Gilbert, P. E. Selective lesions of the dentate
855             gyrus produce disruptions in place learning for adjacent spatial locations. *Neurobiol Learn Mem*
856             **97**, 326-331 (2012). https://doi.org:10.1016/j.nlm.2012.02.005

857    43    McHugh, T. J. *et al.* Dentate gyrus NMDA receptors mediate rapid pattern separation in the
858             hippocampal network. *Science* **317**, 94-99 (2007). https://doi.org:10.1126/science.1140263

859    44    Leutgeb, J. K. *et al.* Progressive transformation of hippocampal neuronal representations in
860             "morphed" environments. *Neuron* **48**, 345-358 (2005).
861             https://doi.org:10.1016/j.neuron.2005.09.007

862    45    Shuman, T. *et al.* Breakdown of spatial coding and interneuron synchronization in epileptic mice.
863             *Nat Neurosci* **23**, 229-238 (2020). https://doi.org:10.1038/s41593-019-0559-0

864    46    Dinocourt, C., Petanjek, Z., Freund, T. F., Ben-Ari, Y. & Esclapez, M. Loss of interneurons
865             innervating pyramidal cell dendrites and axon initial segments in the CA1 region of the
866             hippocampus following pilocarpine-induced seizures. *J Comp Neurol* **459**, 407-425 (2003).
867             https://doi.org:10.1002/cne.10622

868    47    Bouilleret, V., Loup, F., Kiener, T., Marescaux, C. & Fritschy, J. M. Early loss of interneurons and
869             delayed subunit-specific changes in GABA(A)-receptor expression in a mouse model of mesial
870             temporal lobe epilepsy. *Hippocampus* **10**, 305-324 (2000). https://doi.org:10.1002/1098-
871             1063(2000)10:3<305::AID-HIPO11>3.0.CO;2-I

872    48    Hofmann, G., Balgooyen, L., Mattis, J., Deisseroth, K. & Buckmaster, P. S. Hilar somatostatin
873             interneuron loss reduces dentate gyrus inhibition in a mouse model of temporal lobe epilepsy.
874             *Epilepsia* **57**, 977-983 (2016). https://doi.org:10.1111/epi.13376

875    49    McKenzie, S. *et al.* Preexisting hippocampal network dynamics constrain optogenetically induced
876             place fields. *Neuron* **109**, 1040-1054 e1047 (2021).
877             https://doi.org:10.1016/j.neuron.2021.01.011

50 Rolotti, S. V. *et al.* Local feedback inhibition tightly controls rapid formation of hippocampal
place fields. *Neuron* **110**, 783-794 e786 (2022). https://doi.org:10.1016/j.neuron.2021.12.003
51 Royer, S. *et al.* Control of timing, rate and bursts of hippocampal place cells by dendritic and
somatic inhibition. *Nat Neurosci* **15**, 769-775 (2012). https://doi.org:10.1038/nn.3077
52 Chavlis, S., Petrantonakis, P. C. & Poirazi, P. Dendrites of dentate gyrus granule cells contribute
to pattern separation by controlling sparsity. *Hippocampus* **27**, 89-110 (2017).
https://doi.org:10.1002/hipo.22675
53 Galloni, A. R., Samadzelkava, A., Hiremath, K., Oumnov, R. & Milstein, A. D. Recurrent Excitatory
Feedback From Mossy Cells Enhances Sparsity and Pattern Separation in the Dentate Gyrus via
Indirect Feedback Inhibition. *Front Comput Neurosci* **16**, 826278 (2022).
https://doi.org:10.3389/fncom.2022.826278
54 Freund, T. F. & Buzsaki, G. Interneurons of the hippocampus. *Hippocampus* **6**, 347-470 (1996).
https://doi.org:10.1002/(SICI)1098-1063(1996)6:4<347::AID-HIPO1>3.0.CO;2-I
55 Witter, M. P. Organization of the entorhinal-hippocampal system: a review of current
anatomical data. *Hippocampus* **3 Spec No**, 33-44 (1993).
56 Hargreaves, E. L., Rao, G., Lee, I. & Knierim, J. J. Major dissociation between medial and lateral
entorhinal input to dorsal hippocampus. *Science* **308**, 1792-1794 (2005).
https://doi.org:10.1126/science.1110449
57 Tsao, A. *et al.* Integrating time from experience in the lateral entorhinal cortex. *Nature* **561**, 57-
62 (2018). https://doi.org:10.1038/s41586-018-0459-6
58 Lu, L. *et al.* Impaired hippocampal rate coding after lesions of the lateral entorhinal cortex. *Nat*
*Neurosci* **16**, 1085-1093 (2013). https://doi.org:10.1038/nn.3462
59 Santhakumar, V., Aradi, I. & Soltesz, I. Role of mossy fiber sprouting and mossy cell loss in
hyperexcitability: a network model of the dentate gyrus incorporating cell types and axonal
topography. *J Neurophysiol* **93**, 437-453 (2005). https://doi.org:10.1152/jn.00777.2004
60 Liu, X. *et al.* Seizure-induced changes in place cell physiology: relationship to spatial memory. *J*
*Neurosci* **23**, 11505-11515 (2003). https://doi.org:10.1523/JNEUROSCI.23-37-11505.2003
61 Ewell, L. A., Fischer, K. B., Leibold, C., Leutgeb, S. & Leutgeb, J. K. The impact of pathological
high-frequency oscillations on hippocampal network activity in rats with chronic epilepsy. *Elife* **8**
(2019). https://doi.org:10.7554/eLife.42148
62 Hunt, R. F., Girskis, K. M., Rubenstein, J. L., Alvarez-Buylla, A. & Baraban, S. C. GABA progenitors
grafted into the adult epileptic brain control seizures and abnormal behavior. *Nat Neurosci* **16**,
692-697 (2013). https://doi.org:10.1038/nn.3392
63 Zhu, B., Eom, J. & Hunt, R. F. Transplanted interneurons improve memory precision after
traumatic brain injury. *Nat Commun* **10**, 5156 (2019). https://doi.org:10.1038/s41467-019-
13170-w
64 Ewell, L. A. *et al.* Brain State Is a Major Factor in Preseizure Hippocampal Network Activity and
Influences Success of Seizure Intervention. *J Neurosci* **35**, 15635-15648 (2015).
https://doi.org:10.1523/JNEUROSCI.5112-14.2015
65 Hellier, J. L., Patrylo, P. R., Buckmaster, P. S. & Dudek, F. E. Recurrent spontaneous motor
seizures after repeated low-dose systemic treatment with kainate: assessment of a rat model of
temporal lobe epilepsy. *Epilepsy Res* **31**, 73-84 (1998). https://doi.org:Doi 10.1016/S0920-
1211(98)00017-5
66 Racine, R. J. Modification of seizure activity by electrical stimulation. II. Motor seizure.
*Electroencephalogr Clin Neurophysiol* **32**, 281-294 (1972). https://doi.org:10.1016/0013-
4694(72)90177-0
67 Snyder, J. S. *et al.* A Transgenic Rat for Specifically Inhibiting Adult Neurogenesis. *eNeuro* **3**
(2016). https://doi.org:10.1523/ENEURO.0064-16.2016

68    McNaughton, B. L. Neuronal 'Ensemble' Recording and the Search for the Cell Assembly: A
927           Personal History. *Hippocampus* **35**, e23669 (2025). https://doi.org:10.1002/hipo.23669
928    69    Sasaki, T. *et al.* Dentate network activity is necessary for spatial working memory by supporting
929           CA3 sharp-wave ripple generation and prospective firing of CA3 neurons. *Nat Neurosci* **21**, 258-
930           269 (2018). https://doi.org:10.1038/s41593-017-0061-5
931    70    Walsh, T. J., Schulz, D. W., Tilson, H. A. & Schmechel, D. E. Colchicine-induced granule cell loss in
932           rat hippocampus: selective behavioral and histological alterations. *Brain Res* **398**, 23-36 (1986).
933           https://doi.org:10.1016/0006-8993(86)91246-1
934    71    Goldschmidt, R. B. & Steward, O. Preferential neurotoxicity of colchicine for granule cells of the
935           dentate gyrus of the adult rat. *Proc Natl Acad Sci U S A* **77**, 3047-3051 (1980).
936           https://doi.org:10.1073/pnas.77.5.3047
937    72    Mankin, E. A. *et al.* Neuronal code for extended time in the hippocampus. *Proc Natl Acad Sci U S*
938           *A* **109**, 19462-19467 (2012). https://doi.org:10.1073/pnas.1214107109
939    73    Skaggs WE, M. B., Gothard KM, Markus EJ. . *An information-theoretic approach to deciphering*
940           *the hippocampal code*. (1993).
941    74    Mankin, E. A., Diehl, G. W., Sparks, F. T., Leutgeb, S. & Leutgeb, J. K. Hippocampal CA2 activity
942           patterns change over time to a larger extent than between spatial contexts. *Neuron* **85**, 190-201
943           (2015). https://doi.org:10.1016/j.neuron.2014.12.001
944    75    Tauck, D. L. & Nadler, J. V. Evidence of functional mossy fiber sprouting in hippocampal
945           formation of kainic acid-treated rats. *J Neurosci* **5**, 1016-1022 (1985).

# Figure 1

**A**
TIMM, 1    CA1
DG    CA3
GCL

TIMM, 2    CA1
DG    CA3
GCL

**B**
Control
DG    H    CA3
GCL

KA-treated
no seizure
DG    H    CA3
GCL

KA-treated
seizure
DG    H    CA3
GCL

**C**

P = 0.0006
P = 3.9 x 10⁻⁷

Timm Score

3
2
1
0

Ctrl    no
seiz    KA
seiz

**D**

P = 0.005
P = 0.0009

# SST Positive Neurons

Ctrl    no
seiz    KA
seiz

**E**
Dentate-Dependent
Memory Task

or

| walls
★ reward

**F**

P = 0.03
P = 0.009

# Trials to Criterion

Ctrl    KA    DG Lesion

**Figure S1**

**TIMM Score**

**Figure 2**

**A**

DG

**B**

Prop. of Neurons

Peak Firing Rate Hz

2 Hz cutoff

Ctrl
KA

**C**

Ave. Rate (Hz)

P = 0.13

Ctrl KA

**D**

Control

2.2    2.1    1.9    1.9    1.8

3 Hz    3 Hz    2 Hz    3 Hz    8 Hz

1.3    1.3    1.2    1.1    0.5

2 Hz    5 Hz    10 Hz    4 Hz    4 Hz

KA-treated

2.7    2.1    1.7    1.3    1.2

2 Hz    3 Hz    3 Hz    8 Hz    4 Hz

0.4    0.2    0.2    0.2    0.1

6 Hz    5 Hz    3 Hz    6 Hz    3 Hz

Peak Rate (Hz)    max    0

**E**

Spatial Info. (bits/spike)

P = 0.0056

Ctrl KA

Spatial Non-selectivity

P= 0.017

Ctrl KA

**F**

Number of Fields

P= 0.08

Ctrl KA

Field Size (cm²)

P = 0.15

Ctrl KA

Ave. Field Rate (Hz)

P = 0.014

Ctrl KA

**Figure S2**

# Figure 3