A comprehensive AMR genotype-phenotype database (CABBAGE)

Emily Dickens[*1], Romain Derelle[*1], Robert Beardmore[2], Anita Suresh[4], Swapna Uplekar[5], Andrew D Yates[3], Jon Keatley[3], Andrea Winterbottom[3], Andrey G Azov[3], Bilal El Houdaigui[3], Sofiia Ochkalova[3], Tatiana A Gurbich[3], Anu Shivalikanjli[3], Galabina Yordanova[3], John A Lees[3], and Leonid Chindelevitch[1]

[1]*MRC Center for Global Infectious Disease Analysis, School of Public Health, Imperial College London, London, United Kingdom*

[2]*Biosciences, College of Life and Environmental Sciences, University of Exeter, Exeter, United Kingdom*

[3]*European Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Genome Campus, Hinxton, United Kingdom*

[4]*University Hospital Heidelberg, Heidelberg, Germany*

[5]*Foundation for Innovative New Diagnostics (FINDDx), Geneva, Switzerland*

[*]These authors contributed equally to this work.

Corresponding author : l.chindelevitch@imperial.ac.uk (Leonid Chindelevitch)

ABSTRACT

Addressing the growing threat of antimicrobial resistance (AMR) requires the development of large-scale resources that link bacterial genomic data with phenotypic antimicrobial resistance profiles. Such datasets are essential for advancing genotype-based predictions of resistance to uncover novel resistance mechanisms, as well as identifying and tracking global trends. Here, we describe the development of the 'Comprehensive Assessment of Bacterial-Based AMR prediction from GEnotypes' (CABBAGE) database, linking bacterial genomes to associated antibiotic susceptibility data and relevant metadata across WHO Bacterial Priority Pathogens, sourced from both publications and existing databases, and curated into a format that is compatible with and extends both NCBI and ENA's formats. The resulting CABBAGE database, comprising over 170,000 unique sequenced isolates and approximately 1.7 million genome-phenotype pairs linked to extensive metadata, represents the largest of its kind, consolidating existing AMR phenotype-genotype data into a single unified format. The database encompasses a broad range of antimicrobials, facilitating the analysis of global resistance trends as well as benchmarks of genotype-to-phenotype predictive methods, and empowering further research uses. The database is freely accessible at https://www.ebi.ac.uk/amr and is currently being integrated with the BioSample database for easy access.

1 INTRODUCTION

Antimicrobial resistance poses a substantial risk to global health through the reduction of antibiotic efficacy, resulting in increased morbidity, mortality, and healthcare associated costs. Addressing this challenge requires a multifaceted approach, ranging from clinical interventions such as enhanced antibiotic stewardship and diagnostics, to broader initiatives, including the development of novel antimicrobials. Among these, advances in whole-genome sequencing (WGS) have raised the possibility of using genomics-based diagnostics alongside, or as a replacement for, traditional antibiotic susceptibility tests (ASTs). However, while numerous antimicrobial resistance genes have been identified, discordance between genotypic predictions and phenotypic resistance is observed across numerous studies [1, 2, 3]. A key step in addressing this discordance is the development of large-scale, curated datasets that integrate AMR phenotypes with corresponding genomic data, with the aim of characterising resistance mechanisms, including those not yet genetically defined.

The World Health Organisation (WHO) has identified a list of bacterial pathogens which are a priority for new antibiotic development, encompassing 26 bacterial pathogen-antibiotic combinations in 2017 [4], revised to 24 in 2024 [5]. However, while the development of novel antimicrobials remains critical in addressing the growing threat of AMR, this strategy is often viewed as unsustainable due to the substantial costs associated with drug development and the risk of resistance developing soon after new antimicrobials are introduced into clinical use. An alternative approach is to maximise the effectiveness of existing antimicrobials with appropriate diagnostics and antimicrobial stewardship efforts [6, 7]. This could involve making treatment decisions informed by aggregated phenotypic and genotypic AMR data, however, clinically-derived AST results are not routinely shared with the scientific community, and genomes are not sequenced as part of standard practice for the majority of pathogens. Nevertheless, a wealth of genotypic and phenotypic information relevant to AMR already exists in various formats within public repositories.

A number of AMR databases have been developed to support research and surveillance efforts. Key resources include the Pathosystems Resource Integration Center (PATRIC) [8], the NCBI BioSample Antibiograms database [9] and the National Database of Antibiotic Resistant Organisms (NDARO) [10], all of which provide extensive bacterial genomic data linked to antimicrobial susceptibility data. Similarly, PubMLST [11], although primarily focusing on the provision of MLST information, also integrates bacterial population genomic data with phenotypic information, while platforms such as Microreact [12] and PathogenWatch [13] enable the visualisation of global AMR data alongside accompanying genotypic data when available. National level surveillance is also supported by agencies in the U.S. such as the

Centres for Disease Control and Prevention (CDC) [14], which reports AMR trends, and the National Antimicrobial Resistance Monitoring System (NARMS) [15], which focuses on foodborne bacteria. Additionally, the European COMPARE Consortium [16] has created an AMR data hub consisting of over 2500 sequenced isolates linked to structured phenotypic data. However, despite the availability of these resources, current AMR data remains largely fragmented across various institutions and regions, limiting its collective influence. There is therefore a need to develop a single, harmonised AMR database that integrates existing data, standardises data formats, and incorporates expert curation. Publications often also contain genotype-phenotype data and metadata that has not been included in any of the aforementioned resources, and can therefore provide an additional valuable source of information on AMR phenotypes and matching genotypes. A key challenge is that they can rarely be processed in a fully automated way.

The work presented in this study aimed to combine all available isolates of WHO Priority Pathogens with both genomic and antibiotic susceptibility testing data, drawing on two sources: an analysis of existing databases containing such datasets, as well as additional large AMR genotype-phenotype datasets extracted from the literature, followed by a process of data standardisation and curation. The end result is the 'Comprehensive Assessment of Bacterial-Based AMR prediction from GEnotypes' (CABBAGE) database, comprising over 170,000 unique sequenced isolates and approximately 1.7 million genome-phenotype pairs linked to extensive metadata. Here, we present an overview of CABBAGE, including the availability of data among different antibiotics and pathogens, the spatio-temporal data distribution, as well as a comparative analysis with ATLAS [17], a global AMR dataset encompassing over 900k isolates with associated antimicrobial susceptibility data, but no genotypic data aside from the presence/absence of a limited number of β-lactamase genes.

2   METHODS

2.1   Data collection

A literature review was carried out, with the aim of identifying and extracting data from publications containing at least 100 sequenced isolates on the WHO Bacterial Priority Pathogens List, with antimicrobial resistance phenotypes available for at least one drug. Note that this 100 isolate threshold was adjusted for some pathogens to account for their over- or under-representation in the literature, as detailed in the Supplementary Methods (S1.1.1). For each dataset with publicly available data, tables containing phenotypes as well as the accession numbers for the genotypes were extracted and standardised into a CSV format. These files were then processed via custom scripts, available at https://github.com/Leonardini/CABBAGE, to identify all relevant information on the genotype, pheno-

type, and metadata.

In addition to the datasets extracted from the literature, we identified nine public databases (Microreact, NARMS, NDARO, Pathogenwatch, PATRIC, PubMLST, COMPARE, CDC and NCBI antibiograms) as containing relevant information and therefore extracted their data. The number of isolates contributed by each database are listed in Supplementary Table S3 and summarised in Supplementary Table S2.

As the genomic component of CABBAGE entries is structured around BioSample IDs, all sequencing run IDs, sample IDs, and genome accession IDs were converted to BioSample IDs. Missing values in these genomic fields were then retrieved using BioSample IDs as the central reference. In addition, to improve the completeness of metadata fields, isolate metadata were retrieved for entries with missing values, again using BioSample IDs as the reference.

2.2 Data standardisation

To ensure comparability across datasets obtained from different sources, the data was mapped to a set of 29 standardised fields (Table 1), with each entry being defined by a pair of Biosample ID (genotype) and antibiotic name (phenotype). A key step in the data standardisation process involved the harmonisation of antibiotic names and their extraction from abbreviations as, for instance, a single abbreviation may refer to numerous different antibiotics across different datasets. To address this, antibiotic names were derived from the antibiotic abbreviations provided in the source dataset, referring to the paper's full text when necessary, and standardised using the name list employed by NCBI antibiograms.

Further data standardisation steps include, but are not limited to, the conversion of numerical phenotypes (e.g. 0, 1) to their corresponding classification (e.g. susceptible, resistant), the conversion of country names to ISO 3166-1 alpha-3 codes, standardising both broth microdilution and macrodilution to broth dilution, and standardising the host to either 'Homo sapiens' or 'other (non-clinical isolate)'. A full description of the data standardisation process is included in the Supplementary methods (S1.1.13).

2.3 Data curation

A series of curation steps were performed to remove or correct inconsistent entries. Only those entries that were inconsistent or conflicted have been manually curated. Pathogens not recorded in the WHO Bacterial Pathogen Priority List were excluded, and several species were renamed for consistency (e.g. Salmonella typhimurium was changed to Salmonella enterica). Particular attention was paid to the cu-

ration of genomic identifiers (the genomic component of CABBAGE), as well as measurement values and typing methods (the phenotypic component of CABBAGE).

2.3.1 Genomic data curation

During data collection and the subsequent conversion of genomic identifiers to BioSample IDs, we identified cases where multiple genomic identifiers within a single entry did not point to the same BioSample ID (e.g., a sequencing run ID and a genome accession ID corresponding to different BioSample IDs). As the correct BioSample ID could not be determined, these entries were excluded. We also identified multiple BioSample IDs pointing to the same sequencing run ID. The corresponding entries were also excluded as these cases likely correspond to pooled isolate sequencing.

2.3.2 Phenotypic data curation

Minimum inhibitory concentration (MIC) values were examined for potential errors that were subsequently removed, for example, large values inconsistent with standard doubling dilution ranges. Typing methods associated with anomalous diffusion zone diameter values (e.g., small decimal values) or other irregular MIC values (e.g., series of integers outside standard dilution ranges) were manually checked against the original data sources. When errors were confirmed, both the method and the measurement unit were corrected. Original data sources were also checked to retrieve corresponding typing methods in cases where a measurement value was present without a unit and method. Further curation steps addressed anomalies in metadata fields, such as high values of host age (e.g 999) and improbable collection years (e.g. 1800) that were removed.

The final curation stage involved the de-duplication of entries, including entries that were either identical or shared the same pair Biosample ID/antibiotic name but differed in measurement value or phenotype. In the case of identical entries, duplicates were removed, and conflicting entries that could not be resolved by checking the original sources were also excluded. A full description of the data curation process is included in the Supplementary methods (S1.1.14).

2.4 Final processing

In cases where both a measurement value and categorical phenotype (e.g. R, I or S) are recorded, the phenotype is often derived using outdated or custom breakpoints. Additionally, many entries include a measurement value, but lack a corresponding phenotype. To address this, we incorporated updated phenotypes by applying the current 2025 breakpoint criteria from both CLSI and EUCAST, inferred using the AMR package v3.0.0 [18].

The CABBAGE database then underwent a series of consistency checks to ensure that values within each entry were consistent across fields. For instance, a testing standard can only be reported in the presence of a phenotype, a platform value should have a corresponding typing method, a measurement unit is required if a measurement value is provided alongside a typing method, and each entry must contain both a Biosample ID as well as either a measurement value or a categorical resistance phenotype.

2.5 Database accessibility

CABBAGE is available via the Antimicrobial resistance portal at EMBL-EBI (https://www.ebi.ac.uk/amr). The database will be updated on a regular basis to incorporate new datasets.

3 RESULTS

3.1 Antimicrobial resistance portal

The Antimicrobial resistance portal provides a user-friendly interface to explore CABBAGE and consists of three main data resources: 'Experimental AMR', consisting of antibiogram AMR records; '*in silico* AMR prediction', consisting of *in silico* predictions of AMR from AMRFinderPlus [19]; and finally 'Experimental and *in silico* AMR', a merged dataset of experimental evidence and *in silico* predictions of AMR. Within each resource, data can be filtered based on species, genus, resistance phenotype, isolation source category, testing method, collection year, geographical subregion, and country (Figure 1).

Figure 1: Screenshot of the Antimicrobial resistance portal, showing the CABBAGE database filtered by antibiotic, species, isolation source category and geographical region.

3.2 Overview of the CABBAGE database

In total, CABBAGE contains antibiotic susceptibility data and associated metadata for 170,750 sequenced isolates, with an average of 10 ±6 antibiotic susceptibility test results per isolate. The database contains >1.7 million entries, each entry being defined by a Biosample-antibiotic combination with an associated AST result. The AST result can be represented by a measurement value and/or a categorical phenotype (e.g. S, I or R). Both a measurement value and a categorical phenotype are recorded for >964k entries, and both a measurement value and an updated phenotype, which we derived by comparing the measurement value to the latest breakpoints, are recorded for >1.1 million entries (Table S4). Table 1 summarises field completeness alongside two examples of CABBAGE entries. While the fields defining entries are 100% complete, some metadata fields, such as host sex and age, are sparsely populated but are retained in the anticipation that coverage may improve as the database expands.

The number of entries obtained from, and shared between, different data sources is summarised in Figure 2a. PATRIC is the largest single data source, contributing just over 764k entries, followed by datasets derived from our literature review (89 publications), with over 560k entries. We observed high redundancy between data sources, which allowed us to identify and remove problematic entries during the database deduplication process. PATRIC and the data sourced from publications have the largest number of entries in common with just each other (>142k), followed by NCBI antibiograms and NDARO (>126k).

Finally, the data is not evenly distributed across pathogens, with three species representing more than half of isolates: *Mycobacterium tuberculosis* (24.4%), *Salmonella enterica* (19.6%) and *Escherichia coli* (13.9%) (Figure 2b). In contrast, other pathogens are under-represented, with some species being represented by a single unique isolate: *Enterobacter cancerogenus*, *Serratia fonticola*, *Enterobacter mori*, *Enterobacter soli*, *Serratia nematodiphila*, *Providencia alcalifaciens*, *Proteus alimentorum* and *Enterobacter sichuanensis*.

Data type	Field	Example 1	Example 2
Genome	Biosample ID	SAMN06437146	SAMEA2335670
	INSDC secondary accession	SRS2079188	ERS393864
	Assembly ID	GCA_022066305.1	GCA_901213915.1
Phenotype	Measurement sign	>	==
	Measurement	2/38	6
	Measurement unit	mg/L	mm
	Resistance phenotype	resistant	resistant
	Updated phenotype CLSI	resistant	resistant
	Updated phenotype EUCAST	intermediate	resistant
	Used ECOFF	no	no
Metadata	Isolate	KPN123ec	12291_1#11
	Genus	*Klebsiella*	*Streptococcus*
	Organism	*Klebsiella pneumoniae*	*Streptococcus pneumoniae*
	Antibiotic abbreviation	SXT	ERY
	Antibiotic name	Trimethoprim-sulfamethoxazole	Erythromycin
	Laboratory typing method	broth dilution	disk diffusion
	Platform	BD Phoenix	
	AST standard	CLSI	CLSI
	Host	Homo sapiens	Homo sapiens
	Isolation source	wound	csf
	Isolation source category	skin or soft tissue	central nervous system
	Host sex		Female
	Host age		52
	ISO country code	USA	PER
	Isolation longitude	-95.400586	-75.0152
	Isolation latitude	29.710689	-9.18997
	Collection year	2011	2011
	AMR associated publications	29323230;33075053	28756964;31003929;31184299;32375991
	Database	PATRIC;CABBAGE_PubMed_data	PATRIC;microreact;pubMLST

% Completeness

Table 1: CABBAGE database fields, grouped into genomic, phenotypic and metadata categories. Two examples of CABBAGE entries are provided, as well as the percentage of completeness of each field on the right side.

Figure 2: A) Sources of CABBAGE data, in numbers of entries. Only sets with a minimum intersection size of 2,000 are displayed. The full UpSet plot is provided in Supplementary Figure S1. B) Distribution of the 10 most common pathogens in CABBAGE. All remaining species are grouped under 'Other'. Percentages represent the number of unique isolates (Biosample ID's).

3.3 Spatio-temporal distribution of CABBAGE data

As expected, we observed substantial geographical disparities in the AMR data, with some regions, such as Central Africa and Central America, being sparsely sampled (Figure 3a). The US has the largest data coverage (>698k entries), in part due to their formidable AMR surveillance programs and associated databases (NARMS, NDARO and CDC). Variations are also seen in the frequency of resistance phenotypes between different countries, as illustrated in Figure S4, however caution should be taken when comparing resistance frequencies between countries due to varying levels of data coverage and ascertainment bias due to the fact that not all isolates are sequenced, and in some settings a priority is placed on sequencing those isolates that are phenotypically resistant to at least one antibiotic [20].

New AMR datasets are being published at an increasingly rapid pace, as illustrated in Figure 3b. There is a marked increase in the number of entries from approximately the year 2000 onwards, likely reflecting advances in WGS, AMR surveillance and data sharing. However, we also observed a sharp decline in the number of CABBAGE entries corresponding to recent years. This decline may be purely technical, reflecting a lag in the incorporation of recent data into public databases. Alternatively, it could indicate a genuine reduction in global AMR data collection, resulting from the substantial impact of the COVID-19 pandemic on AMR surveillance programs and academic research [21, 22, 23]. Future updates of the CABBAGE database, incorporating more recent data sources, will help disentangle the relative contributions of these two, not mutually exclusive, hypotheses.

Figure 3: A) Geographical distribution of entries ($\log_{10}$). Countries that are not represented in CABBAGE are shown in white. B) Distribution of CABBAGE entries per collection years. The category 'other' includes susceptible-dose dependent and non-susceptible. Frequencies of each categorical phenotype per year are shown in Supplementary Figure S5b.

3.4 Antibiotic resistance frequency in CABBAGE

The percentage of resistant isolates varies widely between drugs (Figure 4a). Among antibiotics with the greatest data coverage (at least 10,000 isolates), ampicillin showed the highest resistance frequency (40%), whereas bedaquiline had the lowest (1.2%). These differences likely reflect both the time since each drugs introduction and their overall levels of clinical use. However, the CABBAGE dataset as a whole is expected to be biased toward resistant phenotypes (22% of CABBAGE entries with a categorical phenotype), since such isolates are more often sequenced and reported.

To evaluate this potential bias, we compared the resistance frequencies of pathogen-antibiotic pairs shared between CABBAGE and the ATLAS database [17] (103 pairs; Figure 4b). ATLAS serves as a useful comparator because it compiles results from standardized global surveillance studies designed to capture a representative range of susceptibility profiles, rather than being enriched for resistant isolates. Logistic regression analyses revealed that resistance was significantly more frequent in CABBAGE than in ATLAS (OR = 1.6, p < 0.001). Nonetheless, this trend was not universal: 66 pathogen-antibiotic pairs showed significantly higher resistance in CABBAGE, while 29 showed greater susceptibility (Fishers exact test, FDR adjusted p < 0.05).

Figure 4: A) Frequencies of resistant (red bars) versus susceptible (green bars) phenotypes across the CABBAGE dataset, by antibiotic. We used here the phenotypes extracted from the original data source, not the updated phenotypes, and the data was restricted to antibiotics with at least 10,000 entries in CABBAGE. Absolute numbers of susceptible and resistant entries are displayed in Supplementary figure S7. B) Frequencies of resistance in CABBAGE versus ATLAS. Each data point represents a unique pathogen-antibiotic pair present in both datasets, restricted to data collected between 2004-2023 to reflect the range of collection dates in ATLAS. A scatter plot without this date restriction is presented in Supplementary Figure S8. Pathogen-antibiotic pairs with at least 100 entries in both CABBAGE and ATLAS are shown. Error bars represent 95% CI and colours are representative of different antibiotic classes. The CABBAGE phenotypes presented here represent the phenotypes derived from the original datasets, rather than the updated phenotypes.

4   DISCUSSION

Comprehensive, large-scale datasets integrating bacterial genome sequences with associated phenotypic antibiotic susceptibility data are essential for the development of genotype-to-phenotype predictive models to further our understanding of AMR mechanisms. Although resources such as PATRIC provide such information, a substantial volume of relevant data outside of PATRIC remains available in the public domain across multiple databases and publications, with more being added on a weekly basis. To address this, we set out to collect, combine, and curate all publicly available genotype-phenotype data in WHO Bacterial Priority Pathogens, with the aim of creating CABBAGE, a unique, comprehensive resource, representing the largest database of its kind.

Through an extensive literature review and examination of public databases, we extracted data on >170,000 unique sequenced isolates with associated phenotypic AMR data, totalling >1.7 million genome-phenotype pairs. In addition, each entry is associated with metadata where available, such as the isolate collection date, isolation source and geographic location. This constitutes a wealth of information that can be used to better understand the genetic mechanisms of AMR, detect trends for AMR surveillance, and develop more accurate prediction methods for AMR diagnostics.

The data within CABBAGE exhibits multiple layers of heterogeneity. First, the genotypes are available in various formats, from raw reads to contigs and scaffolds, and finishing with assemblies. Since the pre-processing pipeline may influence the downstream inference [24, 25], we recommend that genotype data be made available in its rawest format, with the understanding that any prediction methods would likely need to first pre-process this data into genomic features. Second, the phenotypes vary from quantitative MICs and disk diffusion data obtained by a variety of AST methods, to binary S/I/R classifications obtained using outdated guidelines. We sought to address such limitations in the database, by providing updated phenotypes based on current EUCAST and CLSI breakpoints.

In addition, the species in CABBAGE are unevenly distributed, with *M. tuberculosis* alone accounting for just over 24% of all isolates, and at the other extreme, a number of species are only represented by a single isolate. Similar heterogeneities, roughly following Zipf's law, are found for the antibiotic classes, the publications, the countries represented, and the years of collection. It is possible that CABBAGE could be used to systematically identify gaps in surveillance, even though, depending on the sampling methodology and the resources available, such gaps may or may not be easy to fill. Lastly, we note that the fraction of resistant isolates is likely to be over-estimated, as many groups tend to perform AST on all

isolates of interest, but only sequence those that are resistant. We illustrate this bias towards resistance through a comparative analysis of CABBAGE against the AMR dataset ATLAS, showing that overall there is a greater likelihood of resistance in CABBAGE relative to ATLAS.

In summary, CABBAGE consolidates data from existing resources into a single, unified database. Through the integration, standardisation and curation of information that was previously fragmented, CABBAGE provides a strong foundation to advance research and innovation in the AMR space. Additionally, it highlights the need to adopt a standard format for sharing genotype-phenotype data [26], which will facilitate, and aid the automation of, its extension as additional data is generated.

5   DECLARATION OF INTERESTS

The authors declare no competing interests.

6   ACKNOWLEDGMENTS

This work is supported by funding from the UK Medical Research Council (MR/Z505547/1). LC is supported in part by the NIHR Imperial Biomedical Research Centre (BRC). RD, ED and LC acknowledge funding from the MRC Centre for Global Infectious Disease Analysis (reference MR/X020258/1), funded by the UK Medical Research Council (MRC). This UK funded award is carried out in the frame of the Global Health EDCTP3 Joint Undertaking. The authors would like to acknowledge Tim Dallman, Elita Jauneikaite, Nicholas Croucher, Yonatan Grad, Catrin Moore, Gus Hamilton, Gwen Knight, Till Bachmann, Edward Feil, Sarah Fortune, Xavier Didelot, Sean Nachtrab, the PHA4GE data structures working group, the ESGEM-AMR working group and the B2B2B AMR diagnostics network for their ongoing support of the project.

REFERENCES

[1] Rose, R., et al.: Comparing antimicrobial resistant genes and phenotypes across multiple sequencing platforms and assays for enterobacterales clinical isolates. BMC Microbiology **23**(1), 225 (2023). DOI 10.1186/s12866-023-02975-x. URL https://doi.org/10.1186/s12866-023-02975-x

[2] Rebelo, A.R., et al.: One day in denmark: Comparison of phenotypic and genotypic antimicrobial susceptibility testing in bacterial isolates from clinical settings. Frontiers in Microbiology **Volume 13 - 2022** (2022). DOI 10.3389/fmicb.2022.804627. URL https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2022.804627

[3] Domanovich-Asor, T., et al.: Genomic analysis of antimicrobial resistance genotype-to-phenotype agreement in *Helicobacter pylori*. Microorganisms **9**(1) (2021). URL https://www.mdpi.com/2076-2607/9/1/2

[4] Tacconelli, E., et al.: Discovery, research, and development of new antibiotics: the who priority list of antibiotic-resistant bacteria and tuberculosis. The Lancet Infectious Diseases **18**(3), 318–327 (2018). DOI https://doi.org/10.1016/S1473-3099(17)30753-3. URL https://www.sciencedirect.com/science/article/pii/S1473309917307533

[5] World Health Organization (WHO): Who bacterial priority pathogens list, 2024: Bacterial pathogens of public health importance to guide research, development and strategies to prevent and control antimicrobial resistance (2024). URL https://www.who.int/publications/i/item/9789240093461

[6] Bissonnette, L., Bergeron, M.G.: Infectious disease management through point-of-care personalized medicine molecular diagnostic technologies. Journal of Personalized Medicine **2**(2), 50–70 (2012). DOI 10.3390/jpm2020050. URL https://www.mdpi.com/2075-4426/2/2/50

[7] Kaprou, G.D., et al.: Rapid methods for antimicrobial resistance diagnostics. Antibiotics **10**(2) (2021). DOI 10.3390/antibiotics10020209. URL https://www.mdpi.com/2079-6382/10/2/209

[8] Antonopoulos, D.A., et al.: Patric as a unique resource for studying antimicrobial resistance. Briefings in Bioinformatics **20**(4), 1094–1102 (2017). DOI 10.1093/bib/bbx083. URL https://doi.org/10.1093/bib/bbx083

[9] National Center for Biotechnology Information (NCBI): Ncbi biosample antibiograms (2025). URL https://www.ncbi.nlm.nih.gov/biosample/docs/antibiogram/

[10] National Center for Biotechnology Information (NCBI): National database of antibiotic resistant organisms (ndaro) (2025). URL https://www.ncbi.nlm.nih.gov/pathogens/antimicrobial-resistance/

[11] Jolley, K., Bray, J., Maiden, M.: Open-access bacterial population genomics: Bigsdb software, the pubmlst.org website and their applications. Wellcome Open Research **3**(124) (2018). DOI 10.12688/wellcomeopenres.14826.1

[12] Argimn, S., et al.: Microreact: visualizing and sharing data for genomic epidemiology and phylogeography. Microbial Genomics **2**(11), e000093 (2016). DOI https://doi.org/10.1099/mgen.0.000093. URL https://www.microbiologyresearch.org/content/journal/mgen/10.1099/mgen.0.000093

[13] Centre for Genomic Pathogen Surveillance: Pathogenwatch URL https://pathogen.watch

[14] Centers for Disease Control and Prevention (CDC): Centers for disease control and prevention (u.s.), antibiotic resistance threats in the united states, 2019 (Nov 2019). URL http://dx.doi.org/10.15620/cdc:82532

[15] Karp, B.E., et al.: National antimicrobial resistance monitoring system: Two decades of advancing public health through integrated surveillance of antimicrobial resistance. Foodborne Pathogens and Disease **14**(10), 545–557 (2017). DOI 10.1089/fpd.2017.2283. URL https://doi.org/10.1089/fpd.2017.2283. PMID: 28792800

[16] Matamoros, S., et al.: Accelerating surveillance and research of antimicrobial resistance an online repository for sharing of antimicrobial susceptibility data associated with whole-genome sequences. Microbial Genomics **6**(5), e000342 (2020). DOI https://doi.org/10.1099/mgen.0.000342. URL https://www.microbiologyresearch.org/content/journal/mgen/10.1099/mgen.0.000342

[17] Pfizer: Atlas: Antimicrobial testing leadership and surveillance (2025). URL https://atlas-surveillance.com

[18] Berends, M.S., et al.: Amr: An r package for working with antimicrobial resistance data. Journal of Statistical Software **104**(3), 131 (2022). DOI 10.18637/jss.v104.i03. URL https://www.jstatsoft.org/index.php/jss/article/view/v104i03

[19] Feldgarden, M., et al.: Amrfinderplus and the reference gene catalog facilitate examination of the genomic links among antimicrobial resistance, stress response, and virulence. Scientific Reports **11**(1), 12,728 (2021). DOI 10.1038/s41598-021-91456-0. URL https://doi.org/10.1038/s41598-021-91456-0

[20] Ngabonziza, J.C.S., et al.: Multidrug-resistant tuberculosis control in rwanda overcomes a successful clone that causes most disease over a quarter century. Journal of Clinical Tuberculosis and Other Mycobacterial Diseases **27**, 100,299 (2022). DOI https://doi.org/10.1016/j.jctube.2022.100299. URL https://www.sciencedirect.com/science/article/pii/S2405579422000043

[21] Raynaud, M., et al.: Impact of the covid-19 pandemic on publication dynamics and non-covid-19 research production. BMC Medical Research Methodology **21**(1), 255 (2021). DOI 10.1186/s12874-021-01404-9. URL https://doi.org/10.1186/s12874-021-01404-9

[22] Tomczyk, S., et al.: Impact of the covid-19 pandemic on the surveillance, prevention and control of antimicrobial resistance: a global survey. Journal of Antimicrobial Chemotherapy **76**(11), 3045–3058 (2021). DOI 10.1093/jac/dkab300. URL https://doi.org/10.1093/jac/dkab300

[23] Rodríguez-Baño, J., et al.: Key considerations on the potential impacts of the covid-19 pandemic on antimicrobial resistance research and surveillance. Transactions of The Royal Society of Tropical Medicine and Hygiene **115**(10), 1122–1129 (2021). DOI 10.1093/trstmh/trab048. URL https://doi.org/10.1093/trstmh/trab048

[24] Stotzem, N., Guntoro, F., Chindelevitch, L.: BenchmarkDR: A modular and expandable benchmarking pipeline for machine learning based antimicrobial resistance prediction (2022). URL https://mlmg2022.github.io/assets/papers/Stotzem.pdf

[25] Walter, K.S., et al.: Genomic variant-identification methods may alter mycobacterium tuberculosis transmission inferences. Microbial Genomics **6**(8), e000418 (2020). DOI https://doi.org/10.1099/mgen.0.000418. URL https://www.microbiologyresearch.org/content/journal/mgen/10.1099/mgen.0.000418

[26] Chindelevitch, L., van Dongen, M., Graz, H., Pedrotta, A., Suresh, A., Uplekar, S., Jauneikaite, E., Wheeler, N.: Ten simple rules for the sharing of bacterial genotypephenotype data on antimicrobial resistance. PLOS Computational Biology **19**(6), e1011,129 (2023). DOI 10.1371/journal.pcbi.1011129. URL http://dx.doi.org/10.1371/journal.pcbi.1011129