Bioplastic Production Potential of *Azospira suillum* PS: Growth-Associated PHB Production Under Aerobic and Anaerobic Conditions

David A.O. Meier[1], Benjamin Glazer[1], V. Celeste Lanclos[1], Hans K. Carlson[2], John D. Coates[1*]

[1] Department of Plant and Microbial Biology, University of California, Berkeley, Berkeley, CA,
United States

[2] Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory,
Berkeley, CA, United States

* Correspondence: John D. Coates (jdcoates@berkeley.edu), Department of Plant and
Microbiology, 241 Koshland Hall, University of California, Berkeley, CA; Tel +1-510-643-8455

Competing Interests Statement:

The authors do not declare any competing interests.

Abstract

Plastics are indispensable in modern society, yet less than 10% of the 380 million tons
produced annually are recycled. This accumulation of non-degradable waste has intensified the
search for sustainable alternatives. Polyhydroxyalkanoates (PHAs), particularly polyhydroxybutyrate
(PHB), are biodegradable polymers produced by microorganisms. Here, we investigate the capacity
of *Azospira suillum* PS, a model facultative perchlorate-reducing bacterium, to synthesize PHB. We
show that unlike most organisms, PHB production is growth-associated both aerobically, and
anaerobically with perchlorate as an electron acceptor under moderate nitrogen deplete conditions.
Genomic analysis revealed four *phaC* homologs, with one, *phaC4,* co-localized with *phaB* and *phaR*,
suggesting a conserved operon. While three of the *phaC* genes were successfully deleted, *phaC4*
could not be disrupted, indicating an essential role for growth. Redox profiling revealed a positive
correlation between PHB accumulation and NADPH/NADP$^+$ ratios, supporting the role of
cytoplasmic reducing equivalents in PHB synthesis. This study presents the first demonstration of
PHA production under perchlorate-respiring conditions and one of the few documented cases of
growth-associated PHA synthesis. It offers new insights into the genetic and metabolic basis of PHB
production in *A. suillum* PS, underscoring its potential as a platform organism for sustainable,
growth-coupled bioplastic production.

Introduction

The use of processed polymers by humans dates to at least 1600 B.C.E. (Hosler et al., 1999),
but the invention of modern thermoplastics by Charles Goodyear in 1839 revolutionized their
industrial and environmental impact. Today, thermoplastics are indispensable across nearly every
sector due to their lightweight, durable, and versatile properties (Andrady & Neal, 2009). This utility
has driven global plastic production to over 380 million tons annually (Geyer et al., 2017). However,
nearly all conventional plastics are derived from fossil hydrocarbons and are non-biodegradable,
leading to long-term environmental accumulation. This persistent pollution, along with the carbon
footprint associated with petrochemical feedstocks, has accelerated interest in sustainable alternatives
such as bioplastics—materials synthesized from renewable sources.

Polyhydroxyalkanoates (PHAs) are one such family of bioplastics that are fully compostable,
produced by microorganisms as intracellular carbon and energy storage polymers (Macrae &
Wilkinson, 1958). PHAs are classified by monomer length into short-chain-length (scl-PHA, C3–C5)
and medium-chain-length (mcl-PHA, C6–C14) types, with copolymers offering diverse material
properties (Steinbüchel, 1991; Chen et al., 2015; McAdam et al., 2020). This chemical versatility
allows tailoring of rigidity, flexibility, and degradability for various applications (Gao et al., 2011).
However, commercial PHA adoption is constrained by cost of production—typically $3–8 per kg in
2025, compared to ~$1.40 for petrochemical polyethylene—with global production representing less
than 0.5% of total plastics (Rosenboom et al., 2022). Reducing cost of production through
improvement of microbial titers and yields requires deeper insight into PHA biosynthetic regulation
and physiology.

Among PHAs, polyhydroxybutyrate (PHB) is the best studied, first identified in *Bacillus*
subtilis in 1926 (Lemoigne). Since then, PHAs have been detected in diverse bacteria (both
autotrophs and heterotrophs), archaea (Poli et al., 2011), and some yeasts (Thu et al., 2023), where
they typically accumulate under nutrient-limiting, carbon-rich conditions. PHB biosynthesis involves
a core pathway encoded by *phaA* (β-ketothiolase), *phaB* (acetoacetyl-CoA reductase), and *phaC*
(PHA synthase), which polymerizes 3-hydroxybutyryl-CoA. Additional regulators such as *phaR* and
granule-associated *phaP* (phasin) influence polymer accumulation and granule morphology (Pötter et
al., 2005). PHA synthases (PhaC enzymes) are grouped into four classes (I–IV) based on their
substrate specificity and subunit composition, rather than their evolutionary lineage. Classes I, III,
and IV primarily synthesize short-chain-length PHAs like PHB but differ in enzyme architecture:
Class I enzymes are homodimers composed of two identical PhaC subunits; Class III enzymes
function as heterodimers consisting of PhaC and a partner protein, PhaE; and Class IV enzymes also
form heterodimers, pairing PhaC with PhaR. In contrast, Class II enzymes are specialized for
producing medium-chain-length PHAs and typically include multiple PhaC isoforms (e.g., PhaC1
and PhaC2), without accessory subunits like PhaE or PhaR. A deeper understanding of how
feedstock composition and PHA synthase structure influence polymer diversity and accumulation
remains a key research goal.

Cupriavidus necator H16 is a model PHA producer and has been extensively engineered for
high PHB yields (Zhang et al., 2022). Strategies include promoter and pathway optimization (Tang et
al., 2020), substrate engineering (Mifune et al., 2010), and process development (Atlić et al., 2011).
However, a key limitation is that PHB production in *C. necator* is not growth-associated, requiring
nutrient shifts or batch processes, which hinders continuous cultivation (Henderson & Collins, 1997)
and thus loss in industrial efficiency. Broader surveys have revealed PHA production in
extremophiles, anaerobes, and oligotrophs (Wang et al., 2019a; Bennett et al., 2024), suggesting that
novel chassis organisms with more favorable production dynamics remain to be explored.

Dissimilatory perchlorate-reducing bacteria (DPRB) represent a promising but understudied
group in this context. Perchlorate (ClO_4^-) is a highly oxidized, water-soluble oxyanion with a
biological redox potential comparable to oxygen ($E°' = +0.797$ V) (Youngblut et al., 2016). In
DPRB, perchlorate is reduced to chlorite by the perchlorate reductase complex (PcrAB), followed by
dismutation into chloride and molecular oxygen via chlorite dismutase (Cld). The oxygen is
subsequently respired by the microorganism (Youngblut et al., 2016). This pathway enables the
endogenous generation of O_2 under anoxic conditions, supporting oxygenase-dependent metabolism
without external aeration (Carlström et al., 2015). The conserved, mobile perchlorate reduction island
(PRI) allows this metabolism to be transferred across diverse taxa, supporting bioengineering efforts
(Melnyk et al., 2015). *Azospira suillum* PS is a facultative anaerobe and genetically tractable DPRB

capable of coupling perchlorate reduction to energy conservation and growth. Despite its unique
metabolism and flexible redox physiology, its potential for bioproduct synthesis remains unexplored.

In this study, we investigate the genetic and physiological basis of PHB biosynthesis in *A.*
suillum PS. Using comparative genomics, gene deletion, redox analysis, and polymer quantification,
we characterize the pathways supporting PHB production during both aerobic and perchlorate-
respiring growth. Our findings highlight *A. suillum* PS as a unique candidate for sustainable, growth-
coupled bioplastic production in anaerobic biotechnological systems.

Materials and Methods

Bacterial Strains and Plasmids

92   *Azospira suillum* PS (ATCC BAA-33/DSMZ 13638) was revived from laboratory freezer
93   stocks and used as the wild-type strain for all genetic manipulations. *Escherichia coli* XL1-Blue was
94   used for plasmid propagation and cloning. A full list of primers (Supplementary Table 1), strains
95   (Supplementary Table 2), and plasmids (Supplementary Table 3) is provided in the supplementary
96   materials. Prior to all experiments, strains were streaked from freezer stocks to obtain single colonies.

Mutant Construction

Gene deletion mutants in *A. suillum* PS were generated using a sacB-based counterselection
approach (Gay et al., 1983). Electrocompetent cells were transformed with either integrative (~500
ng) or replicative (~50 ng) plasmids in 1 mm electroporation cuvettes using a Gene Pulser Xcell
electroporator (Bio-Rad, USA) with the following settings: 1.8 kV, 25 µF, and 200 Ω. Cells were
chilled on ice before electroporation; cuvettes were kept at room temperature. After pulsing, cells
were recovered in pre-warmed ALP medium at 37°C for 6 h, then 100 µL of culture was plated
directly on ALP agar containing 50 µg/mL kanamycin. Colonies appearing within 48 h were
considered valid transformants and picked into 500 µL ALP broth with 50 µg/mL kanamycin. After 4
h of growth, cultures were plated on ALP agar with 6% sucrose for counterselection. Representative
colonies were screened by colony PCR to confirm deletions. Glycerol stocks were prepared at 15%
final concentration for storage at −80°C.

Culture Conditions and Growth Media

E. coli strains were cultured in LB medium with 50 µg/mL kanamycin. Wild-type and mutant
A. suillum PS strains were grown in acetate minimal medium (AMM). One liter of AMM contained:
0.49 g sodium phosphate monobasic dihydrate, 0.97 g sodium phosphate dibasic anhydrous, 0.1 g
KCl, 0.25 g NH_4Cl, and 10 mL each of vitamin and mineral mix (Bruce et al., 1999). The sodium
acetate concentration in AMM was either 0.82g/L acetate for 10mM AMM or 2.05 g acetate for
25mM AMM. To prepare mixed carbon source media (acetate, lactate, pyruvate), AMM was
supplemented with 2.0 g yeast extract, 7.6 mL of a 60% (w/w) sodium lactate solution, and 1.10 g
sodium pyruvate. For solid media, 15 g/L agar was added. *A. suillum* PS was cultured in 250 mL
baffled Erlenmeyer flasks containing 50 mL of medium at 37°C and 250 rpm. Optical density at 600
nm (OD_{600}) was measured using a 1 mm pathlength cuvette on a Genesys 20 spectrophotometer
(Thermo Scientific, USA).

PHA Extraction and GC-MS Analysis

PHB was quantified from 40 mL of culture. Samples were harvested by centrifugation at 7,197 × g for 10 min at 4°C, and pellets were stored at −80°C. Acidic methanolysis was performed as described by Juengert et al. (2018). Extracts were analyzed by GC-MS using a 7890A GC system (Agilent Technologies, USA) equipped with a DB-WAX UI column and coupled to a 5975C XL EI/CI Mass Selective Detector. The temperature program was: 80°C for 2 min, ramp to 210°C at 10°C/min, then to 250°C at 50°C/min with a 1 min hold. Poly[(R)-3-hydroxybutyric acid] (Santa Cruz Biotechnology, USA) was used for the standard curve, and methyl benzoate (Sigma-Aldrich, USA) served as the internal standard. Compound identity was confirmed by total ion count (TIC) spectra matching the NIST EI database. The presence of polyhydroxyvalerate (PHV) and polyhydroxyhexanoate PHHX was assessed using a commercial PHA standard (Sigma-Aldrich, USA), and TIC peaks were compared with NIST spectra for 3-hydroxypentanoic acid methyl ester and 3-hydroxyhexanoic acid methyl ester.

NAD(P)$^+$/H Quantification

Redox metabolite analysis was performed as described by Kern et al. (2014). Samples were kept on ice and protected from light until incubated at 37°C. To prevent redox degradation, only three samples were processed at a time. Quantification was performed using a Varian Cary 50 MPR Microplate Reader (Agilent Technologies, USA).

Fluorescence Microscopy

Microscopy was conducted as described by Juengert et al. (2018). Cells were harvested in exponential phase, washed once in PBS (8 g NaCl, 201 mg KCl, 1.42 g Na_2HPO_4, 245 mg KH_2PO_4 per liter, pH 7.4), and stained with 1 µg/mL Nile Red for 10 min in the dark. Samples were mounted on 1% agarose pads and visualized on a Zeiss LSM880 confocal microscope (Zeiss, Germany) using a 100× Plan-Apochromat NA 1.4 objective. Nile Red was excited at 561 nm, and emission was detected from 570–620 nm using a GaAsP photon-counting detector. Differential interference contrast (DIC) images were acquired simultaneously. Image acquisition settings were consistent across all samples and processed in ZEN software v2.3 SP1 (Zeiss).

Phylogenetic Tree Construction

Each *phaC* gene from *A. suillum* PS was queried against the NCBI nr database using BLASTP (Altschul et al., 1990). The top 250 unique hits (excluding "multispecies" entries) were retained per gene. PhaC homologs from *Azospira* species were manually reviewed and validated using HMMER v3.4 with TIGRfam-specific models (Finn et al., 2011). Alignments were generated with MUSCLE v3.8.31 and trimmed using trimAl v1.4 (Edgar, 2004; Capella-Gutiérrez et al., 2009). Phylogenies were inferred using IQ-TREE v2.0.6 with 1,000 ultrafast bootstraps and the Q.pfam+I+R7 model. Trees were visualized using iTOL v6 (Minh et al., 2020; Letunic and Bork, 2020).

In Silico Comparative Genomics

The genome of *A. suillum* PS was downloaded from NCBI (Sayers et al., 2024); *Dechloromonas agitata* CKB and additional *Azospira* genomes were retrieved from JGI-IMG (Markowitz et al., 2006). Genomes were searched using HMMER v3.4 with a custom set of

160 TIGRfams relevant to PHA metabolism (Supplementary Table 4). Presence/absence patterns of PHA
161 biosynthesis genes were visualized using R v4.4.2 and RStudio v2024.12.0+467.

Results

Genomic and microscopic evidence for PHB granule formation in *Azospira suillum* PS and other perchlorate-reducing bacteria

Previous work on the perchlorate-reducing bacterium *Dechloromonas agitata* CKB revealed
electron-dense intracellular structures that were hypothesized to be PHA granules (Fig. 1a) (Bruce et
al., 1999). *In silico* analysis (Fig. 1) confirmed that *D. agitata* CKB possesses all the core genes
necessary for PHA biosynthesis, including *phaA*, *phaB*, *phaC*, *phaP*, and *phaR*, which together enable
the conversion of acetyl-CoA to PHB and the regulation and granule-associated assembly of PHA. To
further investigate this capacity in a more genetically tractable system, we turned to *Azospira suillum*
PS, a model perchlorate-reducing bacterium with established genetic tools (Melnyk et al., 2014). Strain
PS also contains the requisite genes for PHA biosynthesis (Fig. 1); however, unlike CKB, it
additionally encodes the PHA depolymerase *phaZ*. Both organisms possess multiple copies of the *phaC*
gene—three in CKB and four in PS (see Supplementary Table 5 for locus tags in PS). In both strains,
one *phaC* copy is co-localized with *phaB* and *phaR* (Fig. 1), suggesting an operon structure, while the
remaining *phaC* copies are dispersed throughout the genome. Of these, only one has a neighboring
phaP gene, and no other adjacent genes associated with PHA biosynthesis were observed.

To determine whether *A. suillum* PS produced PHB granules, we stained cultured cells with
Nile Red, a lipophilic dye commonly used to visualize PHB granules and examined them using
confocal microscopy (Fig. 1b). PS cells exhibited visible intracellular PHB granules with an
approximate diameter of 0.32 µm. Given the ~2 µm cell diameter of PS, the clustered granules span
much of the intracellular space, corresponding to ~64% of the estimated cell volume being occupied
by PHB granules. Cells stained positive for PHB granules under both aerobic and perchlorate-respiring
conditions, providing the first clear evidence of PHA production linked to perchlorate respiration. This
observation confirmed the genetic potential for PHB accumulation; however, it remained unclear
whether polymer synthesis occurred constitutively or only in response to specific nutrient conditions.

To establish whether PHB accumulation in *A. suillum* PS is constitutive or nutrient-dependent,
we first evaluated growth under nitrogen-replete conditions. Cultures were grown in standard acetate
minimal medium (AMM) containing 10 mM sodium acetate and 4.68 mM NH_4Cl, corresponding to a
replete C:N ratio of approximately 1:0.23 relative to the canonical Redfield ratio (C:N = 1:0.15)
(Redfield, 1934). Under these conditions, no detectable PHB was observed by GC–MS analysis (Fig.
1d), confirming that PHB synthesis does not occur under nitrogen-replete growth. The culture exhibited
a specific growth rate (µ) of 0.60 h^{-1} and a doubling time of 1.16 h, indicating robust biomass formation
despite the absence of polymer accumulation.

Given prior genomic and microscopic evidence suggesting PHB-producing potential, we next
examined whether altering the carbon-to-nitrogen ratio could induce polymer synthesis. When the
acetate concentration was increased to 25 mM while maintaining 4.68 mM NH_4Cl (C:N= 1:0.094),
PHB production was observed under both aerobic and perchlorate-respiring conditions (Fig. 2a–d).
Under these carbon-enriched conditions, cell experience ~38% less nitrogen per unit carbon than
under the canonical Redfield ratio, experiencing moderate nitrogen limitation. Under these conditions
A. suillum PS exhibited a slightly reduced specific growth rate (µ = 0.50 h^{-1}; doubling time = 1.39 h)
but accumulated measurable intracellular PHB. These results demonstrate that PHB synthesis in *A.*

suillum PS is triggered by a shift toward carbon surplus relative to nitrogen, rather than being a constitutive feature of its metabolism.

Following this finding, we conducted a detailed time-course analysis of PHB accumulation under the carbon-enriched condition to quantify the extent and dynamics of polymer synthesis during aerobic and perchlorate-respiring growth.

Growth-associated PHB production in *Azospira suillum* PS under aerobic and perchlorate-respiring conditions

To quantify PHA production by *A. suillum* PS, gas chromatography–mass spectrometry (GC-MS) was performed on samples extracted via acidic methanolysis. In addition to quantifying total PHB produced by the culture, this method also enabled identification of the different monomers present. When PS was grown on acetate under either aerobic or perchlorate-respiring conditions 3-hydroxybutyric acid (3-HB) (the monomer of PHB) was detected. PHB was quantified across the growth curve and was found to be produced in a growth-associated manner (Fig 2a-d). While growth-associated PHB production has been previously reported (Scott et al., 2021; Amos & McInerney, 1989; Ackermann & Babel, 1997); PHB biosynthesis is more commonly observed during stationary phase, in response to nutrient limitation (Anderson & Dawes, 1990).

When grown aerobically, *A. suillum* PS reaches a maximum OD_{600nm} of 0.747 ± 0.008 during late exponential phase, at which point it had produced a maximum of 0.077 ± 0.006 mg/mL PHB providing a total yield of 0.10 g/L per OD_{600nm} (Fig 2a). Under perchlorate-respiring conditions with 10 mM perchlorate, PS reaches a lower maximum OD_{600nm} of 0.397 ± 0.009 and produces a maximum of 0.024 ± 0.008 mg/mL PHB or 0.06 g/L per OD_{600nm} (Fig 2b). When grown on perchlorate, PHB levels slightly decline as the culture enters stationary phase (Fig 2d). There is no significant reduction in PHB levels between peak and late-stage culture under aerobic conditions, despite a drop in OD_{600nm} of approximately 0.21. Based on the 1.6-fold higher PHB yield in aerobic versus perchlorate conditions, we further characterized PHB production by PS under aerobic growth.

This growth-associated pattern under carbon-enriched conditions contrasts with the nitrogen-replete culture, where no PHB was detected, reinforcing that polymer synthesis in *A. suillum* PS is activated only when the C:N ratio exceeds a physiological threshold. Under both mild and severe nitrogen limitation, PHB production remained unchanged, which is consistent with observations in other anaerobic organisms such as *Syntrophomonas wolfei*, which produced PHB independently of ammonium chloride concentration (Amos & McInerney, 1989).

When examining normalized PHB levels relative to optical density, both aerobic and perchlorate-grown cultures show evidence of carryover PHB from the inoculum (Fig 2c&d). In aerobic cultures, this residual PHB is rapidly degraded during lag phase prior to new PHB accumulation during log phase, suggesting that the cells may utilize the carryover PHB as an additional electron donor alongside the supplied acetate during the initial phase of growth (Fig 2c). In contrast, perchlorate-respiring cultures do not exhibit this initial degradation phase; instead, normalized PHB/OD_{600} remains stable during early growth and begins increasing only during exponential phase.

Redox cofactor dynamics link NADPH availability to PHB biosynthesis in *Azospira suillum* PS

To assess the redox influence on PHB biosynthesis, we measured intracellular NADH/NAD$^+$
and NADPH/NADP$^+$ ratios over a 24-hour culture period and compared these values to PHB
accumulation (Fig. 3a–d). When grown aerobically, *Azospira suillum* PS exhibited an average
NADH/NAD$^+$ ratio of 0.491—more than twice the reported aerobic values for *E. coli* (0.226) and
Klebsiella aerogenes (0.195)—indicating a relatively reduced intracellular environment. Notably,
this ratio also exceeded those observed under anaerobic conditions for both comparative organisms
(0.396 and 0.219, respectively) (Wimpenny & Firth, 1972). Only *Pseudomonas aeruginosa*, a known
respiratory generalist, exhibited a higher aerobic NADH/NAD$^+$ ratio (0.571). While *E. coli* lacks the
genetic capacity for native PHA production, some *Pseudomonas* spp. are known PHB producers,
although not typically in a growth-associated fashion (Bertrand et al., 1990).

Spearman correlation analysis revealed a strong and statistically significant monotonic
relationship between PHB levels and NADPH/NADP$^+$ ($\rho = 0.829$, $p = 0.042$), while no significant
correlation was observed with NADH/NAD$^+$ ($\rho = 0.600$, $p = 0.208$). This is consistent with the
known cofactor specificity of the PhaB acetoacetyl-CoA reductase enzyme. In *Cupriavidus necator*
H16—a well-studied model organism for PHB production—this enzyme has been shown to
preferentially use NADPH rather than NADH as the reducing equivalent source (Steinbüchel &
Schlegel, 1991).

Phylogenomic analysis reveals conserved, diversified *phaC* loci across the *Azospira* genus

A maximum-likelihood phylogenetic tree (Fig 4) constructed from the four unique *phaC* gene
products identified in *Azospira suillum* PS, along with their 250 closest homologs, revealed that each
PhaC protein is nested within a monophyletic clade composed of *phaC* homologs from other
Azospira species. Despite this shared genus-level affiliation, the four *phaC* variants from strain PS
occupy distinct and divergent branches within the tree, suggesting that each *phaC* gene may have
evolved independently. Furthermore, this analysis revealed that multiple *phaC* gene copies are
consistently present in *Azospira* genomes. Comparative genomic analysis across the genus showed
that each *Azospira* species encodes at least three distinct copies of *phaC*, alongside a single copy of
the transcriptional regulator *phaR*, suggesting that PHA biosynthesis is a conserved metabolic feature
within the genus. To assess whether these genes may have been acquired via horizontal gene transfer,
we used IslandViewer (Bertelli et al., 2017), which predicts genomic islands based on sequence
composition and comparative genomics. This analysis showed no evidence of horizontal gene
transfer near the *phaC* loci in strain PS (data not shown).

To assess conservation of gene neighborhoods, we used CAGE-CAT and Clinker to compare
the genomic loci surrounding *phaC* across *Azospira* genomes (Gilchrist & Chooi, 2021). These tools
align and visualize syntenic regions based on protein sequence identity and gene order, enabling
rapid identification of conserved operons and structural rearrangements. This analysis (Fig 5)
revealed that the local gene context of *phaC* is highly conserved, with pairwise amino acid identity of
phaC homologs exceeding 70% across the genus, further supporting their functional importance in
Azospira. However, while average nucleotide identity remains high across *Azospira* species, they do
not share syntenic conservation of the broader PHA biosynthesis gene cluster.

Functional characterization of *phaC* genes reveals essential role of *phaC4* in PHB biosynthesis

Our analysis found that *Azospira suillum* PS encodes only class I *phaC* genes, which are
typically associated with the production of short-chain-length (scl) PHAs such as PHB. To
investigate the four *phaC* genes in *A. suillum* PS responsible for PHA production, we attempted to

generate deletion mutants for each gene. Knockout strains were successfully constructed for three out
of the four *phaC* genes. However, under all tested growth conditions, we were unable to generate a
mutant lacking the *phaC* gene with the locus tag *dsui_2537* (*phaC*4). Notably, this gene is co-
localized with *phaB* and *phaR*. Previous transposon mutagenesis studies in strain PS also identified
*phaC*4 as essential under all conditions tested (Supplementary Figure 1).

To further assess the functional role of each *phaC*, we compared the PHA monomer profiles
of all knockout strains under both defined and complex media conditions. As shown in Table 1, none
of the knockout strains displayed a PHA profile distinct from wild type. All strains consistently
produced PHB, while polyhydroxyvalerate (PHV) and polyhydroxyhexanoate (PHHX) were absent
(data not shown). These results suggest that the essential *phaC*4 is playing an important physiological
role during normal growth in nutrient replete media.

Discussion

This study provides new insight into the biosynthesis and regulation of polyhydroxybutyrate
(PHB) in *Azospira suillum* PS, a model perchlorate-reducing bacterium. We demonstrate for the first
time that PHB biosynthesis occurs under perchlorate-respiring conditions, expanding the known
metabolic range for polyhydroxyalkanoate (PHA) production. Notably, *A. suillum* PS accumulates
PHB in a growth-associated manner under both aerobic and anaerobic conditions, with aerobic
cultures producing 1.6-fold more PHB. PHB synthesis was absent under nitrogen-replete conditions
and was only observed when the carbon-to-nitrogen ratio was altered to a ~40 % reduction in
nitrogen availability relative to the canonical Redfield ratio. This establishes that PHB production in
A. suillum PS is triggered by moderate nutrient imbalance. This pattern contrasts with the stress-
induced, stationary-phase accumulation commonly observed in other bacteria, such as *C. necator*,
suggesting that PHB plays a role in the central metabolism of *A. suillum* PS under nitrogen limitation
where it is coupled to growth under both aerobic and perchlorate reducing conditions.

Temporal analysis of PHB accumulation revealed distinct regulatory patterns between
respiratory modes. Under aerobic growth, normalized PHB/OD$_{600}$ initially declines during lag phase,
likely reflecting mobilization of stored carbon and reducing equivalents. In contrast, perchlorate-
respiring cultures do not exhibit this early decrease; instead, PHB levels remain stable in early
growth and begin to rise during exponential phase.

Our redox profiling data support this hypothesis. A strong and statistically significant
correlation was observed between intracellular NADPH/NADP$^+$ ratios and PHB content (Spearman ρ
= 0.829, p = 0.042), implicating NADPH availability as a potential driver of PHB accumulation.
Although this observation suggests a role for NADPH in fueling PHB biosynthesis, this correlation
does not confirm cofactor specificity of the biosynthetic enzymes. NADH and NADPH levels are
often tightly coupled and metabolically interconvertible, and the observed association may reflect a
broader redox state rather than direct NADPH usage. Further biochemical characterization will be
required to determine whether PhaB or related enzymes in *A. suillum* PS preferentially utilize
NADPH over NADH.

Genomic analysis revealed four *phaC* genes encoding class I PHA synthases in *A. suillum* PS.
Of these, *phaC4* is particularly notable for its synteny with *phaB* and *phaR*, suggesting a conserved
operon structure. No evidence of horizontal gene transfer was detected near any *phaC* loci, indicating
that these genes are native and vertically inherited. Functional analysis supports the essentiality of
phaC4, as multiple attempts to generate a knockout mutant were unsuccessful under all tested

conditions. These findings further reinforce the hypothesis that PHB biosynthesis plays a central
physiological role in this organism, potentially as a buffer for intracellular reducing equivalents.

From an applied perspective, *A. suillum* PS demonstrates rapid and efficient PHB accumulation
on simple substrates. Under aerobic conditions, wild-type PS produces 8.5 mg/L/h PHB, reaching
77 mg/L in just 9 hours using a single 25 mM acetate dose (600 mg/L carbon). While this yield is
lower than that of *Cupriavidus necator*—which reaches up to 79.2 mg/L/h on soybean oil after 72
hours—*C. necator* requires multiple feedings totaling ~7,000 mg/L carbon (Mifune et al., 2010),
greater than tenfold more than the PS culture. In contrast, PS grows in minimal medium without
supplementation, avoiding the cost and handling challenges of plant oils. These characteristics
position *A. suillum* PS as a promising chassis for continuous, low-input bioplastic production.

With targeted metabolic engineering—such as overexpression of *phaCBR* and optimization of
carbon feed strategies—*A. suillum* PS may achieve higher titers while retaining its growth-associated
PHB phenotype. This production strategy is particularly well-suited for bioprocesses favoring
continuous synthesis and redox-balanced metabolism. Overall, our findings position *A. suillum* PS as
a novel and sustainable chassis for industrial PHA production, especially in systems where
simplicity, growth-coupling, and cost-effective carbon utilization are critical design goals.

Industrial relevance

This study identifies *Azospira suillum* PS as a promising new chassis for sustainable
bioplastic production. Unlike most known PHA producers, *A. suillum* PS accumulates PHB in a
growth-associated manner under both aerobic and anaerobic (perchlorate-respiring) conditions,
enabling simpler, continuous fermentation processes. Its ability to produce PHB rapidly on low-cost
substrates in minimal media, combined with native genetic tractability and redox-linked regulation,
makes it an attractive platform for scalable, low-input, and flexible biomanufacturing—particularly in
resource-limited or oxygen-poor environments.

Conflict of Interest

The authors declare no competing commercial, or financial interests in relation to the work presented
here.

Author Contributions

DAOM, HKC and JDC designed research. DAOM and BG optimized the genetic protocol and BG
constructed the mutants. DAOM and VCL generated the phylogenetic analysis. DAOM performed all
physiology experiments and measurements and comparative genomic analysis. DAOM wrote the
draft manuscript and created the figures with guidance from JDC. All authors contributed to data
analysis, reviewed the manuscript and approved of its publication.

Funding

Funding for this research was supported in part by generous sponsorship from the American Pacific
Corporation and the Baydin (Boomerang) Corporation.

Acknowledgments

The authors acknowledge Yi Liu for guidance on instrumentation and analyses, Dr. Denise Schichnes
at the UC Berkeley Biological Imaging Facility for training and access to microscopy, and Dr.
Ruiwen Hu for his advice on computational analyses.

References

Achenbach, L A, U Michaelidou, R A Bruce, J Fryman, and J D Coates. "Dechloromonas Agitata
Gen. Nov., Sp. Nov. and Dechlorosoma Suillum Gen. Nov., Sp. Nov., Two Novel Environmentally
Dominant (per)Chlorate-Reducing Bacteria and Their Phylogenetic Position." *International Journal*
of Systematic and Evolutionary Microbiology 51, no. 2 (March 1, 2001): 527–33.
https://doi.org/10.1099/00207713-51-2-527.

Ackermann, J.-U., and W. Babel. "Growth-Associated Synthesis of Poly(Hydroxybutyric Acid) in
Methylobacterium Rhodesianum as an Expression of an Internal Bottleneck." *Applied Microbiology*
and Biotechnology 47, no. 2 (1997): 144–49. https://doi.org/10.1007/s002530050903.

Altschul, Stephen F., Warren Gish, Webb Miller, Eugene W. Myers, and David J. Lipman. "Basic
Local Alignment Search Tool." *Journal of Molecular Biology* 215, no. 3 (October 5, 1990): 403–10.
https://doi.org/10.1016/S0022-2836(05)80360-2.

Amos, Dale A., and Michael J. McInerney. "Poly-β-Hydroxyalkanoate in Syntrophomonas Wolfei."
Archives of Microbiology 152, no. 2 (July 1989): 172–77. https://doi.org/10.1007/BF00456097.

Anderson, A. J., and E. A. Dawes. "Occurrence, Metabolism, Metabolic Role, and Industrial Uses of
Bacterial Polyhydroxyalkanoates." *Microbiological Reviews* 54, no. 4 (December 1990): 450–72.
https://doi.org/10.1128/mr.54.4.450-472.1990.

Atlić, Aid, Martin Koller, Dietrich Scherzer, Christoph Kutschera, Elizabeth Grillo-Fernandes,
Predrag Horvat, Emo Chiellini, and Gerhart Braunegg. "Continuous Production of Poly([R]-3-
Hydroxybutyrate) by Cupriavidus Necator in a Multistage Bioreactor Cascade." *Applied*
Microbiology and Biotechnology 91, no. 2 (July 2011): 295–304. https://doi.org/10.1007/s00253-
011-3260-0.

Bennett, Brittany D., David A.O. Meier, V. Celeste Lanclos, Hasti Asrari, John D. Coates, and J.
Cameron Thrash. "Polyhydroxybutyrate Production in Freshwater SAR11 (LD12)." Microbiology,
November 8, 2024. https://doi.org/10.1101/2024.11.08.622676.

Bertelli, Claire, Matthew R Laird, Kelly P Williams,
Simon Fraser University Research Computing Group, Britney Y Lau, Gemma Hoad, Geoffrey L
Winsor, and Fiona SL Brinkman. "IslandViewer 4: Expanded Prediction of Genomic Islands for
Larger-Scale Datasets." *Nucleic Acids Research* 45, no. W1 (July 3, 2017): W30–35.
https://doi.org/10.1093/nar/gkx343.

Bertrand, J.-L., B. A. Ramsay, J. A. Ramsay, and C. Chavarie. "Biosynthesis of Poly-β-
Hydroxyalkanoates from Pentoses by *Pseudomonas Pseudoflava*." *Applied and Environmental*
Microbiology 56, no. 10 (October 1990): 3133–38. https://doi.org/10.1128/aem.56.10.3133-
3138.1990.

Bruce, Royce A., Laurie A. Achenbach, and John D. Coates. "Reduction of (per)Chlorate by a Novel Organism Isolated from Paper Mill Waste." *Environmental Microbiology* 1, no. 4 (August 1999): 319–29. https://doi.org/10.1046/j.1462-2920.1999.00042.x.

Byrne-Bailey, Kathryne G., and John D. Coates. "Complete Genome Sequence of the Anaerobic Perchlorate-Reducing Bacterium Azospira Suillum Strain PS." *Journal of Bacteriology* 194, no. 10 (April 25, 2012): 2767–68. https://doi.org/10.1128/jb.00124-12.

Capella-Gutiérrez, Salvador, José M. Silla-Martínez, and Toni Gabaldón. "trimAl: A Tool for Automated Alignment Trimming in Large-Scale Phylogenetic Analyses." *Bioinformatics (Oxford, England)* 25, no. 15 (August 1, 2009): 1972–73. https://doi.org/10.1093/bioinformatics/btp348.

Carlström, Charlotte I., Dana Loutey, Stefan Bauer, Iain C. Clark, Robert A. Rohde, Anthony T. Iavarone, Lauren Lucas, and John D. Coates. "(Per)Chlorate-Reducing Bacteria Can Utilize Aerobic and Anaerobic Pathways of Aromatic Degradation with (Per)Chlorate as an Electron Acceptor." Edited by Victor De Lorenzo and Sang Yup Lee. *mBio* 6, no. 2 (May 2015): e02287-14. https://doi.org/10.1128/mBio.02287-14.

Chen, Guo-Qiang, Juanyu Zhang, and Ying Wang. "White Biotechnology for Biopolymers." In *Industrial Biorefineries & White Biotechnology*, 555–74. Elsevier, 2015. https://doi.org/10.1016/B978-0-444-63453-5.00018-5.

Coates, John D., and Laurie A. Achenbach. "Microbial Perchlorate Reduction: Rocket-Fuelled Metabolism." *Nature Reviews Microbiology* 2, no. 7 (July 2004): 569–80. https://doi.org/10.1038/nrmicro926.

Edgar, Robert C. "MUSCLE: Multiple Sequence Alignment with High Accuracy and High Throughput." *Nucleic Acids Research* 32, no. 5 (2004): 1792–97. https://doi.org/10.1093/nar/gkh340.

Finn, Robert D., Jody Clements, and Sean R. Eddy. "HMMER Web Server: Interactive Sequence Similarity Searching." *Nucleic Acids Research* 39, no. suppl_2 (July 1, 2011): W29–37. https://doi.org/10.1093/nar/gkr367.

Gao, Xue, Jin-Chun Chen, Qiong Wu, and Guo-Qiang Chen. "Polyhydroxyalkanoates as a Source of Chemicals, Polymers, and Biofuels." *Current Opinion in Biotechnology*, 22/6 Chemical biotechnology and Pharmaceutical biotechnology, 22, no. 6 (December 1, 2011): 768–74. https://doi.org/10.1016/j.copbio.2011.06.005.

Gay, P, D Le Coq, M Steinmetz, E Ferrari, and J A Hoch. "Cloning Structural Gene sacB, Which Codes for Exoenzyme Levansucrase of Bacillus Subtilis: Expression of the Gene in Escherichia Coli." *Journal of Bacteriology* 153, no. 3 (March 1983): 1424–31. https://doi.org/10.1128/jb.153.3.1424-1431.1983.

Geyer, Roland, Jenna R. Jambeck, and Kara Lavender Law. "Production, Use, and Fate of All Plastics Ever Made." *Science Advances* 3, no. 7 (July 19, 2017): e1700782. https://doi.org/10.1126/sciadv.1700782.

Gilchrist, Cameron L M, and Yit-Heng Chooi. "Clinker & Clustermap.Js: Automatic Generation of Gene Cluster Comparison Figures." *Bioinformatics* 37, no. 16 (August 25, 2021): 2473–75. https://doi.org/10.1093/bioinformatics/btab007.

Henderson, Robin A., and Colin W. Jones. "Physiology of Poly-3-Hydroxybutyrate (PHB) Production by Alcaligenes Eutrophus Growing in Continuous Culture." *Microbiology* 143, no. 7 (July 1, 1997): 2361–71. https://doi.org/10.1099/00221287-143-7-2361.

Hosler, D., S. L. Burkett, and M. J. Tarkanian. "Prehistoric Polymers: Rubber Processing in Ancient Mesoamerica." *Science (New York, N.Y.)* 284, no. 5422 (June 18, 1999): 1988–91. https://doi.org/10.1126/science.284.5422.1988.

Juengert, Janina, Stephanie Bresan, and Dieter Jendrossek. "Determination of Polyhydroxybutyrate (PHB) Content in Ralstonia Eutropha Using Gas Chromatography and Nile Red Staining." *BIO-PROTOCOL* 8, no. 5 (2018). https://doi.org/10.21769/BioProtoc.2748.

Kern, Suzanne E., Alexa Price-Whelan, and Dianne K. Newman. "Extraction and Measurement of NAD(P)(+) and NAD(P)H." *Methods in Molecular Biology (Clifton, N.J.)* 1149 (2014): 311–23. https://doi.org/10.1007/978-1-4939-0473-0_26.

Lee, So Jeong, Jiwon Kim, Jung Ho Ahn, Gyeongtaek Gong, Youngsoon Um, Sun-Mi Lee, Kyoung Heon Kim, and Ja Kyong Ko. "Engineering Xylose Utilization in Cupriavidus Necator for Enhanced Poly(3-Hydroxybutyrate) Production from Mixed Sugars." *Bioresource Technology* 418 (February 2025): 131996. https://doi.org/10.1016/j.biortech.2024.131996.

Lemoigne, M. "Produit de Deshydratation et de Polymerisation de l'acide β-Oxybutyrique." *Bull. Soc. Chim. Biol.* 8 (1926): 770–82.

Letunic, Ivica, and Peer Bork. "Interactive Tree Of Life (iTOL) v5: An Online Tool for Phylogenetic Tree Display and Annotation." *Nucleic Acids Research* 49, no. W1 (July 2, 2021): W293–96. https://doi.org/10.1093/nar/gkab301.

Ling, Chen, Guan-Qing Qiao, Bo-Wen Shuai, Karel Olavarria, Jin Yin, Rui-Juan Xiang, Kun-Nan Song, Yun-Hao Shen, Yingying Guo, and Guo-Qiang Chen. "Engineering NADH/NAD+ Ratio in *Halomonas Bluephagenesis* for Enhanced Production of Polyhydroxyalkanoates (PHA)." *Metabolic Engineering* 49 (September 1, 2018): 275–86. https://doi.org/10.1016/j.ymben.2018.09.007.

Macrae, R. M., and J. F. Wilkinson. "Poly- -Hyroxybutyrate Metabolism in Washed Suspensions of Bacillus Cereus and Bacillus Megaterium." *Journal of General Microbiology* 19, no. 1 (August 1, 1958): 210–22. https://doi.org/10.1099/00221287-19-1-210.

Markowitz, Victor M., Frank Korzeniewski, Krishna Palaniappan, Ernest Szeto, Greg Werner, Anu Padki, Xueling Zhao, et al. "The Integrated Microbial Genomes (IMG) System." *Nucleic Acids Research* 34, no. suppl_1 (January 1, 2006): D344–48. https://doi.org/10.1093/nar/gkj024.

McAdam, Blaithín, Margaret Brennan Fournet, Paul McDonald, and Marija Mojicevic. "Production of Polyhydroxybutyrate (PHB) and Factors Impacting Its Chemical and Mechanical Characteristics." *Polymers* 12, no. 12 (December 4, 2020): 2908. https://doi.org/10.3390/polym12122908.

Melnyk, Ryan A., Iain C. Clark, Annette Liao, and John D. Coates. "Transposon and Deletion Mutagenesis of Genes Involved in Perchlorate Reduction in *Azospira Suillum* PS." Edited by Kenneth Nealson and Douglas G. Capone. *mBio* 5, no. 1 (February 28, 2014): e00769-13. https://doi.org/10.1128/mBio.00769-13.

Mifune, Jun, Satoshi Nakamura, and Toshiaki Fukui. "Engineering of Pha Operon on Cupriavidus
Necator Chromosome for Efficient Biosynthesis of Poly(3-Hydroxybutyrate-Co-3-
Hydroxyhexanoate) from Vegetable Oil." *Polymer Degradation and Stability* 95, no. 8 (August
2010): 1305–12. https://doi.org/10.1016/j.polymdegradstab.2010.02.026.

Minh, Bui Quang, Heiko A. Schmidt, Olga Chernomor, Dominik Schrempf, Michael D. Woodhams,
Arndt von Haeseler, and Robert Lanfear. "IQ-TREE 2: New Models and Efficient Methods for
Phylogenetic Inference in the Genomic Era." *Molecular Biology and Evolution* 37, no. 5 (May 1,
2020): 1530–34. https://doi.org/10.1093/molbev/msaa015.

Obruca, Stanislav, Petr Sedlacek, and Martin Koller. "The Underexplored Role of Diverse Stress
Factors in Microbial Biopolymer Synthesis." *Bioresource Technology* 326 (April 1, 2021): 124767.
https://doi.org/10.1016/j.biortech.2021.124767.

Oxley, Jimmie C., James L. Smith, Carolyn Higgins, Patrick Bowden, Jesse Moran, Joe Brady, Carol
E. Aziz, and Evan Cox. "Efficiency of Perchlorate Consumption in Road Flares, Propellants and
Explosives." *Journal of Environmental Management* 90, no. 11 (August 1, 2009): 3629–34.
https://doi.org/10.1016/j.jenvman.2009.06.014.

Poli, Annarita, Paola Di Donato, Gennaro Roberto Abbamondi, and Barbara Nicolaus. "Synthesis,
Production, and Biotechnological Applications of Exopolysaccharides and Polyhydroxyalkanoates by
Archaea." *Archaea* 2011 (2011): 1–13. https://doi.org/10.1155/2011/693253.

Pötter, Markus, Helena Müller, and Alexander Steinbüchel. "Influence of Homologous Phasins
(PhaP) on PHA Accumulation and Regulation of Their Expression by the Transcriptional Repressor
PhaR in Ralstonia Eutropha H16." *Microbiology* 151, no. 3 (March 1, 2005): 825–33.
https://doi.org/10.1099/mic.0.27613-0.

Redfield, A. C. (1934). On The Proportions of Organic Derivatives in Sea Water and their Relation to
the Composition of Plankton. *James Johnson Memorial Volume*, 177–19.

Reyes-Umana, Victor, Sophia D. Ewens, David A. O. Meier, and John D. Coates. "Integration of
Molecular and Computational Approaches Paints a Holistic Portrait of Obscure Metabolisms." Edited
by Vinayaka R. Prasad. *mBio* 14, no. 6 (December 19, 2023): e00431-23.
https://doi.org/10.1128/mbio.00431-23.

Rosenboom, Jan-Georg, Robert Langer, and Giovanni Traverso. "Bioplastics for a Circular
Economy." *Nature Reviews Materials* 7, no. 2 (January 20, 2022): 117–37.
https://doi.org/10.1038/s41578-021-00407-8.

Sayers, Eric W., Jeff Beck, Evan E. Bolton, J. Rodney Brister, Jessica Chan, Donald C. Comeau,
Ryan Connor, et al. "Database Resources of the National Center for Biotechnology Information."
Nucleic Acids Research 52, no. D1 (January 5, 2024): D33–43. https://doi.org/10.1093/nar/gkad1044.

Scott, Felipe, Luz Yañez, Raúl Conejeros, Blanca Araya, and Alberto Vergara-Fernández. "Two
Internal Bottlenecks Cause the Overflow Metabolism Leading to Poly(3-Hydroxybutyrate)
Production in Azohydromonas Lata DSM1123." *Journal of Environmental Chemical Engineering* 9,
no. 4 (August 2021): 105665. https://doi.org/10.1016/j.jece.2021.105665.

Steinbüchel, A., and H. G. Schlegel. "Physiology and Molecular Genetics of Poly(β-
Hydroxyalkanoic Acid) Synthesis in Alcaligenes Eutrophus." *Molecular Microbiology* 5, no. 3
(1991): 535–42. https://doi.org/10.1111/j.1365-2958.1991.tb00725.x.

Steinbüchel, Alexander, and Henry E. Valentin. "Diversity of Bacterial Polyhydroxyalkanoic Acids."
FEMS Microbiology Letters 128, no. 3 (May 1995): 219–28. https://doi.org/10.1111/j.1574-
6968.1995.tb07528.x.

Steinbüchel, Alexander. "3 Polyhydroxyalkanoic Acids" Biomaterials (1991).

Tang, Ruohao, Caihong Weng, Xiaowei Peng, and Yejun Han. "Metabolic Engineering of
Cupriavidus Necator H16 for Improved Chemoautotrophic Growth and PHB Production under
Oxygen-Limiting Conditions." *Metabolic Engineering* 61 (September 2020): 11–23.
https://doi.org/10.1016/j.ymben.2020.04.009.

Thu, Nguyen Thi Tam, Le Huy Hoang, Pham Kien Cuong, Nguyen Viet-Linh, Tran Thi Huyen Nga,
Dang Dinh Kim, Yoong Kit Leong, and Le Thi Nhi-Cong. "Evaluation of Polyhydroxyalkanoate
(PHA) Synthesis by Pichia Sp. TSLS24 Yeast Isolated in Vietnam." *Scientific Reports* 13, no. 1
(February 23, 2023): 3137. https://doi.org/10.1038/s41598-023-28220-z.

Wang, Liang, Qinghua Liu, Xiang Wu, Yue Huang, Michael J. Wise, Zhanzhong Liu, Wei Wang,
Junfeng Hu, and Chunying Wang. "Bioinformatics Analysis of Metabolism Pathways of Archaeal
Energy Reserves." *Scientific Reports* 9, no. 1 (January 31, 2019): 1034.
https://doi.org/10.1038/s41598-018-37768-0.

Wang, Liang, Jianye Yang, Yue Huang, Qinghua Liu, Yaping Xu, Xue Piao, and Michael J Wise.
"Systematic Analysis of Metabolic Pathway Distributions of Bacterial Energy Reserves." *G3
Genes|Genomes|Genetics* 9, no. 8 (August 1, 2019): 2489–96.
https://doi.org/10.1534/g3.119.400123.

Wimpenny, Julian W. T., and Anne Firth. "Levels of Nicotinamide Adenine Dinucleotide and
Reduced Nicotinamide Adenine Dinucleotide in Facultative Bacteria and the Effect of Oxygen."
Journal of Bacteriology 111, no. 1 (July 1972): 24–32. https://doi.org/10.1128/jb.111.1.24-32.1972.

Youngblut, Matthew D., Ouwei Wang, Tyler P. Barnum, and John D. Coates. "(Per)Chlorate in
Biology on Earth and Beyond." *Annual Review of Microbiology* 70, no. 1 (September 8, 2016): 435–
57. https://doi.org/10.1146/annurev-micro-102215-095406.

Zhang, Le, Zicheng Jiang, To-Hung Tsui, Kai-Chee Loh, Yanjun Dai, and Yen Wah Tong. "A
Review on Enhancing Cupriavidus Necator Fermentation for Poly(3-Hydroxybutyrate) (PHB)
Production From Low-Cost Carbon Sources." *Frontiers in Bioengineering and Biotechnology* 10
(July 19, 2022): 946085. https://doi.org/10.3389/fbioe.2022.946085.

Zheng, Jiajia, and Sangwon Suh. "Strategies to Reduce the Global Carbon Footprint of Plastics."
Nature Climate Change 9, no. 5 (May 2019): 374–78. https://doi.org/10.1038/s41558-019-0459-z.

Figure 1 | Microscopic and comparative genomic analysis of PHA production in *Dechloromonas agitata* CKB and *Azospira suillum* PS. (**A**) Transmission electron micrograph (TEM) of *D. agitata* CKB adapted from Bruce et al., 1995, showing electron-dense inclusions consistent with PHA granules. (**B**) Confocal fluorescence microscopy image of Nile Red-stained A. suillum PS overlaid with transmitted light, highlighting intracellular PHA granules. (**C**) Gene neighborhood analysis of phaC loci in CKB and PS, with neighboring PHA biosynthesis genes highlighted where present in the red box. (**D**) Growth (OD$_{600}$) and absolute PHB production (mg/mL) by *A. suillum* PS during aerobic cultivation under nitrogen replete growth conditions.

Figure 2 | PHA production by *Azospira suillum* PS across electron acceptors with acetate as the electron donor under moderate nitrogen deplete conditions. **(A)** Growth (OD_{600}) and absolute PHB production (mg/mL) during aerobic cultivation. **(B)** Growth (OD_{600}) and absolute PHB production (mg/mL) during perchlorate-respiring growth. **(C)** Growth (OD_{600}) and normalized PHB production ($mg\,mL^{-1}\,OD_{600}^{-1}$) under aerobic conditions. **(D)** Growth (OD_{600}) and normalized PHB production ($mg\,mL^{-1}\,OD_{600}^{-1}$) during perchlorate respiration.

Figure 3 | Normalized PHA production and intracellular redox state of *Azospira* suillum PS grown aerobically on acetate. **(A)** Growth (OD_{600}) and absolute intracellular concentrations of NADH and NAD^+ (μM). **(B)** Growth and absolute concentrations of NADPH and $NADP^+$ (μM). **(C)** Growth, normalized PHB production (mg mL^{-1} OD_{600}^{-1}), and the NADH/NAD^+ ratio. **(D)** Growth, normalized PHB production, and the NADPH/$NADP^+$ ratio.

580
581
582

Figure 4 | PhaC protein phylogeny within the genus *Azospira*. Maximum likelihood phylogenetic tree of PhaC protein sequences from members of the genus *Azospira*. Bootstrap support values (n = 1000 replicates) are shown at each node. Branches corresponding to *Azospira* species are highlighted in yellow, and those from A. suillum PS are further annotated in blue. Tip labels include the GenBank protein accession number followed by the corresponding NCBI organism name.

589
590
591
592

Figure 5 | Distribution of PHA-associated genes across the genus *Azospira*. **(A)** Presence and distribution of all identified PHA-associated genes across sequenced genomes within the genus *Azospira*. **(B)** Nucleotide identity percentages for the phaC biosynthetic cluster co-localized with phaB and phaR across *Azospira* species, highlighting variation in conservation of this operon.

610 **Table 1 |** Polyhydroxyalkanoate methyl esters produced by *Azospira suillum* PS when grown on
611 complex or defined media. Summary of PHA monomers identified in *A. suillum* PS when cultivated
612 on complex media (acetate, lactate, pyruvate with yeast extract) or defined acetate minimal medium.
613 PS locus tags corresponding to each *phaC* gene are listed in Supplementary Table 5.

Strains	PHB	PHV	PHHX
WT	+	-	-
ΔphaC1	+	-	-
ΔphaC2	+	-	-
ΔphaC3	+	-	-
Δ(phaC1-phaC2)	+	-	-
Δ(phaC1-phaC3)	+	-	-
Δ(phaC2-phaC3)	+	-	-
Δ(phaC1-phaC2-phaC3)	+	-	-