CycleGRN: Inferring Gene Regulatory Networks from Cyclic Flow Dynamics in Single-Cell RNA-seq

Wenjun Zhao, Elana J. Fertig, Genevieve Stein-O'Brien

November 13, 2025

Abstract

Oscillatory processes such as the cell cycle play critical roles in cell fate determination and disease development, yet existing gene regulatory network (GRN) inference methods often fail to account for their dynamic nature. We propose *CycleGRN*, a novel framework that treats cell cycle gene expression observations as an invariant measure of a stochastic differential equation and learns from data a dynamical system that fits cycling biological processes. Using a directed graph constructed along the inferred flow field in the cell space, we estimate Lie derivatives for all genes, enabling velocity inference beyond the cell cycle subspace. To quantify regulatory interactions, we introduce a time-lagged correlation operator between any pair of genes supported on the flow-aligned directed graph, which respects the intrinsic geometry of the data manifold and allows temporal ordering consistent with the underlying oscillatory process. The method requires only raw gene expression data at single-cell resolution and a list of cycle genes, without temporal binning or splicing dynamics. We evaluate our method on four synthetic datasets generated from mechanistic models with known network structures with oscillatory subnetworks, and on a mouse retinal progenitor single-cell RNA-seq dataset spanning three cell types and a knockout condition. Across all settings, our method outperforms existing methods, demonstrating its ability to recover causal and oscillatory interactions from static single-cell RNA-seq datasets.

1 Introduction

Oscillatory processes such as the cell cycle play a central role in cell fate determination, tissue homeostasis, and disease progression, including cancer and developmental disorders [16, 26, 32]. Beyond cellular phenotypes, even individual pathways, such as *Notch* signaling, can demonstrate oscillatory dynamics. These processes induce structured temporal patterns in gene expression, often reflecting periodic or quasi-periodic dynamics. While such oscillations are readily detectable at the population level, they are typically only indirectly captured in single-cell RNA-sequencing experiments, where each cell is measured at a single time point and thus requires additional methods to infer its developmental or phase state [24].

Despite their biological importance, oscillatory processes have not been fully leveraged in current gene regulatory network (GRN) inference methods. For instance, a common practice in single-cell analysis is to treat cell cycle effects as unwanted confounders [7, 20], removing or regressing out cell cycle–associated genes prior to downstream analyses such as trajectory inference. While this can reduce technical variability, it also discards valuable information about intrinsic temporal structure that could inform regulatory relationships. Numerous algorithms have been developed for recovering temporal labels from static single-cell data [19], with many focusing specifically on cell cycle dynamics [11]. For instance, Tricycle [30] identifies a template with clear cyclic structure, and uses transfer learning trained on large real datasets to map new datasets onto this template.

---

[0]Code with a simple notebook example available at https://github.com/WenjunZHAOwO/CycleGRN/tree/main

Figure 1: *Learning dynamic gene regulation with CycleGRN. (A) Input: scRNA-seq data and a list of cycle genes. (B) A continuous flow in the subspace of cycle genes is learned on this cycle manifold to reproduce the observed cell density. Each gene's directional change along the flow is estimated. (C) Comparing these directional changes across cell transitions yields time-lagged correlations that reconstruct directed regulatory interactions.*

We hypothesize that by constraining GRN inference using an estimated invariant oscillatory process, we can better recover the temporal ordering of regulatory events and improve causal inference in oscillatory systems. Intuitively, the invariant process acts as a latent clock shared across cells, capturing the periodic structure underlying gene expression variation. Empirical evidence using pseudotime learned by Tricycle [30] showed improvement in accuracy [29], but required dividing data into discrete bins, which may fail to capture the continuous and periodic nature of the cell cycle.

To address this limitation, we introduce a framework that learns an autonomous flow field representing the dynamics of a set of invariant genes (e.g., cell cycle genes) directly from single-cell data, without explicit time labels. This flow is estimated by solving an optimization problem that matches the steady-state distribution induced by the flow to the observed distribution of these genes [5, 27]. The learned flow defines a local velocity field that respects the cyclic structure of the underlying process. We then extend this invariant flow to all genes by computing cycle-aware Lie derivatives, which estimate gene-wise rates of change along the learned flow using a flow-aligned transition operator. Existing velocity methods can require spliced and unspliced reads as in RNA velocity [4], which is not always available, and can yield noisy velocity estimates depending on the hyperparameters chosen [31]. Our method also substantially differs from traditional optimal transport-based methods (e.g. [28]) which require discrete temporal binning and suffer from fundamental limitations for recurrent processes like the cell cycle, as commented by Klein et al [12].

Finally, we define a family of time-lagged gene-gene correlation operators built on the graph that follows the invariant flow of cells, enabling the estimation of directional dependencies between genes without explicit temporal binning or splicing information. This results in a signed, asymmetric gene–gene correlation matrix that reflects potential causal ordering along the oscillatory process. An illustration is provided in Figure 1. We demonstrate the performance of our method on four simulated datasets with specified oscillatory subnetworks from HARISSA [25], and three real datasets of different cell types with active cell cycles [8], showing superior performance across all tasks, compared to traditional methods that do not leverage knowledge on the oscillatory process.

2 Methodology

Given a single-cell gene expression matrix $X \in \mathbb{R}^{G \times C}$, where rows correspond to genes and columns to cells, we partition the gene set into two disjoint subsets, $G = G^{\text{inv}} \cup G^{\text{other}}$, where

- G^{inv}: genes associated with an invariant process (e.g., cell cycle or Notch signaling);

- G^{other}: all remaining genes reflecting other dynamics (e.g., differentiation or reprogramming).

Let $X^{\text{inv}} \in \mathbb{R}^{|G^{\text{inv}}| \times C}$ denote the submatrix of X restricted to G^{inv}. The ultimate goal is to use the invariant dynamics in the space of G^{inv} to constrain the temporal order of regulatory events, and infer causality that respects the cell cycle – for the full set of genes including both cell cycle genes G^{inv} and G^{other}. We describe the pipeline to achieve this in the following subsections.

2.1 Learning the dynamics of cell cycle genes

In single-cell datasets, it has been shown that sampling a population of unsynchronized cells allows the dataset to effectively capture temporal dynamics. Numerous trajectory inference methods and RNA-velocity methods have been developed to infer these dynamics [19], and shown to be successful at uncovering trajectories associated with cell fate decisions. However, these methods often rely on knowing a pre-defined initial state and are fundamentally manifold dependent, limiting the ability to infer more complex dynamics in cyclic processes that lack a unidirectional trajectory.

In the context of chaotic dynamical systems, it was first proposed by Yang et al [5, 27] that even without knowing the actual time label, one can still learn from data the dynamical system by matching the observed density and the invariant measure of a flow. Given a noisy point cloud representing the expression of the set of genes in G^{inv}, denoted as $x_c^{\text{inv}} \in \mathbb{R}^{|G^{\text{inv}}|}$ with $c = 1, ..., C$ for C cells in total, we represent the distribution of samples as a histogram over predefined bins of expression values. For each bin B, we count how many samples fall inside it and normalize by the total number of cells:

$$\rho^*(B) := \frac{1}{C} \sum_{c=1}^{C} \chi_B(x_c^{\text{inv}}), \quad \chi_B(x) = \mathbf{1}_{\{x \in B\}}. \tag{2.1}$$

We aim to construct a flow $\nu_\theta(x^{\text{inv}})$ where $\theta \in \Theta$ denotes the parameters in a simple neural network parameterizing the flow in the cell space. The optimal flow is solved through the optimization:

$$\inf_{\theta \in \Theta} \mathcal{D}(\rho_{\nu_\theta}, \rho^*), \tag{2.2}$$

where ρ^* is the density obtained from the point cloud representing cell cycle genes, ρ_{ν_θ} is the steady-state density induced by the flow ν_θ, and $\mathcal{D}$ is a metric that quantifies the discrepancy between two distributions, such as the quadratic Wasserstein distance and Kullback–Leibler divergence.

Derived from the Fokker–Planck equation of a stochastic differential equation $\mathrm{d}x_t = \nu(x_t)\mathrm{d}t + \sqrt{2D}\mathrm{d}W_t$, the steady-state solution $\rho_\nu(x)$ induced by the flow $\nu(x)$ satisfies:

$$\nabla \cdot (\rho_\nu(x)\nu(x)) = D\nabla^2 \rho_\nu(x), \tag{2.3}$$

where $\rho_\nu(x)$ represents the density that is invariant under flow $\nu(x)$, and D is a positive constant representing the scale of the diffusion. Following the approach by [5], we solve the equation using a finite volume numerical solver, compute the gradient of the objective based on the adjoint-state method and backpropagation for the parameters in $\nu(x)$, and use gradient-based optimization algorithms to iteratively update them.

Remark on the existence and non-uniqueness of optimal solution. The existence of a solution depends on the parameterization class of ν [5]. When ν is parameterized by a smooth (i.e. C^∞) neural network, the existence of a regular solution is guaranteed. To this end, we followed the recommendation and used a two-layer fully connected neural network with tanh activation throughout all experiments. However, such a system is not uniquely determined unless further information about the temporal ordering is known. For example, [6] leveraged delayed coordinates when historical values of the time series are available and proved the uniqueness via Takens' theorem [23]. In our context, we can determine the directionality by leveraging additional biological domain knowledge, such as marker genes (e.g. the Schwabe method [20]).

2.2 Cycle-aware Lie derivatives of all genes

The learned flow $\nu(x)$ is defined only on the cycle subspace and therefore must be lifted to the full G-dimensional gene space. We approximate this lifting by constructing a cell-based operator that acts on expression profiles and captures directional changes along the flow field. Conceptually, this corresponds to computing a discrete Lie derivative of each gene's expression with respect to the flow ν.

We first construct a transition matrix $L \in \mathbb{R}_+^{C \times C}$ representing a directed K-nearest neighbor graph [3, 9]. Each entry $L(i,j)$ measures how likely cell i is to transition toward cell j following the local flow direction $\nu(x_i^{\mathrm{inv}})$:

$$L_{ij} \;=\; \frac{\mathbf{1}\{\, j \in \mathcal{N}_K(i) \text{ and } \cos(x_j^{\mathrm{inv}} - x_i^{\mathrm{inv}}, \; \nu(x_i^{\mathrm{inv}})) > \delta \,\}}{\displaystyle\sum_{k \in \mathcal{N}_K(i)} \mathbf{1}\{\cos(x_k^{\mathrm{inv}} - x_i^{\mathrm{inv}}, \; \nu(x_i^{\mathrm{inv}})) > \delta\}},$$

where $\cos(a, b)$ denotes the cosine similarity, and $\delta \geq 0$ is a tolerance threshold. This construction ensures that L_{ij} is nonzero only for neighbors j whose displacement vectors are well aligned with the local flow direction $\nu(x_i^{\mathrm{inv}})$. The denominator normalizes rows of L to sum up to 1.

Given this directed adjacency matrix, the velocity of other genes can be estimated by a discrete forward difference:

$$\mathcal{L}_\nu(X) \approx V(X) \;=\; X(L^\top - I), \quad V_{g,c} = \sum_{j=1}^{C} X_{g,j} L_{cj} - X_{g,c}, \quad g = 1, ..., G, \quad c = 1, ..., C, \tag{2.4}$$

where $X \in \mathbb{R}^{G \times C}$ is the full gene-by-cell matrix. Multiplying X by $L^\top$ propagates expression values one step forward along the directed cell-cell graph, analogous to a Lie derivative along the flow field, in the operator-theoretic view [9, 18]. The term $X(L^\top - I)$ thus represents the expected forward change of each gene's expression across cells. Each element $V_{g,c}$ quantifies the rate of change of gene g in cell c.

Remark on comparison to other velocity methods. Most existing velocity methods, such as RNA velocity [4], rely on spliced and unspliced read counts, which may not be available when sequencing depth is limited. Other approaches incorporate additional information, for example, gene regulatory interactions [13], or rely on binning cells into experimental or pseudotime intervals [28]. In contrast, our method does not require splicing dynamics or explicit temporal binning. It operates solely on a predefined list of cell cycle genes to construct a flow field, and propagates this information to all genes while preserving the continuous granularity of cellular dynamics without discretizing time, under the assumption that the underlying dynamics are truly cyclic, a notoriously hard case for modern trajectory inference tools such as optimal transport (e.g., commented by Supplementary Material of Klein et al [12], "On the limitations of optimal transport for recovering cell cycles"). A comparison between flows found by optimal transport and our method can be found in Supplementary Figure S7, showing that optimal transport may still fail to identify a coherent cyclic flow even with suitable temporal binning.

2.3 Time-lagged correlation on the directed graph

The idea of approximating the manifold through a transition matrix can be extended beyond a single step to capture multi-step propagation along the velocity field ν. To propagate information along the local flow, we construct a lagged neighborhood operator built from L. Given a scalar $\alpha \in [0, 1)$, we define the propagation matrix

$$P_\alpha \;=\; \left(I - \alpha L\right)^{-1} L \;=\; \sum_{t=0}^{\infty} \alpha^t L^{t+1}, \tag{2.5}$$

where the latter expansion as a Neumann series corresponds to repeatedly applying the local flow-aligned transition matrix with exponentially decaying weights α raised to the power of t. When $\alpha = 0$, the propagation operator reduces to $P_0 = L$, which corresponds to a single-step update along the directed neighborhood graph.

Because each row of L sums to one, this is equivalent to applying one step of a Markov chain whose transition probabilities are supported on the K-nearest neighbors aligned with the local flow direction.

We then follow the approach of [28] to define a generalized notion of time-lagged correlation using the propagation matrix P_α, after normalizing each gene velocity V_g to have unit variance. For any pair of genes (g_1, g_2), we define

$$C_\alpha(g_1, g_2) \; = \; \langle V_{g_1}, P_\alpha V_{g_2} \rangle \; = \; \sum_{c_1=1}^{C} \sum_{c_2=1}^{C} V_{g_1,c_1} V_{g_2,c_2} P_{\alpha,c_1,c_2}, \tag{2.6}$$

where $V_{g,c}$ denotes the estimated velocity of gene g in cell c as in Equation (2.4). The resulting matrix $C_\alpha \in \mathbb{R}^{G \times G}$ thus quantifies the empirical correlation between gene-wise rates of change across cell pairs that are likely to be related as ancestor and descendant along the cycle.

Remark on the properties of C_α. The matrix C_α is signed: positive values indicate that two genes tend to change in the same direction along the flow, whereas negative values indicate opposing changes. Moreover, because P_α encodes time-lagged propagation, C_α is generally asymmetric, allowing one to infer potential directional relationships. Specifically, if changes in g_1 consistently precede those in g_2, then $C_\alpha(g_1, g_2)$ will differ from $C_\alpha(g_2, g_1)$, suggesting a source–target (causal) ordering between the two.

3   Results

We benchmark our algorithm on two types of datasets: (1) synthetic datasets simulated from mechanistic models [25], and (2) real mouse retinal progenitor cells undergoing cell cycle transitions during development [8] in three cell types and one knockout condition. We compare against the following representative algorithms:

- HARISSA and CARDAMOM [25]: mechanistic models for RNA count data that infer regulatory networks based on discretized temporal bins and expression matrices. CARDAMOM extends HARISSA by better exploring the energy landscape and metastable states.

- GENIE3 [1] and GRNBoost2 [14]: decision tree–based algorithms that do not use temporal information. These were ranked among the top-performing methods by BEELINE [17] for both curated and real datasets, with GRNBoost2 serving as a computationally efficient implementation of GENIE3.

- SINCERITIES [15]: a time-aware algorithm that quantifies Granger causality through a linear regression model on Kolmogorov–Smirnov statistic between consecutive time pairs.

For those methods that require time coordinates, we use either the Tricycle coordinates generated by [30], which use transfer learning to project cell cycle genes onto a common periodic template, or the simulation/experimental times when available. Performance is quantified by the area under precision curve (AUPR) and its variants.

Because the current implementation of the finite-volume solver for the Fokker–Planck equation is limited to two dimensions, in our proposed method *CycleGRN*, we learn the flow ν based on the first two principal components of cycle genes for synthetic data. For real datasets, we report results using both the first two principal components and the two-dimensional embedding produced by Tricycle [30]. Gene velocities are then computed using Equation (2.4) and normalized to unit variance across cells prior to computing time-lagged correlations. The lag parameter α in Equation (2.5) is set to 0. More details can be found in Supplementary Information Section 1.

3.1   Results on simulations

We first evaluate the performance of our method on simulated datasets where the ground truth regulatory network is known and can be directly compared with the inferred results. HARISSA [25] is a software framework

based on a mechanistic model that simulates RNA counts given any user-specified network. We used three benchmark networks: (1) FN4, a 4-gene network with a feedback loop; (2) CN5, a 5-gene network exhibiting cyclic dynamics; and (3) FN8, an 8-gene network with a feedback loop. To further mimic the complexity of real signaling pathways, we also simulated additional datasets inspired by the *Notch* pathway, including the motif between *Notch1*, *Dll1*, and *Hes1*. For each network, we generated 10 replicates with different random seeds, each containing 1,000 cells. Ground-truth networks and example simulated datasets are shown in the top two rows of Figure 2, with additional visualizations in Supplementary Figure S1.

Figure 2: *Results on oscillatory networks simulated by HARISSA [25]. Top row: ground-truth gene regulatory networks, where solid lines indicate edges associated with cycle genes used to learn the invariant flow, and edge color represents activation (blue) or repression (red). Second row: invariant flow $\nu_\theta(x)$ learned by our algorithm on the oscillatory subspace. Third row: area under the precision–recall curve (AUPR) for directed regulation. Bottom row: signed AUPR that additionally accounts for the regulation type (activation vs. inhibition). GENIE3 and GRNBoost2 are omitted because they do not predict regulation type.*

Each dataset also records the simulated time point for every cell, visualized by color in Supplementary Figure S1. To mimic experimental conditions, we intentionally introduced long lags between consecutive time points so that simulated "experimental time" does not align perfectly with the internal cell-cycle clock. These simulated times were provided to algorithms that require temporal information, including SINCERITIES [15], HARISSA, and CARDAMOM [25]. HARISSA and CARDAMOM both infer networks by directly fitting the mechanistic model used to generate the data, with CARDAMOM offering better exploration of the energy landscape, metastability, and computational efficiency. In contrast, our algorithm does not use simulation time but instead leverages prior knowledge of oscillatory genes (last row in Supplementary Figure S1). We determined the overall flow directionality based on the prior knowledge that the stimulus gene acts as an activator rather than a target. All other algorithms were only given the full count matrix without this information.

We first learn the flow fields on cycle genes and visualize them in the second row of Figure 2. The learned vector field shows smooth transitions in the space of cycle genes and aligns well with the data distribution. We set $K = 200$ and $\delta = 0.95$ to construct the transition matrix L, and infer the gene regulatory network using the time-lagged correlation operator described above.

We evaluate accuracy on the directed network using two metrics: (1) the area under the precision–recall curve (AUPR) computed on the absolute values of predicted edges, and (2) a signed AUPR that additionally requires the correct regulation type (activation or inhibition). All metrics are scaled between 0 and 1, with higher values indicating better performance. Results are shown in the bottom two rows of Figure 2; results for undirected prediction are in Supplementary Figure S2. For the two smaller networks, FN4 and CN5, our method consistently ranks top in all metrics compared to any other method. This is not surprising, since the subnetwork used as cycle genes takes a significant portion of the original true network and almost dominates the dynamics. For the two larger and more complex networks, FN8 and Notch, we show comparable or marginally better performance even compared to HARISSA and CARDAMOM, which have access to the original mechanistic model used for simulating the data. We also note that all the metrics are compared directly against the ground truth network, which is directed (distinguishes source and target genes) and signed (distinguishes activation and inhibition), demonstrating that our method can infer the temporal information even not used as input of the algorithm. This is particularly advantageous in real single-cell datasets, which typically contain only one or a few asynchronous snapshots that do not align with biological time.

3.2   Results on mouse retinal progenitor cells

We further demonstrate the performance of our algorithm on real datasets of mouse retinal progenitor cells (RPCs) [8]. The cells were subset into 27,764 early RPCs, 16,422 late RPCs, and 6,929 neurogenic cells, respectively. We ran the algorithm using 205 cell cycle genes curated from the Gene Ontology term for mouse cell cycle (GO:0007049) [2], and then infer GRNs using $3,164$ highly variable genes. We benchmarked our method against GENIE3 and GRNBoost2 (which do not use temporal information), as well as SINCERITIES with experimental time and with Tricycle positions. Due to the extensive computational cost, we randomly subsampled 1,000 cells for GENIE3 and GRNBoost2 for each cell type. Following BEELINE [17], due to low absolute accuracy, we evaluated performance using the AUPR ratio (AUPR divided by network density) and AUROC.

The cell cycle embeddings are shown in Figure 3 and Supplementary Figure S3, where the point clouds appear as amorphous blobs without clear cyclic structure. Experimental time also misaligns with the cell cycle phase revealed by Tricycle coordinates as shown in Supplementary Figure S3. Nevertheless, our method successfully recovers a dynamical system consistent with the Tricycle positions. The recovered flow fields of each cell type are shown in panel (a) of Figure 3. We then compute gene-wise velocities for all genes, where red indicates increasing and blue indicates decreasing expression. Velocities for *Top2a* are shown in panel (b) of Figure 3, peaking at the mid-cycle phase, consistent with rapid transcriptional dynamics during G2 [26], and is markedly reduced in neurogenic RPCs, reflecting their diminished proliferative activity and exit from the cell cycle.

Next, we apply the lagged correlation operator to 3,164 genes, selected as transcription factors and pattern markers, to infer directed gene–gene relationships. We benchmark the inferred networks against the STRING database [22], restricted to 80 pattern markers genes curated from non-negative matrix factorization analysis of the complete retinal dataset [8] with CoGAPS [10, 21] within each cell type performed previously on the same dataset. Only regulatory links with at least 80% confidence in STRING were retained for evaluation.

The overall performance, measured by AUPR ratio, is shown in panel (c) of Figure 3. For early RPCs, our method outperforms all others across both metrics, with a small margin over SINCERITIES using Tricycle coordinates with 10 bins. In early and late RPCs, where cell cycle activity is particularly pronounced, incorporating this temporal structure yields substantial improvements compared to methods that use either no temporal information

Figure 3: *Results on mouse retinal cells in 3 cell types: early retinal progenitor (first column), late retinal progenitor (second column), and neurogenic (third column) cells. Top row: invariant flow learned for each cell type, overlaid on the first two principal components of 205 cell cycle genes. Second row: gene velocity associated with Top2a, estimated by equation 2.4, consistent with the fact that the gene starts to peak in the middle of cycle rather than the beginning. Third row: accuracy of resulting network benchmarked against STRING, quantified by AUPR ratio (AUPR divided by density of network). Our method (pink) consistently ranks the first in RPCs, and second in neurogenic cells when cell cycle activities are weakened. Bottom row: top 10 highest weighted edges associated with Top2a, with dark blue edges indicating evidence supported by STRING. Our method correctly nominated 9, 8, and 7 for each cell types, respectively.*

or only experimental time. Even in neurogenic cells, our method reaches comparable accuracy compared to GENIE3, while all other methods performs marginally better than random baseline. We also remark that alternatively, the flow can be learned directly on the 2-dimensional tricycle embedding, yet the performance is slightly worse compared to flow learned on the cell cycle genes, possibly due to the distortion of neighbors after transfer learning is applied as in Tricycle [30]. The performance is relatively robust with the choice of k (nearest neighbor) and τ (threshold) used for constructing the transition matrix, and we have picked $k = 400$ and $\tau = 0$ for all cell types, and we performed extensive sensitivity analysis in Supplementary Figure S4.

We further examine whether the method improves predictions for cell cycle genes, as hypothesized. For the representative cell cycle gene *Top2a*, we compare the inferred network edges to those predicted by GENIE3 (Supplementary Figure S5), focusing on the top ten edges by weight. GENIE3 correctly predicts 8, 8, and 6 edges across the three cell types, whereas our method predicts 9, 8, and 7, respectively. Notably, our method identified the edge pointing from *Top2a* to *Cenpa*, a G2/M gene that peaks after *Top2a* [26], while GENIE3 missed it. This is likely because this relationship is phase-dependent and directional within the cell-cycle trajectory, and can be blurred by static networks inferred from regression-based methods such as GENIE3. This result is consistent with our hypothesis and highlights the potential of our method for improving regulatory inference in cell types exhibiting strong oscillatory signals.

The methodology is time-lagged correlation–based; therefore, it is crucial to verify whether the inferred networks capture directional causal relationships rather than static associations. To this end, we performed differential GRN inference on (1) regular late retinal progenitor cells (RPCs) and (2) *Nfia/b/x* triple-knockout RPCs, in which the transcriptional regulators *Nfia*, *Nfib*, and *Nfix* are simultaneously disrupted [8]. This knockout is known to impair the specification of late-born retinal cell types and to cause a loss of Müller glia and bipolar differentiation. The results are summarized in Figure 4.

Figure 4 presents (a) the velocity field learned from cell-cycle genes, (b) the differential incoming and outgoing hub strengths, and (c) the subnetwork formed by the 50 most rewired edges. The inferred network differences recapitulate the experimental phenotype: in the knockout, a reinforced core of cell-cycle regulators (*Hmgb2*, *Top2a*, *Ube2c*, *Cdc20*) gains mutual connectivity and forms a self-sustaining proliferative module, while genes associated with differentiation and late mitotic execution (*Cenpf*, *Cenpa*, *Ccnb1*, *Tubb4b*) lose both incoming and outgoing regulatory edges, reflecting a collapse of coordinated lineage and mitotic control.

This asymmetric rewiring, where upstream proliferative drivers persist while downstream effectors and differentiation targets become decoupled, demonstrates that the inferred network structure reflects directional causality consistent with biological perturbation. The method therefore captures not only known regulatory phenotypes but also distinguishes causal persistence (sustained drivers) from causal disconnection (lost regulation) within perturbed developmental programs.

4 Discussion

We have presented *CycleGRN*, a framework that learns an invariant flow in the space of genes corresponding to oscillatory processes such as the cell cycle and leverages it for gene regulatory network inference. The method is conceptually simple yet highly effective, and achieves superior performance on both simulated and real datasets, enhancing inference for cell cycle genes particularly.

There are several avenues for further improvement. First, the current implementation relies on a two-dimensional embedding prior to learning the dynamics. This assumption may fail in settings with substantial cell type heterogeneity, where a simple 2D projection does not reveal a clean cyclic structure. Extending the flow learning procedure to high-dimensional representations could improve robustness across diverse datasets. Second, as discussed in the methodology, the learned flow is not guaranteed to be unique. Prior work has leveraged time-delay coordinates and Takens' theorem [5, 23] to address this issue. Although true lagged measurements are not

Figure 4: *Differential GRN inference on regular late RPCs versus Nfia/b/x triple-knockout RPCs. (a) Vector field learned from cell cycle genes, (b) incoming/outgoing hub genes, and (c) subnetwork for top 50 differential edges.*

directly available in single-cell experiments, experimental time points or pseudotime estimates could potentially be used to construct suitable surrogate variables. Third, our algorithm is naturally biased toward oscillatory genes, particularly cell cycle genes, and may not perform optimally for regulatory inference involving other processes. For example, we observed suboptimal recovery of interactions involving *Hes1* (Supplementary Figure S6). An important future direction is to integrate our framework with complementary approaches that emphasize other biological processes such as differentiation.

More broadly, biological systems exhibit regulation across multiple temporal and functional scales, from fast cell-cycle oscillations to slower developmental and differentiation programs, as well as pathway-specific signaling modules such as Notch. Capturing these processes within a unified modeling framework remains a key challenge. We envision extending *CycleGRN* toward multi-scale inference, where dynamic regulatory flows corresponding to different timescales and pathways are jointly identified and integrated. Such an approach could bridge cyclic and hierarchical modes of regulation, providing a more comprehensive view of gene control across cellular states.

Overall, *CycleGRN* offers a powerful and flexible framework for leveraging oscillatory structure in single-cell data, and we expect it to enable new insights into regulatory dynamics across diverse biological systems, with potential extensions to circadian, metabolic, or developmental oscillations.

References

[1] Sara Aibar, Carmen Bravo González-Blas, Thomas Moerman, Vân Anh Huynh-Thu, Hana Imrichova, Gert Hulselmans, Florian Rambow, Jean-Christophe Marine, Pierre Geurts, Jan Aerts, Joost van den Oord, Zeynep Kalender Atak, Jasper Wouters, and Stein Aerts. Scenic: single-cell regulatory network inference and clustering. *Nature Methods*, 14(11):1083–1086, October 2017.

[2] Michael Ashburner, Catherine A. Ball, Judith A. Blake, David Botstein, Heather Butler, J. Michael Cherry, Allan P. Davis, Kara Dolinski, Selina S. Dwight, Janan T. Eppig, Midori A. Harris, David P. Hill, Laurie Issel-Tarver, Andrew Kasarskis, Suzanna Lewis, John C. Matese, Joel E. Richardson, Martin Ringwald, Gerald M. Rubin, and Gavin Sherlock. Gene ontology: tool for the unification of biology. *Nature Genetics*, 25(1):25–29, May 2000.

[3] Ralf Banisch and Péter Koltai. Understanding the geometry of transport: Diffusion maps for lagrangian trajectory data unravel coherent sets. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 27(3), February 2017.

[4] Volker Bergen, Marius Lange, Stefan Peidli, F. Alexander Wolf, and Fabian J. Theis. Generalizing rna velocity to transient cell states through dynamical modeling. *Nature Biotechnology*, 38(12):1408–1414, August 2020.

[5] Jonah Botvinick-Greenhouse, Robert Martin, and Yunan Yang. Learning dynamics on invariant measures using pde-constrained optimization. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 33(6), June 2023.

[6] Jonah Botvinick-Greenhouse, Maria Oprea, Romit Maulik, and Yunan Yang. Measure-theoretic time-delay embedding, 2024.

[7] Mengjie Chen and Xiang Zhou. Controlling for confounding effects in single cell rna sequencing studies using both control and target genes. *Scientific Reports*, 7(1), October 2017.

[8] Brian S. Clark, Genevieve L. Stein-O'Brien, Fion Shiau, Gabrielle H. Cannon, Emily Davis-Marcisak, Thomas Sherman, Clayton P. Santiago, Thanh V. Hoang, Fatemeh Rajaii, Rebecca E. James-Esposito, Richard M. Gronostajski, Elana J. Fertig, Loyal A. Goff, and Seth Blackshaw. Single-cell rna-seq analysis of retinal development identifies nfi factors as regulating mitotic exit and late-born cell specification. *Neuron*, 102(6):1111–1126.e5, June 2019.

[9] Ronald R. Coifman and Stéphane Lafon. Diffusion maps. *Applied and Computational Harmonic Analysis*, 21(1):5–30, 2006. Special Issue: Diffusion Maps and Wavelets.

[10] Elana J. Fertig, Jie Ding, Alexander V. Favorov, Giovanni Parmigiani, and Michael F. Ochs. Cogaps: an r/c++ package to identify patterns and biological process activity in transcriptomic data. *Bioinformatics*, 26(21):2792–2793, September 2010.

[11] Xinyu Guo and Liang Chen. From g1 to m: a comparative study of methods for identifying cell cycle phases. *Briefings in Bioinformatics*, 25(2), January 2024.

[12] Dominik Klein, Giovanni Palla, Marius Lange, Michal Klein, Zoe Piran, Manuel Gander, Laetitia Meng-Papaxanthos, Michael Sterr, Lama Saber, Changying Jing, Aimée Bastidas-Ponce, Perla Cota, Marta Tarquis-Medina, Shrey Parikh, Ilan Gold, Heiko Lickert, Mostafa Bakhti, Mor Nitzan, Marco Cuturi, and Fabian J. Theis. Mapping cells through time and space with moscot. *Nature*, 638(8052):1065–1075, January 2025.

[13] Jiachen Li, Xiaoyong Pan, Ye Yuan, and Hong-Bin Shen. Tfvelo: gene regulation inspired rna velocity estimation. *Nature Communications*, 15(1), February 2024.

[14] Thomas Moerman, Sara Aibar Santos, Carmen Bravo González-Blas, Jaak Simm, Yves Moreau, Jan Aerts, and Stein Aerts. Grnboost2 and arboreto: efficient and scalable inference of gene regulatory networks. *Bioinformatics*, 35(12):2159–2161, November 2018.

[15] Nan Papili Gao, S M Minhaz Ud-Dean, Olivier Gandrillon, and Rudiyanto Gunawan. Sincerities: inferring gene regulatory networks from time-stamped single cell transcriptional expression profiles. *Bioinformatics*, 34(2):258–266, September 2017.

[16] Siim Pauklin and Ludovic Vallier. The cell-cycle state of stem cells determines cell fate propensity. *Cell*, 155(1):135–147, September 2013.

[17] Aditya Pratapa, Amogh P. Jalihal, Jeffrey N. Law, Aditya Bharadwaj, and T. M. Murali. Benchmarking algorithms for gene regulatory network inference from single-cell transcriptomic data. *Nature Methods*, 17(2):147–154, January 2020.

[18] Xiaojie Qiu, Yan Zhang, Jorge D. Martin-Rufino, Chen Weng, Shayan Hosseinzadeh, Dian Yang, Angela N. Pogson, Marco Y. Hein, Kyung Hoi (Joseph) Min, Li Wang, Emanuelle I. Grody, Matthew J. Shurtleff, Ruoshi Yuan, Song Xu, Yian Ma, Joseph M. Replogle, Eric S. Lander, Spyros Darmanis, Ivet Bahar, Vijay G. Sankaran, Jianhua Xing, and Jonathan S. Weissman. Mapping transcriptomic vector fields of single cells. *Cell*, 185(4):690–711.e45, February 2022.

[19] Wouter Saelens, Robrecht Cannoodt, Helena Todorov, and Yvan Saeys. A comparison of single-cell trajectory inference methods. *Nature Biotechnology*, 37(5):547–554, April 2019.

[20] Daniel Schwabe, Sara Formichetti, Jan Philipp Junker, Martin Falcke, and Nikolaus Rajewsky. The transcriptome dynamics of single cells during the cell cycle. *Molecular Systems Biology*, 16(11), November 2020.

[21] Genevieve L Stein-O'Brien, Jacob L Carey, Wai Shing Lee, Michael Considine, Alexander V Favorov, Emily Flam, Theresa Guo, Sijia Li, Luigi Marchionni, Thomas Sherman, Shawn Sivy, Daria A Gaykalova, Ronald D McKay, Michael F Ochs, Carlo Colantuoni, and Elana J Fertig. Patternmarkers &; gwcogaps for novel data-driven biomarkers via whole transcriptome nmf. *Bioinformatics*, 33(12):1892–1894, February 2017.

[22] Damian Szklarczyk, Rebecca Kirsch, Mikaela Koutrouli, Katerina Nastou, Farrokh Mehryary, Radja Hachilif, Annika L Gable, Tao Fang, Nadezhda T Doncheva, Sampo Pyysalo, Peer Bork, Lars J Jensen, and Christian von Mering. The string database in 2023: protein–protein association networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids Research*, 51(D1):D638–D646, November 2022.

[23] Floris Takens. *Detecting strange attractors in turbulence*, page 366–381. Springer Berlin Heidelberg, 1981.

[24] Cole Trapnell, Davide Cacchiarelli, Jonna Grimsby, Prapti Pokharel, Shuqiang Li, Michael Morse, Niall J Lennon, Kenneth J Livak, Tarjei S Mikkelsen, and John L Rinn. The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nature Biotechnology*, 32(4):381–386, March 2014.

[25] Elias Ventre, Ulysse Herbach, Thibault Espinasse, Gérard Benoit, and Olivier Gandrillon. One model fits all: Combining inference and simulation of gene regulatory networks. *PLOS Computational Biology*, 19(3):e1010962, March 2023.

[26] Michael L. Whitfield, Gavin Sherlock, Alok J. Saldanha, John I. Murray, Catherine A. Ball, Karen E. Alexander, John C. Matese, Charles M. Perou, Myra M. Hurt, Patrick O. Brown, and David Botstein. Identification of genes periodically expressed in the human cell cycle and their expression in tumors. *Molecular Biology of the Cell*, 13(6):1977–2000, June 2002.

[27] Yunan Yang, Levon Nurbekyan, Elisa Negrini, Robert Martin, and Mirjeta Pasha. Optimal transport for parameter identification of chaotic dynamics via invariant measures. *SIAM Journal on Applied Dynamical Systems*, 22(1):269–310, March 2023.

[28] Wenjun Zhao, Erica Larschan, Björn Sandstede, and Ritambhara Singh. Optimal transport reveals dynamic gene regulatory networks via gene velocity estimation. *PLOS Computational Biology*, 21(5):e1012476, May 2025.

[29] Wenjun Zhao, Alma Plaza-Rodriguez, Pichayathida Luanpaisanon, Elena Xinyi Wang, Linnea Gyllingberg, Elana Fertig, and Genevieve L Stein-O'Brien. Inferring the regulation dynamics of oscillatory networks from scrna-seq data. November 2025.

[30] Shijie C. Zheng, Genevieve Stein-O'Brien, Jonathan J. Augustin, Jared Slosberg, Giovanni A. Carosso, Briana Winer, Gloria Shin, Hans T. Bjornsson, Loyal A. Goff, and Kasper D. Hansen. Universal prediction of cell-cycle position using transfer learning. *Genome Biology*, 23(1), January 2022.

[31] Shijie C. Zheng, Genevieve Stein-O'Brien, Leandros Boukas, Loyal A. Goff, and Kasper D. Hansen. Pumping the brakes on rna velocity by understanding and interpreting rna velocity estimates. *Genome Biology*, 24(1), October 2023.

[32] Binghan Zhou, Wanling Lin, Yaling Long, Yunkai Yang, Huan Zhang, Kongming Wu, and Qian Chu. Notch signaling pathway: architecture, disease, and therapeutics. *Signal Transduction and Targeted Therapy*, 7(1), March 2022.