Title

Mycobacterial surface shedding drives bystander cells response during early intracellular infection

Authors

D'Amico Davide[1], Raykov Lyudmil[1], Soldati Thierry[1,$] & Foulon Mélanie[1,$]

Affiliations

[1] Département de Biochimie, Faculté des Sciences, Université de Genève, Sciences II, 30 quai

Ernest Ansermet, CH-1211 Genève, Switzerland

[$] Co-corresponding authors: thierry.soldati@unige.ch & melanie.foulon@unige.ch

Abstract

Mycobacteria possess a complex cell envelope whose outermost layer plays an underexplored role

in host-pathogen interactions. Using *Dictyostelium discoideum* as a model host phagocyte, we

show that *Mycobacterium marinum* rapidly sheds its envelope components, including surface

proteins, carbohydrates, and virulence-associated lipids, within minutes of uptake. Shed material is

actively trafficking within host endocytic pathways and disseminates to neighboring bystander cells,

where it accumulates and triggers several responses. Notably, bystander cells exposed to shed

material exhibit a transient delay in G1/S cell cycle progression, an activation of membrane

damage-response pathways, and an enhanced resistance to subsequent mycobacterial infection.

These phenotypes are recapitulated by infection-free conditioning with purified envelope extracts,

demonstrating that superficial components of the envelope alone are sufficient to modulate host

cell responses. Moreover, this priming effect is independent of bacterial viability or the Esx-1

secretion systems, underscoring the intrinsic immunomodulatory capacity of the envelope.

Interestingly, bacteria that lose their outer layer are more frequently ubiquitinated, suggesting that

host-driven stripping exposes molecules that are recognized by cytosolic sensors to mount a cell-

autonomous defense. Together, our findings reveal that mycobacterial envelope shedding is a

widespread, early event during intracellular infection that impacts both infected and bystander

cells. These findings suggest that mycobacterial outermost envelope components can influence

host cell physiology and contribute to early innate immune modulation, with implications for

understanding the initial determinants of infection outcomes.

Keywords

Mycobacterium marinum, *Dictyostelium discoideum*, phagocytosis, cell-autonomous defense,

envelope

Introduction

Mycobacteria are a highly diverse group of bacteria that have evolved from environmental organisms to strict human pathogens through reductive evolution (Han & Silva, 2014). The most well-known pathogenic species for human include *Mycobacterium tuberculosis*, the causative agent of tuberculosis, *M. leprae*, which causes leprosy or *M. ulcerans*, responsible for Buruli ulcer. While these pathogens are characterized by their extremely slow growth, the rising emergence of non-tuberculous mycobacteria (NTM) infections, caused by faster-growing species such as *M. abscessus*, demonstrates that mycobacterial pathogenicity is not only linked to a slow metabolism. One of the distinctive features of *Mycobacterium* species is their unique cell envelope, which differs from that of Gram-positive or Gram-negative bacteria. This complex protection contributes to the bacteria's ability to survive in hostile environments (Daffé & Marrakchi, 2019). The envelope is composed of several layers, including peptidoglycan, covalently linked to arabinogalactan, which is esterified by long-chain mycolic acids. Mycolic acids form the inner leaflet of an outer membrane bilayer, contributing to the bacterial cell's impermeability and rigidity. In addition, mycobacteria possess an outermost layer, often referred to as the "capsule". This layer, primarily thought to be a specific feature of pathogenic *Mycobacterium* species (Daffé & Etienne, 1999; Fréhel, 1988), has been identified in a broader spectrum of pathogenic and non-pathogenic, slow or fast-growing mycobacteria (Sani et al., 2010). This labile layer, composed of a mixture of glycans, lipids and proteins, can vary in thickness, reaching up to 40 nm in *M. marinum* (Sani et al., 2010). Despite the recognition of its diversity in biogenesis and composition, the role of the outermost cell envelope layer in mycobacterial interactions with the host remains poorly understood.

Over the past two decades, a growing body of evidence has revealed that mycobacteria release and shed their cell wall components during host infection, contributing to both immune modulation and pathogenesis. Beatty et al. (Beatty et al., 2000; Beatty et al., 2001) were first to demonstrate that fluorescently labeled cell wall surface components (glycoconjugates and proteins) traffic from the phagosome into host endocytic compartments and are ultimately released through exocytosis. Complementary findings showed more specifically that mycobacterial lipids such as lysocardiolipin (Fischer et al., 2001) and lipoproteins, such as the 19-kDa Ag lipoprotein (Neyrolles et al., 2001), dissociate from intra-phagosomal mycobacteria and follow independent

trafficking routes within macrophages. While all these studies were performed with the attenuated

M. bovis BCG strain, it was later shown that also virulent intracellular mycobacteria, such as *M.*

marinum, can shed cell wall fragments after escape from the phagosome to the cytosol (Carlsson

et al., 2009). These foundational insights were later extended by studies showing that

mycobacteria release not only structural components but also nucleic acids, such as DNA sensed

by the cytosolic DNA sensor cyclic GMP-AMP synthase (cGAS) (Manzanillo et al., 2012; Watson et

al., 2015), and that certain cell wall lipids like PDIM can transfer between infected and uninfected

host cells (Cambier et al., 2020). Together, these findings sporadically outline an interplay between

the pathogen and the host beyond the confines of the mycobacteria-containing vacuole. But the

dynamic and mechanistic aspects of envelope components shedding remained to be elucidated.

In this context, we used *Dictyostelium discoideum* as a simple cellular model to investigate

the early stages of mycobacteria phagocytosis. As a professional phagocyte, it shares similarities

with mammalian innate immune cells in both morphology and behavior, making it a suitable tool for

investigating host-pathogen interactions, including mycobacterial infections (Cardenal-Muñoz et al.,

2018; Dunn et al., 2017; Guallar-Garrido & Soldati, 2024). In recent years, *D. discoideum* has

emerged as a powerful and experimentally versatile model host organism to decipher complex

human diseases like Alzheimer's or Parkinson's (Chen et al., 2017; Frej et al., 2017; Gilsbach et

al., 2012; Ludtmann et al., 2014), to investigate infections by several pathogens, such as

Legionella pneumophila, *Mycobacterium spp.*, *Vibrio cholera*, *Francisella noatunensis*,

Pseudomonas aeruginosa, *Salmonella enterica*, yeasts, and fungi like *Cryptococcus neoformans*

and *Aspergillus fumigatus* (reviewed in Cardenal-Muñoz et al., 2018; Dunn et al., 2017), as well as

to perform high-throughput genetic and drug screens (Hanna et al., 2020; Liao et al., 2016;

Nitschke et al., 2024; Ouertatani-Sakouhi et al., 2017).

In this study, we show that *M. marinum* outermost layer is stripped from its surface upon

internalization in *D. discoideum*, filling the phagosomal lumen. We observed that the superficial

components of the *M. marinum* envelope, including surface-exposed carbohydrates, proteins, and

virulence-associated factors are rapidly released within the first 4 hours post-infection. This passive

shedding is conserved across different mycobacterial species and occurs concurrently with

significant remodeling of the MCV. Furthermore, we demonstrate that shed material actively traffics

within the infected cell's endosomal compartment in an ESCRT-dependent manner and is

disseminated to bystander cells, where it delays the cell cycle, particularly at the G1 to S transition,

and primes them to better resist subsequent infection. These findings highlight the dynamic role of

cell wall shedding in host-pathogen interactions and indicate that mycobacterial envelope material

can influence the host cell autonomous immune response by affecting both infected and bystander

cells.

Results

The superficial layer of the *M. marinum* cell wall is shed early during phagocytosis

To investigate the dynamics of *M. marinum* cell wall during infection, surface-exposed

carbohydrates and proteins were respectively labeled with AlexaFluor-hydrazide (after Schmidt

reaction to form reactive aldehyde residues) or AlexaFluor-NHS (by ester bioconjugation to amine

groups), following procedures established in (Beatty et al., 2000) (**Figure 1A**). Labeling efficiencies

were monitored by flow-cytometry (**Suppl. Figure 1A-B**), and the viability of labeled bacilli was

verified by counting colony-forming units (CFU) (**Suppl. Figure 1C**). Early events of phagocytosis

were observed by high-resolution microscopy after spinoculating pre-labeled bacteria onto *D.*

discoideum cells. As shown in **Figure 1B**, 10 min post-contact between cells and bacteria,

AlexaFluor-488 NHS already fills the bacilli-containing compartment, indicating a fast release of

bacterial surface components into the phagosomal environment. Similar observations were made

with BV-2 microglial cells (another phagocyte model), for which the very first step of bacilli entry

was captured (**Figure 1C-D**). In those cells, in the first 40 min of phagocytosis, *M. marinum*

surface-labeled components appeared to be stripped and trafficked inside the host cell. When

observed slightly later during the infection in *D. discoideum* (2 hours post-infection (hpi), early

phase of the MCV genesis), both surface-labeled carbohydrates and proteins were found

accumulating in enclosed compartments at some distance from the MCV (**Figure 1E**).

Immunofluorescence staining of fixed infected cells at 2 hpi highlighted a co-localization of

AlexaFluor-488 hydrazide labeled shed material with an anti-PGL antibody (**Figure 2A**). Phenolic

GlycoLipid (PGL) is a mycobacterial lipid known to play a role in the virulence of several

pathogenic mycobacteria, including *M. leprae* and *M. tuberculosis*. To further confirm its shedding

from *M. marinum* cell wall during phagocytosis, we performed a metabolic pre-labeling of the bacteria (according to (Guzmán et al., 2024)) and infected *D. discoideum* cells as above. PGL labeling was verified by thin-layer chromatography (TLC), to exclude any aspecific labeling of closely related lipid species, such as PDIM (**Suppl. Figure 1D-E-F**). As for surface-exposed carbohydrates or proteins, prelabeled-PGL was released from *M. marinum* surface and accumulated in small intracellular compartments away from the MCV (**Figure 2B**). Altogether, these observations show that shed envelope material, containing virulence-associated factors, diffuses away from the bacteria and is transported from the MCV in the earliest phase of phagocytosis.

M. marinum cell wall superficial layer shedding happens in the first 6hpi

The dynamic of *M. marinum* cell wall superficial components shedding during the infection in *D. discoideum* was monitored by high-content microscopy over 12 hpi, and shedding rates were quantified accordingly (**Figure 2C**). During this time-lapse, *M. marinum* is not replicating yet as reflected by CFU counting or RFU measurement (Hagedorn & Soldati, 2007; **Suppl. Figure 1G**), ensuring no dilution of the labeled signal due to bacterial division. Cells were infected with *M. marinum* labeled for PGL (**Figure 2D-G**), surface-exposed carbohydrates (**Figure 2E-H**) or proteins (**Figure 2F-I**) and the intensity of fluorescence (corresponding to the labeled components) was measured in the extended bacterial mask (**Figure 2C**). Shedding kinetics, reflected by the loss of fluorescence intensity in the mask, was overall similar for the three types of labeling, with a first phase of drastic decrease from 1 to 4 hpi followed by a second phase showing a milder decrease up to 12 hpi (**Figure 2D-E-F**). During this first phase, a loss in intensity of approximately 35% was observed for PGL (**Figure 2D**), with a normalized rate of 0.08 RFU.h^{-1}.Similarly, a loss of intensity of around 22% (**Figure 2E**), with 0.06 RFU.h^{-1} was quantified for surface-exposed carbohydrates. The release of surface-exposed proteins occurred the fastest, with a loss of almost 70% (**Figure 2G**), with a normalized rate of 0.18 RFU.h^{-1} up to 4 hpi (**Figure 2F**). A certain heterogeneity was observed for the three types of pre-labeling at 1 hpi. This heterogeneity tended to diminish by 6 hpi (**Figure 2G-H-I**). These results show that *M. marinum* sheds its cell wall rapidly within the first 4-6 hpi. This early phase coincides with significant remodeling of the phagosomal compartment

(Cardenal-Muñoz et al., 2018; **Suppl. Figure 1H**). Therefore, cell envelope superficial shedding

may play a crucial role in the formation of the MCV, which is essential for successful infection with

M. marinum.

Cell wall shedding is conserved among mycobacteria species, but the dynamics depends

on mycobacterial intracellular compartmentalization

We showed that the shedding of superficial components of mycobacterial cell wall is conserved

among two evolutionary distant phagocyte models (**Figure 1**). We then asked whether this

phenomenon was also conserved between mycobacteria species. Several pathogenic (*M.*

kansasii, *M. abscessus* and *M. tuberculosis*) and attenuated or non-pathogenic strains (*M. bovis*

BCG, *M. smegmatis* and *M. brumae*) were assessed. Their peptidoglycan was metabolically pre-

labeled with the fluorescent precursor NADA (3-[7-nitrobenzofurazan]-carboxamide-D-Alanine) and

surface-exposed proteins were labeled with AlexaFluor-594 NHS following the procedure used for

M. marinum labeling. Those pre-labeled mycobacteria were used to infect *D. discoideum*, and

infected cells were observed by high-resolution microscopy at 2 hpi (**Figure 3A to C**). Shedding of

labeled surface-exposed proteins was observed for all species, pathogenic or not, fast or slow

growers. This shedding was specific for the outermost cell wall layer because no release of NADA-

prelabeled peptidoglycan was observed for any *Mycobacterium* species. We then assessed

whether shedding dynamics could be affected by mycobacterial virulence. To that end, shedding

rates were measured by high-content microscopy during the infection, comparing *M. marinum* WT

and ΔRD1. *M. marinum* ΔRD1 (Region of Difference 1) is an attenuated mutant that does not

assemble its Esx-1 system involved in phagosomal damage via the secretion of the membrane-

damaging toxin EsxA. Damage is essential for *M. marinum* to efficiently establish its MCV and later

escape to the cytosol. Interestingly, while the first minutes after uptake were similar (**Suppl. Figure**

2F), *M. marinum* ΔRD1 sheds its superficial prelabeled proteins and carbohydrates faster than WT,

as reflected by higher shedding rates in the first phase (**Figure 3D to F and 3G to I**). Importantly,

pre-labeling efficiency was identical for both strains and both types of labeling (**Suppl. Figure 2A-**

C for surface-exposed carbohydrate labeling, i.e. AlexaFluor-488 hydrazide, and **Suppl. Figure**

2B-D for surface-exposed proteins labeling, i.e. AlexaFluor-594 NHS) as well as their viability

(**Suppl. Figure 2E**). More specifically, bacteria unable to inflict membrane damage occupy MCV

that remain efficiently connected to endosomal membrane trafficking (Kreibich et *al.* 2015) favoring

the release and trafficking of their superficial layer. To test this hypothesis, we therefore similarly

monitored shedding rates during *M. marinum* WT infection in *D. discoideum* Δatg1, fully impaired in

autophagy, or Δalix, affected in ESCRT-mediated mechanisms. These cells are unable to

effectively repair the damage induced by *M. marinum* to the MCV membrane, leading to an early

escape of the bacilli to the cytosol (Guallar-Garrido & Soldati, 2024). For both surface-exposed

proteins (**Figure 3J to L**) or carbohydrates (**Figure 3M to O**), shedding rates were slower in *D.*

discoideum Δatg1 and Δalix cells, with the most drastic effect being observed for surface-exposed

proteins in Δalix cells with a shedding rate of 0.04 $RFU.h^{-1}$ against 0.23 $RFU.h^{-1}$ in WT cells.

Altogether, the results indicate that intracellular compartmentalization and endolysosomal

membrane trafficking drive the release dynamics of *M. marinum* surface components.

Shed mycobacterial material traffics in endosomal compartments

In mammalian macrophages, mycobacterial cell wall components were shown to traffick

throughout the endocytic compartment (Beatty et al., 2001; **Suppl. Figure 3C-D**). In *D. discoideum*

phagocytes, *M. marinum* labeled carbohydrates and proteins were observed in enclosed

compartments budding from the MCV at 2 hpi (**Figure 4A-B**). Immunofluorescence staining on

fixed cells showed that this shed material can be found in p80-positive compartments (**Figure 4C**),

a ubiquitous marker of endosomes and lysosomes, confirming the observation above. Moreover,

the use of an anti-ubiquitin antibody (FK2) showed that while shed material was not significantly

positive for ubiquitin, bacteria poorly or not anymore labeled for their surface-exposed proteins

tended to be highly ubiquitinated (**Figure 4D**). In contrast, *M. marinum* still covered with their pre-

labeled superficial layer were not ubiquitinated (**Figure 4D**). This might reflect either bacteria still

confined in their MCV, or cytosolic bacteria having lost their surface-exposed proteins normally

sensed and ubiquitinated. Together, these observations support an active dissemination of the

mycobacterial superficial layer driven by host's membrane trafficking. This removal might allow a

better access to components of the cell wall recognizable by the host defense machinery.

The shed material is actively disseminated to bystander cells

During live-microscopy recording of *D. discoideum* cells infected with AlexaFluor-594-NHS-labeled *M. marinum*, transfer of envelope material was observed from infected to non-infected cells (**Figure 5A**). This observation suggested that this material disseminates to bystander cells. To test this hypothesis quantitatively, we designed a dissemination assay (**Figure 5B**), in which an infected *D. discoideum* population (donor) expressing cytosolic mCherry (act5::mCherry) was mixed with an unmarked non-infected population (acceptor) at a 1:1 ratio. In this assay, AlexaFluor-488-NHS-labeled *M. marinum* also expressing cytosolic mCherry was used to infect the donor population. Accumulation of AlexaFluor-488-NHS in acceptor cells was measured by high-content microscopy over 6 hours post-contact (**Figure 5C**). The proportion of 488-NHS positive acceptor cells (reflected by a signal intensity at 510 nm above baseline, therefore named as proportion of positive-cells) increased over time, reaching 8% 6 hours after contact (**Figure 5D-E**), indicating net transfer of shed material without associated mCherry bacteria to non-infected cells early after contact. Another observation was made during *D. discoideum* infection with pre-labeled *M. marinum*. When looking at large fields of view, some bystanders not only received but seemed to accumulate 594-NHS labeled shed material (**Figure 6A**). To quantify the trafficking and dissemination of shed material within the host cells, the fluorescence intensity, reflecting labeled *M. marinum* shed material, was measured in the cell mask excluding the extended bacterial mask (**Figure 6B**), and normalized to the median fluorescence intensity in mock-treated cells (**Figure 6C to F**). Quantification of the accumulation of shed material was applied to the two subsets of the population: bystander and infected cells from *D. discoideum* WT (**Figure 6C**), Δatg1 (**Figure 6D**) or Δalix (**Figure 6E**) cells infected with *M. marinum* WT. For cells infected with *M. marinum* ΔRD1, mock-treated conditions were missing, therefore normalization was applied as the proportion of cells above baseline values (**Figure 6F**). Trafficking rates (i.e. fluorescence intensity in the cell mask excluding bacteria extended mask, over the time) were measured for all the conditions and cell subsets (**Figure 6G**). A fast decrease in fluorescence from 1 to 3 hpi was observed in WT cells infected with *M. marinum* WT or ΔRD1, as well as for Δatg1 cells, happening concomitantly with an increase of the signal in bystander cells. In contrast, little change was observed over the time-course in Δalix cells as reflected by reduced trafficking rates in those cells when compared to WT

cells (**Figures 6H-I**). Dissemination ratio, reflecting the transfer of shed material from infected to

bystander cells (**Figure 6J**) were compared as well. All ratio were positive except for Δalix cells,

showing a defect in transfer of *M. marinum* shed material from these cells. It is to note that similar

quantifications with 488-hydrazide labeled *M. marinum* surface components slightly differed, with

defective trafficking but similar dissemination in both Δatg1 and Δalix when compared to WT cells

(**Suppl. Figure 4**). While interesting, this difference might come from a technical bias due to the

low signal intensity of 488-hydrazide labeling and the high background in the green channel in *D.*

discoideum cells. Overall, this shows that *M. marinum* shed material, more notably proteins, which

actively traffic, disseminate to and accumulate in bystanders early during infection.

The shed envelope material delays the cell cycle of bystander cells

To investigate the impact of this accumulation of shed material in bystander cells, a microfluidic

device (Mottet et al., 2021) was used for long-term single-cell imaging of the two cellular subsets

during infection. Briefly, after infection of *D. discoideum* with GFP-expressing *M. marinum*, cells

were trapped in microchambers, and high-content microscopy was performed to monitor cell

division (**Figure 7A**). Comparisons between mock-treated, bystander and infected cells highlighted

an interdivision time prolonged approximatively two-fold in cells infected with *M. marinum* WT

(17.18 h against 8.24 h for mock cells), and 1.8-fold when infected with *M. marinum* ΔRD1 (15.33

h). Interestingly, bystanders from both infections showed, as well, a delay in their cell cycle, with an

interdivision time of 13.39 h and 13.03 h, respectively. More importantly, cells phagocytosing

antibiotic-killed *M. marinum* also showed an interdivision time prolonged by 1.5-fold, showing that

an active bacterial replication or metabolism is not required to induce cell cycle alterations in host

cells. This correlates with previous observation of cell wall shedding from dead *M. bovis* BCG in

macrophages (Rhoades et al., 2003). In *D. discoideum*, the cell cycle is divided in the standard M,

G1, S and G2 phases with a very short G1 and a prolonged G2 that lasts almost 70% of the total

(Zada-Hames & Ashworth, 1978; **Figure 7B**). To decipher which cell cycle check point is impacted

by the infection, we performed RNA-sequencing in mock-treated and infected populations at 4 hpi

and interrogated different lists of genes known to be associated to the transitions between each

phase. Compared to the strong upregulation of genes in mock-treated cells that divide normally, a

strong down-regulation of gene expression related to the G1 to S transition was observed in *M. marinum* WT infected populations, and, to a milder extent, in *M. marinum* ΔRD1 infected populations and cells in contact with dead bacilli (**Figure 7C**). This trend was also observed in genes related to the M to G1 transition (**Suppl. Figure 5A**). To better investigate these effects in the specific cell subsets, FACS-sorted subpopulations, mock-treated vs infected vs bystander cells, were analyzed at early time-points of infection by RNA-sequencing (Hanna et al., 2019). When compared to mock-treated cells, the G1 to S transition related genes were up-regulated from 1 to 3 hpi in both bystander and infected cells, while they started to be down-regulated from 6 to 12 hpi, again in both cell subsets, with a more marked trend observed in infected cells (**Figure 7D**) likely underlining a delay in G1 to S transition. Again, similar trends were observed for the M to G1 transition related genes (**Suppl. Figure 5B**). When extended to later time-points (from 24 to 48hpi), the expression of these genes was more strongly up-regulated in bystanders than mock-treated cells, or infected cells in which the situation appears "frozen" (**Suppl. Figure 5C-D).** Overall, this suggests that mycobacterial shed material impacts the *D. discoideum* cell cycle, mainly delaying its progression through the G1 to S transition during the earliest phase of the infection, but this effect is relatively transient and does not represent a full block. In addition, the RNA-seq signatures are intermediate for bystander cells compared to mock-treated and infected cells (**Suppl. Figure 6A-B**), corroborating the measured interdivision times.

To attempt at recapitulating the impact of the envelope material on bystander cell cycle progression, we designed an "infection-free" assay. The most superficial layer of the mycobacteria envelope was gently stripped from *M. marinum* grown in broth, without profoundly altering the *M. marinum* morphology and surface structure (**Figure 7E**). After washing and concentrating, this fraction was used to condition *D. discoideum* cells. Incubation of 10^5 cells with increasing concentration from 0.5, 1.25, 2.5 to 5 mg.ml^{-1} of envelope extract for 24 h, revealed a 30% decrease in growth compared to untreated cells (**Figure 7F-G**), validating the direct impact of cell wall components on host cell biology.

The shed material induces a damage-response in bystanders, priming them to better resist subsequent infection

Starting minutes after their uptake and during their intracellular life, virulent mycobacteria damage

the membrane of their compartment to establish their replicative niche and ultimately escape to the

cytosol to further grow and disseminate to naïve cells. Host cells respond to MCV damage by

recruiting the ESCRT and autophagy machineries to repair the membrane and contain bacteria

(reviewed in Guallar-Garrido & Soldati, 2024). This damage-response is also perceptible at the

transcriptomic level and is a hallmark of a mycobacteria intracellular infection. In this context, we

compared the transcriptomic profiles of the ESCRT- and autophagy-related genes, between the

two cell subsets of the infection during the first 12 hpi. A strong up-regulation was observed at 1

hpi, then progressively decreasing until 12 hpi in infected cells compared to mock-treated cells for

both ESCRT- (**Figure 8A**) and autophagy- (**Figure 8B**) related genes. An intermediate profile was

observed in bystander cells when compared to mock-treated cells, with a strong up-regulation

being mostly observed at 1 hpi, underlying a transient induction of the damage response in these

cells. Live microscopy images obtained during infection at 1.5 hpi reflected this induction. *D.*

discoideum cells expressing GFP-Vps32 (as a reporter for the ESCRT machinery) or GFP-Atg8a

(as a reporter for autophagy-mediated repair) were infected with mCherry-expressing *M. marinum*.

As shown in **Figure 8C-D**, GFP-Vps32 and GFP-Atg8a positive structures are not only seen in the

vicinity of bacteria in infected cells but accumulated as well in some bystander cells. While

heterogenous, this phenomenon shows that a damage response can be induced in non-infected

cells. To further investigate whether this is due to shed envelope material, GFP-Vps32 expressing

cells were conditioned with an extract derived from *M. marinum* and monitored by high-content

microscopy (**Figure 8E, Suppl. Figure 7A-B-C**). At 6 h post-conditioning, the median number of

reacting cells per image (i.e. cells with at least one GFP-Vps32 positive dot) went up to 40% with

1.25 mg.ml^{-1} of envelope extract (against 22% for untreated cells) (**Figure 8F**). This increased

reactivity was also reflected in a higher dot intensity (**Figure 8G**) and total area (**Figure 8H**).

Interestingly, extracts derived from *M. marinum* WT or ΔRD1 induced a similar reaction, supporting

an Esx-1-independent damage induction (**Suppl. Figure 7D**). The impact of this cellular

conditioning on the fate of the infection was finally monitored by plate-reader. Conditioned cells

were infected with *M. marinum* expressing a luminescent reporter and luminescence was

quantitated over 72 hpi, as a proxy for intracellular growth over more than one infection cycle. A

constant conditioning induced a drastic reduction of intracellular growth, resulting in a 3-fold

reduction of the intracellular mycobacterial load (**Figure 8I**). When cells were transiently pre-

conditioned for 6 h before infection, a milder intracellular growth defect was observed, with a 25%

reduction. Altogether, these results demonstrate that cell conditioning with a mycobacteria

envelope extract primes the cells to become more resistant to the infection.

Discussion

The surface of the mycobacterial cell wall is known to be labile and responsive to

environmental changes and stresses. Although shedding of surface components has been

reported in some intracellular *Mycobacterium* species, little was known about the fine dynamics of

this process and its direct consequences on host cells. In this study, using *D. discoideum* as a

simple cellular model to investigate the early stages of phagocytosis, we showed that the envelope

of *M. marinum* is rapidly stripped from the bacterial surface upon internalization, accumulating in

the phagosomal lumen. Interestingly, we and others (Beatty et al., 2002), detected the presence of

multilamellar bodies (MLBs) in the phagosomal lumen early during infection (**Suppl. Figure 1H**).

MLBs, consisting of multiple concentric layers of lipid-rich membranes, have been proposed to

reflect the presence of undigested bacterial material (Cooper et al., 1986; Hohl, 1965). Thus, the

presence of MLBs in the mycobacteria-containing vacuole likely reflects the accumulation of a

large amount of shed lipid-rich mycobacterial cell wall components.

The early removal of surface components from mycobacteria was confirmed in a

macrophage model and shown to be a conserved feature across a broad range of both pathogenic

and non-pathogenic mycobacteria. Interestingly, prior work demonstrated that heat-killed *M. bovis*

BCG can also release fluorescently labeled surface glycoconjugates (Rhoades et al., 2003). In our

model, we showed that shedding rates were similar from live or dead bacteria (**Suppl. Figure 3A-**

B). Taken together, these findings strongly suggest that the release of surface materials occurs

through a passive, mechanical process, rather than an active, bacteria-driven mechanism. In line

with this, high membrane trafficking observed shortly after mycobacteria uptake (Schnettger et al.,

2017) might be sufficient to disrupt loosely associated surface components and distribute them

throughout the connected endosomal compartment. Such mechanical stripping may represent a

host-driven strategy to expose microbial pathogen-associated molecular patterns (PAMPs), as *M.*

marinum bacilli that had lost their labeled surface components were more frequently ubiquitinated

(**Figure 4D**).

Previous studies have documented the dissemination of mycobacterial components from

infected to uninfected (bystander) cells, particularly focusing on surface-exposed glycolipids. First

studies showed that shed labeled glycoconjugates traffic through the endosomal pathway,

ultimately being released via extracellular vesicles (EVs), and later appearing in bystander cells

(Beatty et al., 2002; Beatty & Russell, 2000). Similarly, Van den Elzen et al. described a

mechanism by which bacterial glycolipids are secreted, in complex with the apolipoprotein E from

infected macrophages and efficiently taken up by dendritic cells (van den Elzen et al., 2005). Such

findings have shaped our understanding of mycobacterial lipid dissemination as a mechanism for

antigen sharing and immune modulation. However, our observations suggest more than simple

dissemination: a substantial accumulation of mycobacterial shed material occurs in bystander cells,

as reflected by increasing levels of shed NHS-labeled proteins detected in these cells (**Figure 5**).

The mechanisms by which mycobacterial surface components are released from infected

cells and reach bystander cells remained incompletely understood. Previous models have

proposed the role of extracellular vesicles, either bacterial and/or host derived (Layre, 2020).

Beatty and Russell (Beatty & Russell, 2000) speculated that mycobacterial products might be

packaged into vesicles and released extracellularly, thus allowing them to "stimulate" bystander

cells. *M. tuberculosis*-infected macrophages have been shown to release EVs containing both host

and bacterial constituents, including LAM and LM (Athman et al., 2015). Interestingly, while the

authors proposed that LAM release is dependent on mycobacteria viability, their own data show

that substantial LAM release persists when cells are infected with γ-irradiated *M. tuberculosis*,

supporting once again the passive release of mycobacteria surface components. While these

studies provide important insights, most have focused on glycolipid components and at later stages

post-infection. In contrast, our study provides the first quantitative demonstration that the release

and accumulation in bystander cells of mycobacterial surface components, including proteins,

occurs rapidly after phagocytosis initiation. This finding highlights how early, and efficiently,

bacterial material can be exported and accumulate in surrounding cells. The specific mechanisms

involved remain to be fully resolved, but our data suggest that a form of ESCRT-dependent

exocytosis is required, as Δalix host cells failed to disseminate *M. marinum* shed material. In this

model, while autophagy-dependent mechanisms seem to participate only partially to the trafficking

and transfer of *M. marinum* shed material, the ESCRT machinery might be crucial to re-capture the

cytosolic shed material in multivesicular bodies (MVB) and to regurgitate it for dissemination to

neighboring cells.

While the dissemination of mycobacterial surface components to bystanders has been

increasingly recognized, the functional consequences of this transfer, particularly at early time

points, have remained largely unexplored. Here, we show that shed material accumulates in

bystander cells within minutes of a primary infection of neighboring host cells and triggers a

transient cellular response similar to the one observed in infected cells, characterized by an

induction of membrane damage-response and a delay in the G1/S cell cycle transition. Both *M.*

tuberculosis and *M. bovis* BCG have previously been shown to induce G_0/G_1 cell cycle arrest,

apoptosis, and transcriptional reprogramming in infected cells (Cumming et al., 2017; Lai et al.,

2007). Importantly, several purified mycobacterial lipids, such as MAME, PDIM, $PIM_{1,2}$ and SL-1

are sufficient to trigger G_1/S arrest, even in the absence of live bacteria (Cumming et al., 2017).

This aligns with our observation that exposure to *M. marinum* envelope material alone, in the

absence of active infection, can induce a delay in cell cycle progression. Interestingly, *in vivo*

studies have shown that while *M. tuberculosis* infected cells undergo cell cycle arrest, bystanders

are locally proliferating in lung, contributing to tissue remodeling and immune recruitment (Huang

et al., 2018). Consistent with this, we observed an up-regulation of genes involved in G_1/S

transition in bystanders at later stages (**Suppl. Figure 4**). This suggests a dual response:

bystander cells may transiently pause their cell cycle upon sensing shed material but remain

functionally competent and potentially better equipped to control infection upon subsequent bacilli

uptake.

Several mycobacterial surface components are known to possess immunomodulatory

properties that could explain such priming effects. Notably, virulence-associated lipids such as

PDIM, PGL, and SL-1 have been individually proposed to be implicated in immune evasion. For

example, expression of *M. leprae* PGL-1 in *M. bovis* BCG promotes uptake by dendritic cells while

dampening inflammatory gene expression (Tabouret et al., 2010), while *M. tuberculosis* SL-1 has

been shown to enhance lysosomal biogenesis and exocytosis (Umar et al., 2025) . The use of

other purified lipids, including MAME, FAME, and trehalose dimycolate (TDM), can induce

phenotypes associated with lysosomal storage disorders or alter host membrane properties and

fluidity (Fineran et al., 2016; Mishra et al., 2020; Srivatsav & Kapoor, 2023). In our study, we

showed that PGL was found among the shed and disseminating material. The fact that the damage

response was observed in bystander cells, independently of the presence of the major damaging

factor EsxA, indicate that other mycobacterial surface components, such as PGL, may contribute

at high concentrations to the induction of a membrane damage response and help bystanders

sense and respond to the infection. Following this line of thoughts, bystanders, though not initially

infected, can become targets during secondary or tertiary infection cycles following the egress of

bacilli from primary infected cells. Early exposure to shed bacterial components may thus serve to

sensitize these cells, enhancing their ability to recognize and resist the infection at later stages.

Taken together, these observations suggest that mycobacterial surface components, once shed,

act not only as virulence factors within infected cells but also as modulators of bystander cell

biology. Early exposure appears to reprogram bystanders toward a more alert, resistant state, a

form of innate immune "preconditioning" that may shape the outcome of infection spread at the

population level.

Acknowledgments

We gratefully acknowledge Dr. Dimitri Moreau, Dr. Stefania Vossio and Dr. Vincent Mercier, from

the ACCESS Geneva Imaging Facility (University of Geneva) for their assistance and advice with

high-content microscopy imaging and segmentations. A special thanks to Jérôme Bosset, from the

Photonic Bioimaging Center (University of Geneva) for his help in high-resolution microscopy and

SEM images acquisition. We acknowledge Dr. Claire Mulholland and Dr. Michael Berney, from

Albert Einstein College of Medicine, for kindly providing us *M. tuberculosis* BSL-2 strains and

advising us on their cultivation protocols. We thank Céline Michard for kindly providing us the *M.*

marinum Δpks15/1 mutant, as well as the *D. discoideum* Δalix cell line. We thank Manon Mottet for

having performed the microfluidic experiments. We acknowledge the receipt of reagents and

bacteria species from BEI Resources, NIAID, NIH, including purified PGL and PDIM, as well as all

non- *M. marinum* mycobacteria species used in this work. We thank Dr. Aleksandar Salim (from Dr.

Sascha Hoogendoorn lab, University of Geneva) for the synthesis of the azido *p*HB precursor for

PGL metabolic labeling. This work was supported by Swiss National Science Foundation grants

310030_188813 and 310030_219364, and an EMBO post-doctoral fellowship earned by MF (ALTF

715-2021).

Author contributions

447 All authors made substantial contribution for this study, and approved accuracy and integrity of the

448 work presented in this manuscript. DD, MF and TS contributed equally to all conceptual aspects of

449 the project. TS acquired fundings required for this project. DD and MF both worked on all

450 experimental designs, acquisitions, analyses, and interpretations. LR participated in all

451 transcriptomic analyses. MF and TS wrote and revised the manuscript.

Material and methods

D. discoideum and BV-2 cell culture

The *D. discoideum* axenic strain Ax2(ka) was maintained in culture at 22°C in 10 cm culture dishes with HL5c medium including glucose supplemented with vitamins and microelements (ForMedium, ref HLE2), supplemented with 100 U/mL of penicillin and 100 µg/mL of streptomycin (Gibco, ref 15140-122). Plasmids were transfected into *D. discoideum* by electroporation for integration of fluorescent reporters at the safe haven act5 locus (Paschke et al., 2019) and selected with 50 µg/mL of hygromycin. Plasmids and resulting cell lines are listed in **Supplementary table S1**. BV-2 microglial cells were grown at 37°C, 5% CO_2 in DMEM high glucose (Sigma-Aldrich, ref D5796-500ML) complemented with 10% Fetal Bovine Serum (Gibco, ref 10270-106), supplemented with 100 U/mL of penicillin and 100 µg/mL of streptomycin as well.

Mycobacteria strains, plasmids and culture

All mycobacterial strains and species used in this study are listed in **Supplementary table S1**. *M. marinum* M strain was grown in Middlebrook 7H9 (BD Difco, ref 271310) enriched with 10% OADC (Oleic acid Albumin Dextrose Catalase, Becton Dickinson, ref 212351), 0.2% glycerol (Sigma-aldrich, ref G2025-1L) and 0.05% tyloxapol (Sigma-aldrich, ref T8761-50G) at 32°C, in flasks with shaking at 150 rotation per minute (rpm) and in the presence of 5 mm glass beads to minimize bacteria clumping. Other mycobacteria strains were cultured similarly, but at 37°C, except for *M. abscessus* which was grown at 32°C *M. marinum* WT or ΔRD1 expressing cytosolic GFP (pmsp12::GFP for WT, pGFPHYG2 for ΔRD1), pCherry (pMV306::pCherry10) or a bacterial luciferase (pMV306::lux) were used and selected with kanamycin 50 µg/mL or hygromycin 100 µg/mL as appropriate. The BSL-2 approved *M. tuberculosis* mc^{2}6206 strain (genotype H37Rv ΔpanCD ΔleuCD), kindly provided by Prof. Berney, was cultured in 7H9 enriched with 10% OADC, 0.2% glycerol, 0.05% tyloxapol and supplemented with 24 µg/mL pantothenate (Thermoscientific Acros, ref 24330-0050) and 50 µg/mL L-leucine (Sigma-aldrich, ref L-8912) (Mulholland et al., 2024).

Mycobacteria cell wall labeling

o **Labeling of surface-exposed carbohydrates**

Mycobacteria were cultured to reach an optical density (OD_{600}) ~0.8-1 (i.e. mid-log phase, reaching

a biomass of around 1.3 to 1.8 x 10^8 bacteria/mL). 1 x 10^9 bacteria were centrifuged at 3,000 rpm

for 10 min, at room temperature (RT). Resulting pellet was washed 3-times with Phosphate Buffer

Saline (PBS), and re-suspended with 500 µL of 0.1M sodium acetate, pH 5.5 containing 1mM

sodium periodate to oxidize hydroxyl terminal groups and form reactive aldehyde residues. After an

incubation of 20 min at 4°C with gentle rotation, 0.1mM glycerol was added to stop the reaction.

Bacteria suspension was then washed 3-times with PBS, re-suspended in 500 µL PBS containing

1 mM fluorescent-labeled hydrazide (Alexa Fluor™ 488-hydrazide, Invitrogen ref A10436) and

incubated for 1h at RT, in the dark with shaking. After 3 PBS-washes, pellet was finally re-

suspended in PBS, 7H9 or HL5c for infection.

o **Labeling of surface-exposed proteins**

Mycobacteria were cultured as for surface-exposed carbohydrates labeling. 1 x 10^8 bacteria were

centrifuged at 3,000 rpm for 10 min, at RT. Pellet was washed 3-times with PBS-containing 0.2M

sodium bicarbonate pH 8.8, re-suspended in 500 µL PBS-0.2M sodium bicarbonate containing 1

mM fluorescent succinimidyl/NHS ester (Alexa Fluor™ 488-NHS, Invitrogen ref A20000 - Alexa

Fluor™ 594-NHS, Invitrogen ref A20004) and incubated for 1h at RT, in the dark with shaking. After

3 PBS-washes, pellet was finally re-suspended in PBS, 7H9 or HL5c for infection.

o **Metabolic labeling of peptidoglycan**

Metabolic labeling of peptidoglycan was coupled with the two previous labeling protocols. Briefly,

before incubation with fluorescent dyes (hydrazide or NHS), bacterial suspensions were washed 3-

times with 7H9 and incubated for 1h with 500µM HADA (7-Hydroxycoumarin-3-carboxylic-acid-D-

Alanine, Tocris Bioscience ref 6647/5) or 25µM NADA (3-[7-nitrobenzofurazan]-carboxamide-D-

Alanine, Tocris Bioscience ref 6648/5) at 32°C in the dark with shaking. After 3 washes in PBS,

suspensions were processed for surface-exposed carbohydrates or proteins labeling.

o **Metabolic labeling of PGL**

Protocol for metabolic labeling of *M. marinum* PGL was adapted from Guzmán et al., 2024. *M.*

marinum were cultured in 7H9 containing 500µM 3-azido p-hydroxy benzoic acid (*p*HB) for 18h at

32°C with shaking, to reach an OD_{600} ~0.8-1. After incubation, 1 x 10^8 bacteria were pelleted at

3,000 rpm, 10 min RT. Resulting pellet was washed 3-times with PBS and labeled with 30µM

DBCO-AF488 (BaseClick ref BCFA-237) for 1h at RT, in the dark with shaking. Bacteria were then

washed 3-times with PBS and resulting pellet was re-suspended in PBS, 7H9 or HL5c for infection.

Flow cytometry assay

Cell wall labeling efficiency was quantitatively measured by flow cytometry. After labeling protocol,

bacteria re-suspended in PBS were adjusted to a final OD_{600} ~0.1. Samples were analyzed using a

SONY SH800 flow cytometer and data were processed with the free web-based application

Floreada.io.

Colony forming units (CFU) counting

Impact of cell wall labeling protocols on bacterial viability was evaluated by CFU counting. After

labeling protocol, bacteria re-suspended in 7H9 were serially diluted to 10^{-4} and 100µL of each

dilution was plated onto Middlebrook 7H11 plates (Difco, ref 212203). Plates were incubated at

32°C for 1 week, until colony forming.

Total lipid extraction and thin-Layer chromatography (TLC)

PGL effective labeling was controlled by analysis of total lipid extract from labeled *M. marinum*

suspensions. After labeling protocol, bacteria pellets were snap-frozen, and Bligh and Dyer

extraction was done as previously described (Bligh & Dyer, 1959). After lipid extraction, lipid

samples were re-suspended in 100µL chloroform and spotted on silica High performance TLC

plates (Merck Millipore, ref Z740222). Glycolipids were separated in a system containing

chloroform:methanol (60:40, v:v), and PDIM and neutral lipids were separated with a system

consisting of petroleum ether:diethylether (90:10, v:v). Both systems ran to 1cm below the upper

border. After drying plates under a fume hood, fluorescent lipids were visualized with UV

illumination (Fusion Fx device, Vilber Lourmat) and total lipids were detected by charring with a

solution containing 0.63g $MnCl_2.4H_2O$, 60mL water, 60mL methanol and 4mL concentrated sulfuric

acid). Lipid standards were kindly obtained from BEI Resources: purified PGL from *M. canettii* (NR-

36510) and purified PDIM from *M. tuberculosis* (NR-20328).

M. marinum envelope preparation

Protocol for envelope extraction from _M. marinum_ surface was adapted from a protocol previously developed to extract extracellular vesicles embedded in the extracellular matrix produced by _M. ulcerans_ (Foulon et al., 2022). Both fractions are labile, i.e. weakly attached to the myco-membrane, and can be removed by gentle mechanical disruption. Accordingly, _M. marinum_ was cultured in 80 mL 7H9 without beads, to reach an OD$_{600}$ ~0.8-1. Bacteria were centrifuged at 3,000 rpm for 10 min, at RT and pellet was washed 2-times with PBS to remove secreted particles not attached to the envelope/cell wall. After washing, 10 glass beads (5 mm diameter) were added to the pellet with 1 mL cold water, or HL5c, and 4 cycles of 45 sec vortexing were applied (with 1 min resting on ice in between). The suspension was then centrifuged 5 min at 3,000 rpm, RT, and supernatant was collected and filtered on 0.45µm filter units to obtain the capsular extract (approximatively 10 to 15 mg for 80 mL culture). Envelope extracts were lyophilized for long-term storage or directly used to condition _D. discoideum_ cells.

Scanning electron microscopy (SEM)

To image the bacterial surface before and after envelope extraction, 1 x 10^8 bacteria were seeded on poly-L-lysine (1 mg/mL) coated glass coverslips in a 24-well plate. After a brief centrifugation at 600 rpm, coverslips were incubated for 30 min to improve bacteria adhesion, then fixed with 4% paraformaldehyde (Fluka, ref 76240) plus 2% glutaraldehyde (Electron Microscopy Sciences, ref 16120) in PBS overnight at RT in the dark. Samples were then post-fixed with 1% osmium tetroxide (Electron Microscopy Sciences, ref 19190) in 0.1M sodium cacodylate for 1h in the dark. After rinsing 3-times with 0.1M sodium cacodylate (Electron Microscopy Sciences, ref 12310), coverslips were dehydrated in a graded ethanol series (from 30 to 100% ethanol, 5 min incubation each and repeating the incubation in ethanol 100% twice). Ethanol was them replaced with hexamethyldisilazane (HMDS, Milipore ref 8043241000) and coverslips were left to dry overnight, before to be mounted on carbon coated stubs, coated with 5 nm gold-palladium, and imaged on a JEOL SEM (JSM-6510LV) at 30 kV.

Infection assay

In *D. discoideum* cells, infections were performed as previously described (Hagedorn & Soldati. 2007), with few modifications. Mycobacteria suspensions were washed but not disrupted by syringing, to avoid envelope removal. After infection at a multiplicity of infection (MOI) of 10 for *M. marinum* WT and MOI 20 for *M. marinum* ΔRD1, cells were left for 20 min of phagocytosis, extracellular bacteria were washed off with 3 to 5 washes with 5 mL HL5c, and infected cells were re-suspended in filtered HL5c containing a bacteriostatic dose of 5 U/mL of penicillin and 5 μg/mL streptomycin, to prevent extracellular growth of *M. marinum*. Mock cells were treated similarly without adding bacteria. Time post infection was defined relative to the moment when the bacteria were added to the attached cells. For BV-2 cells, the protocol of infection was followed similarly, with slight adaptations: 5×10^{6} cells were plated in DMEM complemented with FBS without antibiotics the day before the infection, and were infected with an MOI of 5 at 32°C.

Plate reader assay

D. discoideum cells infected with *M. marinum* pMV306::lux were seeded (2×10^{5} cells per well) in a white microwell™ 96-well plate (Nunc, ref 136101) and sealed with a gas permeable moisture barrier seal (4titude, ref 4ti-0516/96). Bioluminescence intensity, measured in relative light units (RLU) as a readout for bacterial growth, was measured every 1h for 72h, at a constant temperature of 25°C using a Synergy Mx microplate reader (Biotek).

Live confocal microscopy

After infection, cells were plated in a μ-dish IBIDI 35 mm (ref 81156) for point-scan confocal high-resolution microscopy or in a 96-well IBIDI dish (ref 89626) for high-content spinning disc microscopy. For high-resolution live microscopy, images were recorded with a Leica Stellaris 8 63x1.4 NA oil immersion objective. For high-content live microscopy, images were recorded with a 60x water immersion objective mounted on the ImageXpress Micro XL HC microscope. All segmentations and quantifications were done with the MetaXpress[R] molecular device software.

Immunofluorescence staining

For immunofluorescence staining, infected *D. discoideum* cells were fixed 2hpi with ultra-cold

methanol as previously described (Hagedorn et al., 2006), to achieve optimal preservation and

permeabilization of the cells. After fixation, samples were blocked with PBS containing 0.2% gelatin

(PBG) for 1h at RT. After 3 rinsing, cells were incubated for 1h with primary antibodies directed

against p80 (Lima & Cosson. 2019) or ubiquitin FK2 (Enzo Life Sciences, ref BML-PW8810-0500)

diluted at 1:10 or 1:1000, respectively, in PBG. Cells were rinsing once again 3-times and

incubated with secondary antibodies (goat anti-rabbit or goat anti-mouse, both coupled with Cy5)

diluted at 1:1000 in PBG, mixed with DAPI, diluted 1:100, to mark nuclei. After final rinsing,

coverslips were carefully mounted on Prolong Gold Antifade Mountant (Invitrogen, ref P36934) and

left to dry overnight before imaging. Images were recorded with a Leica Stellaris 8 63x1.4 NA oil

immersion objective.

Cell conditioning

D. discoideum cells (WT or expressing GFP-Vps32) were conditioned in 96-well black plates with

transparent bottom (IBIDI, ref 89626). 1.25×10^4 (for cell multiplication assay) or 5×10^4 (for cell

damaging assay) cells were seeded per well in filtered HL5c and incubated at 22°C half an hour for

cell adhering. Medium was then replaced by envelope extract solubilized in filtered HL5c

(conditioning medium) and images were recorded every 30 min or 1h for 6-12h using a 60x water

immersion objective mounted on the ImageXpress Micro XL HC microscope.

Dissemination assay

This assay was adapted from the dissemination assay previously developed to quantitatively

monitor cell-to-cell spreading of mycobacteria (Hagedorn et *al.* 2009). Here, *D. discoideum*

expressing cytosolic act5::mCherry were infected as described above with *M. marinum* pCherry

prelabeled with 488-NHS, constituting the "donor" population. This population was then added to

D. discoideum not expressing any reporters, being the "receiver" population (already seeded in 96-

well black plate with transparent bottom (IBIDI, ref 89626) at a donor:receiver ratio of 1:1. Images

were recorded with a 60x water immersion objective mounted on the ImageXpress Micro XL HC

microscope.

Long-term single cell analyses by microfluidic

Protocol for long-term single cell imaging of *D. discoideum* was previously described in Mottet et *al.*

2021. Briefly, after infection, 3.5×10^6 cells were added on micropatterned coverslips in a 35 mm

dish, and the whole microfluidic device was carefully mounted and connected to a syringe

containing filtered HL5c medium, loaded to a syringe pump maintaining a constant flow rate of

10µL/min. Images were recorded with a 63x glycerol objective mounted on a spinning disc

confocal microscope (Intelligent Imaging Innovations Marianas SDC mounted on an inverted

microscope (Leica DMIRE2)) at 25°C.

RNA-sequencing

For RNA-sequencing performed on infected populations at 4hpi, *D. discoideum* cells infected with

M. marinum WT, ΔRD1 or dead WT (killed with 10 µM rifabutin overnight) for 4h before to be

washed and re-suspended in TRI-reagent (Sigma-aldrich, ref T9424) for RNA isolation. For cellular

subset characterizations*, D. discoideum* cells infected with *M. marinum* GFP were sorted at

specific time-points (1, 3, 6, 12, 24, 36 and 48 hpi) using an Astrios device (Beckman) at 4°C,

based on GFP-positive (infected cells) or GFP-negative (bystander cells or mock) gating

strategies. For each subset, 5×10^5 cells were collected, centrifuged and re-suspended in TRI-

reagent. RNA isolation was extracted using the Directzol RNA extraction kit (Zymo research, ref

R2053) following manufacturer's instructions. Quality of RNA librairies, sequencing and

bioinformatic analysis were performed as previously described (Hanna et al., 2019).

Statistical analyses

Data are presented as means and SD or SEM (standards deviations or standard errors of the

mean) and were analyzed with GraphPad Prism 7.0 software (GraphPad Software, San Diego, CA,

USA).

References

Athman, J. J., Wang, Y., McDonald, D. J., Boom, W. H., Harding, C. V., & Wearsch, P. A. (2015). Bacterial Membrane Vesicles Mediate the Release of Mycobacterium tuberculosis Lipoglycans and Lipoproteins from Infected Macrophages. *J Immunol*, *195*(3), 1044–1053. https://doi.org/10.4049/jimmunol.1402894

Beatty, W. L., Rhoades, E. R., Hsu, D. K., Liu, F. T., & Russell, D. G. (2002). Association of a macrophage galactoside-binding protein with Mycobacterium-containing phagosomes. *Cell Microbiol*, *4*(3), 167–176. https://doi.org/10.1046/j.1462-5822.2002.00183.x

Beatty, W. L., Rhoades, E. R., Ullrich, H. J., Chatterjee, D., Heuser, J. E., & Russell, D. G. (2000). Trafficking and release of mycobacterial lipids from infected macrophages. *Traffic*, *1*(3), 235–247. https://doi.org/10.1034/j.1600-0854.2000.010306.x

Beatty, W. L., & Russell, D. G. (2000). Identification of mycobacterial surface proteins released into subcellular compartments of infected macrophages. *Infect Immun*, *68*(12), 6997–7002. https://doi.org/10.1128/iai.68.12.6997-7002.2000

Beatty, W. L., Ullrich, H. J., & Russell, D. G. (2001). Mycobacterial surface moieties are released from infected macrophages by a constitutive exocytic event. *Eur J Cell Biol*, *80*(1), 31–40. https://doi.org/10.1078/0171-9335-00131

Bligh, E. G., & Dyer, W. J. (1959). A rapid method of total lipid extraction and purification. *Can J Biochem Physiol*, *37*(8), 911–917. https://doi.org/10.1139/o59-099

Cambier, C. J., Banik, S. M., Buonomo, J. A., & Bertozzi, C. R. (2020). Spreading of a mycobacterial cell-surface lipid into host epithelial membranes promotes infectivity. *Elife*, *9*. https://doi.org/10.7554/eLife.60648

Cardenal-Muñoz, E., Barisch, C., Lefrançois, L. H., López-Jiménez, A. T., & Soldati, T. (2018). When Dicty Met Myco, a (Not So) Romantic Story about One Amoeba and Its Intracellular Pathogen. *Front Cell Infect Microbiol*, *7*, 529. https://doi.org/10.3389/fcimb.2017.00529

Carlsson, F., Joshi, S. A., Rangell, L., & Brown, E. J. (2009). Polar localization of virulence-related Esx-1 secretion in mycobacteria. *PLoS Pathog*, *5*(1), e1000285. https://doi.org/10.1371/journal.ppat.1000285

Chen, S., Annesley, S. J., Jasim, R. A. F., Musco, V. J., Sanislav, O., & Fisher, P. R. (2017). The Parkinson's disease-associated protein DJ-1 plays a positive nonmitochondrial role in endocytosis in Dictyostelium cells. *Dis Model Mech*, *10*(10), 1261–1271. https://doi.org/10.1242/dmm.028084

Cooper, D. N. W., Haywood-Reid, P. L., Springer, W. R., & Barondes, S. H. (1986). Bacterial glycoconjugates are natural ligands for the carbohydrate binding site of discoidin I and influence its cellular compartmentalization. *Developmental Biology*, *114*(2), 416–425. https://doi.org/https://doi.org/10.1016/0012-1606(86)90206-X

Cumming, B. M., Rahman, M. A., Lamprecht, D. A., Rohde, K. H., Saini, V., Adamson, J. H., Russell, D. G., & Steyn, A. J. C. (2017). Mycobacterium tuberculosis arrests host cycle at the G1/S transition to establish long term infection. *PLoS Pathog*, *13*(5), e1006389. https://doi.org/10.1371/journal.ppat.1006389

Daffé, M., & Etienne, G. (1999). The capsule of Mycobacterium tuberculosis and its implications for pathogenicity. *Tuber Lung Dis*, *79*(3), 153–169. https://doi.org/10.1054/tuld.1998.0200

Daffé, M., & Marrakchi, H. (2019). Unraveling the Structure of the Mycobacterial Envelope. *Microbiol Spectr*, *7*(4). https://doi.org/10.1128/microbiolspec.GPP3-0027-2018

Dunn, J. D., Bosmani, C., Barisch, C., Raykov, L., Lefrançois, L. H., Cardenal-Muñoz, E., López-Jiménez, A. T., & Soldati, T. (2017). Eat Prey, Live: Dictyostelium discoideum As a Model for Cell-Autonomous Defenses. *Front Immunol*, *8*, 1906. https://doi.org/10.3389/fimmu.2017.01906

Fineran, P., Lloyd-Evans, E., Lack, N. A., Platt, N., Davis, L. C., Morgan, A. J., Höglinger, D., Tatituri, R. V. V., Clark, S., Williams, I. M., Tynan, P., Al Eisa, N., Nazarova, E., Williams, A., Galione, A., Ory, D. S., Besra, G. S., Russell, D. G., Brenner, M. B.,…Platt, F. M. (2016). Pathogenic mycobacteria achieve cellular persistence by inhibiting the Niemann-Pick Type C disease cellular pathway. *Wellcome Open Res*, *1*, 18. https://doi.org/10.12688/wellcomeopenres.10036.2

Fischer, K., Chatterjee, D., Torrelles, J., Brennan, P. J., Kaufmann, S. H., & Schaible, U. E. (2001). Mycobacterial lysocardiolipin is exported from phagosomes upon cleavage of cardiolipin by

a macrophage-derived lysosomal phospholipase A2. *J Immunol*, *167*(4), 2187–2192. https://doi.org/10.4049/jimmunol.167.4.2187

Foulon, M., Marion, E., & Marsollier, L. (2022). Purification and Characterization of Extracellular Vesicles from Mycobacterium ulcerans Culture. *Methods Mol Biol*, *2387*, 41–51. https://doi.org/10.1007/978-1-0716-1779-3_5

Fréhel, C. R., N.; Bénichou, J.C.; Ryter, A. (1988). Do test tube-grown pathogenic mycobacteria possess a protective capsule? *FEMS Microbiology Letters*, *56*(2), 225–229. https://doi.org/https://doi.org/10.1111/j.1574-6968.1988.tb03182.x

Frej, A. D., Otto, G. P., & Williams, R. S. (2017). Tipping the scales: Lessons from simple model systems on inositol imbalance in neurological disorders. *Eur J Cell Biol*, *96*(2), 154–163. https://doi.org/10.1016/j.ejcb.2017.01.007

Gilsbach, B. K., Ho, F. Y., Vetter, I. R., van Haastert, P. J., Wittinghofer, A., & Kortholt, A. (2012). Roco kinase structures give insights into the mechanism of Parkinson disease-related leucine-rich-repeat kinase 2 mutations. *Proc Natl Acad Sci U S A*, *109*(26), 10322–10327. https://doi.org/10.1073/pnas.1203223109

Guallar-Garrido, S., & Soldati, T. (2024). Exploring host-pathogen interactions in the Dictyostelium discoideum-Mycobacterium marinum infection model of tuberculosis. *Dis Model Mech*, *17*(7). https://doi.org/10.1242/dmm.050698

Guzmán, L. E., Cambier, C. J., Cheng, T. Y., Naqvi, K. F., Shiloh, M. U., Moody, D. B., & Bertozzi, C. R. (2024). Bioorthogonal Metabolic Labeling of the Virulence Factor Phenolic Glycolipid in Mycobacteria. *ACS Chem Biol*, *19*(3), 707–717. https://doi.org/10.1021/acschembio.3c00724

Hagedorn, M., Neuhaus, E. M., & Soldati, T. (2006). Optimized fixation and immunofluorescence staining methods for Dictyostelium cells. *Methods Mol Biol*, *346*, 327–338. https://doi.org/10.1385/1-59745-144-4:327

Hagedorn, M., & Soldati, T. (2007). Flotillin and RacH modulate the intracellular immunity of Dictyostelium to Mycobacterium marinum infection. *Cell Microbiol*, *9*(11), 2716–2733. https://doi.org/10.1111/j.1462-5822.2007.00993.x

Han, X. Y., & Silva, F. J. (2014). On the age of leprosy. *PLoS Negl Trop Dis*, *8*(2), e2544.
https://doi.org/10.1371/journal.pntd.0002544

Hanna, N., Burdet, F., Melotti, A., Bosmani, C., Kicka, S., Hilbi, H., Cosson, P., Pagni, M., & Soldati,
T. (2019). Time-resolved RNA-seq profiling of the infection of Dictyostelium
discoideum by Mycobacterium marinum reveals an integrated host
response to damage and stress. *bioRxiv*, 590810. https://doi.org/10.1101/590810

Hanna, N., Kicka, S., Chiriano, G., Harrison, C., Sakouhi, H. O., Trofimov, V., Kranjc, A., Nitschke,
J., Pagni, M., Cosson, P., Hilbi, H., Scapozza, L., & Soldati, T. (2020). Identification of Anti-
Mycobacterium and Anti-Legionella Compounds With Potential Distinctive Structural
Scaffolds From an HD-PBL Using Phenotypic Screens in Amoebae Host Models. *Front
Microbiol*, *11*, 266. https://doi.org/10.3389/fmicb.2020.00266

Hohl, H. R. (1965). Nature and Development of Membrane Systems in Food Vacuoles of Cellular
Slime   Molds   Predatory   upon   Bacteria.   *J   Bacteriol*,   *90*(3),   755–765.
https://doi.org/10.1128/jb.90.3.755-765.1965

Huang, L., Nazarova, E. V., Tan, S., Liu, Y., & Russell, D. G. (2018). Growth of Mycobacterium
tuberculosis in vivo segregates with host macrophage metabolism and ontogeny. *J Exp
Med*, *215*(4), 1135–1152. https://doi.org/10.1084/jem.20172020

Lai, Y. M., Mohammed, K. A., Nasreen, N., Baumuratov, A., Bellew, B. F., & Antony, V. B. (2007).
Induction of cell cycle arrest and apoptosis by BCG infection in cultured human bronchial
airway   epithelial   cells.   *Am   J   Physiol   Lung   Cell   Mol   Physiol*,   *293*(2),   L393–401.
https://doi.org/10.1152/ajplung.00392.2006

Layre, E. (2020). Trafficking of Mycobacterium tuberculosis Envelope Components and Release
Within Extracellular Vesicles: Host-Pathogen Interactions Beyond the Wall. *Front Immunol*,
11, 1230. https://doi.org/10.3389/fimmu.2020.01230

Liao, X. H., Meena, N. P., Southall, N., Liu, L., Swaroop, M., Zhang, A. L., Xiang, J. J., Parent, C.
A., Zheng, W., & Kimmel, A. R. (2016). A High-Throughput, Multi-Cell Phenotype Assay for
the Identification of Novel Inhibitors of Chemotaxis/Migration. *Sci Rep*, *6*, 22273.
https://doi.org/10.1038/srep22273

Ludtmann, M. H., Otto, G. P., Schilde, C., Chen, Z. H., Allan, C. Y., Brace, S., Beesley, P. W.,

Kimmel, A. R., Fisher, P., Killick, R., & Williams, R. S. (2014). An ancestral non-proteolytic

role for presenilin proteins in multicellular development of the social amoeba Dictyostelium

discoideum. *J Cell Sci*, *127*(Pt 7), 1576–1584. https://doi.org/10.1242/jcs.140939

Manzanillo, P. S., Shiloh, M. U., Portnoy, D. A., & Cox, J. S. (2012). Mycobacterium tuberculosis

activates the DNA-dependent cytosolic surveillance pathway within macrophages. *Cell Host*

Microbe, *11*(5), 469–480. https://doi.org/10.1016/j.chom.2012.03.007

Mishra, M., Dadhich, R., Mogha, P., & Kapoor, S. (2020). Mycobacterium Lipids Modulate Host Cell

Membrane Mechanics, Lipid Diffusivity, and Cytoskeleton in a Virulence-Selective Manner.

ACS Infect Dis, *6*(9), 2386–2399. https://doi.org/10.1021/acsinfecdis.0c00128

Mottet, M., Bosmani, C., Hanna, N., Nitschke, J., Lefrançois, L. H., & Soldati, T. (2021). Novel

Single-Cell and High-Throughput Microscopy Techniques to Monitor Dictyostelium

discoideum-Mycobacterium marinum Infection Dynamics. *Methods Mol Biol*, *2314*, 183–

203. https://doi.org/10.1007/978-1-0716-1460-0_7

Mulholland, C. V., Wiggins, T. J., Cui, J., Vilchèze, C., Rajagopalan, S., Shultis, M. W., Reyes-

Fernández, E. Z., Jacobs, W. R., Jr., & Berney, M. (2024). Propionate prevents loss of the

PDIM virulence lipid in Mycobacterium tuberculosis. *Nat Microbiol*, *9*(6), 1607–1618.

https://doi.org/10.1038/s41564-024-01697-8

Neyrolles, O., Gould, K., Gares, M. P., Brett, S., Janssen, R., O'Gaora, P., Herrmann, J. L.,

Prévost, M. C., Perret, E., Thole, J. E., & Young, D. (2001). Lipoprotein access to MHC

class I presentation during infection of murine macrophages with live mycobacteria. *J*

Immunol, *166*(1), 447–457. https://doi.org/10.4049/jimmunol.166.1.447

Nitschke, J., Huber, R., Vossio, S., Moreau, D., Marcourt, L., Gindro, K., Queiroz, E. F., Soldati, T.,

& Hanna, N. (2024). Discovery of anti-infective compounds against Mycobacterium

marinum after biotransformation of simple natural stilbenes by a fungal secretome. *Front*

Microbiol, *15*, 1439814. https://doi.org/10.3389/fmicb.2024.1439814

Ouertatani-Sakouhi, H., Kicka, S., Chiriano, G., Harrison, C. F., Hilbi, H., Scapozza, L., Soldati, T.,

& Cosson, P. (2017). Inhibitors of Mycobacterium marinum virulence identified in a

Dictyostelium discoideum host model. *PLoS One*, *12*(7), e0181121. https://doi.org/10.1371/journal.pone.0181121

Paschke, P., Knecht, D. A., Williams, T. D., Thomason, P. A., Insall, R. H., Chubb, J. R., Kay, R. R., & Veltman, D. M. (2019). Genetic Engineering of Dictyostelium discoideum Cells Based on Selection and Growth on Bacteria. *J Vis Exp*(143). https://doi.org/10.3791/58981

Rhoades, E., Hsu, F., Torrelles, J. B., Turk, J., Chatterjee, D., & Russell, D. G. (2003). Identification and macrophage-activating activity of glycolipids released from intracellular Mycobacterium bovis BCG. *Mol Microbiol*, *48*(4), 875–888. https://doi.org/10.1046/j.1365-2958.2003.03473.x

Sani, M., Houben, E. N., Geurtsen, J., Pierson, J., de Punder, K., van Zon, M., Wever, B., Piersma, S. R., Jiménez, C. R., Daffé, M., Appelmelk, B. J., Bitter, W., van der Wel, N., & Peters, P. J. (2010). Direct visualization by cryo-EM of the mycobacterial capsular layer: a labile structure containing ESX-1-secreted proteins. *PLoS Pathog*, *6*(3), e1000794. https://doi.org/10.1371/journal.ppat.1000794

Schnettger, L., Rodgers, A., Repnik, U., Lai, R. P., Pei, G., Verdoes, M., Wilkinson, R. J., Young, D. B., & Gutierrez, M. G. (2017). A Rab20-Dependent Membrane Trafficking Pathway Controls M. tuberculosis Replication by Regulating Phagosome Spaciousness and Integrity. *Cell Host Microbe*, *21*(5), 619–628.e615. https://doi.org/10.1016/j.chom.2017.04.004

Srivatsav, A. T., & Kapoor, S. (2023). Biophysical Interaction Landscape of Mycobacterial Mycolic Acids and Phenolic Glycolipids with Host Macrophage Membranes. *ACS Appl Bio Mater*, *6*(12), 5555–5562. https://doi.org/10.1021/acsabm.3c00748

Tabouret, G., Astarie-Dequeker, C., Demangel, C., Malaga, W., Constant, P., Ray, A., Honoré, N., Bello, N. F., Perez, E., Daffé, M., & Guilhot, C. (2010). Mycobacterium leprae phenolglycolipid-1 expressed by engineered M. bovis BCG modulates early interaction with human phagocytes. *PLoS Pathog*, *6*(10), e1001159. https://doi.org/10.1371/journal.ppat.1001159

Umar, I., Gulzar, S. E., & Sundaramurthy, V. (2025). M. tuberculosis surface sulfoglycolipid SL-1 activates the mechanosensitive channel TRPV4 to enhance lysosomal biogenesis and

exocytosis in macrophages. *Mol Biol Cell*, *36*(6), ar76. https://doi.org/10.1091/mbc.E24-12-

0560

van den Elzen, P., Garg, S., León, L., Brigl, M., Leadbetter, E. A., Gumperz, J. E., Dascher, C. C.,

Cheng, T. Y., Sacks, F. M., Illarionov, P. A., Besra, G. S., Kent, S. C., Moody, D. B., &

Brenner, M. B. (2005). Apolipoprotein-mediated pathways of lipid antigen presentation.

Nature, *437*(7060), 906–910. https://doi.org/10.1038/nature04001

Watson, R. O., Bell, S. L., MacDuff, D. A., Kimmey, J. M., Diner, E. J., Olivas, J., Vance, R. E.,

Stallings, C. L., Virgin, H. W., & Cox, J. S. (2015). The Cytosolic Sensor cGAS Detects

Mycobacterium tuberculosis DNA to Induce Type I Interferons and Activate Autophagy. *Cell*

Host Microbe, *17*(6), 811–819. https://doi.org/10.1016/j.chom.2015.05.004

Zada-Hames, I. M., & Ashworth, J. M. (1978). The cell cycle during the vegetative stage of

Dictyostelium discoideum and its response to temperature change. *J Cell Sci*, *32*, 1–20.

https://doi.org/10.1242/jcs.32.1.1

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Time →

Infected cell   Bystander cell

Mm-Containing Vacuole (MCV)

ESCRT machinery

Re-capture in MVB

Mm envelope

M. marinum (*Mm*)

Phagophore

Membrane trafficking

Re-capture by autophagy

Sense and respond

ESCRT-dependent process
ESCRT and autophagy-dependent process

Mm ΔRD1 infection = no MCV damage
↗↗ ENVELOPE RELEASE ↗↗

Infection in autophagy-deficient cells
↓↓ ENVELOPE RELEASE ↓↓

Infection in ESCRT-deficient cells
↓↓ ENVELOPE RELEASE ↓↓
+
DEFECTIVE DISSEMINATION

Bystander cell