Sequence context and methylation interact to shape germline mutation rate variation at CpG sites

Sheel Chandra[1], Ziyue Gao[2*]

[1] Department of Biology, University of Pennsylvania, Philadelphia, Pennsylvania, United States of America

[2] Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, United States of America

* Corresponding author

E-mail: ziyuegao@pennmedicine.upenn.edu

Abstract

A prominent example of sequence context-dependent mutation rate variation is the elevated transition rate at CpG sites, which is largely attributed to cytosine methylation. CpGs with different flanking sequences also exhibit mutation rate variation, but this variation is only partially correlated with context-specific methylation level. Here, we quantify the CpG mutation rate and mutagenic effect of methylation across sequence contexts. Using a regression framework that accounts for recurrent mutations, we analyze human polymorphisms from the gnomAD dataset to estimate mutation rates of unmethylated and methylated CpGs separately in each unique 4-mer or 6-mer context. We find that CpG mutation rate variation in the human genome is shaped by methylation at the focal cytosine, the flanking nucleotides, and interactions between them, suggesting distinct context-dependent mutation patterns for unmethylated and methylated cytosines. Our analysis further reveals that the context effects are driven by largely independent effects of upstream and downstream sequences. Notably, an upstream adenine markedly increases CpG mutation rates regardless of methylation status or downstream sequences. Furthermore, upstream and downstream sequences have similar effects in chimpanzee and rhesus macaque, indicating that some conserved, intrinsic sequence features shape CpG mutability. On the other hand, some inter-species differences, which are especially pronounced at methylated sites on the chimpanzee lineage, point to recent evolutionary changes, possibly in context-specificity of proteins governing DNA demethylation and repair processes.

Author Summary

The DNA sequence surrounding a nucleotide strongly influences how likely it is to mutate. An extreme example is the CpG dinucleotide: cytosines in CpGs mutate far more frequently than other sites in the human genome. This is related to DNA methylation, a chemical modification that occurs almost exclusively at CpGs in vertebrates and makes cytosines more prone to mutations. However, CpGs in different sequence contexts also vary in their mutation rates, and methylation level alone cannot explain this variation. To gain insight into what processes drive this variation, we estimate mutation rates for methylated and unmethylated CpGs in different sequence contexts using human genetic variation data. We find that methylation and neighboring bases interact to influence CpG mutation rates, and that the DNA sequence on either side of the CpG exerts largely independent effects. Extending our analysis to other primates reveals both conserved and species-specific patterns, with differences being especially pronounced at methylated sites on the chimpanzee lineage. Together, our results suggest that while intrinsic DNA sequence features underlie some conserved context effects on CpG mutation rate, inter-species differences may reflect recent evolutionary changes in the mechanisms that regulate DNA demethylation and repair.

Introduction

Mutation rate varies across the human genome, even at the scale of individual nucleotides [1]. Quantifying and accounting for this mutation rate variation is important for making inferences about natural selection [2] and identifying functional coding [3–6] and non-coding regions [7]. Among the many predictors of mutation rate, one of the strongest is the local sequence context—the nucleotides flanking the mutated site [8,9]. Although context effects have been attributed to several mechanisms, including local DNA shape [10,11], biases in DNA polymerase fidelity [12], and differential repair efficiency across sequence environments [13,14], the relative importance of these mechanisms and whether they fully account for observed patterns remain open questions. Addressing these questions requires systematically characterizing how local sequence shapes site-level mutation rate variation, which can help disentangle these contributions and shed light on the mechanistic basis of mutagenesis.

Among sequence contexts, the CpG dinucleotide is both exceptionally mutable and mechanistically informative. Across vertebrates, cytosines at CpG sites undergo C>T transition at rates an order of magnitude higher than those at other genomic sites [1,15,16]. This hypermutability is directly linked to DNA methylation, as supported by two observations: first, in vertebrates, cytosine methylation occurs almost exclusively at CpG sites [17], and second, in humans, de novo C>T mutation rates at CpGs correlate strongly with germline methylation levels [18–20].

The mutagenic effect of methylation is commonly attributed to the spontaneous deamination of 5-methylcytosine (5mC). This process is distinguished from the deamination of unmethylated cytosine in two critical ways: an accelerated reaction rate [21] and a distinct chemical product. Specifically, while deamination of unmethylated cytosine yields uracil—a non-canonical base

efficiently recognized and removed by DNA repair machinery [22]—deamination of 5mC

produces thymine. The resulting T:G mismatch can then be converted to a T:A base pair if the

mismatch is mis-repaired or left unrepaired until DNA replication [23].

However, not all C>T mutations at CpG sites arise from faulty/inefficient repair following

deamination; replication-associated errors independent of deamination likely also contribute

[12,24,25]. For example, analysis of de novo CpG>TpG mutations reveals strong asymmetry

with regard to replication direction, showing up to 33% higher mutation rate on the lagging

strand [24]. In addition, recent work on human DNA polymerase ε suggests that this enzyme

exhibits a bias toward mis-incorporating adenine opposite 5mC and even unmethylated cytosine

in a CpG context [12,25].

CpG mutation rates vary substantially across sequence contexts [8,26]. Aggarwala and Voight

[8] reported that although sperm methylation levels correlate significantly with CpG>TpG

polymorphism rates in 7-mer contexts, the correlation is modest ($R^2 = 0.33$), suggesting that

variation in methylation level alone cannot fully explain CpG mutation rate variation across

sequence contexts. A key question, then, is how the local sequence environment influences

these rates. Potential mechanisms include intrinsic DNA conformational features that shape the

rate of spontaneous deamination [27], the interplay between local sequence and repair

enzymes in resolving T:G mismatches [14,28,29], or sequence-dependent polymerase error

profiles [12,25]. In addition, context-specific DNA-binding proteins such as transcription factors

can also modulate mutation rate by interfering with repair machinery [30–32]. Because 5mC and

unmethylated cytosines possess distinct biochemical properties, quantifying their mutational

profiles separately can help distinguish between these competing mechanisms. This

necessitates a framework capable of decoupling the influence of methylation from sequence

context effects, allowing for the estimation of mutation patterns at both methylated and

unmethylated CpGs.

Here, we develop a modeling framework to investigate the biological determinants of CpG

mutability by explicitly considering both methylation state and sequence context at each CpG

site. We build on existing context-dependent mutation models by incorporating methylation as a

continuous predictor, enabling us to disentangle the contributions of methylation and local

sequence context while leveraging information from CpG sites across varying methylation

levels. Applying this approach to human polymorphism data from the Genome Aggregation

Database (gnomAD v4.0), we estimate mutation rates for both unmethylated and methylated

CpGs in all unique 4-mer and 6-mer contexts. We also extend our analysis to two additional

primate species to identify conserved and divergent patterns in CpG mutation rate variation.

Overall, our results demonstrate that unmethylated and methylated cytosines represent

fundamentally different mutational substrates with distinct context dependencies that together

shape CpG mutation landscape across the genome.

117   Results

118   Interaction between methyl group and flanking nucleotides on CpG

119   mutation rate

We hypothesized that the variation in CpG mutation rates across contexts is shaped by both

differences in baseline mutation rates of unmethylated cytosines and differential mutagenic

effects of methylation (i.e. the difference in mutation rate between methylated and unmethylated

cytosines). We started by calculating the CpG transition polymorphism rate for each

tetranucleotide (4-mer) context at varying methylation levels, using single nucleotide

polymorphisms (SNPs) as a proxy for mutation events in the ancestors of the sampled

genomes. For primary analysis, we used SNPs in the intergenic regions of the human genome

from gnomAD [36], and polarized SNPs based on minor allele frequency (Methods); SNPs from

the 1000 Genomes (1KG) Project were used for replication [37].

To approximate methylation levels in the male germline, we used bisulfite sequencing from

human sperm [38]. We recognize that germline mutations can arise in both the male and female

germlines, as well as during early embryonic development, so the most relevant methylation

metric should be a weighted average of methylation levels across all these stages. However, we

chose to use sperm bisulfite sequencing data as a proxy for germline methylation level for two

reasons. First, the site-level methylation intensity in sperm shows the strongest correlation with

CpG mutation rate and has much greater predictive power than data from any other

developmental stages [39]. Second, whole-genome bisulfite sequencing data for tissues other

than testis and/or sperm are much more limited and typically not corrected for genotypes of the

assayed individuals, which creates biases in methylation level measurement (see Methods).

As expected, the polymorphism rate increases with germline methylation level for all CpG sites

in aggregation and within each 4-mer context (Fig 1A; S1A Fig). However, this increase is

approximately linear only up to ~25% methylation, after which the curve becomes flatter. This

non-linear relationship between polymorphism rate and methylation is likely driven by mutation

saturation of highly mutable sites, where recurrent mutations become frequent in large cohorts

[18,40]. This observation is concordant with prior studies that, at the current sample size of

human genetic variation datasets, the observed polymorphism probability of CpG sites no

longer scales proportionately with the underlying mutation rate [18,33,41,42].

Fig 1. Interaction between sequence context and methylation shapes mutation rate variation at CpG sites. A. Polymorphism rate of CpG sites as a function of methylation levels. CpG sites were partitioned into 20 equal-width bins (ranging from [0, 0.05) to [0.95, 1]), with the polymorphism rate calculated as the proportion of sites harboring SNPs within each bin. Error bars represent 95% confidence intervals assuming binomial sampling. **B.** Proposed mutation rate model where the mutation rate for a given cytosine in a CpG within a 4-mer sequence context is a linear function of its methylation level. is the context-specific baseline rate when methylation is zero, and captures the increase in mutation rate from unmethylated to fully methylated. **C.** Schematic showing how mutation rates are related with the observed methylation level and polymorphism state at every site . **D.** Scaled mutation rate estimates for unmethylated and methylated CpGs in each 4-mer context. Contexts are sorted by descending mutation rate when unmethylated. Error bars represent 95% confidence intervals for the estimates. **E.** Comparison between scaled mutation rates of unmethylated and methylated CpGs estimated from the gnomAD dataset by our model and estimated *de novo* mutation (DNM) rates for lowly methylated (<10% methylation in sperm) and highly methylated sites (>90% methylation), respectively. 95% confidence intervals for DNM rates are estimated assuming binomial sampling.

To account for recurrent mutations, we modeled the relationship between the polymorphism

probability *p* and the underlying per-generation mutation rate *μ* using an exponential

transformation $p = 1 - e^{-\mu T}$, where T represents the total branch length of the coalescent tree

connecting all sampled genomes. This approach adjusts for recurrent mutations to a first

approximation and produces a sample-scaled mutation rate, μT. We treated the observed

polymorphism state (i.e., presence or absence of polymorphism) at each site as a Bernoulli trial

with success probability p (Fig 1C). To model the effects of sequence context, methylation, and

their interaction, we assumed that the mutation rate increases linearly with methylation level,

allowing both the intercept and slope to vary across the 16 4-mer sequence contexts (Fig 1B,C).

We implemented this model in a regression framework and estimated the scaled mutation rates

of unmethylated and methylated CpGs in each 4-mer context using the gnomAD polymorphism

data (S1 Table). In terms of scaled mutation rates, the rank order of the 16 4-mer contexts was

markedly different between unmethylated and methylated states, demonstrating a strong

interaction between the methyl group at the focal cytosine and flanking bases on CpG mutation

rate (Fig 1D; S1B Fig). Similar analysis of polymorphisms from the 1KG project [37] yielded

highly concordant context effects for methylated sites, although unmethylated sites showed

slightly greater divergence (S2B Fig).

Additionally, we compared our estimates with those predicted by previous mutation models

based on similar polymorphisms datasets. Carlson et al. (2018) estimated mutation rates for all

unique 3-mer, 5-mer and 7-mer contexts based on extremely rare variants without considering

methylation level of CpG sites. We found strong concordance between their estimates and our

"methylation-weighted" estimates for each 4mer context (S3A Fig; see Methods). In turn, our

estimates for unmethylated and methylated CpG sites are also in good concordance with 3-mer

estimates from the gnomAD mutation models [7,36] for CpG sites in the lowest and highest

methylation bins (S3B,C Fig).

To further validate our predicted mutation rates, we compared our estimates with mutation rates

estimated using an independent published dataset: de novo mutations (DNM) determined by

genome sequences of 5,420 parent-offspring trios [43]. Next, we identified sites with methylation

level <10% and >90% in sperm as proxies for unmethylated and methylated sites, respectively.

We then compared the observed DNM rates in different 4-mer contexts against our estimated

scaled mutation rates for unmethylated and methylated CpGs. These comparisons showed

strong correlations (Pearson's r=0.82 and 0.97), and the two sets of estimates are

approximately proportional, validating that our framework can reliably recover context-

dependent *de novo* CpG mutation rates from polymorphisms.

Independent effects of upstream and downstream bases

By organizing the 4-mer contexts by their 5' and 3' bases in rows and columns and visualizing

inferred scaled mutation rates in heatmaps, we observed clear marginal effects of each flanking

nucleotide on the mutation rates of unmethylated and methylated CpGs (Fig 2A). Specifically,

we found that a 5'A has a strong, positive marginal effect on CpG mutability regardless of

methylation status, which recapitulates previously reported hypermutable CpG motifs featuring

an A upstream to the focal C [1,8,34,42]. We also observed a negative effect of 5'G on

unmethylated CpG mutability and a strong negative effect of 5'T on methylated sites. For the

downstream base, 3'C is associated with lower mutability at unmethylated CpGs, while 3'T

exerts a negative impact on methylated cytosines. These inferred marginal effects of flanking

bases are all replicated qualitatively using 1KG polymorphism data (S2C Fig).

These consistent marginal effects—where upstream bases have similar effects regardless of

the downstream bases and vice versa— suggested that the flanking bases on the two sides of

the CpG act largely independently. We therefore asked to what degree the mutability of each 4-

mer context could be captured by the combined effect of the upstream and downstream bases.

To test this, we explicitly defined a model where the upstream and downstream nucleotides

independently affect the mutability of the focal cytosine (hereafter referred to as the up_1+down_1

model) (Table 1). This model contrasts with the 4-mer model presented above, which implicitly

allows for interactions between the two flanking bases.

Table 1. Descriptions of the various models specified in our regression framework.

Model	Formula	Variable	Variable description	Assumption
All		*m*	Methylation level as measured in sperm	
4-mer	*4mer * m*	*4mer*	4-mer including the focal CpG and one nucleotide on either side	Each 4-mer context may have unique baseline mutation rate and methylation effect
up_1+$down_1$	*U1 + D1 + m + U1:m + D1:m*	*U1*	Nucleotide one position upstream of focal C	Each nucleotide upstream or downstream to CpG affects the baseline mutation rate and mutagenic effect of methylation independently
		D1	Nucleotide one position downstream of focal G	
6-mer	*6mer * methylation*	*6mer*	6-mer including the focal CpG and two nucleotides on either side	Each 6-mer context may have unique baseline mutation rate and methylation effect
up_{21}+$down_{12}$	*U21 + D12 + m + U21:m + D12:m*	*U21*	Two nucleotides upstream of focal C	Each dimer upstream and downstream to CpG may have a unique baseline mutation rate and methylation effect
		D12	Two nucleotides downstream of focal G	
up_2+up_1+ $down_1$+$down_2$	*U2 + U1 + D1+ D2 + m + U2:m + U1:m + D1:m + D2:m*	*U2*	Nucleotide two positions upstream of focal C	Each nucleotide up to two positions upstream or downstream to CpG affects the baseline mutation rate and mutagenic effect of methylation independently
		U1	Nucleotide one position upstream of focal C	
		D1	Nucleotide one position downstream of focal G	
		D2	Nucleotide two positions downstream of focal G	

We found that the scaled mutation rates predicted by the up_1+$down_1$ model are highly

concordant with those from the 4-mer model (Pearson's r=0.98 for unmethylated and r=0.99 for

methylated CpG; Fig 2B), confirming largely independent effects of the upstream and

downstream nucleotides. Statistically, although the 4-mer model significantly outperforms the

up$_1$+down$_1$ model based on Akaike Information Criterion (AIC) and Bayesian Informational

Criterion (BIC), the gain in variance explained is small (25.18% vs 25.14%) (S2 Table). This

suggests that the simpler up$_1$+down$_1$ model largely captures variation in CpG mutation rates,

despite having fewer than half the parameters (14 vs. 32).

Fig 2. Independent effects of upstream and downstream nucleotides on CpG mutation

rates and shared context effects across primates. **A.** Scaled mutation rates in humans

estimated from the 4-mer model, which assumes interactions between the upstream and

downstream bases. **B.** Concordance between human mutation rates estimated from the 4-mer

model and from the up$_1$+down$_1$ model, which assumes independent effects of upstream and

downstream bases. Dashed line indicates   relationship. Points are labeled by the

nucleotides upstream and downstream of the CpG, with error bars representing 95% confidence

intervals for the estimates on both axes. **C, D.** Same as **A** but for chimpanzee and rhesus

macaque. Each heatmap shows relative mutation rates for that panel; color scales are not

comparable across panels.

Effects of sequence context on CpG mutation rate in other primate

species

We then applied our regression framework to two additional primate species, chimpanzee and

rhesus macaque, and observed patterns highly similar to those found in humans (Fig 2C and

Fig 2D; S1 Table). First, the 4-mer context effects on mutation rate differ between methylated

and unmethylated states (S4 Fig). Second, the mutation rates predicted by the up_1+$down_1$

model and 4-mer model are highly concordant, reaffirming the largely independent effects of

upstream and downstream bases on focal CpG mutability (chimpanzee: Pearson's r=0.96 and

0.99 for unmethylated and methylated CpG, respectively; rhesus macaque: Pearson's r=0.98 for

both; S4 Fig). Furthermore, most marginal effects of flanking bases are conserved across all

three studied primates. In each species, a 5'A increases the mutation rate of both unmethylated

and methylated CpGs, a 5'G or 3'C decreases the mutation rate of unmethylated CpGs (Fig

2A,C,D), and a 5'T or 3'T decreases the mutation rate of methylated CpGs. Together, these

findings suggest strong conservation in context-dependent mutation patterns of CpG sites

across these primate species.

Despite these broad similarities, clear inter-species differences emerge in the context-

dependency of CpG mutation rates. Due to greater estimation uncertainty, inter-species

correlations of point estimates are difficult to interpret for unmethylated sites. However, for

methylated sites where estimation resolution is high, similarity in context effects is surprisingly

highest between human and rhesus macaque, while either species shows lower concordance

with chimpanzee (Fig 3). Specifically, the greatest inter-species divergence occurs between

humans and chimpanzees (Pearson's r = 0.914), with the most pronounced discrepancies

observed in 4mer contexts with a downstream cytosine: a 3'C has a strong positive marginal

effect in humans and rhesus macaques but a relatively lower effect in chimpanzees (Fig 2A, C,

D; Fig 3A). Together, these results suggest that context-specific CpG mutability, particularly at

methylated sites, may have shifted during primate evolution (Fig 3B,C).

Fig 3. Similarity and differences in context effects on CpG mutability across primates. A.
Comparison of scaled mutation rate estimates between human and chimpanzee at
unmethylated and methylated sites, with error bars representing 95% confidence intervals for
the estimates on both axes. **B.** Same as **A.** for comparisons between human and rhesus
macaque. **C.** Same as **A.** for comparison between rhesus macaque and chimpanzee.

Mutational asymmetry within the same CpG dinucleotide

Given that the upstream and downstream nucleotides have asymmetric influences on the
mutability of the focal cytosine, we wondered whether the two adjacent C:G base pairs within
the same CpG dinucleotide differ in mutation rate. To test this, we compared mutation rates of
reverse complement 4-mer contexts (e.g., ACGC and GCGT). Since our modeling and analysis
always measure mutation rate at the cytosine position of each CpG, each pair of reverse
complement contexts correspond to the two adjacent C:G base pairs within the same CpG
dinucleotide (Fig 4).

In humans, we found significant mutational asymmetry between the two adjacent base pairs at

both methylated and unmethylated sites across most context pairs (Fig 4). The direction of this

asymmetry is concordant for unmethylated and methylated CpGs in four out of six 4-mer context

pairs. Notably, a cytosine directly flanked by an upstream A is consistently more mutable than

the neighboring G:C base pair within the same CpG dinucleotide, reflecting the strong

mutagenic effect of 5'A (Fig 2). For the CCGC:GCGG pair, we observed significant asymmetry

for unmethylated CpGs, with the C:G base pair flanked by an upstream C:G being more

mutable; the asymmetry is in the same direction but much weaker for methylated CpGs. For the

CCGA:TCGG and GCGA:TCGC pairs, methylation status determines the direction of mutational

asymmetry: the G:C base pair directly flanked by a downstream A:T is more mutable than its

counterpart when unmethylated but less mutable when methylated.

Consistent with the findings in humans, adjacent C:G base pairs within the same CpG also

differed in mutability in chimpanzees and rhesus macaques for most contexts (Fig 4). When

significant asymmetry is observed for a context pair, the direction of asymmetry is usually

consistent across all three species at both unmethylated and methylated sites, reflecting the

well-conserved effects of certain flanking bases (Fig 2). The only exception occurs at

methylated sites in CCGC:GCGG, where chimpanzee shows asymmetry in the opposite

direction than human and rhesus macaque, although the magnitudes of asymmetry are weak in

all three species. This flip in asymmetry echoes the reduced marginal effect of 3'C in

chimpanzee (Fig 2C).

Fig 4. Mutational asymmetry within a CpG site. Each bar represents a pair of reverse

complement 4-mer contexts (e.g., A**C**GC and G**C**GT) with a different focal cytosine for mutation

rate estimation, with color indicating the species (human, chimpanzee, or rhesus macaque).

Within the same CpG dinucleotide, adjacent C:G base pairs *a* and *b* often show different

mutation rates in both unmethylated and methylated states. Error bars represent 95%

confidence intervals calculated using the context-specific scaled mutation rate estimates from

our model (S1 Table) and assuming propagation of uncertainty.

Effects of expanded sequence context on CpG mutation rate in

primates

Beyond the immediate flanking bases, the expanded sequence context is a known predictor of

mutation rate across the genome [8,33,34]. We leveraged our regression framework to analyze

6-mer contexts, incorporating nucleotides two positions upstream and downstream. The 6-mer

model implicitly assumes all possible interactions between the four flanking nucleotides.

Consistent with our observations for 4-mer contexts, effect of the same 6-mer context differs

between unmethylated and methylated sites, confirming interactions between flanking

nucleotides and the methyl group (Fig 5).

Exploring the marginal effects of upstream and downstream dimers in humans revealed more

refined context-dependencies (Fig 5A). Echoing the strong mutagenic effect of upstream A

detected at the 4-mer scale, an upstream TA dimer has the strongest, positive marginal effect at

unmethylated sites, whereas an upstream GA or TA dimer is mutagenic for methylated CpGs.

Notably, at methylated sites, a T at the downstream +2 position (T_{+2}) is consistently associated

with low mutation rates, regardless of the intermediate +1 nucleotide.

Fig 5. Shared and diverged effects of upstream and downstream sequences on CpG
mutability across primates at 6-mer resolution. A. Scaled mutation rates estimated from the
6-mer model for human at unmethylated (upper) and methylated sites (lower). **B.** Concordance
between scaled mutation rates estimated from the 6-mer model and the $up_{21}+down_{12}$ model,
which assumes interactions only between bases within the upstream and downstream dimers.
Dashed line indicates   relationship. Points are labeled by the dimers upstream and
downstream of the CpG. Only contexts for which the difference between the two model
estimates exceeds 2.5 standard deviations from the mean difference are labeled. Error bars
represent 95% confidence intervals for the estimates. **C, D.** Same as **A** but for chimpanzee and

rhesus macaque. Each heatmap shows relative mutation rates for that panel; color scales are

not comparable across panels.

The row- and column-wise patterns in the 6-mer heatmaps suggested that the upstream and

downstream dimers may act independently. We tested this by comparing the full 6-mer model

against a simpler "$up_{21}+down_{12}$" model, which only includes interactions between adjacent

bases within each upstream or downstream dimer but no interaction between upstream and

downstream sequences. We again found high concordance in inferred scaled mutation rates

predicted by the two models (Pearson's r=0.95 and 0.97 for unmethylated and methylated CpG,

respectively; Fig 5B). Although the 6-mer model significantly outperforms the $up_{21}+down_{12}$ (S2

Table), the additional variance explained is small (25.52% vs 25.43%).

We then tested an even simpler model ($up_2+up_1+down_1+down_2$), where all flanking nucleotides

contributed independently to CpG mutability. This further simplified model showed weaker

concordance with the $up_{21}+down_{12}$ model (Pearson's r=0.93 and 0.94 for unmethylated and

methylated CpG, respectively; S5 Fig), suggesting that interactions within the upstream or

downstream dimers contribute more to CpG mutation rate variation. Although more complex

models statistically outperform simpler ones (S2 Table), they involve many more parameters

that require large datasets to estimate reliably. Interestingly, the $up_2+up_1+down_1+down_2$ model,

despite having fewer parameters and assuming no interactions at all, outperforms the 4mer

model that considers shorter contexts but full interaction among base pairs (variance explained

25.43% vs 25.18%; S2 Table), pointing to the significant influence of extended sequence on

mutation rates [8,33,34]. Overall, the $up_{21}+down_{12}$ model seems to offer the best trade-off by

capturing the bulk of context effects with an order of magnitude fewer parameters than the 6-

mer model (62 vs. 512).

Extending this analysis to include chimpanzee and rhesus macaque identified many conserved

and some species-specific 6-mer effects. We found that the upstream CG dimer and

downstream GC dimer are consistently associated with low mutation rates at unmethylated

sites; a 5'TA dimer has a strong, positive effect on both methylated and unmethylated CpG

mutability in all three species (Fig 5A,C,D). However, at methylated sites, differences in context

effects become more noticeable among the three species. Notably, the striking negative effect

of downstream T_{+2} on methylated CpG in humans is seen in rhesus macaques but slightly

edged by a strong negative effect of the downstream GC dimer in chimpanzees. Across all three

species, the most complex 6-mer model always performs considerably better than simpler

models, with variance explained increasing with model complexity (S2 Table).

Effects of sequence context on CpG mutation rate in an insect

species with minimal DNA methylation

Our analysis of primate polymorphisms treated methylation as a continuous variable, but sites

classified as "unmethylated" based on sperm bisulfite sequencing may still experience

methylation in other tissues or developmental stages. To approximate a truly unmethylated state,

we extended our analysis to the silkworm (*Bombyx mori*), a species with less than 1%

methylation genome-wide, largely in gene bodies [44]. We modified our regression framework to

isolate sequence context effects by eliminating the methylation component entirely (S1 Table).

As in our primate analyses, we also tested simpler additive models without higher-order

interactions between flanking bases.

In *B. mori*, mutation rate estimates from the 4-*mer* model and the up$_1$+down$_1$ model were highly

concordant with nearly identical AIC (Pearson's r = 0.98; S6A Fig; S2 Table), again indicating

that flanking bases influence CpG mutability largely independently. Crucially, some of the

marginal effects of flanking nucleotides in the silkworm mirror those found in primates at

unmethylated sites: a 5'A increases while a 3'C decreases CpG mutability (S6B Fig; Fig 2).

These shared patterns support deep conservation in context effects on CpG mutability that are

independent of DNA methylation.

Discussion

Here, we present a regression framework that enables estimation of CpG mutation rates in

different 4-mer and 6-mer contexts at unmethylated and methylated states, separately. Our

model introduces two key innovations: treating methylation level as a continuous predictor and

explicitly modeling recurrent mutations. This approach avoids the arbitrary binning of

methylation levels into discrete categories (e.g., high, medium, low) and reduces systematic

underestimation of mutation rates at hypermutable sites.

While sperm bisulfite sequencing provides a practical approximation of methylation levels in the

male germline, it has clear limitations. Sperm-based methylation data may not accurately reflect

levels in spermatogonial stem cells, and they fail to capture methylation profiles in other cell

types relevant to germline mutagenesis (e.g., female germ cells, early embryonic tissues). A

particular concern is that some sites classified as "unmethylated" in our analysis may in fact be

methylated in these other contexts. To strengthen our findings on unmethylated sites, we

applied our model to silkworm, a species with very low levels of DNA methylation that is largely

concentrated in gene bodies [44]. The recovery of consistent effects of certain flanking bases

(e.g., positive effect of upstream A) in this system supports that these mutation patterns likely

reflect intrinsic properties of unmethylated CpG in different DNA sequences rather than artifacts

of transient methylation and/or extrapolation.

Unlike several prior mutation models, we explicitly account for recurrent mutations on observed

polymorphisms. At hypermutable sites in large samples, recurrent mutations make the observed

polymorphism rate a poor linear proxy for mutation rate (Fig 1A). To address this, we used a

standard exponential transformation to relate polymorphism probability to mutation rate,

assuming a constant total length of the coalescent tree (T) across the genome. However,

genealogical histories certainly vary among genomic loci due to recombination and coalescent

stochasticity, as well as systematic factors like background selection, all of which cause

heterogeneity in T [45,46]. This assumption is especially problematic for species with very small

sample sizes, such as chimpanzees (S3 Table), where sampling stochasticity in the coalescent

process is greater. The over-simplifying assumption of constant T may distort our quantitative

estimates of context-specific mutation rates and introduce biases in between-species

comparisons. Nonetheless, these factors likely average out across the hundreds of thousands

of CpG sites within each context, so the heterogeneity in T is unlikely to alter the relative ranking

of contexts in the presence of substantial differences in mutability. We thus focus on qualitative

rather than quantitative comparisons across contexts and species. We also note that residual

variance in mutation rate that arise from mis-specification or unmodeled biological determinants

in our modeling framework could introduce uncertainty in the inferred ranking of sequence

contexts, especially those with similar mutation rates.

Despite limitations, our analysis clearly demonstrates that CpG mutation rate variation is shaped

by the interaction between the local sequence context and methylation at the focal cytosine.

Specifically, the effects of flanking sequences are distinct on unmethylated and methylated

cytosines but conserved across species, supporting the view that these two cytosine states act

as fundamentally different mutational substrates. Our analysis also demonstrates that the

upstream and downstream flanking sequences exert largely independent influences on

mutability, a pattern that holds for both 4-mer and 6-mer sequence contexts. Lastly, we identify

several strong marginal effects of adjacent nucleotides on CpG sites, including a strong

mutagenic effect of an upstream A on both unmethylated and methylated CpGs, negative

effects of 5'CG and 3'GC dimers on unmethylated CpGs, and a positive effect of 5'TA dimer on

unmethylated CpGs.

Importantly, the context and methylation effects we report here can help generate hypotheses

and guide interpretations about the underlying mechanisms of germline CpG>TpG

mutagenesis. For example, recent work has shown that transcription factor (TF) binding sites

may act as mutation hotspots by interfering with DNA repair [32], with TF motifs typically

spanning 6-10 nucleotides [47]. By contrast, our results indicate that CpG mutability is primarily

explained by largely independent effects of upstream and downstream sequences. This

suggests that the molecular mechanisms underlying context effects may operate in a modular

fashion—upstream and downstream sequences likely influence distinct steps in mutagenesis

rather than requiring complex long-range structural interactions. This argues against TF

occupancy as a dominant driver of context-dependent CpG mutation rate variation in the

germline. Future work should test whether this modularity extends to CpG transversions or other

single-nucleotide substitutions. If confirmed, this modularity has practical implications: simpler

additive or multiplicative models can achieve nearly the same predictive power as fully

parameterized models while remaining interpretable and computationally tractable.

Notably, an upstream A substantially increases mutability at both unmethylated and methylated

sites relative to other bases, which recapitulates previously reported hypermutable motifs

[1,8,34] and appears conserved across species, including chimpanzee, rhesus macaque, and

even silkworm. This deep conservation suggests that the ACG motif has intrinsic biophysical

properties—perhaps related to DNA shape—that elevate CpG mutation rates independent of

methylation status. Intriguingly, previous work shows that C>T mutation rates are also elevated

at non-CpG sites flanked by an upstream A [34], indicating that hypermutability may be a

general property of ACN sequence, regardless of CpG or non-CpG sites, which calls for

investigation into the mechanism.

In contrast, an upstream G does not elevate mutation rate at methylated sites and consistently

reduces mutation rate at unmethylated CpGs in all three primate species (Fig 2). This

observation contrasts with recent reports of elevated error rate of wild-type DNA polymerase ε in

GCG contexts [12]. Together, these findings suggest that although deamination-independent

replication-related CpG>TpG mutations occur, they cannot explain much of the context-

dependent mutation rate variation and are unlikely to represent the major source of germline

CpG mutations.

Cross-species comparisons of CpG mutation patterns offer additional mechanistic insights,

revealing that divergence in context effects does not strictly follow known phylogenetic

relationships. While correlations between point estimates at unmethylated sites may be difficult

to interpret due to estimation uncertainty, a clearly surprising pattern emerges at methylated

sites, where our model should have high resolution: humans and rhesus macaques exhibit

remarkably high similarity (Pearson's r = 0.986), while either species shows lower concordance

with the chimpanzee (Pearson's r = 0.914 and 0.964, respectively). At the current stage, we

cannot fully rule out technical factors related to the chimpanzee polymorphism data, making it

essential to replicate these findings using independent chimpanzee datasets or alternative

methodologies. If this divergence is indeed robust, the most parsimonious explanation is rapid

evolutionary shifts in the context-dependency of 5mC mutation rates along the chimpanzee

lineage. Such shifts may reflect changes in the context-specificity of enzymes involved in active

demethylation, such as the TET family [43], or in the repair of deamination products by proteins

like thymine DNA glycosylases (TDGs) [44].

Methods

Extracting CpG sites

We retrieved the hard-masked reference genomes for human (hg38), chimpanzee (panTro5),

rhesus macaque (Mmul10), and silkworm [48] from UCSC genome browser and Silkbase (see

Data sources). To minimize the effects of selection, we excluded genic regions based on NCBI

RefSeq annotations for all species studied; for human, we also removed phylogenetically

conserved regions defined by *phastcons* (see Data sources). For the primate species, we also

obtained the inferred reconstructed ancestral genome from the 10-primate EPO alignment [49]

mapped onto the human (hg38), chimpanzee (panTro5), and rhesus macaque (Mmul10)

genome. We then intersected the hard-masked reference genome for each species with the

corresponding ancestral reference and did all downstream analysis on hard-masked sites that

were also present in the ancestral genome.

Then, we scanned the genome to identify every occurrence of "CG" and recorded the position of

the dimer along with the flanking sequences. Each CpG dinucleotide was treated as two

independent cytosines: one on the forward strand in the extracted context and the other on the

reverse strand in the reverse complement sequence context. Based on flanking sequences, we

extracted 4-mer and 6-mer contexts as needed. To test whether any sites lost after intersection

could bias our estimates, we compared results based on all sites in hard-masked reference

genomes to those in intersected regions and observed largely concordant context-specific

mutation rate estimates (S7B Fig, S4 Table). The intersected genome yielded improved model

performance, as measured by the proportion of variance explained. Accordingly, all analyses

and results in the main text are based on the intersected regions.

Data sources

Reference Genome	Human: https://hgdownload.soe.ucsc.edu/goldenPath/hg38/bigZips/hg38.fa.masked.gz Chimpanzee: https://hgdownload.soe.ucsc.edu/goldenPath/panTro5/bigZips/panTro5.fa.masked.gz Rhesus: https://hgdownload.soe.ucsc.edu/goldenPath/rheMac10/bigZips/rheMac10.fa.masked.gz Silkworm: https://silkbase.ab.a.u-tokyo.ac.jp/cgi-bin/download.cgi
Ancestral Reference Genome	Human: https://ftp.ensembl.org/pub/release-106/fasta/ancestral_alleles/homo_sapiens_ancestor_GRCh38.tar.gz Chimpanzee: https://ftp.ensembl.org/pub/release-106/fasta/ancestral_alleles/pan_troglodytes_ancestor_Pan_tro_3.0.tar.gz Rhesus: https://ftp.ensembl.org/pub/release-106/fasta/ancestral_alleles/macaca_mulatta_ancestor_Mmul_10.tar.gz
Conserved Regions	Human: https://hgdownload.cse.ucsc.edu/goldenPath/hg38/database/phastConsEle

	ments30way.txt.gz
Gene Annotation	Human:http://hgdownload.soe.ucsc.edu/goldenPath/hg38/database/ncbiRefSeq.txt.gz Chimpanzee: https://hgdownload.soe.ucsc.edu/goldenPath/panTro5/database/ncbiRefSeq.txt.gz Rhesus: https://hgdownload.soe.ucsc.edu/goldenPath/rheMac10/database/ncbiRefSeq.txt.gz Silkworm: https://silkbase.ab.a.u-tokyo.ac.jp/cgi-bin/download.cgi
Methylation	Human: https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE138598
LiftOver	Human: https://hgdownload.cse.ucsc.edu/goldenpath/hg19/liftOver/hg19ToHg38.over.chain.gz Chimpanzee: https://hgdownload.soe.ucsc.edu/goldenPath/panTro6/liftOver/panTro6ToPanTro5.over.chain.gz https://hgdownload.soe.ucsc.edu/goldenPath/hg38/liftOver/hg38ToPanTro5.over.chain.gz Rhesus: https://hgdownload.soe.ucsc.edu/goldenPath/rheMac8/liftOver/rheMac8ToRheMac10.over.chain.gz

Polymorphism	Human:
	gnomAD v4: https://gnomad.broadinstitute.org/data
	1000 Genomes:
	http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/data_collections/1000G_2504_high_coverage/working/20201028_3202_raw_GT_with_annot/
	Chimpanzee:
	https://phaidra.univie.ac.at/detail/o:2066302
	Rhesus:
	https://zenodo.org/records/3515522
	Silkworm:
	https://ftp.cngb.org/pub/CNSA/data5/CNP0002456/Variation/CVF0092651/

Methylation data

For human, we downloaded bisulfite sequencing data from Chen et al. [38], which reported

sperm WGBS data from nine healthy individuals. When a C/T SNP exists at a CpG site,

methylation estimates are biased in individuals carrying one or two T alleles, as the T allele will

be mistaken for a bisulfite-converted unmethylated C. For instance, even if the CpG allele is

"fully" methylated in sperm, the presence of the TpG allele results in incorrect reporting of only

50% methylation in heterozygotes and 0% methylation in T/T homozygotes. To reduce this bias,

we calculated the average methylation level at each CpG by pooling methylated and

unmethylated read counts across samples. This approach effectively reduces biases for most

CpGs where the derived T allele is rare in the nine individuals but still underestimates

methylation levels at sites where the T allele is at high frequency. Additionally, this pooling

strategy assumes that methylation patterns are relatively consistent across individuals at most

CpG sites, which may not hold for regions with high inter-individual variability or imprinted loci.

We then lifted over the average methylation level from hg19 to hg38.

To test whether hydroxy-methylation (5hmC), which cannot be distinguished from 5mC by

traditional bisulfite sequencing, distorts our estimation of mutation rates for unmethylated

cytosine and 5mC, we used published 5hmC calls from TET-assisted bisulfite sequencing in

human embryonic stem cells [50]. In this dataset, about 2.1% of CpGs in putatively neutral

regions of the human genome have a recorded non-zero 5hmC level, with an average level of

21%. We re-ran our model after excluding all sites with any non-zero 5hmC signal, and found

that the context-dependent mutation rate estimates were nearly perfectly concordant with those

estimated using all sites (S8 Fig), suggesting that hydroxy-methylated sites mis-labeled as

methylated sites do not have an appreciable effect on the estimation of context-specific

mutation rates at unmethylated cytosine and 5mC.

For chimpanzee and rhesus macaque, we retrieved testes EM-Seq methylation calls averaged

between two individuals and corrected for genotype (A. Stolyarova, personal communications)

at each CpG dinucleotide. Chimpanzee methylation data was lifted over from panTro6 to

panTro5.

Filtering and polarization of polymorphism

We downloaded autosomal polymorphism data for human [36,37], rhesus macaque [51],

chimpanzee [52], and silkworm [53] (see Data sources). For rhesus macaque, variant

coordinates were lifted over from Mmul8 to Mmul10. For chimpanzee, variant coordinates were

lifted over from hg38 to panTro5. Singletons were excluded in the silkworm dataset to reduce

false-positive variant calls [54]. For human, we filtered singletons to positions with single-read

Umap100 mappability equal to 1, ensuring that variants were supported by uniquely mappable

100-bp reads and minimizing spurious rare variants due to alignment errors [55].

We polarized SNPs based on allele frequency, defining the major allele as the ancestral state

and the minor allele as the derived state (hereby referred to as MAF polarization). For silkworm,

we used *est-sfs* [56], which estimates the probability that the major allele of the focal species

(*Bombyx mori*) is the ancestral state using polymorphism data from the focal species and

alignment with a closely related outgroup (*Bombyx mandarina*). We annotated each SNP with

the type of substitution and retained only C>T and G>A mutations at CpG sites (extracted as

described above, see S3 Table).

We also used an alternate polarization method for the primate species, by inferring the ancestral

allele based on the 10 primate EPO ancestral genome reconstruction. While we found similar

context-specific mutation rate estimates using this method (S7A Fig), the MAF polarization

method generally yielded better model performance in terms of proportion variance explained

across all primate species, regardless of the CpG annotation method used (S4 Table).

Therefore, we decided to use MAF polarization in our main analysis.

The number of genotyped samples in a polymorphism dataset can vary across sites and can

systematically differ by context thus confounding the estimation of context-specific rates. To

address this, we examined the distribution in sample coverage across contexts in the

polymorphism datasets for the three primate species and found no significant differences (S9

Fig).

In addition, GC-biased gene conversion (gBGC) can have a profound impact on the frequency

of C>T polymorphisms [57]. Since the strength of gBGC depends on population sizes and the

specific effects are also sample-specific, differences in effective population sizes and sample

sizes across species can bias the scaled mutation rates. However, the strength of gBGC is not

expected to differ substantially across contexts and thus unlikely to lead to significant biases in

the rankings of contexts. To address this directly with data, we stratified CpG sites by

recombination rate [58] into four bins covering roughly the same number of CpG sites: high (>

0.389), mid1 (≤ 0.389), mid2 (≤ 0.026), and low (≤ 4.877e-05). We then estimated the mutation

rate for each bin separately. Consistent with our expectation, we find strong correlations

between estimates from all sites and those of each recombination bin for methylated CpGs

(Pearson's r > 0.99) and slightly lower correlation for unmethylated CpGs (Pearson's r>0.97).

The reduction of correlation at unmethylated sites may be explained by noisier point estimates

due to smaller numbers of sites with low methylation levels (S10 Fig).

Modeling effects of methylation and sequence contexts on CpG

mutation rate

To quantify the effects of local sequence context and methylation level on CpG mutability, we

modeled the mutation rate μ_i at a given cytosine i (within a CpG dinucleotide) as a linear

function of its methylation level m_i, allowing both the intercept and slope to vary with sequence

context:

$$\mu_i = \alpha^c + \beta^c m_i$$

where, α^c represents the baseline mutation rate for context c and β^c denotes the effect of full

methylation within that context.

For a site with mutation rate of μ per generation, the probability of having polymorphism in a

sample was modeled as $p = 1 - e^{-\mu T}$, where T is the total coalescent branch length (in

generations) connecting all sampled genomes. For simplicity, we assumed that T was constant

across all CpG sites in non-genic, non-conserved regions. The presence or absence of

polymorphism at each site was then modeled as a Bernoulli trial. i.e., drawn from a binomial

distribution $Binom(1, p)$ with $p = 1 - e^{-\mu T}$.

625   **Regression framework**

To implement this model, we used the *glm* function in R, specifying the binomial family and a

custom link function, which maps the polymorphism probability (p) to a linear predictor $\eta =$

$-\mu T = \log (1 - p)$ as follows:

•   Link function: $link(p) = \log (1 - p)$

•   Inverse link function: $link^{-1}(\eta) = 1 - e^{\eta}$

•   Derivative of the inverse link: $\frac{d\mu}{d\eta} = -e^{\eta}$

We fitted the generalized linear model to site-level data, where each record corresponded to

one base pair in a CpG site annotated with its methylation level, sequence context, and a binary

indicator of polymorphism status (0 = monomorphic, 1 = polymorphic). The regression model

was specified differently depending on the context length and type of interaction, as described in

Table 1.

After fitting the regression models to the polymorphism data, we estimated the scaled mutation

rates (μT) for each sequence context and methylation status (*x*=1 for methylated; *x*=0 for

unmethylated) using the predict() function in R with type = "link" to obtain the linear predictor (η).

To evaluate model performance, we extracted summary statistics from each fitted regression

and compared models using criteria that balance between model fit and complexity, namely the

Akaike Information Criterion (AIC) and the Bayesian Information Criterion (BIC), which is

preferred for larger datasets such as ours. We also measured explanatory power as the

proportion of variance in polymorphism status explained, calculated as the difference between

null and residual deviance, divided by the null deviance (S2 Table).

Acknowledgements

650   We thank Anastasia Stolyarova and Molly Przeworski for kindly sharing sperm methylation data

651   for chimpanzee and rhesus macaque before publication; Iain Mathieson, Fabian Ramos-

652   Almodovar, and other members of the Gao and Mathieson laboratories for helpful discussions.

References

1. Hodgkinson A, Eyre-Walker A. Variation in the mutation rate across mammalian genomes. Nat Rev Genet. 2011;12:756–66. https://doi.org/10.1038/nrg3098

2. McVicker G, Gordon D, Davis C, Green P. Widespread genomic signatures of natural selection in hominid evolution. PLoS Genet. 2009;5:e1000471. https://doi.org/10.1371/journal.pgen.1000471

3. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, et al. The mutational constraint spectrum quantified from variation in 141,456 humans. Nature. 2020;581:434–43. https://doi.org/10.1038/s41586-020-2308-7

4. Cassa CA, Weghorn D, Balick DJ, Jordan DM, Nusinow D, Samocha KE, et al. Estimating the selective effects of heterozygous protein-truncating variants from human exome data. Nat Genet. 2017;49:806–10. https://doi.org/10.1038/ng.3831

5. Agarwal I, Fuller ZL, Myers SR, Przeworski M. Relating pathogenic loss-of-function mutations in humans to their evolutionary fitness costs. eLife. 2023;12:e83172. https://doi.org/10.7554/eLife.83172

6. Zeng T, Spence JP, Mostafavi H, Pritchard JK. Bayesian estimation of gene constraint from an evolutionary model with gene features. Nat Genet. 2024;56:1632–43. https://doi.org/10.1038/s41588-024-01820-9

7. Chen S, Francioli LC, Goodrich JK, Collins RL, Kanai M, Wang Q, et al. A genomic mutational constraint map using variation in 76,156 human genomes. Nature. 2024;625:92–100. https://doi.org/10.1038/s41586-023-06045-0

8. Aggarwala V, Voight BF. An expanded sequence context model broadly explains variability in polymorphism levels across the human genome. Nat Genet. 2016;48:349–55. https://doi.org/10.1038/ng.3511

9. Blake RD, Hess ST, Nicholson-Tuell J. The influence of nearest neighbors on the rate and pattern of spontaneous point mutations. J Mol Evol. 1992;34:189–200. https://doi.org/10.1007/BF00162968

10. Karolak A, Levatić J, Supek F. A framework for mutational signature analysis based on DNA shape parameters. Korolev S, editor. PLOS ONE. 2022;17:e0262495. https://doi.org/10.1371/journal.pone.0262495

11. Liu Z, Samee MAH. Structural underpinnings of mutation rate variations in the human genome. Nucleic Acids Res. 2023;51:7184–97. https://doi.org/10.1093/nar/gkad551

12. Tomkova M, McClellan MJ, Crevel G, Shahid AM, Mozumdar N, Tomek J, et al. Human DNA polymerase ε is a source of C>T mutations at CpG dinucleotides. Nat Genet. 2024;56:2506–16. https://doi.org/10.1038/s41588-024-01945-x

13. Donigan KA, Sweasy JB. Sequence context-specific mutagenesis and base excision repair. Mol Carcinog. 2009;48:362–8. https://doi.org/10.1002/mc.20497

14. Mitra R, Pettitt BM, Blake RD. Conformational states governing the rates of spontaneous
transition mutations. Biopolymers. 1995;36:169–79. https://doi.org/10.1002/bip.360360206

15. Bergeron LA, Besenbacher S, Zheng J, Li P, Bertelsen MF, Quintard B, et al. Evolution of
the germline mutation rate across vertebrates. Nature. 2023;615:285–91.
https://doi.org/10.1038/s41586-023-05752-y

16. Kong A, Frigge ML, Masson G, Besenbacher S, Sulem P, Magnusson G, et al. Rate of de
novo mutations and the importance of father's age to disease risk. Nature. 2012;488:471–5.
https://doi.org/10.1038/nature11396

17. Bird AP. CpG islands as gene markers in the vertebrate nucleus. Trends Genet.
1987;3:342–7. https://doi.org/10.1016/0168-9525(87)90294-0

18. Agarwal I, Przeworski M. Mutation saturation for fitness effects at human CpG sites. eLife.
2021;10:e71513. https://doi.org/10.7554/eLife.71513

19. Chen C, Qi H, Shen Y, Pickrell J, Przeworski M. Contrasting Determinants of Mutation
Rates in Germline and Soma. Genetics. 2017;207:255–67.
https://doi.org/10.1534/genetics.117.1114

20. Gao Z, Moorjani P, Sasani TA, Pedersen BS, Quinlan AR, Jorde LB, et al. Overlooked roles
of DNA damage and maternal age in generating human germline mutations. Proc Natl Acad Sci.
2019;116:9491–500. https://doi.org/10.1073/pnas.1901259116

21. Shen J-C, Rideout WM, Jones PA. The rate of hydrolytic deamination of 5-methylcytosine in
double-stranded DNA. Nucleic Acids Res. 1994;22:972–6. https://doi.org/10.1093/nar/22.6.972

22. Schmutte C, Yang AS, Beart RW, Jones PA. Base excision repair of U:G mismatches at a
mutational hotspot in the p53 gene is more efficient than base excision repair of T:G
mismatches in extracts of human colon tumors. Cancer Res. 1995;55:3742–6.

23. Duncan BK, Miller JH. Mutagenic deamination of cytosine residues in DNA. Nature.
1980;287:560–1. https://doi.org/10.1038/287560a0

24. Seplyarskiy VB, Sunyaev S. The origin of human mutation in light of genomic data. Nat Rev
Genet. 2021;22:672–86. https://doi.org/10.1038/s41576-021-00376-2

25. Tomkova M, McClellan M, Kriaucionis S, Schuster-Böckler B. DNA Replication and
associated repair pathways are involved in the mutagenesis of methylated cytosine. DNA
Repair. 2018;62:1–7. https://doi.org/10.1016/j.dnarep.2017.11.005

26. Harpak A, Bhaskar A, Pritchard JK. Mutation Rate Variation is a Primary Determinant of the
Distribution of Allele Frequencies in Humans. Eyre-Walker A, editor. PLOS Genet.
2016;12:e1006489. https://doi.org/10.1371/journal.pgen.1006489

27. Frederico LA, Kunkel TA, Shaw BR. A sensitive genetic assay for the detection of cytosine
deamination: determination of rate constants and the activation energy. Biochemistry.
1990;29:2532–7. https://doi.org/10.1021/bi00462a015

28. Beard BC, Wilson SH, Smerdon MJ. Suppressed catalytic activity of base excision repair
enzymes on rotationally positioned uracil in nucleosomes. Proc Natl Acad Sci. 2003;100:7465–
70. https://doi.org/10.1073/pnas.1330328100

29. Sibghat-Ullah null, Day RS. DNA-substrate sequence specificity of human G:T mismatch
repair activity. Nucleic Acids Res. 1993;21:1281–7. https://doi.org/10.1093/nar/21.5.1281

30. Hara R, Mo J, Sancar A. DNA Damage in the Nucleosome Core Is Refractory to Repair by
Human Excision Nuclease. Mol Cell Biol. 2000;20:9173–81.
https://doi.org/10.1128/MCB.20.24.9173-9181.2000

31. Polak P, Lawrence MS, Haugen E, Stoletzki N, Stojanov P, Thurman RE, et al. Reduced
local mutation density in regulatory DNA of cancer genomes is linked to DNA repair. Nat
Biotechnol. 2014;32:71–5. https://doi.org/10.1038/nbt.2778

32. Sabarinathan R, Mularoni L, Deu-Pons J, Gonzalez-Perez A, López-Bigas N. Nucleotide
excision repair is impaired by binding of transcription factors to DNA. Nature. 2016;532:264–7.
https://doi.org/10.1038/nature17661

33. Seplyarskiy V, Koch EM, Lee DJ, Lichtman JS, Luan HH, Sunyaev SR. A mutation rate
model at the basepair resolution identifies the mutagenic effect of polymerase III transcription.
Nat Genet. 2023;55:2235–42. https://doi.org/10.1038/s41588-023-01562-0

34. Carlson J, Locke AE, Flickinger M, Zawistowski M, Levy S, Myers RM, et al. Extremely rare
variants reveal patterns of germline mutation rate heterogeneity in humans. Nat Commun.
2018;9:3753. https://doi.org/10.1038/s41467-018-05936-5

35. Adams CJ, Conery M, Auerbach BJ, Jensen ST, Mathieson I, Voight BF. Regularized
sequence-context mutational trees capture variation in mutation rates across the human
genome. Balding D, editor. PLOS Genet. 2023;19:e1010807.
https://doi.org/10.1371/journal.pgen.1010807

36. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, et al. The mutational
constraint spectrum quantified from variation in 141,456 humans. Nature. 2020;581:434–43.
https://doi.org/10.1038/s41586-020-2308-7

37. Fairley S, Lowy-Gallego E, Perry E, Flicek P. The International Genome Sample Resource
(IGSR) collection of open human genomic variation resources. Nucleic Acids Res.
2020;48:D941–7. https://doi.org/10.1093/nar/gkz836

38. Chen X, Lin Q, Wen J, Lin W, Liang J, Huang H, et al. Whole genome bisulfite sequencing
of human spermatozoa reveals differentially methylated patterns from type 2 diabetic patients. J
Diabetes Investig. 2020;11:856–64. https://doi.org/10.1111/jdi.13201

39. Zhou Y, He F, Pu W, Gu X, Wang J, Su Z. The Impact of DNA Methylation Dynamics on the
Mutation Rate During Human Germline Development. G3 GenesGenomesGenetics.
2020;10:3337–46. https://doi.org/10.1534/g3.120.401511

40. Exome Aggregation Consortium, Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E,
et al. Analysis of protein-coding genetic variation in 60,706 humans. Nature. 2016;536:285–91.
https://doi.org/10.1038/nature19057

41. Exome Aggregation Consortium, Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E,
et al. Analysis of protein-coding genetic variation in 60,706 humans. Nature. 2016;536:285–91.
https://doi.org/10.1038/nature19057

42. Schraiber JG, Spence JP, Edge MD. Estimation of demography and mutation rates from
one million haploid genomes. Am J Hum Genet. 2025;112:2152–66.
https://doi.org/10.1016/j.ajhg.2025.07.008

43. Palsson G, Hardarson MT, Jonsson H, Steinthorsdottir V, Stefansson OA, Eggertsson HP,
et al. Complete human recombination maps. Nature. 2025;639:700–7.
https://doi.org/10.1038/s41586-024-08450-5

44. Xiang H, Zhu J, Chen Q, Dai F, Li X, Li M, et al. Single base-resolution methylome of the
silkworm reveals a sparse epigenomic map. Nat Biotechnol. 2010;28:516–20.
https://doi.org/10.1038/nbt.1626

45. Bromham L, Penny D. The modern molecular clock. Nat Rev Genet. 2003;4:216–24.
https://doi.org/10.1038/nrg1020

46. Duchêne DA, Tong KJ, Foster CSP, Duchêne S, Lanfear R, Ho SYW. Linking Branch
Lengths across Sets of Loci Provides the Highest Statistical Support for Phylogenetic Inference.
Kosakovsky Pond S, editor. Mol Biol Evol. 2020;37:1202–10.
https://doi.org/10.1093/molbev/msz291

47. Stewart AJ, Hannenhalli S, Plotkin JB. Why Transcription Factor Binding Sites Are Ten
Nucleotides Long. Genetics. 2012;192:973–85. https://doi.org/10.1534/genetics.112.143370

48. Kawamoto M, Jouraku A, Toyoda A, Yokoi K, Minakuchi Y, Katsuma S, et al. High-quality
genome assembly of the silkworm, Bombyx mori. Insect Biochem Mol Biol. 2019;107:53–62.
https://doi.org/10.1016/j.ibmb.2019.02.002

49. Paten B, Herrero J, Beal K, Fitzgerald S, Birney E. Enredo and Pecan: Genome-wide
mammalian consistency-based multiple alignment with paralogs. Genome Res. 2008;18:1814–
28. https://doi.org/10.1101/gr.076554.108

50. Yu M, Hon GC, Szulwach KE, Song C-X, Zhang L, Kim A, et al. Base-Resolution Analysis of
5-Hydroxymethylcytosine in the Mammalian Genome. Cell. 2012;149:1368–80.
https://doi.org/10.1016/j.cell.2012.04.027

51. Harris RA, Raveendran M, Worley KC, Rogers J. Unusual sequence characteristics of
human chromosome 19 are conserved across 11 nonhuman primates. BMC Evol Biol.
2020;20:33. https://doi.org/10.1186/s12862-020-1595-9

52. Han S, Riyahi S, Huang X, Kuhlwilm M. A curated dataset of great ape genome diversity.
Sci Data. 2025;12:1835. https://doi.org/10.1038/s41597-025-06124-z

53. Tong X, Han M-J, Lu K, Tai S, Liang S, Liu Y, et al. High-resolution silkworm pan-genome
provides genetic insights into artificial selection and ecological adaptation. Nat Commun.
2022;13:5619. https://doi.org/10.1038/s41467-022-33366-x

54. Johnston HR, Hu Y, Cutler DJ. Population Genetics Identifies Challenges in Analyzing Rare
Variants. Genet Epidemiol. 2015;39:145–8. https://doi.org/10.1002/gepi.21881

55. Karimzadeh M, Ernst C, Kundaje A, Hoffman MM. Umap and Bismap: quantifying genome
and methylome mappability. Nucleic Acids Res [Internet]. 2018 [cited 2026 Jan 20];
https://doi.org/10.1093/nar/gky677

56. Keightley PD, Jackson BC. Inferring the Probability of the Derived vs. the Ancestral Allelic
State at a Polymorphic Site. Genetics. 2018;209:897–906.
https://doi.org/10.1534/genetics.118.301120

57. Glémin S, Arndt PF, Messer PW, Petrov D, Galtier N, Duret L. Quantification of GC-biased
gene conversion in the human genome. Genome Res. 2015;25:1215–28.
https://doi.org/10.1101/gr.185488.114

58. Halldorsson BV, Palsson G, Stefansson OA, Jonsson H, Hardarson MT, Eggertsson HP, et
al. Characterizing mutagenic effects of recombination through a sequence-level genetic map.
Science. 2019;363:eaau1043. https://doi.org/10.1126/science.aau1043

Supporting Information

S1 Fig. Interaction between the methyl group at the focal cytosine and flanking bases on

CpG mutation rate. A. Polymorphism rate for each 4-mer context, stratified by methylation

levels. CpG sites were partitioned into 20 equal-width bins (ranging from [0, 0.05) to [0.95, 1]).

Each point represents the proportion of SNPs among all CpG sites within a methylation bin,

along with 95% confidence intervals assuming binomial sampling. Each context is represented

by a color denoting the nucleotide at the 5' position and a shape denoting the nucleotide at the

3' position. **B.** Scaled mutation rates for each 4-mer context as estimated by the model for

unmethylated and methylated sites. Points are labeled by the nucleotides upstream and

downstream of the CpG, with error bars representing 95% confidence intervals for the

estimates.

S2 Fig. Concordance between mutation rates for 4-mer CpG contexts estimated from

gnomAD (v4) and 1000 Genomes (1KG) polymorphism datasets. A. Scaled mutation rate

estimates for unmethylated and methylated sites for each 4-mer context using 1KG data. The

contexts are sorted in descending order of mutability when unmethylated and error bars

represent 95% confidence intervals for the estimates. **B.** Comparison of mutation rates

estimated using polymorphisms from gnomAD and 1KG at unmethylated and methylated sites,

respectively. **C.** Heatmap of scaled mutation rate estimates using the 4-mer model at

unmethylated and methylated sites, using polymorphism data from 1KG. Each heatmap shows

relative mutation rates for that panel; color scales are not comparable between panels. **D.**

Concordance between mutation rates estimated from the 4-mer model and from the up_1+$down_1$

model using 1KG data. Points are labeled by the nucleotides upstream and downstream of the

CpG, with error bars representing 95% confidence intervals for the estimates in panels B and D.

The dashed line indicates $x = y$.

S3 Fig. Context-specific scaled mutation rates are consistent with estimates from other

published context-dependent mutation models. A. Comparison with rates published in

Carlson et al. (2018). To compare Carlson et al.'s estimates for 5-mers and 7-mers with our 4-

mer and 6-mer results, we derived 4-mer and 6-mer rates as weighted averages of the mutation

rate estimates for 5-mer (top) and 7-mer (bottom) contexts reported by Carlson et al., using the

average methylation level of each 5-mer or 7-mer context as weights. For the expanded context

comparison, a linear fit constrained through the origin is shown (dashed line). Only contexts with

residuals exceeding 2.5 standard deviations from the fitted line are labeled. **B, C.** Comparison

with rates published from gnomAD mutation rate models (Karczewski et al., 2020; Chen at al.,

2024) for unmethylated (B) and methylated CpGs (C). To facilitate comparison, we computed

the mutation rate for each 3-mer as the mean of estimates of the four corresponding 4-mer

contexts sharing the same upstream base (e.g., rate for ACG is the mean of the rates of ACGA,

ACGC, ACGG and ACGT). In all panels, points are labeled by the nucleotides upstream and

downstream of the CpG, with horizontal error bars representing 95% confidence intervals for the

model estimates.

S4 Fig. Independent effects of upstream and downstream bases on CpG mutability in

chimpanzee and rhesus macaque. A. Concordance between mutation rates as estimated

from 4-mer model and from up_1 + $down_1$ model which assumes independent effects of upstream

and downstream bases in chimpanzee for unmethylated and methylated sites. **B.** Same as (A),

for rhesus macaque. Points are labeled by the upstream and downstream nucleotide flanking

the CpG, and error bars denote 95% confidence intervals. The dashed line indicates $x = y$.

865
S5 Fig. Concordance in mutation rate estimates between models with and without

interactions within upstream and downstream dimers. Comparison of human mutation rates

as estimated from the up_{21}+$down_{12}$ model, which assumes interactions between bases within

the dimer upstream and downstream to the CpG, and the up_2+up_1+$down_1$+$down_2$ model, which

assumes independent effects of every flanking base. Only contexts for which the difference

between the two models' estimates is more than 2.5 standard deviations away from the average

difference across all contexts are labelled. Error bars represent 95% confidence intervals for the

estimates on both axes. The dashed line indicates $x = y$.

S6 Fig. Independent effects of upstream and downstream bases on CpG mutability in the

insect species, *Bombyx mori* (silkworm). A. Concordance between mutation rates as

estimated from 4-mer model and from up_1+$down_1$ model. Mutation rate is modelled as a function

of sequence context alone**.** Points are labeled by the upstream and downstream nucleotide

flanking the CpG, and error bars denote 95% confidence intervals. The dashed line indicates

$x = y$. **B.** Heatmap of estimated scaled mutation rates using the 4-mer model.

S7 Fig. Context-specific scaled mutation rate estimates are highly concordant regardless

of CpG filtering or SNP polarization method. A. Estimates obtained by polarizing SNPs using

minor allele frequency (MAF) are compared to those based on the inferred ancestral allele

(ANC) for CpG sites present in both the hard-masked reference and inferred ancestral genome.

B. Estimates derived from the intersected reference genome—restricted to CpGs present in

both the hard-masked reference and inferred ancestral genome—are compared to those from

the masked reference genome without intersection ("Masked"); SNPs are polarized by MAF.

Points are labeled by the upstream and downstream nucleotide context flanking the CpG, and

error bars denote 95% confidence intervals around the estimates.

S8 Fig. **Scaled mutation rate estimates are nearly perfectly concordant after excluding**

hydroxy-methylated (5hmC) sites. Comparison of the model's scaled mutation rate estimates

for unmethylated and methylated sites when using all sites and when excluding sites with a non-

zero 5hmC level as measured in human ES-cells. Points are labeled by the upstream and

downstream nucleotide context flanking the CpG, and error bars denote 95% confidence

intervals around the estimates.

S9 Fig. Callable allele coverage across 4-mer contexts in primate polymorphism

datasets. Distribution of the number of callable alleles by 4-mer sequence context in the

human, chimpanzee, and rhesus macaque polymorphism dataset. Boxes show the median and

interquartile range, with whiskers extending to 1.5x the interquartile range.

S10 Fig. Context-specific scaled mutation rate estimates are consistent across recombination rate strata. A. Comparison of scaled mutation rate estimates across all sites versus sites stratified by recombination rate, shown separately for unmethylated (top) and methylated (bottom) CpGs. Panels correspond to recombination rate quartiles: **A.** highest ("High", > 0.389), **B** upper-middle ("Mid1", ≤ 0.389), **C.** lower-middle ("Mid2", ≤ 0.026), and **D.** lowest ("Low", ≤ 4.877×10^{-5}). The dashed line indicates $x = y$. Points are labeled by the upstream and downstream nucleotide flanking the CpG, and error bars denote 95% confidence intervals.

S1 Table. Scaled mutation rate estimates for each 4-mer and 6-mer CpG context at unmethylated and methylated sites (provided as excel file). This table contains separate worksheets for each species and model (e.g., 4-mer, 6-mer), with each sheet reporting the sequence context, methylation status, dataset-specific scaled mutation rate estimate (μT), and its associated standard error. For silkworm, we report estimates from 4-mer and $up_1 + down_1$ models only, and rates are modeled as a function of context alone.

S2 Table. Model comparison for human and non-human species. Summary statistics of each fitted regression model for the four species in our analysis. The proportion variance explained is calculated as the difference between null and residual deviance divided by the null deviance. In silkworm, the mutation rate is modeled as a function of sequence context alone.

S3 Table. Summary of CpG sites and SNPs for each species. Genic regions are excluded in our analysis using NCBI *RefSeq* annotations. For human, phylogenetically conserved regions as defined by *phastcons* are also excluded. We identify C>T SNPs by polarizing the substitution that gave rise to each SNP using minor allele frequencies for human, chimpanzee, and rhesus macaque. For silkworm, we used *est-sfs*, a method that infers the ancestral state probabilities at polymorphic sites using information from a focal and outgroup species.

S4 Table. Model comparison across CpG filtering and SNP polarization methods in human and non-human species. Summary statistics of the 4-mer fitted regression model for each of the three primates in our analysis using the following combinations of CpG filtering and SNP polarization

methods: intersected reference genome (CpGs present in both the hard-masked reference and

inferred ancestral genome) and minor allele frequency (MAF) polarization; hard-masked reference

genome and MAF polarization; intersected reference genome and ancestral allele polarization

(ANC). The proportion variance explained is calculated as the difference between null and residual

deviance, divided by the null deviance. We also show the number of CpG sites extracted from each

genomic build after excluding genic regions and conserved regions (for human) and the number of

CpG>TpG SNPs inferred using the corresponding polarization method.