Sequence context and methylation interact to shape germline mutation rate variation at CpG sites

Sheel Chandra[1], Ziyue Gao[2*]

[1] Department of Biology, University of Pennsylvania, Philadelphia, Pennsylvania, United States of America

[2] Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, United States of America

* Corresponding author

E-mail: ziyuegao@pennmedicine.upenn.edu

Abstract

A prominent example of sequence context-dependent mutation rate variation is the elevated transition rate at CpG sites, which is largely attributed to cytosine methylation. CpGs with different flanking sequences also exhibit mutation rate variation, but this variation is only partially correlated with context-specific methylation level. Here, we quantify the CpG mutation rate and mutagenic effect of methylation across sequence contexts. Using a regression framework that accounts for recurrent mutations, we analyze human polymorphisms from the gnomAD dataset to estimate mutation rates of unmethylated and methylated CpGs in each unique 4-mer or 6-mer context. We find that CpG mutation rate variation in the human genome is shaped by methylation at the focal cytosine, the flanking nucleotides, and interactions between them. Our analysis reveals distinct context-dependent mutation patterns for unmethylated and methylated cytosines, driven by largely independent effects of upstream and downstream sequences. Notably, an upstream adenine markedly increases CpG mutation rates regardless of methylation status or downstream sequences. Furthermore, upstream and downstream sequences have qualitatively similar effects in chimpanzee and rhesus macaque, indicating that some conserved, intrinsic sequence features shape CpG mutability. On the other hand, some inter-species differences, which are especially pronounced at methylated sites, point to recent evolutionary changes, possibly in context-specificity of proteins governing DNA demethylation and repair processes.

Author Summary

The DNA sequence surrounding a nucleotide strongly influences how likely it is to mutate. An extreme example is the CpG dinucleotide: cytosines in CpGs mutate far more frequently than other sites in the human genome. This is related to DNA methylation, a chemical modification that occurs almost exclusively at CpGs in vertebrates and makes cytosines more prone to mutations. However, CpGs in different sequence contexts also vary in their mutation rates, and methylation level alone cannot explain this variation. To gain insight into what processes drive this variation, we estimate mutation rates for methylated and unmethylated CpGs in different sequence contexts using human genetic variation data. We find that methylation and neighboring bases interact to influence CpG mutation rates, and that the DNA sequence on either side of the CpG exerts largely independent effects. Extending our analysis to other primates reveals both conserved and species-specific patterns, with differences being especially pronounced at methylated sites. Together, our results suggest that while intrinsic DNA sequence features underlie some conserved context effects, other differences reflect recent evolutionary changes in the mechanisms that regulate DNA demethylation and repair.

Introduction

Mutation rate varies across the human genome, even at the scale of individual nucleotides [1]. Quantifying and accounting for this mutation rate variation is important for making inferences about natural selection [2] and identifying functional coding [3–6] and non-coding regions [7]. Among the many predictors of mutation rate, one of the strongest is the local sequence context—the nucleotides flanking the mutated site [8,9]. Although context effects have been attributed to several mechanisms, including local DNA shape [10,11], biases in DNA polymerase fidelity [12], and differential repair efficiency across sequence environments [13,14], the relative importance of these mechanisms and whether they fully account for observed patterns remain open questions. Addressing these questions requires systematically characterizing how local sequence shapes site-specific mutation rate variation, which can help disentangle these contributions and shed light on the mechanistic basis of mutagenesis.

Among sequence contexts, the CpG dinucleotide is both exceptionally mutable and mechanistically informative. Across vertebrates, CpG sites experience C>T mutation rates an order of magnitude higher than those at other sites in the genome [1,15,16]. This hypermutability is directly linked to DNA methylation, as supported by two observations: first, in vertebrates, cytosine methylation occurs almost exclusively at CpG sites [17], and second, in humans, de novo C>T mutation rates at CpGs correlate strongly with germline methylation levels [18–20].

A widely accepted mechanism underlying the mutagenic effect of methylation is the spontaneous deamination of 5-methylated cytosine (5mC), which occurs at a much higher rate than for unmethylated cytosine [21]. Deamination of 5mC produces thymine, generating a T:G mismatch that can be converted to a T:A base pair if the mismatch is erroneously repaired or

left unrepaired until the next round of DNA replication. In contrast, deamination of unmethylated

cytosine yields uracil, a non-canonical base that is more reliably recognized and removed [22].

However, not all C>T mutations at CpG sites arise from deamination and faulty/inefficient repair.

Recent work on human DNA polymerase ε suggests that mis-incorporation of nucleotides

during replication can also contribute to CpG>TpG mutations, as this enzyme shows a bias

toward incorporating adenine opposite 5mC and even unmethylated cytosine in a CpG context

[12,23].

CpG sites also exhibit substantial mutation rate variation across sequence contexts [8,24].

Aggarwala and Voight [8] reported such variation and a significant yet modest correlation (R^2 =

0.33) between methylation in sperm and the probability of C>T substitutions in 7-*mer* CpG

contexts, suggesting that variation in methylation level cannot fully explain CpG mutation rate

variation across sequence contexts. A key question, then, is what makes some CpG sites, by

virtue of their sequence context, more mutable than others. Intrinsic features of the local DNA

structure may influence repair efficiency of T:G mismatches [14,25,26], the rate of deamination

[27], or replication fidelity [12,23], offering distinct mechanistic explanations. In addition, context-

specific DNA-binding proteins such as transcription factors can also modulate mutation rate by

interfering with repair machinery [28–30]. To evaluate these mechanisms, we need a mutation

model that explicitly considers both methylation state and sequence context for each CpG site.

Such a model would enable us to disentangle the contribution of a well-defined biochemical

mechanism (i.e., 5mC deamination) from additional context effects that may arise from DNA

replication, damage, repair, or protein binding.

Here, we introduce a framework for modeling CpG mutation rates across sequence contexts

and methylation levels. Unlike existing models that ignore methylation [31,32] or treat it as a

categorical variable [3,33], our approach incorporates methylation as a continuous predictor,

enabling us to disentangle the contributions of methylation and local sequence context while

leveraging information from CpG sites across different methylation levels. Applying this

approach to human polymorphism data from the Genome Aggregation Database (gnomAD

v4.0), we estimate mutation rates for both unmethylated and methylated CpGs in different 4-mer

and 6-mer contexts. We also extend our analysis to two additional primate species to identify

conserved and divergent patterns in CpG mutation rate variation. Overall, our work shows that

unmethylated and methylated cytosines represent fundamentally different mutational substrates

with distinct context dependencies that together shape CpG mutation rate variation across the

genome.

Results

Interaction between methyl group and flanking nucleotides on CpG mutation rate

We hypothesized that the variation in CpG mutation rates across contexts is shaped by both

differences in baseline mutation rates of unmethylated cytosines and differential mutagenic

effects of methylation (i.e. the difference in mutation rate between methylated and unmethylated

cytosines). We started by calculating the CpG transition polymorphism rate for each

tetranucleotide (4-mer) context at varying methylation levels, using single nucleotide

polymorphisms (SNPs) as a proxy for mutation events in the ancestors of the sampled

genomes. For primary analysis, we used SNPs in the inter-genic regions of the human genome

from gnomAD [34], excluding singletons due to concerns of high false positive rates, and

polarized SNPs based on the inferred ancestral human genome (Methods); inter-genic SNPs

from the 1000 Genomes (1KG) Project including singletons were used for replication [35].

To approximate methylation levels in the male germline, we used bisulfite sequencing from

human sperm [36]. We recognize that germline mutations can arise in both the male and female

germlines, as well as during early embryonic development, so the most relevant methylation

metric should be a weighted average of methylation levels across these different stages.

However, we chose to use sperm bisulfite sequencing data as a proxy for germline methylation

level for two reasons. First, the site-level methylation intensity in sperm shows the strongest

correlation with CpG mutation rate and has much greater predictive power than data from any

other developmental stages [37]. Second, whole-genome bisulfite sequencing data for tissues

other than testis and/or sperm are much more limited and not corrected for genotypes of the

assayed individuals, which creates biases in methylation level measurement (see Methods).

As expected, we found that the polymorphism rate increases with germline methylation level for

each 4-mer context (Fig 1A). However, the increase in polymorphism rate is approximately

linear only up to ~25% methylation, after which the curves become flatter. This non-linear

relationship between polymorphism rate and methylation can be explained by higher rates of

recurrent mutations at highly mutable sites in large samples [18,38]. This suggests that, at the

current sample size, the polymorphism probability of CpG sites no longer reflects their mutation

rate proportionately.

Fig 1. Interaction between sequence context and methylation shapes mutation rate

variation at CpG sites. A. Polymorphism rate for each 4-mer context across increasing

methylation level. Polymorphism rate was calculated across 20 bins of methylation level from 0

to 1. Each point represents the proportion of SNPs observed across the total number of sites

within a methylation bin, along with 95% confidence intervals assuming binomial sampling.

Each context is represented by a color denoting the nucleotide at the 5' position and a shape

denoting the nucleotide at the 3' position **B.** Proposed mutation rate model where mutation rate

μ_i for a given cytosine i in a CpG within a 4-mer sequence context c is a linear function of its

methylation level m_i . α^c is the context-specific baseline rate when methylation is zero, and β^c

captures the increase in mutation rate from unmethylated to fully methylated. **C.** Schematic

showing how mutation rates are related with the observed methylation level m_i and

polymorphism state at every site i. **D.** Scaled mutation rate estimates for unmethylated and

methylated sites in each 4-mer context. Contexts are sorted by descending mutation rate when

unmethylated. Shape and color are coded in the same way as in (A). Error bars represent 95%

confidence intervals for the estimates.

To account for recurrent mutations, we related the polymorphism probability p to the underlying

per generation mutation rate μ using an exponential transformation as $p = 1 - e^{-\mu T}$, where T

represents the total branch length of the coalescent tree connecting all sampled genomes. This

approach corrects for recurrent mutations to a first approximation and produces a sample-

scaled mutation rate, μT. We treated the observed polymorphism state (i.e., presence or

absence of polymorphism) at each site as a Bernoulli trial with success probability p (Fig 1C).

To model the effects of sequence context, methylation, and their interaction, we assumed that

the mutation rate increases linearly with methylation level, allowing both the intercept and slope

to vary across sequence contexts (Fig 1B,C).

We implemented this model in a regression framework and estimated the scaled mutation rates

of unmethylated and methylated CpGs in each 4-mer context using the gnomAD polymorphism

data. We observed that, in terms of scaled mutation rates, the order of the 16 4-mer contexts

was completely different between unmethylated and methylated states, demonstrating an

interaction between the methyl group at the focal cytosine and flanking bases on CpG mutation

rate (Fig 1D). Similar analysis of polymorphisms from the 1KG project [35] recovered highly

concordant orders of most to least mutable contexts (S1A Fig).

Independent effects of upstream and downstream bases

178   By organizing the 4-mer contexts by their 5' and 3' bases in rows and columns and visualizing

179   the inferred scaled mutation rates in heatmaps, we observed clear marginal effects of each

180   flanking nucleotide on the mutation rate of unmethylated and methylated CpGs, as

181   demonstrated by the row- and column-wise patterns (Fig 2A). Specifically, we found that 5'A

182   has a strong, positive marginal effect on CpG mutability at both unmethylated and methylated

183 sites, which recapitulates previously reported hypermutable CpG motifs with A upstream to the

184 focal C [1,8,31]. We also observed a negative effect of 5'G on CpG mutability at unmethylated

185 sites and a strong negative effect of 5'T on methylated sites. For the downstream base, 3'C has

186 a negative effect on unmethylated CpGs, while 3'T exerts a negative impact on methylated

187 cytosines. These inferred marginal effects of flanking bases are all replicated qualitatively using

188 1KG polymorphism data (S1B Fig).

190 These consistent marginal effects—where upstream bases have similar effects regardless of

191 the downstream bases and vice versa— suggested that the flanking bases on the two sides of

192 the CpG act largely independently. We therefore asked to what degree the effect of each 4-mer

193 context could be captured by the combined effect of the upstream and downstream bases. To

194 test this, we explicitly defined a model where the upstream and downstream nucleotides

195 independently affect the mutability of the focal cytosine (hereafter referred to as the up_1+down_1

196 model) (Table 1). This model contrasts with the 4-mer model presented above, which implicitly

197 allows for interactions between the two flanking bases.

199 **Table 1. Descriptions of the various models specified in our regression framework.**

Model	Formula	Variable	Variable description	Assumption
All		m	Methylation level as measured in sperm	
4-mer	$4mer * m$	$4mer$	4-mer including the focal CpG and one nucleotide on either side	Each 4-mer context may have unique baseline mutation rate and methylation effect
up_1+down_1	$U1 + D1 + m + U1{:}m + D1{:}m$	$U1$	Nucleotide one position upstream of focal C	Each nucleotide upstream or downstream to CpG affects the baseline mutation rate and mutagenic effect of methylation independently
		$D1$	Nucleotide one position downstream of focal G	

6-mer	*6mer * methylation*	*6mer*	6-mer including the focal CpG and two nucleotides on either side	Each 6-mer context may have unique baseline mutation rate and methylation effect
up$_{21}$+down$_{12}$	*U21 + D12 + m + U21:m + D12:m*	*U21*	Two nucleotides upstream of focal C	Each dimer upstream and downstream to CpG may have a unique baseline mutation rate and methylation effect
		D12	Two nucleotides downstream of focal G	
up$_2$+up$_1$+ down$_1$+down$_2$	*U2 + U1 + D1+ D2 + m + U2:m + U1:m + D1:m + D2:m*	*U2*	Nucleotide two positions upstream of focal C	Each nucleotide up to two positions upstream or downstream to CpG affects the baseline mutation rate and mutagenic effect of methylation independently
		U1	Nucleotide one position upstream of focal C	
		D1	Nucleotide one position downstream of focal G	
		D2	Nucleotide two positions downstream of focal G	

We found that the scaled mutation rates in the 16 unique 4-mer contexts predicted from the

up$_1$+down$_1$ model are highly concordant with those predicted from the 4-mer model (Pearson's

r=0.98 for both methylated and unmethylated CpG; Fig 2A), confirming largely independent

effects of the upstream and downstream nucleotides. Statistically, the 4-mer model is still

significantly better than the up$_1$+down$_1$ model based on Akaike Information Criterion (AIC), but

the gain in variance explained is small (12.148% vs. 12.124%) (S1 Table), suggesting that the

simpler up$_1$+down$_1$ model largely captures variation in CpG mutation rates, albeit with fewer

than half the parameters (14 vs. 32).

Fig 2. Independent effects of upstream and downstream nucleotides on CpG mutation

rates and shared context effects across primates. **A.** Scaled mutation rates in humans

estimated from the 4-mer model, which assumes interactions between the upstream and

downstream bases. **B.** Concordance between human mutation rates estimated from the 4-mer

model and from the up_1+down_1 model, which assumes independent effects of upstream and

downstream bases. Dashed line indicates $x=y$. Points are labeled by the nucleotides upstream

and downstream of the CpG, with error bars representing 95% confidence intervals for the

estimates. **C, D.** Same as **A** but for chimpanzee and rhesus macaque. Each heatmap shows

relative mutation rates for that panel; color scales are not comparable across panels.

Effects of sequence context on CpG mutation rate in other primate

species

We then applied our regression framework to two additional primate species, chimpanzee and

rhesus macaque, and recovered many similar observations as in humans (Fig 2C and Fig 2D).

First, the rank order of 4-mer contexts in mutation rate differs between methylated and

unmethylated states (S2 Fig). Second, the mutation rates predicted by the up_1+down_1 model

and 4-mer model are highly concordant, indicating largely independent effects of upstream and

downstream bases on focal CpG mutability (chimpanzee: Pearson's r=0.97 for unmethylated

and r = 0.98 for methylated CpG; rhesus macaque: Pearson's r=0.98 for both unmethylated and

methylated CpG; S2 Fig). Some of the marginal effects of flanking bases are also conserved

across the studied primates. In all three species, a 5'A increases the mutation rate of both

unmethylated and methylated CpGs, and a 5'G or 3'C decreases the mutation rate of

unmethylated CpGs (Fig 2A,C,D). Together, these findings suggest considerable conservation

in context-dependent mutation patterns of CpG sites across the three primate species.

Despite these similarities in overall patterns, there are clear inter-species differences in the

context-dependency of CpG mutation rate, demonstrated by the imperfect correlation in scaled

mutation rates across contexts (Fig 3). In general, 4-mer contexts show greater between-

species divergence at methylated sites than at unmethylated sites, with the greatest divergence

observed between humans and chimpanzees (Pearson's r = 0.61). Notably, discrepancies in

context effects between humans and chimpanzees are most pronounced in contexts with G/C at

both sites flanking the CpG, especially at methylated CpGs: such "G/C rich" contexts are

associated with relatively high mutation rates in humans but low mutation rates in chimpanzees

(Fig 3A). Together, our results show that context-specific CpG mutability, particularly at

methylated sites, is more diverged between these two closely related species, relative to each

with rhesus macaques (Fig 3B,C).

Fig 3. Similarity and differences in context effects on CpG mutability across primates . A.

Comparison of scaled mutation rate estimates between human and chimpanzee at

unmethylated (upper) and methylated sites (lower). Colors indicate whether the nucleotides

flanking the focal CpG at the 5' and 3' positions are strong (S; C or G) or weak (W; A or T)

bases. Error bars represent 95% confidence intervals for the estimates on both axes. **B.** Same

as **A.** for comparisons between human and rhesus macaque. **C.** Same as **A.** for comparison

between rhesus macaque and chimpanzee.

Mutational asymmetry within the same CpG dinucleotide

Given that the upstream and downstream nucleotides have asymmetric influences on the

mutability of the focal cytosine, we wondered whether the two adjacent C:G base-pairs within

the same CpG dinucleotide differ in mutation rate. To test this, we compared mutation rates of

reverse complement 4-mer contexts (e.g., ACGC and GCGT). Since our modeling and analysis

always measure mutation rate at the C position of each CpG, reverse complement contexts

correspond to the two adjacent C:G base pairs within the same CpG dinucleotide (Fig 4).

For humans, we found that at both methylated and unmethylated sites, all context pairs showed

significant mutational asymmetry between the two adjacent base pairs (Fig 4). The direction of

mutational asymmetry is concordant for unmethylated and methylated CpGs in four out of six 4-

mer context pairs. Notably, a cytosine directly flanked by an upstream A is more mutable than

the neighboring G:C base pair within the same CpG dinucleotide, consistent with the strong

mutagenic effect of 5'A (Fig 2). Two exceptions occur in the CCGA:TCGG and GCGA:TCGC

pairs, where methylation status determines the direction of mutational asymmetry: the C:G base

pair flanked by A:T is more mutable than its counterpart when unmethylated but less mutable

when methylated.

Consistent with the findings in humans, for most contexts, adjacent C:G base pairs within the

same CpG also differed in mutability in chimpanzees and rhesus macaques (Fig 4). Specifically,

in all three species, the C:G base pair flanked by an A:T is more mutable than the neighboring

G:C base pair regardless of methylation state. Importantly, when significant asymmetry is

observed for a context pair, the direction of asymmetry is usually consistent across all three

species at both unmethylated and methylated sites, reflecting the well-conserved effects of

certain flanking bases (Fig 2), except for CCGA:TCGG, where chimpanzee and human show

opposite directionality in asymmetry.

Fig 4. Mutational asymmetry within a CpG site. Each bar represents a pair of reverse

complement 4-mer contexts (e.g., ACGC and GCGT) that contain the same CpG dinucleotide,

with color indicating the species (human, chimpanzee, or rhesus macaque). Within the same

CpG dinucleotide, adjacent C:G base pairs *a* and *b* often show different mutation rates in both

unmethylated and methylated states. Error bars represent 95% confidence intervals calculated

using propagation of uncertainty.

Effects of expanded sequence context on CpG mutation rate in primates

The expanded sequence context, beyond bases immediately adjacent to the mutated site, is

also a known predictor of mutation rate across the genome [8,31,33]. We therefore wanted to

test to what degree expanded contexts could explain variation in CpG mutation rates. We

applied our regression framework to 6-mer contexts, now including the nucleotides two positions

upstream of the cytosine and two positions downstream of the guanine. The 6-mer model

assumes all possible interactions between the four nucleotides flanking the CpG. Consistent

with observations based on a 4-mer context, effect of the same 6-mer context differs between

unmethylated and methylated sites, indicating interactions between flanking nucleotides and the

methyl group (Fig 5).

Using the predicted mutation rates from the 6-mer model, we explored the marginal effects of

upstream and downstream dimers on the mutability of the focal CpG in humans (Fig 5A).

Echoing the strong mutagenic effect of upstream A detected at the 4-mer scale, our 6-mer

model revealed that an upstream TA dimer has the strongest, positive marginal effect on CpG

mutability at unmethylated sites, whereas an upstream GA dimer has the strongest, positive

marginal effect at methylated sites. We also found that at methylated sites, irrespective of the

nucleotide at the +1 downstream position, a T at the downstream +2 position (T_{+2}) is

consistently associated with the lowest mutation rates.

Fig 5. Shared and diverged effects of upstream and downstream bases on CpG mutability

across primates at 6-mer resolution. A. Scaled mutation rates estimated from the 6-mer

model for human at unmethylated (upper) and methylated sites (lower). **B.** Concordance

between scaled mutation rates estimated from the 6-mer model and the $up_{21}+down_{12}$ model,

which assumes interactions only between bases within the upstream and downstream dimers.

Dashed line indicates x=y. Points are labeled by the dimers upstream and downstream of the

CpG. Only contexts for which the difference between the two model estimates exceeds 2.5

standard deviations from the mean difference are labeled. Error bars represent 95% confidence

intervals for the estimates. **C, D.** Same as **A** but for chimpanzee and rhesus macaque. Each

heatmap shows relative mutation rates for that panel; color scales are not comparable across

panels.

Inspired by the distinct row- and column-specific patterns in the heatmaps showing results

generated by the 6-mer model, we tested a simpler model: the $up_{21}+down_{12}$ model, which only

includes interactions between adjacent bases within the upstream and downstream dimers but

no between upstream and downstream sequences. We found high concordance between

scaled mutation rates predicted from the 6-mer model and the $up_{21}+down_{12}$ model (Pearson's

r=0.95 and 0.97 for unmethylated and methylated CpG, respectively; Fig 5B), indicating that

upstream and downstream sequences exert mostly independent effects on mutation rate of the

focal CpG.

We then asked how an even simpler model, where all flanking nucleotides contributed

independently to CpG mutability, would perform. This model, called the $up_2+up_1+down_1+down_2$

model, showed weaker concordance with the $up_{21}+down_{12}$ model (Pearson's r=0.93 and 0.92 for

unmethylated and methylated CpG, respectively; S3 Fig), suggesting that interactions within the

upstream or downstream dimers contribute to CpG mutation rate variation, even though

interactions between upstream and downstream sequences play less important roles. However,

complex models with many more parameters do not perform considerably better than simpler

models (S1 Table). Overall, the $up_{21}+down_{12}$ model seems to offer the best balance by capturing

most context effects without the heavy parametrization of the 6-mer model.

When extending this analysis to include chimpanzee and rhesus macaque, we identified several

conserved context effects at the 6-mer resolution as well as some differences. We found that

the upstream dimer CG and downstream dimer GC were consistently associated with low

mutation rates at unmethylated sites in all three species (Fig 5A,C,D). As in humans, a 5'TA

dimer has a strong, positive effect on unmethylated CpG mutability in chimpanzees and rhesus

macaques. However, at methylated sites, differences in context effects become more noticeable

among the three species. Notably, the striking, negative effect of downstream T_{+2} on methylated

CpG in humans is not seen in chimpanzees. Nonetheless, in chimpanzees and rhesus

macaques, echoing our observations in humans, the most complex 6-mer model did not perform

considerably better than simpler models, and that the $up_{21}+down_{12}$ model provided the best

tradeoff between model complexity and variance explained (S1 Table). Overall, across all

primates, interactions within but not between upstream and downstream dimers can capture

most mutation rate variation at CpG sites.

Effects of sequence context on CpG mutation rate in an insect

species with minimal DNA methylation

Our analysis thus far has inferred the mutagenic effects of methylation by modeling it as a

continuous spectrum. However, sites classified as "unmethylated" in our analysis may still

experience methylation in other tissues or developmental stages that is not captured by sperm

bisulfite sequencing. Therefore, species with minimal methylation can function as a biological

control and can better approximate a truly unmethylated state. We extended our analysis to the

insect *Bombyx mori* (domestic silkworm), where less than 1% of the genome is methylated, with

methylated cytosines largely in gene bodies [39]. We modified our regression framework to

eliminate the effect of methylation, modeling the mutation rate as a function of sequence context

alone. As in our primate analyses, we also tested simpler additive models without higher-order

interactions between flanking bases.

In *B. mori*, we observed high concordance between mutation rate estimates from the 4-*mer*

model and the up_1+down_1 model with nearly identical AIC (Pearson's r = 0.98; S4B Fig; S1

Table), indicating that flanking bases influence CpG mutability mostly independently. Some of

the marginal effects of flanking nucleotides are similar between silkworm and primates at

unmethylated sites. Specifically, silkworms exhibit a positive effect of 5'A and a negative effect

of 3'C on CpG mutability (S4A Fig), mirroring the marginal effects at unmethylated sites in all

three primates (Fig 2). These shared patterns suggest context effects that are independent of

DNA methylation and may instead reflect intrinsic features of the local sequence itself.

However, when we extended the analysis to 6-mer contexts, we found less distinctive column-

or row-wise patterns (S4C Fig) and lower concordance between the 6-mer model and the

$up_{21}+down_{12}$ model (Pearson's r = 0.87, S4D Fig), suggesting that independent effects of

adjacent dimers only partially capture the variation. Interestingly, although the complex 6-mer

model showed a significant increase in the proportion of variance explained compared to the

$up_{21}+down_{12}$ and the $up_2+up_1+down_1+down_2$ models, it has a greater AIC suggesting worse

performance given increased model complexity (S1 Table). Together, these results suggest

diminishing returns with increased model complexity: while more complex models capture more

variance, the improvement in predictive power does not offset the cost of including additional

parameters.

Discussion

Here, we present a regression framework that estimates CpG mutation rates in different 4-mer

and 6-mer contexts at unmethylated and methylated sites. Our model introduces two key

innovations: treating methylation level as a continuous predictor and explicitly modeling

recurrent mutations. This approach avoids the arbitrary binning of methylation levels into

discrete categories (e.g., high, medium, low) and reduces systematic underestimation of

mutation rates at hypermutable sites.

While sperm bisulfite sequencing provides a practical approximation of methylation levels in the

male germline, it has clear limitations. Sperm-based methylation data may not accurately reflect

levels in spermatogonial stem cells, and they fail to capture methylation profiles in other cell

types relevant to germline mutagenesis (e.g., female germ cells, early embryonic tissues). A

particular concern is that some sites classified as "unmethylated" in our analysis may in fact be

methylated in these other contexts. To strengthen our findings on unmethylated sites, we

applied our model to silkworm, a species with very low levels of DNA methylation that is largely

concentrated in gene bodies [39]. The recovery of consistent effects of certain contexts (e.g.,

positive effect of upstream A) in this system supports the robustness of our findings for

unmethylated CpGs.

Unlike several prior mutation models, we explicitly account for effects of recurrent mutations on

observed polymorphisms. At hypermutable sites in large samples, recurrent mutations make the

observed polymorphism rate a poor linear proxy for mutation rate (Fig 1A). To address this, we

used a standard exponential transformation to relate polymorphism probability to mutation rate,

assuming that the total length of the coalescent tree (T) is constant across the genome.

However, genealogical histories certainly vary among genomic loci due to recombination and

coalescent stochasticity, as well as systematic factors like background selection, all of which

cause heterogeneity in T [40,41]. This assumption is especially problematic for species with

very small sample sizes, such as chimpanzees, where sampling stochasticity in the coalescent

process is greater. The over-simplifying assumption of constant T may distort our quantitative

estimates of context-specific mutation rates and introduce biases in between-species

comparisons. Nonetheless, because such heterogeneity should largely average out across CpG

sites within each context, it is unlikely to affect the relative ranking of contexts by mutability. We

thus focus on qualitative rather than quantitative comparisons across contexts and species.

Applying this modeling framework to population polymorphism datasets, we show that CpG

mutation rate variation is shaped by the interaction between the local sequence context and

methylation at the focal cytosine. Specifically, the effects of flanking sequences are often distinct

on unmethylated and methylated cytosines, even though such effects are shared across

species, supporting the view that these two cytosine states act as fundamentally different

mutational substrates. Our analysis also demonstrates that the upstream and downstream

bases flanking the CpG exert largely independent influences on mutability, a pattern that holds

for both 4-mer and 6-mer sequence contexts. Lastly, we identify several strong marginal effects

of adjacent nucleotides on CpG sites, including a strong mutagenic effect of an upstream A on

both unmethylated and methylated CpGs, a negative effect of flanking CG—either upstream or

downstream—on focal unmethylated CpGs, and a positive effect of the 5′TA dimer on

unmethylated CpGs.

Importantly, the context and methylation effects we report here can help generate hypotheses

and guide interpretations about the underlying mechanisms of germline CpG>TpG

mutagenesis. For example, recent work has shown that transcription factor (TF) binding sites

may act as mutation hotspots by interfering with DNA repair [30], with TF motifs typically

spanning 6-10 nucleotides [50]. By contrast, our results indicate that CpG mutability is primarily

explained by largely independent effects of upstream and downstream sequences. This

suggests that the molecular mechanisms underlying context effects may operate in a modular

fashion—upstream and downstream sequences likely influence distinct steps in mutagenesis

rather than requiring complex long-range structural interactions. This argues against TF

occupancy as a dominant driver of context-dependent CpG mutation rate variation in the

germline. Future work should test whether this modularity extends to CpG transversions or other

single-nucleotide substitutions. If confirmed, this modularity has practical implications: simpler

additive or multiplicative models can achieve nearly the same predictive power as fully

parameterized models while remaining interpretable and computationally tractable.

Notably, an upstream A substantially increases mutability at both unmethylated and methylated

sites relative to other bases, which recapitulates previously reported hypermutable motifs

[1,8,31] and appears conserved across species, including chimpanzee, rhesus macaque, and

even silkworm. This deep conservation suggests that the ACG motif has intrinsic biophysical

properties—perhaps related to DNA shape—that elevate CpG mutation rates independent of

methylation status. Intriguingly, previous work shows that C>T mutation rates are also elevated

at non-CpG sites flanked by an upstream A [31], indicating that hypermutability may be a

general property of ACN sequence, regardless of CpG or non-CpG sites, which calls for

investigation into the mechanism.

In contrast, an upstream G does not elevate mutation rate at methylated sites and consistently

reduces mutation rate at unmethylated CpGs in all three primate species (Fig 2). This

observation contrasts with recent reports of elevated error rate of wild-type DNA polymerase ε in

GCG contexts [12]. Together, this suggests that although replication-related CpG>TpG

mutations occur, they cannot explain the observed context-dependent mutation rate variation

and are unlikely to represent the major source of germline CpG mutations .

Cross-species comparisons of the CpG mutation patterns offer additional mechanistic insight.

Within primates, divergence in context effects between species appears more pronounced at

methylated sites than at unmethylated sites, pointing to recent evolutionary changes related to

methylated CpGs. These may reflect shifts in the context-specificity of proteins involved in

demethylation (e.g., TET enzymes [51]) or in DNA repair (e.g., thymine DNA glycosylases [52]).

Methods

Extracting CpG sites

We retrieved the hard-masked reference genomes for human (hg38), chimpanzee (panTro5),

rhesus macaque (Mmul10), and silkworm [53] from UCSC genome browser and Silkbase (see

Data sources). To minimize the effects of selection, we excluded genic regions based on NCBI

RefSeq annotations for all species studied; for human, we also removed phylogenetically

conserved regions defined by *phastcons* (see Data sources). For each species, we scanned the

reference genome to identify every occurrence of "CG" and recorded the position of the dimer

along with four nucleotides upstream and four nucleotides downstream to get the "10-mer"

sequence context for the CpG. Each CpG dinucleotide was treated as two independent

cytosines: one on the forward strand in the extracted context and the other on the reverse

strand in the reverse complement sequence context. From these 10-mers, we extracted 4-mer

and 6-mer contexts as needed.

Since reliable ancestral genome reconstructions were not available for all species studied, we

used each species' reference genome to identify CpG sites. We note that this approach will

miss ancestral CpG sites that are no longer present in the reference due to mutations (e.g.,

ancestral CpG sites where the reference sequences are TpG or CpA). To assess the potential

impact of this limitation, we repeated the analysis by extracting CpG sites from the inferred

human ancestral genome and observed qualitatively similar results (S5 Fig). To test whether

excluding repetitive regions using the hard-masked reference genome could technically bias our

results, we repeated the analysis by extracting CpG sites from the unmasked human reference

genome and again observed qualitatively similar results (S6 Fig), suggesting that the reported

context effects on CpG mutability are likely robust to the choice of reference versus ancestral

context and hard-masked versus unmasked reference genomes.

Data sources

Reference Genome	Human: https://hgdownload.soe.ucsc.edu/goldenPath/hg38/bigZips/hg38.fa.masked.gz Chimpanzee: https://hgdownload.soe.ucsc.edu/goldenPath/panTro5/bigZips/panTro5.fa.masked.gz Rhesus: https://hgdownload.soe.ucsc.edu/goldenPath/rheMac10/bigZips/rheMac10.fa.masked.gz Silkworm: https://silkbase.ab.a.u-tokyo.ac.jp/cgi-bin/download.cgi
Conserved Regions	Human: https://hgdownload.cse.ucsc.edu/goldenPath/hg38/database/phastConsElements30way.txt.gz

Gene Annotation	Human:http://hgdownload.soe.ucsc.edu/goldenPath/hg38/database/ncbiRefSeq.txt.gz Chimpanzee: https://hgdownload.soe.ucsc.edu/goldenPath/panTro5/database/ncbiRefSeq.txt.gz Rhesus: https://hgdownload.soe.ucsc.edu/goldenPath/rheMac10/database/ncbiRefSeq.txt.gz Silkworm: https://silkbase.ab.a.u-tokyo.ac.jp/cgi-bin/download.cgi
Methylation	Human: https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE138598
LiftOver	Human: https://hgdownload.cse.ucsc.edu/goldenpath/hg19/liftOver/hg19ToHg38.over.chain.gz Chimpanzee: https://hgdownload.soe.ucsc.edu/goldenPath/panTro6/liftOver/panTro6ToPanTro5.over.chain.gz Rhesus: https://hgdownload.soe.ucsc.edu/goldenPath/rheMac8/liftOver/rheMac8ToRheMac10.over.chain.gz

Polymorphism	Human:
	gnomAD v4: https://gnomad.broadinstitute.org/data
	1000 Genomes:
	http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/data_collectio ns/1000G_2504_high_coverage/working/20201028_32 02_raw_GT_with_annot/
	Chimpanzee:
	https://www.ebi.ac.uk/ena/browser/view/PRJEB39475? show=analyses
	Rhesus:
	https://zenodo.org/records/3515522
	Silkworm:
	https://ftp.cngb.org/pub/CNSA/data5/CNP0002456/Vari ation/CVF0092651/
Ancestral Reference Genome	Human:
	https://ftp.ensembl.org/pub/release- 106/fasta/ancestral_alleles/homo_sapiens_ancestor_G RCh38.tar.gz
	Chimpanzee:
	https://ftp.ensembl.org/pub/release- 106/fasta/ancestral_alleles/pan_troglodytes_ancestor_ Pan_tro_3.0.tar.gz
	Rhesus:

	https://ftp.ensembl.org/pub/release-106/fasta/ancestral_alleles/macaca_mulatta_ancestor_Mmul_10.tar.gz

Methylation data

For human, we downloaded bisulfite sequencing data from Chen et al. [36], which reported

sperm WGBS data from nine healthy individuals. When a C/T SNP exists at a CpG site,

methylation estimates are biased in individuals carrying one or two T alleles, as the T allele will

be mistaken for a bisulfite-converted unmethylated C. For instance, even if the CpG allele is

"fully" methylated in sperm, the presence of the TpG allele results in incorrect reporting of only

50% methylation in heterozygotes and 0% methylation in T/T homozygotes. To reduce this bias,

we calculated the average methylation level at each CpG by pooling methylated and

unmethylated read counts across samples. This approach effectively reduces biases for most

CpGs where the derived T allele is rare in the nine individuals but still underestimates

methylation levels at sites where the T allele is at high frequency. Additionally, this pooling

strategy assumes that methylation patterns are relatively consistent across individuals at most

CpG sites, which may not hold for regions with high inter-individual variability or imprinted loci.

We then lifted over the average methylation level from hg19 to hg38.

For chimpanzee and rhesus macaque, we retrieved testes EM-Seq methylation calls averaged

between two individuals and corrected for genotype (A. Stoliarova, personal communications) at

each CpG dinucleotide. Chimpanzee methylation data was lifted over from panTro6 to panTro5.

Filtering and polarization of polymorphism

We downloaded autosomal polymorphism data for human [34,35], rhesus macaque [54],

chimpanzee [55], and silkworm [56] (see Data sources). For rhesus macaque, variant

coordinates were lifted over from Mmul8 to Mmul10. Singletons were excluded in gnomAD (v4)

and the silkworm datasets to reduce false-positive variant calls [57]. To polarize the substitution

that gave rise to each SNP, we used ancestral allele reconstructions based on the 10 primate

EPO alignment [58] for human (hg38), chimpanzee (panTro5), and rhesus macaque (Mmul10).

For silkworm, we used *est-sfs* [59], which estimates the probability that the major allele of the

focal species (*Bombyx mori*) is the ancestral state using polymorphism data from the focal

species and alignment with a closely related outgroup (*Bombyx mandarina*). We then annotated

each SNP with the type of substitution and retained only C>T and G>A mutations at CpG sites

in the reference genome (extracted as described above, see S2 Table).

Modeling effects of methylation and sequence contexts on CpG

mutation rate

To quantify the effects of local sequence context and methylation level on CpG mutability, we

modeled the mutation rate μ_i at a given cytosine i (within a CpG dinucleotide) as a linear

function of its methylation level m_i, allowing both the intercept and slope to vary with sequence

context:

$$\mu_i = \alpha^c + \beta^c\, m_i$$

where, α^c represents the baseline mutation rate for context c and β^c denotes the effect of full

methylation within that context.

For a site with mutation rate of μ per generation, the probability of having polymorphism in a

sample was modeled as $p = 1 - e^{-\mu T}$, where T is the total coalescent branch length (in

generations) connecting all sampled genomes. For simplicity, we assumed that T was constant

across all CpG sites in non-genic, non-conserved regions. The presence or absence of

polymorphism at each site was then modeled as a Bernoulli trial. i.e., drawn from a binomial

distribution $Binom(1, p)$ with $p = 1 - e^{-\mu T}$.

Regression framework

To implement this model, we used the *glm* function in R, specifying the binomial family and a

custom link function, which maps the polymorphism probability (p) to a linear predictor $\eta =$

$-\mu T = \log(1 - p)$ as follows:

• Link function: $link(p) = \log(1 - p)$

• Inverse link function: $link^{-1}(\eta) = 1 - e^{\eta}$

• Derivative of the inverse link: $\frac{d\mu}{d\eta} = -e^{\eta}$

We fitted the generalized linear model to site-level data, where each record corresponded to

one base-pair in a CpG site annotated with its methylation level, sequence context, and a binary

indicator of polymorphism status (0 = monomorphic, 1 = polymorphic). The regression model

was specified differently depending on the context length and type of interaction, as described in

Table 1.

After fitting the regression models to the polymorphism data, we estimated the scaled mutation

rates (μT) for each sequence context and methylation status (*x*=1 for methylated; *x*=0 for

unmethylated) using the predict() function in R with type = "link" to obtain the linear predictor (η).

To evaluate model performance, we extracted summary statistics from each fitted regression

and compared models using the Akaike Information Criterion (AIC), which balances between

model fit and complexity. We also measured explanatory power as the proportion of variance in

polymorphism status explained, calculated as the difference between null and residual

deviance, standardized by the null deviance.

Acknowledgments

We thank Anastasia Stoliarova and Molly Przeworski for kindly sharing sperm methylation data

for chimpanzee and rhesus macaque before publication; Iain Mathieson, Fabian Ramos-

Almodovar, and other members of the Gao and Mathieson laboratories for helpful discussions.

References

1. Hodgkinson A, Eyre-Walker A. Variation in the mutation rate across mammalian genomes.
Nat Rev Genet. 2011;12:756–66. https://doi.org/10.1038/nrg3098

2. McVicker G, Gordon D, Davis C, Green P. Widespread genomic signatures of natural
selection in hominid evolution. PLoS Genet. 2009;5:e1000471.
https://doi.org/10.1371/journal.pgen.1000471

3. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, et al. The mutational
constraint spectrum quantified from variation in 141,456 humans. Nature. 2020;581:434–43.
https://doi.org/10.1038/s41586-020-2308-7

4. Cassa CA, Weghorn D, Balick DJ, Jordan DM, Nusinow D, Samocha KE, et al. Estimating the
selective effects of heterozygous protein-truncating variants from human exome data. Nat
Genet. 2017;49:806–10. https://doi.org/10.1038/ng.3831

5. Agarwal I, Fuller ZL, Myers SR, Przeworski M. Relating pathogenic loss-of-function mutations
in humans to their evolutionary fitness costs. eLife. 2023;12:e83172.
https://doi.org/10.7554/eLife.83172

6. Zeng T, Spence JP, Mostafavi H, Pritchard JK. Bayesian estimation of gene constraint from
an evolutionary model with gene features. Nat Genet. 2024;56:1632–43.
https://doi.org/10.1038/s41588-024-01820-9

7. Chen S, Francioli LC, Goodrich JK, Collins RL, Kanai M, Wang Q, et al. A genomic
mutational constraint map using variation in 76,156 human genomes. Nature. 2024;625:92–100.
https://doi.org/10.1038/s41586-023-06045-0

8. Aggarwala V, Voight BF. An expanded sequence context model broadly explains variability in
polymorphism levels across the human genome. Nat Genet. 2016;48:349–55.
https://doi.org/10.1038/ng.3511

9. Blake RD, Hess ST, Nicholson-Tuell J. The influence of nearest neighbors on the rate and
pattern of spontaneous point mutations. J Mol Evol. 1992;34:189–200.
https://doi.org/10.1007/BF00162968

10. Karolak A, Levatić J, Supek F. A framework for mutational signature analysis based on DNA
shape parameters. Korolev S, editor. PLOS ONE. 2022;17:e0262495.
https://doi.org/10.1371/journal.pone.0262495

11. Liu Z, Samee MAH. Structural underpinnings of mutation rate variations in the human
genome. Nucleic Acids Res. 2023;51:7184–97. https://doi.org/10.1093/nar/gkad551

12. Tomkova M, McClellan MJ, Crevel G, Shahid AM, Mozumdar N, Tomek J, et al. Human
DNA polymerase ε is a source of C>T mutations at CpG dinucleotides. Nat Genet.
2024;56:2506–16. https://doi.org/10.1038/s41588-024-01945-x

13. Donigan KA, Sweasy JB. Sequence context-specific mutagenesis and base excision repair.
Mol Carcinog. 2009;48:362–8. https://doi.org/10.1002/mc.20497

14. Mitra R, Pettitt BM, Blake RD. Conformational states governing the rates of spontaneous
transition mutations. Biopolymers. 1995;36:169–79. https://doi.org/10.1002/bip.360360206

15. Bergeron LA, Besenbacher S, Zheng J, Li P, Bertelsen MF, Quintard B, et al. Evolution of
the germline mutation rate across vertebrates. Nature. 2023;615:285–91.
https://doi.org/10.1038/s41586-023-05752-y

16. Kong A, Frigge ML, Masson G, Besenbacher S, Sulem P, Magnusson G, et al. Rate of de
novo mutations and the importance of father's age to disease risk. Nature. 2012;488:471–5.
https://doi.org/10.1038/nature11396

17. Bird AP. CpG islands as gene markers in the vertebrate nucleus. Trends Genet.
1987;3:342–7. https://doi.org/10.1016/0168-9525(87)90294-0

18. Agarwal I, Przeworski M. Mutation saturation for fitness effects at human CpG sites. eLife.
2021;10:e71513. https://doi.org/10.7554/eLife.71513

19. Chen C, Qi H, Shen Y, Pickrell J, Przeworski M. Contrasting Determinants of Mutation
Rates in Germline and Soma. Genetics. 2017;207:255–67.
https://doi.org/10.1534/genetics.117.1114

20. Gao Z, Moorjani P, Sasani TA, Pedersen BS, Quinlan AR, Jorde LB, et al. Overlooked roles
of DNA damage and maternal age in generating human germline mutations. Proc Natl Acad Sci.
2019;116:9491–500. https://doi.org/10.1073/pnas.1901259116

21. Shen J-C, Rideout WM, Jones PA. The rate of hydrolytic deamination of 5-methylcytosine in
double-stranded DNA. Nucleic Acids Res. 1994;22:972–6. https://doi.org/10.1093/nar/22.6.972

22. Schmutte C, Yang AS, Beart RW, Jones PA. Base excision repair of U:G mismatches at a
mutational hotspot in the p53 gene is more efficient than base excision repair of T:G
mismatches in extracts of human colon tumors. Cancer Res. 1995;55:3742–6.

23. Tomkova M, McClellan M, Kriaucionis S, Schuster-Böckler B. DNA Replication and
associated repair pathways are involved in the mutagenesis of methylated cytosine. DNA
Repair. 2018;62:1–7. https://doi.org/10.1016/j.dnarep.2017.11.005

24. Harpak A, Bhaskar A, Pritchard JK. Mutation Rate Variation is a Primary Determinant of the
Distribution of Allele Frequencies in Humans. Eyre-Walker A, editor. PLOS Genet.
2016;12:e1006489. https://doi.org/10.1371/journal.pgen.1006489

25. Beard BC, Wilson SH, Smerdon MJ. Suppressed catalytic activity of base excision repair
enzymes on rotationally positioned uracil in nucleosomes. Proc Natl Acad Sci. 2003;100:7465–
70. https://doi.org/10.1073/pnas.1330328100

26. Sibghat-Ullah null, Day RS. DNA-substrate sequence specificity of human G:T mismatch
repair activity. Nucleic Acids Res. 1993;21:1281–7. https://doi.org/10.1093/nar/21.5.1281

27. Frederico LA, Kunkel TA, Shaw BR. A sensitive genetic assay for the detection of cytosine
deamination: determination of rate constants and the activation energy. Biochemistry.
1990;29:2532–7. https://doi.org/10.1021/bi00462a015

28. Hara R, Mo J, Sancar A. DNA Damage in the Nucleosome Core Is Refractory to Repair by
Human Excision Nuclease. Mol Cell Biol. 2000;20:9173–81.
https://doi.org/10.1128/MCB.20.24.9173-9181.2000

29. Polak P, Lawrence MS, Haugen E, Stoletzki N, Stojanov P, Thurman RE, et al. Reduced
local mutation density in regulatory DNA of cancer genomes is linked to DNA repair. Nat
Biotechnol. 2014;32:71–5. https://doi.org/10.1038/nbt.2778

30. Sabarinathan R, Mularoni L, Deu-Pons J, Gonzalez-Perez A, López-Bigas N. Nucleotide
excision repair is impaired by binding of transcription factors to DNA. Nature. 2016;532:264–7.
https://doi.org/10.1038/nature17661

31. Carlson J, Locke AE, Flickinger M, Zawistowski M, Levy S, Myers RM, et al. Extremely rare
variants reveal patterns of germline mutation rate heterogeneity in humans. Nat Commun.
2018;9:3753. https://doi.org/10.1038/s41467-018-05936-5

32. Adams CJ, Conery M, Auerbach BJ, Jensen ST, Mathieson I, Voight BF. Regularized
sequence-context mutational trees capture variation in mutation rates across the human
genome. Balding D, editor. PLOS Genet. 2023;19:e1010807.
https://doi.org/10.1371/journal.pgen.1010807

33. Seplyarskiy V, Koch EM, Lee DJ, Lichtman JS, Luan HH, Sunyaev SR. A mutation rate
model at the basepair resolution identifies the mutagenic effect of polymerase III transcription.
Nat Genet. 2023;55:2235–42. https://doi.org/10.1038/s41588-023-01562-0

34. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, et al. The mutational
constraint spectrum quantified from variation in 141,456 humans. Nature. 2020;581:434–43.
https://doi.org/10.1038/s41586-020-2308-7

35. Fairley S, Lowy-Gallego E, Perry E, Flicek P. The International Genome Sample Resource
(IGSR) collection of open human genomic variation resources. Nucleic Acids Res.
2020;48:D941–7. https://doi.org/10.1093/nar/gkz836

36. Chen X, Lin Q, Wen J, Lin W, Liang J, Huang H, et al. Whole genome bisulfite sequencing
of human spermatozoa reveals differentially methylated patterns from type 2 diabetic patients. J
Diabetes Investig. 2020;11:856–64. https://doi.org/10.1111/jdi.13201

37. Zhou Y, He F, Pu W, Gu X, Wang J, Su Z. The Impact of DNA Methylation Dynamics on the
Mutation Rate During Human Germline Development. G3 GenesGenomesGenetics.
2020;10:3337–46. https://doi.org/10.1534/g3.120.401511

38. Exome Aggregation Consortium, Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E,
et al. Analysis of protein-coding genetic variation in 60,706 humans. Nature. 2016;536:285–91.
https://doi.org/10.1038/nature19057

39. Xiang H, Zhu J, Chen Q, Dai F, Li X, Li M, et al. Single base-resolution methylome of the
silkworm reveals a sparse epigenomic map. Nat Biotechnol. 2010;28:516–20.
https://doi.org/10.1038/nbt.1626

40. Bromham L, Penny D. The modern molecular clock. Nat Rev Genet. 2003;4:216–24.
https://doi.org/10.1038/nrg1020

41. Duchêne DA, Tong KJ, Foster CSP, Duchêne S, Lanfear R, Ho SYW. Linking Branch
Lengths across Sets of Loci Provides the Highest Statistical Support for Phylogenetic Inference.
Kosakovsky Pond S, editor. Mol Biol Evol. 2020;37:1202–10.
https://doi.org/10.1093/molbev/msz291

42. Kobayashi H, Sakurai T, Miura F, Imai M, Mochiduki K, Yanagisawa E, et al. High-resolution
DNA methylome analysis of primordial germ cells identifies gender-specific reprogramming in
mice. Genome Res. 2013;23:616–27. https://doi.org/10.1101/gr.148023.112

43. Bergeron LA, Besenbacher S, Bakker J, Zheng J, Li P, Pacheco G, et al. The germline
mutational process in rhesus macaque and its implications for phylogenetic dating.
GigaScience. 2021;10:giab029. https://doi.org/10.1093/gigascience/giab029

44. Langergraber KE, Prüfer K, Rowney C, Boesch C, Crockford C, Fawcett K, et al. Generation
times in wild chimpanzees and gorillas suggest earlier divergence times in great ape and human
evolution. Proc Natl Acad Sci. 2012;109:15716–21. https://doi.org/10.1073/pnas.1211740109

45. Xue C, Raveendran M, Harris RA, Fawcett GL, Liu X, White S, et al. The population
genomics of rhesus macaques (Macaca mulatta) based on whole-genome sequences. Genome
Res. 2016;26:1651–62. https://doi.org/10.1101/gr.204255.116

46. Wroblewski EE, Murray CM, Keele BF, Schumacher-Stankey JC, Hahn BH, Pusey AE. Male
dominance rank and reproductive success in chimpanzees, Pan troglodytes schweinfurthii.
Anim Behav. 2009;77:873–85. https://doi.org/10.1016/j.anbehav.2008.12.014

47. Venn O, Turner I, Mathieson I, De Groot N, Bontrop R, McVean G. Strong male bias drives
germline mutation in chimpanzees. Science. 2014;344:1272–5.
https://doi.org/10.1126/science.344.6189.1272

48. Gao Z, Wyman MJ, Sella G, Przeworski M. Interpreting the Dependence of Mutation Rates
on Age and Time. Barton NH, editor. PLOS Biol. 2016;14:e1002355.
https://doi.org/10.1371/journal.pbio.1002355

49. Martinez G, Garcia C. Sexual selection and sperm diversity in primates. Mol Cell Endocrinol.
2020;518:110974. https://doi.org/10.1016/j.mce.2020.110974

50. Stewart AJ, Hannenhalli S, Plotkin JB. Why Transcription Factor Binding Sites Are Ten
Nucleotides Long. Genetics. 2012;192:973–85. https://doi.org/10.1534/genetics.112.143370

51. Ravichandran M, Rafalski D, Davies CI, Ortega-Recalde O, Nan X, Glanfield CR, et al.
Pronounced sequence specificity of the TET enzyme catalytic domain guides its cellular
function. Sci Adv. 2022;8:eabm2427. https://doi.org/10.1126/sciadv.abm2427

52. Maiti A, Morgan MT, Pozharski E, Drohat AC. Crystal structure of human thymine DNA
glycosylase bound to DNA elucidates sequence-specific mismatch recognition. Proc Natl Acad
Sci. 2008;105:8890–5. https://doi.org/10.1073/pnas.0711061105

53. Kawamoto M, Jouraku A, Toyoda A, Yokoi K, Minakuchi Y, Katsuma S, et al. High-quality
genome assembly of the silkworm, Bombyx mori. Insect Biochem Mol Biol. 2019;107:53–62.
https://doi.org/10.1016/j.ibmb.2019.02.002

54. Harris RA, Raveendran M, Worley KC, Rogers J. Unusual sequence characteristics of
human chromosome 19 are conserved across 11 nonhuman primates. BMC Evol Biol.
2020;20:33. https://doi.org/10.1186/s12862-020-1595-9

55. Fair BJ, Blake LE, Sarkar A, Pavlovic BJ, Cuevas C, Gilad Y. Gene expression variability in
human and chimpanzee populations share common determinants. eLife. 2020;9:e59929.
https://doi.org/10.7554/eLife.59929

56. Tong X, Han M-J, Lu K, Tai S, Liang S, Liu Y, et al. High-resolution silkworm pan-genome
provides genetic insights into artificial selection and ecological adaptation. Nat Commun.
2022;13:5619. https://doi.org/10.1038/s41467-022-33366-x

57. Johnston HR, Hu Y, Cutler DJ. Population Genetics Identifies Challenges in Analyzing Rare
Variants. Genet Epidemiol. 2015;39:145–8. https://doi.org/10.1002/gepi.21881

58. Paten B, Herrero J, Beal K, Fitzgerald S, Birney E. Enredo and Pecan: Genome-wide
mammalian consistency-based multiple alignment with paralogs. Genome Res. 2008;18:1814–
28. https://doi.org/10.1101/gr.076554.108

59. Keightley PD, Jackson BC. Inferring the Probability of the Derived vs. the Ancestral Allelic
State at a Polymorphic Site. Genetics. 2018;209:897–906.
https://doi.org/10.1534/genetics.118.301120

Supporting Information

S1 Fig. Concordance between mutation rates for 4-mer CpG contexts estimated from

gnomAD (v4) and 1000 Genomes (1KG) polymorphism datasets. A. Scaled mutation rate

estimates from regression for unmethylated and methylated sites for each 4-mer context using

1KG data. The contexts are sorted in descending order of mutability when unmethylated. **B.**

Heatmap of estimated scaled mutation rates using the 4-mer model, which assumes an

interaction between the upstream and downstream bases, at unmethylated and methylated

sites, using polymorphism data from 1KG. Each heatmap shows relative mutation rates for that

panel; color scales are not comparable across panels. **C.** Comparison of mutation rates

estimated using polymorphisms from gnomAD and 1KG at unmethylated and methylated sites,

respectively. Error bars in panel A and C represent 95% confidence intervals for the estimates.

S2 Fig. Independent effects of upstream and downstream bases on CpG mutability in

chimpanzee and rhesus macaque. A. Concordance between mutation rates as estimated

from 4-mer model and from up_1 + $down_1$ model which independently models the effects of

upstream and downstream bases in chimpanzee. **B.** Same as (A), for rhesus macaque. Error

bars represent 95% confidence intervals for the estimates on both axes.

S3 Fig. Interactions within upstream and downstream dimers are still needed to explain

mutation rate variation at CpG sites for human polymorphisms in gnomAD dataset.

Comparison of mutation rates as estimated from the $up_{21}+down_{12}$ model, which assumes

interactions between bases within the dimer upstream and downstream to the CpG and the

$up_2+up_1+down_1+down_2$ model, which assumes independent effects of every flanking base. Only

contexts for which the difference between the two models' estimates are more than 2.5 standard

deviations away from the average difference across all contexts are labelled. Error bars

represent 95% confidence intervals for the estimates on both axes.

S4 Fig. Independent effects of upstream and downstream bases on CpG mutability in the

insect species, _Bombyx mori_ (silkworm). A. Heatmap of estimated scaled mutation rates

using the 4-mer model, which assumes an interaction between the upstream and downstream

bases. Each heatmap shows relative mutation rates for that panel; color scales are not

comparable across panels. Mutation rate is modelled as a function of sequence context alone.

B. Concordance between mutation rates as estimated from 4-mer model and from up_1+down_1

model which independently models the effects of upstream and downstream. **C.** Heatmap of

scaled mutation rates from the 6-mer model which assumes all possible interactions between

flanking nucleotides for the two species. **D.** Concordance between scaled mutation rates as

estimated from 6-mer model and those estimated from $up_{21}+down_{12}$, which only assumes

interactions between bases within the dimer upstream and downstream to the CpG. Only

contexts for which the difference between the two model estimates are more than 2.5 standard

deviations away from the average difference across all contexts are labelled.

S5 Fig. Order of mutability for 4-mer CpG contexts predicted by the regression is similar

when CpG contexts are extracted from the reference genome or the ancestral genome in

humans. A. Concordance between mutation rates as estimated from 4-mer model using the

human reference genome versus the human ancestral genome inferred from the EPO 10-

primate alignment (see Methods) for identification of CpG sites and flanking sequence. **B.**

Concordance between mutation rates as estimated from 4-mer model and from up_1+down_1

model which independently models the effects of upstream and downstream bases at

unmethylated and methylated sites, using the human ancestral genome. Error bars represent

95% confidence intervals for the estimates on both axes. **C.** Heatmap of estimated scaled

mutation rates using the 4-mer model, which assumes an interaction between the upstream and

downstream bases, at unmethylated and methylated sites, using the human ancestral genome.

Each heatmap shows relative mutation rates for that panel; color scales are not comparable

across panels.

S6 Fig. Order of mutability for 4-mer CpG contexts predicted by the regression is similar

when CpG contexts are extracted from the masked or unmasked human reference

genome. A. Concordance between mutation rates as estimated from 4-mer model using the

hard-masked human reference genome versus the unmasked reference genome for

identification of CpG sites and flanking sequence. **B.** Concordance between mutation rates as

estimated from 4-mer model and from up_1+down_1 model which independently models the effects

of upstream and downstream bases at unmethylated and methylated sites, using the unmasked

human reference genome. Error bars represent 95% confidence intervals for the estimates on

both axes. **C.** Heatmap of estimated scaled mutation rates using the 4-mer model, which

assumes an interaction between the upstream and downstream bases, at unmethylated and

methylated sites, using the unmasked human reference genome. Each heatmap shows relative

mutation rates for that panel; color scales are not comparable across panels.

S1 Table. Model comparison for human and non-human species. Summary statistics of

each fitted regression model for each of the four species in our analysis. The proportion

variance explained is calculated using the difference between null and residual deviance,

standardized by the null deviance. In silkworm, the mutation rate is modeled as a function of

sequence context alone.

S2 Table. Summary of sites and SNPs for each species. Genic sites are excluded in our

analysis using NCBI *RefSeq* annotations. For human, phylogenetically conserved regions as

defined by *phastcons* are also excluded. We identify C>T SNPs by polarizing the substitution

that gave rise to each SNP using ancestral allele reconstruction based on the 10 primate EPO

for human, chimpanzee, and rhesus macaque, and *est-sfs*, a tool that infers the ancestral state

probabilities at polymorphic sites using information from a focal and outgroup species, for

silkworm.