# Regional adaptation to mosquito vectors shapes *Plasmodium falciparum* populations

Duangkamon Loesbanluechai[1], Lauriane Sollelis[2,3], Megan Armstrong[1], Inish Menezes[1], Amelia Cox[1], Lizzie Bridget Tchongwe Divala[2], Anne Lawson[1], Sabyasachi Pradhan[1], Prince Chigozirim Ubiaru[1], Antrea Pallikara[1], Dorothy Armstrong[1], Gillian Parker[2], Chiyun Lee[4], Richard D Pearson[4], Barbara H Stokes[2,3], Lisa Ranford-Cartwright[1], Matthias Marti[2,3], Virginia Howick[1,*]

[1] School of Biodiversity, One Health and Veterinary Medicine, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, Scotland, U.K.

[2] School of Infection and Immunity, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, Scotland, U.K.

[3] Institute of Parasitology Zurich, VetSuisse and Medical Faculties, University of Zurich, Zurich, Switzerland

[4] Genomic Surveillance Unit, Wellcome Sanger Institute, Hinxton, U.K.

[*] Virginia.Howick@glasgow.ac.uk

## Abstract

Transmission of *Plasmodium falciparum* through mosquitoes represents the most severe population bottleneck in the parasite's life cycle, yet the genetic basis of parasite–vector compatibility remains poorly understood. Here, we show that mosquito species–specific transmissibility depends on allelic variation in multiple *P. falciparum* genes expressed during midgut invasion, beyond the well-studied Pfs47. Using an allelic replacement strategy, we targeted highly geographically differentiated SNPs in *P. falciparum* that match regional variation in vector community composition. Transmissibility was compared across four mosquito species representing distinct geographic ranges (*An. gambiae*, *An. stephensi*, *An. minimus*, and *An. albimanus*). Two of five tested polymorphisms showed increased oocyst and sporozoite burdens in sympatric parasite–vector combinations compared to allopatric ones. Both substitutions occurred in ookinete micronemal proteins, CTRP and WARP, within von Willebrand factor A domains, suggesting that regional allelic variation modulates *Plasmodium*–vector compatibility by altering midgut adhesion interactions. These findings reveal that vector compatibility is a polygenic trait shaped by molecular interactions across several loci. Understanding this complexity refines models of parasite adaptation and can inform the design of transmission-blocking interventions effective across diverse vector-parasite combinations.

## Introduction

Malaria remains one of the world's most devastating infectious diseases, causing more than 500,000 deaths each year (*World malaria report 2024: addressing inequity in the global malaria response*, 2024). The disease is caused by *Plasmodium* parasites transmitted through the bites of *Anopheles* mosquitoes. Among these parasites, *Plasmodium falciparum* is the most virulent and is widespread across the tropics. However, the community composition of *Anopheles* vectors varies geographically and may impose strong, regionalized selective pressures on the parasite (Mackinnon and Marsh, 2010; Sinka et al., 2012; Sollelis et al., 2024).

Transmission through the mosquito represents the most severe population bottleneck in the *Plasmodium* life cycle, making it a critical target for transmission blocking interventions. After being ingested during an infectious blood meal, parasites must undergo rapid sexual differentiation, fertilization, and meiosis before forming invasive ookinetes that traverse the mosquito midgut epithelium (Bennink et al., 2016). Only a small fraction of ookinetes successfully cross this barrier: parasite numbers often drop from thousands of gametocytes to fewer than ten oocysts (Smith et al., 2014). In incompatible vectors, parasites are eliminated by the mosquito immune system or fail to exit the midgut lumen (Baton and Ranford-Cartwright, 2012; Molina-Cruz et al., 2012).

A key molecular determinant of this compatibility is the parasite antigen Pfs47, which mediates evasion of the mosquito immune response. Variants of Pfs47 act as the "key" that must match a mosquito "lock" for successful infection (Molina-Cruz et al., 2015). Population genomic analyses reveal strong geographic structuring, with near fixation of distinct Pfs47 haplotypes across major continental regions, consistent with adaptation to regional vector communities (Anthony et al., 2007; Molina-Cruz et al., 2023, 2015, 2013). However, genome-wide analyses indicate that many other *P. falciparum* genes expressed during mosquito transmission show similarly strong geographic differentiation (MalariaGEN et al., 2021; Ukegbu et al., 2020). These patterns suggest that vector compatibility may be a polygenic trait, shaped by multiple interacting loci rather than a single lock-and-key mechanism.

Here, we tested this hypothesis by generating allelic replacement lines for five geographically differentiated transmission-stage genes and assaying their infectivity across four *Anopheles* species with distinct geographic ranges. We found that for two genes, sympatric parasite–vector combinations achieved higher transmission success than allopatric pairings. Both loci encode micronemal proteins, with substitutions localized to the same adhesive functional domain, suggesting direct interactions with mosquito midgut components. These findings support a model of vector compatibility driven by coevolution between *Plasmodium* and local mosquito populations through multiple molecular interfaces.

**Results**

*Identification and allelic replacement of candidate loci*

To identify loci involved in adaptation to regional mosquito vectors, we integrated population genomic and gene expression data to pinpoint variants that may influence parasite infectivity in the mosquito midgut. The MalariaGEN network analysed genomic variation in 7,000 *P. falciparum* samples and ranked genes by a global differentiation score based on the most divergent non-synonymous SNP per gene (MalariaGEN et al., 2021). We selected the top 5% of genes (278) from this list and cross-referenced them with Malaria Cell Atlas expression data (Howick et al., 2019; Real et al., 2021; Reid et al., 2018) to identify those associated with gametocytes- or ookinete- specific gene clusters and/or were upregulated in *P. falciparum* relative to *P. berghei* during the ookinete stage (844 genes). These expression criteria enriched for genes likely involved in midgut traversal, and we hypothesized that transcriptional differences between *P. falciparum* and *P. berghei* reflect vector-specific adaptations to their respective mosquito hosts (Cohuet et al., 2006). This intersection yielded 60 candidates (**Table S1**), from which eight were selected (**Table 1**), on the basis of high expression in mature gametocytes or ookinetes, and presence of high-frequency alternative alleles in geographic regions that coincided with mosquito species available in our insectary (**Table 1, S1; Figure 1A, S1**). Six of the genes (CTRP, Pfs25, Pfs47, SOAP, WARP, P230) have established roles in transmission and are targets for potential transmission-blocking interventions (Healy et al., 2021; Jones et al., 2013; Keleta et al., 2021; Molina-Cruz and Barillas-Mury, 2022). P230, P25, and P47 are surface proteins involved in gamete interaction, midgut traversal, and immune evasion (Eksi et al., 2006; Molina-Cruz et al., 2013; Tomas et al., 2001; van Schaijk et al., 2006; Vermeulen et al., 1985). SOAP, WARP, and CTRP localize to ookinete micronemes and mediate motility and midgut invasion (Dessens et al., 2003, 1999; Li et al., 2004; Yuda et al., 2001, 1999), with CTRP knockout blocking parasite invasion of the mosquito midgut (Templeton et al., 2000). The remaining two loci, HSP101 and Pf3D7_1004600, have not previously been linked to transmission but show higher expression in *P. falciparum* ookinetes compared to *P. berghei* (Real et al., 2021). This species-specific expression suggests possible adaptation to distinct mosquito hosts. HSP101 encodes a PTEX translocon component active in asexual blood stages (Beck et al., 2014), while Pf3D7_1004600 encodes a protein of unknown function.

**Table 1. List of candidate loci.** Eight candidate genes were identified for allelic replacement. Shown are the gene ID, gene name, amino acid (AA) change, RNA expression pattern of the gene, global differentiation score with rank, and whether transfection was successful (✔) or not (✘). Gene expression data were analysed from the Malaria Cell Atlas (Real et al., 2021; Reid et al., 2018) (**Figure S1**). Global differentiation score, a gene-level *Fst*-derived metric, and rank out of the 5,561 *P. falciparum* genes are from the MalariaGEN Pf6 dataset (MalariaGEN et al., 2021).

| Gene ID | Gene name | AA change | Highest gene expression stage | Global differentiation score (rank) | Transfection (3D7) |
|---|---|---|---|---|---|
| PF3D7_1346800 | Pfs47 | S242L | Gametocyte | 1 (1) | ✔ |
| PF3D7_0315200 | CTRP | D319N | Ookinete | 0.98 (5) | ✔ |

| PF3D7_1404300 | SOAP | S95R | Ookinete | 0.62 (120) | ✘ |
|---|---|---|---|---|---|
| PF3D7_0801300 | WARP | F125L | Ookinete | 0.66 (64) | ✔ |
| PF3D7_0209000 | P230 | N1566T | Gametocyte | 0.61 (141) | ✘ |
| PF3D7_1031000 | Pfs25 | G131A | Ookinete | 0.58 (205) | ✔ |
| PF3D7_1116800 | HSP101 | R172S | Ookinete | 0.81 (7) | ✔ |
| PF3D7_1004600 | . | L94I | Ookinete | 0.67 (53) | ✘ |

**Figure 1. Distribution of *Anopheles* species and *P. falciparum* genotypes** (A) The global distributions of *An. albimanus, An. gambiae, An. minimus,* and *An. stephensi* from the Malaria Atlas Project (Hay and Snow, 2006; Sinka et al., 2012)**.** Shaded areas indicate the probability of presence is greater than 50%. (B) Alternative allele frequencies for the five candidate loci across the four broad geographic areas associated with each of the vector species: Africa (AF), South America (SAM), South Asia (SAS), Southeast Asia (SEA). Data are from (Malaria Genomic Epidemiology Network (MalariaGEN) et al., 2025). (C) UMAP of single-cell gene expression data from the Malaria Cell Atlas project (Real et al., 2021) for the five candidate genes. High expression was observed in ookinetes and/or gametocytes for each gene (IDC = Intraerythrocytic Developmental Cycle; M = Male gametocyte; F = Female gametocyte; Ook = Ookinete; Spz = Sporozoite).

To assess the role of these non-synonymous SNPs in mosquito-specific adaptation, we used CRISPR-Cas9 to generate transgenic *P. falciparum* parasites by inserting the alternative allele into the reference parasite strain 3D7, which is of African origin, to generate a single amino acid change in each gene (**Figure S2**). For each gene, we also

generated a synonymous (silent) control line that recoded the gRNA/PAM region to prevent Cas9 re-cutting while leaving the protein sequence unchanged, serving as a transgenesis-handling control for our phenotypic assays (**Table S2**). Successful genome editing was achieved for five of eight genes: CTRP*, Pfs47*, Pfs25*, WARP*, and HSP101. Attempts to edit SOAP*, P230*, and Pf3D7_1004600 using alternative guide RNAs were unsuccessful, as parasites did not recover post-transfection despite multiple efforts.

*Two genes show higher infection with sympatric parasite-allele and vector-species combinations*

Based on our hypothesis that these genes are involved in regional adaptation to vector communities, we expected that replacement of the reference allele with the alternative allele should increase infectivity in species that are sympatric for the alternative allele and decrease infectivity in allopatric ones. To test this, we performed Standard Membrane Feeding Assays (SMFAs) using five allelic-replacement lines and their silent controls across four *Anopheles* species with distinct geographic ranges (**Figure 1A**). Exflagellation rates prior to infectious feeding showed no differences between replacement and silent control lines (**Figure S3**), confirming comparable male gametocyte quality.

In total, 4,708 mosquitoes were infected across all experiments. Infection levels were assessed at the oocyst stage by measuring prevalence (proportion infected) and intensity (infection level among infected mosquitoes) (**Figure 2**). Oocysts were counted and measured 10 days post-infectious feed from dissected midguts. *An. gambiae* infected with the unmodified 3D7 clone was included as a positive control to confirm successful infection in each experimental replicate (**Figure S4**).

Significant mosquito-by-allele interactions were detected for two genes, CTRP and WARP, with sympatric combinations showing higher infection levels compared to allopatric ones. Both genes showed a mosquito-by-allele effect on oocyst prevalence ($p$ < 0.001; **Figure 2B**, **2N**). For CTRP D319N, oocyst prevalence was reduced in *An. gambiae* (allopatric; adj $p$ < 0.001) but elevated in *An. stephensi* (sympatric; adj $p$ < 0.001; **Figure 2A**). Similarly, the WARP F125L allele increased oocyst prevalence in *An. minimus* (sympatric; adj $p$ = 0.018) and decreased it in *An. gambiae* (allopatric; adj $p$ = 0.003; **Figure 2M**).

WARP also showed a significant mosquito-by-allele effect on oocyst intensity ($p$ < 0.001; **Figure 2O**), with the WARP F125L allele yielding higher infection loads in *An. minimus* (sympatric, adj $p$ = 0.047) and *An. stephensi* (sympatric, adj $p$ = 0.040) and lower loads in *An. gambiae* (allopatric, adj $p$ = 0.003) (**Figure 2M**). No significant differences in intensity were observed for CTRP D319N. The remaining alleles (HSP101 R127S, Pfs25 G131A, and Pfs47 S242L) had no measurable effects on oocyst prevalence or intensity (**Figure 2D-L**).

**Figure 2. Infection of different *Anopheles* species by five *P. falciparum* allelic-replacement lines at oocyst stage**. For each gene tested, the oocyst count data, predicted oocyst prevalence, and predicted oocyst intensity are displayed (CTRP: A-C; HSP101: D-F; Pfs25 G-I; Pfs47: J-K; WARP: M-O). The oocyst count data shows the number of oocysts for each mosquito midgut counted 10 days post infectious feed across experimental replicates. The violin plot shows all oocyst data for each species (*An. albimanus, An. gambiae, An. minimus, An. stephensi*) and each parasite allele (reference (Ref) vs alternative (Alt)). The proportion of infected mosquitoes and total number of midguts assessed is displayed below each violin.

Stars represent significant differences between the reference and alternative allele within an individual mosquito species based on post-hoc pairwise comparisons from oocyst prevalence models (stars below) or oocyst intensity models (stars above). Sympatric combinations of alternative allele and vector species are are marked with a ✓above the plot, whereas allopatric combinations are marked with an ✖ (A, D, G, J, M). The predicted prevalence plots show the estimated marginal means from the oocyst prevalence model (logistic regression) for each gene plotted by species and allele. Stars represent a significant Mosquito x Allele interaction for that locus (B, E, H, K, N). The predicted intensity plots show the estimated marginal means from the oocyst intensity model for each gene plotted by species and allele. Stars represent a significant Mosquito x Allele interaction (C, F, I, L, O). Points in predicted prevalence and intensity plots are shaped by whether the alternative allele is sympatric (triangle) or allopatric (circle) for that mosquito species. Shaded regions around each reaction norm represent the standard error.  *** $p < 0.001$, ** $p < 0.01$ * $< 0.05$.

For CTRP and WARP, an additional clone of both the allelic-replacement and silent control was tested to confirm the vector-specific transmission phenotype. We observed the same pattern of the allele in each species as seen with the primary clones (Figure 3A). We also measured oocyst diameter in a subset of the infected midguts and found no mosquito-by-allele effect on oocyst size at day 10 post-infectious feed in CTRP or WARP (**Figure 3B**) or for the other three genes tested (**Figure S5**). We then quantified sporozoite prevalence and intensity in the mosquito salivary glands at day 17 post infection using qPCR for CTRP and WARP and found a significant mosquito-by-allele interaction for WARP ($p$ = 0.034), with the alternative allele (F125L) showing lower sporozoite prevalence compared to the reference in *An. gambiae* (allopatric, adj $p$ = 0.018, **Figure 3C**). Similarly, CTRP D319N was associated with reduced sporozoite prevalence in *An. gambiae* (adj $p$ = 0.027, **Figure 3C**). No significant mosquito-by-allele interaction was found for sporozoite intensity for either gene. Surprisingly, sporozoite prevalences were generally higher than oocyst prevalences, and sporozoites were detected in *An. albimanus* infected with the WARP-F125 parasite, but oocysts were not. These differences may reflect variation in assay sensitivity or specificity, and recent work has seen the same discrepancy (higher sporozoite prevalence compared to oocysts) using the same qPCR primers, and suggests there is a higher false-positive rate in the qPCR assay (Guissou et al., 2023). Despite this discrepancy, we note the trend seen across the two genes in *An. albimanus*, where the alternative allele showed higher sporozoite prevalence in the sympatric combination compared to its reference (CTRP D319N-*An. albimanus*) but allopatric combination (WARP F125L- *An. albimanus*) resulted in a decrease in sporozoite prevalence (**Figure 3C**).

*CTRP D319N and WARP F125L both fall in von Willebrand factor A domains*

Protein structure analysis of WARP and CTRP revealed that CTRP contains six von Willebrand factor A (vWFA) domains, whereas WARP has a single vWFA domain (**Figure 3D**)(Li et al., 2004; Templeton et al., 2000; Trottein et al., 1995; Yuda et al., 2001). Both CTRP D319N and WARP F125L substitutions occur within these domains. The vWFA domains of CTRP are essential for gliding motility and onward transmission through the

mosquito (Ramakrishnan et al., 2011). By homology to integrin-like adhesion modules, these domains are predicted to mediate attachment to host surfaces through metal ion–dependent adhesion sites (MIDAS) (Pihlajamaa et al., 2013; Whittaker and Hynes, 2002).

The D319N substitution in CTRP replaces a negatively charged aspartic acid with an uncharged asparagine (**Figure 3D**). Aspartic acid residues within vWFA domains may coordinate bound cations at or near the MIDAS motif, which could stabilize ligand binding and domain conformation (Lee et al., 1995; Liddington and Ginsberg, 2002). Loss of this negative charge may weaken cation coordination and alter the structural dynamics of the vWFA domain, potentially modifying adhesion to mosquito midgut epithelial ligands or affecting downstream processes required for ookinete motility and invasion (Dessens et al., 1999; Ramakrishnan et al., 2011; Templeton et al., 2000).

Similarly, the WARP F125L substitution replaces phenylalanine with leucine, removing an aromatic side chain (**Figure 3D**). Aromatic residues often contribute to hydrophobic packing within vWFA cores and π-interactions that stabilize protein–protein interfaces (Emsley et al., 1998; Takagi et al., 2002). The loss of this aromatic interaction may subtly affect domain folding or reduce affinity for midgut-associated receptors such as basal lamina components and mosquito-derived lectins (Li et al., 2004).

Given that vWFA domains mediate critical adhesion processes, these substitutions are likely to modulate protein–protein or protein–matrix binding in the midgut. We hypothesize that such alterations could underlie the observed allele-specific differences in transmission efficiency by tuning parasite–vector compatibility at the molecular interface of midgut invasion.

**Figure 3. CTRP D319N and WARP F125L show vector-species dependent transmission levels and are in von Willebrand factor A (vWFA) domains.** (A) Oocysts per midgut are shown by clone. For CTRP, Ref E2 and Alt G9 were the primary clones tested and Ref B5 and Alt E12 were the secondary clones tested. For WARP, Ref E4 and Alt B8 were the primary clones tested, whereas Ref E5 and Alt D6 were the secondary clones tested. Each dot represents oocyst counts from a single mosquito midgut. No significant allele-by-mosquito-by-clone interactions were observed for oocyst prevalence or intensity for either gene. (B) Oocyst diameter for CTRP and WARP alleles across vector species. No significant mosquito-by-allele interaction was observed. (C) Sporozoite infection levels for CTRP and WARP alleles. For each gene, the left panel shows the the violin plot shows all sporozoite data for each species (*An. albimanus*, *An. gambiae*, *An. minimus*, *An. stephensi*) and each parasite allele (reference (Ref) vs alternative (Alt)). The proportion of infected mosquitoes and total number of midguts assessed is displayed below each violin. Stars represent significant differences between the reference and alternative allele within an individual mosquito species based on post-hoc pairwise comparisons from sporozoite prevalence models (stars below) or sporozoite intensity models (stars above) ** p <0.01, * <0.05. The right panel shows the estimated marginal means from the sporozoite prevalence model for each gene is plotted by species and allele. A star represents a

significant Mosquito x Allele interaction (p <0.05). Points are shaped by whether the alternative allele is sympatric (triangle) or allopatric (circle) for that mosquito species. Shaded regions around each reaction norm represent the standard error. (D) The protein structure of CTRP and WARP predicted by UCSF Chimera software (Pettersen et al., 2004). Distinct vWFA domains are shown in colour. The D319N substitution in CTRP reduces negative charge on this residue (negative charge from aspartic acid replaced with uncharged asparagine). The F125L substitution in WARP removes an aromatic side chain (phenylalanine replaced by leucine).

## Discussion

Understanding how spatial variation and local adaptation shape *P. falciparum* genetic diversity and transmission dynamics is fundamental to public health because treatment or intervention in one location may be ineffective in another. Here we have demonstrated how natural allelic variation in *P. falciparum* transmission-stage genes can produce vector-specific effects on parasite infectivity. By combining population genomic data with functional assays across multiple geographically diverse *Anopheles* species, we show that two variants, CTRP D319N and WARP F125L, modulate infection outcomes in a manner consistent with adaptation to sympatric vector hosts. These results provide direct experimental support for the long-standing hypothesis that *P. falciparum* is adapting to regional vector community composition. Additionally, our results challenge the lock-and-key paradigm that compatibility is determined by a single *P. falciparum* locus (Pfs47), but instead indicate that this is a complex, polygenic trait where multiple molecular interactions determine vector specialisation.

CTRP and WARP encode micronemal proteins required for ookinete motility and midgut invasion and both contain vWFA domains that are likely involved in adhesion to the midgut epithelium and contained the polymorphisms of interest. The specific amino acid changes within the vWFA domains altered the charge of the residue for CTRP and removed an aromatic ring for WARP. These changes could potentially modify the strength or the specificity of binding to mosquito midgut ligands such as laminins, lectins or glycans (Dinglasan et al., 2007; Li et al., 2004; Mahairaki et al., 2005). Future work to identify the binding partners of *P. falciparum* WARP and CTRP, and how the strength and specificity of these interactions change with different parasite alleles and sympatric vs allopatric vector midgut binding partners, could help elucidate the underlying molecular mechanisms driving the mosquito-by-allele phenotypes we observed.

We did not observe mosquito-by-allele interactions for Pfs47, HSP101, or Pfs25. We cannot rule out that these loci are also regionally adapted to different vectors, given the limited number of mosquito species tested and the fact that only one parasite genetic background was tested. Additionally, for Pfs47, previous work has shown that the full haplotype with four amino acid changes is needed to match the regional vector (*An. gambiae* vs *An. albimanus*) (Canepa et al., 2016). However, the S242L mutation in an NF54 parasite background has previously been shown to result in immune detection and parasite melanisation by the *An. gambiae* L35 resistant strain (Canepa et al., 2016). This difference from our results is likely explained by the use of different *An. gambiae* strains,

as we do not regularly observe melanisation in our *An. gambiae* Kisumu colony. For all loci tested, future investigations could employ a full haplotype replacement strategy, as we only edited a single amino acid substitution — the variant with the highest *Fst* that contributed to the global differentiation score (MalariaGEN et al., 2021). Several additional non-synonymous SNPs showed similar fixation patterns within each gene (**Figure S6**), and together these may enhance compatibility between parasite and vector (MalariaGEN et al., 2025). Additionally, future work could test multiple genetic backgrounds and combine haplotypes from different genes to understand additive and epistatic effects on vector compatibility. *P. falciparum* lineages from these distinct geographic areas are genomically very different, which is likely due to very different selective pressures beyond vector species including transmission intensity and drug-pressure (Mackinnon and Marsh, 2010; Manske et al., 2012; Rono et al., 2018). Understanding how these selective pressures together shape the parasite genome and the spread of distinct lineages, including drug-resistant *P. falciparum*, is key to understanding the epidemiology of the disease.

Together, these findings demonstrate that naturally occurring amino acid substitutions in key transmission-stage proteins can alter *P. falciparum* infectivity in a vector-dependent manner. The effects of CTRP D319N and WARP F125L across mosquito species provide experimental evidence that parasite–vector compatibility is an evolving, polygenic trait shaped by spatially variable ecological interactions. Such adaptation has important implications for the geographic structuring of malaria transmission and for the design of transmission-blocking interventions. For example, as the distribution of *An. stephensi* expands into East Africa (Emiru et al., 2023), its contribution to transmission could depend on the circulating parasite genotypes in this region. Accounting for natural allelic variation and vector diversity will be essential to developing control strategies that remain effective across the heterogeneous parasite and mosquito populations that sustain global malaria transmission.

**Materials and Methods**

*Generation of Transgenic Parasites*

The pBgC plasmids (Filarsky et al., 2018; Hitz et al., 2021) were constructed by sequentially cloning the gRNA oligo and donor template. The plasmid was first digested with BsaI, then the sgRNA oligo was inserted by In-Fusion cloning (Takara Bio), following the manufacturer's protocol. The construct was transformed into XL-10 Gold competent cells, and colonies containing the sgRNA insert were selected. A donor template (300–600 bp) containing the desired mutation along with silent mutations was synthesized and cloned into the HindIII and EcoRI sites of the plasmid using In-Fusion. After transformation, four colonies were selected for sequencing to confirm insertion of the correct donor template.

*P. falciparum* 3D7 parasites were transfected with pBgC plasmids containing the desired non-synonymous mutation or silent control for each gene (**Table S2**). Synchronized ring-stage parasites (5–10% parasitemia) were transfected with 80–100 µg of plasmid. Transfected parasites were selected with 2.5 µg/ml Blasticidin S for 8 days, then returned to normal 1640 medium supplemented with 5% human serum, 25 mM HEPES, 50 uM hypoxanthine and 0.2% sodium bicarbonate. Cultures were incubated at 37°C in a gas mixture of 1% $O_2$, 3% $CO_2$, and 96% $N_2$. Once parasites recrudesced, genomic DNA was extracted, amplified using the primers in **Table S3** and sequenced to confirm genome editing. Limiting dilution cloning was done to select clones that had the desired mutation. Bulk cultures were diluted to 0.5% prior to serial dilution in 96-well plate. After two weeks, 50 µl from each well was stained with 1:5000 SYBR Green in PBS to detect parasites by flow cytometery (MACSQuant). Clonal lines were expanded and sequenced to confirm successful editing.

*Gametocyte Induction*

Gametocytes were induced at 0.5–0.7% asexual parasitaemia and 6% hematocrit in 25 $cm^2$ flasks with 5 ml of culture following established protocols (Carter et al., 1993). Two flasks were set up for the day 17 gametocytes and three days later two flasks were set up for the day 14 gametocytes. Flasks were incubated at 37°C and media was changed daily. The media was increased to 1.5x when parasitaemia was 3-5%.

*Mosquito rearing*

*An. albimanus* PANAMA, *An. gambiae* KISUMU, *An. minimus* MINIMUS1, and *An. stephensi* SDA500 were reared for transmission experiments at 27 degrees C, 70% relative humidity and 12 hour by 12 hour light/dark cycle conditions. Larvae were fed on ground Tetramin and Tetra Pond pellets. Adults were maintained on a 5% D-glucose solution with 0.05% para-amino benzoic acid (PABA) via cotton pads placed on top of cages. *An. albimanus* were maintained on 10% glucose solution with 0.05% PABA.

*Standard Membrane Feeding Assay and Exflagellation*

The Standard Membrane Feeding Assay (SMFA) (Churcher et al., 2012; Miura et al., 2013) was performed by pooling gametocyte cultures from day 14 and day 17 flasks. The cultures were centrifuged at 500 × g for 7 minutes at 37 °C, then resuspended with an equal volume of human serum and diluted to 1% gametocytemia with 40% hematocrit. Blood meals were loaded in glass membrane feeders covered with goldbeater's skin, connected to a 37°C water bath. Mosquito pots that had 4-7 day old female mosquitoes were put under the glass feeder for 10 minutes to feed and then maintained at 27°C with 70% humidity.

Exflagellation was assessed from an aliquot of the prepared blood meal by diluting to 2.5% haematocrit in ookinete medium containing xanthurenic acid (10.4 g RPMI powder, 2 g of $NaHCO_3$ , 50 mg of hypoxanthine in 1 L of $dH_2O$ and 1 ml of 100 mM xanthurenic acid) (Delves et al., 2016). 10 ul of the dilution was added to a hemocytometer (Neubauer Improved C-Chip). After 10 minutes, exflagellation events were counted under 40X magnification for 10 minutes to estimate the number of events per µl at 1% gametocytemia.

*Oocyst and Sporozoite Quantification*

Ten days post feeding, mosquitoes were killed with 70% ethanol. Midguts were dissected in PBS and stained with 1% mercurochrome (25 µl per sample). Midguts were examined at 20X magnification on an Echo revolve microscope. The number oocysts per midgut was counted and oocyst size was measured.

Mosquitoes were frozen at day 17 post infectious feed at -20 °C for sporozoite measurement using qPCR. DNA extraction was performed on dissected mosquito heads and top of thorax using the DNeasy Blood & Tissue Kit (Qiagen). Quantitative PCR was performed by mixing 1X SYBR Green Master Mix (Applied Biosystems) and 600 nM of each primer targeting the *PfCOX* gene (forward: 5'-TTACATCAGGAATGTTATTGC-3', reverse: 5'-ATATTGGATCTCCTGCAAAT-3', total reaction 15 ul) (Boissière et al., 2013). Quantitative PCR was run on a Viia 7 system with the following cycling conditions: denaturation: 98°C and annealing: 56°C. Melting curve analysis was used to confirm product specificity. Standard curves were generated using known numbers of ring-stage parasites; 100000, 10000, 1000, 100, 10 and 1.

*Statistical Analysis*

Statistical models for oocyst and sporozoite prevalence and intensity, as well as oocyst size, had the following core structure for each gene:

$$Response\ variable \sim Allele + Mosquito + Allele\ x\ Mosquito$$

Where Allele represents the reference or alternative allele for that locus and Mosquito represents the vector species in which the oocysts or sporozoites were measured (*An. albimanus*, *An. gambiae*, *An. minimus*, or *An. stephensi*). For oocyst prevalence, logistic regression (family = binomial(link = "logit")) was fitted with a biased-reduced generalized linear model using the brglm2 package in R with Replicate as a fixed effect in the model (Kosmidis and Firth, 2021). This allowed for analysis of mosquito-allele combinations that had no oocysts detected (Ubiaru and Ranford-Cartwright, 2025). For oocyst intensity, non-zero samples were analysed using a generalized linear mixed model with a negative binomial distribution (family = nbinom2) using the glmmTMB package in R (Brooks et al., 2017). Replicate and Clone were considered as random effects for CTRP

and WARP where two clones were tested, and Replicate alone was considered a random effect for HSP101, Pfs25, and Pfs47. For sporozoite prevalence, glmmTMB was used for logistic regression again with Replicate and Clone as random effects. For sporozoite intensity, glmmTMB was used to fit a generalised linear mixed model with a Gaussian distribution with the same random factors as the sporozoite prevalence model. For oocyst size models, a linear mixed effects model was fitted using the lme4 package in R with Midgut from which the oocyst was measured as a random effect (Bates et al., 2015). Replicate was considered a fixed factor for CTRP, Pfs25, Pfs47 and WARP, but not for HSP101 as oocyst size was only measured for one of the three experimental replicates. To test the effect of clone, Clone and Clone x Allele x Mosquito were considered as fixed effects in the model. A reduced model was tested without with the Clone x Allele x Mosquito interaction and the full and reduced models were compared with an LRT test. For oocysts and sporozoite models, estimated marginal means and pairwise contrasts within species were calculated using the emmeans package in R (Searle et al., 1980).

*Vector distribution mapping*

Rasters downloaded from the Malaria Atlas Project (Hay and Snow, 2006; Sinka et al., 2012) are continuous probability of occurrence surfaces with values from 0 to 1. To produce a clear distribution footprint, surfaces were converted to a presence polygon using a fixed threshold of 0.5. Shaded areas indicate the probability of presence is higher than 50%.

*Protein structure visualisation*

Protein structures were obtained from the AlphaFold database: AF-Q8IAL0-F1 (WARP) and AF-O97267-F1 (CTRP). Domain visualization, amino acid charge analysis, and mutation modelling were performed using UCSF ChimeraX version 1.7.

## Acknowledgements

We thank Luca Nelli for creation of the map in Figure 1A and Victor Tobiasson for support with the protein structure diagram in Figure 3D. We thank Farzana Khaliq, Andy Waters, and Katarzyna Modrzynska for production of the *An. stephensi* mosquitoes. We thank Francesco Baldini, Arthur Talman, and Adam Dobson for helpful feedback on earlier versions of this manuscript. This work was funded by a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal Society (Grant 220185/Z/20/Z) and a Lord Kelvin Adam Smith Leadership Fellowship awarded to VMH.

## References

Anthony TG, Polley SD, Vogler AP, Conway DJ. 2007. Evidence of non-neutral polymorphism in Plasmodium falciparum gamete surface protein genes Pfs47 and Pfs48/45. *Mol Biochem Parasitol* **156**:117–123.
Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using **lme4**. *J Stat Softw* **67**:1–48.

Baton LA, Ranford-Cartwright LC. 2012. Ookinete destruction within the mosquito midgut lumen explains Anopheles albimanus refractoriness to Plasmodium falciparum (3D7A) oocyst infection. *Int J Parasitol* **42**:249–258.

Beck JR, Muralidharan V, Oksman A, Goldberg DE. 2014. PTEX component HSP101 mediates export of diverse malaria effectors into host erythrocytes. *Nature* **511**:592–595.

Bennink S, Kiesow MJ, Pradel G. 2016. The development of malaria parasites in the mosquito midgut. *Cell Microbiol* **18**:905–918.

Boissière A, Gimonneau G, Tchioffo MT, Abate L, Bayibeki A, Awono-Ambéné PH, Nsango SE, Morlais I. 2013. Application of a qPCR assay in the investigation of susceptibility to malaria infection of the M and S molecular forms of An. gambiae s.s. in Cameroon. *PLoS One* **8**:e54820.

Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, Skaug HJ, Mächler M, Bolker BM. 2017. glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal* **9**:378–400.

Canepa GE, Molina-Cruz A, Barillas-Mury C. 2016. Molecular analysis of Pfs47-mediated Plasmodium evasion of mosquito immunity. *PLoS One* **11**:e0168279.

Carter R, Ranford-Cartwright L, Alano P. 1993. The culture and preparation of gametocytes of Plasmodium falciparum for immunochemical, molecular, and mosquito infectivity studies. *Methods Mol Biol* **21**:67–88.

Churcher TS, Blagborough AM, Delves M, Ramakrishnan C, Kapulu MC, Williams AR, Biswas S, Da DF, Cohuet A, Sinden RE. 2012. Measuring the blockade of malaria transmission--an analysis of the Standard Membrane Feeding Assay. *Int J Parasitol* **42**:1037–1044.

Cohuet A, Osta MA, Morlais I, Awono-Ambene PH, Michel K, Simard F, Christophides GK, Fontenille D, Kafatos FC. 2006. Anopheles and Plasmodium: from laboratory models to natural systems in the field. *EMBO Rep* **7**:1285–1289.

Delves MJ, Straschil U, Ruecker A, Miguel-Blanco C, Marques S, Dufour AC, Baum J, Sinden RE. 2016. Routine in vitro culture of P. falciparum gametocytes to evaluate novel transmission-blocking interventions. *Nat Protoc* **11**:1668–1680.

Dessens JT, Beetsma AL, Dimopoulos G, Wengelnik K, Crisanti A, Kafatos FC, Sinden RE. 1999. CTRP is essential for mosquito infection by malaria ookinetes. *EMBO J* **18**:6221–6227.

Dessens JT, Sidén-Kiamos I, Mendoza J, Mahairaki V, Khater E, Vlachou D, Xu X-J, Kafatos FC, Louis C, Dimopoulos G, Sinden RE. 2003. SOAP, a novel malaria ookinete protein involved in mosquito midgut invasion and oocyst development. *Mol Microbiol* **49**:319–329.

Dinglasan RR, Alaganan A, Ghosh AK, Saito A, van Kuppevelt TH, Jacobs-Lorena M. 2007. Plasmodium falciparum ookinetes require mosquito midgut chondroitin sulfate proteoglycans for cell invasion. *Proc Natl Acad Sci U S A* **104**:15882–15887.

Eksi S, Czesny B, van Gemert G-J, Sauerwein RW, Eling W, Williamson KC. 2006. Malaria transmission-blocking antigen, Pfs230, mediates human red blood cell binding to exflagellating male parasites and oocyst production. *Mol Microbiol* **61**:991–998.

Emiru T, Getachew D, Murphy M, Sedda L, Ejigu LA, Bulto MG, Byrne I, Demisse M, Abdo M, Chali W, Elliott A, Vickers EN, Aranda-Díaz A, Alemayehu L, Behaksera SW, Jebessa G, Dinka H, Tsegaye T, Teka H, Chibsa S, Mumba P, Girma S, Hwang J,

Yoshimizu M, Sutcliffe A, Taffese HS, Bayissa GA, Zohdy S, Tongren JE, Drakeley C, Greenhouse B, Bousema T, Tadesse FG. 2023. Evidence for a role of Anopheles stephensi in the spread of drug- and diagnosis-resistant malaria in Africa. *Nat Med* **29**:3203–3211.

Emsley J, Cruz M, Handin R, Liddington R. 1998. Crystal structure of the von Willebrand Factor A1 domain and implications for the binding of platelet glycoprotein Ib. *J Biol Chem* **273**:10396–10401.

Filarsky M, Fraschka SA, Niederwieser I, Brancucci NMB, Carrington E, Carrió E, Moes S, Jenoe P, Bártfai R, Voss TS. 2018. GDV1 induces sexual commitment of malaria parasites by antagonizing HP1-dependent gene silencing. *Science* **359**:1259–1263.

Guissou E, Da DF, Hien DF de S, Yameogo KB, Yerbanga SR, Ouédraogo GA, Dabiré KR, Lefèvre T, Cohuet A. 2023. Intervention reducing malaria parasite load in vector mosquitoes: No impact on Plasmodium falciparum extrinsic incubation period and the survival of Anopheles gambiae. *PLoS Pathog* **19**:e1011084.

Hay SI, Snow RW. 2006. The malaria Atlas Project: developing global maps of malaria risk. *PLoS Med* **3**:e473.

Healy SA, Anderson C, Swihart BJ, Mwakingwe A, Gabriel EE, Decederfelt H, Hobbs CV, Rausch KM, Zhu D, Muratova O, Herrera R, Scaria PV, MacDonald NJ, Lambert LE, Zaidi I, Coelho CH, Renn JP, Wu Y, Narum DL, Duffy PE. 2021. Pfs230 yields higher malaria transmission-blocking vaccine activity than Pfs25 in humans but not mice. *J Clin Invest* **131**. doi:10.1172/JCI146221

Hitz E, Grüninger O, Passecker A, Wyss M, Scheurer C, Wittlin S, Beck H-P, Brancucci NMB, Voss TS. 2021. The catalytic subunit of Plasmodium falciparum casein kinase 2 is essential for gametocytogenesis. *Commun Biol* **4**:336.

Howick VM, Russell AJC, Andrews T, Heaton H, Reid AJ, Natarajan K, Butungi H, Metcalf T, Verzier LH, Rayner JC, Berriman M, Herren JK, Billker O, Hemberg M, Talman AM, Lawniczak MKN. 2019. The Malaria Cell Atlas: Single parasite transcriptomes across the complete Plasmodium life cycle. *Science* **365**. doi:10.1126/science.aaw2619

Jones RM, Chichester JA, Mett Vadim, Jaje J, Tottey S, Manceva S, Casta LJ, Gibbs SK, Musiychuk K, Shamloul M, Norikane J, Mett Valentina, Streatfield SJ, van de Vegte-Bolmer M, Roeffen W, Sauerwein RW, Yusibov V. 2013. A plant-produced Pfs25 VLP malaria vaccine candidate induces persistent transmission blocking antibodies against Plasmodium falciparum in immunized mice. *PLoS One* **8**:e79538.

Keleta Y, Ramelow J, Cui L, Li J. 2021. Molecular interactions between parasite and mosquito during midgut invasion as targets to block malaria transmission. *NPJ Vaccines* **6**:140.

Kosmidis I, Firth D. 2021. Jeffreys-prior penalty, finiteness and shrinkage in binomial-response generalized linear models. *Biometrika* **108**:71–82.

Lee C, Ünlü ES, White NFD, Almagro-Garcia J, Ariani C, Pearson RD. 2024. Pf-HaploAtlas: an interactive web app for spatiotemporal analysis of Plasmodium falciparum genes. *Bioinformatics* **40**:btae673.

Lee JO, Rieu P, Arnaout MA, Liddington R. 1995. Crystal structure of the A domain from the alpha subunit of integrin CR3 (CD11b/CD18). *Cell* **80**:631–638.

Li F, Templeton TJ, Popov V, Comer JE, Tsuboi T, Torii M, Vinetz JM. 2004. Plasmodium ookinete-secreted proteins secreted through a common micronemal pathway are targets of blocking malaria transmission. *J Biol Chem* **279**:26635–26644.

Liddington RC, Ginsberg MH. 2002. Integrin activation takes shape. *J Cell Biol* **158**:833–839.

Mackinnon MJ, Marsh K. 2010. The selection landscape of malaria parasites. *Science* **328**:866–871.

Mahairaki V, Voyatzi T, Sidén-Kiamos I, Louis C. 2005. The Anopheles gambiae gamma1 laminin directly binds the Plasmodium berghei circumsporozoite- and TRAP-related protein (CTRP). *Mol Biochem Parasitol* **140**:119–121.

Malaria Genomic Epidemiology Network (MalariaGEN), Abdel Hamid MM, Abdelraheem MH, Acheampong DO, Adam I, Aide P, Ajibaye O, Ali M, Almagro-Garcia J, Amambua-Ngwa A, Amenga-Etego L, Aniebo I, Aninagyei E, Ansah F, Apinjoh TO, Ariani CV, Auburn S, Awandare GA, Balmer A, Bejon P, Boene S, Bwire G, Candrinho B, Chidimatembue A, Chindavongsa K, Comiche K, Conway D, Dara A, Diakite M, Djimde A, Dondorp A, Doumbia S, Drury E, Fanello CA, Ferdig M, Figueroa K, Gamboa D, Golassa L, Gonçalves S, Guindo MDA, Hamaluba M, Hanboonkunupakarn B, Howe K, Hussien M, Imwong M, Ishengoma D, Jeans J, Kabaghe A, Kamuhabwa A, Kindermans J-M, Konate DS, Kwiatkowski DP, Lee C, Lee SK, Lee SJ, Ley B, Llanos-Cuentas A, Marfurt J, Matambisso G, Maude RR, Maude RJ, Mayor A, Mayxay M, Maïga-Ascofaré O, McCann RS, Miles A, Miotto O, Mohamed AO, Morang'a CM, Murie K, Ngasala BE, Nguyen T-N, Nolasco O, Nosten F, Noviyanti R, O'Connor Í, Oboh M, Ochola-Oyier LI, Olufunke Falade C, Olukosi A, Olumide A, Olusola FI, Onyamboko MA, Oriero EC, Oyibo WA, Pannebaker D, Pearson RD, Phiri K, van der Pluijm RW, Price RN, Quang HH, Rajkumar Devaraju V, Randrianarivelojosia M, Ranford-Cartwright L, Rayner JC, Rovira-Vallbona E, Rowlands K, Ruano-Rubio V, Sanchez JF, Saúte F, Shettima S, da Silva C, Simpson VJ, Suddaby S, Takken W, Thu AM, Toure M, Unlu E, Valdivia HO, van Vugt M, Waithira N, Wellems T, Wendler J, White N, Wuendrich Ogidan R. 2025. Pf8: an open dataset of *Plasmodium falciparum* genome variation in 33,325 worldwide samples. *Wellcome Open Res* **10**:325.

MalariaGEN, Ahouidi A, Ali M, Almagro-Garcia J, Amambua-Ngwa A, Amaratunga C, Amato R, Amenga-Etego L, Andagalu B, Anderson TJC, Andrianaranjaka V, Apinjoh T, Ariani C, Ashley EA, Auburn S, Awandare GA, Ba H, Baraka V, Barry AE, Bejon P, Bertin GI, Boni MF, Borrmann S, Bousema T, Branch O, Bull PC, Busby GBJ, Chookajorn T, Chotivanich K, Claessens A, Conway D, Craig A, D'Alessandro U, Dama S, Day NP, Denis B, Diakite M, Djimdé A, Dolecek C, Dondorp AM, Drakeley C, Drury E, Duffy P, Echeverry DF, Egwang TG, Erko B, Fairhurst RM, Faiz A, Fanello CA, Fukuda MM, Gamboa D, Ghansah A, Golassa L, Goncalves S, Hamilton WL, Harrison GLA, Hart L, Henrichs C, Hien TT, Hill CA, Hodgson A, Hubbart C, Imwong M, Ishengoma DS, Jackson SA, Jacob CG, Jeffery B, Jeffreys AE, Johnson KJ, Jyothi D, Kamaliddin C, Kamau E, Kekre M, Kluczynski K, Kochakarn T, Konaté A, Kwiatkowski DP, Kyaw MP, Lim P, Lon C, Loua KM, Maïga-Ascofaré O, Malangone C, Manske M, Marfurt J, Marsh K, Mayxay M, Miles A, Miotto O, Mobegi V, Mokuolu OA, Montgomery J, Mueller I, Newton PN, Nguyen T, Nguyen T-N, Noedl H, Nosten F, Noviyanti R, Nzila A, Ochola-Oyier LI, Ocholla H, Oduro A, Omedo I, Onyamboko MA, Ouedraogo J-B, Oyebola K, Pearson RD, Peshu N, Phyo AP, Plowe CV, Price

RN, Pukrittayakamee S, Randrianarivelojosia M, Rayner JC, Ringwald P, Rockett KA, Rowlands K, Ruiz L, Saunders D, Shayo A, Siba P, Simpson VJ, Stalker J, Su X-Z, Sutherland C, Takala-Harrison S, Tavul L, Thathy V, Tshefu A, Verra F, Vinetz J, Wellems TE, Wendler J, White NJ, Wright I, Yavo W, Ye H. 2021. An open dataset of Plasmodium falciparum genome variation in 7,000 worldwide samples. *Wellcome Open Res* **6**:42.

Manske M, Miotto O, Campino S, Auburn S, Almagro-Garcia J, Maslen G, O'Brien J, Djimde A, Doumbo O, Zongo I, Ouedraogo J-B, Michon P, Mueller I, Siba P, Nzila A, Borrmann S, Kiara SM, Marsh K, Jiang H, Su X-Z, Amaratunga C, Fairhurst R, Socheat D, Nosten F, Imwong M, White NJ, Sanders M, Anastasi E, Alcock D, Drury E, Oyola S, Quail MA, Turner DJ, Ruano-Rubio V, Jyothi D, Amenga-Etego L, Hubbart C, Jeffreys A, Rowlands K, Sutherland C, Roper C, Mangano V, Modiano D, Tan JC, Ferdig MT, Amambua-Ngwa A, Conway DJ, Takala-Harrison S, Plowe CV, Rayner JC, Rockett KA, Clark TG, Newbold CI, Berriman M, MacInnis B, Kwiatkowski DP. 2012. Analysis of Plasmodium falciparum diversity in natural infections by deep sequencing. *Nature* **487**:375–379.

Miura K, Deng B, Tullo G, Diouf A, Moretz SE, Locke E, Morin M, Fay MP, Long CA. 2013. Qualification of Standard Membrane-Feeding Assay with Plasmodium falciparum Malaria and Potential Improvements for Future Assays. *PLoS One* **8**:e57909.

Molina-Cruz A, Barillas-Mury C. 2022. Pfs47 as a malaria transmission-blocking vaccine target. *Am J Trop Med Hyg* **107**:27–31.

Molina-Cruz A, Canepa GE, Dwivedi A, Liu W, Raytselis N, Antonio-Nkondjio C, Hahn BH, Silva JC, Barillas-Mury C. 2023. Role of *Pfs47* in the dispersal of ancestral *Plasmodium falciparum* malaria through adaptation to different anopheline vectors. *Proc Natl Acad Sci U S A* **120**:e2213626120.

Molina-Cruz A, Canepa GE, Kamath N, Pavlovic NV, Mu J, Ramphul UN, Ramirez JL, Barillas-Mury C. 2015. Plasmodium evasion of mosquito immunity and global malaria transmission: The lock-and-key theory. *Proc Natl Acad Sci U S A* **112**. doi:10.1073/pnas.1520426112

Molina-Cruz A, DeJong RJ, Ortega C, Haile A, Abban E, Rodrigues J, Jaramillo-Gutierrez G, Barillas-Mury C. 2012. Some strains of Plasmodium falciparum, a human malaria parasite, evade the complement-like system of Anopheles gambiae mosquitoes. *Proc Natl Acad Sci U S A* **109**:E1957-62.

Molina-Cruz A, Garver LS, Alabaster A, Bangiolo L, Haile A, Winikor J, Ortega C, van Schaijk BCL, Sauerwein RW, Taylor-Salmon E, Barillas-Mury C. 2013. The human malaria parasite Pfs47 gene mediates evasion of the mosquito immune system. *Science* **340**:984–987.

Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. 2004. UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* **25**:1605–1612.

Pihlajamaa T, Kajander T, Knuuti J, Horkka K, Sharma A, Permi P. 2013. Structure of Plasmodium falciparum TRAP (thrombospondin-related anonymous protein) A domain highlights distinct features in apicomplexan von Willebrand factor A homologues. *Biochem J* **450**:469–476.

Ramakrishnan C, Dessens JT, Armson R, Pinto SB, Talman AM, Blagborough AM, Sinden RE. 2011. Vital functions of the malarial ookinete protein, CTRP, reside in the A domains. *Int J Parasitol* **41**:1029–1039.

Real E, Howick VM, Dahalan FA, Witmer K, Cudini J, Andradi-Brown C, Blight J, Davidson MS, Dogga SK, Reid AJ, Baum J, Lawniczak MKN. 2021. A single-cell atlas of Plasmodium falciparum transmission through the mosquito. *Nat Commun* **12**:3196.

Reid AJ, Talman AM, Bennett HM, Gomes AR, Sanders MJ, Illingworth CJR, Billker O, Berriman M, Lawniczak MKN. 2018. Single-cell RNA-seq reveals hidden transcriptional variation in malaria parasites. *Elife* **7**:e33105.

Rono MK, Nyonda MA, Simam JJ, Ngoi JM, Mok S, Kortok MM, Abdullah AS, Elfaki MM, Waitumbi JN, El-Hassan IM, Marsh K, Bozdech Z, Mackinnon MJ. 2018. Adaptation of Plasmodium falciparum to its transmission environment. *Nature Ecology & Evolution* **2**:377–387.

Searle SR, Speed FM, Milliken GA. 1980. Population marginal means in the linear model: An alternative to least squares means. *Am Stat* **34**:216–221.

Sinka ME, Bangs MJ, Manguin S, Rubio-Palis Y, Chareonviriyaphap T, Coetzee M, Mbogo CM, Hemingway J, Patil AP, Temperley WH, Gething PW, Kabaria CW, Burkot TR, Harbach RE, Hay SI. 2012. A global map of dominant malaria vectors. *Parasit Vectors* **5**:69.

Smith RC, Vega-Rodríguez J, Jacobs-Lorena M. 2014. The Plasmodium bottleneck: malaria parasite losses in the mosquito vector. *Mem Inst Oswaldo Cruz* **109**:644–661.

Sollelis L, Howick VM, Marti M. 2024. Revisiting the determinants of malaria transmission. *Trends Parasitol* **40**:302–312.

Takagi J, Petre BM, Walz T, Springer TA. 2002. Global conformational rearrangements in integrin extracellular domains in outside-in and inside-out signaling. *Cell* **110**:599–511.

Templeton TJ, Kaslow DC, Fidock DA. 2000. Developmental arrest of the human malaria parasite Plasmodium falciparum within the mosquito midgut via CTRP gene disruption. *Mol Microbiol* **36**:1–9.

Tomas AM, Margos G, Dimopoulos G, van Lin LH, de Koning-Ward TF, Sinha R, Lupetti P, Beetsma AL, Rodriguez MC, Karras M, Hager A, Mendoza J, Butcher GA, Kafatos F, Janse CJ, Waters AP, Sinden RE. 2001. P25 and P28 proteins of the malaria ookinete surface have multiple and partially redundant functions. *EMBO J* **20**:3975–3983.

Trottein F, Triglia T, Cowman AF. 1995. Molecular cloning of a gene from Plasmodium falciparum that codes for a protein sharing motifs found in adhesive molecules from mammals and plasmodia. *Mol Biochem Parasitol* **74**:129–141.

Ubiaru PC, Ranford-Cartwright LC. 2025. When a mean can be meaningless: evaluating mosquito infections with Plasmodium parasites. *Parasitology* 1–9.

Ukegbu CV, Giorgalli M, Tapanelli S, Rona LDP, Jaye A, Wyer C, Angrisano F, Blagborough AM, Christophides GK, Vlachou D. 2020. PIMMS43 is required for malaria parasite immune evasion and sporogonic development in the mosquito vector. *Proc Natl Acad Sci U S A* **117**:7363–7373.

van Schaijk BCL, van Dijk MR, van de Vegte-Bolmer M, van Gemert G-J, van Dooren MW, Eksi S, Roeffen WFG, Janse CJ, Waters AP, Sauerwein RW. 2006. Pfs47, paralog of the male fertility factor Pfs48/45, is a female specific surface protein in Plasmodium falciparum. *Mol Biochem Parasitol* **149**:216–222.

Vermeulen AN, Ponnudurai T, Beckers PJ, Verhave JP, Smits MA, Meuwissen JH. 1985. Sequential expression of antigens on sexual stages of Plasmodium falciparum accessible to transmission-blocking antibodies in the mosquito. *J Exp Med* **162**:1460–1476.

Whittaker CA, Hynes RO. 2002. Distribution and evolution of von Willebrand/integrin A domains: widely dispersed domains with roles in cell adhesion and elsewhere. *Mol Biol Cell* **13**:3369–3387.

World malaria report 2024: addressing inequity in the global malaria response. 2024. . Geneva: World Health Organization.

Yuda M, Sakaida H, Chinzei Y. 1999. Targeted disruption of the plasmodium berghei CTRP gene reveals its essential role in malaria infection of the vector mosquito. *J Exp Med* **190**:1711–1716.

Yuda M, Yano K, Tsuboi T, Torii M, Chinzei Y. 2001. von Willebrand Factor A domain-related protein, a novel microneme protein of the malaria ookinete highly conserved throughout Plasmodium parasites. *Mol Biochem Parasitol* **116**:65–72.

**Supplementary Information**

**Supplementary Figures**

Figure S1. Expression of all candidate genes in Malaria Cell Atlas data

Figure S2. Genome and Protein alignment for candidate loci

Figure S3. Exflagellation from five allelic-replacement parasite lines and silent controls

Figure S4. Level of 3D7 infection in *An. gambiae* for each experimental replicate

Figure S5. Oocyst size across parasite lines and vector species for HSP101, Pfs25, Pfs47

Figure S6. CTRP and WARP haplotypes

**Supplementary Tables**

Table S1. List of 60 candidates with ranks and gene expression data

Table S2. Donor template sequence for plasmid construct of mutation and silene control

Table S3. Guide oligonucleotide and primer sequences.

**Supplementary Figures**

**Figure S1. Expression of candidate genes in Malaria Cell Atlas data**

Expression of the eight candidate genes across Malaria Cell Altas Smart-Seq2 *Plasmodium falciparum* data from (Real et al., 2021; Reid et al., 2018). Expression is shown for each gene in each stage (A) based on annotated clusters (B). Candidate genes showed high expression in the mature gametocytes or ookinete stages. Annotated clusters are ordered by developmental progression. Asex = asexual, mg = male gametocyte, fg = female gametocyte, ook=ookinete, and spz = sporozoite. Targeted candidate genes were selected from the intersection of genes in the top 5% of global differentiation scores from MalariaGEN data (MalariaGEN et al., 2021) and genes in the Malaria Cell Atlas data that either fell in gene clusters 7-12 or were more highly expressed in *P. falciparum* ookinetes compared to *P. berghei* (Table S1). Gene clusters 7-12 (g7-g12) are associated with high expression in *P. falciparum* gametocytes or ookinetes (Real et al., 2021).

```
WARP_F125L_chr8:102454
Genomic
PF3D7_0801300          GAATCTATGGTCTTTAATTCGAATGTAAGTAAGGATAAGGCTCATATGTCAGTTGTCCTT          420
PfHB3_080006400        GAATCTATGGTCTTTAATTCGAATGTAAGTAAGGATAAGGCTCATATGTCAGTTGTCCTT          420
Pf7G8_080006600        GAATCTATGGTCTTTAATTCGAATGTAAGTAAGGATAAGGCTCATATGTCAGTTGTCCTT          420
PfDd2_080006700        GAATCTATGGTCCTTAATTCGAATGTAAGTAAGGATAAGGCTCATATGTCAGTTGTCCTT          420
                       *********** *************************************************

Protein
PF3D7_0801300.1-p1         ESMVFNSNVSKDKAHMSVVLFAGKQRVVVPFTDESSQDKNMLLEKINSIDEGIATSSDTL      180
Pf7G8_080006600-t41_1-p1   ESMVFNSNVSKDKAHMSVVLFAGKQRVLVPFTDESSQDKNMLLEKINSIDEGIATSSDTL      180
PfHB3_080006400-t41_1-p1   ESMVFNSNVSKDKAHMSVVLFAGKQRVLVPFTDESSQDKNMLLEKINSIDEGIATSSDTL      180
PfDd2_080006700-t41_1-p1   ESMVLNSNVSKDKAHMSVVLFAGKQRVFVPFTDESSQDKNMLLEKINSIDEGIATSSDTL      180
                           ****:***************.*** .:.***************.****************

CTRP_D319N_chr3:617426
Genomic
PfHB3_030019000        GAGGATTATTACGATTTAACTTTAATTTTAGATGAGTCACGAAGTATAACTCTCAACAAA          960
PF3D7_0315200          GAGGATTATTACGATTTAACTTTAATTTTAGATGAGTCACGAAGTATAACTCTCGACAAA          960
Pf7G8_030021000        GAGGATTATTACGATTTAACTTTAATTTTAGATGAGTCACGAAGTATAACTCTCAACAAA          960
PfDd2_030020200        GAGGATTATTACGATTTAACTTTAATTTTAGATGAGTCACGAAGTATAACTCTCAACAAA          960
                       ******************************************************* *****

Protein
PfHB3_030019000-t41_1-p1   EDYYDLTLILDESRSITLNKWKKDVVPFAEKVLNNLNIDKDKIHVGIMRFAKSMKTDIGY      360
PF3D7_0315200.1-p1         EDYYDLTLILDESRSITLDKWKKDVVPFAEKVLNNLNIDKDKIHVGIMRFAKSMKTDIGY      360
Pf7G8_030021000-t41_1-p1   EDYYDLTLILDESRSITLNKWKKDVVPFAEKVLNNLNIDKDKIHVGIMRFAKSMKTDIGY      360
PfDd2_030020200-t41_1-p1   EDYYDLTLILDESRSITLNKWKKDVVPFAEKVLNNLNIDKDKIHVGIMRFAKSMKTDIGY      360
                           ******************:*****************************************

HSP101_R172S_chr11:638298
Genomic (intron included)
Pf7G8_110020400          GATGAAGCCATTAAAGCAATATTAAAAAGCAAATTTGAAAAAACGAAAAAGGATAAAGAT      1221
PfHB3_110019900          GATGAAGCCATTAAAGCAATATTAAAAAGAAAATTTGAAAAAACGAAAAAGGATAAAGAT      1253
PF3D7_1116800            GATGAAGCCATTAAAGCAATATTAAAAAGAAAATTTGAAAAAACGAAAAAGGATAAAGAT      1243
PfDd2_110019800          GATGAAGCCATTAAAGCAATATTAAAAAGCAAATTTGAAAAAACGAAAAAGGATAAAGAT      1236
                         **************************** *******************************

Protein
PF3D7_1116800.1-p1         TLALAEAAANKYKSPKVDVEHLLSGLSNDELVNEIFNEVYLTDEAIKAILKRKFEKTKKD      180
PfHB3_110019900-t41_1-p1   TLALAEAAANKYKSPKVDVEHLLSGLSNDELVNEIFNEVYLTDEAIKAILKRKFEKTKKD      180
Pf7G8_110020400-t41_1-p1   TLALAEAAANKYKSPKVDVEHLLSGLSNDELVNEIFNEVYLTDEAIKAILKSKFEKTKKD      180
PfDd2_110019800-t41_1-p1   TLALAEAAANKYKSPKVDVEHLLSGLSNDELVNEIFNEVYLTDEAIKAILKSKFEKTKKD      180
                           **************************************************** ********

P47_S242L_chr13:1879470
Genomic
PfDd2_130052600        TTATCAAATAATAATGTGGTTTTAAAAATGCCTTCCATATTTATAAATAATATTAATTGT          780
Pf7G8_130051200        TTATTAAATAATAATGTGGCTTTAAAAATGCCTTCCATATTTATAAATAATATTAATTGT          780
PF3D7_1346800          TTATCAAATAATAATGTGGTTATAAAAATGCCTTCCATATTTATAAATAATATTAATTGT          780
PfHB3_130053100        TTATTAAATAATAATGTGGCTTTAAAAATGCCTTCCATATTTATAAATAATATTAATTGT          780
                       **** *************** *  *************************************

Protein
PfDd2_130052600-t41_1-p1   LSNNNVVLKMPSIFINNINCMLSSQDQNNEKYYINLKADKTKHIDGCDFTKPKGKGIYKN      300
PF3D7_1346800.1-p1         LSNNNVVIKMPSIFINNINCMLSSQDQNNEKNYINLKADKTKHIDGCDFTKPKGKGIYKN      300
Pf7G8_130051200-t41_1-p1   LLNNNVALKMPSIFINNINCMLSSQDQNNEKNYINLKADKTKHIDGCDFTKPKGKGIYKN      300
PfHB3_130053100-t41_1-p1   LLNNNVALKMPSIFINNINCMLSSQDQNNEKNYINLKADKTKHIDGCDFTKPKGKGIYKN      300
                           * ****.:********************** *****************************

Pfs25_G131A_chr10:1253677
Genomic
PF3D7_1031000          ATATTAGATACAAGCAATCCTGTTAAAACTGGAGTTTGCTCATGTAATATAGGCAAAGTT          420
Pf7G8_100035300        ATATTAGATACAAGCAATCCTGTTAAAACTGGAGTTTGCTCATGTAATATAGGCAAAGTT          420
PfHB3_100035300        ATATTAGATACAAGCAATCCTGTTAAAACTGGAGTTTGCTCATGTAATATAGGCAAAGTT          420
PfDd2_100036200        ATATTAGATACAAGCAATCCTGTTAAAACTGCAGTTTGCTCATGTAATATAGGCAAAGTT          420
                       ******************************* ****************************

Protein
PF3D7_1031000.1-p1         ILDTSNPVKTGVCSCNIGKVPNVQDQNKCSKDGETKCSLKCLKENETCKAVDGIYKCDCK      180
Pf7G8_100035300-t41_1-p1   ILDTSNPVKTGVCSCNIGKVPNVQDQNKCSKDGETKCSLKCLKENETCKAVDGIYKCDCK      180
PfHB3_100035300-t41_1-p1   ILDTSNPVKTGVCSCNIGKVPNVQDQNKCSKDGETKCSLKCLKENETCKAVDGIYKCDCK      180
PfDd2_100036200-t41_1-p1   ILDTSNPVKTAVCSCNIGKVPNVQDQNKCSKDGETKCSLKCLKENETCKAVDGIYKCDCK      180
                           **********.*************************************************

P230_N1566T_chr10:375134-375135
Genomic
PfDd2_020011800        TTAAATATAAAATTTGATACATATGTTACTGGCACTGATCAAAATAAATATCTTATGACA          4740
Pf7G8_020013700        TTAAATATAAAATTTGATAATTATGTTACTGGCACTGATCAAAATAAATATCTTATGACA          4737
PF3D7_0209000          TTAAATATAAAATTTGATAATTATGTTACTGGCACTGATCAAAATAAATATCTTATGACA          4737
PfHB3_020013700        TTAAATATAAAATTTGATAATTATGTTACTGGCACTGATCAAAATAAATATCTTATGACA          4725
                       ****************** *****************************************

Protein
PfDd2_020011800-t41_1-p1   LNIKFDTYVTGTDQNKYLMTYMDLHLSHKRNYLKELFHDLGKKKPADTDANPESIIESLS      1620
Pf7G8_020013700-t41_1-p1   LNIKFDNYVTGTDQNKYLMTYMDLHLSHKRNYLKELFHDLGKKKPADTDANPESIIESLS      1619
PF3D7_0209000.1-p1         LNIKFDNYVTGTDQNKYLMTYMDLHLSHKRNYLKELFHDLGKKKPADTDANPESIIESLS      1619
PfHB3_020013700-t41_1-p1   LNIKFDNYVTGTDQNKYLMTYMDLHLSHKRNYLKELFHDLGKKKPADTDANPESIIESLS      1615
                           ******.****************************************************
```

```
 ______________
SOAP_S95R_chr2:150282
Genomic
PF3D7_1404300        TCAATGAAATATAGACATGCCAGTAAGGGATCTAAAGGATATAGTAAAAAAATGACAAAG        300
Pf7G8_140009700      TCAATGAAATATAGACATGCCAGTAAGGGATCTAAAAGATATAGTAAAAAAATGACAAAG        300
PfHB3_140010200      TCAATGAAATATAGACATGCCAGTAAGGGATCTAAAAGATATAGTAAAAAAATGACAAAG        300
PfDd2_140008900      TCAATGAAATATAGACATGCCAGTAAGGGATCTAAAAGATATCGTAAAAAAATGACAAAG        300
                     ********************************** *****  ****************

Protein
PF3D7_1404300.1-p1         KCLPDNFCECECSCKNKTGFSMKYRHASKGSKGYSKKMTKTSSRSRQCNDLPLEKSEQET   120
Pf7G8_140009700-t41_1-p1   KCLPDNFCECECSCKNKTGFSMKYRHASKGSKRYSKKMTKTSSRSRQCNDLPLEKSEQET   120
PfHB3_140010200-t41_1-p1   KCLPDNFCECECSCKNKTGFSMKYRHASKGSKRYSKKMTKTSSRSRQCNDLPLEKSEQET   120
PfDd2_140008900-t41_1-p1   KCLPDNFCECECSCKNKTGFSMKYRHASKGSKRYRKKMTKTSSRSRQCNDLPLEKSEQET   120
                           ********************************** * ************************

 ______________
Unknown protein_L94I_chr10:216319
Genomic
PF3D7_1004600        AATATTCATGGAACATTATTTATTATAAGTTTAATTTGTTATTCTCTTGAAAAAATTATT        616
Pf7G8_100009100      AATATTCATGGAACATTATTTATTATAAGTTTAATTTGTTATTCTCTTGAAAAAATTATT        612
PfDd2_100010000      AATATTCATGGAACATTATTTATTATAAGTTTAATTTGTTATTCTATTGAAAAAATTATT        498
PfHB3_100009100      AATATTCATGGAACATTATTTATTATAAGTTTAATTTGTTATTCTCTTGAAAAAATTATT        660
                     *********************************************  **************

Protein
PF3D7_1004600.1-p1         LFICSGLKEDENHGNITTNIHGTLFIISLICYSLEKIIPFSFCSDFLFIFLLSLSCTLPF   120
PfHB3_100009100-t41_1-p1   LFICSGLKEDENHGNITTNIHGTLFIISLICYSLEKIIPFSFCSDFLFIFLLSLSCTLPF   120
Pf7G8_100009100-t41_1-p1   LFICSGLKEDENHGNITTNIHGTLFIISLICYSLEKIIPFSFCSDFLFIFLLSLSCTLPF   120
PfDd2_100010000-t41_1-p1   LFICSGLKEDENHGNITTNIHGTLFIISLICYSIEKIIPFSFCSDFLFIFLLSLSCTLPF   120
                           *********************************:**************************
```

**Figure S2. Genome and protein alignments for candidate genes.** Non-synonymous SNPs identified in each gene selected for this study in four *P. falciparum* genomes from Africa (PF3D7), America (PfHB3 and Pf 7G8) and Asia (PfDd2). Targeted amino acids are highlighted in green within the protein sequence alignment, while blue indicates the genomic variants.

**Figure S3 Exflagellation from five allelic-replacement parasite lines and silent controls**

Exflagellation was measured for each infectious blood meal. Each point represents a single replicate, and the black bar represents the mean. No difference was observed between reference and alternative alleles in any of the five candidate loci (Linear model: Exflagellation centers ~ Allele, $p > 0.05$ for all loci).

Figure S4. **Oocyst prevalence for wild-type 3D7 in *An. gambiae* in each experimental replicate.** An. gambiae infected with unedited wild-type 3D7 was included as a control in all experimental replicates to confirm that infection was achieved. The number above each bar represents the number of midguts examined.

**Figure S5. Oocyst diameter across parasite lines and vector species**. No significant mosquito-by-allele interaction was found for any locus. However, for HSP101, the alternative allele had larger oocysts for *An. minimus (*adj *p* = 0.003) and *An. stephensi* (adj *p* = 0.040). Only a single replicate was measured for HSP101 compared to two or three replicates for all other loci.

**Figure S6 CTRP and WARP haplotypes.** Images were created using the MalariaGEN Pf-HaploAtlas (Lee et al., 2024). See https://pf-haploatlas.streamlit.app/?gene_id=PF3D7_0315200 and https://pf-haploatlas.streamlit.app/?gene_id=PF3D7_0801300 for more information. Legend terms: OC-NG = Oceania, New Guinea; AS-SE-E = Southeast Asia (East); AS-SE-W (Southeast Asia (West)); AS-S-FE = South Asia (Far East); AS-S-E = South Asia (East); AF-E = Africa (East); AF-NE = Africa (North-East); AF-C = Africa (Central); AF-W = Africa (West); SA = South America.

Table S1. 60 Candidate genes

See sup file

Table S2. Donor template sequences for mutant and silent control plasmids

See sup file

Table S3. Guide oligonucleotide and primer sequences. The guide oligos were cloned into a plasmid construct to facilitate targeted gene editing, and the primer sets were employed to confirm the successful insertion of the SNP.

| Primer name | Sequence | Purpose |
|---|---|---|
| Pf1004600_L94I_g1F | TAAGTATATAATATTAAATGGTAACGTACATGATAGTTTTAGAGCTAGAA | gRNA oligo primer |
| Pf1004600_L94I_g1R | TTCTAGCTCTAAAACTATCATGTACGTTACCATTTAATATTATATACTTA | gRNA oligo primer |
| Pf1004600_L94I_g2F | TAAGTATATAATATTTATGTTCATAAAACACAAGCGTTTTAGAGCTAGAA | gRNA oligo primer |
| Pf1004600_L94I_g2R | TTCTAGCTCTAAAACGCTTGTGTTTTATGAACATAAATATTATATACTTA | gRNA oligo primer |
| PfP230_N1566T_g2F | TAAGTATATAATTATGTTCTGCATCTACATCCCGTTTTAGAGCTAGAA | gRNA oligo primer |
| PfP230_N1566T_g2R | TTCTAGCTCTAAAACGGGATGTAGATGCAGAACATAATATTATATACTTA | gRNA oligo primer |
| PfSOAP_S95R_g1F | TAAGTATATAATATTGAAATATAGACATGCCAGTAGTTTTAGAGCTAGAA | gRNA oligo primer |
| PfSOAP_S95R_g1R | TTCTAGCTCTAAAACTACTGGCATGTCTATATTTCAATATTATATACTTA | gRNA oligo primer |
| PfHSP101_R172S_g1F | TAAGTATATAATATTTATTGCTTTAATGGCTTCATGTTTTAGAGCTAGAA | gRNA oligo primer |
| PfHSP101_R172S_g1R | TTCTAGCTCTAAAACATGAAGCCATTAAAGCAATAAATATTATATACTTA | gRNA oligo primer |
| PfP47_S242L_g1F | TAAGTATATAATATTGAAGAAACTATTGTAGAATCGTTTTAGAGCTAGAA | gRNA oligo primer |
| PfP47_S242L_g1R | TTCTAGCTCTAAAACGATTCTACAATAGTTTCTTCAATATTATATACTTA | gRNA oligo primer |
| PfWARP_F125L_g1F | TAAGTATATAATATTAGAGTCAAGTTTTGTTCAGAGTTTTAGAGCTAGAA | gRNA oligo primer |
| PfWARP_F125L_g1R | TTCTAGCTCTAAAACTCTGAACAAAACTTGACTCTAATATTATATACTTA | gRNA oligo primer |
| PfCTRP_D319N_g1F | TAAGTATATAATATTGAAGTATAACTCTCGACAAAGTTTTAGAGCTAGAA | gRNA oligo primer |
| PfCTRP_D319N_g1R | TTCTAGCTCTAAAACTTTGTCGAGAGTTATACTTCAATATTATATACTTA | gRNA oligo primer |
| Pfs25_G131A_g1F | TAAGTATATAATATTTAAGCATTTTAATGAGCATTGTTTTAGAGCTAGAA | gRNA oligo primer |
| Pfs25_G131A_g1R | TTCTAGCTCTAAAACAATGCTCATTAAAATGCTTAAATATTATATACTTA | gRNA oligo primer |
| For1_out_HSP101 | CAAACAAGAGCGGGAGCGgtatg | primer for testing the insertion of SNP in genome |
| Rev1_out_HSP101 | CGGTCATATCTACTGAAGGAGG | primer for testing the insertion of SNP in genome |
| For1_out_P47 | TGATTCCAGCGCATATCATTTAG | primer for testing the insertion of SNP in genome |
| Rev1_out_P47 | GCGATATGTAATTCCATTACTGC | primer for testing the insertion of SNP in genome |
| For1_out_WARP | CCCGTTTTATCTTTATTATTCG | primer for testing the insertion of SNP in genome |
| Rev1_out_WARP | GTTCAGTCTTGTTCATACATATTG | primer for testing the insertion of SNP in genome |
| For1_out_CTRP | GAGATTACGTCACTTATGGTGATG | primer for testing the insertion of SNP in genome |
| Rev1_out_CTRP | GATCCCGTATTACATATCTTATCCG | primer for testing the insertion of SNP in genome |
| For2_out_WARP | GTTGTTGTATAATAAGAGAGAG | primer for testing the insertion of SNP in genome |
| Rev2_out_WARP | CATCTTATGATTTATTCTTATCAC | primer for testing the insertion of SNP in genome |
| For2_out_HSP101 | gCCATTGGATAATAAAAGCAAAATTG | primer for testing the insertion of SNP in genome |
| Rev2_out_HSP101 | CTTCTTTCGAAAGCTGAACAAC | primer for testing the insertion of SNP in genome |