Unified modeling of cellular responses to diverse perturbation types

Chen Li[1], Lei Wei[1*] and Xuegong Zhang[1,2*]

1 MOE Key Laboratory of Bioinformatics and Bioinformatics Division of BNRIST, Department of
Automation, Tsinghua University, Beijing 100084, China

2 Center for Synthetic and Systems Biology, School of Life Sciences and School of Medicine, Tsinghua
University, Beijing 100084, China

* To whom correspondence should be addressed. Email: zhangxg@tsinghua.edu.cn;
weilei92@tsinghua.edu.cn

Abstract

Predicting cellular responses to perturbations is essential for understanding gene regulation and advancing
drug development. Most existing *in silico* perturbation models treat perturbation–cell interactions with
simplistic fusion strategies that overlook the hierarchical nature of regulatory processes, leading to inaccurate
predictions and poor generalization across perturbation types. We present X-Pert, a universal *in silico*
perturbation model that explicitly captures both gene–perturbation interactions and gene–gene dependencies
through attention mechanisms. X-Pert flexibly handles diverse perturbation inputs across different types and
combinations, and quantitatively models their dosage- and efficacy-dependent effects, all within a unified
representation space. This enables accurate prediction of unseen, combinatorial, and dose- or efficacy-
dependent responses of various perturbation types. Across benchmarks spanning genetic and chemical
perturbations, X-Pert consistently demonstrates superior performance at both gene and pathway levels.
Moreover, its unified representation space enables downstream analyses such as perturbation retrieval and
drug–gene association discovery. By integrating data across perturbation types, experimental platforms, and
cell contexts, X-Pert establishes a versatile and generalizable foundation for *in silico* perturbation, enabling
broad applications in biological and therapeutic discovery.

Introduction

In silico perturbation aims to predict gene expression changes induced by genetic[1-7], chemical[8-12], or protein-based[13-16] interventions, expanding our ability to explore perturbation outcomes at scale and supporting mechanistic discovery, cellular engineering, and drug development. A variety of computational methods have been developed for *in silico* perturbation[17-30], including models based on graph neural networks (GNNs)[17, 18], variational autoencoders (VAEs)[22-24], and transformers[19, 20]. While these approaches aim to capture global expression responses, they fail to reflect how perturbations propagate through hierarchical regulatory cascades that progressively reshape global expression programs and cell states. Most existing models approximate this complex process with simple embedding operations, such as concatenating or adding perturbation and cell representations, to predict transcriptional responses. This oversimplification neglects the context-dependent interaction between perturbations and cell states and overlooks the hierarchical nature of gene regulation, ultimately leading to inaccurate predictions, weak generalization, and limited interpretability.

These limitations not only hinder accurate modeling of intracellular regulatory processes but also restrict the applicability of current methods when extended to broader perturbation types. Most existing *in silico* perturbation methods are designed for a specific perturbation type, such as either genetic or chemical interventions. Although a few approaches[28, 29] have been explored across multiple perturbation types, they generally require training separate models for each dataset rather than learning within a shared representation space. With the rapid expansion of heterogeneous perturbation datasets, this separate training strategy has become a major bottleneck, as it fails to reflect the biological reality that different classes of perturbations often converge on shared regulatory mechanisms to reshape gene expression and cell states. Learning these shared mechanisms in a unified way is therefore critical for achieving cross-perturbation generalization and deeper mechanistic interpretability. A unified framework that can accommodate diverse perturbation types within a single representation space is thus essential for capturing shared principles of cellular responses and enabling more accurate response prediction.

To address these challenges, we propose X-Pert, a unified framework for modeling cellular responses to diverse perturbation types. X-Pert adopts a transformer backbone with latent query tokens that represent different external perturbations in a shared representation space, enabling joint training across both genetic and chemical perturbation datasets as well as extending naturally to additional perturbation types such as cytokine perturbations[31]. Through a combination of cross-attention and self-attention mechanisms, X-Pert explicitly captures gene–perturbation and gene–gene interactions, thereby modeling the hierarchical regulatory processes that determine how perturbations reshape transcriptional programs. We benchmark X-Pert across both genetic and chemical perturbation tasks, using not only conventional metrics that measure prediction accuracy across the entire expression profile but also new biology-aware metrics that assess the prediction on top differentially expressed genes and pathway-level responses. Across these comprehensive benchmarks, X-Pert consistently outperforms existing methods, achieving higher predictive accuracy and

capturing biologically meaningful variation among perturbations. Moreover, perturbations of varying dosage, combination, and type can be consistently embedded into a unified latent space, supporting downstream applications including perturbation retrieval, drug–gene association discovery, combinatorial perturbation prediction, and dose- or efficacy-aware prediction. Together, these capabilities position X-Pert as a versatile and generalizable framework that bridges genetic mechanisms and pharmacological interventions, offering a systematic foundation for understanding and designing cellular perturbations.

Results

X-Pert: a unified model for *in silico* perturbation

We designed X-Pert to predict the gene expression profile of a perturbed cell from the original cell and the corresponding perturbation information. Unlike previous approaches that treat different perturbation types separately, X-Pert integrates perturbation signals and cellular context within a unified architecture to capture their fine-grained interactions.

The model consists of two key components: the Perturbation Perceiver and the Cell Encoder (Fig. 1, Methods). The Perturbation Perceiver maps perturbations of different types into a shared latent representation space. Each perturbation type is first initialized as a fixed-length embedding by a type-specific encoder (in this work, we used text embeddings from GPT-4o[32] for genetic perturbations, and SMILES-based embeddings computed with RDKit[33] for chemical perturbations). Dose or perturbation efficacy information, when available, is then incorporated by augmenting the corresponding perturbation embedding. Depending on the experimental setting, a single perturbation embedding or a set of embeddings for multiple perturbations is then passed through a Perceiver Resampler, where a fixed set of learnable latent query tokens are employed to transform these embeddings into unified perturbation tokens (Methods). These tokens define a shared representation space that enables unified characterization of perturbations and their combinations, which we term the Perturbverse.

The unified perturbation tokens produced by the Perturbation Perceiver serve as conditional context for the Cell Encoder, a transformer-based module designed to predict perturbed gene expression profiles by modeling both gene–perturbation interactions and gene–gene dependencies (Methods). The Cell Encoder builds upon a large cellular model to encode an original cell as a sequence of gene identity tokens and expression value tokens (Methods). To better incorporate perturbation effects, we introduce an Impact Tokenizer that encodes perturbation-specific influences on each gene. The formulation of impact tokens is designed to flexibly accommodate different perturbation types: for genetic perturbations, they are modeled as one-hot vectors marking perturbed genes; for chemical perturbations, they are derived from protein–molecule affinity scores predicted by a pre-trained drug–target binding affinity prediction model[34] (Methods). Each gene token is then summarized by integrating its gene identity, expression value, and impact components. These summarized

gene tokens interact with perturbation tokens through cross-attention to capture perturbation–gene relationships. Subsequently, self-attention layers are employed to model gene–gene dependencies and preserve the intrinsic regulatory structure of the cell. The Cell Encoder outputs normalized gene expression values for the perturbed cell.

X-Pert is trained end-to-end with a mean squared error (MSE) loss between predicted and observed profiles (Methods). Once trained, it serves not only as a robust predictor of cellular responses but also as a versatile framework for downstream biological discovery. The Perturbverse serves as a unified reference space that enable cross-perturbation analysis, supporting perturbation retrieval and gene–drug association discovery. The Cell Encoder flexibly handles diverse datasets and biological contexts, supporting the prediction of single or multiple perturbations as well as dose- or efficacy-dependent responses. These capabilities establish X-Pert as both a powerful *in silico* perturbation predictor and a general foundation for applications in biology and medicine.

Fig. 1. Overview of the model architecture of X-Pert. X-Pert consists of two core components: the Perturbation Perceiver and the Cell Encoder. The Perturbation Perceiver is designed to accommodate different types of external perturbations, including multiple simultaneous perturbations and dose- or efficacy-dependent effects. Through a set of learnable latent queries, it projects perturbations of any type and number into a fixed-length latent representation, termed the Perturbverse. The Cell Encoder models the original cell using three types of tokens, namely identity tokens, value tokens, and impact tokens, which respectively encode gene identity, expression magnitude, and perturbation influence. By combining a gated cross-attention module with transformer encoder layers, the Cell Encoder integrates perturbation information with original gene expression to predict the transcriptional profile of perturbed cells. X-Pert can be applied to

116   a wide range of perturbation-based tasks. Leveraging the Perturbverse, X-Pert enables perturbation retrieval and gene–

117   drug association discovery through similarity analysis. Using the Cell Encoder's predictive capability, X-Pert supports

118   single/combinatorial perturbation prediction as well as modeling of perturbation efficacy and dose–response

119   relationships.

Predicting responses of unseen genetic perturbations

We first evaluated X-Pert's performance on unseen genetic perturbation prediction, representing a real-world

use case in which the model infers cellular responses to perturbations not yet experimentally profiled within

the same cell type (Fig. 2A). To evaluate this capability, we benchmarked X-Pert on three single-cell

perturbation datasets spanning CRISPR interference[4] (CRISPRi, Replogle2022_K562 and

Replogle2022_RPE1) and CRISPR activation[35] (CRISPRa, Norman2019), comprising between 224 and 1,299

unique perturbations (Methods). For each dataset, perturbations were split 8:2 into training and testing

(Methods). We compared X-Pert with six representative methods, including linear models[36] (Linear, Linear-

scGPT), deep learning methods (CellOracle[37], GEARS[17]), and large cellular foundation models (scGPT[20],

scFoundation[19]), and the mean predictor (AveKnown) that have been used as a naive baseline in some

studies[28, 38].

We first evaluated model performance using a commonly used metric, the Pearson correlation coefficient

(PCC) across all genes, referred to as PCC_All, which quantifies the correlation between predicted and

observed gene-expression changes for each perturbation, calculated over all genes as paired values of

predicted and measured expression differences relative to the original cells. As shown in Fig. 2B, X-Pert

achieved the best or highly competitive performance on PCC_All across all three datasets. However, we found

that PCC_All provides little meaningful differentiation among models, as even naïve predictors such as

AveKnown perform similarly. A recent study has revealed a pervasive systematic bias in genetic-perturbation

datasets, where shared global response components dominate gene-expression variance, overshadowing the

true perturbation-specific regulatory signals[39]. Consequently, even simple baselines such as AveKnown can

achieve deceptively high correlation scores by reproducing these shared patterns, a phenomenon consistently

reported across benchmarks[36, 38]. Therefore, while PCC_All remains a convenient and widely used summary

statistic, it primarily reflects the recovery of background structure rather than the accurate modeling of

biological perturbation effects.

To better assess biological relevance responses, we next examined PCC_DEG, which quantifies the

correlation between predicted and observed expression changes only among DEGs (Methods). This metric

filters out nonspecific or noisy variation and provides a clearer view of biologically meaningful transcriptional

responses. For instance, it reached 0.720 on Replogle2022_K562, a 13.2% improvement over the second-best

method, with similar gains observed on Replogle2022_RPE1 (+3.8%). Importantly, most methods showed a

larger increase in performance when evaluated by PCC_DEG compared with PCC_All across datasets. This

consistent widening of the performance gap suggests that PCC_DEG captures perturbation effects more faithfully by highlighting genuine expression changes associated with each perturbation, which are largely averaged out in PCC_All by shared global responses. However, despite this improvement, the PCC_DEG of AveKnown remains high, indicating that this metric still carries dataset-level correlations and is limited in distinguishing biologically meaningful variation from background effects. These limitations underscore the need for evaluation strategies that move beyond correlation-based similarity and instead assess a model's ability to reconstruct both the magnitude and the functional organization of true perturbation effects.

To move beyond correlation-based metrics and directly assess how accurately each model captures real perturbation effects, we focused on perturbations with at least 20 significant DEGs. This subset reflects conditions where transcriptional regulation is strongly altered and thus provides a clearer test of biological fidelity. For each perturbation, we compared the predicted and observed expression changes of the top 20 DEGs, which represent the most strongly responsive genes and are typically related to the downstream functional consequences of the perturbation. This evaluation highlights not only whether a model predicts the correct direction of change but also whether it reproduces the actual magnitude of transcriptional shifts. For example, in the POLA1 perturbation of Replogle2022_K562, X-Pert closely recapitulated both the direction of magnitude of top 20 DEGs' expression changes, whereas other methods tended to compress predictions toward low-magnitude shifts (Fig. 2C). Similar patterns were observed for additional perturbations across datasets (Fig. S1A). We further quantified this agreement using the coefficient of determination (R^2) between the predicted and observed expression changes of the top DEGs (Methods). Across all three datasets, X-Pert consistently achieved the highest R^2 scores (Fig. 2D and Fig. S1B), demonstrating its ability to accurately reproduce the real magnitudes of perturbation-induced transcriptional responses rather than merely capturing overall expression trends.

While single-gene evaluations help assess whether a model can reproduce individual expression changes, cellular responses are in fact mediated by coordinated regulation of gene programs. Therefore, evaluating predictions at the pathway level provides a more biologically meaningful and noise-robust assessment of model performance. To assess pathway-level responses, we computed Hallmark gene set scores[40] for each perturbation by averaging the corresponding signature scores across all cells under that condition (Methods), reflecting the activation state of specific pathways after perturbation. Such evaluation moves beyond conventional correlation-based metrics, as it directly tests whether a model can recover the coordinated activation or repression of biological programs that define a cell's functional state. We quantified the agreement between predicted and observed gene set scores across different perturbations using PCC (PCC_GS). Unlike gene-level metrics that assess agreement within each perturbation, PCC_GS compares model performance across perturbations, evaluating whether the model can correctly rank perturbations by the magnitude and direction of their pathway-level effects. This distinction is important because the ability to prioritize perturbations by their functional impact forms the basis for many biological applications such as perturbation retrieval, target identification, and mechanism inference.

On 260 test perturbations from the Replogle2022_K562 dataset, X-Pert captured this diversity with high fidelity. For example, in the DNA repair gene set, it achieved the highest PCC_GS (0.40), representing a 53.8% improvement over the second-best method, GEARS (PCC_GS = 0.26, Fig. 2E). In contrast, AveKnown completely failed under this setting, producing identical predictions for all perturbations and thus showing no ability to distinguish functional differences between them. Furthermore, X-Pert successfully predicted a cluster of perturbations exhibiting coordinated decreases in DNA repair gene set scores within the UMAP space, a structure that none of the other methods were able to capture (Fig. 2E). Similar advantages were observed for additional gene sets such as adipogenesis, apical junction, and protein secretion (Fig. S1C). Across all 50 Hallmark gene sets, X-Pert consistently outperformed all other methods (Fig. 2F), demonstrating superior ability to model perturbation-specific gene program changes.

These results demonstrate that X-Pert not only achieves accurate performance across multiple evaluation metrics but also provides biologically coherent predictions at both gene and pathway levels. These findings also highlight that pathway-level benchmarks, by comparing relative perturbation effects across perturbations rather than isolated gene-level predictions within a single perturbation, provide a more faithful measure of biological realism. Such benchmarks align with findings from a recent perturbation benchmark[39] that emphasized evaluating performance across the full spectrum of perturbations rather than isolated cases.

Building on X-Pert's strong predictive performance, we next examined what the model has learned about the underlying structure of perturbations. We explored the interpretability of X-Pert's learned latent perturbation space, or Perturbverse, generated by the Perturbation Perceiver (Methods). Because perturbations are represented using gene-related text embeddings, even perturbations absent from training can be directly embedded into the space (Fig. 2G). This provides a structured representation of perturbation effects for revealing functional relationships among genetic perturbations. For instance, 504 genetic perturbations from Replogle2022_K562 revealed clusters corresponding to biological processes such as proteasome, mediator complex, and DNA replication, despite only 209 perturbations being included in training (Fig. 2H and S2A).

The Perturbverse also provides a convenient framework to examine how clusters of perturbation genes relate to downstream expression changes. By mapping all input perturbations into the Perturbverse (Methods), we found that proteasome-related perturbations formed a coherent cluster, and their predicted profiles showed coordinated upregulation of multiple subunits (PSMB5, PSMD1, PSMD13), consistent with the known "bounce-back" transcriptional response[41, 42] (Fig. 2H–I). A similar Perturbverse constructed for Replogle2022_RPE1, despite being trained on only 73 perturbations, also recovered clear and biologically coherent structures (Fig. S2A–B). These results highlight that X-Pert not only achieves accurate prediction of unseen genetic perturbations but also organizes them into an interpretable latent space that reveals functional relationships between perturbation clusters and transcriptional programs.

Fig. 2. X-Pert's predictions of genetic perturbations. (A) Schematic illustration of predictions on unseen genetic

perturbations. **(B)** Comparison of X-Pert with other methods. **(C)** Comparison between predicted and ground-truth

expression changes of the top 20 DEGs following POLA1 perturbation in Replogle2022_K562. Dots represent the mean

predicted expression changes from each method, and a linear regression fit was added to visualize the correlation

between the predicted and true mean values. CellOracle is not included because it only performs prediction on

transcription factors. **(D)** Box plots comparing the R^2 score between predicted and observed expression changes of top 20

DEGs in Replogle2022_K562. **(E)** UMAP visualization of Hallmark DNA repair gene set scores. The leftmost panel

shows ground-truth scores, and each point represents a genetic perturbation. The PCC_GS between predicted and

observed gene set scores is shown in each panel. The dashed circle highlights the ground-truth clustering of perturbations

with reduced gene set scores. **(F)** Box plots comparing the PCC_GS of all 50 Hallmark gene sets across 260 test

perturbations in Replogle2022_K562. For bar plots, bar: mean performance value; error bar: standard deviation across

replicates. For box plots, center line, median; box limits, upper and lower quartiles; whiskers, 1.5× interquartile range.

(G) Schematic illustration of using latent tokens from the Perturbverse to explore the functional landscape of genetic

perturbations. **(H)** UMAP visualization of 504 genetic perturbations from Replogle2022_K562, with pathway

annotations manually curated from the original dataset. **(I)** UMAP visualization of mean gene expressions across

237 504 annotated genetic perturbations from Replogle2022_K562. The ten displayed genes were selected from

238 significant DEGs in proteasome-related perturbations. Dashed circles represent cells perturbed by proteasome-

239 related genes.

Predicting responses under perturbation combinations and considering efficacy

The flexible design of the Perturbation Perceiver enables X-Pert to naturally handle both combinatorial

perturbations and perturbation efficacy. To assess its ability to generalize to unseen combinations (Fig. 3A),

we used the Norman2019 dataset, which includes 148 single-gene and 76 two-gene CRISPRa perturbations.

To simulate the unseen-combination setting, we randomly excluded 30% of the single-gene perturbations

during training and evaluated two types of double perturbations: one-unseen combinations ($n = 61$), where one

perturbed gene was absent from training, and two-unseen combinations ($n = 15$), where both were unseen.

Notably, the evaluation did not include "zero-unseen" combinations, as such cases do not naturally arise from

this data-splitting strategy.

As shown in Fig. 3B, X-Pert substantially outperformed all other methods in the one-unseen setting across all

metrics. Even in the more challenging two-unseen setting, X-Pert maintained the best performance in in R^2

score and PCC_GS, and remained highly competitive in PCC_All and PCC_DEG. Notably, X-Pert was the

only model that achieved a clearly positive median PCC_GS in the two-unseen setting, underscoring its

capacity to generalize to novel perturbations by leveraging gene-related text embeddings for perturbation

representation. Together, these results demonstrate that X-Pert effectively integrates multiple perturbations

and extends robustly to unseen combinatorial conditions.

Modeling perturbation efficacy represents another important challenge (Fig. 3C). For example, In CRISPR-

based experiments, perturbation efficacy depends heavily on sgRNA design but is not directly measurable,

unlike chemical perturbations where dose provides a natural quantitative proxy. To address this, we quantified

genetic perturbation efficacy by measuring target gene downregulation relative to original cells (Methods),

and examined its relationship with the overall transcriptional impact, quantified by the energy distance (E-

distance) between perturbed and original profiles. For the full Replogle2022_K562 dataset, we didn't find

global correlation between E-distance and perturbation efficacy (Fig. 3D). This is likely because many

perturbations induce weak or noisy transcriptional shifts, which obscure the link between efficacy and

downstream effects. However, focusing on a more homogeneous subset of 37 ribosomal protein large subunit

(RPL) perturbations, we observed a significant negative correlation (PCC = -0.43, $p = 0.0085$; Fig. 3D),

indicating that higher efficacy generally drives stronger transcriptional responses. These findings suggest that

efficacy information should be explicitly incorporated into predictive models.

Building on this insight, we extended X-Pert to incorporate efficacy information, yielding X-Pert_efficacy

(Methods). Across all perturbations in Replogle2022_K562, X-Pert_efficacy performed comparably to or

slightly better than the original model (Fig. 3E). When investigating the RPL subset, it showed a clear

advantage: X-Pert_efficacy achieved significantly higher R^2 on eight held-out perturbations (Fig. 3E) and

more faithfully recovered expression changes of top DE genes (Fig. 3F and Fig. S3). For example, in the case

of RPL14 perturbation, it can more accurately predicted expression changes in top DEGs, which were

underestimated by the original model (Fig. 3F). Similar improvements were consistently observed across other

test RPL perturbations (Fig. S3), highlighting the value of incorporating efficacy information into the

predictive framework.

Fig. 3. X-Pert's predictions of perturbation combinations and with efficacy modeling. (A) Schematic illustration of

multi-perturbation prediction. **(B)** Comparison of X-Pert with other methods in the combinatorial perturbation prediction

setting. For bar plots, bar: mean performance value; error bar: standard deviation across replicates. For box plots, center

line, median; box limits, upper and lower quartiles; whiskers, 1.5× interquartile range. **(C)** Schematic illustration of

perturbation prediction incorporating efficacy information. **(D)** Scatter plots with fitted regression lines showing the

relationship between E-distance and perturbation efficacy across all perturbations (left) and RPL-related perturbations

(right) in Replogle2022_K562. **(E)** Bar plots comparing X-Pert and X-Pert_efficacy across all test perturbations (left)

and RPL-related perturbations (right). **(F)** Comparison between predicted and ground-truth expression changes of the top

20 DEGs following RPL14 perturbation in Replogle2022_K562. Dots represent the mean predicted expression changes

from each method, and a linear regression fit was added to visualize the correlation between the predicted and true mean

values.

Predicting responses of chemical perturbations

We next evaluated X-Pert on chemical perturbations (Fig. 4A), We first benchmarked it on the Sciplex-3

dataset[11], which profiles three cell lines (A549, K562, MCF7) treated with 188 drugs across four

concentrations (10 nM–10,000 nM, 758,368 cells after preprocessing). Perturbations were split 8:2 into

training and testing. We selected representative chemical perturbation prediction methods for comparison,

including ChemCPA[24], BioLord[25], and PRnet[26]. As shown in Fig. 4B, X-Pert achieved the best or highly

competitive performance across PCC_All, PCC_DEG, and R^2 score, and reached the highest PCC_GS in

K562 and MCF7. However, the overall predictive performance across all models was modest, with median R^2

scores close to zero. This limitation appears to arise from the intrinsic characteristics of the Sciplex-3 dataset:

most perturbations induced only a small number of significant DEGs, even with higher compound

concentrations (Fig. S4A). Such weak transcriptional responses result in a low signal-to-noise ratio, making it

difficult for models to learn robust perturbation-response relationships.

To test whether X-Pert captures dose-dependent effects, we examined the M344 perturbation in MCF7 cells

as a case study. M344 is a histone deacetylase (HDAC) inhibitor with marked dose-dependent heterogeneity,

currently under investigation for its therapeutic potential in cancer and Alzheimer's disease[43]. Across the top

10 DEGs, X-Pert faithfully reproduced dose–response patterns, capturing both trends and magnitudes of

change (Fig. 4C). Similar results were observed across other compounds (Fig. S4B), confirming the

effectiveness of explicit dose modeling.

We further extended the evaluation to the CMAP (Connectivity Map) dataset[44], a large bulk-level

perturbation resource profiling 17,775 small molecules across 70 cell lines using the L1000 technology

(678,401 profiles after preprocessing). The diversity of CMAP allows testing on unseen molecules and unseen

cell types, posing greater challenges than single-cell datasets. To account for bulk-level measurements, we

evaluated both PCC_All and R^2 (PCC_DEG and PCC_GS were excluded from the evaluation because the

L1000 technology already selected 978 genes for profiling; R^2 was calculated based on all profiled genes). As

shown in Fig. 4D, X-Pert consistently outperformed all other methods under both settings. On unseen

molecules, it exceeded BioLord by 3.1% in PCC_All and 12.0% in R^2. On unseen cell types, the gains were

even larger (PCC_All +11.4%, R^2 +74.3%), underscoring X-Pert's ability to capture drug responses in diverse

cellular contexts.

We then explored whether chemical perturbations form meaningful representations in the Perturbverse. For

851 CMAP molecules with DrugBank ATC level 1 annotations[45], original embeddings lacked discernible

organization, whereas X-Pert grouped molecules into functionally coherent clusters, such as "antineoplastic

and immunomodulating agents" (Fig. 4E and Fig. S4C). This enrichment is biologically reasonable, as nearly

all CMAP profiles were generated from cancer-derived cell lines, likely leading X-Pert to amplify signals

from antineoplastic drugs.

Finally, we tested the utility of X-Pert for drug retrieval (Fig. 4F) by projecting ~30,000 natural compounds

from the natural compound database HERB 2.0[46] into the Perturbverse and ranking them by cosine similarity

to reference antineoplastic drugs (Fig. 4G, Methods). Among the top 20 retrieved candidates, nearly half have

documented anticancer activity (Fig. 4H), including diterpenoid quinones[47] (e.g., Agastaquinone), taxane-type

diterpenoids[48, 49] (Taxezopidines K and G), and caffeoylquinic acid derivatives[50, 51] (3-O-caffeoyl-4-O-

sinapoylquinic acid), all consistent with known pharmacological properties. In addition, structurally related

analogues of paclitaxel (e.g., N-debenzoyl-N-cinnamoyltaxol) emerged, suggesting potential therapeutic

relevance.

Together, these results demonstrate that X-Pert not only predicts chemical perturbation responses with high

accuracy, including dose-dependent effects, but also organizes perturbations into biologically meaningful

embeddings that enable *in silico* drug retrieval and discovery.

Fig. 4. X-Pert predictions on chemical perturbations. (A) Schematic illustration of chemical perturbation prediction.

(B) Model comparison on the Sciplex-3 dataset under the unseen molecule prediction scenario. **(C)** Line plots showing

X-Pert–predicted gene expression levels versus ground-truth measurements across four chemical doses. **(D)** Model

comparison on the CMAP dataset under both unseen molecule and unseen cell type settings. **(E)** UMAP visualization of

chemical perturbations in the original molecular embedding space and the Perturbverse, colored by ATC Level 1

annotations. **(F)** Schematic illustration of the drug retrieval process using latent chemical embeddings derived from the

Perturbverse. **(G)** UMAP visualization showing the cosine similarity distribution of candidate molecules from the HERB

2.0 database. **(H)** Top-ranked molecules retrieved by X-Pert from the HERB 2.0 database, shown with their rankings and

compound names. For bar plots, bar: mean performance value; error bar: standard deviation across replicates. For box

plots, center line, median; box limits, upper and lower quartiles; whiskers, 1.5× interquartile range.

In silico **drug screening using large-scale chemical perturbation data**

To extend X-Pert to *in silico* drug screening, we trained it on Tahoe-100M[12], the largest publicly available

single-cell chemical perturbation dataset to date. Generated with the Mosaic platform, Tahoe-100M comprises

~100 million cells across 379 compounds, each profiled at three doses in 50 cell lines. This breadth of

compounds and contexts makes it an ideal resource for large-scale perturbation modeling. To improve

computational efficiency, we aggregated single-cell profiles into pseudo-bulk measurements (Methods), which

preserved perturbation signals while enabling tractable training. Perturbations were split 8:2 into training and

testing.

X-Pert accurately reconstructed the transcriptomic distributions of different cell lines in the UMAP space,

clearly distinguishing them from one another (Fig. 5A), indicating that X-Pert not only predicts gene

expression but also captures cell line-specific contexts. However, closer inspection of A549 cells revealed that

plate labels introduced a strong batch effect, which masked drug-specific signals (Fig. 5B). To mitigate this,

we incorporated a linear regression-based adjustment to remove batch-associated variation before drug

screening (Methods). After correction, we implemented an *in silico* drug screening pipeline (Fig. 5C,

Methods): for each disease, we curated up- and down-regulated gene sets as disease signatures, computed

gene set scores under candidate perturbations (denoted as up-scores and down-scores), and identified drugs

with lower up-scores and higher down-scores with more potential that could reverse disease signatures. This

strategy allowed drug ranking to reflect genuine therapeutic potential rather than technical noise.

We applied this approach to prioritize drugs for three cancer types: pancreatic ductal adenocarcinoma

(PDAC; Panc 03.27), hepatocellular carcinoma (HCC; HepG2), and lung adenocarcinoma (LUAD; NCI-

H1792). From an FDA-approved drug list, we excluded compounds used for training, yielding 831 candidates.

X-Pert ranked the top ten drugs for each cancer.

As shown in Fig. 5D, for PDAC, top candidates included glucocorticoids (desonide, meprednisone, and

dexamethasone acetate), which have shown unexpected anti-tumor activity in PDAC models[52]. X-Pert also

highlighted Pracinostat, a pan-HDAC inhibitor under clinical investigation[53]. Another strong candidate was

Entospletinib, a selective SYK inhibitor originally developed for B-cell cancers[54], with evidence of reshaping

tumor immunity[55] and overcoming PDAC chemoresistance[56], as supported by related compounds in

preclinical models. For HCC, X-Pert identified nitroxoline, a cathepsin B inhibitor with well-documented

anti-liver cancer effects[57]. Another candidate, clioquinol, an 8-hydroxyquinoline derivative, exerts anticancer

activity through metal chelation, proteasome inhibition, and apoptosis induction[58], and preclinical studies

reported its antiproliferative effects in liver cancer cells[59]. Empagliflozin, an SGLT2 inhibitor approved for

diabetes[60] with emerging evidence of benefit in preclinical HCC models[61], was also prioritized. For LUAD, X-

Pert prioritized sotagliflozin, consistent with the reliance of LUAD on SGLT2-mediated glucose uptake[62] and

clinical signals of benefit from SGLT2 inhibitors in NSCLC patients[63]. It also highlighted ciclopirox, which

suppresses proliferation and induces apoptosis in LUAD models[64], and veliparib, a PARP inhibitor with

reported benefit in NSCLC subgroups[65].

Together, these results show that X-Pert can integrate large perturbation datasets with disease-specific

transcriptional signatures to prioritize candidate drugs across different cancer types, demonstrating its

potential as a practical tool for accelerating drug discovery and repurposing.

Fig. 5. *In silico* **drug screening using the Tahoe-100M dataset. (A)** UMAP visualization of test perturbations predicted by X-Pert, where each point represents a pseudo-bulk cell derived from a specific chemical perturbation in a cell line. Points are colored by cell line. **(B)** UMAP visualization of all A549 samples, colored by sequencing plate and count of all genes, respectively, highlighting potential batch effects. **(C)** Schematic illustration of the *in silico* drug screening pipeline. **(D)** Scatter plots showing the top-ranked candidate drugs and fitted regression lines for three diseases: PDAC, HCC, and LUAD. Lower up-scores and higher down-scores represent drugs with higher potential that could reverse disease signatures. The top ten prioritized drugs are annotated in red.

Aligning genetic and chemical perturbations in Perturbverse

The unified design of the Perturbation Perceiver allows X-Pert to jointly model multiple perturbation types within a unified framework. To evaluate this capability, we used the CMAP dataset, which contains both genetic and chemical perturbations profiled on same cell lines. We focused on three cell lines with the largest number of dual–perturbation-type profiles: A549 (21,685 chemical and 3,620 genetic perturbations), VCAP (34,753 chemical, 3,641 genetic), and HT29 (14,513 chemical, 3,302 genetic), and trained X-Pert independently on each cell line with an additional Maximum Mean Discrepancy (MMD) loss to encourage cross-type alignment (Methods).

We first quantitatively evaluated X-Pert's ability to align genetic and chemical perturbations. Visualization of the Perturbverse showed that genetic and chemical perturbations were well integrated (Fig. 6A). We selected a set of representative genes, including AKT1, AKT2, CTSK, SMAD3, TP53, MTOR, BRAF, and PIK3CA, as they represent diverse, therapeutically relevant signaling pathways and are associated with known or putative small-molecule modulators. For each gene, we queried GDIdb[66] to identify drugs interacting with

it, yielding 7–23 compounds per gene. We then calculated the cosine similarity between the latent embeddings

of these drugs and their corresponding genes learned by X-Pert. As shown in Fig. 6B, ground-truth gene–drug

pairs consistently showed higher cosine similarity than random drug–gene pairs for most genes across all three

cell lines. These results confirm that X-Pert effectively captures functional correspondence between genetic

and chemical perturbations.

To assess its generalization, we focused on the HT29 cell line and examined the test perturbations that were

excluded from model training (Fig. 6C). The low-dimensional visualization revealed that genetic and chemical

perturbations were also well fused. Subgroups in this space revealed biologically plausible relationships. For

example, in subgroup 1, deltaline aligned with NAIP deletion, both reducing muscle contraction despite

distinct mechanisms[67, 68]; tetramethylpyrazine hydrochloride (TMPH) aligned with MAP2K4 knockout, both

suppressing JNK signaling[69, 70]. Other examples included vicriviroc aligning with VIM knockdown, both of

which impair tumor cell invasion[71, 72], and farnesylthioacetic acid aligning with RASGRP4 knockdown, both

suppressing Ras-dependent signal transduction and thus reducing proliferation and promoting apoptosis[73, 74].

Subgroup 2 showed parallels such as aliskiren with MRGPRD loss, both protecting cardiovascular cell

function by attenuating angiotensin II (Ang II) signaling[75, 76]. Likewise, MRGPRD knockout showed

functional similarity to anti-inflammatory drugs such as clocortolone pivalate and limonin[77-79]. Besides, both

PTPRG loss and alpha-linolenic acid supplementation improved insulin sensitivity and reduced

inflammation[80, 81]. Together, these results highlight X-Pert's capacity to align genetic and chemical

perturbations in a unified latent space and to uncover biologically meaningful relationships between

perturbations.

We further evaluated X-Pert's ability to perform cross-type perturbation retrieval within the Perturbverse of

HT29 cells. HT29, a human colon cancer cell line harboring the BRAF V600E mutation, is frequently used to

study the effectiveness of targeted therapies that inhibit the BRAF–MEK–ERK signaling pathway[82]. We

queried the BRAF perturbation embedding of Perturbverse against 1,919 natural compounds from the

Selleckchem database[83], and we found a distinct subset of compounds with exceptionally high similarity

scores to the BRAF perturbation embedding (Fig. 6D). We then identified top 10 candidates with high

similarity (Fig. 6E). The top-one candidate is Astragaloside A, which, though not a direct enzymatic inhibitor

of BRAF, have been shown to modulate the MAPK/ERK pathway[84], thereby producing functional outcomes

comparable to inhibition of RAF/MEK signaling. Another prominent group, including glycyrrhizic acid and

its derivatives (including ammonium glycyrrhizinate and diammonium glycyrrhizinate), has been shown to act

as RAS allosteric inhibitors[85]. We also identified Fargesin which has been reported to inhibit the PKA/CREB

and p38 MAPK pathways, thereby affecting cellular processes regulated by the MAPK family[86]. While its

mechanism does not directly target BRAF, its modulation of MAPK downstream kinases, such as ERK and

p38[87], suggests potential functional convergence with BRAF inhibition.

Taken together, these results demonstrate that X-Pert's latent space can not only align genetic and chemical

perturbations but also retrieve compounds that mimic the effects of gene depletion. This highlights the
model's potential as a framework for flexible drug discovery and repurposing.

Fig. 6. Cross-type alignment with genetic and chemical perturbations. (A) UMAP visualization of Perturbverse
embeddings for all perturbations across A549, VCAP, and HT29 cell lines. **(B)** Radar plots comparing the cosine
similarity between X-Pert–prioritized drugs and randomly selected drugs. Cosine similarity was calculated between the
prioritized compounds and ground-truth drugs known to interact with the corresponding genes, as obtained from the
DGIdb database. Lines denote the mean values, and shaded areas denote standard deviations across samples. **(C)** UMAP
visualization of held-out test perturbations in the HT29 cell line, with the two highlighted subgroups shown with their
corresponding perturbation names. Black lines denote biologically plausible relationships. **(D)** Histogram showing the
distribution of cosine similarities between 1,919 natural compounds from the SelleckChem database and the BRAF
perturbation embedding in the Perturbverse space. (E) Top ten drugs with the highest similarities with BRAF
perturbation, shown with their rankings and compound names.

Discussion

In this study, we introduced X-Pert, a unified framework that not only achieves accurate prediction across
genetic and chemical perturbations, but also constructs a shared perturbation embedding space, the
Perturbverse, that supports systematic interpretation and exploration of perturbation effects. The strong
performance and wide-ranging capability arise from its architectural design. By combining a Perturbation
Perceiver with a transformer-based Cell Encoder, X-Pert explicitly models fine-grained gene–perturbation
interactions while maintaining the intrinsic structure of cell states. This design results in strong performance
across tasks ranging from unseen perturbation prediction to combinatorial and dose/efficacy-aware modeling.
Beyond predictive accuracy, the Perturbverse provides an interpretable latent space in which different types of
perturbations can be compared, aligned, and retrieved, opening new possibilities for drug discovery and gene–

drug association investigation.

Although our work primarily focused on genetic and chemical perturbations, the design of X-Pert renders it

inherently extensible to additional perturbation types. Furthermore, the transformer-based architecture and the

design of Perturbation Perceiver make X-Pert naturally suited for pre-training on large-scale perturbation

corpora. As single-cell perturbation technologies continue to mature, the volume of such datasets will create

opportunities to establish X-Pert as a foundation model for perturbation biology, capable of integrating

heterogeneous data across platforms, cell types, and perturbation types.

Most existing computational approaches to perturbation modeling have focused almost exclusively on RNA

transcriptomic profiles[17-22], reflecting the widespread adoption of single-cell RNA sequencing (scRNA-seq) as

the most mature and accessible technology for large-scale single-cell profiling. However, advances in

sequencing and profiling technologies are rapidly expanding the omics modalities available for perturbation

studies, including epigenomic readouts[88] and protein expression measurements[89] at single-cell resolution.

These developments call for methods that can move beyond RNA-centric prediction to jointly capture multi-

modal perturbation effects, encompassing RNA, DNA regulatory features, and protein-level responses. The

growing availability of such datasets also opens avenues for the development of foundation models, capable

of learning universal representations that generalize across biological contexts[90, 91]. In this work, we adopted a

scRNA-seq foundation model[20] as the backbone of X-Pert, and X-Pert's modular design makes it readily

extensible to additional omics layers, positioning it as a candidate perturbation foundation model that

integrates transcriptomic, epigenomic, and proteomic data to capture the full spectrum of cellular responses to

external perturbations.

A central challenge in advancing perturbation modeling lies in defining what it truly means for a model to

be "accurate". Conventional evaluation metrics, such as correlation across genes, often overestimate

performance by rewarding the recovery of shared expression patterns rather than the reconstruction of causal,

perturbation-specific effects. This issue extends beyond benchmarking, because it fundamentally shapes how

models are trained and optimized. Our work emphasizes that evaluating predictive accuracy from the

perspective of biological function is crucial for capturing genuine regulatory responses, as it reflects how well

a model understands cellular mechanisms rather than statistical associations. Nevertheless, the metrics

proposed in this study represent only an initial step toward evaluating *in silico* perturbation models from a

biological perspective. Future progress will depend on developing a broader suite of complementary metrics

that can more effectively disentangle true causal perturbation responses from confounding factors and

technical artifacts.

Another critical direction concerns data: generating large-scale perturbation corpora while addressing

intrinsic challenges such as batch effects, cellular heterogeneity, and stochasticity. Although our study paired

original and perturbed cells, we recognize that cells are inherently dynamic and noisy, and it remains difficult

to disentangle true signals from background variation. By building on the foundation provided by X-Pert's

framework, future efforts can expand toward a true perturbation foundation model and ultimately a virtual
cell, offering unprecedented power for understanding and engineering cellular responses.

Methods

Perturbation Perceiver

Perturbation Resampler

The Perturbation Perceiver, inspired by the perceiver architecture used in image–text multi-modal models[92], is
designed to integrate heterogeneous perturbations into a unified latent representation. Formally, let $X_p \in$
$\mathbb{R}^{n_p \times d}$ denote the perturbation embeddings, where n_p is the number of perturbation tokens and d is the
latent dimensionality. We additionally introduce a set of learnable latent queries $X_l \in \mathbb{R}^{n_l \times d}$, where n_l is
fixed and independent of the type or number of perturbations. In the attention layer, the latent queries are used
to extract information from both the perturbation inputs and the latent space itself. Specifically, the queries are
defined as $Q = X_l$, while the keys and values are constructed by concatenating the perturbation embeddings
with the latent queries: $K = V = [X_p, X_l]$. The attention operation is then given by

517
$$\text{Attn}(Q, K, V) = \text{softmax}\left(\frac{QK^{\text{T}}}{\sqrt{d}}\right)V.$$

518 The output of the attention block is subsequently passed through a feed-forward network (FFN) with residual
519 connections to enhance representational capacity and stabilize training:

520
$$h = \text{FFN}(\text{Attn}(Q, K, V) + X_l) + X_l.$$

We set the number of latent queries $n_l = 12$ and embedding dimension $d = 512$. The Perturbation
Perceiver consists of 2 attention layers, each employing a multi-head attention mechanism with 8 heads. The
final output of Perturbation Perceiver is denoted as $X \in \mathbb{R}^{n_l \times d}$, which serves as the keys and values for the
downstream Cell Encoder. This design enables Perturbation Perceiver to project perturbations of diverse types
into a shared latent space, providing a unified and flexible representation that can be seamlessly integrated by
the Cell Encoder for subsequent modeling of perturbation–gene interactions.

Perturbation-type–specific encoders

For different perturbation types, we employ type-specific encoders to transform the raw perturbation
embeddings into the input space of the Perturbation Resampler. Specifically, for genetic perturbations we
denote the original input as $X_p^g = [x_1^g, ..., x_{n_p}^g]$, and for chemical perturbations we denote the input as $X_p^m =$
$[x_1^m, ..., x_{n_p}^m]$, where x_i^g and x_i^m represent individual genetic and chemical perturbation embeddings,

respectively. To ensure that perturbations of different types are mapped into a comparable latent space, we use
separate three-layer multilayer perceptrons (MLPs) to project the original embeddings:

$$\boldsymbol{X}_p = \widehat{\boldsymbol{X}}_p^g = f_g(\boldsymbol{X}_p^g), \qquad \boldsymbol{X}_p = \widehat{\boldsymbol{X}}_p^m = f_m(\boldsymbol{X}_p^m),$$

where $f_g(\cdot)$ and $f_m(\cdot)$ denote the genetic and chemical encoders, respectively. These MLP-based encoders
enable flexible handling of heterogeneous perturbation types while maintaining a unified representation that
can be processed by the Perturbation Resampler.

For genetic perturbations, we leverage large language models (LLMs) to generate raw perturbation
embeddings, motivated by the fact that LLMs capture broad biological knowledge across genes through large-
scale pretraining. For each target gene, we constructed a natural language prompt of the form: "Impact of
knocking out the gene {gene_name} on cell phenotype?" and submitted it to the GPT-4o model. The
generated textual response, which encodes functional information about the gene perturbation, was then
transformed into a numerical embedding using the GPT text-embedding-ada-002 model, resulting in a 1536-
dimensional vector that serves as the raw genetic perturbation embedding.

For chemical perturbations, we adopted a cheminformatics-based approach. Each molecule was represented
by its SMILES string, and the RDKit library[33] was used to compute chemical embeddings via Morgan
fingerprints. This procedure generated 1024-dimensional binary vectors, where each dimension (0/1) indicates
the presence or absence of a specific chemical substructure feature.

Finally, to map perturbations from both types into a shared latent space, the perturbation-type–specific
projection networks $f_g(\cdot)$ and $f_m(\cdot)$, were applied. These networks reduce the dimensionality of the raw
embeddings from 1536 (genes) and 1024 (molecules) down to a unified 512-dimensional representation,
which serves as the input for the Perturbation Resampler.

Perturbation strength incorporation

The perturbation strength (dose or efficacy) can also be incorporated into the Perturbation Perceiver
framework, encompassing both the dose information for chemical perturbations and the efficacy information
for genetic perturbations. Formally, for input perturbation embeddings $\boldsymbol{X}_p = [\boldsymbol{x}_1, \dots, \boldsymbol{x}_{n_p}]$, we denote the
strength for each perturbation $\boldsymbol{x}_i$ as $s_i \in \mathbb{R}$. To integrate this factor, we introduce a learnable vector $\boldsymbol{w} \in \mathbb{R}^d$
that modulates the embedding dimensions according to the perturbation strength. When strength information
is included, the original perturbation embedding is modified as:

$$\widetilde{\boldsymbol{x}}_i = \boldsymbol{x}_i \odot (s_i \cdot \boldsymbol{w}),$$

where $\odot$ denotes element-wise multiplication. This formulation allows the perturbation embedding to be
adaptively scaled according to its strength.

For genetic perturbations, the perturbation strength is equivalent to perturbation efficacy, which we estimate

by comparing gene expression levels between original and perturbed cells. Let μ_c denote the average
expression level of the target gene in original cells, and μ_p denote its expression level in perturbed cells. The
perturbation efficacy is then defined as:

$$s_i = \frac{\mu_p - \mu_c}{\mu_c}.$$

An efficacy value of -1 corresponds to complete knockout, meaning the target gene is reduced to zero
expression relative to original cells.

For chemical perturbations, strength information is typically encoded as dose. For instance, in the Sciplex-3
dataset, each molecule is profiled under four concentrations: 10 nM, 100 nM, 1,000 nM, and 10,000 nM. We
apply a $\log_{10}$ transformation to normalize the dose levels, yielding dose-based strength values that are more
consistent for model integration: $s_i = \log_{10} \text{dose}$.

In this way, the Perturbation Perceiver framework can incorporate quantitative measures of perturbation
strength across types, providing a more nuanced and biologically informed representation of perturbations.

Cell Encoder

Original cell input modelling

In the Cell Encoder, the input original cell is represented with three types of tokens: identity tokens, value
tokens, and impact tokens. Following the design philosophy of previous single-cell foundation models, we
treat the gene expression profile of a cell analogously to a sentence, where each gene corresponds to a word.
Identity tokens are defined as gene identifiers, and we adopt the vocabulary from scGPT[20] to represent them as
the scGPT model supports efficient training. For genes not present in the original vocabulary, we extend the
vocabulary by adding the new gene identifiers and initialize their embeddings using a uniform distribution
matched to the mean and variance of the pretrained gene embeddings. For a given cell, the identity token
sequence is thus represented as:

$$\boldsymbol{g} = [\text{id}(g_1), \text{id}(g_2), \dots, \text{id}(g_m)],$$

588 where $\text{id}(g_i)$ denotes the index of g_i in the vocabulary and m is the total number of genes.

589   The value tokens represent the expression levels of the corresponding genes and are denoted as

$$\boldsymbol{v} = [v_1, v_2, \dots, v_m],$$

591 where v_i is typically the log-normalized expression value of gene i.

592   The impact tokens are designed to capture perturbation–gene interactions and vary depending on the
593 perturbation type. For genetic perturbations, they are defined as

594
$$p = p_g = [p_1, p_2, \ldots, p_m],$$

595
$$p_i = \begin{cases} 1, \text{if gene } i \text{ is perturbed} \\ \quad 0, \text{else} \end{cases}$$

596   For chemical perturbations, they are defined as

597
$$p = p_m = [p_1, p_2, \ldots, p_m],$$

598   where p_i represents the affinity score between the protein encoded by gene i and the input molecule,
599   calculated by inputting the protein sequence and the chemical SMILES string into a pretrained GraphDTA
600   model.

601   The final embedding for an original cell is then obtained by summing the three projected components:

602
$$h = \text{embed}_{\text{id}}(g) + \text{embed}_{\text{value}}(v) + \text{embed}_{\text{impact}}(p),$$

where $\text{embed}_{\text{id}}(\cdot)$ projects gene identifiers into embeddings, $\text{embed}_{\text{value}}(\cdot)$ is a nonlinear neural network
mapping continuous expression values into embeddings, and $\text{embed}_{\text{impact}}(\cdot)$ projects either binary
perturbation indicators or affinity scores into embeddings depending on the perturbation type. All embedding
dimensions are set to 512, consistent with the Perturbation Perceiver.

Transformer layers

The transformer layers in the Cell Encoder are designed to integrate the original cell input with perturbation
information. Each layer is composed of two modules: a gated cross-attention layer that incorporates
perturbation signals into the gene-expression representation, and a self-attention layer that models
dependencies among genes.

In the gated cross-attention layer, cross-attention is performed between gene-expression tokens and
perturbation embeddings, with an additional gating mechanism that adaptively regulates the strength of
perturbation influence. Formally, given the perturbation embeddings X and the l-th layer output of gene-
expression tokens $H^{(l)} \in \mathbb{R}^{m \times d}$, the cross-attention is defined as:

616
$$Q_c = H^{(l)} W_Q^c, K_c = X W_Q^c, V_c = X W_V^c,$$

617
$$C = \text{softmax}\left(\frac{Q_c K_c^T}{\sqrt{d}}\right) V_c.$$

618   The gating mechanism is then applied to adaptively regulate the perturbation contribution:

619
$$\widetilde{H} = \text{LN}\left(H^{(l)} + \tanh\left(H^{(l)} W_{g1} + b_{g1}\right) \odot C\right),$$

620   where $\text{LN}(\cdot)$ denotes layer normalization. The gated features are further refined with a gated feed-forward
621   network:

622
$$H_c = \text{LN}\left(\widetilde{H} + \tanh(\widetilde{H}W_{g2} + b_{g2}) \odot \text{FFN}(\widetilde{H})\right).$$

623 The output H_c is then passed through a standard Transformer encoder block to capture gene–gene
624 dependencies. Multi-head self-attention is applied as:

625
$$Q = H_c W_Q, K = H_c W_K, V = H_c W_V,$$

626
$$S = \text{softmax}\left(\frac{QK^{\text{T}}}{\sqrt{d_k}}\right)V.$$

627 Residual connections and a feed-forward layer are applied to update the representation:

628
$$\widehat{H} = \text{LN}(H_c + S),$$

629
$$H^{(l+1)} = \text{LN}\left(\widehat{H} + \text{FFN}(\widehat{H})\right).$$

630 All W_* and b_* are trainable parameters. In summary, the entire encoder layer alternates between gated
631 cross-attention and self-attention modules:

632
$$H = H^{(0)},$$

633
$$H^{(l+1)} = \text{SelfAttnBlock}\left(\text{CrossAttnBlock}(H^{(l)})\right), \forall l \in [0, n_l - 1].$$

In this study, we set the number of encoder layers to $n_l = 12$, with 8 attention heads in each multi-head
attention module.

Gene expression value decoder

The final output of the transformer layers is passed to the gene expression decoder, which predicts the
complete gene expression profile of the perturbed cell. Let the output of the last transformer layer be denoted
as $H^{(n_l)} = [h_1^{(n_l)}, h_2^{(n_l)}, ..., h_m^{(n_l)}]$, where $h_i^{(n_l)}$ represents the latent embedding corresponding to gene i, and
m is the total number of genes. For each output token, the decoder projects the latent representation into a
scalar expression value:

642
$$\hat{y}_i = f_d\left(h_i^{(n_l)}\right),$$

643 where $f_d(\cdot)$ is a nonlinear projection layer implemented as a neural network.

644 Model training is performed by minimizing the mean squared error (MSE) loss between predicted and
645 ground-truth perturbed expression values:

646
$$\mathcal{L}_{\text{MSE}} = \frac{1}{m}\sum_{i=1}^{m}(y_i - \hat{y}_i)^2,$$

where y_i denotes the experimentally measured expression level of gene i under perturbation, and $\hat{y}_i$ is the

corresponding model prediction. This objective directly enforces accurate reconstruction of perturbation-

induced transcriptional responses at the genome-wide level.

Co-training across different perturbation types

For the X-Pert model trained jointly on genetic and chemical perturbations, we denote k as the total number

of samples in a mini-batch, with k_g and k_m representing the number of genetic and chemical perturbation

samples, respectively. To encourage alignment between perturbation types, we incorporate a Maximum Mean

Discrepancy (MMD) loss that minimizes the distributional gap between the latent embeddings of gene

perturbations and drug perturbations. The MMD loss is applied on the outputs of the perturbation-type–

specific encoders, thereby enforcing the latent space to capture perturbation representations in a type-agnostic

manner.

Formally, the MMD loss is defined as the squared distance between the kernel mean embeddings of the two

distributions in a reproducing kernel Hilbert space (RKHS):

661
$$
\mathcal{L}_{\text{MMD}}\left(\widehat{X}_p^g, \widehat{X}_p^m\right) = \frac{1}{k_g(k_g - 1)}\sum_{i \neq i'} f_k\left(x_i^g, x_{i'}^g\right) + \frac{1}{k_m(k_m - 1)}\sum_{i \neq i'} f_k\left(x_i^m, x_{i'}^m\right) - \frac{2}{k_g k_m}\sum_{i=1}^{k_g}\sum_{j=1}^{k_m} f_k\left(x_i^g, x_j^m\right),
$$

662 where $f_k(\cdot,\cdot)$ is the kernel function. We adopt the radial basis function (RBF) kernel, also known as the

663 Gaussian kernel, defined as:

664
$$
f_k(x, y) = \exp\left(-\frac{||x - y||^2}{2\sigma^2}\right),
$$

665 where $||x - y||^2$ denotes the squared Euclidean distance between vectors x and y, and $\sigma > 0$ is the

666 kernel bandwidth controlling the smoothness of the similarity measure. Following common practice, we set σ

667 using the median heuristic, which has been shown to yield stable performance in high-dimensional spaces.

668 The overall training objective combines the MSE loss with the alignment loss:

669
$$
\mathcal{L}_{\text{total}} = \mathcal{L}_{\text{MSE}} + \alpha \mathcal{L}_{\text{MMD}},
$$

where α is a balancing hyperparameter. In this study, we set $\alpha = 0.2$, which we found to provide a good

trade-off between accurate reconstruction and effective cross–perturbation-type alignment.

Perturbverse embedding extraction

We use the output of the Perturbation Perceiver, $X \in \mathbb{R}^{n_l \times d}$, as the Perturbverse representation of the input

perturbations. Since the number of latent queries n_l is fixed at 12, this design ensures that perturbations of

any type, regardless of their original perturbation type or number of input tokens, are consistently projected
into a standardized latent space of dimension 12×512. To obtain a compact perturbation embedding, the latent
tokens are subsequently flattened into a 6,144-dimensional vector, which serves as the unified latent
representation of the perturbation.

Model evaluation

We used several metrics to perform the model evaluation and application.

Pearson correlation coefficient (PCC)

The PCC measures the linear correlation between predicted $(\hat{y})$ and observed (y) values:

685
$$\text{PCC}(y, \hat{y}) = \frac{\sum_{i=1}^{m}(y_i - \bar{y})(\hat{y}_i - \bar{\hat{y}})}{\sqrt{\sum_{i=1}^{m}(y_i - \bar{y})^2}\sqrt{\sum_{i=1}^{m}(\hat{y}_i - \bar{\hat{y}})^2}},$$

686 where $\bar{y}$ and $\bar{\hat{y}}$ denote the means of the true and predicted values, respectively.

687 **Coefficient of determination (R^2)**

688 The R^2 score quantifies the proportion of variance in the observed data explained by the predictions:

689
$$R^2(y, \hat{y}) = 1 - \frac{\sum_{i=1}^{m}(y_i - \hat{y}_i)^2}{\sum_{i=1}^{m}(y_i - \bar{y})^2}.$$

690 Higher values indicate stronger predictive power, with $R^2 = 1$ representing perfect prediction.

691 **Energy distance (E-distance)**

692 The E-distance evaluates the dissimilarity between the distributions of perturbed and original cells, serving as
693 a measure of perturbation effect size. Given two sets of samples $X = \{x_1, \dots, x_n\}$ and $Y = \{y_1, \dots, y_m\}$, the
694 E-distance is defined as:

695
$$\text{E}(X, Y) = \frac{2}{nm}\sum_{i=1}^{n}\sum_{j=1}^{m}||x_i - y_j|| - \frac{1}{n^2}\sum_{i=1}^{n}\sum_{j=1}^{m}||x_i - x_j|| - \frac{1}{m^2}\sum_{i=1}^{n}\sum_{j=1}^{m}||y_i - y_j||.$$

696 We use the scPerturb package to calculate the E-distance.

697 **Cosine similarity**

698 Cosine similarity measures the angle between two vectors and evaluates their directional similarity:

$$699 \quad \mathrm{CosSim}(y, \hat{y}) = \frac{\sum_{i=1}^{m} y_i\, \hat{y}_i}{\sqrt{\sum_{i=1}^{m} y_i^2}\ \sqrt{\sum_{i=1}^{m} \hat{y}_i^2}}$$

700   This metric is scale-invariant and emphasizes relative patterns rather than absolute magnitude.

Perturbation quantification with Hallmark gene sets

For each genetic perturbation, we evaluated pathway-level transcriptional changes by calculating gene set

scores across all single cells. We used the Hallmark gene sets[40] from the Molecular Signatures Database

(MSigDB), which consist of 50 curated collections that capture and summarize specific, well-defined

biological processes and cell states. These gene sets provide a compact and biologically interpretable

representation of transcriptional programs, allowing us to assess whether perturbations induce consistent

pathway-level responses. Gene set scores were computed using the Scanpy package[93], specifically the

'sc.tl.score_genes' function, which calculates a per-cell enrichment score by comparing the expression of

genes within each gene set against a reference background.

KNN accuracy

To quantitatively assess the discriminative ability of the learned Perturbverse embeddings (Fig. S2A), we

employed a *K*-nearest neighbor (KNN) classification framework with stratified five-fold cross-validation.

Specifically, a KNN classifier ($K = 5$) was trained on the PCA-transformed feature space, and the resulting

predictions were evaluated using confusion matrices aggregated across folds. For each class (corresponding to

a pathway label), a per-class accuracy was computed as the ratio of correctly classified samples to the total

number of true samples in that class. This metric provides a fine-grained evaluation of how well the

embedding preserves class-specific separability, independent of class imbalance.

Databases and datasets

Candidate drug databases

We employed three different drug databases as candidate libraries to support perturbation retrieval and *in*

silico drug screening tasks. For the drug perturbation retrieval experiment (Fig. 4G), we used the HERB 2.0

database[46], which contains 30,916 natural compounds with curated annotations of chemical structures and

pharmacological information. For the *in silico* drug screening task (Fig. 5D), we used an FDA-approved drug

library from TargetMol[94], consisting of 935 clinically approved compounds, ensuring the relevance of

predictions to translational applications. Finally, for the cross-type perturbation retrieval analysis (Fig. 6E), we

utilized a unique collection of 1,930 monomeric compounds derived from traditional Chinese medicine

(TCM) curated by SelleckChem[83], which provides structurally diverse natural products not typically

represented in standard drug libraries.

Genetic and chemical perturbation data

We benchmarked X-Pert against multiple competing methods on genetic perturbation prediction using four

datasets: Replogle2022_K562[4], Replogle2022_RPE1[4], and Norman2019[35]. The raw data of all the datasets

were obtained from the scPerturb[16] database.

For chemical perturbation benchmarking, we employed the Sciplex-3[11] and CMAP[44] datasets. The Sciplex-

3 dataset is a single-cell perturbation dataset comprising 758,360 processed cells across three cell lines and

188 small-molecule perturbations after preprocessing. The CMAP dataset, which provides bulk-level gene

expression profiles, contains 678,401 samples spanning 70 cell lines and 17,775 chemical perturbations after

preprocessing.

All the datasets are processed with the same procedure following our previous benchmark[38]:

•   Feature Selection: Top 5,000 highly variable genes (HVGs) were identified from each dataset using

standard procedures. We also included all perturbed genes originally present in the dataset's feature list.

•   Normalization: Raw count data was normalized using the log-normalization method, implemented

through the standard preprocessing workflow of Scanpy[93].

•   Original-cell pairing: For every cell subjected to perturbation, we randomly sampled an original cell from

the same dataset to form a pre- and post-perturbation cell pair. These pairs served as the basis for training

and evaluating models.

•   Identifying differentially expressed genes (DEGs): For each perturbation, we identified DEGs by

comparing perturbed cells to original cells. Genes were ranked based on adjusted p-values computed

using the Wilcoxon test in Scanpy, with those having adjusted p-values < 0.05 designated as DEGs.

Tahoe-100M

Tahoe-100M is an exceptionally large-scale chemical perturbation dataset comprising approximately one

million single cells. To reduce data redundancy and facilitate efficient training, we aggregated single-cell

profiles into pseudo-bulk samples: for each combination for drug, cell line, and dose, the expression values of

all corresponding single cells were averaged to yield a representative profile. After this preprocessing step, a

total of 65,768 pseudo-bulk samples were retained for model training. To ensure that the model focuses on

biologically informative signals, we further selected 10,000 highly variable genes (HVGs) across all samples

as the feature set. This dimensionality reduction strategy retains the genes with the largest expression

variability, thereby capturing the most relevant transcriptional changes induced by perturbations while

reducing noise from less informative features.

***In silico* drug screening with X-Pert**

To perform *in silico* drug screening, we combined disease-associated gene signatures with X-Pert–predicted
gene expression profiles under drug perturbations. Disease signatures were obtained from the CRowd
Extracted Expression of Differential Signatures (CREEDS) database[95], which contains 839 annotated disease-
versus-normal transcriptional signatures. For each disease, CREEDS provides two complementary gene sets:
one consisting of up-regulated genes associated with the disease state, and another consisting of down-
regulated genes. The rationale of our approach is that a drug with therapeutic potential should simultaneously
suppress the disease-associated up-regulated genes while inducing the disease-associated down-regulated
genes, thereby reversing the pathological transcriptional program.

We applied this framework to cell lines with defined disease associations in Tahoe-100M. For example, in
the HepG2 cell line (originating from hepatocellular carcinoma, HCC), we used X-Pert to predict
transcriptional responses to 831 candidate drugs across the 10,000 highly variable genes selected from the
dataset. This produced drug-specific predicted gene expression profiles for HepG2 cells. The corresponding
HCC gene signatures from CREEDS were then used to evaluate these predictions. Specifically, we computed
two complementary enrichment scores using the Scanpy function 'sc.tl.score_genes': the up-regulated gene
set score (up-score) and the down-regulated gene set score (down-score). Intuitively, lower up-scores and
higher down-scores represent drugs with higher potential that could reverse disease signatures.

Large-scale chemical perturbation datasets such as Tahoe-100M are subject to technical batch effects,
which in this case primarily arise from sequencing plate variability (Fig. 5B). These batch effects introduce
systematic biases into X-Pert's predictions, leading to spurious correlations between the calculated up-scores
and down-scores. As a result, high up-scores often co-occur with proportionally high down-scores (Fig. 5D).
This phenomenon is likely attributable to plate-specific biases in total transcript counts across different drug
conditions, which can confound the biological signal and distort the prioritization of candidate drugs.

To mitigate these effects, we implemented a simple yet effective correction strategy based on linear
regression. Specifically, we fit a regression line between the up-scores and down-scores across all candidate
drugs for a given disease. The fitted line captures the dominant technical trend introduced by batch effects. We
then identify drugs with the largest positive residuals (intercepts above the fitted line) as top candidates, since
these drugs deviate most strongly from the batch-driven correlation and therefore likely reflect genuine
biological activity rather than technical artifacts.

Drugs achieving large positive residuals (simultaneously low up-scores and high down-scores) are
prioritized as candidate therapeutic compounds for the corresponding disease. This strategy allows us to
systematically evaluate the potential of existing drugs to reverse disease-specific transcriptional dysregulation
and highlights X-Pert's utility in drug repurposing.

GDIdb drugs filtering

For the cross-type perturbation retrieval evaluation, we selected drugs known to interact with eight
representative genes—AKT1, AKT2, CTSK, SMAD3, TP53, MTOR, BRAF, and PIK3CA—as the ground-
truth references. For each gene, we obtained its interacting drugs from the DGIdb database[66], which provides
curated drug–gene interactions along with an interaction score that quantifies the strength of evidence
supporting each relationship. To ensure reliability, we applied a threshold of 0.2 on the interaction score and
retained only those drug–gene pairs exceeding this cutoff. After filtering, the number of drugs associated with
each gene ranged from 7 to 23, forming the reference sets for evaluation.

Data availability

805 The genetic perturbation data (Replogle2022_K562, Replogle2022_RPE1, and Norman2019) were
806 downloaded from https://zenodo.org/records/10044268. The Sciplex-3 data were accessed from
807 https://figshare.com/articles/dataset/sciplex3/22122701. The CMAP data was downloaded from Gene
808 Expression Omnibus (GEO) with accession number GSE92742. The Tahoe-100M data were downloaded from
809 https://huggingface.co/datasets/tahoebio/Tahoe-100M.

Code availability

812 The code and documents of X-Pert are available at https://github.com/Chen-Li-17/X-Pert.

Acknowledgements

815 The work is sponsored by The National Key R&D Program of China (2025YFC3409300), National Natural
816 Science Foundation of China (62373210, 62433001, 92470105), and Tsinghua-Toyota Joint Research Institute
817 Inter-Disciplinary Program (20243930093).

Author contributions

820 C.L., L.W. and X.Z. conceived and designed the study. X.Z and L.W. supervised the study and provided
821 critical revisions. C.L. collected the data, build the model, conducted the experiments, and analyzed the
822 results. C.L. and L.W. drafted the manuscript and prepared the figures. X.Z. reviewed the manuscript and
823 provided critical insights. All authors contributed to reviewing and approving the final manuscript.

Competing interests

826	The authors declare no competing interests.

Reference

1. Dixit, A. et al. Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell RNA Profiling of Pooled Genetic Screens. *Cell* **167**, 1853-1866 e1817 (2016).

2. Adamson, B. et al. A Multiplexed Single-Cell CRISPR Screening Platform Enables Systematic Dissection of the Unfolded Protein Response. *Cell* **167**, 1867-1882 e1821 (2016).

3. Datlinger, P. et al. Pooled CRISPR screening with single-cell transcriptome readout. *Nat Methods* **14**, 297-301 (2017).

4. Replogle, J.M. et al. Mapping information-rich genotype-phenotype landscapes with genome-scale Perturb-seq. *Cell* **185**, 2559-2575 e2528 (2022).

5. Rubin, A.J. et al. Coupled Single-Cell CRISPR Screening and Epigenomic Profiling Reveals Causal Gene Regulatory Networks. *Cell* **176**, 361-376 e317 (2019).

6. Wessels, H.H. et al. Efficient combinatorial targeting of RNA transcripts in single cells with Cas13 RNA Perturb-seq. *Nat Methods* **20**, 86-94 (2023).

7. Otto, J.E. et al. Structural and functional properties of mSWI/SNF chromatin remodeling complexes revealed through single-cell perturbation screens. *Mol Cell* **83**, 1350-1367 e1357 (2023).

8. Aissa, A.F. et al. Single-cell transcriptional changes associated with drug tolerance and response to combination therapies in cancer. *Nat Commun* **12**, 1628 (2021).

9. Chang, M.T. et al. Identifying transcriptional programs underlying cancer drug response with TraCe-seq. *Nat Biotechnol* **40**, 86-93 (2022).

10. Zhao, W. et al. Deconvolution of cell type-specific drug responses in human tumor tissue with single-cell RNA-seq. *Genome Med* **13**, 82 (2021).

11. Srivatsan, S.R. et al. Massively multiplex chemical transcriptomics at single-cell resolution. *Science* **367**, 45-51 (2020).

12. Zhang, J. et al. Tahoe-100m: A giga-scale single-cell perturbation atlas for context-dependent gene function and cellular modeling. *bioRxiv*, https://doi.org/10.1101/2025.1102.1120.639398 (2025).

13. Papalexi, E. et al. Characterizing the molecular regulation of inhibitory immune checkpoints with multimodal single-cell screens. *Nat Genet* **53**, 322-331 (2021).

14. Frangieh, C.J. et al. Multimodal pooled Perturb-CITE-seq screens in patient models define mechanisms of cancer immune evasion. *Nat Genet* **53**, 332-341 (2021).

15. Yang, B. et al. PerturbDB for unraveling gene functions and regulatory networks. *Nucleic Acids Res* **53**, D1120-D1131 (2025).

16. Peidli, S. et al. scPerturb: harmonized single-cell perturbation data. *Nat Methods* **21**, 531-540 (2024).

17. Roohani, Y., Huang, K. & Leskovec, J. Predicting transcriptional outcomes of novel multigene perturbations with GEARS. *Nat Biotechnol* **42**, 927-935 (2024).

18. Bai, D., Ellington, C.N., Mo, S., Song, L. & Xing, E.P. AttentionPert: accurately modeling multiplexed genetic perturbations with multi-scale effects. *Bioinformatics* **40**, i453-i461 (2024).

19. Hao, M. et al. Large-scale foundation model on single-cell transcriptomics. *Nat Methods* **21**,

1481-1491 (2024).

20.   Cui, H. et al. scGPT: toward building a foundation model for single-cell multi-omics using
generative AI. *Nat Methods* **21**, 1470-1480 (2024).

21.   Schrod, S., Zacharias, H.U., Beissbarth, T., Hauschild, A.C. & Altenbuchinger, M. CODEX:
COunterfactual Deep learning for the in silico EXploration of cancer cell line perturbations.
Bioinformatics **40**, i91-i99 (2024).

22.   Yu, H., Qian, W., Song, Y. & Welch, J.D. PerturbNet predicts single-cell responses to unseen
chemical and genetic perturbations. *Mol Syst Biol* **21**, 960-982 (2025).

23.   Xu, Y., Fleming, S., Tegtmeyer, M., McCarroll, S.A. & Babadi, M. Explainable modeling of
single-cell perturbation data using attention and sparse dictionary learning. *Cell Syst* **16**,
101245 (2025).

24.   Hetzel, L. et al. Predicting Cellular Responses to Novel Drug Perturbations at a Single-Cell
Resolution [C]. Proceedings of the 36th Conference on Neural Information Processing
Systems (NeurIPS). 2022.

25.   Piran, Z., Cohen, N., Hoshen, Y. & Nitzan, M. Disentanglement of single-cell data with
biolord. *Nat Biotechnol* **42**, 1678-1683 (2024).

26.   Qi, X. et al. Predicting transcriptional responses to novel chemical perturbations using deep
generative model for drug discovery. *Nat Commun* **15**, 9256 (2024).

27.   Maleki, S. et al. Efficient Fine-Tuning of Single-Cell Foundation Models Enables Zero-Shot
Molecular Perturbation Prediction. *arXiv preprint arXiv:2412.13478* (2024).

28.   Klein, D. et al. CellFlow enables generative single-cell phenotype modeling with flow
matching. *bioRxiv*, https://doi.org/10.1101/2025.1104.1111.648220 (2025).

29.   Adduri, A.K. et al. Predicting cellular responses to perturbation across diverse contexts with
STATE. *bioRxiv*, https://doi.org/10.1101/2025.1106.1126.661135 (2025).

30.   Li, C. et al. TFcomb identifies transcription factor combinations for cellular reprogramming
based on single-cell multiomics data. *Genome Res* **35**, 1429-1439 (2025).

31.   Cui, A. et al. Dictionary of immune responses to cytokines at single-cell resolution. *Nature*
625, 377-384 (2024).

32.   OpenAI GPT-4 Technical Report. *arXiv:2303.08774* (2023).

33.   Bento, A.P. et al. An open source chemical structure curation pipeline using RDKit. *J
Cheminform* **12**, 51 (2020).

34.   Nguyen, T. et al. GraphDTA: predicting drug-target binding affinity with graph neural
networks. *Bioinformatics* **37**, 1140-1147 (2021).

35.   Norman, T.M. et al. Exploring genetic interaction manifolds constructed from rich single-cell
phenotypes. *Science* **365**, 786-793 (2019).

36.   Ahlmann-Eltze, C., Huber, W. & Anders, S.J.N.M. Deep-learning-based gene perturbation
effect prediction does not yet outperform simple linear baselines. *Nat Methods*, 1-5 (2025).

37.   Kamimoto, K. et al. Dissecting cell identity via network inference and in silico gene
perturbation. *Nature* **614**, 742-751 (2023).

38.   Li, C. et al. Benchmarking AI models for in silico gene perturbation of cells. *bioRxiv*,
https://doi.org/10.1101/2024.1112.1120.629581 (2024).

39.   Vinas Torne, R. et al. Systema: a framework for evaluating genetic perturbation response

prediction beyond systematic variation. *Nat Biotechnol* (2025).

40.   Liberzon, A. et al. The Molecular Signatures Database (MSigDB) hallmark gene set
collection. *Cell Syst* **1**, 417-425 (2015).

41.   Radhakrishnan, S.K. et al. Transcription factor Nrf1 mediates the proteasome recovery
pathway after proteasome inhibition in mammalian cells. *Mol Cell* **38**, 17-28 (2010).

42.   Oron, M. et al. The molecular network of the proteasome machinery inhibition response is
orchestrated by HSP70, revealing vulnerabilities in cancer cells. *Cell Rep* **40**, 111428 (2022).

43.   Volmar, C.H. et al. M344 promotes nonamyloidogenic amyloid precursor protein processing
while normalizing Alzheimer's disease genes and improving memory. *Proc Natl Acad Sci U S*
A **114**, E9135-E9144 (2017).

44.   Subramanian, A. et al. A Next Generation Connectivity Map: L1000 Platform and the First
1,000,000 Profiles. *Cell* **171**, 1437-1452 e1417 (2017).

45.   Knox, C. et al. DrugBank 6.0: the DrugBank Knowledgebase for 2024. *Nucleic Acids Res* **52**,
D1265-D1275 (2024).

46.   Gao, K. et al. HERB 2.0: an updated database integrating clinical and experimental evidence
for traditional Chinese medicine. *Nucleic Acids Res* **53**, D1404-D1414 (2025).

47.   Lee, H.K., Oh, S.R., Kim, J.I., Kim, J.W. & Lee, C.O. Agastaquinone, a new cytotoxic
diterpenoid quinone from Agastache rugosa. *J Nat Prod* **58**, 1718-1721 (1995).

48.   Mosca, L., Ilari, A., Fazi, F., Assaraf, Y.G. & Colotti, G. Taxanes in cancer treatment: Activity,
chemoresistance and its overcoming. *Drug Resist Updat* **54**, 100742 (2021).

49.   Schneider, F., Samarin, K., Zanella, S. & Gaich, T. Total synthesis of the complex taxane
diterpene canataxpropellane. *Science* **367**, 676-681 (2020).

50.   Gupta, A., Atanasov, A.G., Li, Y., Kumar, N. & Bishayee, A. Chlorogenic acid for cancer
prevention and therapy: Current status on efficacy and mechanisms of action. *Pharmacol*
Res **186**, 106505 (2022).

51.   Huang, S. et al. Chlorogenic acid effectively treats cancers through induction of cancer cell
differentiation. *Theranostics* **9**, 6745-6763 (2019).

52.   Ibello, E. et al. Three-dimensional environment sensitizes pancreatic cancer cells to the anti-
proliferative effect of budesonide by reprogramming energy metabolism. *J Exp Clin Cancer*
Res **43**, 165 (2024).

53.   Yalniz, F.F. et al. A phase II study of addition of pracinostat to a hypomethylating agent in
patients with myelodysplastic syndromes who have not responded to previous
hypomethylating agent therapy. *British journal of haematology* **188**, 404-412 (2020).

54.   Sharman, J. et al. An open-label phase 2 trial of entospletinib (GS-9973), a selective spleen
tyrosine kinase inhibitor, in chronic lymphocytic leukemia. *Blood* **125**, 2336-2343 (2015).

55.   Joshi, S. New insights into SYK targeting in solid tumors. *Trends Pharmacol Sci* **45**, 904-918
(2024).

56.   Rohila, D. et al. Syk Inhibition Reprograms Tumor-Associated Macrophages and Overcomes
Gemcitabine-Induced Immunosuppression in Pancreatic Ductal Adenocarcinoma. *Cancer*
Res **83**, 2675-2689 (2023).

57.   Varshosaz, J., Fard, M.M., Mirian, M. & Hassanzadeh, F. Targeted Nanoparticles for Co-
delivery of 5-FU and Nitroxoline, a Cathepsin B Inhibitor, in HepG2 Cells of Hepatocellular
Carcinoma. *Anticancer Agents Med Chem* **20**, 346-358 (2020).

958 58. Choi, S.M. et al. Clioquinol, a Cu(II)/Zn(II) chelator, inhibits both ubiquitination and
asparagine hydroxylation of hypoxia-inducible factor-1alpha, leading to expression of
vascular endothelial growth factor and erythropoietin in normoxic cells. *J Biol Chem* **281**,
34056-34063 (2006).

59. Huang, Z., Wang, L., Chen, L., Zhang, Y. & Shi, P. Induction of cell cycle arrest via the p21,
p27-cyclin E,A/Cdk2 pathway in SMMC-7721 hepatoma cells by clioquinol. *Acta Pharm* **65**,
463-471 (2015).

60. White, J.R., Jr. Empagliflozin, an SGLT2 inhibitor for the treatment of type 2 diabetes
mellitus: a review of the evidence. *Ann Pharmacother* **49**, 582-598 (2015).

61. Abdelhamid, A.M. et al. Empagliflozin adjunct with metformin for the inhibition of
hepatocellular carcinoma progression: Emerging approach for new application. *Biomed*
Pharmacother **145**, 112455 (2022).

62. Scafoglio, C.R. et al. Sodium-glucose transporter 2 is a diagnostic and therapeutic target for
early-stage lung adenocarcinoma. *Sci Transl Med* **10** (2018).

63. Luo, J., Hendryx, M. & Dong, Y. Sodium-glucose cotransporter 2 (SGLT2) inhibitors and non-
small cell lung cancer survival. *Br J Cancer* **128**, 1541-1547 (2023).

64. Lu, J. et al. Ciclopirox targets cellular bioenergetics and activates ER stress to induce
apoptosis in non-small cell lung cancer cells. *Cell Commun Signal* **20**, 37 (2022).

65. Ramalingam, S. et al. A Randomized, Double-Blind, Phase 2 Trial of Veliparib (ABT-888)
With Carboplatin and Paclitaxel in Previously Untreated Metastatic or Advanced Non-Small
Cell Lung Cancer. *International Journal of Radiation Oncology*Biology*Physics* **90**, S4-S5
(2014).

66. Cannon, M. et al. DGIdb 5.0: rebuilding the drug-gene interaction database for precision
medicine and drug discovery platforms. *Nucleic Acids Res* **52**, D1227-D1235 (2024).

67. Dobelis, P., Madl, J.E., Pfister, J.A., Manners, G.D. & Walrond, J.P. Effects of Delphinium
alkaloids on neuromuscular transmission. *J Pharmacol Exp Ther* **291**, 538-546 (1999).

68. Ahn, E.J. et al. Genotype-Phenotype Correlation of SMN1 and NAIP Deletions in Korean
Patients with Spinal Muscular Atrophy. *J Clin Neurol* **13**, 27-31 (2017).

69. Lin, J., Wang, Q., Zhou, S., Xu, S. & Yao, K. Tetramethylpyrazine: A review on its
mechanisms and functions. *Biomed Pharmacother* **150**, 113005 (2022).

70. Xue, Z. et al. MAP3K1 and MAP2K4 mutations are associated with sensitivity to MEK
inhibitors in multiple cancer models. *Cell Res* **28**, 719-729 (2018).

71. Velasco-Velazquez, M. et al. CCR5 antagonist blocks metastasis of basal breast cancer
cells. *Cancer Res* **72**, 3839-3850 (2012).

72. Berr, A.L. et al. Vimentin is required for tumor progression and metastasis in a mouse model
of non-small cell lung cancer. *Oncogene* **42**, 2074-2087 (2023).

73. Perez-Sala, D., Gilbert, B.A., Rando, R.R. & Canada, F.J. Analogs of farnesylcysteine induce
apoptosis in HL-60 cells. *FEBS Lett* **426**, 319-324 (1998).

74. Huang, S. et al. Ras guanine nucleotide-releasing protein-4 promotes renal inflammatory
injury in type 2 diabetes mellitus. *Metabolism* **131**, 155177 (2022).

75. Weng, L.Q. et al. Aliskiren ameliorates pressure overload-induced heart hypertrophy and
fibrosis in mice. *Acta Pharmacol Sin* **35**, 1005-1014 (2014).

76. Zhao, K. et al. Nuclear import of Mas-related G protein-coupled receptor member D induces
pathological cardiac remodeling. *Cell Commun Signal* **21**, 181 (2023).

77. Lan, L. et al. Mas-related G protein-coupled receptor D participates in inflammatory pain by promoting NF-kappaB activation through interaction with TAK1 and IKK complex. *Cell Signal* **76**, 109813 (2020).

78. Song, C. et al. Limonin ameliorates dextran sulfate sodium-induced chronic colitis in mice by inhibiting PERK-ATF4-CHOP pathway of ER stress and NF-kappaB signaling. *Int Immunopharmacol* **90**, 107161 (2021).

79. Scheinman, R.I., Gualberto, A., Jewell, C.M., Cidlowski, J.A. & Baldwin, A.S., Jr. Characterization of mechanisms involved in transrepression of NF-kappa B by activated glucocorticoid receptors. *Mol Cell Biol* **15**, 943-953 (1995).

80. Brenachot, X. et al. Hepatic protein tyrosine phosphatase receptor gamma links obesity-induced inflammation to insulin resistance. *Nat Commun* **8**, 1820 (2017).

81. Lin, J., Li, Q., Lei, X. & Zhao, H. The emerging roles of GPR158 in the regulation of the endocrine system. *Front Cell Dev Biol* **10**, 1034348 (2022).

82. Chapdelaine, A.G., Ku, G.C., Sun, G. & Ayrapetov, M.K. The Targeted Degradation of BRAF V600E Reveals the Mechanisms of Resistance to BRAF-Targeted Treatments in Colorectal Cancer Cells. *Cancers (Basel)* **15** (2023).

83. Selleck products. https://www.selleckchem.com/ (2025).

84. Chen, T., Yang, P. & Jia, Y. Molecular mechanisms of astragaloside-IV in cancer therapy (Review). *Int J Mol Med* **47** (2021).

85. Zhang, Y. et al. Glycyrrhetinic acid binds to the conserved P-loop region and interferes with the interaction of RAS-effector proteins. *Acta Pharm Sin B* **9**, 294-303 (2019).

86. Fu, T., Chai, B., Shi, Y., Dang, Y. & Ye, X. Fargesin inhibits melanin synthesis in murine malignant and immortalized melanocytes by regulating PKA/CREB and P38/MAPK signaling pathways. *J Dermatol Sci* **94**, 213-219 (2019).

87. Shi, A., Liu, L., Li, S. & Qi, B. Natural products targeting the MAPK-signaling pathway in cancer: overview. *J Cancer Res Clin Oncol* **150**, 6 (2024).

88. Metzner, E., Southard, K.M. & Norman, T.M. Multiome Perturb-seq unlocks scalable discovery of integrated perturbation effects on the transcriptome and epigenome. *Cell Syst* **16**, 101161 (2025).

89. Saunders, R.A. et al. Perturb-Multimodal: A platform for pooled genetic screens with imaging and sequencing in intact mammalian tissue. *Cell* **188**, 4790-4809 e4722 (2025).

90. Wu, J., Wan, C., Ji, Z., Zhou, Y. & Hou, W. EpiFoundation: A Foundation Model for Single-Cell ATAC-seq via Peak-to-Gene Alignment. *bioRxiv* (2025).

91. Li, W. et al. scPROTEIN: a versatile deep graph contrastive learning framework for single-cell proteomics embedding. *Nat Methods* **21**, 623-634 (2024).

92. Alayrac, J.B. et al. Flamingo: a Visual Language Model for Few-Shot Learning [C]. Proceedings of the 36th Conference on Neural Information Processing Systems (NeurIPS). 2022.

93. Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol* **19**, 15 (2018).

94. TargetMol. https://www.targetmol.com (2025).

95. Wang, Z. et al. Extraction and analysis of signatures from the Gene Expression Omnibus by the crowd. *Nat Commun* **7**, 12846 (2016).

1041