Tools for genetic manipulation of the endemic fungal pathogen, *Emergomyces africanus,* and the application of a fluorescent reporter strain in infection models

Lucian Duvenage[1,2,3]# , Alisha Chetty[1,2,3], Darren D. Thomson[4], Elizabeth R. Ballou[4], Nelesh P. Govender[5,6], Chad A. Rappleye[7]*, J. Claire Hoving[1,2,3]*

1. CMM AFRICA Medical Mycology Research Unit, Institute of Infectious Disease and Molecular Medicine, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa
2. Wellcome Centre for Infectious Diseases Research in Africa, Institute of Infectious Diseases and Molecular Medicine, University of Cape Town, Cape Town, South Africa
3. Division of Immunology, Department of Pathology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa
4. MRC Centre for Medical Mycology, The University of Exeter, Exeter, United Kingdom.
5. Wits Mycology Division, School of Pathology, Faculty of Health Sciences, University of the Witwatersrand, Parktown, Gauteng, South Africa.
6. National Institute for Communicable Diseases, A Division of the National Health Laboratory Service, Johannesburg, Gauteng, South Africa
7. Department of Microbiology, Ohio State University, Columbus, OH, USA

*(CAR and JCH are co-senior authors)

Address correspondence to: Lucian Duvenage, lucian.duvenage@uct.ac.za

Running title: Molecular tools for Emergomyces africanus

Abstract

*Emergomyce*s *africanus* is a thermally dimorphic fungal pathogen endemic to Southern Africa, which can cause fatal systemic infections in persons with advanced HIV disease. Its mechanisms of pathogenesis are not well understood. Characterisation of virulence traits in this pathogen requires appropriate molecular tools for genetic manipulation. Molecular technologies developed for the transformation of *H. capsulatum* were adapted for use in *E. africanus. Agrobacterium*-mediated transformation was used to generate a reporter strain expressing green fluorescent protein (GFP). The *E. africanus* GFP reporter strain facilitated the study of yeast interaction with macrophages *in vitro* and allowed the identification of infected phagocyte cell types in the mouse lung by flow cytometry. *E. africanus* could also maintain episomal plasmids with telomere-like sequences to introduce expression constructs without genome modification. Using this plasmid system, RNA interference constructs were used to knock down the expression of cell wall α(1,3)-glucan by targeting the transcripts of the α-glucan synthase (*AGS1*). An episomal CRISPR/Cas9 system was evaluated for *E. africanus*, which effectively disrupted *GFP* in a reporter strain and enabled the generation of a *URA5* uracil auxotroph. These tools and strains will facilitate future studies to elucidate the mechanisms of pathogenesis of *E. africanus.*

Importance

Emergomyces africanus is an opportunistic fungal pathogen affecting persons with advanced HIV disease in South Africa. The biology and pathogenesis of *E. africanus*

are not well understood, as the importance of the disease caused by this fungus

(emergomycosis) has only been recognised in recent years, and molecular studies

have been impaired by the lack of genetic technologies. In this work, we describe tools

and methods for the genetic modification of this pathogen, which will accelerate future

studies investigating how the fungus causes disease in the human host. These

essential tools include (1) the ability to create fluorescent reporter strains, such as the

green fluorescent protein *E. africanus* strain described here, which facilitates tracking

the spread of the fungus during infection and enhances microscopy studies, (2)

methods for knocking down gene expression in *E. africanus*, and (3) the permanent

disruption of genes through CRISPR/Cas9 gene editing.

Introduction

Emergomyces africanus, part of the recently classified genus *Emergomyces* (1), is a

fungal pathogen of humans endemic to Southern Africa (2). The fungus can cause life-

threatening systemic infection in immunocompromised individuals, which, in South

Africa, has exclusively been persons with advanced HIV disease (3). Because this

pathogen is less often suspected by clinicians, patients affected by emergomycosis

often are diagnosed only at the late stages of infection, characterised by widespread

skin lesions. An understanding of its biology and host-pathogen interaction is needed

to develop effective diagnostics and treatment strategies for this infection in high-risk

populations.

E. africanus is a thermally dimorphic fungus. It exists in a saprobic mycelial phase in

soil and releases airborne conidia (4), which, when inhaled by the host, transition to a

yeast phase triggered by body temperature (5). The mechanisms by which these yeast

cells infect host cells are unknown. The ability to genetically manipulate a fungal

pathogen can open new avenues of research, whether by examining the effects of

gene deletions or introducing reporter genes for a wide range of applications, such as

facilitating the detection of the pathogen in infection models. New tools or protocols

must be developed or adapted for non-model fungi such as *E. africanus.* In this study,

we used molecular tools originally developed for *Histoplasma capsulatum* and other

dimorphic fungi (6-11) and adapted them to the genetic modification of *E. africanus*.

In some fungi, notably the dimorphic fungal pathogens, homologous recombination is

inefficient, which prevents gene replacement strategies. In these cases,

Agrobacterium-mediated gene transfer (ATMT) has been employed to introduce genes

of interest (9, 11) or fluorescent transcriptional reporters or to create random insertion

libraries for phenotypic screening. Co-culture of *A. tumefaciens,* harbouring the shuttle

plasmid, with the fungus results in the insertion of a single copy (in the majority of

cases) of the T-DNA randomly in the genome (12). This method has been used

successfully for several of the thermally dimorphic pathogens, including *Histoplasma*

capsulatum, Blastomyces dermatitidis, Paracoccidioides brasiliensis and *Talaromyces*

marneffei (as reviewed in (13))*.* We used ATMT to introduce a GFP transgene into *E.*

africanus, using shuttle vectors and a protocol previously developed for the

transformation of *H. capsulatum* (14)*.* The *E. africanus* GFP reporter strain facilitated

imaging of yeast-macrophage interactions and identification of phagocyte cell types in

the lung associated with the yeast in the mouse model of infection (15) by flow

cytometry analysis.

Shuttle vectors which replicate in *E. coli* can be maintained by *H. capsulatum* as linear

plasmids through the addition of telomere sequences (16-18). These vectors are

useful for introducing expression constructs without genome modification. For

example, RNA interference (RNAi) constructs can be used to knock down expression

of virulence-associated genes (7, 18-22). These plasmids are rapidly lost without

selection pressure, which is useful when transient expression is desired, e.g.

expression of the genome-modifying enzyme Cas9. Gene deletion/mutation is a

powerful tool to study putative virulence-associated traits. CRISPR-Cas9 genome

editing has been applied to study these traits in thermally pathogenic fungi, including

Blastomyces dermatitidis (23) and *Histoplasma capsulatum (6, 24).* In this study, we

applied an episomal plasmid system designed for *H. capsulatum* for CRISPR-Cas9

genome editing (6) in *E. africanus* to delete distinct genes in the genome.

With effective molecular genetic tools in place, and their application optimised for *E.*

africanus, its virulence traits and pathogenesis can be investigated in future work. This

knowledge could facilitate the development of more specific diagnostic tests for early

screening of at-risk patients or inform clinical management of the disease.

Results and Discussion

Generation of fluorescent strains using *Agrobacterium*-mediated transformation of a GFP transgene

We used *Agrobacterium tumefaciens*-mediated transformation (ATMT) to transform *E.*

africanus using the shuttle vector pAG22, which encodes a GFP transgene. GFP and

hygromycin phosphotransferase are expressed under the control of *H. capsulatum*

regulatory elements: *TEF1* (translation elongation factor) constitutive promoter and

TUB2 (tubulin) terminator, and *RPL1B* (large ribosomal subunit protein) promoter and

RPL7 (large ribosomal subunit protein) terminator, respectively. Transformation of *E.*

africanus using *A. tumefaciens* LBA1100 or EHA105 strains carrying pAG22 yielded

similar numbers of transformants. Following transformation, individual transformants

were screened for GFP expression levels, as insertion of the T-DNA in different loci

can result in varying levels of GFP expression (12). To identify reporter strains with

high GFP expression, transformant colonies were grown in liquid BHI broth and

inspected by fluorescence microscopy. Figure 1 shows an example of differences in

GFP fluorescence levels between two transformants.

One pAG22 transformant with high fluorescence was selected for further applications.

In preliminary experiments with the mouse model of infection, it was noted that the

virulence of laboratory-passaged *E. africanus* (including untransformed wild-type

strains) was reduced, as higher doses of yeast were required to replicate pathology

compared to previous work. Therefore, all wild-type and control strains used in this

study were passaged under the same conditions in the laboratory as would be required

to generate respective transformants. An intranasal dose of 1×10^7 yeast cells was

then used to infect mice, as this could replicate the pathology seen in previous work

(15). With these conditions, mice infected with the GFP fluorescent strain did not show

a difference in mortality compared to the passage-matched wild-type strain in survival

studies.

Fluorescent E. africanus yeast interaction with macrophages

The GFP-fluorescent yeast facilitated imaging of *E. africanus* interaction with

macrophages. J774A.1 macrophages in chambered imaging slides were infected with

E. africanus yeast (MOI=1) and imaged every 20 minutes. Un-opsonised yeast cells

are readily phagocytosed, and a high proportion of yeast cells were intracellular at the

start of imaging, one hour following yeast addition. We hypothesise that *E. africanus*

may be taken up by macrophages by the same mechanism as *H. capsulatum*, namely,

recognition of heat shock protein 60 (Hsp60) on the surface of yeast cells by

complement receptor 3 (CD11b/CD18) of macrophages (25), as Hsp60 is highly

conserved between the two species. Intracellular yeast could easily be observed by

its GFP fluorescence (Figure 2A). Budding of intracellular yeast could be measured in

the time-lapse data, demonstrating phagocytosed *E. africanus* yeasts remained

viable. The observed budding time of intracellular yeast was not significantly different

to that of extracellular yeast in the same proximity (Figure 2B). Yeast cells continued

to proliferate intracellularly over time, with some macrophages appearing heavily

infected and unable to control yeast proliferation. These data indicate that the

intracellular niche is permissive to *E. africanus* replication, as it is for *H. capsulatum.*

Previous work showed that bone marrow-derived macrophages are not actively lysed

by *E. africanus*, unlike *H. capsulatum*, which may be due to the non-lytic activity of the

E. africanus Cbp1 homologue (26). These data suggest that, similar to *H. capsulatum*

and *P. brasiliensis,* non-activated phagocytes are unable to control *E. africanus,* which

may aid in fungal systemic dissemination during infection (27, 28). Further studies

using the GFP reporter strain and live cell imaging could provide more information on

the dynamics of intracellular replication and macrophage cell death.

Identification of infected phagocyte populations in the mouse lung following

infection with the GFP reporter strain

We applied the GFP fluorescent *E. africanus* strain to examine its association with

phagocytes *in vivo.* Mice were infected intranasally with the fluorescent *E. africanus*

strain, and after 7 days, myeloid cell recruitment in the lungs was analysed by flow

cytometry. Lungs from uninfected mice were analysed as a control. Gating first on

GFP+ cells (i.e., cells with internalised E. africanus) showed that <1% of GFP+ cells

were CD45-negative (Figure 3A), suggesting that nearly all phagocytosed *E. africanus*

yeasts resided in CD45+ phagocytes. Similar results were reported in a study with an

H. capsulatum GFP-expressing strain (29), suggesting that both pathogens persist

well within this intracellular niche *in vivo*. However, these results only reflect the

intracellular fungal burden at the day 7 timepoint, and do not account for fungi that

may have been cleared by this time. Most GFP-fluorescent yeasts were associated

with neutrophils (CD11b^{hi}, Ly6G^{hi}) (66.4±17.4% of GFP+ cells; Figure 3B). The

remainder of cell types that contained GFP yeast were interstitial macrophages

(10.3±5.3%), dendritic cells (MHCII^{hi}, CD11c+) (2.1±0.7%) and a very small subset of

CD11b-, CD11c- cells (Figure 3B).

Alternatively, gating first on all live host cells showed what proportion of total cells were

infected with *E. africanus* (GFP+). Of CD45+ cells within the lung, approximately

16.4±4% were infected (GFP+). We then characterised the GFP-fluorescent (i.e.,

infected) proportion of each myeloid cell type, using a similar gating strategy. The

mean proportion of DCs that were infected (GFP-positive) was 57.3±4.3% of DCs

detected (MHCII^{hi}, CD11c^{hi}), with smaller proportions of total lung macrophages and

neutrophils infected (Figure 3C). Similar proportions were reported for *H. capsulatum*

with the distribution of GFP-fluorescent *H. capsulatum*-infected phagocytes at day 7

post-infection: macrophages: 25.1%, PMNs: 51.2% and DCs: 24.3% for the G217B

strain (29). DCs are pivotal in the adaptive immune response to fungi, including *H.*

capsulatum (30, 31). The high proportion of DCs infected with *E. africanus* at the 7-

day timepoint suggests the initiation of adaptive immunity. The cellular recruitment (of

CD45+ cells) to the lung in response to the GFP-fluorescent strain was not significantly

different to CD45+ cell populations in animals infected with the non-fluorescent wild-

type control (Figure 3D), indicating the GFP-fluorescent E. africanus strain stimulates

a similar host immune response. Previous work examined myeloid cell recruitment in

E. africanus-infected mouse lung at 14 days post-infection, although a lower dose of

yeast was used for infection (15). A time course of infection was not performed in the

current study; this could be included in future experiments to gain more information on

cell recruitment and infected phagocyte proportions over time.

SiglecF+ alveolar macrophages were detected in lungs from uninfected mice (1.5-4%

of total cells), but the proportion was reduced in infected mice (0.01-0.16% of total

cells). Similarly, the number of interstitial macrophages and dendritic cells was higher

in uninfected mice (Figure 3D). This could be due to the proportional increase in

neutrophils recruited or apoptosis of leukocytes in the lung, a known feature of

infection with *H. capsulatum* and a critical element of protective immunity (32).

Transformation of *E. africanus* with episomal vectors and reduction of cell wall

α(1,3)-glucan by RNA interference

E. africanus could be transformed with episomal plasmids based on pCR115 (18) for

which linear plasmids with telomeres are maintained without chromosome integration.

This was first tested by constructing a modified plasmid with hygromycin B selection

(pHygR-GFP). The number of colonies obtained using electroporation of episomal

plasmids was typically lower than for ATMT. Hygromycin selection pressure was

maintained when working with these transformants, as these plasmids are rapidly lost

from the cell population without selection.

To silence production of α(1,3)-glucan in *E. africanus*, we expressed an AGS1-

targeting RNA hairpin from an episomal plasmid based on the AGS1-RNAi plasmid

pCR115. The RNA hairpin is designed for RNAi-induced knockdown of the *H.*

capsulatum AGS1 gene, with homology to a 678-base pair region of exon 3 of this

gene (18). This region is 84% similar overall in *H. capsulatum* and *E. africanus,* with

several continuous sections of at least 20 nucleotides or with 100% identity. Therefore,

we hypothesised that effective RNAi silencing using the same RNA hairpin could be

achieved in *E. africanus,* given that several 21-22 small interfering RNA molecules

with complete sequence match are produced following Dicer processing of the long

double-stranded RNA. We replaced the GFP in pHygR-GFP with the *AGS1* inverted

repeat sequence from pCR115 and used this plasmid (pHygR-HcAGS1-RNAi) to

transform *E. africanus.* Yeast transformed with pHygR-HcAGS1-RNAi did not differ in

growth rate compared to control yeast, which was transformed with a similar plasmid

containing a GFP inverted repeat.

Reduction of cell wall α(1,3)-glucan through RNA interference was assessed by

staining the AGS1-RNAi plasmid-transformed yeasts with a monoclonal antibody

recognising α(1,3)-glucan. It was noted that not all yeast cells were uniformly stained

(Figure 4A), suggesting variable amounts of α(1,3)-glucan. *H. capsulatum* also

displays variation in cell wall α(1,3)-glucan levels, as assessed by staining with the

same antibody, during broth culture growth (33). Similarly, *E. africanus* cell wall α(1,3)-

glucan levels may increase at different time points during culture, or in different growth

media. Nevertheless, yeasts transformed with the *AGS1* RNAi plasmid showed an

overall lower level of cell wall α(1,3)-glucan staining in the population compared to the

control (Figure 4A and 4B). RNAi-induced gene silencing does not typically lead to a

complete knockdown of the target, and due to variability in dsRNA expression, a small

proportion of cells do not exhibit the full RNAi effect (18). Therefore, some α(1,3)-

glucan staining was still observed in *E. africanus* yeasts maintaining the RNAi plasmid.

Gene mutation in *E. africanus* using an episomal CRISPR/Cas9 system

To test whether CRISPR-Cas9 gene editing could facilitate gene mutation in E.

africanus, we transformed a GFP-expressing strain of *E. africanus* with a linear

plasmid expressing Cas9 and a GFP-targeting CRISPR gRNA. The GFP-fluorescent

E. africanus strain was generated through ATMT using pAG21 (GFP-expression and

G418 selection). The CRISPR-Cas9 plasmid provided for hygromycin resistance and

expressed a gRNA targeting the GFP gene. Hygromycin-resistant transformant

colonies were passaged in BHI broth with hygromycin B selection, and the yeasts were

examined by fluorescence microscopy for loss of the GFP fluorescence. After

successive passages, an increasing proportion of the population was no longer GFP

fluorescent (Figure 5A). Sequencing of the GFP gene in the transformant population

confirmed a mixed population with indels at the expected Cas9 cut site (Figure 5B),

showing efficient CRISPR/Cas9-based genome editing in *E. africanus*.

Auxotroph strains are useful for genetic manipulations because they facilitate the

selection of plasmids carrying genes that rescue the auxotrophy. A previous

CRISPR/Cas9 workflow used for *H. capsulatum* (6) was used to disrupt *URA5* in *E.*

africanus to generate a uracil auxotroph strain. The *E. africanus URA5* homologue

(Gene: ACJ72_02173) was identified through a BLAST search. The protein sequence

is highly similar to that of *H. capsulatum* (88.1% identity, 93.0% similarity as

determined by pairwise alignment using EMBOSS Water (34)). A guide RNA sequence

was designed using CRISPOR (https://crispor.gi.ucsc.edu/) (35) to maximise the

chance of indel generation in the *URA5* gene while minimising off-target effects.

Following transformation of wild-type *E. africanus* with the URA5-targeting CRISPR-

Cas9 plasmid, followed by four passages under hygromycin selection, uracil

auxotrophs were selected using 5-FOA-resistance screening. Several 5-FOA-resistant

clones were identified, and their inability to grow without uracil was validated by failure

to grow in media lacking uracil. Sequencing of four clones showed a single nucleotide

T insertion in all clones (Figure 5C), causing a frameshift mutation, preventing

production of a functional *URA5* gene product. This *E. africanus ura5*-mutant can now

be used as the host for *URA5*-based expression vectors or RNAi vectors. These

vectors are also stable *in vivo,* as evidenced by continued virulence*; H. capsulatum*

uracil auxotrophs are avirulent in mice, which is likely to be the case for *E. africanus*

as well (36).

Conclusions

In this study, we demonstrate that molecular tools developed for genetic modification

of *H. capsulatum* could be adapted for use in *E. africanus,* suggesting that the two

species use similar regulatory elements for high-level gene expression.   Our results

include the use of ATMT to transform *E. africanus* with expression plasmids,

generating a GFP-fluorescent strain that can be used for tracking infections both in

cultured macrophages and *in vivo*. In addition, we provide results showing the efficacy

of episomal plasmids designed to induce RNA-interference-based gene knockdown

and CRISPR-Cas9-based mutation of target genes. Thus, by adapting technologies,

we have put into place tools and methodologies to functionally study genes in E.

africanus through expression as well as loss-of-function.   Furthermore, we have

created an *E. africanus* uracil auxotroph strain enabling the use of URA*5*-based

complementation vectors instead of hygromycin selection. Such URA5-based vectors

have the advantage of providing for selection *in vivo*, which cannot be done with

hygromycin. Together, these tools make *E. africanus* amenable to genetic modification

to investigate virulence traits and how they influence the host-pathogen interaction.

Materials and Methods

E. africanus and *A. tumefaciens* strains and growth conditions

Emergomyces africanus (clinical isolate CBS 136260 (4)) was grown in brain-heart

infusion broth (BHI) (Cat No. 110493, Merck, USA), at 37 °C, 180 rpm, or maintained

on BHI agar plates (Cat. No. 70138, Merck) at 37 °C. *A. tumefaciens* strains LBA1100

and EHA105 were maintained on LB agar plates at 25 °C. Transformation of

Agrobacterium binary vectors into *A. tumefaciens* was done by electroporation as

described in (37) and transformants selected with 100 µg/mL kanamycin.

Transformation using *Agrobacterium tumefaciens*

Transformation of *E. africanus* yeast by *Agrobacterium* was done using previously

established methods for *H. capsulatum* (14), with the exception that BHI broth or BHI

agar plates were used instead of *Histoplasma*-specific growth medium. *Agrobacterium*

tumefaciens strains LBA1100 and EHA105 were used to transform *E. africanus* with

pAG22 (GFP-expression, hygromycin resistance), yielding similar results. BHI was

inoculated with a colony of *E. africanus* from a fresh streak plate, and the culture was

grown for three days at 37 °C with shaking. Colonies from a streak plate of *A.*

tumefaciens were used to inoculate 5 ml minimal glucose medium (14) for 24 h at 25

°C, then 1 ml of this culture was used to inoculate 5 ml induction medium containing

0.1 mM acetosyringone (Sigma-Aldrich) for a further 16 h of growth. *E. africanus* yeast

cells were washed with PBS by centrifugation and resuspension, then resuspended in

induction medium. The optical density (600 nm) of both *E. africanus* and *A.*

tumefaciens suspensions was determined, then combined in 900 µl to give final optical

densities of 1.0 for each. Suspensions OD600=1.0 have approximately 5×10^8 cells/ml

and 2.5×10^8 cells/ml of *A. tumefaciens* and *E. africanus,* respectively, giving an MOI

of 2:1. The co-culture was applied dropwise to sterile Whatman No. 5 qualitative filter

papers placed on co-cultivation plates. The co-cultivation plates were incubated at 25

°C for 2 days, then transferred to BHI agar plates containing 300 µg/ml hygromycin B

and 200 µM cefotaxime. The plates were incubated at 37 °C for 10-14 days until *E.*

africanus transformant colonies appeared. Single colonies were picked and streaked

on fresh BHI plates containing 300 µg/ml hygromycin B. From these plates, single

colonies were used to inoculate BHI broth to grow liquid cultures of individual

transformants. Glycerol stocks (20%) of liquid cultures were preserved at -80 °C.

Infection of macrophages with *E. africanus* and live-cell imaging

J774A.1 murine macrophage cells (ATCC: TIB-7) were seeded at a density of 2×10^5

cells in 0.3 ml macrophage medium (DMEM high glucose GlutaMAX®, 10% FBS,

penicillin/streptomycin [Gibco]) in 8-well imaging slides (ibidi GMBH, Cat. No. 80826)

24 h prior to the addition of *E. africanus* yeast. *E. africanus* yeast was grown for 3 days

in BHI broth, then washed with PBS and counted using a haemocytometer. 2×10^5

yeast cells in 300 µl macrophage media were added to macrophages in imaging slides

to achieve a ratio of one yeast cell per macrophage (MOI=1). The slides were then set

up for time-lapse imaging under the conditions of 37 °C, 5% CO_2. Imaging was started

1 hour post-yeast addition. Imaging was done using a Zeiss AxioObserver

fluorescence microscope at 40X magnification. Analysis of observed budding time was

done by examining yeast cells that were intracellular or extracellular, at t=0, with no

visible buds.

Mouse model of infection and flow cytometry

To examine the responses to *E. africanus* in the mouse lung, three groups of mice

were analysed (n=6 per group): (1) uninfected mice, (2) mice infected with the wild-

type clinical isolate and (3) mice infected with the GFP fluorescent *E. africanus* strain.

Eight- to twelve-week-old male CL57/B6 mice were obtained from the University of

Cape Town Animal Research Specific Pathogen Free Facility. *E. africanus* wild-type

and GFP fluorescent strains were grown in BHI broth for 3 days at 37 °C until OD600

reached approximately 1.0. Yeast cells were washed twice with PBS before counting

using a haemocytometer. Mice were anaesthetised with ketamine and xylazine (80

mg/kg and 16 mg/kg, respectively), then infected intranasally with 50 µl of yeast cell

suspension containing 1×10^7 yeast cells. Animal welfare was monitored daily. After 7

days, mice were euthanised by halothane inhalation and the lungs were removed and

stored in DMEM + 5% FBS on ice. Lungs were digested in digestion buffer (15)

containing 50 units/ml Collagenase Type I and 13 µg/ml DNAse I for 1h at 37 °C. The

digested lungs were passed through a 70 µm cell strainer using a sterile syringe

plunger. This process was repeated using a 40 µm cell strainer. Red blood cells were

lysed using 1x RBC lysis buffer (eBioscience), then remaining cells collected by

centrifugation and resuspension in DMEM + 5% FBS, then counted using a

haemocytometer. 2×10^6 cells per well were added to a V-bottom 96-well plate.

Myeloid cells were stained in MACS buffer (0.5% BSA, 2 mM EDTA in PBS) using the

antibodies in Table 3, together with 1% Fc block and 2% heat-inactivated rat serum,

for 20 min at 4 °C in the dark, then washed twice with MACS buffer. 7-AAD (Biolegend)

was used as the cell viability stain. Single-stained and unstained controls were used

to compensate for spectral overlap. Stained cells were analysed by flow cytometry

using an LSR Fortessa™ (BD Immunocytometry Systems, San Jose, CA, USA) and

BD FACS Diva software (v6.0). Flowjo software (v10.0.7) (Treestar, Ashland, OR,

USA) was used for post-acquisition analysis and cell population determination. The

gating strategy used is shown in Figure 3A.

Plasmid construction

Plasmids pAG21, pAG22, pCR115 (18), pSL01 and pCR745 (6) were provided by

CAR. Isolation of plasmid DNA, restriction enzyme cloning and PCR/gel purification

were done using standard methods, and plasmids were maintained in *E. coli* DH5α.

Restriction enzymes and ligation kits were purchased from New England Biolabs. An

episomal GFP expression plasmid (pHygR-GFP) was constructed by digestion of

pAG22 with EcoRI and HindIII, and ligating the fragment containing the GFP- and *hph*

expression cassettes into the similarly-digested pCR115 vector backbone. To

construct pHyg-HcAGS1-RNAi (AGS1 inverted repeat) used for *AGS1* RNAi, the *GFP*

sequence from plasmid pHygR-GFP was exchanged with the *AGS1* inverted repeat

sequence from pCR115 by digestion with AscI and SpeI. A control *GFP* RNAi plasmid

(pHygR-GFP-RNAi) was constructed by inserting an inverted repeat of *GFP* into

pHygR-GFP by PCR with GFP-SpeI-F and XhoI-GFP-R primers and restriction

enzyme cloning. For CRISPR/Cas9-mediated disruption of GFP, plasmid pCR115-

sgGFP was created: a portion of plasmid pCR745 (6) containing the guide RNA

targeting GFP was digested with SwaI and AvrII and ligated into similarly digested

pSL01 (6), resulting in plasmid pSL01-sgGFP. For CRISPR/Cas9-mediated disruption

of *URA5,* a synthetic DNA sequence containing the guide RNA sequence,

hammerhead and HDV ribozyme sequences, and part of the tracr RNA sequence,

including 15 bp up- and downstream-homology regions, was designed for cloning into

pSL01 as described in (6). The guide RNA sequence was designed using CRISPOR

(https://crispor.gi.ucsc.edu/) (35) to maximise the chance of indels while preventing

off-target effects. This sequence was synthesised by Integrated DNA Technologies

(Leuven, Belgium). The synthetic DNA sequence was cloned into SwaI-digested

pSL01 and transformed into competent *E. coli* cells using In-Fusion® HD Cloning Kit

(Takara Bio) using the manufacturer's recommendations.

414   Transformation of yeast with episomal plasmids by electroporation

Transformation of *E. africanus* with plasmids by electroporation of was based on

previously established methods for *H. capsulatum* (17). Briefly, *E. africanus* yeasts

were grown in 5 ml BHI broth for 3 days until OD600 reached approximately 1.0. Yeast

cells were collected by centrifugation and resuspended in 2 ml 10% mannitol.

Plasmids were linearised by digestion with PmeI to expose the telomeres, then

column-purified and eluted with nuclease-free water. Three hundred microlitres of *E.*

africanus yeast suspension (1×10^7 yeast cells) was mixed with 200 ng purified

plasmid in a chilled electroporation cuvette (0.2 cm gap), then electroporated in a

single pulse using a Bio-Rad Gene Pulser® II system with the following settings: 1.25

kV, 50 mF, 600 Ω, which gave a time constant of 8-12 ms. Following electroporation,

500 µl of BHI broth was mixed with cells and this cell suspension was applied dropwise

to sterile Whatman No. 5 qualitative filter papers placed on BHI agar plates with no

selection. These plates were incubated for 48 h at 37 °C to allow for recovery of

transformants, before transferring the membranes to BHI agar plates containing 300

µg/ml hygromycin B. The plates were incubated at 37 °C for 10-14 days until

transformant colonies appeared. Single colonies were picked and streaked onto fresh

BHI plates (300 µg/ml hygromycin B). Selection with hygromycin B (300 µg/ml) was

maintained for growing liquid cultures, after which -80 °C glycerol stocks were

prepared for long-term storage of strains, as described above.

Cell wall alpha(1,3)-glucan staining of RNAi strains

Yeasts harbouring the *AGS1* RNAi plasmid pHygR-HcAGS1-RNAi or the control

plasmid pHygR-GFP-RNAi were grown for 3 days in BHI broth with 300 µg/ml

hygromycin B until cell density reached an OD600 of 1.0. Cells from 300 µl of culture

were collected by centrifugation, then washed twice with PBS, and fixed in 4%

paraformaldehyde overnight at 4 °C. Cells were washed twice with PBS, then blocked

with 1% BSA, PBS for 1 h. Cells were then stained with 1:150 mouse IgM MOPC-

104E antibody (M5170, Sigma-Aldrich) in 1% BSA, PBS for 2 h at ambient

temperature. The cells were washed twice by centrifugation and resuspended in PBS

and stained with 1:400 anti-mouse IgM secondary antibody conjugated to

AlexaFluor488 (Invitrogen, A-21042) for 1 h at ambient temperature. To stain total

yeast cells, calcofluor white was added at a final concentration of 100 μg/ml and

incubated for 15 min at ambient temperature in the dark. Cells were washed twice with

PBS and resuspended in 20 μl PBS. A 5μl sample of stained cells was spotted onto a

microscope slide, allowed to air-dry and mounted with ProLong™ Glass Antifade

Mountant (P36984, Invitrogen). The slides were examined using a Zeiss Axiovert

200M fluorescence microscope with a monochrome Zeiss AxioCam HRm camera at

100x magnification. Exposure times were kept constant for all samples when capturing

images for mean fluorescence analysis. Fluorescence of the population was

determined by measuring the background-corrected mean grey values for 10 fields of

view each using ImageJ, for two pooled independent experiments, then dividing this

value by the number of total cells (as determined by calcofluor white staining).

Targeted mutagenesis using CRISPR/Cas9-plasmids

A GFP-expressing strain of *E. africanus* was generated through ATMT using pAG21

(equivalent to pAG22 but with a G418 resistance marker instead of hygromycin B

resistance). The GFP-fluorescent strain was subsequently transformed with PmeI-

linearised plasmid pSL01-sgGFP by electroporation. Hygromycin-resistant

transformant colonies were picked and grown in BHI broth for successive passages

(1:50 dilution), maintaining hygromycin selection every 3 days. Samples of these

cultures were examined by fluorescence microscopy for GFP expression. For DNA

sequencing, yeast cells from 1 ml of culture was collected by centrifugation, lysed with

200 μl of lysis buffer (2% Triton X-100, 20 mM EDTA) and beating (2 min) with

approximately 100 μl of glass beads (0.5 μm diameter). 100 μL of phenol and 100 μl

chloroform-isoamyl alcohol (25:24:1) were added, and the phases were separated by

centrifugation. DNA was recovered by transferring 50 µl of the aqueous (upper) phase

into 200 µl distilled water. This crude DNA-containing solution was used as the

template (2% vol/vol) for PCR using Taq polymerase and primers flanking the

expected gene editing sites: GFP-SpeI-F and GFP-Xho-R, or Ura seq-F and Ura seq-

R.

For mutation of *URA5* to create a uracil auxotroph, a guide RNA sequence targeting

URA5 was designed using CRISPOR (https://crispor.gi.ucsc.edu/) (35). A DNA

construct containing the crRNA, hammerhead- and HDV ribozyme sequences, and

part of the tracrRNA sequence, with 15 bp flanking homology arms, was synthesised

as described in (6) for cloning into plasmid pSL01, by Integrated DNA Technologies

(Leuven, Belgium). The sequence of this construct is shown in Table 2. Cloning of this

synthetic DNA into SwaI-digested plasmid pSL01 was done through InFusion Cloning

(Takara Biotec) as per the manufacturer's recommendations. Constructs were verified

through Sanger sequencing. *E. africanus* was transformed with this plasmid by

electroporation as described above, followed by selection with hygromycin B. After

four passages in liquid culture, cultures were streaked onto BHI plates supplemented

with 1mg/ml 5-Fluoroorotic Acid (5-FOA, R0811, ThermoFisher) and 100 µg/ml uracil

to isolate uracil auxotrophs. Uracil auxotrophy was confirmed by the inability to grow

in DMEM/F12 media (11320033, Gibco), which lacks uracil, compared to growth in

parallel in DMEM/F12 media supplemented with 100 µg/ml uracil. Independent clones

were picked for Sanger sequencing.

Ethics statement

All work with mice was done with ethical clearance from the University of Cape Town Animal Research Ethics Committee, reference number 020/017.

Statistical analysis and software

Values are reported as means ± standard deviation. Differences between groups were determined using a two-tailed Student's t-test or one-way ANOVA using GraphPad Prism v10.6.1. Plasmid map generation and alignments of DNA sequence data were done using SnapGene v8.02. ImageJ 1.54p was used for image analysis (mean fluorescence intensities and total cell counting).

Funding statement

This work was supported by the National Research Foundation of South Africa (to JCH), Carnegie Corporation (to JCH and LD), National Institute of Allergy and Infectious Diseases (NIAID) AI148561 to CAR, and Wellcome Trust: 209293/Z/17/Z to JCH and 310933/Z/24/Z to ERB and JCH. The funders played no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interest Statement

The authors have declared no competing interest.

517 Table 1: Plasmids used in the study

Plasmid	Application	Selection
pAG22	ATMT	Hygromycin B (300 µg/ml)
pAG21	ATMT	G418 (1 mg/ml)
pCR115	*AGS1* RNAi	[Uracil auxotrophy]
pHygR-GFP	GFP episomal plasmid	Hygromycin B (300 µg/ml)
pHygR-GFP-RNAi	GFP inverted repeat plasmid (RNAi control)	Hygromycin B (300 µg/ml)
pHygR-HcAGS1-RNAi	*AGS1* RNAi	Hygromycin B (300 µg/ml)
pCR745	CRISPR/Cas9	[Uracil auxotrophy]
pSL01	CRISPR/Cas9	Hygromycin B (300 µg/ml)
pSL01-sgGFP	CRISPR/Cas9	Hygromycin B (300 µg/ml)
pSL01-sgURA	CRISPR/Cas9	Hygromycin B (300 µg/ml)

522 Table 2: Primers and oligonucleotides

GFP-SpeIF	GAGAACTAGTATGAGCAAGGGCGAGGAGC
GFP-XhoIR	GAGACTCGAGCTTGTACAGCTCGTCCATGC
Synthesised sequence for *URA5* CRISPR/Cas9 editing (guide sequence underlined)	CTCTCCCAATTTTCACATTTAAATCCCGCC ACCCGATGTCTGATGAGTCCGTGAGGAC GAAACGAGTAAGCTCGTC<u>ACATCGGGTT TCGGAATGG</u>AGTTTTAGAGCTAGAAATAG CAAGTTAAAATAAGGCTAGTCCGTTAT
Ura seq F	TTGACACAGCCCATAAACG
Ura seq R	ATCGCCTTCTATATTCCTCC

531   Table 3: Flow cytometry antibodies

Target (Clone)	Conjugated fluorophore	Manufacturer (catalogue number)	Dilution
CD45 (30-F11)	BV510	BD Biosciences (567800)	1:100
CD11b (M1/70)	BV421	BD Biosciences (566313)	1:100
CD11c (HL3)	Alexa Fluor 700	BD Biosciences (560583)	1:100
Ly-6G (1A8)	APC	BD Biosciences (560599)	1:100
F4/80 (BM8)	PE/Cy7	Biolegend (123113)	1:100
Siglec-F (E50-2440)	APC/Cy7	BD Biosciences (565527)	1:50
I-A/I-E (MHCII) (M5/114.15.2)	PE/Cy5	Biolegend (107611)	1:100

References

1. Dukik K, Munoz JF, Jiang Y, Feng P, Sigler L, Stielow JB, Freeke J, Jamalian A, Gerrits van den Ende B, McEwen JG, Clay OK, Schwartz IS, Govender NP, Maphanga TG, Cuomo CA, Moreno LF, Kenyon C, Borman AM, de Hoog S. 2017. Novel taxa of thermally dimorphic systemic pathogens in the Ajellomycetaceae (Onygenales). Mycoses 60:296-309.

2. Schwartz IS, Govender NP, Sigler L, Jiang Y, Maphanga TG, Toplis B, Botha A, Dukik K, Hoving JC, Munoz JF, de Hoog S, Cuomo CA, Colebunders R, Kenyon C. 2019. Emergomyces: The global rise of new dimorphic fungal pathogens. PLoS Pathog 15:e1007977.

3. Schwartz IS, Kenyon C, Lehloenya R, Claasens S, Spengane Z, Prozesky H, Burton R, Parker A, Wasserman S, Meintjes G, Mendelson M, Taljaard J, Schneider JW, Beylis N, Maloba B, Govender NP, Colebunders R, Dlamini S. 2017. AIDS-Related Endemic Mycoses in Western Cape, South Africa, and Clinical Mimics: A Cross-Sectional Study of Adults With Advanced HIV and Recent-Onset, Widespread Skin Lesions. Open Forum Infect Dis 4:ofx186.

4. Schwartz IS, McLoud JD, Berman D, Botha A, Lerm B, Colebunders R, Levetin E, Kenyon C. 2018. Molecular detection of airborne Emergomyces africanus, a thermally dimorphic fungal pathogen, in Cape Town, South Africa. PLoS Negl Trop Dis 12:e0006174.

5. Jiang Y, Dukik K, Muñoz JF, Sigler L, Schwartz IS, Govender NP, Kenyon C, Feng P, van den Ende BG, Stielow JB, Stchigel AM, Lu H, de Hoog S. 2018. Phylogeny, ecology and taxonomy of systemic pathogens and their relatives in Ajellomycetaceae (Onygenales): Blastomyces, Emergomyces, Emmonsia, Emmonsiellopsis. Fungal Diversity 90:245-291.

6. Rappleye CA. 2023. Targeted gene deletions in the dimorphic fungal pathogen Histoplasma using an optimized episomal CRISPR/Cas9 system. mSphere 8:e0017823.

7. Brechting PJ, Shah C, Rakotondraibe L, Shen Q, Rappleye CA. 2023. Histoplasma capsulatum requires peroxisomes for multiple virulence functions including siderophore biosynthesis. mBio 14:e0328422.

8. Kujoth GC, Sullivan TD, Klein BS. 2020. Gene Editing in Dimorphic Fungi Using CRISPR/Cas9. Curr Protoc Microbiol 59:e132.

9. Kemski MM, Stevens B, Rappleye CA. 2013. Spectrum of T-DNA integrations for insertional mutagenesis of Histoplasma capsulatum. Fungal Biol 117:41-51.

10. Youseff BH, Rappleye CA. 2012. RNAi-based gene silencing using a GFP sentinel system in Histoplasma capsulatum. Methods Mol Biol 845:151-64.

11. Sullivan TD, Rooney PJ, Klein BS. 2002. Agrobacterium tumefaciens integrates transfer DNA into single chromosomal sites of dimorphic fungi and yields homokaryotic progeny from multinucleate yeast. Eukaryot Cell 1:895-905.

12. Idnurm A, Bailey AM, Cairns TC, Elliott CE, Foster GD, Ianiri G, Jeon J. 2017. A silver bullet in a golden age of functional genomics: the impact of Agrobacterium-mediated transformation of fungi. Fungal Biol Biotechnol 4:6.

13.   Sil A, Andrianopoulos A. 2014. Thermally Dimorphic Human Fungal
Pathogens--Polyphyletic Pathogens with a Convergent Pathogenicity Trait.
Cold Spring Harb Perspect Med 5:a019794.
14.   Zemska O, Rappleye CA. 2012. Agrobacterium-Mediated Insertional
Mutagenesis in Histoplasma capsulatum, p 51-66, Host-Fungus Interactions
doi:10.1007/978-1-61779-539-8_4. Humana Press.
15.   Hoft MA, Duvenage L, Salie S, Keeton R, Botha A, Schwartz IS, Govender
NP, Brown GD, Hoving JC. 2024. The pathogenesis of experimental
Emergomycosis in mice. PLoS Negl Trop Dis 18:e0011850.
16.   Woods JP, Goldman WE. 1993. Autonomous replication of foreign DNA in
Histoplasma capsulatum: role of native telomeric sequences. J Bacteriol
175:636-41.
17.   Woods JP, Heinecke EL, Goldman WE. 1998. Electrotransformation and
expression of bacterial genes encoding hygromycin phosphotransferase and
beta-galactosidase in the pathogenic fungus Histoplasma capsulatum. Infect
Immun 66:1697-707.
18.   Rappleye CA, Engle JT, Goldman WE. 2004. RNA interference in Histoplasma
capsulatum demonstrates a role for alpha-(1,3)-glucan in virulence. Mol
Microbiol 53:153-65.
19.   Ray SC, Rappleye CA. 2022. Mac1-Dependent Copper Sensing Promotes
Histoplasma Adaptation to the Phagosome during Adaptive Immunity. mBio
13:e0377321.
20.   Shen Q, Ray SC, Evans HM, Deepe GS, Jr., Rappleye CA. 2020. Metabolism
of Gluconeogenic Substrates by an Intracellular Fungal Pathogen
Circumvents Nutritional Limitations within Macrophages. mBio 11.
21.   Dade J, DuBois JC, Pasula R, Donnell AM, Caruso JA, Smulian AG, Deepe
GS, Jr. 2016. HcZrt2, a zinc responsive gene, is indispensable for the survival
of Histoplasma capsulatum in vivo. Med Mycol 54:865-75.
22.   Hilty J, George Smulian A, Newman SL. 2011. Histoplasma capsulatum
utilizes siderophores for intracellular iron acquisition in macrophages. Med
Mycol 49:633-42.
23.   Kujoth GC, Sullivan TD, Merkhofer R, Lee TJ, Wang H, Brandhorst T,
Wuthrich M, Klein BS. 2018. CRISPR/Cas9-Mediated Gene Disruption
Reveals the Importance of Zinc Metabolism for Fitness of the Dimorphic
Fungal Pathogen Blastomyces dermatitidis. mBio 9.
24.   Joehnk B, Ali N, Voorhies M, Walcott K, Sil A. 2023. Recyclable
CRISPR/Cas9-mediated gene disruption and deletions in Histoplasma.
mSphere 8:e0037023.
25.   Long KH, Gomez FJ, Morris RE, Newman SL. 2003. Identification of heat
shock protein 60 as the ligand on Histoplasma capsulatum that mediates
binding to CD18 receptors on human macrophages. J Immunol 170:487-94.
26.   Azimova D, Herrera N, Duvenage L, Voorhies M, Rodriguez RA, English BC,
Hoving JC, Rosenberg O, Sil A. 2022. Cbp1, a fungal virulence factor under
positive selection, forms an effector complex that drives macrophage lysis.
PLoS Pathog 18:e1010417.
27.   Allendoerfer R, Deepe GS, Jr. 1997. Intrapulmonary response to Histoplasma
capsulatum in gamma interferon knockout mice. Infect Immun 65:2564-9.
28.   Calich VL, da Costa TA, Felonato M, Arruda C, Bernardino S, Loures FV,
Ribeiro LR, de Cássia Valente-Ferreira R, Pina A. 2008. Innate immunity to
Paracoccidioides brasiliensis infection. Mycopathologia 165:223-36.

29. Deepe GS, Jr., Gibbons RS, Smulian AG. 2008. Histoplasma capsulatum
manifests preferential invasion of phagocytic subpopulations in murine lungs.
J Leukoc Biol 84:669-78.
30. Van Prooyen N, Henderson CA, Hocking Murray D, Sil A. 2016. CD103+
Conventional Dendritic Cells Are Critical for TLR7/9-Dependent Host Defense
against Histoplasma capsulatum, an Endemic Fungal Pathogen of Humans.
PLOS Pathogens 12:e1005749.
31. Szymczak WA, Deepe GS, Jr. 2010. Antigen-presenting dendritic cells rescue
CD4-depleted CCR2-/- mice from lethal Histoplasma capsulatum infection.
Infect Immun 78:2125-37.
32. Allen HL, Deepe GS, Jr. 2005. Apoptosis modulates protective immunity to the
pathogenic fungus Histoplasma capsulatum. J Clin Invest 115:2875-85.
33. Kugler S, Schurtz Sebghati T, Groppe Eissenberg L, Goldman WE. 2000.
Phenotypic variation and intracellular parasitism by Histoplasma capsulatum.
Proc Natl Acad Sci U S A 97:8794-8.
34. Rice P, Longden I, Bleasby A. 2000. EMBOSS: the European Molecular
Biology Open Software Suite. Trends Genet 16:276-7.
35. Concordet JP, Haeussler M. 2018. CRISPOR: intuitive guide selection for
CRISPR/Cas9 genome editing experiments and screens. Nucleic Acids Res
46:W242-W245.
36. Retallack DM, Heinecke EL, Gibbons R, Deepe GS, Jr., Woods JP. 1999. The
URA5 gene is necessary for histoplasma capsulatum growth during infection
of mouse and human cells. Infect Immun 67:624-9.
37. den Dulk-Ras A, Hooykaas PJ. 1995. Electroporation of Agrobacterium
tumefaciens. Methods Mol Biol 55:63-72.

Transformant 1   Transformant 2

Figure 1: GFP-expressing E. africanus derived through ATMT. E. africanus GFP-expressing transformants were derived by transformation with A. tumefaciens harbouring pAG22. Individual transformant colonies were evaluated by fluorescence microscopy, showing variable expression levels of GFP between transformants under constant imaging conditions. Bar=10μm

Figure 2: An *E. africanus* GFP fluorescent reporter strain facilitates the study of intracellular yeast growth
GFP fluorescent reporter yeast was added to J774.1 macrophage cells in chambered imaging slides. Time-lapse imaging was started 1 hour after yeast addition (t=0), at which point several yeast cells had been internalised (a representative example is shown in A, left). The emergence of new buds (A, right, indicated by arrows) was recorded. Observed budding time was summarised for several fields of view that contained a mix of both extracellular and intracellular yeasts, P>0.05 (mean ± SD). Bar=10µm.

D

Neutrophils

Interstitial macrophages

Dendritic cells

Figure 3: Identification of myeloid cell types associated with GFP-expressing E. africanus in mouse lung at 7 days post-infection. Scatter plots showing the strategy of gating on GFP first (A), which was used to identify myeloid cell types in the mouse lung that were GFP-positive (associated with the GFP reporter yeast), as summarised in (B). (C) Alternate gating on the various cell types first determined the proportions of each cell type associated with GFP yeast. (D) Myeloid cell distribution in the WT-and and GFP strain-infected mouse lung and in uninfected mouse lung. Differences in cell recruitment between WT- and GFP-strain-infected mice were not significant, P>0.05 (mean ± SD). n=6 mice per group, data is representative of two independent experiments.

Figure 4: (A) Cell wall staining and fluorescence microscopy of wild-type and AGS1 RNAi strains of E. africanus. E. africanus harbouring the AGS1 RNAi plasmid or the control GFP RNAi plasmid were stained with a monoclonal antibody specific for $\alpha(1,3)$-glucan and calcofluor white (CFW) and examined by fluorescence microscopy. Imaging settings were kept constant for all samples. Bar=10µm (B) Background-corrected mean fluorescence intensity of microscopy images was measured and normalised by the number of cells per image. Ten images each from two independent experiments were analysed, Student's unpaired t-test, $p < 0.001$.

A

parent

passage=2

passage=4

B

Sequencing: GFP-F

expected Cas9 cut site

CCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCGGCAARS SKGCCC GGGCCC TGCCCCCCCCTCGKGCCCCCCC TGC

C

guide sequence

PAM

CCGAAGATTACATCGGGTTTCGGAATTGGAAGGATTCGAGGA

Consensus   CCGAAGATTACATCGGGTTTCGGAATTGGAAGGATTCGAGGA

H08_WT_ura_R___1.seq   CCGAAGATTACATCGGGTTTCGGAA-TGGAAGGATTCGAGGA
H07_P1a_ura_R___1.seq   CCGAAGATTACATCGGGTTTCGGAATTGGAAGGATTCGAGGA
B08_P1b_ura_R___1.seq   CCGAAGATTACATCGGGTTTCGGAATTGGAAGGATTCGAGGA
D08_P4a_ura_R___1.seq   CCGAAGATTACATCGGGTTTCGGAATTGGAAGGATTCGAGGA
F08_P4b_ura_R___1.seq   CCGAAGATTACATCGGGTTTCGGAATTGGAAGGATTCGAGGA

T insertion

Figure 5: Evaluation of a CRISPR/Cas9 episomal system in E. africanus. (A) Merged fluorescence microscopy and brightfield images showing a decline in the proportion of fluorescent yeast cells with subsequent passages following transformation of GFP-expressing E. africanus with the CRISPR/Cas9 plasmid pSL01-sgGFP. Bar=10μm. (B) Sanger sequencing of a mixed population showed a mixture of signals following the expected Cas9 cut site, indicating the presence of various indels in the population. (C) Sequencing data of individually selected clones, aligned with the WT strain, following CRISPR/Cas9-mediated gene disruption of URA5. A T-insertion was observed in all clones.