Paternal over- and under-nutrition program fetal and placental development in a sex-specific manner in mice

Short title: Poor paternal diet and sex-specific placental programming

Hannah L. Morgan[1, 2], Nader Eid[1], Nadine Holmes[3], Matthew Carlile[3] Sonal Henson[3], Fei Sang[3], Victoria Wright[3,] Marcos Castellanos-Uribe[4], Iqbal Khan[4], Nazia Nazar[4], Sean T. May[4], Rod T. Mitchell[5], Federica Lopes[6], Robert S. Robinson[7], A. Augusto Coppi[8], Vipul Batra[9] and Adam J. Watkins[9]

[1]Lifespan and Population Health, School of Medicine, University of Nottingham, Nottingham, NG7 2UH, United Kingdom.

[2]Maternal and Fetal Health Research Centre, Faculty of Biology Medicine and Health, University of Manchester, St Mary's Hospital, Oxford Road, Manchester, M13 9WL, United Kingdom.

[3]Deep Seq, School of Life Sciences, Queen's Medical Centre, University of Nottingham, Nottingham, NG7 2UH, United Kingdom.

[4] Nottingham Arabidopsis Stock Centre (NASC), University of Nottingham, Sutton Bonington Campus, Plant Science Building, School of Biosciences, Loughborough, LE12 5RD, United Kingdom.

[5]Centre for Reproductive Health, Institute for Regeneration and Repair, University of

Edinburgh, Edinburgh, EH16 4UU, United Kingdom.

[6] School of Medicine, University of Dundee, Dundee, DD1 9SY, United Kingdom.

[7] School of Veterinary Medicine and Science, University of Nottingham, Loughborough,

LE12 5RD, United Kingdom.

[8]Faculty of Health and Life Sciences, University of Bristol, Bristol, United Kingdom

[9]Division of Clinical Medicine, School of Medicine and Population Health, University of

Sheffield, Beech Hill Road, Sheffield, S10 2RX, United Kingdom

Corresponding author: Adam Watkins. Division of Clinical Medicine, School of Medicine

and Population Health, University of Sheffield, Beech Hill Road, Sheffield. S10 2RX, United

Kingdom. a.watkins@sheffield.ac.uk

Keywords: Paternal diet; metabolic disorder; placental development; fetal programming;

sexual dimorphism

Abstract

The association between sub-optimal paternal diet and offspring well-being is becoming established. However, the underlying mechanisms are yet to be fully defined. The aim of this study was to establish the impact of over- and under-nutrition, with or without macronutrient supplementation, on male reproductive fitness and post-fertilisation development. Male C57/BL6J mice were fed either control diet (CD), isocaloric low protein diet (LPD), high fat/sugar 'Western' diet (WD) or LPD or WD supplemented with methyl-donors and carriers (MD-LPD or MD-WD respectively) for 8 weeks before mating with virgin C57/BL6J females. Placental tissue was collected at embryonic day (E)8.5, to assess early placental (ectoplacental cone) morphology and metabolism and E17.5 for sex-specific transcriptomic profiling. Post-mating, stud male tissues were harvested for assessment of testicular morphology and gene expression, gut microbiota composition and metabolic status. WD and MD-WD males displayed increased adiposity, hepatic cholesterol and free fatty acids and gut microbiota dysbiosis when compared to CD fed males. In the testes, WD and MD-WD perturbed the expression of genes associated with metabolism and transcription regulation. Additionally, we observed differential expression of multiple genes within the Wnt signalling pathway, central in the regulation of cellular proliferation, migration, survival, and cell fate determination during development. Despite no impact on fundamental male fertility, significant changes in ectoplacental cone metabolism, fetal growth, and placental gene expression were observed in response to specific dietary regimens. Interestingly, while CD male and female placentas displayed 301 genome-wide, sexually-dimorphic genes, LPD, MD-LPD, WD and MD-WD male and female placentas possessed only 13, 0, 14 and 15 sexually-dimorphic genes respectively. Our data show that while sub-optimal paternal diet has minimal impact on male fertility, fetal and placental development are perturbed in a sex-specific manner.

Introduction

Under- and over-nutrition are directly linked to poor physiological, metabolic and

reproductive health. The prevalence of poor dietary habits is increasing worldwide, with both

under- and over-nutrition rising dramatically. Current estimates suggest that the combined

prevalence of under- and overweight has increased in over 160 countries in both adults and

school-age children (Collaboration, 2024). Therefore, there has been significant interest in the

impact of sub-optimal nutrition and dietary supplementation on physiological, metabolic, and

reproductive health (Jawaid et al., 2021).

The consumption of diets high in fat and/or sugar are central to the development of various

metabolic disorders including obesity, hypertension, non-alcoholic fatty liver disease

(NAFLD; also referred to as Metabolic Dysfunction-Associated Steatotic Liver Disease,

MASLD) (Ludwig et al., 2001, Nielsen et al., 2010, von Frankenberg et al., 2017) and

metabolic syndrome (O'Neill and O'Driscoll, 2015, Saklayen, 2018). Poor quality diet has

also been linked to impairments in fertility and reproductive fitness. Obesity has been linked

to a reduction in testosterone levels, sperm concentration, motility and viability, and damage

to the seminiferous tubules of the testes (Crean et al., 2023, Fan et al., 2015, Leisegang et al.,

2014, Lotti et al., 2013). Undernutrition has also been linked to multiple metabolic and

reproductive impairments. In rodents, a low protein diet (LPD) significantly reduces body

weight, weight of the testes, epididymis and seminal vesicles, and serum testosterone and

follicle-stimulating hormone levels (Ajuogu et al., 2020).

Poor paternal diet also impacts negatively on fetal development and adult offspring health

(Batra et al., 2022, Capobianco and Pirrone, 2023, Fleming et al., 2018). Impairments in

rodent fetal and placental development have been reported in response to diets low in protein

(Morgan et al., 2021, Watkins et al., 2017), folate (Lambrot et al., 2013) and high in fat.

(Jazwiec et al., 2022). Separately, children born to fathers with early onset type-2 diabetes are

leaner and have a higher insulin sensitivity than children from healthy men (Penesova et al.,

2010). Additionally, children from men with type-2 diabetes at the time of conception have a

lower weight at birth, but with increased BMI in later life and a greater risk of developing

type-2 diabetes and hyperlipidaemia themselves (Moss and Harris, 2015, Penesova et al.,

2010, Silva et al., 2017). In rodent models, paternal high fat and sugar diets perturb offspring

insulin and glucose homeostasis (Cesar et al., 2022), behaviour and gut microbiota

composition (Bodden et al., 2022). Interestingly, poor paternal diet impacts on offspring

health in a sex-specific manner. Sex-specific changes in offspring tissue lipid abundance have

been reported following a paternal low protein diet (Furse et al., 2023, Morgan et al., 2022,

Morgan et al., 2020b, Watkins et al., 2018). Similarly, paternal high fat diet in rats programs

pancreatic β-cell dysfunction and impaired glucose metabolism in female offspring (Ng et al.,

2010) and adult-onset obesity in males (Sanchez-Garrido et al., 2018).

Separate to macro-nutrient modification, the significance of dietary micro-nutrients such as

vitamins, minerals and antioxidants has also been assessed. Studies have focused on

supplementation as a potential means to reverse or alleviate the effects of sub-optimal diets.

In rats, the detrimental effects of a high fat/sucrose diet on liver steatosis were ameliorated

through the supplementation with a range of methyl donors and carriers such as folate,

choline, betaine, and vitamin B12 (Cordero et al., 2013a, Cordero et al., 2013b) In mice,

supplementing calorically restricted males with vitamins and antioxidants prevented sperm

oxidative damage and normalized offspring growth and adiposity (McPherson et al., 2016).

Similarly, supplementation of a paternal LPD with methyl donors and carriers has been shown

to normalise patterns of fetal over-growth in mice (Morgan et al., 2020a).

While the association between paternal diet and offspring well-being is becoming established,

the underlying mechanisms are yet to be fully defined. The aim of this study was to establish

and compare the impact of over- and under-nutrition on male reproductive fitness and post-

fertilisation development. As poor micronutrient status has also been associated with

perturbed fetal development (Lambrot et al., 2013), we additionally explored the role that

supplementation with specific methyl donors and carriers may play in ameliorating any

detrimental effects of poor paternal diet.

Results

Male physiology and metabolic status

To investigate the impact of over- and under-nutrition, with or without methyl-donor and

carrier supplementation, on male reproductive fitness, we fed male C57BL/6J mice one of

five diets comprising a control diet (CD), low protein diet (LPD), LPD supplemented with

methyl donors (MD-LPD), Western diet (WD) or WD supplemented with methyl donors

(MD-WD) for up to 24 weeks. Over the course of the feeding period, all experimental groups

became lighter than CD males (Figure 1A). However, no significant differences were

observed when analysed at individual time points or as average growth profile (area under the

growth curve). At the time of cull, there was no difference in mean testicular (Figure 1B) or

seminal vesicle (Figure 1C) weights. MD-WD had an increased total fat mass (combined

weight of gonadal, peri-renal, inguinal and interscapular fat pads) when compared to CD fed

males (Figure 1D, P = 0.01). Analysis of white (WAT, combined weight of gonadal, peri-

renal and inguinal fat) to brown adipose tissue (BAT; interscapular) ratio revealed an increase

in WD and MW-WD males when compared to CD fed males (Figure 1E, P <0.05). Next, we

assessed serum profiles of central indicators of inflammatory status (Tnf) and testicular

regulation (Inhibin β-A chain). There was no difference in the concentrations of Tnf (Figure

1F) between males. However, WD fed males displayed an elevated concentration of serum

Inhibin β-A chain, a paracrine regulator of germ cell proliferation, spermatogenesis and

Leydig cell steroidogenesis, when compared to LPD and MD-LPD fed males (Figure 1G, P

<0.01).

As we observed significant changes in adiposity in response to our diets, we assessed stud

male metabolic status. There were no differences in the concentrations of serum glucose

(Figure 2A) or insulin (Figure 2B) between groups. In contrast, WD males displayed an

elevated concentration of hepatic cholesterol when compared to CD males (Fig, 2C) while

MD-WD males displayed elevated levels when compared to CD, LPD and MD-LPD males

(Fig, 2C, P <0.05). WD males also displayed elevated levels of free fatty acids (FFAs) when

compared to CD, LPD and MD-LPD males (Figure 2D, P <0.05), while the levels of FFA in

MD-WD males was not significantly different to that of either CD or WD males. No

differences in the concentration of hepatic triglyceride were observed between groups (Figure

2E).

Analysis of gut bacterial profiles revealed no difference in overall bacterial diversity (Figure

3A) or species evenness (Figure 3B) between groups. When analysed at the phylum level

(Figure 3C), MD-WD males displayed a significant increase in the abundance of

Deferribacteres (Figure 3D; P = 0.042), while WD and MD-WD males displayed an increased

abundance of Proteobacteria (Figure 3E; P <0.05) when compared to CD males. Furthermore,

the abundance of TM7 (Saccharibacteria) was decreased in MD-WD males when compared to

LPD and MD-LPD fed males (Figure 3F; P <0.05). Analysis of bacteria at the family level

identified significant reductions in the abundance of Turicibacteraceae and Lachnospiraceae

in both WD and MD-WD males when compared to CD fed males (Figure 3G, P <0.05).

Additionally, the abundance of S24-7 (Muribaculaceae), Clostridiaceae, Dehalobacteriaceae,

Ruminococcaceae and Desulfovibrionaceae were all significantly altered (both increased and

decreased) in MD-WD males when compared to CD males (Figure 3G; P <0.05).

Sub-optimal diet alters seminiferous tubule cytoarchitecture and cellular distribution

As diet and nutritional status have been directly linked to reproductive health, we next

assessed testicular and epididymal morphology. Gross histological analysis (Figure 4A) of

testicular tissue revealed an increased number of tubules displaying abnormalities such as

vacuoles, loss of the epithelium or complete loss of the tubular architecture (Figure 4B) in

WD and MD-WD males when compared to CD, LPD and MD-LPD males (Figure 4C,

P<0.01). To analyse seminiferous tubule composition further, we stained testicular sections

from the same males to determine Sertoli (Sox-9), spermatocytes and spermatids (Ddx4) and

spermatogonial stem (Plzf) cells (Figure 4D) numbers. There were no differences in the mean

number of nuclei per tubule (as determined via DAPI staining; Figure 4E), the number of

Sertoli cells per tubule (Figure 4F) or the number of spermatocytes and spermatids per tubule

(Figure 4G). However, a significant reduction in the number of Plzf[+] spermatogonial stem

cells was detected in tubules from LPD and WD males when compared to CD males (Figure

4H, P <0.05). In contrast, no difference in the number of Plzf[+] cells were observed in MD-

LPD and MD-WD testes when compared to CD males (Figure 4H).

Testicular gene expression is perturbed in a diet specific manner

To define the impact of diet on testicular gene expression profiles, we performed a micro-

array transcriptomic analysis using testicular tissue from the same males. We identified 0, 0,

402 (267 down-regulated, 135 upregulated) and 285 (187 down-regulated, 98 up-regulated)

differentially expressed (fold change <-1 or >1; Padj <0.05) genes in LPD, MD-LPD, WD

and MD-WD testes when compared to CD testes (Figure 5A-C, F). In WD testes, we

identified 267 downregulated and 135 upregulated genes (Figure 5C) associated with an up

regulation of fatty acid metabolism (GO:0006631) and a down regulation of chromatin

binding (GO:0003682), RNA binding (GO:0003723) and actin filament organization

(GO:0007015) pathways (Figure 5D). Network analysis of the 402 differentially expressed

genes within the WD testes identified several modular relationships between genes (Figure

5E). Specifically, we observed significant (Padj <0.05) gene nodes associated with the

expression of *Ywhae* (FC -1.14), *Rara* (FC -1.35), *Dnmt1 (*FC -1.36), *H3f3a* (FC 1.12), and

Wdr5 (FC -1.3). In MD-WD testes, we identified 187 down-regulated, 98 up-regulated genes

(Figure 5F) associated with an up-regulation of RNA splicing (GO:0008380) and RNA

binding (GO:0003723), and a down-regulation of DNA damage (GO:0006974), protein

transport (GO:0030154), cell differentiation (GO:0015031) and RNA binding (GO:0003723)

pathways (Figure 5G). In contrast to WD testes, we observed an integrated gene network

based around the expression of *Snrpg* (FC -1.27), *Top1* (FC 1.38) and *Hnrnph1* (FC -1.3)

(Figure 5H) in MD-WD testes. We also observed networks linked to *H3f3a* (FC -1.14), *Mdm2*

(FC 1.24) and *Pcgf2* (FC -1.52) in MD-WD testes. As we identified no differentially

expressed genes in LPD (Figure 5A) or MD-LPD (Figure 5B) testes, we conducted gene

ontology analysis of genes displaying a fold change <-1 or >1 and a P <0.05. In LPD testes

we identified 910 genes (490 down-regulated, 420 up-regulated) associated with an

upregulation of metabolic processes and a down regulation of RNA binding and transcription

regulation. In MD-LPD testes, we identified 3425 (2095 down-regulated, 1330 up-regulated)

genes associated with an upregulation of RNA splicing (GO:0008380) and a down regulation

of extracellular structure organization (GO:0043062), T cell activation (GO:0046631) and

MAPK regulation (GO:0043410).

Paternal low-protein diet alters early placental morphology and metabolism

Recently, we have shown that while our LPD, MD-LPD, WD and MD-WD diets do not

detrimentally affect male fertility, they do have a significant impact on post-fertilisation

embryonic development (Morgan et al., 2024). To establish whether sub-optimal paternal diet

influenced early placental development, whole conceptus morphology was analysed at

embryonic (E) day 8.5. We observed no difference in the mean number of implantation sites

between groups (Figure 6A). However, morphological assessment of whole embryo

implantation sites (Figure 6B) revealed a significant reduction of invasion depth in LPD

embryos compared to MD-LPD embryos (Figure 6C). Furthermore, LPD embryos

demonstrated a trend towards a smaller EPC area (Figure 6D) and an increased angle of

misalignment from the central maternal channel (Figure 6E) when compared to CD embryos

(P <0.1). To explore the impact of paternal diet on early placental dynamics further,

individual ectoplacental cones (EPCs) were isolated and cultured for either 24 or 48 hours

(Figure 6F). There was no difference in mean trophoblast outgrowth (Figure 6G) or central

EPC area (Figure 5H) in either the 24- or 48-hour explants.

Sup-optimal paternal diet alters fetal weights in late gestation in a sex-specific manner

Next, we examined the impact of paternal diets on late gestation (E17.5) fetal and placental

development. There was no difference in mean fetal weight between groups. However, in

both LPD and MD-LPD groups, a significantly higher proportion of fetuses displayed a

weight below the 10th centile when compared to CD fetuses (15% and 19% respectively, P

<0.05, Figure 7A). No differences in fetal weight distribution were observed in either WD or

MD-WD groups (Figure 7A). When fetal weights were separated by fetal sex, LPD, MD-LPD

and MD-WD female fetusess were significantly lighter than CD females (P <0.05; Figure

7B). However, there were no differences in mean fetal weight between any of the male groups

(Figure 7C). When assessing the influence of fetal sex on weight, we observed that males

were typically heavier than females, with the difference being significant for CD, LPD, WD

and MD-WD males when compared to their respective females (P <0.05, Figure 7D).

Analysis of mean placental weight showed no difference between groups. However, placentas

from LPD fed males showed a higher proportion above the 90^{th} centile for weight when

compared to CD males (Figure 7E, P <0.05). Analysis of placental weight by fetal sex showed

no difference for either females (Figure 7F), or males (Figure 7G) in any groups. However, in

line with fetal weight, males had heavier placentas than females in all groups (P <0.001,

Figure 7H).

Finally, while there was no difference in mean fetal:placental ratio across all groups, a

significantly higher proportion of LPD fetuses displayed a ratio below the 10^{th} centile, and

MD-WD fetuses showed a reduction in the number above the 90^{th} centile (Figure 7I; P <0.05),

when compared to CD fetuses. Analysis of fetal:placental ratio revealed a significantly lower

ratio in LPD females when compared to CD and WD females (P <0.05, Figure 7J). In

contrast, no differences between any of the male groups was observed (Figure 7K). When

comparing females to males, we observed a lower fetal:placental ratio in CD, WD and MD-

WD males (P <0.05, Figure 7L) when compared to their respective females. However, no

differences were observed between LPD or MD-LPD males and females (Figure 7L).

Paternal sub-optimal diet has minimal impact on placental morphology

As fetal placental ratio appeared altered in response to paternal diet, we next assessed placental morphology, measuring the relative areas of the decidua, junctional zone, labyrinth zone and chorionic plate (Figure 8A). There was no difference in mean cross sectional area (Figure 8B) or in the relative proportion of each placental compartment (Figure 8C) between diets. Additionally, we undertook a stereological analysis of placenta morphology (Figure 8D-F) observing no difference in mean placental volume (Figure 8G), labyrinth zone volume (Figure 8H), junctional zone volume (Figure 8I), maternal blood volume (Figure 8J), maternal blood surface area (Figure 8K) or fetal capillary surface area (Figure 8L).

Paternal sub-optimal diets influence sex-specific placental transcriptome

Finally, we performed a transcriptomic analysis of the late gestation placenta by RNA-Seq using 4 male and 4 female placentas from each dietary group. Initial analysis, separating the results by diet and sex, identified 0, 102, 34 and 77 differentially expressed genes (Padj <0.05) between LPD, MD-LPD, WD, MD-WD and CD male placentas respectively (Figure 9A). Pathway analysis of the 102 differentially expressed genes in MD-LPD male placentas identified pathways involved in metabolism (Type 1 Diabetes mellitus), immune system regulation (antigen processing, complement and coagulation cascades), cellular adhesion, signalling and senescence (Figure 9B). No significant pathway enrichment was found for the differentially expressed genes in WD or MD-WD male placentas when compared to CD males. Similarly, when comparing female placentas, we observed 0, 3, 97 and 52 differentially expressed genes (Padj <0.05) when comparing LPD, MD-LPD, WD and MD-WD with CD placentas respectively (Figure 9C). Pathway analysis of the 97 differentially expressed genes in WD female placentas identified pathways involved in lipid metabolism and signalling (RAS, GnRH) pathways (Figure 9D). No significant pathways were identified

for either MD-LPD or MD-WD female placenta differentially expressed genes. These data

suggest that paternal diet may be influencing placental gene expression in a sex-dependent

manner, with more differentially expressed genes observed within the male MD-LPD and

MD-WD groups, while the WD and MD-WD females shower a greater change in gene

expression profiles. Principal component analysis of expression between CD males and

females suggested the existence of a sex-specific profile (Figure 9E). Additional analysis

identified 301 differentially expressed placental genes between CD males and females (Padj

<0.05, Figure 9F) (See Supplemental Data S1 for full list). Of these 301 genes, 43 were

upregulated in males and 258 were upregulated in females. Gene ontology analysis identified

pathways involved in vascular development, developmental processes and signal transduction

(Figure 9G). Interestingly, comparison of these same 301 genes in LPD, MD-LPD, WD and

MD-WD placentas showed no such sexual dimorphism (Figure 9H-K). In total, we identified

13, 0, 14 and 15 genes to be differentially expressed between males and females in the LPD,

MD-LPD, WD and MD-WD groups respectively (Figure 9L-O). In total, we identified just 9

differentially expressed genes, and one unprocessed pseudogene, present in all 5 dietary

groups (Figure 9P). Of the 9 conserved differentially expressed mRNAs, 5 were located on

the X chromosome (*Eif2s3x, Taf1, Otg, Xist, Kdm5c*), while 4 were on the Y chromosome

(*Kdm5d, Uty, Ddx3y, Eif2s3y*) (Figure 9Q). Separate to these, LPD placentas displayed

sexually dimorphic gene expression on chromosomes 5 and 13, WD placentas had sexually

dimorphic gene expression on chromosomes 3, 5 and 15, and MD-WD demonstrated sexually

dimorphic gene expression on chromosomes 3, 5, 6 and 14 (see Supplementary Data S1).

Discussion

Consumption of an unbalanced or unhealthy diet is a major risk factor for a range of diseases

such as obesity, diabetes and cardiovascular disease. Poor parental diet prior to conception

also impacts on patterns of post-fertilisation development, shaping fetal and placental

dynamics and increases the risk for a range of non-communicable disorders in adult life

(Ozanne et al., 2004). Under the current study, we explored the impact of both paternal under-

(LPD) and over-nutrition (WD) on male reproductive health, fetal growth and placental

development. We also investigated the impact of supplementing these diets with a range of

methyl-donors and carriers to define if their use could negate any detrimental influences

imposed by the diets. We observed that males consuming a high fat/sugar diet (WD and MD-

WD), displayed a series of physiological changes in their metabolic health, gut bacterial

profiles and testicular morphology and gene expression. While both over- and under-

nutritional regimens had no impact on fundamental male fertility, we observed significant

changes in fetal and placental weights in late gestation. Furthermore, we observed a

significant loss in sexual dimorphism in the late gestation placenta. While sex-specific

differences in fetal and adult offspring are reported widely in response to both maternal and

paternal diet (Dearden et al., 2018, Rosenfeld, 2015), our data provide additional insight into

how sex-specific paternal programming may be mediated during development.

While we observed minimal impact of the LPD and MD-LPD on male metabolic health, mice

fed the WD and MD-WD displayed significant changes in adiposity and hepatic cholesterol

and FFAs. Excessive accumulation of cholesterol and FFAs within the liver are central

hallmarks of metabolic conditions such NAFLD and MASLD. While elevated hepatic FFAs

have been linked to insulin resistance and elevated levels of proinflammatory mediators, we

observed no change in circulating glucose, insulin or Tnf levels. This is in line with studies

showing that nearly half of patients with NAFLD/MASLD also do not present with insulin

resistance (Singh et al., 2015), highlighting it heterogenous nature. Changes in metabolic

status have also been linked to gut dysbiosis. We observed differences in the abundance of

Defferibacteres, Proteobacteria and TM7 in fecal samples from WD and MD-WD males.

Similar increases in Deferribacteres have been observed in mice maintained on high fat diets

(Liu et al., 2019, Serino et al., 2012), while the abundance of TM7 has been linked with

elevated BMI, fat mass and inflammatory status in humans (Gomes et al., 2020). Separately,

negative links between TM7 abundance and body weight and adiposity in rodents have been

identified (de la Garza et al., 2022, Hu et al., 2022). While our study indicates significant

shifts in male metabolic homeostasis, further studies are needed to define how metabolic,

inflammatory and microbiota status interact and their impact on male reproductive health. For

example, bacterial metabolites such as 5-hydroxytryptamine (5-HT) have been shown to

promote sperm hyperactivation and the acrosome reaction in mice (Sugiyama et al., 2019),

hamsters (Fujinoki, 2011) and men (Omote et al., 2023). Additionally, many polyunsaturated

fatty acids (PUFAs) produced by the gut microbiota have significant role in stabilising the

sperm plasma membrane (Lv et al., 2024). Finally, additional microbiota derived metabolites

such as androgens (Collden et al., 2019) and hormones including GLP-1 and peptide YY

(PYY) (Cannarella et al., 2021) have been associated with changes in sperm production.

Interestingly, studies have also shown that fecal transplants can be used to treat infertility and

improve sperm quality in mice (Hao et al., 2022, Zhang et al., 2021). Therefore, additional

studies, with the inclusion of a methyl donor CD (MD-CD), are required to understand better

the connections between dietary-induced gut dysbiosis and male reproductive fitness.

Within the testis, LPD and WD males displayed a significant reduction in the number of

undifferentiated spermatogonial stem (Plzf+) cells when compared to CD males, while no

difference was observed in MD-LPD and MD-WD males. While the loss of Plzf is not

detrimental for spermatogenesis, Plzf ensures spermatogonial stem remain in a state of

undifferentiation, interacting with other transcription factors such as SRY-box TF 3 (SOX3)

and Spalt like TF 4 (SALL4) to establish the chromatin and transcription landscape in

spermatogonial stem cells (Yi et al., 2025). However, the regulation of Plzf by methyl donors

is currently undetermined. Separately, WD and MD-WD males displayed an increased rate of

tubule abnormalities when compared to CD males. In rodents, exposure to a high fat diet has

been shown to induce tubule vacuolization atrophy (Ghosh and Mukherjee, 2018) as well as

disrupting the essential adhesion between spermatogenic and Sertoli cells, resulting in Sertoli

cell atrophy (Liu et al., 2014). Sertoli cells are essential for supporting spermatogenesis and

their loss is associated with male infertility. Sertoli cells also produce inhibin β and its serum

levels correlate positively with testicular size and sperm production rates (Luisi et al., 2005).

Inhibin production is also tightly linked to the secretion of FSH from the pituitary, providing a

negative feedback signal to regulate spermatogenesis. As we observed a significant elevation

in serum inhibin β-A in WD males, but no change in Sertoli (Sox9$^+$) cell number or

fundamental male fertility, the increase in inhibin β might indicate a dysfunction in normal

hypothalamic-pituitary-testis axis regulation. However, further studies are needed to confirm

this.

Analysis of testicular gene expression revealed minimal changes in response to LPD or MD-

LPD. In contrast, testes from WD and MD-WD-fed males displayed an upregulation of genes

involved in fatty acid metabolism, mRNA processing, Wnt signalling and protein transport

processes, and a downregulation of transcription related genes. The increase in genes

associated with fatty acid metabolism (*Acadsb*, *Crat, Pla2g5*, *Daglb, Mgll*) and glycolytic

components (*Pdha1, Hk2*) link with the changes observed in metabolic status in these males.

Furthermore, we observed up regulation of *Paf1* and *Ctr9,* both components of the Paf1C

complex. PAF1C associates with the promoter and coding sequences of active genes (Pokholok et al., 2002) and interacts with a series of transcriptional elongation factors (Pokholok et al., 2002, Squazzo et al., 2002). Studies have identified Paf1C as crucial for the acquisition of transcription-associated histone modifications such as H3 methylation at K4, K36, and K79 (Cao et al., 2015, Mbogning et al., 2013). Interestingly, we identified an increase in the expression of the histone variants *H3f3a* and *H3f3b*. Unlike most other canonical histones, H3.3 is not completely removed from the chromatin during spermatogenesis and is enriched at genomic regions important for zygotic development in mice and men (Brykczynska et al., 2010, Hammoud et al., 2011). Studies have shown that disruption of H3K4me2 marks in mouse sperm have significant consequences for offspring health over multiple generations (Siklenka et al., 2015) and levels of sperm H3K4me2 may serve as a metabolic sensor linking paternal diet with offspring metabolic ill-health predisposition (Pepin et al., 2022). The downregulation of transcription regulating genes such as *Dnmt1, Mta1, Phf1, Pwwp2a, Wdr5, Hsf1, Nfrkb. Pelp1, Prox1* and *Rara* suggest paternal HFD might influence differential patterns of testicular transcriptional regulation, which could affect sperm composition and post-fertilisation development (Cui et al., 2025). Finally, we observed a down regulation of multiple RNA binding genes in both WD and MD-WD testes. Recently, the significance of RNA-binding proteins as critical regulators of spermatogenesis and sperm function has been recognised (Li et al., 2024). As many RNA-binding proteins play central roles in orchestrating intricate transcriptional and post-transcriptional regulatory networks, their function as essential regulators of testicular germ cell development is becoming defined (Gao et al., 2026).

Together, these findings suggest that sub-optimal male diets have the potential to alter sperm modifications, which are not simply corrected by methyl-donor supplementation. To

determine whether these paternal diet-driven changes to male reproductive physiology

impacted on post-fertilisation dynamics, we analysed early (E8.5) and late (E17.5) fetal and

placental development. Analysis of E8.5 placental invasion revealed poor paternal diet had

minimal impact on overall morphology or invasion depth *in vivo*. However, we observed

significant sex-specific influences of paternal diet on late gestation fetal and placental growth.

During fetal development, differences in patterns of male and female growth have been

described from as early as the first trimester with males growing at a faster rate (Broere-

Brown et al., 2016). We identified over 300 sexually dimorphic, differentially expressed

genes between CD males and females, with the majority being upregulated in females. Gene

ontology analysis identified vascular, immunological, extracellular matrix organisation and

cell-signalling pathways as being differentially expressed, mirroring observations from male

and female term placentas in humans (Buckberry et al., 2014). While female human term

placentas have increased expression of multiple immune regulation genes when compared to

males (Sood et al., 2006), Y-chromosome genes such as *Ddx3y, Uty,* and *Kdm5d,* encode

epitopes that contribute to minor histocompatibility antigens present throughout the placenta

(Cvitic et al., 2013). Studies have also shown that cultured trophoblast tissues isolated from

human healthy male placentas secrete more TNF and less IL-10 in response to

lipopolysaccharide than female tissues (Yang et al., 2021). In contrast, genes such as *Tgfb1,*

which was upregulated in CD females, are critical in regulating fetal-maternal immune

tolerance (Yang et al., 2021). Furthermore, studies have shown female placentas to be more

responsive to shifts in maternal inflammation and dietary status than males (Osei-Kumah et

al., 2011, Sedlmeier et al., 2014). Similar to our data, studies have also reported upregulation

of receptor-ligand signalling in human female placentas including several collagens, laminins

and integrins, suggestive of an enhanced invasion and placentation potential (Braun et al.,

2021). Finally, we observed up regulation of genes involved in steroid synthesis and

endocrine regulation in female placentas. Placental endocrine function operates to regulate

both maternal physiological response and fetal development (Stern et al., 2021). In contrast to

our CD males and females, all other groups displayed a dramatic loss of sexual dimorphism.

While the underlying mechanism(s) for our observed loss of sexual-dimorphism are unknown,

perturbed paternal sperm epigenetic status could modulate early embryo X-chromosome

inactivation and/or X-chromosome dosage in females, resulting in differential profiles of

autosomal gene expression (Gonzalez et al., 2018, Petropoulos et al., 2016).

Collectively, we observe that paternal LPD or WD, with or without supplementation, had no

impact on overall male fertility, but have influences on male reproductive physiology. While

paternal LPD induced minor changes in early placental morphology and metabolism, these

effects were small, and the consequences of these changes remain to be defined. In late

gestation, paternal diet affected fetal development in a sex- and diet-specific manner. While

we observed significant sexual dimorphism in gene expression patterns between CD males

and females, these differences were largely removed in response to our experimental diets.

Such loss of sexual- dimorphism could provide one process through which poor paternal diet

programs offspring ill-health in adulthood. However, the precise mechanisms through which

this occurs remain to be determined.

Materials and Methods

Mice diet regime and matings

All animal procedures were conducted in accordance with the UK Home Office Animals

(Scientific Procedures) Act 1986 and carried out under Project License PP8899264 with local

ethical approval at University of Nottingham. C57BL/6J mice (Charles River, UK) were

housed in controlled 12/12-hour light/dark conditions with a constant temperature ($21° C \pm 3°$

C) and *ad libitum* access to water. Virgin 8-week old males (n = 8 per treatment group) were

fed either control diet (CD; 18% casein, 21% sucrose, 0% milk fat, 0% cholesterol), isocaloric

low protein diet (LPD; 9% casein, 24% sucrose, 0% milk fat, 0% cholesterol), 'Western' diet

(WD; 19% casein, 34% sucrose, 20% milk fat, 0.15% cholesterol) or LPD or WD

supplemented with methyl-donors and carriers (5 g/kg diet choline chloride, 15 g/kg diet

betaine, 7.5 g/kg diet methionine, 15 mg/kg diet folic acid, 1.5 mg/kg diet vitamin B12;

termed MD-LPD or MD-WD respectively), for up to 24 weeks. Exact dietary formulations

are outlined in Supplementary Table S1. Virgin female C57BL/6J mice were maintained on

standard rodent chow (rat/mouse No.1 maintenance diet, Special Diet Services) throughout

study the study. Females were mated at 9-weeks old (+/-7 days) with stud males who had

been fed their respective diet for a minimum of 8 weeks. Successful mating was confirmed by

the presence of a copulation plug and denoted as embryonic day (E)0.5. Dams were

euthanized via cervical dislocation on either E8.5, for the collection of ectoplacental cones

(EPCs), or E17.5 for fetal and placental tissues. Stud males were culled by cervical

dislocation for the collection and storage of tissues. Blood was collected via heart puncture,

allowed to clot on ice and centrifuged at $10,000 \times g$, 4 °C for 10 minutes before storage of the

serum at -80 °C. Heart, kidney, liver, testis, and caudal epididymis were dissected, weighed

and either snap frozen prior to storage at -80 °C, or fixed overnight in 10 % neutral buffered

formalin (Sigma, UK) at 4 °C before processing and embedding in paraffin wax. Fecal pellets

were collected from the descending colon using sterile forceps and placed in 2 ml

DNase/RNase-free collection tubes, snap frozen and stored at -80 ℃. A second batch of males

(n = 6) were fed using the same experimental protocol and utilized for the analysis of fat pad

mass.

Stud male metabolic status assessment

Stud male non-fasting serum glucose was measured using the Glucose Colorimetric Detection

Kit (EIAGLUC, Thermo Fisher Scientific, UK), non-fasting insulin was measured using a

Rat/Mouse Insulin ELISA Kit (EZRMI-13K; Millipore, UK), Tnf was measured using the

TNF alpha SimpleStep ELISA® Kit (ab208348, Abcam®), inhibin β-A chain was measured

using the FineTest® Mouse Inhba ELISA Kit (EM0273, Wuhan Fine Biotech Co. China),

non-fasting liver tissue total cholesterol was measured using the Cholesterol Quantification

Kit (MAK043, Sigma-Aldrich, UK), non-fasting hepatic free fatty acids (FFA) were

measured using the Free Fatty Acid Quantitation Kit (MAK044, Sigma-Aldrich, UK) and

non-fasting hepatic triglycerides were measured using the Triglyceride Quantification Kit

(MAK266, Sigma-Aldrich, UK), all in accordance with the manufacturers' instructions.

Gut microbiota sequencing

DNA was extracted from stool pellets using the QIAamp DNA Stool Mini Kit (Qiagen, UK)

following the manufacturer's instructions and stored at -80 °C. 16S rDNA sequencing (V3-V4

region) was conducted as described previously (Morgan et al., 2022). The extracted stool

DNA samples (n = 6 per diet group) were sequenced using the Illumina MiSeq Reagent kit v3

on the Illumina MiSeq platform (Illumina), in accordance with Illumina 16S Metagenomic

Sequencing Library Preparation protocol. Briefly, 16S rRNA amplicons were generated using

the forward 5'

(TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGGNGGCWGCAG) and

Reverse   5'

(GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACTACHVGGGTATCTAATC

C) primers, flanked by Illumina adapter-overhang sequences. Illumina dual index barcodes

(Illumina XT Index Kit v2, Set A: FC-131-2001) were attached to each amplicon. PCR clean-

up was conducted using AMpure XP beads (Beckman; A63882). Library fragment-length

distributions were analysed using the Agilent TapeStation 4200 and the Agilent D1000

ScreenTape Assay (Agilent; 5067-5582 and 5067-5583). The generated libraries were pooled

in equimolar amounts and the pool was size selected using the Blue Pippin (Sage Science)

and a 1.5 % Pippin Gel Cassette (Sage Science; BDF2010). The samples were then run over a

shared 300 paired end (PE) MiSeq run to deliver about 60-80,000 PE reads per sample. The

sequencing run additionally had a 20 % PhiX library spike-in as an internal quality control.

Raw reads were processed by Qiime2 pipeline and trimmed. Greengenes version 13.8 was

used in the classification (Caporaso et al. 2010).

Testicular histology

Parafin embedded testis tissue (n = 8 per group) was sectioned at 5 μm using a Leitz 1512

rotary microtome (Leica). For analysis of seminiferous tubule morphology, sections were

processed and stained with Haematoxylin-Eosin prior to imaging using a Leica DMRB

microscope with an Oasis glide scanner. Image analysis was carried out using Fiji. For each

testis sample, an average of 50 seminiferous tubule cross-sections were analysed.

For the determination of seminiferous tubule cell types, additional testis sections were

dewaxed, rehydrated into PBS, and equilibrated in 100 mM citrate buffer (pH 6.0) at room

temperature (RT) prior to being microwaved in 0.1M tri-sodium citrate buffer (pH 6.0) for 5

minutes. The sections were incubated in blocking solution (5 % bovine serum albumin, 20 %

normal goat serum in 0.1 % Triton X-100 in PBS; Sigma, UK) for 1 hour before incubating

with anti-Ddx4 antibody (to identify germ cells; ab27591, 1:100; Abcam, UK) or Rabbit anti-

Sox9 Antibody (to identify Sertoli cells: ab5535, 1:500; Chemicon, UK) in blocking solution,

and incubated at 4° C overnight. Slides were washed (three times in 0.1% Triton X-100 in

PBS; PBST) prior to incubation in Alexa Fluor 568 (A-21124, 1:200; Invitrogen, UK) in

blocking solution for the detection of Ddx4. For the detection of Sox9, slides were incubated

with goat anti-rabbit biotinylated antibody (E0432, 1:200; DakoCytomation, UK) for 1 hour

in the dark in blocking solution prior to Alexa Fluor 488 streptavidin conjugate, washed in

PBST prior to incubation with Alexa Fluor 488 (S-32354, 1:200; Invitrogen, UK) in blocking

solution for 30 minutes at room temperature. For the detection of Plzf, sections were

incubated overnight with anti-Plzf antibody (to identify spermatogonial stem cells: ab189849,

1:3000; Abcam, UK) at 4 $^{\circ}$C. Sections were washed in PBST prior to incubation with Alexa

Fluor 568 (ab175471, 1:200; Abcam) in blocking solution for 1 hour at room temperature.

Negative controls (absence of primary or secondary antibodies) were also conducted to

confirm staining specificity (see Supplemental Figure 1). Slides were washed (PBST) before

mounting with Antifade Mounting Media with DAPI (Vectashield, UK). Sections were

imaged using a Nikon Eclipse 90i fluorescent microscope with a mercury-fiber illuminator

Nikon Intensilight C-HGFI and a Hamamatsu ORCA-ER Digital camera (C4742-80) at 10

and 20 x magnification. Images were analysed using Volocity (Quorum Technologies Inc)

with an average of 50 seminiferous tubules counted per male.

Testicular micro-array

Total RNA was extracted from frozen testicular tissues using the RNeasy Plus Mini Kit

(Qiagen; UK) following the manufacturer's instructions. Testicular RNA was diluted to 100

ng / μl using RNAse-free water (Qiagen, UK) prior to RNA integrity assessment on a

Bioanalyzer 2100 platform (Agilent, UK). Samples with an RIN >7 were used for analysis.

The GeneChip™ WT PLUS Reagent Kit (ThermoFisher Scientific, UK) was used to prepare

RNA samples for whole transcriptome expression analysis with GeneChip™ Whole

Transcript (WT) Expression Arrays, as per the manufacturer's instructions. First-strand cDNA

was synthesized from the extracted RNA using Reverse Transcriptase, followed by a second-

strand cDNA synthesis using DNA Polymerase and RNAse H. Complementary RNA (cRNA)

was synthesized by *in vitro* transcription of the second-strand cDNA using T7 RNA

polymerase and purified. The purified ss-cDNA was enzymatically fragmented and labelled

by terminal deoxynucleotidyl transferase (TdT) using the provided DNA Labelling Reagent

following the manufacturer's instructions prior to hybridisation to onto the Clariom™ S

Assay Mouse GeneChip™ (ThermoFisher Scientific, UK). Data analysis was performed

using the Partek Genomics Suite 6.6 analysis software. Comparative analysis between the five

diet groups was performed using standard One-way ANOVAs. Dysregulated transcripts were

classified based on False Discovery Rate (FDR) = 0.05, fold change < +/-1.1, and $p \leq 0.05$.

The functional annotation and enrichment analysis web tool WEB-based GEneSeTAnaLysis

Toolkit (WebGestalt) (Liao et al., 2019) was used for Gene set enrichment analysis (GSEA).

Network analysis was conducted using Network Analyst

(https://www.networkanalyst.ca/NetworkAnalyst/home.xhtml). All gene expression data is

publicly available via Gene Expression Omnibus (GSE279868).

In vivo **EPC invasion**

Formalin-fixed, paraffin-embedded E8.5 whole implantation sites were sectioned to the mid-

point as indicated by appearance of the maternal channel. Haematoxylin-Eosin staining was

conducted on 5 µm sections and whole sites were imaged at 20x magnification using Leica a

DMRB microscope with an Oasis glide scanner. Images were imported into Fiji for the

assessment of total EPC, trophoblast specific area, depth of invasion (measured from chorion

to the furthest trophoblast going in the direction of the maternal channel) and alignment of

invasion measured according to (Wilkinson et al., 2021) within the implant site and was

conducted by a blinded observer.

Isolation and culture of EPC explants

The isolation and culture of E8.5 EPCs was conducted as previously described (Watkins et

al., 2015). Twenty-four hours prior to dissection, sterile coverslips were coated with BD

Matrigel basement membrane matrix (BD Biosciences, Oxford, UK) diluted to 6 mg/ml in

RPMI 1640 medium (Life Technologies, UK) and kept at 4° C in four-well plates. On the

morning of EPC isolation, the Matrigel-RPMI medium was replaced with 500 μl sterile

filtered RPMI-1640 medium containing 2% KnockOut Serum Replacement (Life

Technologies, UK), penicillin–streptomycin–glutamine mix (Life Technologies, UK) and

26.2 μmol 2-mercaptoethanol (Sigma–Aldrich, UK) and allowed to equilibrate at 37 ° C in

5% CO2 for at least 1 hour. Whole uteri were excised and placed in pre-warmed (37° C)

DMEM (Fisher Scientific, UK) supplemented with 10% (v/v) fetal calf serum (Life

Technologies, UK). Individual implantation sites were dissected from the uterine tissue and

placed in RPMI-1640 medium supplemented with 2% KO-serum replacement (Fisher

Scientific, UK), 1% penicillin–streptomycin–glutamine mix (Fisher Scientific, UK) and

26.2μmol 2-mercaptoethanol (Sigma–Aldrich, UK). The EPCs was then isolated from the

decidual capsules and separated from the embryo. The isolated EPCs were then placed onto

the Matrigel-coated coverslips and cultured individually at 37 ° C, 5% CO_2 for either 24 or

48 hours, with a 50% medium change after 24 hours.

After 24 or 48 hours in culture, EPC outgrowths were fixed in 4% neutral buffered formalin

(Sigma–Aldrich, UK) for 15 minutes at room temperature, washed (1x PBS) and stored at 4°

C in PBS. Fixed EPCs were permeabilised with 0.1% Triton-X100 (Sigma–Aldrich, UK) and

autofluorescence quenched with ammonium chloride (2.6 mg/ml in PBS; Sigma-Aldrich

UK). Non-specific binding was blocked with 2% BSA in PBS containing 0.1% Triton-X100

for 30 minutes at room temperature before overnight incubation at 4 ° C with a primary

antibody for α-tubulin (1:2,000; Cell Signaling, Danvers, MA, USA) in PBS with 1% BSA

and 0.1% Triton-X100. Following washing (PBS, 0.1% Triton-X100), outgrowths were

incubated with the appropriate Alexa Fluor conjugated secondary antibody (1:10,000;

Molecular Probes, Life Technologies, Paisley, UK) for 1 hour at room temperature and

counterstained with DAPI (Sigma–Aldrich, UK) for 10 minutes before mounting on slides in

DPX (Fisher Scientific, Loughborough, UK). All outgrowths were imaged using a Nikon

Eclipse 90i fluorescent microscope fitted with a mercury-fiber illuminator, Nikon Intensilight

C-HGFI and a Hamamatsu ORCA-ER Digital camera (C4742-80) and analysed using the

Volocity Software. The central EPC was defined as the central, single mass within which

individual cell nuclei could not be distinguished from those neighbouring them. Proliferative

trophoblast nuclei were defined as those located within the proximity of the central EPC as

described previously (Watkins et al., 2015).

Placental stereology

At E17.5, dams were culled for the collection of placental tissues. Fetuses and placentas were

weighted prior to fixation in Karnovsky's fixative (2.5% glutaraldehyde, 2% formaldehyde in

0.1M sodium phosphate buffer) for 16 hours at 4^o C. Fixed placentas were weighed (wet

weight, g) and converted into pre-embedding placental volumes (V_{pre}) by dividing wet weight

by tissue density (1.05 g/cm^3) prior to multistage systematic and uniformly-random sampling

(SURS) (Veras et al., 2008). Briefly, placentas were cut into ~six 2-mm slices perpendicular

to the chorionic plate. Slices were randomised to generate isotropic uniformly random (IUR)

orientation and embedded in paraffin for stereological estimation of placental compartment

volumes, maternal blood volume and surface area, and fetal capillary surface area. Shrinkage

correction (performed per placenta). Shrinkage introduced during dehydration/clearing and

paraffin embedding was quantified for each placenta individually by estimating the post-

embedding placental volume (V_{post}) using the Cavalieri Principle (Howard et al., 2005,

Ribeiro et al., 2008). Each paraffin-embedded 2-mm slice was serially sectioned at 5 µm, and

~10 evenly spaced sections per slice were selected systematic and uniformly-random through

the block, mounted, and stained with Haematoxylin and Eosin (H&E). A point grid was

applied to the sampled sections to estimate V_{post} by Cavalieri. The volume shrinkage

coefficient for each placenta was calculated as:

$$k_V = \frac{V_{\text{post}}}{V_{\text{pre}}}$$

653   (with fractional shrinkage $= 1 - k_V$). Absolute compartment volumes estimated from

654   paraffin sections were corrected back to pre-embedding dimensions as:

$$V_{\text{corrected}} = \frac{V_{\text{estimated}}}{k_V}$$

656   Surface area was estimated as the product of surface density and the corresponding shrinkage-

657   corrected placenta volume:

$$S = S_v \times V_{\text{corrected}}$$

(Where S_v is the surface density estimated by intersection counting and $V_{\text{corrected}}$ is the

placenta volume corrected after the paraffin-embedding shrinkage.

Placental RNA extraction and sequencing

Placental RNA was extracted from one quatre of an intact frozen placenta using the

miRNeasy mini kit (Qiagen, UK), following tissue disruption in Qiazol (Qiagen, UK) using

the Qiagen TissueLyserII (25Hz, 4x30 seconds), according to manufacturer's instructions.

RNA concentrations were assessed using Nanodrop and integrity was examined using Agilent

4200 TapeStation and the RNA ScreenTape Assay (Agilent; 5067-5576 and 5067-5577) with

an acceptable RIN>8. Stranded RNA-seq libraries were prepared from 1000 ng of total RNA

per sample, using the NEBNext Ultra II Directional RNA Library Preparation Kit for Illumina

(NEB; E7760) and NEBNext Multiplex Oligos for Illumina (96 Unique Dual Index Pairs)

(NEBNext; E6440). Prior to library preparation, total RNA was treated with QIAseq

FastSelect -rRNA HMR, to prevent any rRNA present from being converted into sequencing

library. Libraries were quantified using the Qubit Fluorometer and the Qubit dsDNA HS Kit

(ThermoFisher Scientific; Q32854). Library fragment-length distributions were analysed

using the Agilent 4200 TapeStation and the Agilent High Sensitivity D1000 ScreenTape

Assay (Agilent; 5067-5584 and 5067-5585). Libraries were pooled in equimolar amounts and

final library quantification was performed using the KAPA Library Quantification Kit for

Illumina (Roche; KK4824). The library pool was sequenced on the Illlumina NextSeq500

over three NextSeq500 High Output 150 cycle kits (Illumina; 20024907), to generate over 40

million pairs of 75-bp paired-end reads per sample. Raw reads were trimmed of Illumina

adapters and low quality (Q<20) nucleotides using TrimGalore (v 0.6.7)

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/). Reads shorter than 15 bp

were discarded. Trimmed reads were aligned to Mus musculus reference genome GRCm39

using HISAT2 (v 2.2.1) (Kim et al., 2019). StringTie (v 2.2.1) (Kovaka et al., 2019) was used

to assemble genes and calculate gene abundance. Differential expression analysis was

performed using DESeq2. Data was visualised using ShinyGo, ClustVis, Network Analyst

and Venny2.1 free web-based software using a pAdj<0.05, FDR <0.05 and no fold change cut

off. All gene expression data is publicly available via Gene Expression Omnibus

(GSE301011).

Fetal sex determination

DNA was extracted from fetal tail tissue using the DNeasy Blood and Tissue kit (Qiagen,

UK) according to the manufacturer's instruction. For PCR determination of fetal sex, 2 μl

(100 ng) of template DNA was added to a mastermix comprising 10 μl mastermix (2 ×

GoTaq® Green Master Mix, Promega UK), 1 μl primer mix (25 μM forward and reverse

primers) and 5 μl water per reaction. Amplification was performed using an Techgene

thermocycler using 3 primer sets specific for regions on the Y (Sry and Zfy) and X (Dxnds3)

chromosomes (primer sequences are provided in Supplementary Table S2).

Statistical Analysis

All data were assessed for normality with GraphPad Prism (version 10) or SPSS (version 28).

Early placental (EPC) data were analysed using a One-way ANOVA for normally distributed

data or a Kruskal-Wallis test for non-normally distributed data with appropriate post-hoc test.

For the analysis of stud male growth, an additional FDR correction was applied to account for

the multiple testing of weight at each individual week. Late gestation (E17.5) fetal and

placental data were analysed using a generalised linear mixed model analysis with paternal

origin of litter, and paternal duration on their respective diet incorporated as random effects

(Watkins et al., 2018). Significance was taken at $P < 0.05$.

Conflicts of interest:

The authors declare no known financial or personal interests which could have influenced the

work reported in this paper.

Author Contributions:

Conceptualization: HLM, NE, NH, MC, SH, FS, MCU, IK, NN, RTM, FL, AAC, VB, AJW

Methodology: HLM, NE, NH, MC, SH, FS, MCU, IK, NN, RTM, FL, AAC, VB, AJW

Investigation: HLM, NE, NH, MC, SH, FS, MCU, IK, NN, AAC, VB, AJW

Data curation: HLM, NE, NH, MC, SH, FS, MCU, IK, NN, RTM, FL, AAC, VB, AJW

Formal analysis: HLM, NE, NH, MC, SH, FS, MCU, IK, NN, RTM, FL, AAC, VB, AJW

Funding acquisition: AJW, RTM

Writing – original draft: HLM, VB, AJW

Writing – review & editing, HLM, NE, NH, MC, SH, FS, VW, MCU, IK, NN, STM, RTM,

FL, RSR, AAC, VB, AJW.

All authors have read and agreed to the published version of the manuscript.

Funding:

This work was supported by Biotechnology and Biological Sciences Research Council

(BBSRC) grants (BB/R003556/1 and BB/V006711/1) to AJW. This work was also supported

by a Covid support grant from the Society for Reproduction and Fertility awarded to NE.

RTM was supported by a United Kingdom Research and Innovation (UKRI) Future Leaders

Fellowship (MR/Y011783/1).

Data Availability Statement:

All data are either contained within the Supplemental Materials or are available upon

reasonable request sent to the corresponding author. Placental and testicular transcriptomic

data have been deposited with in the Gene Expression Omnibus under accession numbers

GSE301011 and GSE279868 respectively.

Acknowledgments:

The authors would like to thank the staff within the Bio Support Unit at the University of

Nottingham for animal provision and maintenance.

References

AJUOGU, P. K., AL-AQBI, M. A., HART, R. A., WOLDEN, M., SMART, N. A. & MCFARLANE,
J. R. 2020. The effect of dietary protein intake on factors associated with male
infertility: A systematic literature review and meta-analysis of animal clinical trials
in rats. *Nutr Health,* 26**,** 53-64.
BATRA, V., NORMAN, E., MORGAN, H. L. & WATKINS, A. J. 2022. Parental Programming
of Offspring Health: The Intricate Interplay between Diet, Environment,
Reproduction and Development. *Biomolecules,* 12.
BODDEN, C., PANG, T. Y., FENG, Y., MRIDHA, F., KONG, G., LI, S., WATT, M. J.,
REICHELT, A. C. & HANNAN, A. J. 2022. Intergenerational effects of a paternal
Western diet during adolescence on offspring gut microbiota, stress reactivity, and
social behavior. *FASEB J,* 36**,** e21981.
BRAUN, A. E., MUENCH, K. L., ROBINSON, B. G., WANG, A., PALMER, T. D. & WINN, V.
D. 2021. Examining Sex Differences in the Human Placental Transcriptome During
the First Fetal Androgen Peak. *Reprod Sci,* 28**,** 801-818.
BROERE-BROWN, Z. A., BAAN, E., SCHALEKAMP-TIMMERMANS, S., VERBURG, B. O.,
JADDOE, V. W. & STEEGERS, E. A. 2016. Sex-specific differences in fetal and
infant growth patterns: a prospective population-based cohort study. *Biol Sex*
Differ, 7**,** 65.
BRYKCZYNSKA, U., HISANO, M., ERKEK, S., RAMOS, L., OAKELEY, E. J., ROLOFF, T. C.,
BEISEL, C., SCHUBELER, D., STADLER, M. B. & PETERS, A. H. 2010. Repressive
and active histone methylation mark distinct promoters in human and mouse
spermatozoa. *Nat Struct Mol Biol,* 17**,** 679-87.
BUCKBERRY, S., BIANCO-MIOTTO, T., BENT, S. J., DEKKER, G. A. & ROBERTS, C. T.
2014. Integrative transcriptome meta-analysis reveals widespread sex-biased
gene expression at the human fetal-maternal interface. *Mol Hum Reprod,* 20**,** 810-
9.
CANNARELLA, R., CALOGERO, A. E., CONDORELLI, R. A., GRECO, E. A., AVERSA, A. & LA
VIGNERA, S. 2021. Is there a role for glucagon-like peptide-1 receptor agonists in
the treatment of male infertility? *Andrology,* 9**,** 1499-1503.
CAO, Q. F., YAMAMOTO, J., ISOBE, T., TATENO, S., MURASE, Y., CHEN, Y., HANDA, H. &
YAMAGUCHI, Y. 2015. Characterization of the Human Transcription Elongation
Factor Rtf1: Evidence for Nonoverlapping Functions of Rtf1 and the Paf1 Complex.
Mol Cell Biol, 35**,** 3459-70.
CAPOBIANCO, E. & PIRRONE, I. 2023. Paternal programming of fetoplacental and
offspring metabolic disorders. *Placenta,* 141**,** 71-77.
CESAR, H., SERTORIO, M. N., DE SOUZA, E. A., JAMAR, G., SANTAMARINA, A., JUCA, A.,
CASAGRANDE, B. P. & PISANI, L. P. 2022. Parental high-fat high-sugar diet
programming and hypothalamus adipose tissue axis in male Wistar rats. *Eur J*
Nutr, 61**,** 523-537.
COLLABORATION, N. C. D. R. F. 2024. Worldwide trends in underweight and obesity from
1990 to 2022: a pooled analysis of 3663 population-representative studies with
222 million children, adolescents, and adults. *Lancet,* 403**,** 1027-1050.
COLLDEN, H., LANDIN, A., WALLENIUS, V., ELEBRING, E., FANDRIKS, L., NILSSON, M.
E., RYBERG, H., POUTANEN, M., SJOGREN, K., VANDENPUT, L. & OHLSSON, C.
2019. The gut microbiota is a major regulator of androgen metabolism in
intestinal contents. *Am J Physiol Endocrinol Metab,* 317**,** E1182-E1192.
CORDERO, P., CAMPION, J., MILAGRO, F. I. & MARTINEZ, J. A. 2013a. Transcriptomic
and epigenetic changes in early liver steatosis associated to obesity: effect of
dietary methyl donor supplementation. *Mol Genet Metab,* 110**,** 388-95.
CORDERO, P., GOMEZ-URIZ, A. M., CAMPION, J., MILAGRO, F. I. & MARTINEZ, J. A.
2013b. Dietary supplementation with methyl donors reduces fatty liver and
modifies the fatty acid synthase DNA methylation profile in rats fed an obesogenic
diet. *Genes Nutr,* 8**,** 105-13.
CREAN, A. J., AFRIN, S., NIRANJAN, H., PULPITEL, T. J., AHMAD, G., SENIOR, A. M.,
FREIRE, T., MACKAY, F., NOBREGA, M. A., BARRES, R., SIMPSON, S. J. & PINI, T.

2023. Male reproductive traits are differentially affected by dietary macronutrient balance but unrelated to adiposity. *Nat Commun,* 14**,** 2566.

CUI, L., NIE, X., GUO, Y., REN, P., GUO, Y., WANG, X., LI, R., HOTALING, J. M., CAIRNS, B. R. & GUO, J. 2025. Single-cell transcriptomic atlas of the human testis across the reproductive lifespan. *Nat Aging.*

CVITIC, S., LONGTINE, M. S., HACKL, H., WAGNER, K., NELSON, M. D., DESOYE, G. & HIDEN, U. 2013. The human placental sexome differs between trophoblast epithelium and villous vessel endothelium. *PLoS One,* 8**,** e79233.

DE LA GARZA, A. L., MARTINEZ-TAMEZ, A. M., MELLADO-NEGRETE, A., ARJONILLA-BECERRA, S., PENA-VAZQUEZ, G. I., MARIN-OBISPO, L. M. & HERNANDEZ-BRENES, C. 2022. Characterization of the Cafeteria Diet as Simulation of the Human Western Diet and Its Impact on the Lipidomic Profile and Gut Microbiota in Obese Rats. *Nutrients,* 15.

DEARDEN, L., BOURET, S. G. & OZANNE, S. E. 2018. Sex and gender differences in developmental programming of metabolism. *Mol Metab,* 15**,** 8-19.

FAN, Y., LIU, Y., XUE, K., GU, G., FAN, W., XU, Y. & DING, Z. 2015. Diet-induced obesity in male C57BL/6 mice decreases fertility as a consequence of disrupted blood-testis barrier. *PLoS One,* 10**,** e0120775.

FLEMING, T. P., WATKINS, A. J., VELAZQUEZ, M. A., MATHERS, J. C., PRENTICE, A. M., STEPHENSON, J., BARKER, M., SAFFERY, R., YAJNIK, C. S., ECKERT, J. J., HANSON, M. A., FORRESTER, T., GLUCKMAN, P. D. & GODFREY, K. M. 2018. Origins of lifetime health around the time of conception: causes and consequences. *Lancet,* 391**,** 1842-1852.

FUJINOKI, M. 2011. Serotonin-enhanced hyperactivation of hamster sperm. *Reproduction,* 142**,** 255-66.

FURSE, S., MORGAN, H. L., KOULMAN, A. & WATKINS, A. J. 2023. Characterisation of the Paternal Influence on Intergenerational Offspring Cardiac and Brain Lipid Homeostasis in Mice. *Int J Mol Sci,* 24.

GAO, Y., ZHOU, Y., HONG, Z., MA, B., WANG, X., NIE, L., MA, L., ZHANG, Y., ZHANG, M. & WANG, M. 2026. The intricate dance of RNA-binding proteins: unveiling the mechanisms behind male infertility. *Hum Reprod Update,* 32**,** 58-104.

GHOSH, S. & MUKHERJEE, S. 2018. Testicular germ cell apoptosis and sperm defects in mice upon long-term high fat diet feeding. *J Cell Physiol,* 233**,** 6896-6909.

GOMES, A. C., HOFFMANN, C. & MOTA, J. F. 2020. Gut microbiota is associated with adiposity markers and probiotics may impact specific genera. *Eur J Nutr,* 59**,** 1751-1762.

GONZALEZ, T. L., SUN, T., KOEPPEL, A. F., LEE, B., WANG, E. T., FARBER, C. R., RICH, S. S., SUNDHEIMER, L. W., BUTTLE, R. A., CHEN, Y. I., ROTTER, J. I., TURNER, S. D., WILLIAMS, J., 3RD, GOODARZI, M. O. & PISARSKA, M. D. 2018. Sex differences in the late first trimester human placenta transcriptome. *Biol Sex Differ,* 9**,** 4.

HAMMOUD, S. S., NIX, D. A., HAMMOUD, A. O., GIBSON, M., CAIRNS, B. R. & CARRELL, D. T. 2011. Genome-wide analysis identifies changes in histone retention and epigenetic modifications at developmental and imprinted gene loci in the sperm of infertile men. *Hum Reprod,* 26**,** 2558-69.

HAO, Y., FENG, Y., YAN, X., CHEN, L., ZHONG, R., TANG, X., SHEN, W., SUN, Q., SUN, Z., REN, Y., ZHANG, H. & ZHAO, Y. 2022. Gut microbiota-testis axis: FMT improves systemic and testicular micro-environment to increase semen quality in type 1 diabetes. *Mol Med,* 28**,** 45.

HOWARD, V., REED, M. G. & NETLIBRARY, I. 2005. *Unbiased stereology : three-dimensional measurement in microscopy,* New York, Garland Science/BIOS Scientific Publishers.

HU, Y., XU, J., SHENG, Y., LIU, J., LI, H., GUO, M., XU, W., LUO, Y., HUANG, K. & HE, X. 2022. Pleurotus Ostreatus Ameliorates Obesity by Modulating the Gut Microbiota in Obese Mice Induced by High-Fat Diet. *Nutrients,* 14.

JAWAID, A., JEHLE, K. L. & MANSUY, I. M. 2021. Impact of Parental Exposure on Offspring Health in Humans. *Trends Genet,* 37**,** 373-388.

JAZWIEC, P. A., PATTERSON, V. S., RIBEIRO, T. A., YEO, E., KENNEDY, K. M., MATHIAS, P. C. F., PETRIK, J. J. & SLOBODA, D. M. 2022. Paternal obesity induces placental hypoxia and sex-specific impairments in placental vascularization and offspring metabolismdagger. *Biol Reprod,* 107**,** 574-589.

KIM, D., PAGGI, J. M., PARK, C., BENNETT, C. & SALZBERG, S. L. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol,* 37**,** 907-915.

KOVAKA, S., ZIMIN, A. V., PERTEA, G. M., RAZAGHI, R., SALZBERG, S. L. & PERTEA, M. 2019. Transcriptome assembly from long-read RNA-seq alignments with StringTie2. *Genome Biol,* 20**,** 278.

LAMBROT, R., XU, C., SAINT-PHAR, S., CHOUNTALOS, G., COHEN, T., PAQUET, M., SUDERMAN, M., HALLETT, M. & KIMMINS, S. 2013. Low paternal dietary folate alters the mouse sperm epigenome and is associated with negative pregnancy outcomes. *Nat Commun,* 4**,** 2889.

LEISEGANG, K., UDODONG, A., BOUIC, P. J. & HENKEL, R. R. 2014. Effect of the metabolic syndrome on male reproductive function: a case-controlled pilot study. *Andrologia,* 46**,** 167-76.

LI, Y., WANG, Y., TAN, Y. Q., YUE, Q., GUO, Y., YAN, R., MENG, L., ZHAI, H., TONG, L., YUAN, Z., LI, W., WANG, C., HAN, S., REN, S., YAN, Y., WANG, W., GAO, L., TAN, C., HU, T., ZHANG, H., LIU, L., YANG, P., JIANG, W., YE, Y., TAN, H., WANG, Y., LU, C., LI, X., XIE, J., YUAN, G., CUI, Y., SHEN, B., WANG, C., GUAN, Y., LI, W., SHI, Q., LIN, G., NI, T., SUN, Z., YE, L., VOUREKAS, A., GUO, X., LIN, M. & ZHENG, K. 2024. The landscape of RNA binding proteins in mammalian spermatogenesis. *Science,* 386**,** eadj8172.

LIAO, Y., WANG, J., JAEHNIG, E. J., SHI, Z. & ZHANG, B. 2019. WebGestalt 2019: gene set analysis toolkit with revamped UIs and APIs. *Nucleic Acids Res,* 47**,** W199-W205.

LIU, S., QIN, P. & WANG, J. 2019. High-Fat Diet Alters the Intestinal Microbiota in Streptozotocin-Induced Type 2 Diabetic Mice. *Microorganisms,* 7.

LIU, Y., ZHAO, W., GU, G., LU, L., FENG, J., GUO, Q. & DING, Z. 2014. Palmitoyl-protein thioesterase 1 (PPT1): an obesity-induced rat testicular marker of reduced fertility. *Mol Reprod Dev,* 81**,** 55-65.

LOTTI, F., CORONA, G., DEGLI INNOCENTI, S., FILIMBERTI, E., SCOGNAMIGLIO, V., VIGNOZZI, L., FORTI, G. & MAGGI, M. 2013. Seminal, ultrasound and psychobiological parameters correlate with metabolic syndrome in male members of infertile couples. *Andrology,* 1**,** 229-39.

LUDWIG, D. S., PETERSON, K. E. & GORTMAKER, S. L. 2001. Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet,* 357**,** 505-8.

LUISI, S., FLORIO, P., REIS, F. M. & PETRAGLIA, F. 2005. Inhibins in female and male reproductive physiology: role in gametogenesis, conception, implantation and early pregnancy. *Hum Reprod Update,* 11**,** 123-35.

LV, S., HUANG, J., LUO, Y., WEN, Y., CHEN, B., QIU, H., CHEN, H., YUE, T., HE, L., FENG, B., YU, Z., ZHAO, M., YANG, Q., HE, M., XIAO, W., ZOU, X., GU, C. & LU, R. 2024. Gut microbiota is involved in male reproductive function: a review. *Front Microbiol,* 15**,** 1371667.

MBOGNING, J., NAGY, S., PAGE, V., SCHWER, B., SHUMAN, S., FISHER, R. P. & TANNY, J. C. 2013. The PAF complex and Prf1/Rtf1 delineate distinct Cdk9-dependent pathways regulating transcription elongation in fission yeast. *PLoS Genet,* 9**,** e1004029.

MCPHERSON, N. O., FULLSTON, T., KANG, W. X., SANDEMAN, L. Y., CORBETT, M. A., OWENS, J. A. & LANE, M. 2016. Paternal under-nutrition programs metabolic syndrome in offspring which can be reversed by antioxidant/vitamin food fortification in fathers. *Sci Rep,* 6**,** 27010.

MORGAN, H. L., ALJUMAH, A., ROUILLON, C. & WATKINS, A. J. 2021. Paternal low protein diet and the supplementation of methyl-donors impact fetal growth and placental development in mice. *Placenta,* 103**,** 124-133.

MORGAN, H. L., AMPONG, I., EID, N., ROUILLON, C., GRIFFITHS, H. R. & WATKINS, A. J. 2020a. Low protein diet and methyl-donor supplements modify testicular physiology in mice. *Reproduction,* 159**,** 627-641.

MORGAN, H. L., EID, N., HOLMES, N., HENSON, S., WRIGHT, V., COVENEY, C., WINDER, C., O'NEIL, D. M., DUNN, W. B., BOOCOCK, D. J. & WATKINS, A. J. 2024. Paternal undernutrition and overnutrition modify semen composition and preimplantation embryo developmental kinetics in mice. *BMC Biol,* 22**,** 207.

MORGAN, H. L., FURSE, S., DIAS, I. H. K., SHABIR, K., CASTELLANOS, M., KHAN, I., MAY, S. T., HOLMES, N., CARLILE, M., SANG, F., WRIGHT, V., KOULMAN, A. & WATKINS, A. J. 2022. Paternal low protein diet perturbs inter-generational metabolic homeostasis in a tissue-specific manner in mice. *Commun Biol,* 5**,** 929.

MORGAN, H. L., PAGANOPOULOU, P., AKHTAR, S., URQUHART, N., PHILOMIN, R., DICKINSON, Y. & WATKINS, A. J. 2020b. Paternal diet impairs F1 and F2 offspring vascular function through sperm and seminal plasma specific mechanisms in mice. *J Physiol,* 598**,** 699-715.

MOSS, J. L. & HARRIS, K. M. 2015. Impact of maternal and paternal preconception health on birth outcomes using prospective couples' data in Add Health. *Arch Gynecol Obstet,* 291**,** 287-98.

NG, S. F., LIN, R. C., LAYBUTT, D. R., BARRES, R., OWENS, J. A. & MORRIS, M. J. 2010. Chronic high-fat diet in fathers programs beta-cell dysfunction in female rat offspring. *Nature,* 467**,** 963-6.

NIELSEN, J. E., LINDEGAARD, M. L., FRIIS-HANSEN, L., ALMSTRUP, K., LEFFERS, H., NIELSEN, L. B. & RAJPERT-DE MEYTS, E. 2010. Lipoprotein lipase and endothelial lipase in human testis and in germ cell neoplasms. *Int J Androl,* 33**,** e207-15.

O'NEILL, S. & O'DRISCOLL, L. 2015. Metabolic syndrome: a closer look at the growing epidemic and its associated pathologies. *Obes Rev,* 16**,** 1-12.

OMOTE, M., WAKIMOTO, Y. & SHIBAHARA, H. 2023. Possible Role of 5-Hydroxytryptamine (5-HT) Receptor on Human Sperm Motility Regulation. *Cureus,* 15**,** e49530.

OSEI-KUMAH, A., SMITH, R., JURISICA, I., CANIGGIA, I. & CLIFTON, V. L. 2011. Sex-specific differences in placental global gene expression in pregnancies complicated by asthma. *Placenta,* 32**,** 570-8.

OZANNE, S. E., FERNANDEZ-TWINN, D. & HALES, C. N. 2004. Fetal growth and adult diseases. *Semin Perinatol,* 28**,** 81-7.

PENESOVA, A., BUNT, J. C., BOGARDUS, C. & KRAKOFF, J. 2010. Effect of paternal diabetes on pre-diabetic phenotypes in adult offspring. *Diabetes Care,* 33**,** 1823-8.

PEPIN, A. S., LAFLEUR, C., LAMBROT, R., DUMEAUX, V. & KIMMINS, S. 2022. Sperm histone H3 lysine 4 tri-methylation serves as a metabolic sensor of paternal obesity and is associated with the inheritance of metabolic dysfunction. *Mol Metab,* 59**,** 101463.

PETROPOULOS, S., EDSGARD, D., REINIUS, B., DENG, Q., PANULA, S. P., CODELUPPI, S., PLAZA REYES, A., LINNARSSON, S., SANDBERG, R. & LANNER, F. 2016. Single-Cell RNA-Seq Reveals Lineage and X Chromosome Dynamics in Human Preimplantation Embryos. *Cell,* 165**,** 1012-26.

POKHOLOK, D. K., HANNETT, N. M. & YOUNG, R. A. 2002. Exchange of RNA polymerase II initiation and elongation factors during gene expression in vivo. *Mol Cell,* 9**,** 799-809.

RIBEIRO, A. A., DE OLIVEIRA LACERDA, P. M., PEREIRA DE MELO, M., DE CARVALHO BALIEIRO, J. C. & DE SOUZA, R. R. 2008. Placental microstructure and efficiency in cloned bovines: a design-based stereological approach. *Cell Tissue Res,* 333**,** 105-14.

ROSENFELD, C. S. 2015. Sex-Specific Placental Responses in Fetal Development. *Endocrinology,* 156**,** 3422-34.

SAKLAYEN, M. G. 2018. The Global Epidemic of the Metabolic Syndrome. *Curr Hypertens Rep,* 20**,** 12.

SANCHEZ-GARRIDO, M. A., RUIZ-PINO, F., VELASCO, I., BARROSO, A., FERNANDOIS, D., HERAS, V., MANFREDI-LOZANO, M., VAZQUEZ, M. J., CASTELLANO, J. M.,

ROA, J., PINILLA, L. & TENA-SEMPERE, M. 2018. Intergenerational Influence of
Paternal Obesity on Metabolic and Reproductive Health Parameters of the
Offspring: Male-Preferential Impact and Involvement of Kiss1-Mediated Pathways.
Endocrinology, 159**,** 1005-1018.
SEDLMEIER, E. M., BRUNNER, S., MUCH, D., PAGEL, P., ULBRICH, S. E., MEYER, H. H.,
AMANN-GASSNER, U., HAUNER, H. & BADER, B. L. 2014. Human placental
transcriptome shows sexually dimorphic gene expression and responsiveness to
maternal dietary n-3 long-chain polyunsaturated fatty acid intervention during
pregnancy. *BMC Genomics,* 15**,** 941.
SERINO, M., LUCHE, E., GRES, S., BAYLAC, A., BERGE, M., CENAC, C., WAGET, A.,
KLOPP, P., IACOVONI, J., KLOPP, C., MARIETTE, J., BOUCHEZ, O., LLUCH, J.,
OUARNE, F., MONSAN, P., VALET, P., ROQUES, C., AMAR, J., BOULOUMIE, A.,
THEODOROU, V. & BURCELIN, R. 2012. Metabolic adaptation to a high-fat diet is
associated with a change in the gut microbiota. *Gut,* 61**,** 543-53.
SIKLENKA, K., ERKEK, S., GODMANN, M., LAMBROT, R., MCGRAW, S., LAFLEUR, C.,
COHEN, T., XIA, J., SUDERMAN, M., HALLETT, M., TRASLER, J., PETERS, A. H. &
KIMMINS, S. 2015. Disruption of histone methylation in developing sperm impairs
offspring health transgenerationally. *Science,* 350**,** aab2006.
SILVA, D. R., WERNECK, A. O., COLLINGS, P. J., FERNANDES, R. A., BARBOSA, D. S.,
RONQUE, E. R. V., SARDINHA, L. B. & CYRINO, E. S. 2017. Family history of
cardiovascular disease and parental lifestyle behaviors are associated with
offspring cardiovascular disease risk markers in childhood. *Am J Hum Biol,* 29.
SINGH, S. P., MISRA, B., KAR, S. K., PANIGRAHI, M. K., MISRA, D., BHUYAN, P.,
PATTNAIK, K., MEHER, C., AGRAWAL, O., ROUT, N. & SWAIN, M. 2015.
Nonalcoholic fatty liver disease (NAFLD) without insulin resistance: Is it different?
Clin Res Hepatol Gastroenterol, 39**,** 482-8.
SOOD, R., ZEHNDER, J. L., DRUZIN, M. L. & BROWN, P. O. 2006. Gene expression
patterns in human placenta. *Proc Natl Acad Sci U S A,* 103**,** 5478-83.
SQUAZZO, S. L., COSTA, P. J., LINDSTROM, D. L., KUMER, K. E., SIMIC, R., JENNINGS,
J. L., LINK, A. J., ARNDT, K. M. & HARTZOG, G. A. 2002. The Paf1 complex
physically and functionally associates with transcription elongation factors in vivo.
EMBO J, 21**,** 1764-74.
STERN, C., SCHWARZ, S., MOSER, G., CVITIC, S., JANTSCHER-KRENN, E., GAUSTER, M.
& HIDEN, U. 2021. Placental Endocrine Activity: Adaptation and Disruption of
Maternal Glucose Metabolism in Pregnancy and the Influence of Fetal Sex. *Int J*
Mol Sci, 22.
SUGIYAMA, Y., FUJINOKI, M. & SHIBAHARA, H. 2019. Effects of 5-hydroxytryptamine on
spermatozoal hyperactivation and in vitro fertilization in mice. *J Reprod Dev,* 65**,**
541-550.
VERAS, M. M., DAMACENO-RODRIGUES, N. R., CALDINI, E. G., MACIEL RIBEIRO, A. A.,
MAYHEW, T. M., SALDIVA, P. H. & DOLHNIKOFF, M. 2008. Particulate urban air
pollution affects the functional morphology of mouse placenta. *Biol Reprod,* 79**,**
578-84.
VON FRANKENBERG, A. D., MARINA, A., SONG, X., CALLAHAN, H. S., KRATZ, M. &
UTZSCHNEIDER, K. M. 2017. A high-fat, high-saturated fat diet decreases insulin
sensitivity without changing intra-abdominal fat in weight-stable overweight and
obese adults. *Eur J Nutr,* 56**,** 431-443.
WATKINS, A. J., DIAS, I., TSURO, H., ALLEN, D., EMES, R. D., MORETON, J., WILSON,
R., INGRAM, R. J. M. & SINCLAIR, K. D. 2018. Paternal diet programs offspring
health through sperm- and seminal plasma-specific pathways in mice. *Proceedings*
of the National Academy of Sciences of the United States of America, 115**,** 10064-
10069.
WATKINS, A. J., LUCAS, E. S., MARFY-SMITH, S., BATES, N., KIMBER, S. J. & FLEMING,
T. P. 2015. Maternal nutrition modifies trophoblast giant cell phenotype and fetal
growth in mice. *Reproduction,* 149**,** 563-75.
WATKINS, A. J., SIROVICA, S., STOKES, B., ISAACS, M., ADDISON, O. & MARTIN, R. A.
2017. Paternal low protein diet programs preimplantation embryo gene

expression, fetal growth and skeletal development in mice. *Biochimica Et*
Biophysica Acta-Molecular Basis of Disease, 1863**,** 1371-1381.
WILKINSON, A. L., MENELAOU, K., RAKOCZY, J., TAN, X. S. & WATSON, E. D. 2021.
Disruption of Folate Metabolism Causes Poor Alignment and Spacing of Mouse
Conceptuses for Multiple Generations. *Front Cell Dev Biol,* 9**,** 723978.
YANG, D., DAI, F., YUAN, M., ZHENG, Y., LIU, S., DENG, Z., TAN, W., CHEN, L., ZHANG,
Q., ZHAO, X. & CHENG, Y. 2021. Role of Transforming Growth Factor-beta1 in
Regulating Fetal-Maternal Immune Tolerance in Normal and Pathological
Pregnancy. *Front Immunol,* 12**,** 689181.
YI, C., KITAMURA, Y., MAEZAWA, S., NAMEKAWA, S. H. & CAIRNS, B. R. 2025.
ZBTB16/PLZF regulates juvenile spermatogonial stem cell development through
an extensive transcription factor poising network. *Nat Struct Mol Biol,* 32**,** 1213-
1226.
ZHANG, P., FENG, Y., LI, L., GE, W., YU, S., HAO, Y., SHEN, W., HAN, X., MA, D., YIN,
S., TIAN, Y., MIN, L., SUN, Z., SUN, Q., ZHANG, H. & ZHAO, Y. 2021.
Improvement in sperm quality and spermatogenesis following faecal microbiota
transplantation from alginate oligosaccharide dosed mice. *Gut,* 70**,** 222-225.

1047

1048

 Figure legends

 Figure 1. Impact of diet on male growth. (A) Mean weekly body weight of males fed either

 a control diet (CD), low protein diet (LPD), methyl donor-supplemented LPD (MD-LPD),

 Western diet (WD) or methyl donor-supplemented Western Diet (MD-WD). Mean (B)

 testicular weight, (C) seminal vesicle weight (D) total fat weight (combined weight of

 individual fat pads) and (E) ratio of white adipose tissue (WAT) to brown adipose tissue

 (BAT). Mean serum (F) Tnf and (G) inhibin β-A chain levels. N = 6-8 males in each group.

 Data were analysed using a one-way ANOVA with Holm-Sidak post hoc tests for multiple

 comparison. * P <0.05, ** P <0.01.

 Figure 2. Impact of diet on male non-fasting metabolic status. Serum (A) glucose and (B)

 insulin in males fed either a control diet (CD), low protein diet (LPD), methyl donor-

 supplemented LPD diet (MD-LPD), Western diet (WD) or methyl donor-supplemented WD

 (MD-WD). Hepatic (C) cholesterol, (D) free fatty acids (FFAs) and (E) triglyceride (TG)

 concentrations. N = 8 males in each group. Data were analysed using either a one-way

 ANOVA with Holm-Sidak post hoc tests for multiple comparison respectively. * P <0.05, **

 P <0.01.

 Figure 3. Impact of diet on male non-fasting metabolic status and gut microbiota. (A)

 Faith's phylogenetic diversity and (B) Pielou's measure of species evenness. (C) Overall

 bacterial abundance at the phylum level and relative abundance of (D) Deferribacteres, (E)

 Protobacteria and (F) TM7 phylum. (G) Relative (Z-score) abundance of bacteria at the

 family level in males fed either a control diet (CD), low protein diet (LPD), methyl donor-

 supplemented LPD diet (MD-LPD), Western diet (WD) or methyl donor-supplemented WD

 (MD-WD). N = 8 males in each group. Data were analysed using either a one-way ANOVA

1074 (panels D, E, F and G) or Kruskal–Wallis test (panels A, B and C) with Holm-Sidak or

1075 Dunn's post hoc tests for multiple comparison respectively. * P <0.05, ** P <0.01.

1076

Figure 4. Impact of diet on male testicular morphology. (A) Representative images

showing tubule morphology in males fed either a control diet (CD), low protein diet (LPD),

methyl donor-supplemented LPD (MD-LPD), Western diet (WD) or methyl donor-

supplemented WD (MD-WD). (B) Examples of tubule anomalies identified in WD and MD-

WD males including separation of the epithelium from the tubule basement membrane (as

indicated by an arrow in i), appearance of vacuoles (as indicated by an arrow in ii), and loss of

the germinal epithelium (as indicated by arrows in iii and iv). (C) Frequency of abnormal

tubules per testis in males fed either CD, LPD, MD-LPD, WD or MD-WD. (D)

Representative staining patterns for DAPI (nuclear counterstain), Sox9 (marker of Sertoli

cells), Ddx4 (marker of spermatocytes and spermatids) and Plzf (marker of spermatogonial

stem cells) in testes from males fed CD, LPD, MD-LPD, WD and MD-WD. (E) Number of

DAPI, (F) Sox9, (G) Ddx4 and (H) Plzf$^+$ cells. N = 7-8 males in each group. Data were

analysed using either a one-way ANOVA (panels E, F, G and H) or Kruskal–Wallis test

(panel C) with Holm-Sidak or Dunn's post hoc tests for multiple comparison respectively. * P

<0.05, ** P< 0.01.

1092

Figure 5. Impact of diet on testicular gene expression. Volcano plots comparing

differential gene expression between males fed a control diet (CD) and males fed (A) a low

protein diet (LPD), (B) methyl donor-supplemented LPD (MD-LPD) or (C) Western diet

(WD). (D) Pathway and (E) network analysis of differentially expressed genes between CD

and WD fed males. (F) Volcano plots comparing differential gene expression between males

fed a control diet (CD) and males fed a methyl donor-supplemented WD (MD-WD). (G)

Pathway and (H) network analysis of differentially expressed genes between CD and MD-

WD fed males. N = 8 males in each group.

1101

Figure 6. Impact of paternal diet on early (E8.5) placental development. (A) Litter size at

E8.5 from males fed either a control diet (CD), low protein diet (LPD), methyl donor-

supplemented LPD (MD-LPD), Western diet (WD) or methyl-donor supplemented WD (MD-

WD) prior to mating. (B) Representative images of H&E stained whole E8.5 conceptuses.

Ectoplacental cone (EPC) (C) invasion dept, (D) area and (I) alignment. (F) Representative

EPC outgrowth after 48 hours in culture, stained for alpha-tubulin. (G) Trophoblast and (H)

EPC area at 24 and 48 hour timepoints. N = 10-13 litters in A, derived from a minimum of 8

separate stud males per group, and 4-9 conceptuses in C-H, each from a separate litter and

stud male. Data were analysed using either a one-way ANOVA or Kruskal–Wallis test with

Holm-Sidak or Dunn's post hoc tests for multiple comparison respectively. * P <0.05.

1112

Figure 7. Impact of paternal diet on fetal and placental weight in late (E17.5) gestation.

(A) Late gestation fetal weight and distributions above the 90^{th} centile and below the 10^{th}

centile of control diet (CD) fetal weights from males fed either a CD, low protein diet (LPD),

methyl donor-supplemented LPD (MD-LPD), Western diet (WD) or methyl donor-

supplemented WD (MD-WD) prior to mating. Late gestation (B) female, (C) male and (D)

intra-diet comparison of fetal weight. (E) Late gestation placental weight and distributions

above the 90^{th} centile and below the 10^{th} centile of CD placental weights. Late gestation (F)

female, (G) male and (H) intra-diet comparison of placental weight. (I) Late gestation

fetal:placental ratio and distributions above the 90^{th} centile and below the 10^{th} centile of CD

fetal:placental ratio. Late gestation (G) female, (H) male and (I) intra-diet comparison of

fetal:placental weight ratio. N = 10-13 litters, derived from a minimum of 8 separate stud

males per group. Data are presented as mean ± SEM or as box plots with the mean and

individual data points outside of the 10-90[th] percentile. Data were analysed using a

generalised linear mixed model analysis with paternal origin of litter and duration on

respective diet incorporated as random effects (panels A, B, C, E, F, G, I, J and K) or t-test

(panels D, H and L) following assessment for normality using a Shapiro-Wilk test. * P <0.05,

** P <0.01, ***P <0.001.

1130

Figure 8. Analysis of late gestation (E17.5) placental morphology in response to paternal

diet. (A) Representative image of a cross section through the late gestation (E17.5) placenta

highlighting the decidua, junctional zone, labyrinth zone and chorionic plate regions. (B)

Cross section area and (C) relative regional composition of placentas derived from males fed

either a control diet (CD), low protein diet (LPD), methyl donor-supplemented LPD (MD-

LPD), Western diet (WD) or methyl-donor supplemented WD (MD-WD) prior to mating. (D)

Representative images of a E17.5 placenta at 20X magnification prior to the application of (E)

estimation of compartment global volumes using Cavalieri's Principle with a grid comprising

a test system of points and (F) stereological method for estimating the surface density of

maternal blood vessels and fetal capillaries using a grid comprising isotropic lines. E17.5

placental (G) volume, (H) labyrinth zone volume, (I) junctional zone volume, (J) maternal

blood space volume, (K) maternal blood surface area and (L) fetal capillary surface area. N =

5 placentas per group, each from a separate litters and stud males. Data are presented as mean

± SEM and were analysed using a one-way ANOVA (panels B, G-L) or Kruskal–Wallis test

(panel B) with post-hoc correction where appropriate.

1146

Figure 9. Late gestation (E17.5) placental gene expression is altered in a diet and sex-

specific manner. (A) Differential gene expression and (B) pathway analysis from male

placentas derived from males fed either a control diet (CD), low protein diet (LPD), methyl

donor-supplemented LPD (MD-LPD), Western diet (WD) or methyl-donor supplemented WD

(MD-WD) prior to mating. (C) Differential gene expression and (D) pathway analysis from

female placentas. (E) Principal component analysis comparing CD male and female placental

gene expression profiles. (F) Heat map of the 301 sexually dimorphic genes between male

and female CD placentas and (G) the pathway analysis. Heatmaps showing the relative level

of expression and clustering of the same 301 genes in (H) LPD, (I) MD-LPD, (J) WD and (K)

MD-WD male and female placentas. Volcano plots highlighting the significantly (pAdj

<0.05) upregulated (red) and downregulated (blue) differentially expressed genes between

male and female (L) LPD, (M) MD-LPD, (N) WD and (O) MD-WD placentas. (P) Overlap of

sexually dimorphic genes between male and female CD, LPD, WD and MD-WD placentas.

(Q) Chromosomal locations of common X and Y chromosome sexually dimorphic genes

between CD, LPD, WD and MD-WD placentas. N = 4 male and 4 female placentas, each

from separate litters and stud males.

1163

1164   **Supplemental Figure 1:** Representative (A) no primary and (B) no secondary negative

1165   controls of testicular sections sained for Plzf and Ddx4. Representative (C) no primary and

1166   (D) no secondary negative controls of testicular sections sained for Sox9.