1 Title: Genome assembly and diagnostic DNA markers for sex of the largest freshwater fish in
2 North America, the white sturgeon (*Acipenser transmontanus*).
3

4 Authors: Stuart C. Willis, Jeremiah Smith, Shawn R. Narum
5
6 Addresses
7
8 SCW, SRN: Hagerman Genetics Lab, Columbia River Inter-Tribal Fish Commission, 3059F
9 National Fish Hatchery Rd, Hagerman, Idaho, 83332, USA
10
11 JS: Department of Biology, University of Kentucky, 195 Huguelet Dr, Lexington, KY 40508, USA
12
13 Corresponding author's email: swillis@critfc.org

14
15 Running title: Genome assembly of white sturgeon
16

17 Keywords: polyploid, autopolyploidy, multivalent

Abstract

Sturgeon and paddlefish represent some of the most early diverging branches of ray-finned fishes and have undergone at least one global and several lineage-specific whole genome duplication events. White sturgeon (*Acipenser transmontanus*), the largest freshwater fish in North America, have experienced at least two rounds of whole genome duplication, and may exhibit both di- and multi-valent meiotic segregation. Moreover, they exhibit contemporary ploidy variants due to spontaneous autopolyploidy, particularly in aquaculture. Nonetheless, as a species with several population segments that exhibit chronic recruitment failure, conservation aquaculture is an important part of their management. To facilitate the development of genetic tools to aid white sturgeon conservation, as well as a basis to understand how ploidy changes and variation historically and contemporarily shape the evolution of this species, we present a genome assembly for a white sturgeon from the Snake River, Idaho, USA. Analysis of sequence data used for assembly indicated a haploid genome size of approximately 1.5Gbp, implying tetraploidy (4N), while analysis of heterozygous k-mers from 21 to 41 bp suggest the genome reflects both 4N and 8N variants. The final genome assembly, scaffolded using linkage maps constructed from Fraser River and Snake River F1 families, contained 6.26Gbp in 832,145 scaffolds, consistent with published genome size estimates. Conserved ortholog completeness for this genome (90.5%; 22.8% single-copy and 67.7% duplicated) was similar to the putatively diploid sterlet sturgeon, and the largest linkage map-based scaffold was 55.2Gbp, though the N50 for this assembly was only 416Kbp, indicating the assembly remains fragmented. We demonstrate the utility of this assembly by identifying genomic regions significantly associated with sex. Genetic markers, designed for inclusion in an amplicon genotyping panel, predicted sex 96.6% and 81.5% correctly in females and males, respectively, providing a strong overall association (χ^2 p-value $< 2.7 \times 10^{-37}$) with some variation by geographic region.

Article summary:

Functional polysomes (more than two chromosomes that pair in meiosis) are rare among vertebrates. White sturgeon, the largest freshwater fish in North America, are tetraploid and occasionally hexaploid. Several populations of this species are stagnant or declining, requiring aquaculture to bolster reproduction. We analyzed whole genomic data and assembled the genome of an individual from the unique Snake River, Idaho, population. Results indicate most genetic variants are consistent with tetraploidy, and the genome assembly, while fragmented, is largely complete. We demonstrate its utility by designing genetic markers for sex for use in conservation and commercial aquaculture.

Introduction

Sturgeons and paddlefishes, fishes in the order Acipenseriformes, are often referred to as 'living fossils'. While somewhat misleading, this moniker reflects that they were an early-diverging branch within the Actinopterygii or clade of 'ray-finned' fishes, have exhibited conservative morphological change relative to ancient acipenseriform fossils, and unlike most other ray-finned fishes, exhibit only a weakly ossified, cartilaginous skeleton (Peng et al., 2007). In contrast, relative to the estimated ancestral Actinoperygiian karyotype, sturgeon and paddlefish have undergone several whole genome duplications (Peng et al., 2007), and while many species appear to have made considerable if not complete evolution back to functional diploidy, some remain polyploid and potentially multivalent across much of the genome (Fontana, 2002). Moreover, in some sturgeon species, spontaneous autopolyploidy results in ploidy variation within a single population, although the origin and long-term demographic importance of these variants is not well-understood (Drauch Schreier et al., 2011). Nonetheless, these observations make sturgeons some of the few contemporarily polyploid, fully sexually reproducing vertebrate species (Jighly et al., 2018; Spoelhof et al., 2017).

Sturgeon and paddlefishes are often known for their size and unique, even bizarre morphology, which makes them a conspicuous target for commercial and recreational fisheries, as well as for being the source of roe for caviar. Most contemporary caviar and sturgeon meat, however, comes from commercial aquaculture operations, since most natural populations were overharvested if not threatened or endangered by the mid-twentieth century, and at least one acipenseriform, the Chinese paddlefish, was declared extinct in the wild in 2019 (Lobanov et al., 2024). Indeed, the 'periodic' life history strategy of these fishes makes them extremely vulnerable to overfishing (Kopf et al., 2025; Winemiller, 2005). They exhibit individuals that mature after a decade or more, experience very low rates of natural mortality with adults that live for decades, if not more than a century, and have high fecundity. Notably, however, survivorship among early life stages is low and depends on environmental conditions that may occur unpredictably in time and space, including flow rates and substrate composition (Parsley et al., 1993; Parsley & Beckman, 1994). In addition to overharvest, anthropogenic alteration of natural flow patterns and habitat fragmentation have resulted in species with reduced population sizes that now must also contend with diminished or failed recruitment (Mallette, 2014). As such, there are numerous population management programs for sturgeons and paddlefish across their range, including the development of conservation aquaculture programs to bolster threatened populations (Korman et al., 2025). The success of these aquaculture programs, however, depends on the efficient production of hatchery progeny which reflect the genetic diversity and sex-ratios of natural-origin populations, an endeavor which can be greatly aided by molecular markers and especially markers for sex, notwithstanding the ancestrally and possibly functionally polyploid nature of the genomes.

The white sturgeon (*Acipenser transmontanus*) typifies all of the challenges exhibited by acipenseriform fishes. White sturgeon are the largest 'freshwater' fish in North America, historically reaching lengths over 4m. Although they are obligate freshwater spawners, white sturgeon are considered to be quasi-amphidromous and range along the Pacific coast from Baja California to Alaska (Hildebrand et al., 2016). Despite their extensive range, there are only three known self-sustaining populations, in the Sacramento-San Joaquin, Fraser, and Columbia River

Basins, with the latter estimated to be the largest (Hildebrand et al., 2016), despite the strong
segmentation of the basin by hydropower facilities. Within the Columbia Basin, as elsewhere,
white sturgeon populations were severely overharvested into the twentieth century and, despite
modern protections, are imperiled by limited recruitment in many areas.
White sturgeon have a karyotype of ~270 chromosomes (Van Eenennaam et al., 1998),
although there is considerable uncertainty of their genome size (C-value), with estimates
ranging from 5.1 to 9.6pg (Blacklidge & Bidwell, 1993; Hinegardner, 1976). Relative to the
Actinopterygiian median (2N ~48), this karyotype indicates considerable genome expansion
within Acipenseriformes, though contemporary congeners with ~120 chromosomes are
considered functional diploids (Fontana, 2002). Indeed evidence for white sturgeon suggests
they exhibit tetravalent or bivalent segregation and inheritance patterns (Delomas et al., 2021;
Van Eenennaam et al., 1998), though the proportions of each across the genome are difficult to
estimate, and for simplicity they are considered tetraploid (4N) relative to sturgeon species with
~120 chromosomes (Fontana et al., 2008; Gille et al., 2015). Moreover, spontaneous
autopolyploidy is not uncommon in hatchery white sturgeon, and occasionally among natural-
origin fish as well, producing hexaploid offspring (6N, relative to 2N≈120), which themselves
produce viable pentaploid (5N) offspring when backcrossed (Drauch Schreier et al., 2011; Gille
et al., 2015). Notably, female white sturgeon are predicted to be the heterogametic sex (ZW sex
determination), like other sturgeons (Fopp-Bayat et al., 2007; Van Eenennaam et al., 1999).
However, if autopolyploids are produced through the retention of the second polar body of the
egg, as hypothesized (Gille et al., 2015), meaning maternal genetic contributions are doubled,
sex ratios, fertility, and perhaps viability may commonly be compromised among spontaneous
autopolyploids and backcross offspring (Wertheim et al., 2013). Despite these potential
consequences, ploidy variation clearly has long played a role in sturgeon evolution (Redmond et
al., 2023), but deciphering how rediploidization following repeated genome duplications has
resulted in both susceptibility and resilience to contemporary ploidy variation in white sturgeon
has been limited by genomic resources for it. Here, we present a draft genome assembly for
white sturgeon produced from chromatin proximity ligation, microfluidic fragment barcoding, and
linkage disequilibrium mapping. We demonstrate the utility of this genomic assembly by
identifying regions of the assembly that are homologous to sex-associated regions of other
sturgeon and testing the efficacy of molecular markers for them at predicting phenotypic sex in
white sturgeon.
132
133
134   **Methods**
135
136   **Initial genome assembly**
137
Whole blood from an adult female white sturgeon originating from the Snake River
population (a tributary to the Columbia River) near Hagerman, Idaho, USA was submitted to
NRGene (Ness Ziona, Israel) for paired-end, mate-pair (proximity-ligation) and 10X Chromium
(microfluidic fragment barcoding) sequencing (Table 1). To estimate the genome size and ploidy
status of the sequenced individual, we utilized the ≤470bp-insert, 2x250bp library with
Genomescope and Smudgeplot (Ranallo-Benavidez et al., 2020). For each analysis, kmers

were tallied with jellyfish (Marçais & Kingsford, 2011) and FastK (Myers, 2020), respectively,
with kmer lengths of 21, 23, 31, and 41 bp. Genomescope analysis was run with assumed
ploidy of 2N and 4N (maximum for this analysis is 6N).
Sequence data were utilized with the DeNovoMAGIC v3 assembler to produce a phased
(allelic) genome assembly (NRGene). Completeness of the phased assembly was assessed
using Compleasm v0.2.7 (Huang & Li, 2023) based on the Actinopertygii ortholog set from
orthoDB v12, which contains 7,207 orthologous genes. For comparison, similar evaluations of
completeness were made on the NCBI RefSeq (haploid) assembly for the sterlet sturgeon
(*Acipenser ruthenus*), a putatively diploid European sturgeon with 2N ~120 chromosomes
(GCF_902713425.1) (Du et al., 2020).
154
155 **F1 family sequencing and variant calling**
156
Linkage mapping was made from two outbred F1 families. The first used restriction-site
associated DNA sequencing (RADseq) of a mixed reduced- and normal-ploidy family of white
sturgeon originating from the Fraser River population (British Columbia, Canada) and the
second used messenger RNA sequencing (RNAseq) of a normal-ploidy family from the Snake
River population.
For the Fraser River F1 family, ultraviolet light irradiated sperm were used to fertilize
eggs from a female white sturgeon at Vancouver Island University (British Columbia, Canada).
Whole genomic DNA from somatic (fin) tissue from 226 offspring plus 2 parents was utilized in a
restriction digest with a single enzyme with a 8bp recognition sequence (SbfI) following
published protocols (Baird et al., 2008; Micheletti et al., 2018). Barcoded libraries were
sequenced on an Illumina NextSeq 500 (2x150bp). Reads were mapped to the genome
assembly using Hisat2 v2.2.1 (Kim et al., 2019) with the --no-spliced-alignment option enabled,
and SNPs were called using bcftools v1.22 (Danecek et al., 2021) jointly using all individuals
and prospective parents, then filtered using vcftools v0.1.17 (Danacek et al., 2011) to separate
diploid vs haploid mapping families and retain SNPs with minor allele frequencies greater than
0.1 and with missing genotypes in fewer than 50% of segregants within a mapping family.
Genotypes in VCF format were converted to joinmap format using the populations pipeline from
STACKS v2.68 (Catchen et al., 2011).
For the Snake River F1 family, sampling occurred at the Idaho Springs Foods sturgeon
hatchery near Hagerman, ID, USA. Dorsal fin punches were collected from the and preserved in
RNAlater. Whole bodies of 92 offspring were collected and preserved in RNAlater and stored at
-20C until RNA isolation. Larvae were ~3cm in length at the time of collection and were
euthanized through cervical dislocation. Total RNA was isolated from a mixed-tissue portion of
the head of each offspring and fin clips from the parents using the Direct-zol RNA Miniprep Plus
Kit using manufacturer protocols (Zymo Research, Irvine, CA, USA). Quality for each individual
extract was assessed utilizing High Sensitivity RNA ScreenTape Analysis (Agilent Technologies
INC, Clara, CA, USA). Individuals with RNA integrity numbers ≥ 7 were used in downstream
library preparation with a total of 92 offspring and 2 parents used. We isolated messenger RNA
(mRNA) from the total RNA using Poly(A) mRNA Magnetic Isolation Module (New England
Biolabs, MA, USA).

RNA-seq library preparations were performed using half volume reactions with the
NEBNext Ultra II RNA Library Prep Kit for Illumina (New England Biolabs, MA, USA), which
produces non-stranded cDNA fragments. We used an 8-minute mRNA fragmentation time to
ensure fragment sizes were ≥ 350bp prior to adapter ligation and used 13X cycles during PCR
enrichment. Bead cleaning was omitted prior to PCR enrichment but two or three bead cleans
were performed post-PCR to ensure dimer removal. Equimolar pools of two sets of 46 offspring
plus the two parents were created for sequencing on an Illumina NextSeq2000 using a P3 300
cycle kit with 2x150bp chemistry (two libraries of 46 unique offspring; each parent sequenced
twice). Data were quality-trimmed with BBmap (Bushnell, 2015) then mapped to the phased
assembly using HiSat2 as unstranded cDNA and retaining secondary alignments. Only properly
paired alignments were retained and used to score variants with bcftools, which assumed
diploidy. Variants were filtered for a minimum quality of 20, minimum depth of 10 reads,
maximum missing data of 10%, and minimum allele frequency of 0.1.
Parents and offspring from both F1 families were also genotyped for a panel of 291
anonymous, autosomal SNPs developed for white sturgeon (Delomas et al., 2021). These were
amplified and genotyped using GT-seq (Campbell et al., 2015), with a minimum of 20 reads,
using sequences from an Illumina NextSeq 500 (1x75bp). These SNP data, as well as the
RADseq data, were used to distinguish Fraser River F1 family members that were reduced
versus normal ploidy (see Delomas et al., 2021 for details), since not all sperm were effectively
irradiated to eliminate the male contribution.
207
208   **Linkage map construction and phased assembly scaffolding**
209
Further scaffolding of the phased assembly was performed using genetic linkage maps
from segregating polymorphisms within the three sibships described above. Specifically, 1)
polymorphisms from RAD-seq data from reduced-ploidy (hereafter 'haploid') Fraser River
sibships, 2) polymorphisms from RAD-seq data from normal-ploidy (hereafter 'diploid') Fraser
River sibships, and 3) polymorphisms from RNA-seq data from the normal-ploidy Snake River
sibships. For each dataset, we initially constructed a high-confidence linkage map using a
subset of markers with highly replicated genotypes (those wherein the same segregation
patterns across all offspring were observed 5 or more times).
Linkage analysis was performed for each mapping family separately using JoinMap5
using maximum likelihood mapping under the appropriate crossing design for each sibship:
"Haploid" for the haploid sibship and "outbred F1" (CP: cross pollinator) for the other two (Stam,
1993). Additional markers were added to this map if >80% of individuals had identical
genotypes to framework markers when considering segregation in both coupling and repulsion
phases. We then used these three maps as a framework to further scaffold the initial assembly
using Allmaps from JCVI utility libraries 1.4.2 (Tang et al., 2015), with weights for each position
scaled to the number of genotype mismatches relative to a framework marker
(mismatches/weight: 0/20, 1/8, 2/4, 3/2, 4/1) within each map, initially prioritizing contigs that
were captured by the haploid map. Subsequent scaffolding rounds captured additional scaffolds
that were absent from the haploid map, prioritizing the diploid map, and then those that were
represented only in the RNA-seq-derived map. We evaluated the resulting scaffolds using the
geneticmap.py LD function of JCVI.

231
Comparison to the sterlet assembly

To assess the large-scale structure of the scaffolded assembly and identify orthologous
(sturgeon/sterlet) or paralogous (sturgeon/sturgeon) chromosomes, we aligned our white
sturgeon assembly to the sterlet genome assembly (GCF_902713425.1) using minimap v2.28-
r1209 (Li, 2016) with the -asm20 option enables for divergence assembly to reference mapping.
Alignments were visualized using DGENIES v1.5 (Cabanettes & Klopp, 2018). An alignment of
white sturgeon scaffolds to themselves was similarly made.

Genetic markers for phenotypic sex

In comparison of the published female and male assemblies of sterlet sturgeon (*A.
ruthenus*), Kuhl et al. (2021) identified two sex-specific, "unalignable" regions (~10Kbp for
males, ~16Kbp for females) embedded in an otherwise homologous region, with males showing
little to no coverage for the female-specific region (putative W chromosome). We extracted the
sex-specific and flanking regions from these assemblies and searched for similar sequences in
our phased white sturgeon assembly using NCBI *blastn* (Camacho et al., 2009) with minimum
e-value 1×10^{-5} and 50% similarity. This identified three phased scaffolds containing homologous
sequences: two putative Z and 1 putative W scaffolds. Alignments made with Mummer v4
(Marçais et al., 2018) (*nucmer --mum --nosimplify -c 100 -g 1000*) confirmed the homology of
the flanking regions, as well as the sex-specific regions for the respective scaffolds. From
alignments of all of these, made in Geneious (Biomatters), we identified a small (~5Kbp) island
of homology within the otherwise unalignable, sex-specific regions which contained SNP
variants distinguishing the putative Z and W scaffolds (named *zwhr*, for 'Z/W homologous
region'). We developed primers for two small amplicons (<100bp), one covering a subset of
these SNPs in the nested, homologous region, and the other a putative-W-specific amplicon
from the adjacent, unalignable region (named *p455160ex*, for 'phased scaffold 455160
extraction'). We tested the utility of these markers to predict sex phenotype in known sex
sturgeon from the Columbia River below McNary Dam (Washington/Oregon, USA), natural- and
hatchery-origin individuals from the Snake River (Hagerman, Idaho), natural-origin individuals
from Kootenai River (near Bonners Ferry, Idaho, USA), hatchery broodstock from the lower
Fraser River population (British Columbia, Canada), and hatchery broodstock from the
Sacramento River population (California, USA). Sex from these individuals was identified
through abdominal biopsy and/or through hatchery cross records. Markers were optimized for
inclusion into the GT-seq panel (Supplemental File 1), sequenced on an Illumina NextSeq 2000
(1x100bp), and genotyped with a custom pipeline (Delomas et al., 2021).

Results

Estimated coverage depth of the sequencing data used to construct the genome
assembly depended on which published genome size (5.1pg vs. 9.6pg; Blacklidge & Bidwell,
1993; Hinegardner, 1976) was used for calculation (Table 1). The Genomescope estimated
haploid genome size of the individual used for sequencing ranged from 1.514 Gbp to 1.622

Gbp, depending on assumed ploidy (2N or 4N) and kmer length utilized (Supplemental Table 1),
indicating published genome sizes for white sturgeon reflect tetraploidy or hexaploidy.
Consistent with this, Smudgeplot analysis of heterozygous kmers (hetmers) indicated that the
genome of the sequenced individual largely reflected tetraploidy, with an equal or smaller
portion reflective of octoploidy with smaller or larger kmer lengths, respectively (Figure 1,
Supplemental Figure 1). The signal of tetraploidy at longer lengths of k may reflect the products
of a more derived duplication event within Acipenseriformes, while shorter values of k appear to
capture sequence matches resulting from an older duplication within the sturgeon stem lineage.
The total size of the assembled contigs represented 5.96Gbp of genomic sequence, and
these were assembled into initial scaffolds reflecting 6.26Gbp of genomic sequence with 4.86%
gaps (Table 2). An additional 826Kbp of sequence data could not be incorporated into the
assembly, which likely represents complex regions such as those with high GC content or highly
repetitive sequences, as well as possible microbial contamination. Moreover, the scaffolded
assembly, despite the robust structural sequence information, was notably fragmented, with an
N50 of only 339Kbp for 3,870 sequences. Similarly, the largest scaffold, only 8.59Mbp, was
small relative to the sterlet RefSeq assembly, which contained 52 chromosome-level scaffolds
larger than this. In contrast, assessments of genome completeness based on orthologous gene
content using Compleasm indicated the white sturgeon assembly was only marginally less
complete than the female sterlet RefSeq assembly, with 6.1% of orthologs missing vs. 3.3% in
sterlet (Table 3). Notably, both assemblies exhibited similar proportions of single copy (22.8%
and 27.0%, respectively) and duplicated (67.7% and 68.9%, respectively) orthologs, despite the
higher ploidy of white sturgeon.
The mapping families used for scaffolding varied with respect to the number of
individuals passing quality thresholds, numbers of markers captured, and the estimated
recombination rates (Table 4). The map based on haploid RAD-seq data was considered to be
the most accurate because it yielded a number of linkage groups that is close to the expected
number of chromosomes based on karyotypic analyses (1n ~ 135; Van Eenennaam et al.,
1998), with an average rate of recombination of 0.5 per LG per meiosis. The RNA-seq-based
linkage map appeared to overestimate recombination rates, possibly due to lower genotyping
accuracy. Based on the anticipated and realized properties of these maps, we preferentially
used the haploid map for scaffolding, supplementing with the diploid and RNA-seq-based maps
where they did not conflict with the haploid map.
Joining of the initial scaffolds using mapped linkage group markers increased the size of
the largest scaffolds in the assembly, with the largest scaffold becoming 55.2Mbp and an N50 of
416Kbp for 1,976 scaffolds (Table 2). The largest scaffolds of the final, linked assembly showed
clear homology to chromosome-level scaffolds of the sterlet sturgeon, including numerous 1:2
and 1:many relationships (Figure 2). Notably, several white sturgeon scaffolds showed
contiguous homology to large regions of sterlet chromosomes, with a few breaks that either
reflect fission/fusion events or assembly fragmentation (e.g. white sturgeon scaffolds 1 and 3 to
sterlet chromosome 3). Other scaffolds show lower degrees of linear conservation (e.g.
homologs of sterlet chr. 1 and 2), reflecting higher rates of intrachromosomal rearrangement,
potential non-diploid segregation, or errors in scaffolding. Similar patterns of conservation and
synteny disruption are also observed between   divergent paralogous chromosomes within the
sterlet genome, suggesting that   patterns of homology observed   between white sturgeon

and sterlet might reflect true   differences in white sturgeon chromosomal structure. For
example, sterlet chr. 7 has segments that are paralogous to chr. 13 and chr. 14, whereas each
ancestral sequence corresponds to two large scaffolds in sturgeon (scaf. 7, 10, 11 and 16),
indicating that sterlet chr. 7 is likely the product of a derived fission event. Overall   , most
sterlet chromosomes showed similar patterns of homology to two white sturgeon scaffolds,
reflecting the derived paralogy from sublineage-specific whole genome duplications (e.g. white
sturgeon scaf. 1 and 5 to sterlet chr. 3 in Figure 2; see also Supplemental Figure 2). Closer
examination of LD patterns showed that these paralogous white sturgeon chromosomes appear
to segregate independently, with a few notable exceptions (Figure 3).
The white sturgeon scaffolds identified through *blastn* searches with the sterlet sex-
specific regions showed contiguous homology to those sterlet extractions, including in the
respective sex-specific regions (Supplemental Figure 3). Notably, the regions corresponding to
the sterlet male extraction (putative-Z) were joined into the final white sturgeon scaffolds 1 and
5, while the much shorter female-corresponding scaffold (putative W), only 108Kbp, was not
incorporated into the linkage map, exhibiting segregations patterns that were not correlated with
portions of the genome that were captured by linkage analyses. As noted above, these two final
white sturgeon scaffolds are part of a group which form contiguous and duplicated homology
with sterlet chromosome 3, wherein the sex-specific regions reside. Notably, these two joined
scaffolds show evidence of latent linkage disequilibrium (Figure 3), suggesting the possibility
that they continue to pair at meiosis and exhibit ongoing recombination.
The two amplicons developed from these homologous sex-specific regions showed strict
linkage in genotyping, with only individuals exhibiting the putative-W-specific SNP haplotype
also exhibiting successful amplification of the putative-W-specific presence/absence marker
positioned in the adjacent, unalignable region. Also notable, the putative-W-specific haplotype,
when present in individuals with >80% overall genotyping success, reflected approximately
1/4th of the sequenced reads for that locus (mean 0.251, stdev 0.023). The overall efficacy of
the presence or absence of the putative-W-specific haplotype or amplicon at predicting
phenotypic females and males, respectively, was high (96.6% and 81.5%, respectively),
providing a strong overall association with sex (χ^2 p-value < 2.7x10^{-37}). However, the efficacy
varied across stocks (Table 5): for individuals in the lower Columbia River, whose sex was
confirmed both by abdominal biopsy and reproductive success (parentage), accuracy was 100%
in both sexes, but for males in the Sacramento Basin, presence of the putative-W-specific
markers was higher than expected (66%). Other stocks showed intermediate accuracy for one
or both sexes.

Discussion

This study provides a novel genomic resource for white sturgeon which will aid in the
conservation of this charismatic and imperiled species as well as help facilitate research into the
process of genome evolution for polyploid vertebrates. However, this resource remains in a
notably fragmented state compared with the sterlet assembly, which partially reflects the higher
ploidy and inherently higher paralogy within the white sturgeon genome. We utilized a pipeline
that has been successfully applied previously for species with autopolyploid genomes, but it
could well be that the complex nature of the white sturgeon genome, with compounded

autopolyploid events, will require the use more advanced sequencing approaches such as long-
read and Hi-C libraries. Despite the residual fragmentation in the assembly provided here,
orthology assessments indicate that the current assembly contains most of the gene content of
other sturgeon genomes. Thus, the present white sturgeon genome assembly will be useful for
a variety of genome-dependent genetic studies.
To demonstrate the utility of the present genome assembly, we have utilized it in pursuit
of genetic markers for phenotypic sex in white sturgeon. White sturgeon are widely used in
aquaculture for both meat and caviar production, with the latter relying on the production of
mature, female fish. Having molecular markers to predict the sex of fish prior to maturity would
assist commercial operations in managing limited resources most effectively. Further,
conservation aquaculture programs have been implemented to bolster declining populations or
limited recruitment for a number of white sturgeon populations (Mallette, 2014). These programs
are at times forced to selectively cull slower growing young in order to maximize resource
efficiency but without knowing the impact of these events on residual sex ratios. Moreover, while
hatchery fertilization procedures have been optimized in order to minimize the incidence of
spontaneous autopolyploidy in hatchery offspring, selective culling for higher ploidy continues to
be necessary but without knowledge of the dynamics of sex among autopolyploids.
Within the present white sturgeon genome assembly, we identified regions which show
strong homology to the sex-specific regions of other sturgeon (Kuhl et al., 2021). Moreover,
markers developed for two sequences within this region showed strong but variable association
with sex. While the listed primers for these markers have been optimized for inclusion in a
multiplex, GT-seq panel, they could be adapted to be used as a standalone qPCR or visual
assay (e.g., Kanefsky et al., 2022), e.g. presence/absence of the female-specific marker,
p455160ex, possibly with the shared marker, *zwhr*, as a positive control (in which case the
primers of one marker should be 5' elongated to distinguish amplicons based on size). PCR
efficiency or primer concentrations notwithstanding, the expected amplicon concentration for
females would be ~1:4 *p455160ex*:*zwhr*.
However, like any new technology, we urge caution in implementing this assay for
phenotypic sex and suggest validating the specific association with sex using known sex fish in
individual white sturgeon stocks. As indicated by our tests, while the association with sex for
these markers is strong, it varies significantly among the stocks tested here. There are several
explanations for variable association. First, while the association of these sex-specific,
unalignable regions appears clear for white, sterlet, and other sturgeons (Degani et al., 2022;
Kuhl et al., 2021; Scribner & Kanefsky, 2021; this study) it may only be linked to variants that
contribute to rather than being directly involved in sex determination *per se*. The RefSeq
annotation for the female sterlet assembly indicates that the sex-specific region resides in the
first intron of the *ofcc1* gene ("orofacial cleft 1 candidate 1"; LOC117394325), a transcribed but
non-coding region with a possible regulatory role. While this gene is largely uncharacterized in
fishes and has not been directly proposed as a sex determining gene, we note that it is strongly
expressed in embryos as well as the testis and ovaries of male and female zebrafish (*Danio*
rerio) (expression for ENSDARG00000104791 on Bgee, https://www.bgee.org). Whether this
gene, and the sex-associated region within it, is causal or simply linked to sex-determining
variation remains to be determined. Second, patterns of residual linkage suggest that two
scaffolds containing sequences homologous to the sterlet male-specific region (putative Z

chromosomes) experience some level of ongoing recombination. In many species,
recombination is suppressed between the sex chromosomes, or sex regions of pseudo-
autosomes. Thus, it remains to be resolved how distantly situated from the actual locus or
variant(s) of determination these markers lie and how much even low levels of recombination
may diminish association in any given stock. Third, while most species with genetic sex
determination exhibit a single, 'master' sex-determining gene, sex determination could be
polygenic in sturgeon. Indeed, Degani et al. (2022) identified a number of sex-associated SNP
variants in the Russian sturgeon (*A. gueldenstaedtii*), a distantly related, octoploid species. It is
possible that there are additional, unlinked variants which play a larger role in mediating sex
determination in some stocks of white sturgeon. Finally, although it has been accepted for some
time that sturgeon exhibit a female-heterogametic sex determination system (ZW), it is not clear
that sex determination is fully genetic in white or other sturgeon. If not, environmental conditions
that vary among stocks, or between hatchery and natural-origin fish, could diminish the
association of molecular markers for sex despite strong genetic linkage.
Beyond the use of additional sequencing technologies to further resolve the polyploid
white sturgeon assembly, greater utility of a white sturgeon genome resource will come from a
'pan-genome' of several assemblies that represents a compendium of sexual, population
genetic, and other relevant variation from across a species (e.g., Gong et al., 2023). While this
term was originally applied to microbial species which exhibit a 'core' set of genes present
across all individuals and a large library of 'accessory' genes, it has been increasingly applied to
vertebrate species as the unexpected degree of genomic variation among individuals has
become apparent. As has been shown for species of salmonids, gene content and structural
variation among individuals can play a critical role in mediating major life history variations
relevant for both aquaculture and population management (e.g. Pearse 2019, Bertolotti 2020).
For an ancestrally octoploid and functionally tetraploid species with strong genetic structure
among some stocks, the potential for important structural variation to segregate is great, but
beyond even knowing of its existence, representing and making use of that variation requires a
genomic resource that can reflect genomic variation of multidimensional complexity.
Representation and bioinformatic tools making use of pan-genomes is an active area of
research, but the complexity reflected in a single white sturgeon genome assembly highlights
the opportunities created by progress in this area.
438
439 **Data Availability**
440
441 Raw data for linkage map creation are available from the NCBI Short Read Archive
442 (PRJNA1271479). The final assembly and raw data used to construct it are available from NCBI
443 Genome and Short Read Archive, respectively (PRJNA1358253).
444

445 **Acknowledgements**
446
The authors acknowledge Ben Sutherland and Dan Baker, and Gary and Linda Lemmon, who
provided the Fraser River and Snake River white sturgeon families used to construct linkage
maps. Samples from the focal Snake River white sturgeon were collected by Nate Campbell
and Andrew Matala. Laboratory staff at the Hagerman Genetics Lab, in particular Stephanie

Harmon, Kim Vertacnik, and Rachael Kane, facilitated library construction and sequencing of
the linkage map data.
453

454 **Funding**
455

456 Funding for this work was provided by the Bonneville Power Administration (Grant no. 2008-
457 907-00).

459 **Conflicts of Interest**
460

461 The authors declare no conflicts of interest.
462

463 **Table and Figure Legends**
464

465 Table 1. Sequence data statistics. Coverage (depth) is listed for published genome estimates of
466 (1) 5.1pg; (Hinegardner 1976) and (2) 9.6pg (Blacklidge et al. 1993).
467

468 Table 2. Statistics for the white sturgeon genome assemblies.
469

470 Table 3. Results from Compleasm evaluation for otholog content using Actinopterygii_od12 of
471 sterlet and white sturgeon genome assemblies.
472

473 Table 4. Statistics for the linkage maps constructed from RAD-seq (Fraser River family) and
474 RNA-seq (Snake River family) data.
475

476 Table 5. Proportions of known sex white sturgeon exhibiting the predicted genotypes for
477 molecular markers for phenotypic sex. In parentheses, correctly predicted and all samples for a
478 given sex. For the Snake River, natural-origin (NOR) and hatchery-origin (HOR) individuals are
479 distinguished.
480

481 Supplemental Table 1. Haploid genome sizes estimated with Genomescope from various kmer
482 lengths and diploid or tetraploid status.
483

484 Figure 1. Genome content estimated from Smudgeplot analysis of heterozygous kmers with
485 k=41bp.
486

Figure 2. Alignment of white sturgeon scaffolds to the sterlet genome. Plots reflect alignments
between the largest 50 white sturgeon and sterlet scaffolds (left). Three sets of sterlet homologs
are outlined in red, orange and pink correspond to outlined chromosomes in Figure 3. Sterlet
self-alignments (right) are shown to highlight older paralogy derived from a common ancestral
duplication in the sturgeon lineage. A red asterisk marks the approximate location of the sterlet
sex determining region.
493

Figure 3. Linkage disequilibrium (ld) among segregating maternal polymorphisms from haploid segregants. A) Patterns of ld within and between the largest 25 scaffolds with segregating maternal markers. Chromosomes outlined in red, orange and pink correspond to outlined chromosomes in Figure 1. B) Patterns of ld within and between four scaffolds that are homologous to the sterlet sex chromosome.

Supplemental Figure 1. Genome content analysis using kmers. A) Representative Genomescope results, with k=21bp and ploidy=4N. B) Smudgeplot analysis with k=21bp. C) Smudgeplot analysis with k=23bp. D) Smudgeplot analysis with k=31bp. E) Smudgeplot analysis with k=41bp.

Supplemental Figure 2. Alignment of white sturgeon largest 50 scaffolds to themselves. Scaffold ordering follows Figure 2.

Supplemental Figure 3. Mummer alignments of the sex-specific and flanking regions of sterlet sturgeon (from Du et al. 20XX) and three homologous scaffolds from the white sturgeon phased assembly.

Supplemental Files.

Supplemental File 1. GT-seq panel for white sturgeon, revised Sept. 2025

References

Baird, N. A., Etter, P. D., Atwood, T. S., Currey, M. C., Shiver, A. L., Lewis, Z. A., Selker, E. U., Cresko, W. A., & Johnson, E. A. (2008). Rapid SNP discovery and genetic mapping using sequenced RAD markers. *PLoS ONE, 3*(e3376).

Blacklidge, K. H., & Bidwell, C. A. (1993). Three ploidy levels indicated by genome quantification in acipenseriformes of North America. *Journal of Heredity, 84*(6). https://doi.org/10.1093/oxfordjournals.jhered.a111367

Bushnell, B. (2015). *BBMap*. Https://Sourceforge.Net/Projects/Bbmap/.

Cabanettes, F., & Klopp, C. (2018). D-GENIES: Dot plot large genomes in an interactive, efficient and simple way. *PeerJ, 2018*(6). https://doi.org/10.7717/peerj.4958

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics, 10.* https://doi.org/10.1186/1471-2105-10-421

Campbell, N. R., Harmon, S. A., & Narum, S. R. (2015). Genotyping-in-Thousands by sequencing (GT-seq): A cost effective SNP genotyping method based on custom amplicon sequencing. *Molecular Ecology Resources, 15*(4), 855–867. https://doi.org/10.1111/1755-0998.12357

Catchen, J. M., Amores, A., Hohenlohe, P. A., Cresko, W. A., & Postlethwait, J. H. (2011). Stacks: Building and Genotyping Loci De Novo From Short-Read Sequences. *G3: Genes, Genomes, Genetics`, 1*, 3171–3182.

Danacek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R.,
Lunter, G., Marth, G., Sherry, S. T., McVean, G., Durbin, R., & Group, 1000 Genomes
Project Analysis. (2011). The Variant Call Format and VCFtools. *Bioinformatics*, *27*(15),
2156–2158.
Danacek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A.,
Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools and
BCFtools. *GigaScience*, *10*(2), giab008. https://doi.org/10.1093/gigascience/giab008
Degani, G., Nevo Sarel, M., Hajouj, A., Hurvitz, A., Veksler-Lublinsky, I., & Meerson, A. (2022).
Whole-Genome Inter-Sex Variation in Russian Sturgeon (Acipenser gueldenstaedtii).
International Journal of Molecular Sciences, *23*(16). https://doi.org/10.3390/ijms23169469
Delomas, T. A., Willis, S. C., Parker, B. L., Miller, D., Anders, P., Schreier, A., & Narum, S.
(2021). Genotyping single nucleotide polymorphisms and inferring ploidy by amplicon
sequencing for polyploid, ploidy-variable organisms. *Molecular Ecology Resources*, *21*(7).
https://doi.org/10.1111/1755-0998.13431
Drauch Schreier, A., Gille, D., Mahardja, B., & May, B. (2011). Neutral markers confirm the
octoploid origin and reveal spontaneous autopolyploidy in white sturgeon, Acipenser
transmontanus. *Journal of Applied Ichthyology*. https://doi.org/10.1111/j.1439-
0426.2011.01873.x
Du, K., Stöck, M., Kneitz, S., Klopp, C., Woltering, J. M., Adolfi, M. C., Feron, R., Prokopov, D.,
Makunin, A., Kichigin, I., Schmidt, C., Fischer, P., Kuhl, H., Wuertz, S., Gessner, J., Kloas,
W., Cabau, C., Iampietro, C., Parrinello, H., … Schartl, M. (2020). The sterlet sturgeon
genome sequence and the mechanisms of segmental rediploidization. *Nature Ecology and
Evolution*. https://doi.org/10.1038/s41559-020-1166-x
Fontana, F. (2002). A cytogenetic approach to the study of taxonomy and evolution in
sturgeons. *Journal of Applied Ichthyology*, *18*(4–6). https://doi.org/10.1046/j.1439-
0426.2002.00360.x
Fontana, F., Lanfredi, M., Kirschbaum, F., Garrido-Ramos, M. A., Robles, F., Forlani, A., &
Congiu, L. (2008). Comparison of karyotypes of Acipenser oxyrinchus and A. sturio by
chromosome banding and fluorescent in situ hybridization. *Genetica*, *132*(3).
https://doi.org/10.1007/s10709-007-9171-4
Fopp-Bayat, D., Kolman, R., & Woznicki, P. (2007). Induction of meiotic gynogenesis in sterlet
(Acipenser ruthenus) using UV-irradiated bester sperm. *Aquaculture*, *264*(1–4).
https://doi.org/10.1016/j.aquaculture.2006.12.006
Gille, D. A., Famula, T. R., May, B. P., & Schreier, A. D. (2015). Evidence for a maternal origin
of spontaneous autopolyploidy in cultured white sturgeon (Acipenser transmontanus).
Aquaculture. https://doi.org/10.1016/j.aquaculture.2014.10.002
Gong, Y., Li, Y., Liu, X., Ma, Y., & Jiang, L. (2023). A review of the pangenome: how it affects
our understanding of genomic variation, selection and breeding in domestic animals? In
Journal of Animal Science and Biotechnology (Vol. 14, Issue 1).
https://doi.org/10.1186/s40104-023-00860-1
Hildebrand, L. R., Drauch Schreier, A., Lepla, K., McAdam, S. O., McLellan, J., Parsley, M. J.,
Paragamian, V. L., & Young, S. P. (2016). Status of White Sturgeon (Acipenser
transmontanus Richardson, 1863) throughout the species range, threats to survival, and
prognosis for the future. *Journal of Applied Ichthyology*. https://doi.org/10.1111/jai.13243

Hinegardner, R. (1976). The cellular DNA content of sharks, rays and some other fishes.
Comparative Biochemistry and Physiology -- Part B: Biochemistry And, *55*(3).
https://doi.org/10.1016/0305-0491(76)90305-9
Huang, N., & Li, H. (2023). compleasm: a faster and more accurate reimplementation of
BUSCO. *Bioinformatics*, *39*(10). https://doi.org/10.1093/bioinformatics/btad595
Jighly, A., Lin, Z., Forster, J. W., Spangenberg, G. C., Hayes, B. J., & Daetwyler, H. D. (2018).
Insights into population genetics and evolution of polyploids and their ancestors. *Molecular*
Ecology Resources, *18*(5). https://doi.org/10.1111/1755-0998.12896
Kanefsky, J., Smith, S., & Scribner, K. T. (2022). Real-Time PCR-Based Method for Sex
Determination in Lake Sturgeon (Acipenser fulvescens). *Diversity*, *14*(10).
https://doi.org/10.3390/d14100839
Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*, *37*(8).
https://doi.org/10.1038/s41587-019-0201-4
Kopf, R., Banks, S., Brent, L., Humphries, P., Jolly, C., Lee, P., Luiz, O., Nimmo, D., &
Winemiller, K. O. (2025). Loss of Earth's old, wise, and large animals. *Science*, *387*(6729).
Korman, J., Crossman, J. A., McLellan, J. G., & Miller, A. L. (2025). The effectiveness of
conservation aquaculture in meeting recovery objectives for an endangered White
Sturgeon population. *Transactions of the American Fisheries Society*, *154*, 537–558.
Kuhl, H., Guiguen, Y., Höhne, C., Kreuz, E., Du, K., Klopp, C., Lopez-Roques, C., Yebra-
Pimentel, E. S., Ciorpac, M., Gessner, J., Holostenco, D., Kleiner, W., Kohlmann, K.,
Lamatsch, D. K., Prokopov, D., Bestin, A., Bonpunt, E., Debeuf, B., Haffray, P., … Stöck,
M. (2021). A 180 Myr-old female-specific genome region in sturgeon reveals the oldest
known vertebrate sex determining system with undifferentiated sex chromosomes.
Philosophical Transactions of the Royal Society B: Biological Sciences, *376*(1832).
https://doi.org/10.1098/rstb.2020.0089
Li, H. (2016). Minimap and miniasm: Fast mapping and de novo assembly for noisy long
sequences. *Bioinformatics*, *32*(14). https://doi.org/10.1093/bioinformatics/btw152
Lobanov, V. P., Pate, J., & Joyce, A. (2024). Sturgeon and paddlefish: Review of research on
broodstock and early life stage management. In *Aquaculture and Fisheries* (Vol. 9, Issue
6). https://doi.org/10.1016/j.aaf.2023.04.001
Mallette, C. (2014). *White Sturgeon mitigation and restoration in the Columbia and Snake*
Rivers upstream from Bonneville Dam. Annual progress report (2012) to the Bonneville
Power Administration.
Marçais, G., Delcher, A. L., Phillippy, A. M., Coston, R., Salzberg, S. L., & Zimin, A. (2018).
MUMmer4: A fast and versatile genome alignment system. *PLoS Computational Biology*,
14(1). https://doi.org/10.1371/journal.pcbi.1005944
Marçais, G., & Kingsford, C. (2011). A fast, lock-free approach for efficient parallel counting of
occurrences of k-mers. *Bioinformatics*, *27*(6). https://doi.org/10.1093/bioinformatics/btr011
Micheletti, S. J., Matala, A. R., Matala, A. P., & Narum, S. R. (2018). Landscape features along
migratory routes influence adaptive genomic variation in anadromous steelhead
(Oncorhynchus mykiss). *Molecular Ecology*, *27*(1), 128–145.
https://doi.org/10.1111/mec.14407

Myers, G. (2020). *A fast K-mer counter for high-fidelity shotgun datasets*.
https://github.com/thegenemyers/FASTK.
Parsley, M. J., & Beckman, L. G. (1994). White Sturgeon Spawning and Rearing Habitat in the
Lower Columbia River. *North American Journal of Fisheries Management*, *14*(4).
https://doi.org/10.1577/1548-8675(1994)014<0812:wssarh>2.3.co;2
Parsley, M. J., Beckman, L. G., & McCabe, G. T. (1993). Spawning and Rearing Habitat Use by
White Sturgeons in the Columbia River Downstream from McNary Dam. *Transactions of*
the American Fisheries Society, *122*(2). https://doi.org/10.1577/1548-
8659(1993)122<0217:sarhub>2.3.co;2
Peng, Z., Ludwig, A., Wang, D., Diogo, R., Wei, Q., & He, S. (2007). Age and biogeography of
major clades in sturgeons and paddlefishes (Pisces: Acipenseriformes). *Molecular*
Phylogenetics and Evolution, *42*(3). https://doi.org/10.1016/j.ympev.2006.09.008
Ranallo-Benavidez, T. R., Jaron, K. S., & Schatz, M. C. (2020). GenomeScope 2.0 and
Smudgeplot for reference-free profiling of polyploid genomes. *Nature Communications*,
11(1). https://doi.org/10.1038/s41467-020-14998-3
Redmond, A. K., Casey, D., Gundappa, M. K., Macqueen, D. J., & McLysaght, A. (2023).
Independent rediploidization masks shared whole genome duplication in the sturgeon-
paddlefish ancestor. *Nature Communications*, *14*(1). https://doi.org/10.1038/s41467-023-
38714-z
Scribner, K. T., & Kanefsky, J. (2021). Molecular sexing of lake sturgeon. *Journal of Great*
Lakes Research, *47*(3). https://doi.org/10.1016/j.jglr.2021.03.015
Spoelhof, J. P., Soltis, P. S., & Soltis, D. E. (2017). Pure polyploidy: Closing the gaps in
autopolyploid research. In *Journal of Systematics and Evolution*.
https://doi.org/10.1111/jse.12253
Stam, P. (1993). Construction of integrated genetic linkage maps by means of a new computer
package: JOINMAP. *Plant Journal*, *3*(5). https://doi.org/10.1111/j.1365-313X.1993.00739.x
Tang, H., Zhang, X., Miao, C., Zhang, J., Ming, R., Schnable, J. C., Schnable, P. S., Lyons, E.,
& Lu, J. (2015). ALLMAPS: Robust scaffold ordering based on multiple maps. *Genome*
Biology, *16*(1). https://doi.org/10.1186/s13059-014-0573-1
Van Eenennaam, A. L., Murray, J. D., & Medrano, J. F. (1998). Synaptonemal complex analysis
in spermatocytes of white sturgeon, Acipenser transmontanus richardson (pisces,
acipenseridae), a fish with a very high chromosome number. *Genome*, *41*, 51–61.
https://doi.org/10.1139/g97-101
Van Eenennaam, A. L., Van Eenennaam, J. P., Medrano, J. F., & Doroshov, S. I. (1999).
Evidence of female heterogametic genetic sex determination in white sturgeon. *Journal of*
Heredity, *90*(1). https://doi.org/10.1093/jhered/90.1.231
Wertheim, B., Beukeboom, L. W., & Van De Zande, L. (2013). Polyploidy in animals: Effects of
gene expression on sex determination, evolution and ecology. *Cytogenetic and Genome*
Research, *140*(2–4). https://doi.org/10.1159/000351998
Winemiller, K. O. (2005). Life history strategies, population regulation, and implications for
fisheries management. *Canadian Journal of Fisheries and Aquatic Sciences*, *62*(4), 872–
885.
666

Pairwise Linkage Disequilibrium Between Markers

0.0   0.5   1.0

Library type	Insert size	Run Type	Sequencing data (Gbp)	Depth(1)	Depth(2)
Paired-end	450-470bp	2x250bp	365.8	73.3	39.0
Paired-end	700-800bp	2x150bp	236.3	47.4	25.2
Mate Pair	2-4Kbp	2x150bp	162.0	32.5	17.3
Mate Pair	5-7Kbp	2x150bp	185.7	37.2	19.8
Mate Pair	8-10Kbp	2x150bp	160.5	32.2	17.1
10X Chromium	N/A	2x150bp	169.3	33.9	18.0

	Contigs	Scaffolds	Linked Scaffolds
Total contigs/scaffolds	1,347,142	835,784	832,145
Assembly size (bp)	5,960,370,858	6,264,925,766	6,265,289,666
Gaps size (bp)	N/A	304,554,908	304,918,808
Gaps (%)	N/A	4.861	4.867
N50 (bp)	13,367	339,349	415,779
N50 (#seq)	90,723	3,870	1,976
N90 (bp)	1,565	2,015	2,016
N90 (#seq)	761,174	402,904	399,337
Largest contig/scaffold (bp)	359,327	8,590,638	55,262,319

	sterlet RefSeq (GCF_902713425.1)	white sturgeon phased scaffolds
Complete, single copy	27.00% (1946)	22.80% (1643)
Complete, duplicated	68.91% (4966)	67.67% (4877)
Fragmented	0.82% (59)	3.42% (247)
Missing	3.27% (236)	6.11% (440)

	'haploid' (RAD)	'diploid' (RAD)	RNA
N indivduals	113	37	92
total LGs	168	156	435
total cM	9651.818	3815.993	186319.269
total variants	1678	951	11359

	female ZZZW	male ZZZZ
Fraser River	85% (6/7)	80% (8/10)
Sacramento River	100% (9/9)	44% (4/9)
Columbia River (McNary)	100% (40/40)	100% (36/36)
Snake River NOR	100% (21/21)	75% (15/20)
Snake River HOR	100% (10/10)	90% (9/10)
Kootenai River	92% (23/25)	100% (24/24)

kmer	ploidy	min (bp)	max (bp)
21	2N	1,514,830,630	1,519,678,665
21	4N	1,515,246,697	1,519,912,333
23	2N	1,530,386,796	1,535,475,021
23	4N	1,532,477,521	1,536,954,560
31	2N	1,574,211,122	1,580,387,873
31	4N	1,582,577,639	1,586,632,609
41	2N	1,604,596,808	1,612,141,825
41	4N	1,618,902,487	1,622,808,454