Context-specific configuration of orthogonal integrator dynamics for flexible foraging decisions

Lyle Kingsbury[1,2,*], Grace Zhang[1,2], Juan Ignacio Sanguinetti-Scheck[1,2,3], Naoshige Uchida[1,2,*]

1. Dept. of Molecular and Cellular Biology, Harvard University, Cambridge, MA, USA
2. Center for Brain Science, Harvard University, Cambridge, MA, USA
3. Current address: Dept. of Psychology, University of Pennsylvania, Philadelphia, PA, USA

* Correspondence: lylekingsbury@fas.harvard.edu, uchida@mcb.harvard.edu

Abstract

Adaptation of cognition and behavior across different contexts is a fundamental feature of intelligence, yet how neural circuits are reconfigured to support such flexibility remains poorly understood. Foraging, a ubiquitous and ecological behavior, provides a natural window into the neural basis of decision making and cognitive flexibility. We developed an experimental paradigm in which mice forage across two environment contexts defined by distinct reward dynamics. Mice adopted context-specific decision strategies that can be described as different configurations of integrator dynamics for "stay or leave" foraging decisions. Neural recordings reveal brain-wide activity states which organize largely separate neural subpopulations encoding context-specific configurations of integrator decision variables. In the dorsal frontal cortex, this orthogonal decision coding is preserved across free-moving and virtual task conditions and is necessary for flexible decision making. Together, these findings identify orthogonalization of integrator dynamics as a general neural mechanism that enables rapid, volitional switching between decision strategies.

Introduction

The capacity to adapt cognition and behavior across different contexts is a core feature of natural intelligence and is crucial for surviving in dynamic natural environments. A fundamental goal of neuroscience is to understand how cognitive processes such as flexible decision making arise from underlying cell and circuit mechanisms. In mammals, the frontal cortex is critical for cognitive flexibility, which is thought to involve top-down regulation of neural network modes for information processing and behavioral control.[1–5] Theory and modeling studies give insight into how network modes may be organized by neuron physiology, connectivity, and circuit architecture to perform context-specific computations for flexible behavior.[6–10]

Despite these advances, experimental progress in linking neural dynamics models with cell/circuit-level mechanisms has been limited. The cornerstone of investigations in animals has been rule-based choice tasks, which elucidate how learned context cues can configure perceptual decision processes.[11–15] Most of these studies identify mechanisms that locate context effects upstream of decision dynamics, incorporating context contingencies through attentional shifting (e.g. across modalities),[13,14,16–18] sensorimotor re-mapping,[12] or altering how sensory inputs perturb decision activity.[11,15,19] However, population coding theories, modeling, and other studies also raise the possibility that decision activity could itself be parsed into different modes (e.g. encoded in separate activity subspaces) to enable flexible selection across contexts.[7,10,20–25]

It is now appreciated that neural mechanisms for flexible computation are diverse, sometimes idiosyncratic, and likely variable across problem domains,[8,19] yet the extant literature has focused mostly on the narrow domain of perceptual decision making. While deeply informative and foundational, this program has left open the question of how flexibility is achieved for complex economic and social problems that are more central to animal's native ecology.[26,27] Indeed, it is unknown how neural mechanisms tuned for perceptual choice tasks may generalize outside that domain, or whether (and what) computational and neural solutions are used for ecological problems that flexible cognition likely evolved to solve.

In this vein, recent years have seen a conceptual shift in neuroscience toward natural behaviors and task designs that better preserve the native regime of brain functions for ecological challenges.[26,27] Foraging – how to source and harvest resources from the environment – is a challenge faced across the animal kingdom.[28,29] Animals routinely make decisions about where to forage and when to leave depleting resource locations ("patches") in search of alternatives (the "patch leaving problem"),[30,31] and can adjust foraging strategies across changing environments or contingencies. In contrast to alternative-choice perceptual decisions, patch foraging is immediately intuitive for animals and requires little training. Inspired by this, recent studies have begun to uncover the neural basis of foraging behaviors.[32,33] Evidence suggests that patch leaving decisions can be solved through a neural process akin to bounded evidence accumulation, linking natural cognition to classical models.[34–38] The flexibility and ecological grounding of foraging thus open a new window onto the computational logic and neural basis of context-dependent decision making that meets animal subjects closer to "where they are."

We developed an experimental paradigm in which mice flexibly toggle decision strategies for patch foraging across environment contexts using distinct configurations of a neural integrator process. Using an equivalent virtual task for large-scale neural recordings, we find that environment contexts recruit distinct, brain-wide activity states. In contrast to context-invariant

decision dynamics observed in perceptual choice tasks, these activity states organize integrator dynamics for foraging decisions into context-specific, orthogonal coding dimensions realized in separate neuronal subpopulations. In the dorsal frontal cortex (dFC), this coding structure is shaped by experience and preserved across experimental settings, even when mice make repeated, volitional context switches. Inactivation experiments demonstrate causal necessity of dFC for context-specific configuration of decision modes. Together, our results reveal a cellular-level neural basis for context-dependent decision making in orthogonal integrator dynamics and open a new avenue to investigate cell and circuit mechanisms of cognition in the mammalian brain.

Results

A patch foraging task for context-dependent decision making in mice

In patch foraging,[30,31] animals encounter resource patches that deplete over time and must decide when to leave in search of alternatives (**Figure 1A**). We hypothesized that mice foraging in environments with different resource distributions could learn and deploy context-specific patch leaving decision strategies. We developed a task in which mice freely navigate an arena with two sub-chambers ("environment" contexts) for water rewards from multiple ports ("patches," **Figures 1B-C**). Mice nose-poke a port to initiate a trial and immediately (t = 0s) receive a 2µL water drop, after which they can stay for more rewards. While patches in the "Stochastic" (Stc, **Figure 1B**, top) context (the left chamber) yield probabilistic rewards each second with a decaying probability (**Figure S1A**), patches in the "Deterministic" (Det, **Figure 1B**, bottom) context (the left chamber) always yield three rewards at fixed times (t = 0, 1, 2s). 15% of the trials in the Stc environment were programmed with this Det reward sequence ("Stc:012" trials) to create reward-matched trials for behavior analysis.

Mice first foraged in the Det context to learn a decision strategy for the deterministic distribution. In Phase 1 (**Figure 1C**, left), mice quickly learned to harvest water, performing an average of 64 trials on session 1 within the first three minutes (**Figures S1B-C**). Though initially impatient, mice learned to collect all rewards within the first session (80% of trials, **Figure S1D**). Examination of patch residence time (PRT, how long mice stay at a patch) distributions across Phase 1 revealed calibration of the decision process over time (**Figures S1E-G**) to Det rewards.

Mice then proceeded to Phase 2 (**Figure 1C**, middle), in which they explored both Det and Stc contexts once per session in blocks, allowing them to learn context-specific decision strategies. In Phase 3, the door separating the chambers was left open and mice could roam freely and switch contexts volitionally (**Figure 1C**, right). Overall, mice collected more rewards per trial on Det patches and preferred them (**Figure S1H-I**), possibly reflecting risk aversion or small differences in expected value. Nonetheless, mice explored both contexts extensively, switching on average 2.25 times per minute and sampling all patches multiple times in every session (**Figure S1J-K**).

Mice exhibit context-specific decision strategies for different foraging environments

To characterize context-specific decision strategies, we compared PRTs across different trial types. Stc trials included four hidden types with different reward probability levels (**Figure S1A**) which correlated positively with PRTs, confirming that mice use reward information to make

Figure 1

A flexible foraging task for context-dependent decision making

A configurable integrator explains context-specific foraging decisions

Figure 1: Mice use rapidly flexible foraging strategies across different environments.

A) Illustration of the classical patch leaving problem (when to leave a patch with depleting resources); animals must make patch leaving decisions across two environment contexts with different resource distributions.
B) Reward returns (2µL water drops with probability each second) for Stochastic (Stc, orange) and Deterministic environment (Det, purple) patches.
C) Progression through flexible foraging task, from Det environment (Phase 1, 7-10 days) to block structure (Phase 2, 14 days) and unconstrained volitional transitions (door open, Phase 3).
D) Mean patch residence times (PRT, mean ± s.d.) for Det and Stc:012 patch trials (SR test, p = 0.218, n = 20 mice). Stc:012 trials are contained in the Stc environment but programmed to give the Det reward return (rewards at t = 0, 1, 2s).
E) CDF of leaving probabilities P(Leave) over time for Det and Stc:012 patches from example Mouse 227 (KS test, p=3.25e-5).
F) Difference in stay probability P(Stay) on Det vs. Stc:012 patches for each mouse (gray lines) and average across mice (black line, n = 20 mice).
G) Comparison of P(Stay) at t = 2 seconds for Det vs. Stc:012 patches (mean ± s.d., SR test, p = 0.0013, n = 20 mice).
H) Comparison of time on patch when P(Leave) = 0.9 for Det vs. Stc:012 trials (mean ± s.d., SR test, p = 0.0038, n = 20 mice).
I) KS statistics comparing different trial type PRT distributions (SR tests, p = 2.19e-4, Det/Det vs. Det/Stc:012, p = 0.0522, Det/Stc:012 vs. Det/Stc, n = 20 mice).
J) Schematic of drift-diffusion integrator model for patch-leaving decisions (top). Decision variable (DV) X_t is incremented each time step by drift rate δ and perturbed by an ρ upon reward receipt ($\rho = 0$ otherwise), accumulating to a boundary to trigger a leaving decision. pLeave$_t$ is generated by a logistic function transform of X_t with temperature T (see Methods). Parameter configurations (δ, ρ) implement different computations for leaving decisions (Models 1-4).
K) Performance (cross-validated Bayesian information criterion, cv-BIC) of integrator models 1-4 fit to mouse PRT behavior and tested on held-out data for Stc patch trials (SR test between models 1v3, 2v3, 3v4, p=0.857e-5 for each, n = 20 mice).
L) Model performance as in **(K)** for Det trials (SR test between models 1v3, 2v3, 3v4; p=0.857e-5 for each, n = 20 mice).
M) Performance of "reward integration" Model 3 (correlation r between predicted and true PRTs) for Stc and Det patch trials. Each mouse is one dot; red circles indicate independent statistical significance (p < 0.01) of regression on predicted vs. true PRTs for that mouse.
N) Model 3 prediction vs. true PRTs for example mouse 219; red line indicates linear regression.
O) Comparison of fit drift rate parameter δ to Model 3 for Stc vs. Det trials (mean ± s.d., SR test, p = 0.0036, n = 20 mice).
P) Comparison of fit reward sensitivity parameter ρ to Model 3 for Stc vs. Det trials (mean ± s.d., SR test, p = 0.023, n = 20 mice).
Q) Left, Rollout of Model 3 DV for integrator parameters fit to Stc or Det trials (example mouse 251). Right, example of P(Leave) CDFs generated from simulated PRTs using mouse 251 parameters (see Methods), recapitulating the typical pattern of flexible decision behavior.

Standard deviation (s.d.); Kolmogorov-Smirnov test (KS test); Wilcoxon signed-rank test (SR test); ***p<0.001; **p<0.01; *p<0.05; n.s. p>0.05.

leaving decisions (**Figure S2A**).[37,38] Examining Det vs. Stc:012 trials, we observed no difference
in average PRTs (**Figure 1D**). However, examination of PRT distributions with the CDF of leaving
probabilities P(Leave) revealed clear differences in decision behavior across contexts. For
example, while Mouse 227 typically waited to collect all Det rewards and then left (after t = 2s),
it did not exhibit this "step-like" pattern on Stc:012 patches, often leaving before the final reward
or staying after (**Figure 1E**). This pattern was consistent across mice (**Figure 1F**), and indeed,
mice were significantly more likely to collect all rewards on Det vs. Stc:012 trials (**Figure 1G**)
and faster to leave on Det trials after collecting the final reward at t = 2s (**Figure 1H**).

To quantify context-specific decision flexibility, we measured the difference between Det
and Stc:012 PRT distributions using the Kolmogorov-Smirnov (KS) statistic, which measures the
maximum vertical difference in P(Leave) CDFs (**Figures 1E, S2B**). We found that PRT
distributions for Det vs. Stc:012 trials were significantly more different than subsampled Det trials,
despite identical reward structure (**Figure 1I**). Divergence of PRT distributions was detectable
early in Phase 2 and increased over time, indicating calibration of decision strategies that are
maintained in Phase 3 (**Figures S2C, S1G**). While this suggests that mice learn distinct strategies,
they may also rely on recent trial outcomes to continually update their strategy. To test this, we
analyzed PRTs on trial sequences in which mice performed a Stc:012 followed by a Det trial
(**Figure S2D**). We observed a similar difference in PRT distributions (**Figure S2E**), even for paired
trial comparisons (**Figure S2F**). This shows that mice immediately adapt to context transitions and
exhibit context-specific behavior without needing to sample recent outcomes. Taken together,
these results demonstrate that mice calibrate their decision process to rewards in different
environment contexts and can flexibly and rapidly toggle context-specific decision strategies.

Configurations of a common integrator process explain decision flexibility

We previously showed that patch leaving decisions in mice are well described by a bounded
accumulator process instantiated by neural dynamics that integrate time and reward as "evidence"
for, or against, patch leaving (**Figure 1J**).[34,37] We hypothesized that an integrator process may
explain decisions in the present task and may account for decision flexibility. To test this, we
modeled leaving decisions as a drift-diffusion decision variable (DV) which increments each time
step (by drift rate δ) and is perturbed on reward receipt (by reward sensitivity ρ), and then passed
through a logistic function, generating a leaving probability pLeave$_t$ (**Figure 1J**, top**)**. In this
model, the parameters δ and ρ govern decision behavior and, as in classical models of perceptual
decisions,[35,36] can be thought of as weights of evidence for or against the leaving decision.
Different configurations produce different algorithms for solving the patch leaving problem
(**Figure 1J**, bottom). For example, while ρ = 0 yields a reward-insensitive integrator that only
tracks time on the patch (equivalent to the "giving up time" heuristic in classical foraging
theories),[30] ρ > 0 yields a counting strategy that accumulates reward quantity.

In principle, mice could use different algorithms for Stc vs. Det leaving decisions. To test
which algorithm best explained behavior, we fit different models of the leaving decision to
empirical PRTs and measured predictive performance on held-out trials. Consistent with our
previous report,[37] we found that the "reward integration" algorithm ("Model 3") outperformed
others (**Figures 1K-L**) and was the best model for all mice for both patch types, suggesting that
mice integrate time and reward as decision evidence. Despite distinct patterns of behavior (**Figures**

1E-F), the reward integration model captured PRT variability for both Det and Stc patches consistently across mice (**Figures 1M-N**). As these results indicate a common algorithm, this raises the possibility that context-specific decision strategies could be driven by different parameter configurations (i.e. δ and ρ) of the integrator process.

To test this, we compared δ and ρ for models fit to Stc vs. Det patches and found consistent differences in parameterization: while Det trials had significantly larger drift rates δ (**Figure 1O**), they also had larger negative values for reward sensitivity ρ (**Figure 1P**). To see how these configurations could produce the observed behavior, we plotted DVs following a Det reward return from example mice (**Figure 1Q**, left). These produced distinct DV rollouts, and sampling simulated PRTs qualitatively replicated the pattern of behavior observed in most mice (**Figure 1Q,** right). Taken together, these results suggest that context-specific behavior may be driven by different configurations of a common integrator process underlying patch leaving decisions.

An equivalent head-fixed virtual task for wide sampling of neural activity

We next aimed to investigate the neural mechanisms underlying contextual configuration of the decision process. While the free-moving (FM) task design was critical for understanding naturalistic behavior,[26,27] constraint with head-fixation enables large-scale neural recordings and can reduce some confounds such as unmeasured movements.[39] We therefore developed a structural equivalent virtual-reality[37] (VR) version of the flexible foraging task (**Figure 2B**, left). Mice run on a treadmill to navigate a virtual corridor and stop at patches – spaced landmarks indicated with visual and odor cues – to collect 2µL water rewards. While stopped, they can receive water each second after an initial reward. As in the FM task, mice encountered Stc and Det patches in distinct virtual environments. Det patches always yield three rewards at fixed times (at t = 0, 2, 4s, **Figure 2B**, right), and 10% of Stc trials had this same return ("Stc:024" trials). Otherwise, Stc patches gave rewards each second with a decaying probability (**Figure 2B**, **S3A**). After a habituation phase (**Figure 2C**), animals foraged in the Det context for 7-10 days (Phase 1), then proceeded to Phase 2, where they foraged in Det and Stc contexts in 20-25-minute blocks once per session with one context transition (a black screen shown for 10 seconds).

While all mice foraged in VR, we observed wide individual variability in task engagement, waiting behavior, and decision behavior, consistent with our previous report.[37] On average, mice performed 71 trials/session with a range of engagement rates (**Figure S3B**). While mice invested an average of 9.5 sec/trial, median PRTs also varied substantially, indicating individual differences in patience levels (**Figure S3C**). PRTs on Stc patches were again correlated positively with reward frequency (**Figure 2D**) and were similar between Det vs. Stc:024 trials (**Figure 2E**).

Despite this variability, P(Leave) CDFs revealed a tendency for mice to stay more at early times on Det patches and to over-stay on Stc:024 patches (**Figure 2F**), as in the FM task. Analysis of PRT distributions confirmed that decision behavior was more distinct across Det vs. Stc:024 patches than among sub-sampled Det patches (**Figure 2G**), and many animals (e.g. Mouse 124, **Figure 2H**) showed dramatic contextual shifts in behavior consistent with this group trend.

We further tested whether decision flexibility could similarly be captured by different configurations of a common integrator process. As before, we fit four versions of the integrator model to Stc and Det PRTs. Model comparisons showed that time/reward integration ("Model 3")

Figure 2

Figure 2. Mice exhibit decision flexibility in an equivalent virtual-reality task.

A) Cartoon of "experiment space" tradeoff illustrating how experimental constraint (such as head-fixation) can improve mechanistic interpretability/access at the expense of ecological validity.

B) Left, illustration of structural equivalent virtual-reality (VR) task in which mice do patch foraging for water rewards and encounter Stochastic (Stc) and Deterministic (Det) patches in different environment contexts cued with visuals and odors. Right, reward returns for Stc and Det patches in the VR task (also see **Figure S3A**).

C) Training timeline for VR task. Mice are habituated to head fixation and running wheel, first exposed to the Det context (Phase 1), then forage in block-structured contexts (Phase 2).

D) Mean patch residence times (PRT, mean ± SEM) for Stc patches with different initial reward probabilities (P_i = 0.1, 0.25, 0.57, 0.95, SR tests, p = 0.0315 [0.1 vs. 0.95], p = 0.01746 [0.25 vs. 0.95], p = 0.0996 [0.57 vs. 0.95], n = 40 mice).

E) Comparison of PRTs (mean ± SEM) for Det vs. Stc:024 (Stc context patches programmed with the Det reward return) trials (SR test, p=0.6653, n = 40 mice).

F) Average difference in P(Stay) on Det vs. Stc:024 patches (mean ± SEM, n = 36 mice).

G) KS statistics for differences in PRT distributions across trial type comparisons (SR tests, p = 2.1661e-7 [Det/Det vs. Det/Stc:024], p = 0.3379 [Det/Stc:024 vs. Det/Stc], n = 36 mice).

H) PRT distributions (CDF of P(Leave)) over time for Det and Stc:024 patches from from example mouse 124 (KS test, p=0.002).

I) Performance (cross-validated Bayesian information criterion, cv-BIC) of integrator models fit to mouse PRT behavior and tested on held-out data for Stc patches (mean ± SEM, SR tests between models 1v3, 2v3, 3v4, p = 1.35e-5, p = 1.35e-5, p = 5.78e-5, n = 36 mice).

J) Model performance and best models as in **(I)** for Det patches (mean ± SEM, SR test between models 1v3, 2v3, 3v4, p = 2.07e-5, p = 1.92e-5, p = 5.05e-5, n = 36 mice).

K) Comparison of Model 3 drift rate parameter δ fit to Stc vs. Det trials (mean ± s.d., SR test, p = 5.85e-7, n = 36 mice).

L) Comparison of Model 3 reward sensitivity parameter ρ fit to Stc vs. Det trials (mean ± s.d., SR test, p = 4.22e-7, n = 36 mice).

M) Conceptual model of toggling between context-specific decision modes. Different configurations of the integrator process (parameters δ and ρ) produce computations for patch leaving decisions calibrated to different reward distributions.

Standard deviation (s.d.); Standard error of mean (SEM); Kolmogorov-Smirnov test (KS test); Wilcoxon signed-rank test (SR test); ***p<0.001; **p<0.01; *p<0.05; n.s. p>0.05.

was the best fit algorithm, both in terms of predictive performance (**Figures 2I-J**, left) and as the
best fit for each mouse (**Figures 2I-J**, right). We also observed similar differences in fits of δ and
ρ to Det vs. Stc behavior as in the FM task (**Figures 2K-M**). Male and female mice showed no
differences in any behavior in either task (**Figure S4**). These results establish that mice use similar
decision strategies across FM and VR task designs, opening a window to rigorously investigate
the structure of neural dynamics for flexible decision making.

Foraging contexts recruit distinct brain-wide neural activity states

To enable context-specific decision strategies, context information from the environment
must configure neural processes that instantiate or feed into decision computations (**Figure
3A**).[8,9,11] To investigate how environment context shapes neural activity, we performed high-
density *in vivo* electrophysiological recordings using Neuropixels[40] probes targeting a range of
brain regions likely to encode sensory context or decision dynamics (**Figures 3B-C, S5A-D**).
Following our previous study,[37] we densely sampled the frontal cortex, as well as the basal ganglia,
subcortical and thalamic nuclei, and sensory cortex spanning olfactory, somatosensory, visual and
associative domains (**Figure 3D**). After clustering and curation of spiking data (see Methods), we
included a total of 7804 units recorded from 47 sessions across 10 mice for analysis (**Figure S6A**).

We first explored large-scale activity patterns by splitting the dataset into seven "macro-
region" categories: dorsal frontal cortex (dFC), ventral frontal cortex (vFC), sensory cortex (SC),
hippocampal (HC), basal ganglia (BG), thalamic (THL), and subcortical (SUB) areas (**Figure 3D,
Table S1**). We visualized single-neuron activity in a heatmap, centering a 20-minute window at
the time of the context switch in each session (**Figure 3E**). Coding of sensory context (i.e. identity
of the Stc vs. Det environment) in single neurons was evident in every region, with persistent
changes in activity aligned to the context switch. To analyze the time-course of this state transition,
we measured population vector distances between timepoints following the context switch and a
preceding window (**Figure 3F**). Population activity exhibited a transient deviation which decayed
over 1-2 minutes and stabilized away from the pre-transition state. While this response was largest
in dFC, we observed similar dynamics in all regions, suggesting that environment contexts recruit
distinct activity states represented across the brain.

To examine more local dynamics structure, we plotted the correlation matrix of population
vectors for individual sessions (**Figure 3G**), which a revealed clear "block" pattern aligned to the
context switch, indicating context-specific activity patterns. We also projected population activity
onto principal components (**Figure 3H**), which confirmed a dynamical trajectory spanning two
distinct states. To rule out the possibility that context coding is spuriously driven by slow-timescale
fluctuations or drift artifacts, we quantified the population state change using the angle θ between
population vectors sampled across contexts and compared these to within-context samples.
Population vectors were nearly orthogonal ($\theta = 97°$) across contexts, and significantly more
aligned within ($\theta = 66°, 62°$), indicating distinct context-specific states (**Figure 3I**).

Finally, we characterized the finer regional distribution of context coding by identifying
single neurons that exhibit significant and persistent changes in activity at the context switch
("context coding" neurons). Overall, 29.6% of cells encoded context, and although non-uniform,
context information was widespread across the brain. In fact, while sensory areas (e.g. tenia tecta
and visual cortex) exhibited substantial context coding, likely attributable to visual and odor cues,

Figure 3

A Context controls network mode

B Probe track / Probe dye (DiI)

C Major insertion sites for Neuropixels 1.0 or 2.0 probes

D Number of neurons

E Context switch (Change VR environment)

F Context switch

G Context switch (VIS, RSC)

H

I

J *** (one-way anova)

K Shared subspaces / Orthogonal subspaces

Environment contexts recruit separate activity subspaces

L ***

M Context configurability heat-map

N

Figure 3. Foraging environments recruit distinct brain-wide activity states.

A) Cartoon illustrating configuration of neural network modes by control signals for context-dependent decision computations (concept adapted from Mante et al. 2013).
B) Example histological section showing location of a Neuropixels probe insertion through frontal cortex (dotted red line, probe tract, yellow, probe dye; see also **Figure S5** for examples).
C) Schematic showing major insertion locations with coordinates for acute recording experiments.
D) Summary of recording dataset; for each region, bar indicates the number of recorded neurons, and number of (sessions, mice) to the right (see **Figure S6** for full recording summaries).
E) Heat map showing activity of all recorded neurons (n = 7804, rows) over a 20-minute window centered around the context transition (change of virtual environment, t = 0, σ = s.d.).
F) Time course of population vector distances (euclidean norm) from pre-transition time window (-10 to -8 minutes) for pooled pseudo-populations of neurons from each macro-region.
G) Correlation matrix of population vectors from an example recording in visual/retrosplenial cortex. Block structure indicates stable within-context activity patterns that are reorganized at the time of context transition with greater within- vs. across-context similarity.
H) Principal component (PC) projections of activity from session in **(G)** showing transient deviation followed by persistent change in population activity at the time of context switch (marked with arrow; gray arrow indicates direction of time).
I) Angle measurements (θ in degrees) comparing similarity of within- vs. between-context population vectors (mean ± SEM, SR tests, p = 9.09e-8 [within context 1 vs. between], p = 3.13e-8 [between vs. within context 2], n = 47 sessions).
J) Single neuron context coding across sampled regions (one-way anova, p = 5.07e-4, n = 187 samples, see Methods for description of statistical tests).
K) Schematic illustrating relationship between Det- and Stc-context population activity subspaces. Canonical correlation analysis (CCA, also see **Figure S7** and Methods) its used to quantify distance between subspaces, which could be largely shared (top) or orthogonal (bottom).
L) Normalized (between - within context) subspace distance (the "chordal distance" $\sqrt{\sum \sin^2(\theta)}$ of angles between canonical vectors) for Det- vs. Stc-context subspaces (mean ± s.d., t-test, p = 7.23e-5, n = 47 sessions).
M) Left, correlation across regions between single-neuron context coding and subspace difference (chordal distance, linear regression, p = 4.95e-4, n = 32 regions); right, heat-map visualizing normalized chordal distance across brain regions (scaled min-max in blue-red colormap).
N) Correlation across mice between subspace difference (chordal distance) and flexible decision behavior, measured using the KS statistic of Stc vs. Det PRT distributions (linear regression, p = 0.0112, n = 10 mice).

Standard deviation (s.d.); Standard error of mean (SEM); Wilcoxon signed-rank test (SR test); ***p<0.001; **p<0.01; *p<0.05; n.s. p>0.05.

neurons in subcortical (e.g. septal nuclei) and most frontal areas were also modulated by context.
This indicates that environment contexts organize distributed activity states which could structure
context-specific dynamics for decision computations (**Figure 3A**).

Foraging contexts organize dynamics into separate activity subspaces

Defined in this way, "context coding" indicates sensitivity to environment identity, but does
not mean that the structure of neural dynamics is distinct across contexts, as might be expected for
different configurations of a decision process. For example, neural dynamics for each context
could lie within shared activity subspaces, indicating similar structure (**Figure 3K**). Alternatively,
they could be organized into separate or orthogonal subspaces, creating context-specific activity
modes.[7,22,41,42] To examine how neural dynamics are configured by context, we quantified the
alignment of Det vs. Stc activity subspaces. We used PCA to identify orthonormal bases (subspaces
S_{Det} and S_{Stc}) capturing within-context activity, and then applied canonical correlation analysis[43,44]
(**Figure S7A**, see Methods) to obtain vector pairs, and their canonical correlations, which capture
the alignment between S_{Det} and S_{Stc}. This method can be used to quantify the number of orthogonal
dimensions between subspaces or to measure overall subspace distance (e.g. the "chordal distance"
$\sqrt{\sum sin^2(\theta)}$ using angles θ from the canonical correlations, **Figure S7A-E**). We quantified these
for each session, subtracting within-context values to control for baseline drift.

We found that subspace distances (**Figure 3L**) and the number of orthogonal vectors
(**Figure S7H**) were significantly higher than within-context comparisons, suggesting that
environment contexts not only drive linear changes in firing rate (i.e. "context coding") but also
structure context-specific population dynamics. Interestingly, we observed substantial variability
in subspace distances across brain regions, which was itself correlated with context coding (**Figure
3M**) and much greater than expected from the presence of context coding (**Figure S7F-I**). This
indicates that brain regions which are more sensitive to sensory context also tend to exhibit more
context-specific dynamics, potentially supporting distinct functional modes for cognition and
behavior. Consistent with this idea, subspace distance was also correlated, across mice and
sessions, with decision flexibility in the task (**Figures 3N, S7J**), suggesting a connection between
dynamics modes and behavior. Thus, we next focused our investigation specifically on how
environment contexts shape decision activity for foraging.

Context-specific decision variables are encoded in separate neuronal subpopulations

Previous studies of context-dependent decisions have mostly observed context effects that
operate upstream of decision activity.[11–14,19] These can be thought of as "**input configuration**"
mechanisms (**Figure 4A**, top), which affect decision making by configuring inputs to a general
(i.e. context-invariant) decision process. Another model is that decision activity could be parsed
into distinct modes occupying different subspaces ("**subspace configuration**", **Figure 4A**,
bottom). Such orthogonalization of context-specific decision dynamics could offer several
computational and learning advantages (see Discussion).[7,8,22,24] In our framework, these models
describe the population geometry of decision activity, and in terms of cellular-level instantiation,
relate to the correlation structure of decision coding across contexts (**Figure 4B**). That is, while
correlated DV coding corresponds to aligned, context-invariant coding vectors (**Figure 4B**, right),

Figure 4

Figure 4. Subspace dynamics structure context-specific configurations of decision activity.

A) Illustration of input (top) vs. subspace (bottom) configuration models for how context structures integrator dynamics for foraging decisions. Input configuration preserves context-invariant decision dynamics, and subspace configuration exhibits context-specific, potentially orthogonal, Det vs. Stc decision coding vectors.

B) Cartoon of different possibilities for cellular-level coding structure underlying population geometry of decision activity. Orthogonal coding vectors (population level) correspond to uncorrelated or non-overlapping single-neuron decision coding.

C) Top, percentage breakdown of Stc-DV, Det-DV, both, and non-coding neurons from among all recorded; bottom, breakdown of positive and negative coding cells within each category. For cells encoding both Stc and Det DVs, "opp" indicates opposite-sign coding for Det vs. Stc.

D) Traces of trial-average activity for example Stc-DV cells (top) and population average on Stc or Det patch trials (bottom, n = 575 cells, σ = s.d.).

E) Traces of trial-average activity for example Det-DV cells (top) and population average (bottom, n = 708 cells, σ = s.d.).

F) Predictions of distinct activity dynamics for Stc-DV and Det-DV populations on patch trials based on parameter fits from integrator models (**Figures 2K-L**).

G) Left, average responses of Stc-DV (orange) and Det-DV (purple) cells at patch initiation (mean ± SEM); middle, comparison of response magnitude (t-test, p = 4.08e-5); right, comparison of ramping slope following response (t-test, p = 6.87e-7).

H) Left, average responses of Stc- and Det-DV cells at additional patch rewards beyond trial initiation (mean ± SEM); middle, comparison of response magnitude (t-test, p = 3.37e-7); right, comparison of ramping slope following response (t-test, p = 2.59e-4).

I) Left, average responses of Stc- and Det-DV cells around patch leave (mean ± SEM); right, comparison of ramping slope before patch leave (t-test, p = 0.0392).

J) Breakdown of single-neuron DV coding (fraction significant for either or both category at α = 0.05 - shuffle) across regions (one-way Anova, p = 0.0006).

K) DV coding similarity (|correlation r| between Stc-DV and Det-DV coding measures) for neurons from sampled regions (* indicates independent correlation for that region at α = 0.05).

L) Correlation (r) of Stc- and Det-DV coding across single neurons from two example regions (top, CP, Caudoputamen; bottom, M2, secondary motor cortex).

M) Correlation across regions between subspace difference (from **Figure 3M**) and DV coding as in **(J)**; grey lines indicate average of these metrics across all regions (linear regression, R^2 = 0.215, p = 7.56e-3).

N) Correlation across regions between subspace difference (from **Figure 3M**) and coding similarity as in **(K)**; grey lines indicate average metrics across all regions (linear regression, R^2 = 0.208, p = 8.71e-3).

Standard error of mean (SEM); ***p<0.001; **p<0.01; *p<0.05; n.s. p>0.05.

uncorrelated DV coding corresponds to divergent (potentially orthogonal) coding vectors, as pictured in the subspace configuration model (**Figure 4B**, left).

To test which model is best supported by our data, we first quantified the correlation between single-neuron activity and model-fit DVs against shuffled data to identify "DV-coding" neurons (see Methods). Many neurons (38.9% of all) encoded the integrator DV on either Stc patches, Det patches, or both (**Figure 4C**, top). Interestingly, while a small subset (5.7% of all) encoded DVs across both contexts, most cells were specific to either Stc (12.9%) or Det (20.3%) patches with significantly less overlap than within-context control comparisons (**Figure S8A**). Both positive and negative DV coding was present (**Figure 4C**, bottom). From this, we identified neurons that closely tracked the DV in one context yet were nearly unresponsive in the non-preferred context (**Figures 4D-E**, top), and averaging these yielded context-specific coding of the Stc or Det DV in completely non-overlapping subpopulations (**Figures 4D-E**, bottom).

This single-neuron analysis suggests that context signals organize decision activity into separate subpopulations, indicating subspace configuration (**Figure 4B**, left). However, this does not necessitate that context-specific DV neurons track distinct configurations of the decision, as predicted from the behavior analysis (**Figure 2M**). Based on our modeling results, we reasoned that neurons encoding configured DVs should exhibit distinct dynamics during patch leaving decisions (**Figure 4F**). Specifically, we hypothesized that: 1) Det-DV neurons exhibit faster ramping activity than Stc-DV neurons (reflecting a larger drift rate δ), and 2) Det-DV neurons exhibit larger reward responses (reflecting a larger reward sensitivity ρ).

To test these predictions, we examined activity of Stc- and Det-DV neurons during patch trials. At trial onset (**Figure 4G**), we observed negative responses in both populations, with a larger response in Stc-DV neurons reflecting the trial onset and trial-initial reward at t = 0s. Following this, both populations exhibited ramping, with Det-DV neurons ramping faster (**Figure 4G**, right). As this was confounded with trial initiation, we also examined responses around additional patch rewards (at t > 0s). As predicted, rewards after t = 0s elicited stronger responses in Det- vs. Stc-DV neurons as well as faster ramping after the reward (**Figure 4H**).

Finally, to test more directly whether ramping is faster in Det-DV cells, we examined activity leading up to the leaving decision (**Figure 4I**) in trials with no preceding rewards, eliminating the potential confound of reward responses. Det-DV neurons also exhibited faster ramping before leaving decisions. This shows that neurons encoding context-specific DVs exhibit distinct activity patterns reflecting distinct integrator parameter configurations: Det-DV neurons exhibit both faster ramping (δ) and larger reward deflections (ρ) during leaving decisions.

Activity subspaces structure context-specific decision coding

We next examined how coding of configured DVs was represented across the brain. Quantifying the fraction of neurons encoding either or both DVs, we found that most regions contained substantial single-neuron DV coding (**Figure 4J**), consistent with other reports of nearly brain-wide integrator dynamics during perceptual decisions.[45,46] Regions also exhibited broadly similar coding strength (**Figure S8B**).

However, while integrator dynamics are widespread, it is possible context-specific DV coding may be more localized.[8] For each region, we quantified DV similarity as the correlation

between single-neuron Stc- and Det-DV coding strength. Intriguingly, we found substantial regional heterogeneity in coding similarity (**Figure 4K**): while some regions exhibited highly correlated DV coding (e.g. Cautoputamen, **Figure 4L**, top), others exhibited completely uncorrelated, context-specific DV coding (e.g. secondary motor cortex, **Figure 4L**, bottom).

How might this heterogeneity arise? One possibility is that contextual configuration of activity subspaces decorrelates Stc- and Det-DV coding into orthogonal dimensions. To test this idea, we examined the relationship between subspace distance for each region ("chordal distance" from **Figure 3N**) and single-neuron coding properties **(Figure M-N)**. Strikingly, DV coding similarity was negatively correlated with subspace distance, suggesting that context-specific subspace configurations parse decision activity into uncorrelated modes. A subset of regions (bottom-right quadrant in **Figure 4N**), including parts of the dFC, exhibited strong subspace divergence and context-specific DVs. To further investigate this coding structure and its role in behavior, we thus narrowed our focus to dFC regions (including M2, ACC, and PL), which have been previously linked to foraging decisions and behavioral flexibility.[37,38,47–49]

dFC neurons encode orthogonalized context-specific decision variables

To examine decision coding in dFC, we first used a linear decoder to confirm that the Stc and Det DVs could be read out from population activity (**Figure 5A**). Decision coding within each context was stable enough to visualize single-trial DVs by projecting activity onto either Det- or Stc-DV coding vectors (**Figure 5B**). If dFC decision coding is orthogonal, as in the subspace configuration model, these coding dimensions should be: 1) geometrically perpendicular (i.e. angle $\theta \approx 90°$), and 2) mutually uninformative of the DV. We measured the angle θ between DV coding vectors and found these to be close to 90° and greater than within-context comparisons (**Figure 5C**), indicating orthogonality. Further, while within-context decoder performance was above chance for dFC neurons (**Figure 5A**), decoders could not predict DVs from the non-preferred context (**Figure 5D**), indicating that DV coding is context-specific and mutually uninformative.

As dFC receives direct input from sensory areas, we wondered whether orthogonal DV coding was shaped by task demands or was simply inherited from upstream signals conveying sensory context. We reasoned that if context specificity is independent of cognitive demand, then it should also be present in task-naïve mice experiencing the same sensory conditions. To test this, we recorded dFC neurons in a separate cohort of mice on the first 3 days of the task (**Figures 5E, S6B**) and, first, repeated the subspace alignment analysis on early (days 1-3) and experienced (days 10+) sessions. We observed a significant increase in subspace distances and orthogonal vectors over days (**Figures 5F-G**), suggesting that dFC dynamics are organized into context-specific subspaces with experience. We also quantified similarity in Stc vs. Det DVs using the correlation of single-neuron coding. Notably, while DV coding was initially correlated on day 1 (**Figure 5H**) across dFC, it was successively decorrelated across days and completely uncorrelated in later sessions (**Figure 5I**), consistent with the progressive divergence of activity subspaces (**Figure 5J**). Together, these results show that while task variables including context and decision coding are generally present in naïve animals, task experience organizes separate activity subspaces and orthogonalizes decision dynamics into context-specific modes.

Figure 5

dFC neurons learn orthogonal representations of the integrator decision variable

dFC activity causally contributes to decision flexibility

Figure 5. Orthogonalized integrator dynamics in dFC support decision flexibility.

A) Performance of linear decoders for Stc or Det decision variables (DVs, cross-validated correlation (*cv-r*) between predicted and model-fit DVs on held out trials) from dFC activity (SR tests, p = 0.0084, Stc, p = 0.0151, Det, n = 16 sessions).
B) dFC activity projected onto Det-DV or Stc-DV coding vectors (defined using regression coefficients, see Methods) from an example session (Mouse 116, session 10), showing population-level DV coding on individual Stc and Det trials.
C) Angle (θ in degrees) between Det-DV and Stc-DV coding vectors for dFC neurons (including secondary motor cortex M2, anterior cingulate ACC, and prelimbic cortex PL) or for each region separately (mean ± s.d., t-tests against θ = 90°, p = 0.1444 [dFC], p = 0.2244 [M2], p = 0.0276 [ACC], p = 0.3598 [PL]; 2-way anova, p = 0 (within vs. between), p = 0.7308, region, d.f. = 49).
D) Performance of Stc-DV and Det-DV linear decoders to predict DVs from patch trials in the other context (dashed lines indicate within-context performance from (**A**), SR tests, p = 0.8767, Stc, p = 0.7174, Det, n = 16 sessions).
E) Schematic of experiment to measure neural activity in task-naive mice; After habituation to running wheel and head-fixation, acute Neuropixels recordings targeting dFC were performed on the first 3 days of Phase 2 of the VR foraging task (Phase 1 was omitted).
F) Subspace difference (chordal distance $\sqrt{\sum \sin^2(\theta)}$) in dFC neurons from task-naive (days 1-3) and sessions (mean ± SEM, 2-way anova, p = 0.0083, region, p = 0.0124, time, d.f. = 57).
G) Number of orthogonal vectors (θ > 80°) in dFC neurons from task-naive (days 1-3) and trained sessions (mean ± SEM, 2-way anova, p = 0.0083, region, p = 0.0137, time, d.f. = 57).
H) DV coding similarity (correlation between Stc-DV and Det-DV coding metric *r*) across days (mean ± SEM, 2-way anova, p = 0.8174, region, p = 0, time, d.f. = 53).
I) Left, correlation across dFC neurons between Stc-DV and Det-DV coding in day 1 sessions (linear regression, R^2 = 0.282, p = 2.01e-9, n = 111 cells); right, correlation between Stc-DV and Det-DV coding in experienced sessions (days 10+, R^2=8.83e-4, p=0.271, n = 1372 cells).
J) Cartoon illustration of progressive subspace divergence in dFC across days and de-correlation/ orthogonalization of context-specific decision coding as shown quantitatively in (**I**).
K) Approach for dFC inactivation experiments. After 2 weeks of training in the VR task, mice were injected unilaterally in the M2 region of dFC with saline or Muscimol (alternating across four days) and then tested in the task each day (see Methods).
L) Comparison of overall patch residence time (median PRT for each mouse) behavior under Muscimol vs. saline control sessions (mean ± s.d., SR test, p = 0.9219, n = 10 mice).
M) Comparison of difference between Det and Stc PRT distributions (KS statistic) under Muscimol vs. saline control (mean ± s.d., SR test, p = 0.0195, n = 10 mice).
N) Det and Stc PRT distributions (CDF of P(Leave) over time) for example mouse 231 under saline (left) and Muscimol (right) showing collapse of context-specific PRT behavior with inactivation.

Standard deviation (s.d.); Standard error of mean (SEM; Wilcoxon signed-rank test (SR test).
***p<0.001; **p<0.01; *p<0.05; n.s. p>0.05.

dFC activity causally contributes to context-specific foraging decisions

This coding structure positions dFC as a potentially key node for contextual configuration of foraging decisions. Thus, we targeted dFC to test its causal contribution to decision flexibility in our task (**Figure 5K**). After training mice, we injected either saline or the GABA$_A$ receptor agonist Muscimol into the M2 region of dFC to locally silence neural activity. While inactivation had no effect on average PRTs (**Figure 5L**) or task engagement (**Figure S9A**), indicating that foraging behavior remained intact, we observed a significant decrease in decision flexibility (**Figures 5M, S9B**). In some mice, Stc and Det PRT distributions were indistinguishable under Muscimol (Mouse 231, **Figure 5N**), indicating complete abolition of context-specificity. This demonstrates that dFC activity, while apparently redundant for the core behavior, contributes to contextual configuration of foraging decisions.

dFC coding structure is preserved across virtual and naturalistic settings

The VR task enabled us to sample neural activity widely, revealing distributed activity states that organize context-specific, orthogonal decision coding, with a functional role in dFC. However, we noted that despite similar tendencies, behavior in VR was less consistent than in the FM task (**Figure S3**), possibly due to the artificiality of head fixation and/or virtual reality. Thus, we next sought to explore how this coding structure, identified under strong experimental constraint, might generalize to the naturalistic setting of free-moving foraging (**Figure 6A**).[26,27,29] For this, we returned to the FM task we initially developed (**Figures 1C, 6B**). We implanted chronic Neuropixels probes[50] in four mice targeting dFC, recorded across daily sessions in the task, and retained 822 dFC neurons from 16 sessions (Phase 2) for analysis (**Figures 6C, S6C**).

We first tested whether dFC neurons encoded context (i.e. Stc vs. Det chamber) in the FM task. Binary decoders could accurately predict Stc vs. Det context from population activity on both patch trials and inter-trial intervals (ITI, **Figure 6D**), indicating stable context coding. Visualizing neural activity in a heatmap centered around the "door open" time, when mice transition between contexts (**Figure 6E**), we observed a large-scale reorganization of population activity, with many neurons exhibiting sustained changes in firing rate. We confirmed this dynamics structure in individual sessions by examining population vector correlation matrices (**Figure 6F**) and trajectories (**Figure 6G**), which exhibited similar state transition patterns as in the VR task (**Figures 3G-H**). We also observed a transient excitation of population activity immediately on context transition (**Figure 6H**, top), which reflected a sum over bidirectional, single-neuron responses (**Figures 6H-6I**). We ruled out contributions from drift artifacts by comparing population vector angles within vs. between contexts (**Figure 6J**). Consistent with the VR task, these were significantly larger between contexts, indicating that activity states are more distinct than expected from baseline fluctuations. Context identity was encoded in single dFC neurons (**Figure 6K**) and drove separation of activity subspaces (**Figure 6L**).

To examine the structure of decision coding in the FM task, we repeated our previous analysis to identify neurons that encoded model-fit DVs. Overall, 33% encoded the DV (**Figure 6M**), the majority of which (91%) were context specific. Stc- and Det-DV neurons exhibited ramping activity and reward deflections specifically on Stc or Det patches (**Figure 6N**). While 9% of DV-coding neurons were context-invariant, the dominant pattern of context-specificity suggests subspace configuration. Consistent with this, DVs could be independently decoded from dFC

Figure 6

A Free-moving foraging task (FM)
Ecological validity
Naturalistic setting
Head-fixed virtual task (VR)
Mechanistic access/interpretability

B Phase 1 7d · Phase 2 14d expert state · Phase 3 4d
Door
Chronic Implanted Neuropixels 1.0 probe
Phase 2
Door open (context switch)
Stc context · Det context
or
30 min

C Example probe implant
DiI DAPI · M2 · ACC · Probe tract · Probe dye · PL · OFC

Recording summary
M2 — n=412 (12,3)
ACC — n=155 (13,4)
PL — n=255 (15,4)
IL — n=81 (9,3)
OLF — n=45 (4,1)
CP — n=91 (8,2)
Number of units 0 200 400 600

D Binary decoder (Stc vs. Det)
** **
Context decoder (cv-auROC)
1 0.8 0.6 0.4
Patches Shuffle ITIs Shuffle

Context coding and configuration of neural activity states

E Door open (context switch)
Firing rate (z-score) 2σ / -2σ
M2 (n=412)
ACC (n=155)
PL (n=255)
Time from door open (min) -10 -5 0 5 10

F Door open
Pop vector corr (r) 1 / -1
Time (min) 0 5 10 15 20 25 30

G Det context · Stc context
Context switch · t · Door open
PC 2 (score, a.u.)
-10min · +10min
PC 1 (score, a.u.) -10 0 10

H Door open
all dFC
Activity (σ) 0.6 0.4 0.2 0
switch + · switch -
0 -1
Time from door open (min) -5 -4 -3 -2 -1 0 1 2 3 4 5

I Switch-responsive neurons (%)
40 30 20 10
M2 ACC PL

J ** (θ≠90°)
Δ pop vector (θ°)
** · **
150 100 50
w/n ctx 1 · b/n ctx · w/n ctx 2

K n.s.
% context coding
100 50 0
M2 ACC PL

L **
Subspace distance √Σsin²(θ)
2 1 0 -1

Orthogonal decision coding for naturalistic foraging

M
14.0%
2.9%
16.1%
67.0%
Stc DV · Det DV · Stc & Det DV · Not DV coding (α=0.05 against shuffle)

N Stc-DV neurons
Stc patch · Det patch
2µL
Δ activity 1σ
Time on patch (s) -1 0 1 2 3 4

Det-DV neurons
Stc patch · Det patch
2µL
Δ activity 1σ
Time on patch (s) -1 0 1 2 3 4

O ** **
DV decoder (cv-r)
0.8 0.6 0.4 0.2 0
shuffle
Stc DV · Det DV

P **
DV coding vector angle (θ°)
120 90 60 30
90°
Across Within

Figure 6. Structure of dFC dynamics is preserved across experimental settings.

A) Cartoon illustration of tradeoffs in experimental approach. Return to the free-moving (FM) task design can test generalization of orthogonal decision coding and functionality with less constraint and greater ecological validity (green arrow).
B) Schematic of FM foraging task Phase 2, in which mice forage in Det and Stc contexts across two 15-minute blocks with one transition in the middle of the session through a controllable door.
C) Top, example histology showing location of chronic Neuropixels probe implant targeting the dorsal frontal cortex (dFC); bottom, summary of single neuron recordings from Phase 2. A total of n = 822 neurons were recorded from secondary motor cortex (M2), anterior cingulate (ACC), and prelimbic cortex (PL) across 16 sessions in 4 mice.
D) Performance of Fisher's Linear Discriminant decoders to predict Det vs. Stc context identity from dFC population activity during patch trials (left) or inter-trial intervals (ITI, right), quantified using the area under ROC (*cv-auROC*) curve for predicted vs. true labels on held-out trials (SR test, p = 0.0023, trials, p = 4.3778e-4, ITIs, n = 12 sessions).
E) Heat map of single-neuron activity centered around the time of door opening/context transition.
F) Correlation matrix of dFC population vectors sampled across an example session. Block structure indicates within-context similarity reorganized at the context switch (door open time).
G) Principal component (PC) projection of activity from example session in **(F)** showing persistent change in population activity upon context transition.
H) Top, transient response in mean population activity of dFC neurons centered around the door open time; bottom, averages of subpopulations that are excited (light blue, n = 93) or inhibited (dark blue, n = 41) transiently at context transition ("switch-responsive" neurons, permutation tests at $\alpha = 0.01$).
I) Breakdown of switch-responsive neurons across dFC regions (M2, ACC, PL, light, positive; dark, negative responses).
J) Angles (θ in degrees) comparing within- vs. between-context population vectors (t-tests against $\theta = 90°$, p = 0.0227 [within context 1], p = 6.2025e-7 [between], p = 9.2935e-9 [within context 2], n = 12 sessions).
K) Context coding neurons across dFC regions (one-way Anova, p=0.0362, n = 12 sessions).
L) Normalized (between context - within context) chordal distance $\sqrt{\sum \sin^2(\theta)}$ for Det- vs. Stc-context dFC subspaces (mean ± s.d., t-test, p = 0.0132, n = 12 sessions).
M) Breakdown of Stc-DV (orange), Det-DV (purple), Stc and Det DV (red), and non-DV coding neurons across dFC (n = 822 neurons, permutation test at $\alpha = 0.05$, as in **Figure 4C**).
N) Trial-averaged Stc and Det patch activity for example Stc-DV (left) and Det-DV (right) neurons.
O) Performance of linear decoders to predict Stc or Det DVs from dFC activity on held-out trials (SR tests against shuffle, p = 9.76e-4 [Stc trials], p = 4.88e-4 [Det trials], n = 12 sessions).
P) Angle θ between DV-coding vectors for across-context DVs vs. within-context control comparisons (SR test, p = 4.88e-4, n = 12 sessions).

Standard error of mean (SEM); Wilcoxon signed-rank test (SR test). ***p<0.001; **p<0.01; *p<0.05; n.s. p>0.05.

activity (**Figure 6O**), and the angle between Stc-DV and Det-DV coding vectors was close to 90°
(**Figure 6P**), indicating orthogonal geometry. This shows that context-specific configuration of
orthogonal integrator dynamics is a general neural solution for flexible foraging decisions,
preserved across experimental settings with different levels of constraint and ecological validity.

Decision dynamics are reconfigured immediately on context transitions

As most task designs impose block structure or rule cues to indicate context, it remains
unclear how neural dynamics are reconfigured to enable flexible behavior when animals are free
to adjust cognitive modes of their own volition, as would be the case in the natural world. We took
advantage of our task design to address this, continuing to record dFC neurons into Phase 3 of the
task ("volitional transitions", **Figures 7A-B**) and retaining 712 neurons across 16 sessions from
the same mice (**Figure 7B**). We again confirmed encoding of context identity (**Figure 7C**) and
context-specific DVs (**Figure 7D**) from dFC activity, indicating that these variables are stably
represented even when block structure is removed and mice are unconstrained. Despite switching
contexts repeatedly (2.25 times/min, **Figure S1K**), orthogonal DV coding was preserved (**Figure
7E**), suggesting that mice "toggle" decision modes on context transitions and do not require blocks
of repeated trial types to stabilize coding structure.

To explore this more directly, we analyzed neural population trajectories (visualized with
PC projections) during patch trials within and across contexts. Taking the Euclidean distance
between trajectories (ΔTraj, **Figure 7F**), we found that average Stc vs. Det trajectories were more
distinct than same-context trajectories (**Figure 7G**), consistent with observed context-specific DV
coding (**Figures 5C, 6P, 7E**). To see the immediate effect of context transitions, we examined
trajectories for sequential within-context (e.g. Det→Det) or across-context (e.g. Det→Stc) trial
pairs. Visualization in example sessions suggested that single-trial trajectories were more distinct
across transitions (**Figure 7H**), and quantitatively, ΔTraj for cross-context trials sequences was
greater than within-context comparisons (**Figure 7I**). This indicates that context-specific neural
dynamics for foraging decisions are immediately reconfigured upon volitional context switches
and that, once learned, these decision modes are flexibly and rapidly deployable.

dFC is necessary for integrator reconfiguration and volitional transitions

The striking consistency of orthogonal DV coding across levels of experimental constraint
led us to hypothesize that dFC is generally required for decision flexibility, even without imposed
task structure. To test this, we repeated the Muscimol inactivation experiment to silence dFC
neurons during Phase 3 of the FM task (**Figure 7J**). Like the previous result (**Figure 5M**), decision
flexibility was significantly reduced under Muscimol (**Figures 7K-L**). In the integrator model, this
behavioral pattern should reflect a collapse of parameter configurations which underlie distinct
decision modes. Indeed, fitting the model as before, we found that δ and ρ were reduced in
magnitude under dFC inactivation (**Figure 7M**), and that the difference in parameter fits across
Stc vs. Det patches was significantly smaller (**Figure 7N**). Plots of example DV rollouts confirmed
the interpretation that Muscimol disrupted context-specific DV configurations (**Figure 7O**).

Finally, we observed some additional effects of dFC inactivation in the FM task. While
mice still performed well, they were significantly less likely to make volitional patch or context

Figure 7

dFC dynamics exhibit rapid reconfiguration for decision flexibility

dFC activity is necessary for integrator configurations under volitional mode switching

Figure 7. dFC supports volitional mode switching and rapid configuration of foraging decisions.

A) Schematic of chronic Neuropixels recording in Phase 3 of the free-moving (FM) foraging task when mice are free to switch environments volitionally through the open door.
B) Cartoon of volitional transitions in Phase 3 and recording summary (n = 712 dFC neurons, 16 sessions, 4 mice).
C) Performance (cross-validated *auROC*) of binary decoders for context identity on patch trials (left) and inter-trial interval (ITI) periods (right) (mean ± SEM, SR tests against shuffle, p = 4.88e-4 [trials], p = 0.0015 [ITIs], n = 12 sessions).
D) Performance (cross-validated *r*) of Stc-DV and Det-DV decoders compared to shuffle controls (mean ± SEM, SR tests against shuffle, p = 4.88e-4 [Stc DV], p =4 .88e-4 [Det DV], n = 12 sessions).
E) Angle θ between DV-coding vectors for across-context DVs vs. within-context control comparisons (SR test, p = 0.0049, n = 11 sessions, see Methods).
F) Schematic illustrating neural trajectory analysis and quantification of ΔTraj metric (the sum of time point-matched euclidean distances between patch trial trajectories).
G) Comparison of ΔTraj across within- and between-context trial-average trajectories (U tests, p = 4.77e-4 [within Det v.s. within Stc], p = 0.0035 [within Stc v.s. across contexts]).
H) Average of trajectories for Det→Det (left) and Det→Stc (right) sequential trial pairs from one example session (Mouse 251, session 17); arrow indicates direction of time evolution.
I) Comparison of session average ΔTraj for within- vs. between-context sequence pairs as visualized in **(H)** (median ± SEM, SR test, p = 9.76e-4, n = 12 sessions).
J) Schematic of dFC Muscimol inactivation experiments and timeline.
K) Comparison of Det vs. Stc:012 PRT distribution KS statistics for saline vs. Muscimol sessions (SR test, p=0.0398, n = 13 mice).
L) Det and Stc:012 patch PRTs (CDF of P(Leave) over time) from example Mouse 220 under saline (left) and Muscimol (right) sessions, showing collapsed PRT distributions and reduced KS statistic under dFC inactivation.
M) Scatters of average parameter fits (ρ and δ) for Stc (orange) and Det (purple) patches under saline (grey outline, circles) and Muscimol (green outline, squares, mean ± SEM).
N) Comparison of parameter differences (euclidean distance of z-scored ρ and δ values) for saline vs. Muscimol sessions (SR test, p=0.0398, n = 13 mice).
O) DV rollouts for a Det patch reward sequences (reward at t = 0, 1, 2s) using parameter fits from example Mouse 219 for saline (top) or Muscimol (bottom) sessions.

Standard error of mean (SEM); Mann-Whitney U test (U test); Wilcoxon signed-rank test (SR test). ***p<0.001; **p<0.01; *p<0.05; n.s. p>0.05.

transitions under Muscimol (**Figures S9C-D**) and exhibited longer PRTs in general (**Figure S9E**).
While this may indicate perseverative behavior, it is also consistent with the established role for
dFC in tracking task performance and regulating explore-exploit modes,[18,51–53] which may be
relevant for context transitions in our foraging task. Longer PRTs under Muscimol are also broadly
consistent with frontal cortex involvement in biasing waiting behavior and sustaining persistent
ramping dynamics underlying decision computations.[38,45,47,48,54–57]

Discussion

The question of how neural processes are contextually reconfigured is fundamental to our
understanding of flexible cognition and behavior.[1,4] We investigated how the brain organizes neural
dynamics to support context-dependent decision making in the ecological setting of patch foraging.
Our approach stands in contrast to classical task designs, which require animals to learn artificial
rule cues for perceptual choices and mostly locate context effects upstream of decision dynamics,
operating as "input configuration" mechanisms that shift attention or gate sensory inputs.[11–13,17,19]
Instead, we observed a strikingly different neural solution for context-dependency: different
foraging environments recruit brain-wide activity states, which organize decision dynamics into
orthogonal modes instantiated in separate neural subpopulations. These neurons encoded two
distinct configurations of an integrator decision variable (DV) that computes patch leaving
decisions through a process akin to bounded evidence accumulation. In the dorsal frontal cortex
(dFC), decision modes are rapidly toggled as mice transition between contexts, and activity in dFC
is required for expression of context-specific decision strategies.

Configurations of neural integrator dynamics explain context-specific decision behavior

Recent efforts to interpret classical foraging problems, such as the patch leaving problem,
through the lens of decision theory have led to new proposals for biologically realistic
computational models and neural mechanisms.[34,35] We recently demonstrated that foraging
decisions in mice are well described by an integrator process that weighs time and rewards as
decision evidence.[37] Ours and other studies have also identified neural dynamics encoding
integrator DVs in the frontal cortex of rodents.[38,45,47,48,58,59] This burgeoning literature begins to
clarify the algorithmic logic and neural implementation of foraging decisions,[28,29,60] raising timely
questions about how the brain can adapt or reconfigure neural dynamics to express flexible
cognitive strategies for ecological problems. For example, different configurations of a neural
integrator can track time investment, time since last reward, reward rate, and reward quantity
(**Figure 1J**, Models 1-4) for patch foraging.[34,61]

How do animals adapt decision strategies across different foraging environments? Here,
we addressed this by studying mice foraging for water rewards across two environment contexts
defined by distinct reward dynamics: a Stochastic (Stc) context, yielding decaying probabilistic
rewards, and a Deterministic (Det) context yielding a fixed number of rewards (**Figure 1A-C**). An
intuitive initial hypothesis is that mice may switch between a "reward rate" computation (i.e.
"Model 3") for Stc patches, and a "reward accumulation" computation (i.e. "Model 4") for Det
patches. Instead, we found that mice solve this problem using the *same algorithm* (**Figures 1K-L,**
2I-J) under *different parameter configurations* (**Figures SO-P, 2K-L**). Model fits to Det patches

had larger values for both the drift rate $\boldsymbol{\delta}$ and reward sensitivity $\boldsymbol{\rho}$, and these were reflected in
neural correlates of DVs (**Figure 4G-I**). In our task, many mice adapt decisions to leave Det
patches soon after the final reward, suggesting a "counting" algorithm, yet integrator models and
neural DVs did not exhibit features of reward accumulation (i.e. $\boldsymbol{\rho} > 0$, "Model 4"). This suggests
that reconfiguration of a common evidence accumulation computation may be a general solution
to calibrate foraging decisions to different reward dynamics.[34,61]

What mechanisms underlie changes in configuration of the neural integrator? The drift rate
$\boldsymbol{\delta}$ in our model effectively captures the weight of passing time as evidence *for* patch leaving, and
is analogous to the same parameter in drift-diffusion models for perceptual decisions, where it can
be modulated by the quality of continuous sensory evidence, confidence, and value.[36,37,62–65]
Mechanistically, this may be driven by top-down cortical signals which encode drift rates,
organization of recurrent interactions sustaining persistent activity, or neuromodulator influences.
For patch foraging, promising candidates include frontal or thalamic inputs to dFC, which bias
waiting times for reward seeking[48,54] and modulate recurrent dynamics,[14,66] and 5-HT signaling,
which has been linked to behavioral "patience" for uncertain rewards.[67–69]

The (negative) reward sensitivity $\boldsymbol{\rho}$ in our model can be thought of as a weight of pulsatile
evidence *against* patch leaving. As rewards have rapid influence on integrator dynamics (**Figure
4H**), this signal is likely mediated, at least in part, by detection of water in the mouth through
somatosensory pathways. The gain (or sign) of this signal onto the integrator could be modulated
by top-down sensory gating mechanisms and/or local interneuron interactions. In general, what
mechanisms control the parameters of integrator dynamics for foraging, and how these are learned
and calibrated from reward feedback, are important and ripe domains for future investigation.

489 Orthogonal coding of context-specific integrator dynamics in separate neuron populations

A deeper aim of our study was to understand how neural dynamics are structured to support
context-dependent decision making. To express flexible (i.e. context-specific) behavior, context
signals must organize different functional modes which alter how sensory information (e.g. water
rewards) affect the decision process (**Figures 3A, 4A**).[8,9] As previous studies of rule-based
perceptual decisions have shown, this can be achieved mechanistically by adjusting how sensory
signals are processed (e.g. through sensory gating or attentional shifting) or how they perturb
decision dynamics.[11,12,16,17,19] Our framework treats such mechanisms which operate upstream of
the decision process as aspects of "input configuration" (**Figure 4A**). Indeed, this is important for
context-dependency, especially for perceptual choice problems, and has recently been described
in a unified model which can explain individual variability in rule-based sensory discrimination as
idiosyncratic expression of different input configuration mechanisms.[19]

However, the mechanistic details of input configuration are agnostic as to how decision
activity itself is structured. Alternatively (or additionally), the brain could use a "subspace
configuration" strategy to structure different instantiations of a decision process to operate in
different contexts.[7,20,22,70] Subspace coding has been explored and studied in other domains,
including to decorrelate sensory representations,[22,71] for cortical communication,[20] and to stay
premotor activity from motor expression.[45,72] Context-specific DVs represented in orthogonal
coding subspaces could be separately tuned without interference, read out independently by
downstream circuits, and flexibly toggled by control signals.[7,24,41,73]

In contrast to the input configuration solutions for perceptual choice tasks, we find evidence of widespread subspace configuration and orthogonal coding of context-specific DVs in the mouse brain. Furthermore, this orthogonal DV coding is instantiated at the cellular level in largely separate neuron subpopulations. In principle, this simple structure allows the brain to calibrate and toggle context-specific decision dynamics by activating subsets of neurons in different contexts, expressing and exposing them selectively to teaching signals and plasticity.

Why do we observe orthogonal, rather than context-invariant, integrator dynamics? One factor is that neural solutions for flexible computation are likely task dependent.[8] For perceptual choice tasks which have dominated previous research, filtering or gating sensory inputs is likely sufficient to embed context contingencies, and this may be especially so if context/rule cues are transient arbitrary stimuli (e.g. lights or tones) or must be inferred from task performance.[11–13,18,19] In our foraging task, however, context is not only continuously given through stable cues, but the notion of context is also less abstract, and perhaps more ecologically intuitive (for a mouse) as a spatially extended environment.[26] We speculate that when "context" is synonymous with "environment," which carries intrinsic motivational meaning (as in a place to explore and forage), the brain treats it differently than an artificial rule cue, and uses it to organize brain-wide activity states that can shift behavior and cognition between discrete modes. Indeed, we found that environment context was not only encoded in sensory areas but widely throughout the brain, including in frontal, subcortical, and thalamic structures.[8,71,74,75] Beyond linear context coding, foraging environments also recruit separate activity subspaces in many regions, which organize orthogonal coding of integrator dynamics (**Figure 4N**). In terms of neural activity, these observations are consistent with a picture of context-sensitivity and "configurability" as distributed or graded rather than localized.[8] At the same time, some circuits, including in the frontal cortex, may be of particular functional importance for regulating behavioral and cognitive modes.[1,3]

Placement of dFC in a network of configurable circuitry

In this study, we sampled widely across 32 brain regions to characterize how context shapes neural dynamics and integrator DVs (**Figures 3-4**). We then focused on the dFC to analyze orthogonal coding structure (**Figure 5**) and test causality and consistency across experimental settings (**Figures 6-7**). We observed orthogonal coding of context-specific DVs in dFC regions (including M2, ACC, and PL) and found that inactivation of M2 disrupted decision flexibility. Interestingly, unilateral M2 inactivation did not disrupt foraging in the VR task, and only mildly affected engagement in the FM task, suggesting that M2 may be redundant for the core decision process. This is perhaps not surprising given the distributed nature of integrator dynamics for decision and movement timing[45,46,57,76,77] (**Figure 4J**), which are sustained through multi-region recurrent interactions. However, this result suggests a more specific function for M2 in *selecting context-specific cognitive modes*. Indeed, previous studies support a role for M2 in shaping flexible behavior and action policies based on sensory contingencies.[78–80] The anatomy also supports this picture, as M2 receives direct input from sensory areas and projects broadly to motor control circuits in the cortex, midbrain, and basal ganglia.[47,49,57,58,76]

However, M2 (and dFC in general) do not exist in isolation, and their contributions to behavior must depend on interactions with a wide network of interconnected circuits. Indeed, our subspace alignment and coding analyses identified several other regions outside dFC that exhibit context-specific dynamics and DV coding (**Figure 4N**), including sensory and subcortical areas,

553 raising questions about how these interact, how distributed network dynamics are structured, and
554 whether M2 or dFC play a specialized role in contextually shaping network activity. Leveraging
555 high-density recordings[40,50] to sample neural populations across regions simultaneously, including
556 with targeted perturbations in specific circuits, will be illuminating.

Methodological consideration of ecological validity and complementary task designs

Finally, our approach respects the limitations of both controlled/reduced and unconstrained
naturalistic experiments (**Figures 2A, 6A**), as well as the importance of ecological relevance to
drive interpretable behavior.[26,27,39] Motivated to investigate the neural basis of flexible cognition,
we designed tasks that encourage decision flexibility for self-paced foraging instead of perceptual
discrimination as in classical designs.[28,29,39] In this ecological setting, mice naturally calibrated
context-specific decision strategies within days instead of weeks or months. Practically, this
allowed us to characterize behavior across dozens of animals and widely sample population
activity with single-neuron resolution, leading to new insights that have eluded previous studies.

Our approach also demonstrates how complementary task designs can be used
synergistically to elucidate and validate models of circuit function. We began with a naturalistic
foraging task and studied behavior to understand the computational logic driving expression of
flexible decision strategies. We then used a virtual, head-fixed task to screen neural activity across
32 brain regions and characterize coding structures supporting decision flexibility. Finding clear
evidence for subspace configuration and orthogonal integrator dynamics, we returned to the free-
moving setting to study how this neural solution is used under conditions that better match real-
world experience (e.g. for volitional and repeated mode switching). This dual-task approach not
only gave us multiple perspectives on the problem, but also allowed us to rigorously replicate and
test consistency of our results across experimental conditions. Taken together, our work identifies
orthogonal integrator dynamics as a general neural mechanism for context-dependent computation
enabling rapid toggling between cognitive modes for decision making.

Figure S1

Figure S1. Additional analysis of behavior through foraging task progression.

A) Left, reward probabilities over time for stochastic patches. Each trial yields one reward at t=0s, and then one each second after with decaying probability starting with random P_i = 0.66, 0.81, 0.93, or 0.99. Right, cumulative expected reward over time for stochastic patches.
B) Number of patch trials performed by naive mice in the first session on Phase 1 of the foraging task (each dot one mouse, mean ± s.d., n = 16 mice).
C) Latency (minutes) to the first patch trial in the first session of the task (each dot one mouse, mean ± s.d., n = 16 mice).
D) Fraction of trials among the first 5 in the first session (left) or the last 5 in the first session (right) in which mice collected all three deterministic rewards (data pooled across mice).
E) CDF of leaving probabilities for trials on each session (days 1-10) in Phase 1 (data pooled across mice).
F) Fraction of trials where mice collected all three rewards for each session (days 1-10) in Phase 1.
G) CDF of leaving probabilities for deterministic trials on day 1 in Phase 1 (cyan), day 10 in Phase 1 (purple), late (last 4 days) in Phase 2 (grey), and Phase 3 (green) showing similar PRT distributions across all phases.
H) Average number of rewards collected per trial for Det and Stc trials in Phase 2 and Phase 3 (each dot one mouse, mean ± s.d., SR test, p = 8.86e-5 (P2 Stc vs. Det), p = 8.86e-5 (P3 Stc vs. Det), p = 0.0522 (P2 Det vs. P3 Det), n = 20 mice).
I) Choice probability for Det patches in Phase 3 of the task (each dot one mouse, mean ± s.d., t-test > 0.5, p = 2.4e-6, n = 20 mice).
J) Patch transition matrix for Det and Stc patches in Phase 3 of the task (data pooled across mice).
K) Volitional context transitions per minute in Phase 3 (each dot one mouse, mean ± s.d.).

Wilcoxon signed rank test (SR test); ***p<0.001; **p<0.01; *p<0.05; n.s. p>0.05.

Figure S2

Figure S2. Additional analysis of decision behavior and context effects.

A) Mean patch residence times (PRT, seconds) for hidden Stc trial types with different reward probabilities P_i = 0.66, 0.81, 0.93, or 0.99 (**Figure S1A**, mean ± s.d.; SR test, p = 8.85e-5 (0.66 vs. 0.99), p = 0.004, (0.81 vs. 0.99), p = 0.0479 (0.93 vs. 0.99), n = 20 mice).

B) Diagram showing difference metrics for comparing PRT distributions. The Kolmogorov-Smirnov (KS) statistic measures the maximum vertical distance δ between CDFs (left), and ΔCDF (also the 1D Wasserstein metric) measures the total/average absolute difference between CDFs (right).

C) KS statistic comparing Det vs. Stc:012 PRT distributions across early (days 1-5), middle (days 6-10), and late (days 11+) Phase 2 progression and Phase 3 (SR test, p = 0.0438 (early P2 vs. P3), p =0 .3703 (late P2 vs. P3), n = 20 mice).

D) Cartoon showing example of a Stc:012 (trial t) followed by Det (trial t+1) sequence in Phase 3.

E) P(Leave) CDF comparing PRT distributions for Stc:012→Det trial sequences illustrated in **(D)** (KS test, p = 8.83e-8, n = 271 trials).

F) Scatter plot showing PRTs for Stc:012→Det trial sequences in Phase 3 and histograms of PRT distributions.

Kolmogorov-Smirnov test (KS test); Wilcoxon signed-rank test (SR test); ***p<0.001; **p<0.01; *p<0.05; n.s. p>0.05.

Figure S3

Figure S3. Additional details of virtual foraging task and behavior.

A) Left, reward probability over time for stochastic patches. Each trial yields one reward at t=0s, and then one reward each second with exponentially decaying probability starting at t = 1s with random P_i = 0.15, 0.255, 0.575, 0.947. Right, cumulative expected reward over time for stochastic patches.
B) Patch engagement rate (trials per minute) for virtual (VR) foraging task (each dot one mouse, mean ± s.d.).
C) Median patch residence time (PRT) for virtual foraging task (each dot one mouse, mean ± s.d.).

Figure S4

Free-moving (FM) task

Virtual (VR) head-fixed task

Figure S4. Lack of sex differences in foraging behavior and decision flexibility.

A) Comparison of patch engagement rate (trials per minute) between male vs. female mice in the free-moving (FM) foraging task (mean ± s.d., U test, p = 0.1965, n = 9 males, n = 11 females).
B) Comparison of median patch residence time (PRT, seconds) between male vs. female mice in the FM task (mean ± s.d., U test, p = 0.9697).
C) Comparison of KS statistic for Stc:012 vs. Det patch PRTs between male vs. female mice in the FM task (mean ± s.d., U test, p = 1).
D) Comparison of ΔCDF for Stc:012 vs. Det patch PRTs between male vs. female mice in the FM task (mean ± s.d., U test, p = 0.4941).
E) Correlation between KS **(C)** statistic and ΔCDF **(D)** for all mice in the FM task (linear regression, R^2 = 0.69, p = 4.73e-6, n = 20 mice).
F) Comparison of patch engagement rate between male vs. female mice in the virtual (VR) foraging task (mean ± s.d., U test, p = 0.9296, n = 21 males, n = 19 females).
G) Comparison of median patch residence time between male vs. female mice in the VR task (mean ± s.d., U test, p = 0.5513).
H) Comparison of KS statistic for Stc vs. Det patch PRTs between male vs. female mice in the VR task (mean ± s.d., U test, p = 0.203).
I) Comparison of ΔCDF for Stc vs. Det patch PRTs between male vs. female mice in the VR task (mean ± s.d., U test, p = 0.5157).
J) Correlation between KS statistic **(H)** and ΔCDF **(I)** for all mice in the VR task (linear regression, R^2 = 0.77; p = 1.16e-13, n = 40 mice).

For all plots, each dot is one mouse (blue, males; red, females). Mann-Whitney U test (U test); ***p<0.001; **p<0.01; *p<0.05; n.s. p>0.05.

Figure S5

A

Mouse 160; Bregma +1.35 AP

Sampled regions:
Secondary motor cortex (M2)
Anterior cingulate (ACC)
Striatum (STR)
Nucleus Accumbens (NAC)

B

Mouse 257; Bregma -0.3 AP

Sampled regions:
Somatosensory areas (SS)
Caudoputamen (CP)
Nucleus Accumbens (NAC)

C

Mouse 283; Bregma -1.85 AP

Sampled regions:
Retrosplenial cortex (RSC)
Hippocampus CA1
Dentate Gyrus (DG)
Habenula (HAB)
Mediodorsal thalamus (MDT)
Centromedial thalamus (CMT)

D

Mouse 283; Bregma -2.65 AP

Sampled regions:
Posterior parietal cortex (PPC)
Hippocampus CA1
Dentate Gyrus (DG)
Lateral geniculate (LGN)

Figure S5. Examples of Neuropixels probe insertions.

A) Example histology (Mouse 160) showing a 1-shank Neuropixels probe insertion targeting frontal and basal ganglia regions (M2, ACC, STR, NAC).
B) Example histology (Mouse 257) showing a probe insertion targeting sensory cortex and basal ganglia regions (SS, CP, NAC).
C) Example histology (Mouse 283) showing a probe insertion targeting sensory cortex, hippocampal, subcortical and anterior thalamus regions (RSC, CA1, DG, HAB, MDT, CMT).
D) Example histology (Mouse 283) showing a probe insertion targeting sensory cortex, hippocampal, and posterior thalamus regions (PPC, CA1, DG, LGN).

For all images, blue = DAPI, orange/red = probe dye (DiI/DiD), green dashes = probe tract); insertion coordinates in millimeters (mm) from Bregma. See **Table S1** for more detail on region identification.

Figure S6

Acute recordings in virtual (VR) task (n = 10 mice, 47 sessions)

A

Mouse	Exp #	Probe	dFC			vFC			SC						HC			BG			THL								SUB						
			M2	ACC	PL	IL	OFC	DP	TT	OLF	SS	VIS	PPC	RSC	DG	CA1	CA3	STR	CP	NAC	LGN	MDT	CMT	VPT	PNT	PVT	PFS	LSR	LSV	MSP	PCN	HAB	HYP	MID
83	16	1 shank	.	31	.	.	.	.	.	.	.	.	.	.	.	.	.	31	.	78	.	.	.	.	.	.	.	51	4	.	.	.	.	.
83	17	1 shank	24	123	.	.	.	.	.	.	.	.	.	.	.	.	.	6	.	.	.	.	.	.	.	.	.	24	.	.	.	.	.	.
83	18	1 shank	3	15	.	.	.	.	.	.	.	.	.	.	.	.	.	15	.	.	.	.	.	.	.	.	.	44	23	.	.	.	26	.
83	19	1 shank	9	15	.	.	.	.	.	.	.	.	.	.	.	.	.	3	.	.	.	.	.	.	.	.	.	51	17	.	.	.	.	.
88	14	1 shank	.	.	1	.	.	3	10	2	.	.	.	.	.	.	.	13	.	.	.	.	.	.	.	.	.	17	.	34	.	.	.	.
88	15	1 shank	.	12	4	9	.	46	41	1	.	.	.	.	.	.	.	26	.	.	.	.	.	.	.	.	.	59	.	10	.	.	.	.
88	16	1 shank	.	4	25	31	.	30	27	4	.	.	.	.	.	.	.	31	.	.	.	.	.	.	.	.	.	41	.	14	.	.	.	.
88	17	1 shank	.	9	.	.	.	.	.	.	.	.	.	.	.	.	.	27	.	.	.	.	.	.	.	.	.	60	21	.	.	.	59	.
88	18	1 shank	.	13	.	.	.	.	.	.	.	.	.	.	.	.	.	20	.	.	.	.	.	.	.	.	.	24	37	.	.	.	13	.
88	19	1 shank	.	53	.	.	.	.	.	.	.	.	.	.	.	.	.	12	.	.	.	.	.	.	.	.	.	58	9	.	.	.	42	.
116	10	1 shank	96	2	14	.	25	.	.	19	.	.	.	.	.	.	.	12	.	110	.	.	.	.	.	.	.	.	.	.	.	.	.	.
116	11	1 shank	57	8	22	.	44	.	.	24	.	.	.	.	.	.	.	32	.	80	.	.	.	.	.	.	.	.	.	.	.	.	.	.
116	12	1 shank	.	12	5	.	.	3	13	1	.	.	.	.	.	.	.	48	.	.	.	.	.	.	.	.	.	129	.	19	.	.	.	.
116	14	1 shank	.	53	95	33	.	55	40	4	.	.	.	.	.	.	.	31	.	2	.	.	.	.	.	.	.	44	.	.	.	.	.	.
116	15	1 shank	203	.	.	.	30	.	.	30	.	.	.	.	.	.	.	20	.	24	.	.	.	.	.	.	.	.	.	.	.	.	.	.
119	13	1 shank	.	5	140	181	.	38	54	8	.	.	.	.	.	.	.	26	.	5	.	.	.	.	.	.	.	.	.	.	.	.	.	.
119	14	1 shank	2	50	141	163	.	26	64	4	.	.	.	.	.	.	.	25	.	2	.	.	.	.	.	.	.	.	.	.	.	.	.	.
119	15	1 shank	39	42	132	129	.	66	58	5	.	.	.	.	.	.	.	11	.	.	.	.	.	.	.	.	.	15	.	.	.	.	.	.
257	19	4 shank	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	59	.	.	.	.	6	31	.	.	.	.	2	.	.
257	20	4 shank	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	106	.	.	.	.	8	.	.	.	.	20	7	.	42
257	21	4 shank	.	.	.	.	.	.	.	.	79	7	73	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
257	22	1 shank	.	.	.	.	.	.	13	.	.	.	.	.	.	.	.	.	182	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
258	18	4 shank	.	.	.	.	.	.	.	.	.	.	40	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
258	19	4 shank	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	15	.	.	.	.	.	.	.	.	.	.	.	.	1
258	20	4 shank	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	30	.	.	.	13	16	.	.	.	.	17	.	.	.
258	22	1 shank	.	.	.	.	.	.	25	.	.	.	.	.	.	.	.	.	48	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
261	17	4 shank	.	.	.	.	.	.	.	.	.	.	145	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
261	19	4 shank	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	71	.	.	.	12	11	.	.	.	.	.	.	.	13
261	20	4 shank	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	32	.	2	.	.	15	.	.	.	.	.	.	.	27
261	21	4 shank	.	.	.	.	.	.	.	27	.	21	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
261	22	1 shank	.	.	.	.	.	.	7	.	.	23	.	.	.	.	3	111	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
262	18	4 shank	.	.	.	.	.	.	.	.	.	23	.	1	1	4	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
262	19	4 shank	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	77	10	.	.	2	1	.	.	.	.	.	.	.	.
262	20	4 shank	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	34	.	.	.	.	29	.	.	.	.	.	.	.	.
262	21	4 shank	.	.	.	.	.	.	.	2	5	17	.	10	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
262	22	1 shank	.	.	.	.	.	.	25	.	.	.	.	.	.	.	.	156	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
283	17	4 shank	.	.	.	.	.	.	7	.	.	.	.	.	.	.	.	143	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
283	18	4 shank	.	.	.	.	.	.	.	.	.	28	.	24	26	20	.	.	.	103	3	.	.	.	.	.	.	.	.	.	.	.	.	.
283	19	4 shank	.	.	.	.	.	.	.	.	.	.	.	12	74	.	.	.	.	.	.	.	.	203	.	.	.	.	.	.	.	.	.	.
283	21	4 shank	.	.	.	.	.	.	.	.	.	26	.	.	.	.	.	.	.	42	25	.	.	1	.	.	.	.	.	.	.	.	.	.
283	22	4 shank	.	.	.	.	.	.	.	.	.	4	.	20	.	9	.	.	.	5	55	.	.	.	.	.	.	.	.	.	12	.	.	.
283	23	4 shank	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	47	.	.	.	10	.	.	.	.	.	.	6	.	.	.
283	24	4 shank	.	.	.	.	.	.	.	.	.	.	.	7	23	.	.	.	.	36	.	.	.	.	.	.	.	.	.	.	28	.	.	.
283	25	4 shank	.	.	.	.	.	.	.	.	.	.	.	31	27	.	.	.	.	4	.	.	.	.	41	.	.	.	.	.	.	.	.	.
284	17	4 shank	.	.	.	.	.	.	118	.	.	.	.	.	.	.	.	51	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
284	18	4 shank	.	.	.	.	.	.	.	.	.	.	.	12	10	2	.	.	.	27	3	.	77	18	.	.	.	.	.	.	.	.	.	.
284	19	4 shank	.	.	.	.	.	.	.	.	5	.	.	18	29	.	.	.	.	.	.	.	5	177	.	.	.	.	.	.	.	.	.	.

Acute recordings in VR task (n = 6 naive nice, 17 sessions)

B

Mouse	Exp #	Probe	dFC			vFC			SC						HC			BG			THL								SUB						
			M2	ACC	PL	IL	OFC	DP	TT	OLF	SS	VIS	PPC	RSC	DG	CA1	CA3	STR	CP	NAC	LGN	MDT	CMT	VPT	PNT	PVT	PFS	LSR	LSV	MSP	PCN	HAB	HYP	MID
166	2	1 shank	65	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	107	21	.	.	.	.	.	.	.	.	.	.	.	.	.	.
166	3	1 shank	87	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	87	11	.	.	.	.	.	.	.	.	.	.	.	.	.	.
167	1	1 shank	.	7	7	.	.	.	.	.	.	.	.	.	.	.	.	.	102	12	.	.	.	.	.	.	.	.	.	.	.	.	.	.
167	2	1 shank	1	.	11	.	.	.	.	.	.	.	.	.	.	.	.	.	74	5	.	.	.	.	.	.	.	.	.	.	.	.	.	.
167	3	1 shank	.	11	19	.	.	.	.	.	.	.	.	.	.	.	.	.	102	18	.	.	.	.	.	.	.	.	.	.	.	.	.	.
168	1	1 shank	11	17	29	.	.	.	.	.	.	.	.	.	.	.	.	.	90	32	.	.	.	.	.	.	.	.	.	.	.	.	.	.
168	2	1 shank	8	18	19	.	.	.	.	.	.	.	.	.	.	.	.	.	76	37	.	.	.	.	.	.	.	.	.	.	.	.	.	.
168	3	1 shank	7	2	4	.	.	.	.	.	.	.	.	.	.	.	.	.	57	13	.	.	.	.	.	.	.	.	.	.	.	.	.	.
169	1	1 shank	23	7	1	.	.	.	.	.	.	.	.	.	.	.	.	5	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.
169	2	1 shank	37	12	2	.	.	.	.	.	.	.	.	.	.	.	.	31	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
169	3	1 shank	28	22	1	.	.	.	.	.	.	.	.	.	.	.	.	51	.	5	.	.	.	.	.	.	.	.	.	.	.	.	.	.
170	1	1 shank	25	6	.	.	.	.	.	.	.	.	.	.	.	.	.	40	.	55	.	.	.	.	.	.	.	.	.	.	.	.	.	.
170	2	1 shank	32	8	13	.	.	.	.	.	.	.	.	.	.	.	.	31	.	7	.	.	.	.	.	.	.	.	.	.	.	.	.	.
170	3	1 shank	2	1	7	.	.	.	.	.	.	.	.	.	.	.	.	35	.	9	.	.	.	.	.	.	.	.	.	.	.	.	.	.
171	1	1 shank	40	.	.	.	66	.	.	21	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
171	2	1 shank	48	.	2	.	112	.	.	23	.	.	.	.	.	.	.	.	.	18	.	.	.	.	.	.	.	.	.	.	.	.	.	.
171	3	1 shank	60	.	5	.	109	.	.	35	.	.	.	.	.	.	.	.	.	33	.	.	.	.	.	.	.	.	.	.	.	.	.	.

Chronic recordings in free-moving (FM) task (n = 4 animals, 32 sessions)

C

Mouse	Exp #	Probe	dFC			vFC			SC						HC			BG			THL								SUB						
			M2	ACC	PL	IL	OFC	DP	TT	OLF	SS	VIS	PPC	RSC	DG	CA1	CA3	STR	CP	NAC	LGN	MDT	CMT	VPT	PNT	PVT	PFS	LSR	LSV	MSP	PCN	HAB	HYP	MID
248	11	1 chronic	40	11	12	.	.	.	.	.	.	.	.	.	.	.	.	.	11	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
248	12	1 chronic	37	12	10	.	.	.	.	.	.	.	.	.	.	.	.	.	11	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
248	13	1 chronic	46	9	10	.	.	.	.	.	.	.	.	.	.	.	.	.	13	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
248	14	1 chronic	41	12	9	.	.	.	.	.	.	.	.	.	.	.	.	.	10	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
248	15	1 chronic	46	11	13	.	.	.	.	.	.	.	.	.	.	.	.	.	15	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
248	16	1 chronic	37	11	9	.	.	.	.	.	.	.	.	.	.	.	.	.	10	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
248	17	1 chronic	41	13	9	.	.	.	.	.	.	.	.	.	.	.	.	.	10	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
248	18	1 chronic	31	17	13	.	.	.	.	.	.	.	.	.	.	.	.	.	7	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
249	11	1 chronic	1	.	.	2	.	.	12	.	.	.	.	.	.	.	.	.	15	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
249	12	1 chronic	1	1	2	5	.	.	12	.	.	.	.	.	.	.	.	.	9	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
249	13	1 chronic	.	.	2	4	.	.	11	.	.	.	.	.	.	.	.	.	13	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
249	14	1 chronic	2	.	2	5	.	.	10	.	.	.	.	.	.	.	.	.	9	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
249	15	1 chronic	.	.	2	5	.	.	9	.	.	.	.	.	.	.	.	.	11	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
249	16	1 chronic	.	.	.	4	.	.	11	.	.	.	.	.	.	.	.	.	12	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
249	17	1 chronic	.	.	2	3	.	.	13	.	.	.	.	.	.	.	.	.	10	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
249	18	1 chronic	.	.	2	5	.	.	12	.	.	.	.	.	.	.	.	.	11	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
250	11	1 chronic	.	22	45	11	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
250	12	1 chronic	.	24	54	18	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
250	13	1 chronic	.	26	55	18	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
250	14	1 chronic	.	16	46	17	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
250	15	1 chronic	.	13	62	12	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
250	16	1 chronic	.	14	42	6	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
250	17	1 chronic	.	14	50	22	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
250	18	1 chronic	.	7	55	22	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
251	11	1 chronic	61	5	2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
251	12	1 chronic	60	5	2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
251	13	1 chronic	68	8	2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
251	14	1 chronic	54	4	2	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
251	15	1 chronic	47	6	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
251	16	1 chronic	34	4	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
251	17	1 chronic	45	4	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
251	18	1 chronic	47	4	2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.

Figure S6. Additional information about Neuropixels recordings.

A) Extended summary table for acute Neuropixels recordings in trained mice performing the virtual (VR) foraging task, showing mouse, session number, probe type (1-shank/4-shanks), and number of recorded neurons from each region (n = 47 recording sessions from 10 mice).
B) Extended summary table for acute Neuropixels recordings in task-naive (first 3 days) mice performing the VR task as in **(A)** (n = 17 recording sessions from 6 mice).
C) Extended summary table for acute Neuropixels recordings in mice performing the free-moving (FM) foraging task showing details as in **(A-B)** (n = 32 recording sessions from 4 mice).

Figure S7

A

Subspace alignment analysis
using canonical correlation analysis (CCA)

1) PCA to define subspaces

$X_{Det} \xrightarrow{PCA} S_{Det}$
$X_{Stc} \rightarrow S_{Stc}$

(Keep PCs that capture 80% of variance)

2) Compute cross-covariance C

$C = (S_{Det}^T S_{Stc})$

3) Compute SVD of C

$U\sum V^T = C$

U, V
Singular vectors
(canonical directions)

diag($\sum$)
Singular values σ
(canonical correlations)

4) Extract angles between canonical directions

$(\theta_1, \theta_2, \ldots, \theta_n) = \cos^{-1}(\sigma_1, \sigma_2, \ldots, \sigma_n)$

5) Compute distance metrics

$D_{chord} = \sqrt{\sum \sin^2(\theta)}$
ortho vecs = sum(find(θ>80°))

B Identical subspaces

$\sqrt{\sum \sin^2(\theta)} = 0$
Ortho vecs = 0
$\Theta_1 = 0$
$\Theta_2 = 0$

C Separate subspaces (rotate 30°) (no orthogonal vector pairs)

$\sqrt{\sum \sin^2(\theta)} = 0.5$
Ortho vecs = 0
$\Theta_1 = 0$
$\Theta_2 = 30$

Subspace_A
Subspace_B
Vector pair 1 (θ_1)
Vector pair 2 (θ_2)

D Separate subspaces (one orthogonal vector pair)

$\sqrt{\sum \sin^2(\theta)} = 1$
Ortho vecs = 1
$\Theta_1 = 0$
$\Theta_2 = 90$

E Separate subspaces (one orthogonal vector pair)

$\sqrt{\sum \sin^2(\theta)} = 1$
Ortho vecs = 1
$\Theta_1 = 0$
$\Theta_2 = 90$

rotate 30° around X_1

F

$\beta = 0.4363$
$R^2 = 0.998$
$p = 1.85e-6$

Δ $\sqrt{\sum \sin^2(\theta)}$ (above baseline Frac=0)

Fraction context coding (synthetic activity)

G

$\beta = -0.1519$
$R^2 = 0.199$
$p = 0.375$

Δ orthogonal vectors (above baseline Frac=0)

Fraction context coding (synthetic activity)

H

**

orthogonal vectors (across - within context)

I

n = 32 regions
$R^2 = 0.275$
$p = 0.002$

orthogonal vectors (across-within)

Context coding (fraction neurons)

J

n = 47 sessions
$R^2 = 0.091$
$p = 0.0393$

KS statistic (Det vs. Stc PRT)

Subspace distance $\sqrt{\sum \sin^2(\theta)}$

Figure S7. Description of subspace alignment analysis and additional analytical controls.

A) Graphical description of subspace alignment analysis pipeline. For each session, principal component analysis (PCA) is used on neuron spiking data within each context to determine subspaces S_{Det} and S_{Stc}, and canonical correlation analysis (CCA) is used to find canonical correlations and compute subspace distance metrics between S_{Det} and S_{Stc} (see Methods).

B) Example visualization of subspace alignment analysis applied to identical 2D subspaces (green = 1st pair of canonical directions; red = 2nd pair of canonical directions; θ = angle between vector pairs).

C) Visualization of analysis applied to separate 2D subspaces with 30° angle between them and one shared canonical direction.

D) Visualization of analysis applied to separate 2D subspaces with one orthogonal pair of canonical directions and one shared canonical direction.

E) Visualization after rotating (30°) one subspace in **(D)**, maintaining the same subspace distance.

F) Relationship between chordal distance $\sqrt{\sum \sin^2(\theta)}$ and addition of synthetic context coding activity (see Methods) in single neurons (fraction of context-coding neurons, linear regression, $\beta = 0.44$, $R^2 = 0.99$, $p = 1.85\mathrm{e}\text{-}6$).

G) Relationship between orthogonal vectors and addition of synthetic context coding activity in single neurons (linear regression, $\beta = -0.1519$, $R^2 = 0.199$, $p = 0.375$).

H) Normalized (between - within context) number of orthogonal vectors for Det- vs. Stc-context subspaces (mean ± s.d., t-test, $p = 0.0034$, $n = 47$ sessions).

I) Correlation across regions between context coding (fraction of neurons) and number of orthogonal vectors between context subspaces ($R^2 = 0.275$, $p = 0.002$, $n = 32$ regions).

J) Correlation across sessions between subspace distance $\sqrt{\sum \sin^2(\theta)}$ and decision flexibility, indexed as the KS statistic between Det vs. Stc PRT distributions ($R^2 = 0.091$, $p = 0.039$, $n = 47$ sessions).

***$p<0.001$; **$p<0.01$; *$p<0.05$; n.s. $p>0.05$.

Figure S8

A

Single-neuron DV coding overlap (% of DV coding cells)

B

Figure S8. Additional analysis of neural data from virtual foraging task.

A) Comparison of single-neuron DV coding and overlap across within-context control conditions. DV coding for individual neurons was computed using correlation analysis as in **Figure 4C** (see Methods), except using even splits of within- or across-context trial data as schematized at bottom, and overlap of DV coding is quantified in pie charts. Across-context DV coding (left) overlaps significantly less than within-context comparisons (middle, right, Fisher's exact tests, p = 1.1967e-27 [across vs. within-Stc], p = 1.842e-12 [across vs. within-Det], p = 8.079e-6 [within-Stc vs. within-Det]).
B) Comparison of decision variable (DV) coding strength in DV-coding neurons across sampled brain regions, measuring (for each neuron) the absolute-value correlation |r| between its firing rate and the model-fit DV (median, IQR, range of session samples for each region).

Figure S9

Virtual reality (VR) patch foraging task
dFC inactivation (Muscimol → M2)

Free-moving (FM) patch foraging task
dFC inactivation (Muscimol → M2)

Figure S9. Additional data for dFC inactivation experiments.

A) Comparison of patch engagement (trials/min) for mice in the virtual (VR) foraging task under muscimol vs. saline (SR test, p = 0.0645, n = 10 mice).
B) Comparison of ΔCDF of Det vs. Stc PRT distributions for mice in the VR task under Muscimol vs. saline (SR test, p = 0.0273, n = 10 mice, alternative to KS statistic, see **Figure S2B**).
C) Comparison of context transition rate (switches/min) for mice in the free-moving (FM) foraging task under Muscimol vs. saline (SR test, p = 2.44e-4, n = 10 mice).
D) Comparison of port switch rate (average probability of port transition) for mice in the FM task under Muscimol vs. saline (SR test, p = 2.44e-4, n = 10 mice).
E) Comparison of average patch residence times (PRT) for mice in the FM task under Muscimol vs. saline (SR test, p = 2.44e-4, n = 10 mice).

For all plots, each dot or line is one mouse; Wilcoxon signed rank test (SR test); ***p<0.001; **p<0.01; *p<0.05; n.s. p>0.05.

Table S1. Brain region categories and Allen atlas correspondence.

Region name	Tag	Macro-region category	Allen atlas labels
Secondary motor cortex	M2	Dorsal frontal cortex (dFC)	'Secondary motor area'
Anterior Cingulate cortex	ACC	Dorsal frontal cortex (dFC)	'Anterior cingulate area'
Prelimbic cortex	PL	Dorsal frontal cortex (dFC)	'Prelimbic'
Infralimbic cortex	IL	Ventral frontal cortex (vFC)	'Infralimbic'
Orbitofrontal cortex	OFC	Ventral frontal cortex (vFC)	'Orbital'
Dorsal peduncle	DP	Ventral frontal cortex (vFC)	'Dorsal peduncular'
Tenia tecta	TT	Sensory cortex (SC)	'Tenia tecta'
Olfactory areas	OLF	Sensory cortex (SC)	'Olfactory areas'
Somatosensory areas	SS	Sensory cortex (SC)	'somatosensory'
Visual areas	VIS	Sensory cortex (SC)	'visual areas'
Posterior parietal cortex	PPC	Sensory cortex (SC)	'anterior area'
Retrosplenial cortex	RSC	Sensory cortex (SC)	'Retrosplenial'
Dentate gyrus	DG	Hippocampal (HC)	'Dentate gyrus'
Hippocampus CA1	CA1	Hippocampal (HC)	'Field CA1'
Hippocampus CA3	CA3	Hippocampal (HC)	'Field CA3'
Striatum	STR	Basal ganglia (BG)	'Striatum'
Caudoputamen	CP	Basal ganglia (BG)	'Caudoputamen'
Nucleus accumbens	NAC	Basal ganglia (BG)	'accumbens'
Lateral geniculate nucleus	LGN	Thalamic (THL)	'lateral geniculate'
Mediodorsal thalamus	MDT	Thalamic (THL)	'Mediodorsal nucleus of the thalamus'
Centromedial thalamus	CMT	Thalamic (THL)	'Central medial nucleus of the thalamus' 'Rhomboid nucleus'
Ventral thalamus	VPT	Thalamic (THL)	'Ventral posteromedial nucleus of the thalamus' 'Ventral posterolateral nucleus of the thalamus'
Posterior thalamus	PNT	Thalamic (THL)	'posterior nucleus of the thalamus' 'Posterior complex of the thalamus' 'Posterior triangular thalamic nucleus' 'Suprageniculate nucelus' 'Ethmoid nucleus of the thalamus'
Paraventricular thalamus	PVT	Thalamic (THL)	'Paraventricular nucleus of the thalamus'
Parafascicular nucleus	PFS	Thalamic (THL)	'Subparafascicular area'
Lateral septum rostral	LSR	Subcortical (SUB)	'Lateral septal nucleus rostral'
Lateral septum ventral	LSV	Subcortical (SUB)	'Lateral septal nucleus ventral'
Medial septum	MSP	Subcortical (SUB)	'Medial septal nucleus'
Precommissural nucleus	PSN	Subcortical (SUB)	'Precommissural nucleus'
Hebnula	HAB	Subcortical (SUB)	'habenula'
Hypothalamus	HYP	Subcortical (SUB)	'Hypothalamus' 'Preoptic area'
Midbrain	MID	Subcortical (SUB)	'Midbrain'

Methods

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mice

A total of 67 adult mice (C57/BL6j, 33 male, 34 female, age 2-6 months) were used in the experiments. Mice were housed on a 12h dark/12h light cycle (dark from 07:00 to 19:00) and performed behavior tasks in the dark period. All procedures were performed in accordance with the National Institute of Health Guide for the Care and Use of Laboratory Animals and approved by the Harvard Animal Care and Use Committee.

METHOD DETAILS

Surgical procedures

For virtual-reality (VR) Muscimol inactivation experiments, mice were anaesthetized using isoflurane (4% induction, 1-2% maintenance). For experiments requiring head-fixation, a custom titanium head-plate was attached to the skull with Metabond (Parkell). For both acute and chronic Neuropixels recordings, fiducial marks were made at the target sites for probe insertion using a fine-tipped pen, and a ground pin was inserted into the skull above the cerebellum and attached with Metabond. For Muscimol inactivation experiments, a craniotomy was placed above the M2 region of the dorsal frontal cortex (Bregma +2.35mm AP, +1.0mm ML) and Muscimol or saline injected 1.0mm ventral to the skull surface.

For chronic Neuropixels implants, mice were anaesthetized using Ketamine/medetomidine (Ketamine 10 mg/mL, Dexdomitor 0.04 mg/mL, 6 mL/kg). We used a custom adapted version of the Apollo implant described in Bimbard et al.,[50] which uses separable "docking" and "payload" modules to secure and cement the probe to the skull after insertion and recover the probe after completion of experiments. Neuropixels 1.0 single-shank probes were used for chronic recordings targeted in dorsal frontal cortex, and we tracked electrophysiology signals online using SpikeGLX to guide probe placement. After implanting and securing the probe, we injected Antisedan (Atipamezole hydrochloride, 5mg/mL) to reverse the effects of Ketamine and allowed animals to fully recover for 1-2 weeks before behavior experiments.

Foraging behavior experiments

For all experiments, mice were water restricted to maintain body weight > 85% of their baseline weight before training to ensure task motivation and maintain health. Additional water supplementation was provided following training sessions with a total of task and post-task water ranging from 1.0-2.4mL per day, determined individually per mouse. Mice that showed task motivation at higher weights were given greater amounts of water supplementation accordingly to minimize stress from water restriction. Body weights ranged from 85-105% of baseline.

38   Male and female mice were used in approximately equal proportions for all behavior and
39   recording experiments. We observed no significant differences in task engagement, decision
40   behavior, or contextual flexibility in foraging decisions across sexes (**Figure S4**).

Virtual reality setup for VR foraging task

Virtual reality setups were identical to those used in Bukwich et al.[37] Three monitors (width 53cm,
height 30cm) were placed in front and on either side of the animal. Virtual reality scenes were
generated using VirMEn software in MATLAB on a workstation computer (DELL Precision 5810).
Mice were head-fixed on a cylindrical Styrofoam treadmill (diameter 20.3cm, width 10.4cm) and
the rotational velocity of the treadmill was recorded using a rotary encoder. The output pulses of
the encoder were converted into continuous velocity signals using custom Arduino code running
on a microprocessor (Teensy 3.2). Velocity was integrated within VirMEn to compute position.
Water was delivered to the mouse from a spout placed in front of the mouse's mouth, and licks
were monitored using an infrared sensor (OPB819Z, TT electronics). Voltage signals from the
rotary encoder were digitized and recorded on the virtual reality computer using a data acquisition
system (PCIe-6323, National Instruments). Water timing and amount was controlled using a
solenoid valve (LHDA 1221111H, The Lee Company) and a switch (2N7000, On Semiconductor),
with TTL pulses generated by the virtual reality computer via the PCIe-6323 data acquisition card.

Virtual reality (VR) patch foraging task

Habituation: Mice were habituated to the treadmill prior to initial head-fixation with delivery of
occasional water droplets from the reward spout while mice were free to explore on top of the
treadmill, which was held in place to prevent rotation. Once mice showed consistent interest in
consuming water droplets (usually 2 days with 15-minute exposures), head fixation began on the
following session. Initial head-fixation habituation sessions were run for durations of 15 minutes,
with gradual increases in session duration across consecutive sessions reaching a maximum
duration of 50-60 minutes. During these sessions, water rewards (2μL) were delivered in 10 second
intervals with additional rewards given during running bouts to reinforce running behavior. Once
starting head-fixation, mice performed daily sessions with 1 day off per week (6 days on, 1 day
off) for the remainder of the task. Once mice were consistently running for most of the session
(usually 4-6 days), they proceeded on the next session to the first day of the foraging task.

Phase 1: Once mice demonstrated consistent running behavior during habituation (above),
they began training on the linear virtual track. Mice navigated forward down the virtual corridor
and could stop at visually indicated patch locations to receive water rewards (they never received
water between patches). The VR gain was set so that the travel distance between patches was
50cm. On patch entry, mice received an odor puff indicating that they were on the patch, and if
they stopped moving for 0.05s, could receive an initial water reward (2μL). Following this reward,
they could stay on the patch to receive additional rewards following patch-specific statistics. For
the first 7-10 days, mice were only exposed to the Deterministic ("Det") environment/context,
which was indicated with a unique visual pattern on the corridor walls, unique visuals for patch
locations, and a unique patch odor (isoamyl acetate or ethyl butyrate). Det patches always yield
the patch-initial reward (at t = 0s) followed by rewards at t = 2s and t = 4s for a total of three

guaranteed rewards. Mice could leave the patch at any time by moving forward on the track. The
immobile time required to initiate the patch trial (starting at 0.05s) was gradually increased by
0.05s each day to a maximum of 0.5s. Occasionally, this was reduced for animals that showed
faltering engagement rates and then increased again to 0.5s as engagement behavior recovered.

Phase 2: Mice then proceeded to Phase 2 of the task, where they foraged in the Det
environment and the Stochastic (Stc) environment once per session in two blocks (20-25 minutes
each) with one context transition in between. The Stc environment was indicated with distinct
visuals and odor (isoamyl acetate or ethyl butyrate) on patch entry. On Stc patches, after an initial
reward at t = 0s, mice could receive additional rewards each second with probably P(Reward) that
decayed over time (**Figure S3A**). Initial P(Reward) P_i at t = 1s for Stc patches was drawn randomly
(uniformly) from P_i = [0.1, 0.26, 0.58, 0.95] and followed an exponential decay with τ = 0.19s
($P_t = P_t e^{(-\tau t)}$), with P = 0 after 30 seconds. 10% of the patches in the Stc environment were
programmed with the Det patch return ("Stc:024 patches" with rewards at t = 0, 2, 4s) but were
otherwise indistinguishable from other Stc patches. Initial context identity alternated each day, and
the context transition in the middle of each session was only trigged between patches and included
a 10 second blank screen with no visual graphics before initiating the second environment. All
rewards were 2µL water droplets.

Arena setup for free-moving foraging task

Mice navigated a hexagonal arena (inner radius 26.35 cm) with waterspouts placed at four
locations in the center of the arena walls (see **Figure 1C**) and a wall with a controllable door
separating the arena into two chambers (the "Stc" and "Det" environment contexts, each with two
waterspouts). Mice were detected at waterspouts using IR beam breaks (3mm LED IR beam
sensors, Adafruit), and water timing and amount was controlled using solenoid valves (LHDA
1221111H, The Lee Company). Task events (water delivery and door separating the chambers)
were controlled and recorded using custom Arduino code running on a microprocessor (Teensy
4.1). Transparent acrylic barriers were placed between the patches to increase and equalize the
effective path length between patches (approximately 30 inches, 76.2 cm) and encourage greater
time investment for patch trials. Mice could approach and nose-poke a reward port to initiate a
trial, upon which they would immediately receive a 2µL water droplet (marking t = 0s in the patch
trial). If they remained with their nose inserted, they could continue to receive water rewards each
second with decreasing probability based on patch statistics (see below) and were free to leave the
patch at any time by removing their nose. After leaving a patch, that reward port became "locked
out" and would not initiate another trial for 6 seconds. All rewards were 2µL water droplets.

Free-moving (FM) patch foraging task

Habituation: Mice began water restriction one day prior to the task and were handled for 15
minutes prior to starting experiments. Mice were not habituated to the experiment chamber before
exposure on day 1.

Phase 1: For the first 10 sessions (30 minutes), mice were exposed only to the right chamber
(the Det environment) which included two reward ports yielding Det reward returns (the door
separating the chambers remained closed). After the initial reward at t = 0s, mice could receive

additional rewards at t = 1 and t = 2s (for a total of three guaranteed rewards). Mice shaped their
patch leaving decision strategies to this deterministic structure during Phase 1 (**Figure S1**).

Phase 2: Mice then proceeded to Phase 2, in which they were exposed to the Det environment
(the right chamber) and the Stochastic ("Stc") environment (the left chamber) once per session in
15-minute blocks, alternating the starting environment each day. After 15 minutes in the session,
the door separating the chambers opened, and once the mouse explored the other chamber and
completed a patch trial, the door was automatically shut, imposing a block structure. On Stc
patches, after an initial reward at t = 0s, mice could receive additional rewards each second with
decaying probability. Initial P(Reward) P_i at t = 1s for Stc patches was drawn randomly (uniformly)
from P_i = [0.66, 0.81, 0.93, 0.99] and followed exponential decay with $\boldsymbol{\tau} = 0.325s$ ($\boldsymbol{P_t = P_t e^{(-\tau t)}}$),
with P = 0 after 20 seconds. 15% of the patches in the Stc environment were programmed with the
Det patch return ("Stc:012 patches" with rewards at t = 0, 1, 2s) but were otherwise
indistinguishable. Mice experienced the Phase 2 task structure for 2 weeks, typically with 1 day
off per week (6 days on, 1 day off).

Phase 3: After Phase 2, mice transitioned to the final Phase 3 of the task, in which the
controllable door separating the environments/chambers was always left open. Mice were
therefore free to engage context transitions and forage completely under their own volition.
Although mice gained on average more rewards on Det compared to Stc patches (**Figure S1H**)
and preferred to exploit patches in the Det context (**Figure S1I**), they still explored both contexts
extensively in every session, typically making 2-3 context transitions each minute. Despite the
lack of block structure or specific context cues, mice exhibited the same differences in decision
strategies across contexts as observed in Phase 2 (**Figures S2C-F**).

Muscimol inactivation experiments

To test the functional role of dorsal frontal cortex (dFC) activity in context-specific decision
making, we inactivated dFC neurons using local infusion of the GABA$_A$ receptor agonist
Muscimol (Sigma-Aldrich M1523) targeting the secondary motor cortex (M2) subregion of dFC.
For the VR task, mice were trained as described above for two weeks in Phase 2. Then, once per
day about 2 hours before experiments, they were briefly (~15 minutes) anesthetized and injected
with either Muscimol (1ng/nL, 150 nL) or saline (150 nL) over 5 minutes targeting the right side
M2 using the coordinates (Bregma +2.35mm AP, +1.0 MLmm, -1.0mm DV from the skull
surface). Mice were completely recovered from anesthesia before experiments and performed the
task as usual. We injected Muscimol or saline for 2 sessions each on alternating days. Mice showed
no major deficits in task engagement (**Figure S9A**).

For inactivation experiments in the FM task, mice were trained as described above through
Phase 1 and Phase 2, and for four days in Phase 3. We then injected either Muscimol or saline into
M2 on alternating days and performed behavior experiments (FM task Phase 3) about 2 hours after
injection. Mice exhibited several behavioral changes under Muscimol (**Figures 7J-O, S9C-E**) but
did not show obvious deficits in motor behavior and still completed 10s-100s of trials per session.

Electrophysiological recording and spike sorting

Acute recordings in the VR task: Spiking data were collected using Neuropixels 1.0 single shank
or Neuropixels 2.0 4-shank probes and acquired using SpikeGLX
(https://billkarsh.github.io/SpikeGLX/). Craniotomies were performed 12-24 hours before the
recording using a dental drill and covered with Kwik-cast (World precision Instruments), and mice
recovered fully before experiments. On the day of recording, a 3D-printed shield was placed above
the animal's head to block light and sight of the experimenter. Prior to insertion, probes were dyed
with DiI, DiD, or DiO (Thermofisher) to label probe tracts from each recording session in tissue.
Neuropixels probes were lowered using a Thorlabs micromanipulator (PT1-Z8) at 12-14 µm/sec.
After reaching the target depth, the probe was allowed to settle in the tissue for 20 minutes prior
to the start of the recording. Recordings typically lasted 50-60 minutes.

Chronic recordings in the FM task: Spiking data were collected using Neuropixels 1.0 single
shank probes targeting the dorsal frontal cortex (+2.25mm AP, +0.8mm ML) and recorded with
Bonsai software (OpenEphys.Onix1 software library) using the ONIX PCIe acquisition system
(open Ephys). Craniotomies and probe implantation were performed at least one week before
behavior and recording sessions began. Artificial dura (Dura-Gel, Cambridge Neurotech) was
applied during probe implantation to protect the site of the craniotomy. Probes were mounted with
dental cement using a custom adaptation of the "Apollo" implant system to enable probe recovery
and were dyed using DiI to mark the tissue trajectory of the probe.[50]

For all neural recording experiments, Neuropixels data were spike sorted offline using
Kilosort 4 (https://github.com/MouseLand/Kilosort) followed by manual curation in Phy
(https://github.com/cortex-lab/phy) to ensure quality spike waveforms and recording stability for
isolated units and to split doublet clusters. Multi-unit and LFP signals were not used for any
analysis. See **Figure S6** for full recording summaries.

Histology and probe tract analysis

After the last recording session, mice were transcardially perfused with phosphate buffered saline
(PBS) followed by 4% paraformaldehyde in PBS. The brains were sliced in 100 µm coronal
sections using a vibratome (Leica). Brain sections were mounted on glass slides and stained with
4', 6-diamidino-2-phenylindole (DAPI, Vectashield) in mounting medium. Slides were then
imaged with a fluorescence microscope (Zeiss, Axio Scan.Z1). Probe tracts (labeled with
fluorescence from DiI, DiD, and DiO) were identified in histology images (processed using Zen
3.3, Carl Zeiss Microscopy) and aligned using histological and electrophysiological landmarks to
a reference atlas (Allen brain atlas CCF) using MATLAB software written by Dr. Andrew Peters
(https://github.com/petersaj/AP_histology). Single units were accordingly sorted into 32 region
and 7 macro-region categories using region identities from the reference atlas (see **Table S1**).
Single units localized in white-matter structures within 50mm (vertical) of an identified region
were assigned to it. Units localized in white-matter structures outside of 50mm from other regions,
or those assigned to regions outside the 32 categories, were not used for analysis.

Mouse and trial inclusion criteria

For basic behavior and recording experiments in the VR head-fixed foraging task, 41 mice (22
males and 19 females) were trained and used for analysis, of which 10 were recorded from. 47
recording sessions were used for neural analysis. For analysis of decision behavior, trials from
individual sessions were only included if at least 10 trials were performed in the session
(70548/70827, 99.61% of all trials). Mice occasionally disengaged from the task for brief periods
of time. Therefore, to prevent contamination of decision behavior by the disengaged state, we did
not include trials with PRT (patch residence time) greater than 60 seconds for behavior analysis
(2243/70827, 3.17% of all trials dropped).

For task-naïve recordings in the VR task, 6 mice (4 females, 2 males) were habituated to head
fixation as described above and then proceeded directly to Phase 2 of the foraging task for 3 days,
during which acute Neuropixels recordings were performed. 17 recording sessions were used for
neural analysis.

For Muscimol inactivation experiments in the VR task, mice were required to engage at least
10 trials of each type (Det and Stc:024) during both Muscimol and saline sessions to be used for
analysis. 10/14 mice trained on the task passed this inclusion criterion and were used for behavior
analysis.

For basic behavior and recording experiments in the FM foraging task, 20 mice (9 males and
11 females) were trained and used for analysis, of which 4 were recorded from. 32 recording
sessions (16 from Phase 2 and 16 from Phase 3) were used for neural analysis. All sessions were
used for behavior analysis. We did not include trials with PRT greater than 6 seconds for behavior
analysis.

For Muscimol inactivation experiments in the FM task, mice were required to engage at least
10 trials of each type (Det and Stc:012) during both Muscimol and saline sessions to be used for
analysis. 13/13 mice trained on the task passed this inclusion criterion and were used for behavioral
analysis.

Analysis of patch residence time (PRT) decision behavior

We analyzed decision behavior for patch foraging trials using the patch residence time (PRT) for
different trial types. To evaluate whether mice made patch leaving decisions in a value-sensitive
manner, we compared PRTs across four hidden trial types within Stc patches which had different
initial reward probabilities (P_i). P_i was not cued to animals but was drawn randomly on trial
initiation (see task descriptions above). In both the VR and FM tasks, average PRTs scaled
positively with P_i (**Figures S2A, 2D**), indicating value sensitivity as we previously reported.[37]

We compared distributions of PRTs across patch types as a measure of contextual flexibility in
foraging decisions, visualizing them using the cumulative density function (CDF) of leaving
probabilities over time (as in **Figure 1E**). We quantified their divergence using the Kolmogorov-
Smirnov statistic (KS statistic), which measures the difference in CDFs at their point of maximum
vertical difference. We compared the KS statistic across Det and reward-matched Stc:012 or
Stc:024 trial types, or across subsets of Det trials (**Figures 1I, 2G**), and found that PRT
distributions across contexts were more distinct than within-context Det comparisons. However,
the KS statistic can be sensitive to large, local deviations in the shape of a distribution and can be
skewed by small sample sizes. Thus, for some analyses which required comparing distributions

with very different sample sizes (e.g. comparing PRTs on Det vs. Stc trials for Muscimol
experiments), we estimated the KS statistic by down-sampling trials to equalize sample sizes and
taking the average resampled KS statistic across 1000 draws. We validated behavioral results by
additional measuring distribution differences using the Wasserstein distance (see **Figure S2B**),
which in the 1D case is equivalent to the normalized absolute difference between CDFs (ΔCDF)
and is less sensitive than the KS statistic to localized deviations. These measures were highly
correlated in both task designs (**Figures S4E, S4J**), indicating that they robustly capture
differences in decision behavior implicit in the PRT distributions.

Integrator models of patch leaving decisions

Following our previous report, we modeled patch leaving decisions in both the VR and FM
foraging tasks as a drift-diffusion decision variable (DV) that integrates time on patch (δ) and
reward receipt (ρ) toward a boundary as evidence for or against the leaving decision.[34,37] In the
model (schematized in **Figure 1J**), a decision variable X_t is initialized to 0 at t = 0s (for each patch
trial), incremented by a "drift rate" δ each time step, and is perturbed upon reward receipt by ρ
(the "reward sensitivity"). At each time step, X_t is converted into a leaving probability ($0 <= P_{Leave}$
$<= 1$) using a sigmoid function centered at 0.5 with temperature T ($P_{Leave} = \frac{1}{e^{(-T(X_t - 0.5))}}$). Thus,
the model contains three parameters (δ, ρ, T), and different algorithms for the decision
computation ("Models 1-4") are produced by setting different constraints and definitions for δ and
ρ. Specifically, the "reward insensitive" Model 1 has ($\delta > 0, \rho = 0$); the "reward reset" Model 2
has ($\delta > 0, \rho = -X_t$); the "reward integration" Model 3 has ($\delta > 0, \rho < 0$); and the "reward
accumulation" model 4 has ($\delta > 0, \rho > 0$). Integrator models were fit by using empirical reward
returns and mouse PRT behavior to generate DV rollouts, and then estimating the parameters (δ,
ρ, T) using the fmincon function in MATLAB (model fitting toolbox) to maximize the log-
likelihood between P_{Leave} and the empirical patch leave times (PRTs). For parameter comparisons
across patch types (**Figures 1O-P, 2K-L**), full models were used to estimate δ and ρ using trials
with 0.5s <= PRT < 30s (for VR behavior) and 0.5s <= PRT < 10s (for FM behavior). For
comparison and quantification of model performance, we fit models on 90% of the trials, computed
the Bayesian Information Criterion (BIC) between predicted P_{Leave} and ground truth leave times
for held-out trials, and compared average (across folds) cross-validated BIC (*cv BIC*) for Models
1-4 (**Figures 1K-L, 2I-J**).

We reported in a previous study (Bukwich et al.)[37] that, in a similar VR patch foraging task,
patch leaving decisions were best described using an integrator model that is scaled by a latent
"patience" state ϕ, which captures individual variability in waiting behavior as well as within-
session slow fluctuations in PRTs that are not explained by patch rewards. For behavior in the VR
task, we used a similar scaling parameter in the models to improve fits to behavior while preserving
interpretability of the evidence weight parameters δ and ρ. For each patch trial, we computed the
current value of a "patience" variable ϕ by taking the average of gaussian smoothed PRTs of the
two adjacent trials and then scaled these to the average ϕ across all trials used for model fitting
($\phi = \phi/mean(\phi)$).Then, to compute the DV X_t in the model fitting procedure, we used δ/ϕ as
the "drift rate." As smaller drift rates yield longer PRTs, this captures the effect of the patience
state on leaving decisions by scaling the drift rate to bias trial PRTs in the direction of those for

neighboring trials. As we did not observe substantial variability in FM task behavior, we did not
include this "patience" scaling by ϕ for fits of the integrator models for the FM task.

Generating predicted PRTs from model outputs

To simulate PRTs from specific DV rollouts fit for example mice (**Figure 1Q**), we took the
deterministic DVs $\mathbf{X_t}$ generated from model fits (**Figure 1Q**, left), added gaussian noise ($\boldsymbol{\sigma = 0.1}$),
computed $\mathbf{P_{Leave}}$ as described above, took the first timepoint where $P_{Leave} > 0.5$ as one simulated
PRT, and repeated this procedure 100 times. We used this distribution of simulated PRTs to
compute $\mathbf{P_{Leave}}$ CDFs for different parameterizations of the DV (fit to behavior from Stc and Det
patch trials) and plotted these.

Normalization of single-neuron firing rate traces

300 Firing rate traces for each unit were generated by binning spikes into 50ms time bins and then z-
301 scoring across the entire time series. Throughout (e.g. in **Figures 3E and 4D-E**), σ indicates 1
302 standard deviation (s.d.) of firing rate activity. For some analyses, local single-neuron activity was
303 further normalized by subtracting activity within a pre-event (e.g. such as pre-patch trial) window
304 and shown as "Δ activity" (still in units of σ).

Smoothing and de-noising for principal component projections

Where principal component analysis (PCA) was used to visualize low-dimensional projections of
population activity, single-unit traces were further smoothed or averaged to reduce noise
contributions to PC identification.[42] For PC projections of activity across long timescales capturing
the context transition (**Figures 3H and 6G**), single-neuron traces were smoothed using a 10-
second left moving-mean filter (i.e. only using previous, and not future, timepoints). For PC
projections of single-trial activity, single-neuron activity was averaged within trial types to reduce
noise from trial variability and then smoothed with a 1-second left moving-mean filter.

Analysis of single-neuron and population coding of sensory context

We identified individual neurons as encoding context identity by measuring consistent changes in
firing rate corresponding to environment identity ("Stc" or "Det") that occur at the time of context
transition. For each neuron, we sampled activity in 60-second bins across the entire session and
performed a 2-sample t-test at $\alpha = 0.05$ comparing the mean activity across samples from context
1 and context 2. Then, we measured the change in mean activity "Δ activity" across 5-minute
windows before and after the context transition and obtained a p-value (testing at $\alpha = 0.01$) by
comparison to a null distribution generated from sampling Δ activity around random within-
context timepoints. Neurons that independently passed both statistical tests and exhibited the same
direction of activity change (i.e. consistent global and transient changes in firing rate) were
considered "context coding." For regional comparisons of single-neuron context coding (**Figure
3J**), only session/region samples with at least 5 neurons were used for analysis.

For quantification of population-level context discriminability (**Figure 3I and 6J**), we split the
single-neuron traces into rough quarter bins (for each session) corresponding to the first and second
half of each context block, and then took the average activity of each neuron in each of these bins,
forming population vectors for context 1 (half 1 and half 2) and for context 2 (half 1 and half 2).
We then computed three angles comparing these population vectors by first normalizing their
lengths and then taking the inverse cosine of their dot product ($\theta = cos^{-1}(V_1 \cdot V_2)$). For within-
context comparisons, we computed the angle between the two halves within each context, and for
across context comparison, we took the angle between vectors for the second half of context 1 and
the first half of context 2.

Analysis of subspace divergence

To analyze how context reconfigures population dynamics, we quantified the divergence of
activity subspaces that separately capture population activity for Det and Stc contexts. To do this,
we used canonical correlation analysis (CCA), which finds directions of maximum covariance
between two sets of data and can be used to measure their subspace overlap (see **Figure S7** for
graphical description and examples).[43] For each session or session/region sample of population
activity, we first separately applied PCA to epochs corresponding to the Stc and Det context to
define directions in the state space (the principal components) which capture within-context
dynamics, and retained the top N_{PC} components for each, where N_{PC} is the number of PCs
computed for the full dataset that capture 80% of total variance. These sets of PCs (S_{Stc} and S_{Det})
are orthonormal bases that define subspaces of the full state space.[22,42] To measure their alignment,
we first computed their cross-covariance matrix ($C = S_{Stc}{}^T S_{Det}$) and then performed a singular
value decomposition of C. ($U\Sigma V^T = C$). This yields two sets of singular vectors (or canonical
directions) **U** and **V** which are successively maximally correlated with each other under the
constraint that **U** and **V** are themselves orthogonal.[43,44] The diagonal of $\sum$ contains the
corresponding singular values (or canonical correlations) for each vector pair, which are related to
their alignment and the angle between them. These angles (θ) can be used to quantify subspace
alignment; completely overlapping subspaces yield all angles $\theta = 0°$, while completely orthogonal
subspaces yield all $\theta = 90°$ (see **Figures S7B-E** for illustrations). We measured overall subspace
divergence as $\sqrt{\sum sin^2(\theta)}$, the "chordal distance" between the subspaces, as well as the number
of singular vector pairs with $\theta > 80°$ ("orthogonal vectors"), which can be interpreted as the
dimensional size of the components of the subspaces that are orthogonal to one another.

To evaluate whether subspace divergence was greater than expected from within-context
neural variability (**Figures 3L-M**), we performed this analysis across epochs corresponding to the
second half of context 1 and the first half of context 2 (across-context) and compared these by
subtraction to the first and second halves of context 2 (within-context) as direct within-session
controls. We measured overall subspace difference metrics for each brain region as the median of
all region/session samples with at least 5 neurons.

Because we observed a strong correlation between subspace difference and context coding
across regions (**Figures 3M, S7I**), we analyzed how much these alignment metrics were obligately
correlated with linear firing rate changes along one dimension (e.g. along a "context coding"
vector), as this could yield a spurious association. To test this, we repeated the subspace alignment
analysis using data from each session taken from two halves of one context and synthetically added

activity with gaussian noise (μ = 0-2 s.d., σ = 0-0.1 s.d.) to different sized subpopulations of
neurons to replicate the presence of linear "context coding." Although adding synthetic context
coding to real neural activity slightly increased chordal distance measures (**Figure S7F**), the
relationship (measured using linear regression) was far weaker than that observed in the real data
(β = 0.436 synthetic, β = 1.185 real). Additionally, adding synthetic context coding did not increase
the number of orthogonal vectors between subspaces (**Figure S7G**), while real context coding was
significantly correlated with the number of orthogonal vectors (**Figure S7I**).

Analysis of single-neuron decision variable (DV) coding

To identify single-neuron correlates of the foraging decision variable (DV), we used a similar
approach as in Bukwich et al.[37] We first fit the cognitive DV to decision behavior for each mouse
using the integrator model as described, extracted the DV for trials corresponding to each trial type
(e.g. Det trials or Stc trials), and concatenated these separately. For each neuron, we then extracted
the (0.5s mean smoothed) firing rate corresponding to the same patch trials, concatenated these,
and computed the correlation (*r*) between the unit activity and the DV (separately for Stc and Det
trials), which we took as a measure of DV "coding strength." To test statistical significance, we
repeated this process 1000 times, shuffling the trial order before concatenating, and computed a
null distribution of *r* to test the observed value against. Neurons that exceeded either 0.5% tails (α
= 0.01) of this distribution were considered "DV coding" for either the Stc or Det DV.

Analysis of reward responses and ramping activity in DV coding neurons

To assess whether DV coding neurons represented differently parameterized decision variables as
predicted from fits of the integrator models, we measured local reward responses and ramping
activity in neurons encoding the Stochastic DV (Stc-DV) or Deterministic DV (Det-DV). We
included all neurons (pooled across all sampled regions) which passed our bootstrap statistical test
at α = 0.01 for only one (Det or Stc) DV with a positive correlation. We measured average activity
in these Stc-DV and Det-DV neurons around three trial events: the reward marking trial initiation
(t = 0s), additional patch rewards (at t > 0s), and patch leave (the time the animal moves off the
patch location) (**Figures 4G-I**). For the trial initiation response, we included activity from all trials
with PRT >= 1s, measured the reward response as the local change in mean activity across
surrounding 0.5s windows, and measured ramping activity as the slope (β) for regression of
activity against time in the interval [0.25 to 1s] after trial initiation (reported as change in
activity/sec, or $\sigma \cdot s^{-1}$. Patch reward responses were analyzed the same way, except centering the
window to rewards that occur after t = 0s. For ramping activity before patch leave, we included
trials with PRT >= 7s and no rewards at t > 4s so that neural activity was not contaminated by
recent reward responses. We measured ramping activity as the slope β for regression of activity
against time in the interval [-2 to 0.5s], reported in activity/sec, or $\sigma \cdot s^{-1}$.

Analysis of population DV coding and DV coding vectors

To define DV coding vectors for measurements of population coding geometry, we concatenated
Stc or Det DVs as described above for the single-neuron analysis, extracted corresponding single-

neuron activity for all neurons in the population (within sessions), and performed linear regression
of the neuron traces ($\mathbf{X}$) against the DV (i.e. $\boldsymbol{DV = X^T\beta}$). The normalized coefficients $\beta/\mathbf{norm(\beta)}$
can be interpreted as a direction in the population activity space that best captures how neural
activity linearly covaries with the DV and thus defines a coding vector (or coding dimension) $\beta_{\mathbf{Stc}}$
(Stc-DV) or $\beta_{\mathbf{Det}}$ (Det-DV).[11,42] The angle between these coding vectors can be measured as the
inverse cosine of their dot product ($\boldsymbol{\theta = cos^{-1}(\beta_{Det} \cdot \beta_{Stc})}$).

Within-context comparisons of the angle between DV coding vectors were computed in the
same way but instead using trial data corresponding to two subsampled halves of trial data within
each context rather than across contexts.

Population decoder analysis of decision variables

For linear decoding of DVs, we fit DVs to mouse behavior and concatenated trial DVs and
corresponding neural activity (for region/session populations) as described above for the coding
vector analysis. However, instead of concatenating all trials, we performed leave-one-out cross-
validation by iterating through trials, leaving each trial out once, computing the DV coding vector
using the remaining trial data, projecting the neural data ($\mathbf{X}$) for the held-out trial onto the coding
vector using $\mathbf{DV_{pred} = X^T\beta}$, and computing the correlation r between the ground truth DV and
$\mathbf{DV_{pred}}$. For each session/region sample, we took the average r across all held-out trial predictions
as a summary measure of decoder performance (*cv-r*). For cross-context decoder analyses, we
projected trial data from one context onto the coding vector for the other context and computed
cv-r. For shuffle comparisons, we first randomly circularly permuted each neuron activity trace
and then repeated the analysis as described. Trial data (DVs and corresponding activity) were all
time-warped to common length so that, once concatenated, they would contribute equally to the
regression regardless of variable trial lengths. For DV decoders for the VR task, trials with 0.5s <
PRT < 30s were used, and for the FM task, trials with 0.5s < PRT < 10s were used.

Population decoder analysis of context identity

For linear binary decoding of context identity, we took the average activity for each patch trial
(trial decoder) or each inter-trial-interval (ITI) period in the session as population vector samples,
then randomly up-sampled the group (i.e. Stc or Det trials/ITIs) with fewer samples to equalize
sample sizes and remove bias from the decoder. Iterating 10 times, we then held out 10% of the
samples (non-overlapping across iterations), fit a Fisher's Linear Discriminant (FLD) model on
the remaining 90% to discriminate Stc vs. Det identity (fitcdiscr in MATLAB), and tested predicted
labels against ground truth, using the area under the receiver operating characteristic curve
(auROC) as a performance metric. The average across auROCs for the 10 folds (*cv-auROC*) was
used as an overall summary measure. For shuffle comparisons, class labels (Det vs. Stc context
identity) were randomly shuffled before model fitting.

QUANTIFICATION AND STATISTICAL ANALYSIS

Data were analyzed with custom code written in MATLAB (version 2019b, Mathworks). All box and whisker plots represent the median, interquartile range (box), and full range (whiskers) of the underlying distribution. For all statistical tests throughout, normality of data and equal variance of groups was not assumed, and non-parametric (Wilcoxon rank-sum and signed-rank) tests were used for unpaired and paired group comparisons, respectively. Test for normality was used before applying parametric tests (t-tests, Anova), and for all significance testing we used two-tailed tests at $\alpha = 0.05$ or $\alpha = 0.01$ (noted in figure legends). For statistical comparisons of PRT distributions, two-sample Kolmogorov-Smirnov tests were used. Resampling methods based on temporally permuted neuron activity traces or shuffled time points were used to assess significance of single-neuron coding (described in detail above). Statistical significance of population decoder analyses was assessed by comparison to null distributions of performance statistics generated from shuffled trial sequences or class labels. The sizes of mouse groups were not pre-specified and approximated those of previous studies.

ACKNOWLEDGEMENTS

We thank Mike Bukwich and Malcolm Campbell for help setting up the VR foraging task and early discussions on design of experiments; Edward Soucy and Yuwei Li of the Harvard Neurotechnology Core for engineering assistance; the Harvard Center for Biological Imaging for support with microscopy; Mitsuko Watabe-Uchida and Uchida lab members for critical input and feedback on the manuscript. This work was supported by grants from the National Institutes of Health (NIH) (5U19NS113201 to N.U. and 5R01DA05975 to N.U. and M.W.-U.) and the Simons Collaboration on Global Brain (to N.U.). L.K. is supported by a Helen Hay Whitney Fellowship.

AUHOR CONTRIBUTIONS

L.K. designed the foraging tasks, basing the virtual-reality (VR) task on a previous design reported in Bukwich et al.[37] L.K. adapted and built behavioral and recording setups. L.K. performed all behavior and recording experiments for the VR task, L.K. performed all behavior experiments for the free-moving (FM) task, and L.K. and N.S.S. set up and performed chronic Neuropixels recordings for the FM task. L.K. performed Muscimol inactivation experiments with assistance from G.Z. L.K. analyzed all behavioral and neural data with feedback from N.U. L.K. wrote the manuscript with feedback from N.U.

DECLARATION OF INTERESTS

The authors declare no competing interests.

DATA AND CODE AVAILABILITY

All code and data will be uploaded to a public repository upon publication

References

1.	Miller, E. K. & Cohen, J. D. An integrative theory of prefrontal cortex function. *Annu. Rev. Neurosci.* **24**, 167–202 (2001).

2.	Diamond, A. Executive functions. *Annu. Rev. Psychol.* **64**, 135–168 (2013).

3.	Passingham, R. *Understanding the Prefrontal Cortex*. (Oxford University Press (OUP), 2021).

4.	Stokes, M. G. *et al.* Dynamic Coding for Cognitive Control in Prefrontal Cortex. *Neuron* **78**, 364 (2013).

5.	Cole, M. W. *et al.* Multi-task connectivity reveals flexible hubs for adaptive task control. *Nat. Neurosci.* **16**, 1348 (2013).

6.	Rigotti, M. *et al.* The importance of mixed selectivity in complex cognitive tasks. *Nature* **497**, 585–590 (2013).

7.	Liu, Y. & Wang, X. J. Flexible gating between subspaces in a neural network model of internally guided task switching. *Nat. Commun.* **15**, 1–20 (2024).

8.	Okazawa, G. & Kiani, R. Neural Mechanisms That Make Perceptual Decisions Flexible. *Annu. Rev. Physiol.* **85**, 191–215 (2023).

9.	Nakajima, M. Neuronal identity and cognitive control dynamics in the PFC. *Semin. Cell Dev. Biol.* **129**, 14–21 (2022).

10.	Rigotti, M., Rubin, D. B. D., Wang, X. J. & Fusi, S. Internal representation of task rules by recurrent dynamics: the importance of the diversity of neural responses. *Front. Comput. Neurosci.* **4**, (2010).

11.	Mante, V., Sussillo, D., Shenoy, K. V & Newsome, W. T. Context-dependent computation by recurrent dynamics in prefrontal cortex. *Nature* **503**, 78–84 (2013).

12.	Wu, Z. *et al.* Context-Dependent Decision Making in a Premotor Circuit. *Neuron* **106**, 316-328.e6 (2020).

13.	Rikhye, R. V., Gilra, A. & Halassa, M. M. Thalamic regulation of switching between cortical representations enables cognitive flexibility. *Nat. Neurosci.* **21**, 1753–1763 (2018).

14.	Schmitt, L. I. *et al.* Thalamic amplification of cortical connectivity sustains attentional control. *Nature* **545**, 219–223 (2017).

15.	Kumano, H., Suda, Y. & Uka, T. Context-Dependent Accumulation of Sensory Evidence in the Parietal Cortex Underlies Flexible Task Switching. *J. Neurosci.* **36**, 12192–12202 (2016).

16.	Wimmer, R. D. *et al.* Thalamic control of sensory selection in divided attention. *Nature* **526**, 705–709 (2015).

17.	Nakajima, M., Schmitt, L. I. & Halassa, M. M. Prefrontal Cortex Regulates Sensory

Filtering through a Basal Ganglia-to-Thalamus Pathway. *Neuron* **103**, 445-458.e10 (2019).

18. Lam, N. H. *et al.* Prefrontal transthalamic uncertainty processing drives flexible switching. *Nature* **637**, 127–136 (2025).

19. Pagan, M. *et al.* Individual variability of neural computations underlying flexible decisions. *Nature* **639**, 421–429 (2025).

20. MacDowell, C. J. *et al.* Multiplexed subspaces route neural activity across brain-wide networks. *Nat. Commun.* **16**, 1–21 (2025).

21. Ebitz, R. B. & Hayden, B. Y. The population doctrine in cognitive neuroscience. *Neuron* **109**, 3055–3068 (2021).

22. Langdon, C., Genkin, M. & Engel, T. A. A unifying perspective on neural manifolds and circuits for cognition. *Nat. Rev. Neurosci. 2023* 1–15 (2023). doi:10.1038/s41583-023-00693-x

23. Yang, G. R., Joglekar, M. R., Song, H. F., Newsome, W. T. & Wang, X. J. Task representations in neural networks trained to perform many cognitive tasks. *Nat. Neurosci.* **22**, 297 (2019).

24. Flesch, T., Juechems, K., Dumbalska, T., Saxe, A. & Summerfield, C. Orthogonal representations for robust context-dependent task performance in brains and neural networks. *Neuron* **110**, 1258 (2022).

25. Okazawa, G., Hatch, C. E., Mancoo, A., Machens, C. K. & Kiani, R. Representational geometry of perceptual decisions in the monkey parietal cortex. *Cell* **184**, 3748-3761.e18 (2021).

26. Ulanovsky, N. Natural neuroscience : toward a systems neuroscience of natural behaviors. (2025).

27. Miller, C. T. *et al.* Natural behavior is the language of the brain. *Curr. Biol.* **32**, R482–R493 (2022).

28. Mobbs, D., Trimmer, P. C., Blumstein, D. T. & Dayan, P. Foraging for foundations in decision neuroscience: insights from ethology. *Nat. Rev. Neurosci.* **19**, 419 (2018).

29. Pearson, J. M., Watson, K. K. & Platt, M. L. Decision making: the neuroethological turn. *Neuron* **82**, 950–965 (2014).

30. Stephens, D. & Krebs, J. *Foraging Theory*. (Princeton University Press, 1987).

31. Charnov, E. L. Optimal foraging, the marginal value theorem. *Theor. Popul. Biol.* **9**, 129–136 (1976).

32. Hall-McMaster, S. & Luyckx, F. Revisiting foraging approaches in neuroscience. *Cogn. Affect. Behav. Neurosci.* **19**, 225 (2019).

33. Hayden, B. Y., Pearson, J. M. & Platt, M. L. Neuronal basis of sequential foraging decisions in a patchy environment. *Nat. Neurosci. 2011 147* **14**, 933–939 (2011).

34. Davidson, J. D. & El Hady, A. Foraging as an evidence accumulation process. *PLoS Comput. Biol.* **15**, (2019).

35. Gold, J. I. & Shadlen, M. N. The neural basis of decision making. *Annu. Rev. Neurosci.* **30**, 535–574 (2007).

36. Shadlen, M. N. & Kiani, R. Decision making as a window on cognition. *Neuron* **80**, 791–806 (2013).

37. Bukwich, M. *et al.* Competitive integration of time and reward explains value-sensitive foraging decisions and frontal cortex ramping dynamics. *Neuron* **113**, 3458-3475.e12 (2025).

38. Cazettes, F. *et al.* A reservoir of foraging decision variables in the mouse brain. *Nat. Neurosci. 2023* 1–10 (2023). doi:10.1038/s41593-023-01305-8

39. Fetsch, C. R. The importance of task design and behavioral control for understanding the neural basis of cognitive functions. *Curr. Opin. Neurobiol.* **37**, 16 (2016).

40. Jun, J. J. *et al.* Fully integrated silicon probes for high-density recording of neural activity. *Nature* **551**, 232–236 (2017).

41. Maggi, S. & Humphries, M. D. Activity Subspaces in Medial Prefrontal Cortex Distinguish States of the World. *J. Neurosci.* **42**, 4131–4146 (2022).

42. Cunningham, J. P. & Yu, B. M. *Dimensionality reduction for large-scale neural recordings*. *Nature Neuroscience* **17**, 1500–1509 (Nature Publishing Group, 2014).

43. Wang, H. T. *et al.* Finding the needle in a high-dimensional haystack: Canonical correlation analysis for neuroscientists. *Neuroimage* **216**, 116745 (2020).

44. Zhang, X. *et al.* Inter-brain neural dynamics in biological and artificial intelligence systems. *Nature* **645**, 991–1001 (2025).

45. Khilkevich, A. *et al.* Brain-wide dynamics linking sensation to action during decision-making. *Nature* **634**, 890 (2024).

46. Zhang, Y. *et al.* A brain-wide map of neural activity during complex behaviour. *Nature* **645**, 177–191 (2025).

47. Murakami, M., Vicente, M. I., Costa, G. M. & Mainen, Z. F. Neural antecedents of self-initiated actions in secondary motor cortex. *Nat. Neurosci.* **17**, 1574–1582 (2014).

48. Murakami, M., Shteingart, H., Loewenstein, Y. & Mainen, Z. F. Distinct Sources of Deterministic and Stochastic Components of Action Timing Decisions in Rodent Frontal Cortex. *Neuron* **94**, 908-919.e7 (2017).

49. Adam, E. M., Johns, T. & Sur, M. Dynamic control of visually guided locomotion through corticosubthalamic projections. *Cell Rep.* **40**, (2022).

50. Bimbard, C. *et al.* An adaptable, reusable, and light implant for chronic Neuropixels probes. *Elife* **13**, (2025).

51. Karlsson, M. P., Tervo, D. G. R. & Karpova, A. Y. Network resets in medial prefrontal

cortex mark the onset of behavioral uncertainty. *Science* **338**, 135–139 (2012).

52. Akam, T. *et al.* The Anterior Cingulate Cortex Predicts Future States to Mediate Model-Based Action Selection. *Neuron* **109**, 149-163.e7 (2021).

53. Durstewitz, D., Vittoz, N. M., Floresco, S. B. & Seamans, J. K. Abrupt transitions between prefrontal neural ensemble states accompany behavioral transitions during rule learning. *Neuron* **66**, 438–448 (2010).

54. Lak, A. *et al.* Orbitofrontal Cortex Is Required for Optimal Waiting Based on Decision Confidence. *Neuron* **84**, 190–201 (2014).

55. Yang, W., Tipparaju, S. L., Chen, G. & Li, N. Thalamus-driven functional populations in frontal cortex support decision-making. *Nat. Neurosci. 2022 2510* **25**, 1339–1352 (2022).

56. Finkelstein, A. *et al.* Attractor dynamics gate cortical information flow during decision-making. *Nat. Neurosci.* **24**, 843–850 (2021).

57. Yang, Z., Inagaki, M., Gerfen, C. R., Fontolan, L. & Inagaki, H. K. Integrator dynamics in the cortico-basal ganglia loop for flexible motor timing. *Nat. 2025* 1–10 (2025). doi:10.1038/s41586-025-09778-2

58. Li, N., Chen, T. W., Guo, Z. V., Gerfen, C. R. & Svoboda, K. A motor cortex circuit for motor planning and movement. *Nat. 2015 5197541* **519**, 51–56 (2015).

59. Inagaki, H. K., Fontolan, L., Romani, S. & Svoboda, K. Discrete attractor dynamics underlies persistent activity in the frontal cortex. *Nature* **566**, 212–217 (2019).

60. Rudebeck, P. H. & Izquierdo, A. Foraging with the frontal cortex: A cross-species evaluation of reward-guided behavior. *Neuropsychopharmacol. 2021* 1–13 (2021). doi:10.1038/s41386-021-01140-0

61. Kilpatrick, Z. P., Davidson, J. D. & El Hady, A. Uncertainty drives deviations in normative foraging decision strategies. *J. R. Soc. Interface* **18**, (2021).

62. Hanks, T. D., Kiani, R. & Shadlen, M. N. A neural mechanism of speed-accuracy tradeoff in macaque area LIP. *Elife* **3**, e02260 (2014).

63. Hanks, T. D., Mazurek, M. E., Kiani, R., Hopp, E. & Shadlen, M. N. Elapsed Decision Time Affects the Weighting of Prior Probability in a Perceptual Decision Task. *J. Neurosci.* **31**, 6339 (2011).

64. Mochol, G., Kiani, R. & Moreno-Bote, R. Prefrontal cortex represents heuristics that shape choice bias and its integration into future behavior. *Curr. Biol.* **31**, 1234 (2021).

65. Lo, C. C. & Wang, X. J. Cortico-basal ganglia circuit mechanism for a decision threshold in reaction time tasks. *Nat. Neurosci.* **9**, 956–963 (2006).

66. Scott, D. N., Mukherjee, A., Nassar, M. R. & Halassa, M. M. Thalamocortical architectures for flexible cognition and efficient learning. *Trends Cogn. Sci.* **28**, 739–756 (2024).

67. Luo, M., Zhou, J. & Liu, Z. Reward processing by the dorsal raphe nucleus: 5-HT and

beyond. *Learn. Mem.* **22**, 452–60 (2015).

68.	Miyazaki, K., Miyazaki, K. W. & Doya, K. The role of serotonin in the regulation of patience and impulsivity. *Mol. Neurobiol.* **45**, 213–224 (2012).

69.	Miyazaki, K., Miyazaki, K. W. & Doya, K. Activation of dorsal raphe serotonin neurons underlies waiting for delayed rewards. *J. Neurosci.* **31**, 469–479 (2011).

70.	Miller, E. K. The prefontral cortex and cognitive control. *Nat. Rev. Neurosci.* **1**, 59–65 (2000).

71.	Avitan, L. & Stringer, C. Not so spontaneous: Multi-dimensional representations of behaviors and context in sensory areas. *Neuron* **110**, 3064–3075 (2022).

72.	Churchland, M. M. & Shenoy, K. V. Preparatory activity and the expansive null-space. *Nat. Rev. Neurosci.* **25**, 213–236 (2024).

73.	Weber, J. *et al.* Subspace partitioning in the human prefrontal cortex resolves cognitive interference. *Proc. Natl. Acad. Sci. U. S. A.* **120**, (2023).

74.	Hyman, J. M., Ma, L., Balaguer-Ballester, E., Durstewitz, D. & Seamans, J. K. Contextual encoding by ensembles of medial prefrontal cortex neurons. *Proc. Natl. Acad. Sci.* **109**, 5086–5091 (2012).

75.	Siegel, M., Buschman, T. J. & Miller, E. K. Cortical Information Flow During Flexible Sensorimotor Decisions. *Science* **348**, 1352 (2015).

76.	Inagaki, H. K. *et al.* A midbrain-thalamus-cortex circuit reorganizes cortical dynamics to initiate movement. *Cell* **185**, 1065-1081.e23 (2022).

77.	Paton, J. J. & Buonomano, D. V. The Neural Basis of Timing: Distributed Mechanisms for Diverse Functions. *Neuron* **98**, 687 (2018).

78.	Siniscalchi, M. J., Phoumthipphavong, V., Ali, F., Lozano, M. & Kwan, A. C. Fast and slow transitions in frontal ensemble activity during flexible sensorimotor behavior. *Nat. Neurosci.* **19**, 1234–1242 (2016).

79.	Barthas, F. & Kwan, A. C. Secondary Motor Cortex: Where 'Sensory' Meets 'Motor' in the Rodent Frontal Cortex. *Trends Neurosci.* **40**, 181–193 (2017).

80.	Kim, J. H., Daie, K. & Li, N. A combinatorial neural code for long-term motor memory. *Nature* **637**, 663–672 (2025).