Butterfly host plant use driven by microclimate not nutritional quality

William B.V. Langdon, Department of Biology, University of Oxford, Oxford, UK and Epping

Forest Office, Loughton, Essex, UK. ORCID: 0000-0001-8695-403X

Richard Fox, Butterfly Conservation, Manor Yard, East Lulworth, Dorset, UK. ORCID: 0000-

0001-6992-3522

Owen T. Lewis, Department of Biology, University of Oxford, Oxford, UK. ORCID: 0000-0001-

7935-6111

Corresponding author: Owen T. Lewis (owen.lewis@biology.ox.ac.uk)

Acknowledgements

We would like to thank Julian Cooper (Oxford District Council) for permission to carry out

fieldwork at Shotover Country Park and François Bouchereau (Forest Research) for plant

chemistry analyses. Lisbeth Hordley generously provided statistical advice. WL was supported

by the Natural Environmental Research Council through the University of Oxford's

Environmental Research Doctoral Training Programme (NE/L002612/1). RF was supported by

the Heather Corrie Fund from Butterfly Conservation and OL by Brasenose College, University

of Oxford.

Author contributions

Willam B.V. Langdon Conceptualization (equal); Investigation (lead); Formal Analysis (lead);

Writing – Original Draft Preparation (lead); Writing – Review & Editing (equal). **Richard Fox**

Conceptualization (supporting); Supervision (equal); Writing – Original Draft Preparation

(supporting); Writing – Review & Editing (equal). **Owen T. Lewis** Conceptualization (equal);

Supervision (equal); Writing – Review & Editing (equal).

Conflict of interest statement

The authors declare that they have no conflicts of interest.

Insect host plant use driven by microclimate not nutritional quality

Abstract

1. Abundance of insect herbivores often depends on host plant suitability for their specialised

immature stages. Suitability can be strongly influenced by both microclimate and the

nutritional quality of the plants themselves. Where soil nitrogen is high, host plants tend to

have high nutritional quality, but vigorous growth of surrounding vegetation reduces

microclimatic temperatures. Thus, thermophilous insects may face a choice between host

plants with optimal microclimates and those with optimal nutritional quality.

2. We investigated how microclimate and nitrogen content influence oviposition choices by the

declining Small Copper butterfly, *Lycaena phlaeas,* on its host plant, *Rumex acetosa*. We

predicted that warmer plants would have lower nitrogen content, and that butterflies would

choose cooler, high-nitrogen plants during warmer ambient conditions.

3. Although warmer *R. acetosa* plants had lower nitrogen content, *L. phlaeas* consistently

chose to lay eggs on plants in warm microclimates, implying a trade-off between

temperature and the nutritional quality of host plants.

4. Patches of bare ground created by *Talpa europaea* (European Mole) near *R. acetosa* plants

increased microclimatic temperatures and decoupled the negative association between

nutritional quality and thermal suitability.

5. Our results have implications for the conservation of thermophilous insect herbivores,

especially close to their range margins and in the context of climate change. Rather than

maximising host plant abundance or nutritional quality, management that creates suitable

microclimatic conditions is likely to be critical. Our findings also suggest that, while nitrogen

pollution may increase host plant nutritional quality, its negative impacts on microclimate will

likely further reduce breeding habitat for *L. phlaeas* and other insects in grassland habitats.

Keywords: Lepidoptera, nitrogen, oviposition preference, insect decline

INTRODUCTION

The quality of habitat for insects is dictated primarily by the availability of locatons where their least mobile and most specialised early stages can survive (Bourn & Thomas, 2002; Thomas et al., 2011). For many herbivorous insects of open habitats a key factor is the availability of suitably warm microhabitats around host plants, particularly for species close to their cool range margins (Oliver et al., 2009; Thomas et al., 1999). Here, ambient temperatures are often lower than those required to complete development (van Wingerden et al., 1991). With limited ability to thermoregulate behaviourally (Ashe-Jepson et al., 2023) or move between feeding and thermoregulating sites, this means that immature stages can often only survive on host plants growing in particularly warm microhabitats (Curtis & Isaac, 2015; WallisDeVries, 2006). As a result, the availability of host plants in such microhabitats often dictates the abundance of thermophilous insect species (Curtis & Isaac, 2015; J.A. Thomas et al., 2001).

Microhabitat aside, high inter- and intra-specific variability in plant chemistry means that not all host plants are of equal nutritional quality for insects (Persson et al., 2010). Insects require a wide variety of nutrients, each in specific amounts (Raubenheimer et al., 2009; Raubenheimer & Simpson, 1993, 2004). When the chemistry of food does not match these requirements, insects are forced to overeat and excrete excess nutrients to make up for deficiencies of others, at a cost of time and energy (Raubenheimer et al., 2005; Simpson et al., 2004). Thus, host plant nutritional quality is defined as the match between insect requirements and host chemistry (Sterner & Elser, 2002; Vogels et al., 2023), and large mis-matches can have strong effects on fitness that scale up to population-level impacts (Cease et al., 2012; Clissold et al., 2006).

Insects are, therefore, strongly constrained by their thermal and nutritional requirements, and

their fitness will be highest on host plants where both of these are optimal. Such conditions may

be rare, however, if thermally and nutritionally suitable conditions tend not to overlap. For

example, the host plants of many herbivorous insects have a much higher carbon content and a

much lower nitrogen content than the insects themselves (Elser et al., 2000). Thus, to attain

sufficient nitrogen to complete their development, insects must generally overeat large

quantities of carbon. Plant nitrogen content is, therefore, often regarded as a key indicator of

host nutritional quality for insects, with performance generally improved on higher-nitrogen host

plants (Mattson, 1980; Throop & Lerdau, 2004; White, 1993). Plant nitrogen content typically

increases where soil nitrogen levels are higher (Fujita et al., 2013; Sardans et al., 2012; Throop

& Lerdau, 2004) but, in open habitats, higher soil nitrogen also enhances vegetation

productivity, leading to taller vegetation, with reduced penetration of sunlight to ground level

(Bobbink, 1991; DeMalach et al., 2017; Hautier et al., 2009). Microhabitat temperatures in such

situations are lower than in more open locations where bare ground and dead plant material

exposed to sunlight warm up rapidly and radiate heat (Curtis & Isaac, 2015; WallisDeVries,

2006). Therefore, in general, host plants with higher nutritional quality (nitrogen content) tend

not to overlap with warm microhabitats, creating a potential trade-off for thermophilous insects in

open habitats.

Such trade-offs have important implications for the effective conservation of insect populations,

especially in the context of long-term declines in abundance of many terrestrial insect taxa

around the world (Edwards et al., 2025; Van Klink et al., 2020; Warren et al., 2021). Both local

habitat management and wider-scale policy instruments can influence insects' exposure to

changes in microclimate and nitrogen availability, so understanding species' responses to these

environmental variables will directly inform effective conservation. Furthermore, anthropogenic

climate change and increasing diffuse nitrogen pollution are acknowledged global and regional

drivers of insect population change (Carvalheiro et al., 2020; Hill et al., 2021; Roth et al., 2021;

Wilson & Fox, 2021).

The roles of thermal and nutritional requirements in dictating insect microhabitat use are rarely

studied simultaneously, however (e.g. Bourn & Thomas, 1993; Ellis, 2003). As a result, it is

difficult to understand the relative strength of thermal and nutritional constraints for many insect

herbivores, and whether our hypothesised trade-off exists. Such an understanding is important

when climate change and nitrogen pollution are simultaneously altering the thermal and

nutritional conditions experienced by insects (Rashid et al., 2023). Rising ambient temperatures

under climate change may lessen thermal constraints on some insect species, as microhabitats

that were previously too cool become sufficiently warm (Roy & Thomas, 2003; C.D. Thomas et

al., 2001). On the other hand, warming extends the growing season (Menzel et al., 2020),

enhancing plant productivity and potentially leading to cooler microhabitats overall

(WallisDeVries & van Swaay, 2006). Rising levels of soil nitrogen under nitrogen pollution are

expected to enhance plant productivity and alter vegetation structure alongside warming

(WallisDeVries & van Swaay, 2006), while also leading to increases in plant nitrogen content

and changes in host nutritional quality for insect herbivores (Vogels et al., 2023).

Thus, the impacts of two major anthropogenic stressors on insect populations largely depend on

the interplay of changing host nutritional quality, vegetation structure and microclimate with

insect thermal and nutritional requirements. If there is a trade-off between using host plants with

high nutritional quality and those growing in warm microhabitats, species requiring warmer

conditions may be unable to access the potential benefits of higher host nutritional quality driven

by nitrogen pollution, unless higher macroclimatic temperatures enable them to use cooler

microhabitats.

Here, we investigate how host plant nutritional quality (nitrogen content) and microclimate shape

habitat use by a grassland butterfly, *Lycaena phlaeas* (Small Copper), and the relationship

between these two aspects of habitat quality in this system. Specifically, by studying the

distribution of eggs and larvae in a *L. phlaeas* population, we assess the evidence for our

hypothesised trade-off between high nutritional quality host plants and high temperatures

around host plants, and how differences in vegetation structure bring this about. Female insects

generally select host plants that maximise larval survival (Gripenberg et al., 2010; Jaenike,

1978), and the availability of hosts that meet larval requirements has consistently been shown to

be the main factor constraining butterfly populations in similar systems (Bourn & Thomas, 2002;

Thomas et al., 2011). Thus, we expect female oviposition choice, measured by the distribution

of eggs, to indicate how *L. phlaeas* populations are constrained by thermal and nutritional

requirements. By replicating sampling across seasons, we also attempt to understand how

these variables are related as macroclimate changes.

Specifically, we test the following hypotheses:

1.　The nutritional quality of host plants is negatively correlated with the temperature under

them (microclimate), presenting a potential trade-off for *L. phlaeas* oviposition.

2.　In early summer, when macroclimatic temperatures are relatively high, microclimate is less

important in dictating egg-laying patterns, and host plants with higher nutritional quality are

preferentially selected.

3.　In autumn, when macroclimatic temperatures are lower than in early summer, *L. phlaeas*

egg-laying is determined more by microclimate around host plants than by their nutritional

quality.

METHODS

Study system

Lycaena phlaeas is a widespread lycaenid butterfly in open habitats across Eurasia and North America. It is oligophagous on *Rumex* spp., but in north-west Europe it mostly uses *Rumex acetosa* (Common Sorrel) and *Rumex acetosella* (Sheep's Sorrel) (Thomas & Lewington, 2014). In the United Kingdom, *L. phlaeas* undergoes a winter diapause at the larval stage and usually has between one and three generations per year depending on latitude (Thomas & Lewington, 2014). While still widespread and not categorised as threatened, it declined in abundance by 39% in the UK between 1976 and 2019 (Fox et al., 2022). Like many temperate butterfly species, *L. phlaeas* requires warm microhabitats (Streitberger et al., 2014) but will also respond strongly to changes in host nutritional quality in laboratory experiments. In oligotrophic and mesotrophic environments, *L. phlaeas* larvae appear to be nitrogen-limited, and benefit from moderate increases in plant nitrogen content, before performance declines at high nitrogen levels (Kurze et al., 2018; Langdon, 2024). Thus, they have high potential to be impacted by changes in vegetation structure, which alter microclimates, but also by changes in plant nitrogen content. Predicting the likely outcome of these changes requires an understanding of the relative strength of their thermal and nutritional constraints.

Data collection

Our study took place at Shotover Country Park (Oxfordshire, United Kingdom, 51°44′42″N, 001°11′01″W). This site is a typical habitat for *L. phlaeas* in lowland UK, a mesotrophic grassland where the butterfly uses *R. acetosa* as its sole larval host plant and has three generations per year. *Rumex acetosa* plants were searched systematically across the study area for *L. phlaeas* eggs or larvae. For each occupied plant a paired control plant was then selected. The control plant was the nearest unoccupied *R. acetosa* plant in a randomly selected cardinal direction (based on randomly generated number between 1 and 4 corresponding to

North, East, South or West respectively). For both the occupied and control plant in each pair, a

50cm quadrat with 10cm grids was used to measure the surrounding percentage cover of living

vegetation, bare ground and leaf litter.

Sward height around each plant in each pair was measured following Stewart et al. (2001),

using a ruler to read the height below which an estimated 80% of vegetation was growing. The

area of each plant was estimated as the dimensions of the tightest-fitting rectangle, i.e. the

longest dimension of the plant viewed from above, and the maximum dimension perpendicular

to this. The three largest leaves without *L. phlaeas* eggs or larvae were then removed from each

plant for chemical analysis, and their length and breadth measured and multiplied to

approximate their area, which was averaged across the three leaves as a measure of leaf size.

Leaf samples were then frozen in glassine envelopes at -10°C until further analysis. The

microclimate of each plant was measured as the temperature at the soil surface directly under

the plant in sunny conditions between 12:00 and 14:30 (the hottest part of the day, when

differences between plants were expected to be most apparent) using an ultrafine thermocouple

(TC Direct Ultra Fine Wire PFA type K Thermocouple 0.08mm diameter, with handheld Type K

Thermocouple Indicator; TC Direct, Uxbridge, United Kingdom). The same thermocouple was

also used to measure ambient temperature at approximately 1.8m above the ground. This

fieldwork was carried out both in early summer (10-16 June 2023) and early autumn (21-28

September 2023), when thermal constraints were expected to differ due to differences in

ambient (macroclimatic) temperature.

Chemical analysis

Cost limitations meant that we were unable to analyse the nutritional quality (in terms of nitrogen

content) of every plant sample. Instead, we aimed to identify reliable proxies for plant nitrogen

content that we could use in our analysis. Langdon (2024) found that the area of *R. acetosa*

plants and the size of their leaves were correlated with plant nitrogen content. We therefore

performed a principal component analysis of average leaf size and plant area across all

sampled plants and separated plants into 30 groups based on the first principal component.

Within these, we made a further two groups based on sampling season (i.e. 60 groups in total,

one for each season by principal component group combination). From each of these we

randomly selected one plant for chemical analysis. This gave a total of 58 plants (as two group x

season combinations lacked plants). We expected that analysing leaf samples across the full

range of plant morphology (plant area and leaf size) would provide the best chance of detecting

relationships between these proxy measurements and plant nitrogen content.

Selected leaf samples were dried in individual glassine envelopes at 70°C for 48 hours before

being transferred to microtubes and ground to a fine powder via bead beating (MP Bio

FastPrep-24 5G, MP Biomedicals, Irvine, USA). Leaf samples were beaten for two minutes with

three ceramic beads of diameter 2.3mm (Zirconia/Silica, Thistle Scientific, Warwickshire,

England). Approximately 10mg of each leaf sample was then analysed for carbon and nitrogen

content via combustion using a Thermo Flash Carlo Erba CN analyser (EA 1112 series)

(Thermo Fisher Scientific, Waltham, Massachusetts, USA) by Forest Research (Alice Holt,

United Kingdom).

Statistical analysis

To identify which of plant area, average leaf size, or a combination of the two was the most

appropriate proxy for plant nitrogen content, we ran models of plant nitrogen content as a

function of average leaf size and plant area (separately and together) and the first and second

principal components of their PCA (alone, but not together, since this would be the equivalent to

using both unmodified variables), all with a dummy 'season' (summer or autumn) variable to

control for differences between seasons. We compared the performance of these models via

their AIC, BIC and R^2, as well as calculating the Pearson correlation coefficients between each

individual proxy and nitrogen content, looking for a combination of high explanatory power (a

robust proxy) and parsimony (i.e. one variable that performs almost as well as two would be

preferred, for easier interpretation of later modelling steps).

Lycaena phlaeas habitat use was analysed across seasons using two linear models comparing

the measurements of plant temperature and host quality for occupied and unoccupied plants in

early summer and autumn. For each variable we thus ran a model with the following structure:

*Variable for Plant $_i$ ~ Occupancy Status of Plant $_i$ * Season in which plant $_i$ was sampled.*

Autumn was used as the reference level for season, so that estimates for the seasonal

interaction indicate the change in the environmental variables of occupied plants from autumn to

early summer.

Temperature measurements were made over several days with varying weather conditions in

each field season. To account for the effect of daily variation in ambient temperature, and its

influence on the microclimate temperatures measured under plants, we centred and scaled our

temperature measurements by date. Thus, microclimate temperature was relative to others

measured on that date, and our model indicated whether *L. phlaeas* was using the warmest

plants, but not how warm these actually were. Our host nutritional quality proxy was also

centred and scaled to match this. We compared the results of this model to one using the

scaled predicted nitrogen content of plants instead of our proxy for this, with the inverse of the

scaled standard error of predicted nitrogen content as a weight.

To test the hypothesis that host plant microclimate and nutritional quality are negatively related,

we modelled the relationship between temperature and host nutritional quality using a linear

model. During data collection we noted that many of the plants used for *L. phlaeas* egg-laying

were on bare ground created by *Talpa europaea* (European Mole) activity (as also observed by

Streitberger et al., (2014)). This could produce warm microclimates under plants that would

otherwise have had cool microclimates and high plant nitrogen content (in areas with high soil

nitrogen and dense vegetation), thereby disrupting the hypothesised relationship. We therefore

included an interaction between presence or absence of bare ground and host nutritional quality

in this model. As for the egg-laying model, we also ran this model using the scaled predicted

nitrogen content of plants instead of our proxy measure, with the inverse of the scaled standard

error of predicted nitrogen content as a weight.

To understand what might drive the hypothesised relationship between temperature and host

nutritional quality we then examined the effect of environmental variables (bare ground,

vegetation cover, leaf litter cover and sward height) on relative microclimate temperature using

linear models. Finally, we tested the effect of sward height and litter cover on host nutritional

quality in the same way (omitting bare ground and vegetation cover and dropping plants with

bare ground from the analysis since these mostly reflected *T. europaea* disturbance rather than

underlying soil conditions).

All analyses were carried out in R (R Core Team, 2024) using the 'tidyverse', 'readr' and 'Hmisc'

packages for data manipulation (Harrel, 2024; Wickham et al., 2019, 2024), 'performance' for

model comparison (Lüdecke et al., 2021), and 'ggplot2', 'cowplot' and 'ggeffects' for plotting

(Lüdecke, 2018; Wickham, 2016; Wilke, 2024). Model fit was assessed via inspection of plots of

the residuals and fitted values.

RESULTS

Host nutritional quality and proxies

Among our 58 *R. acetosa* leaf samples, nitrogen content (by mass) averaged 2.54% (sd = 0.76)

with a maximum of 4.39% and a minimum of 0.98%. This translated into molar content

averaging 1815.79 micromoles g^{-1} (sd = 539.03) with a maximum of 3130.64 micromoles g^{-1} and

a minimum of 701.09 micromoles g^{-1}. This led to molar C:N ratios of between 11.24 and 51.98,

averaging 22.41 (sd = 7.67), which is higher than the optimal value for *L. phlaeas* larvae of 18.1

estimated in previous research (Langdon, 2024). The distribution of these measurements is

plotted in Figure S1, although the stratified nature of our sampling means that these are not

representative of the whole population of *R. acetosa* at the study site.

Table 1: Results of Principal Components analysis of host nutritional quality proxies.

	PC1	PC2
Eigenvalue	1.59	0.42
% Variance Explained	79.25	20.75
Average Leaf Size (mm^2) loading	-0.71	-0.71
Plant Area (cm^2) loading	-0.71	0.71

Plant area and leaf size were positively correlated (Pearson's R = 0.61, t = 13.54, df = 316, p <

0.001) and the first principal component of their PCA explained 79.25% of the variance in the

two measurements (Table 1). Models of plant nitrogen content as a function of these proxies

and principal components (results in Table S1 and Figure S2) indicated that leaf size was the

best proxy for nitrogen content. Of the individual options, it showed the strongest positive

correlation with plant nitrogen content (Table 2), and models with just leaf size performed better

than either principal component alone (Table 3). The model with leaf size and plant area

performed only marginally better than leaf size alone (Table 3). Therefore, on the basis of

parsimony and predictive power, leaf size was chosen as our proxy for plant nitrogen content.

The results were qualitatively the same when C:N ratio was used instead of N content as the

'true' measure of host nutritional quality (Table S2 and Table S3).

Table 2: Pearson's correlation coefficients and significance tests for correlation of possible plant

proxies with plant nitrogen content (measured in micromoles g^{-1}).

Proxy	R	t	DF	p	Lower Confidence Interval	Upper Confidence Interval
Average Leaf Size (mm^2)	0.62	5.94	56	<0.001	0.43	0.76
Plant Area (cm^2)	0.37	2.96	56	<0.001	0.12	0.57
Quality PC1	-0.55	-4.98	56	<0.001	-0.71	-0.34
Quality PC2	-0.23	-1.76	56	0.08	-0.46	0.03

288   **Table 3:** Comparison of models relating plant nitrogen content to different possible proxies.

Proxies	AIC	BIC	R^2
Average Leaf Size (mm^2) + Plant Area (cm^2)	873	883	0.41
Average Leaf Size (mm^2)	871	879	0.41
Plant Area (cm^2)	883	891	0.27
Quality PC1	874	882	0.37
Quality PC2	892	900	0.15

Microclimate under _R. acetosa_ plants

Ambient temperatures in early summer averaged 24.74°C (sd = 0.77), and in autumn 16.34°C

(sd = 1.47), approximately 8.40°C lower (t = 60.42, df = 185.49, p < 0.001) (Figure S3).

Matching this, microclimate temperatures measured under _R. acetosa_ plants were higher in

early summer, averaging 36.68°C (sd = 6.17) compared to 18.42°C (sd = 1.64) in autumn, a

difference of 18.26°C that was also statistically significant (t = 38.34, df = 217.53, p < 0.001)

(Figure S3). The maximum temperature recorded under _R. acetosa_ plants in early summer was

56°C and in autumn 23°C.

Impacts of host plant nutritional quality and microclimate on _L. phlaeas_ egg-laying

Overall, 318 plants were sampled, 184 in early summer and 134 in the autumn (evenly divided

into occupied plants and control plants). Occupied host plants were significantly warmer than

unoccupied host plants (β = 1.22 ± 0.14, t = 8.83, df = 314, p < 0.001), but there was no

difference in their nutritional quality (β = 0.15 ± 0.16, t = 0.89, df = 314, p = 0.37) (Figure 1,

Table 4). This pattern was observed in both early summer and autumn, and the differences in

thermal and nutritional quality of occupied and unoccupied plants were constant across seasons

(Figure 1, Table 4). Results for plant quality were qualitatively the same when scaled predicted

nitrogen content of plants was used instead of the proxy of leaf size (Table S4).

On occupied plants, the mean number of eggs in early summer was 2.32 (sd = 2.82) with a

maximum of 16 and a total (across all plants) of 213. In autumn, these figures were 1.90 (sd =

1.54), 8 and 127 eggs respectively. The median in both seasons was 1 egg.

315 **Table 4:** Outputs from models assessing differences in nutritional quality (scaled leaf size) and

316 microclimate (scaled soil surface temperature) of occupied and unoccupied *L. phlaeas* host

317 plants in early summer and autumn.

Measurement	Term	Estimate	Standard Error	t	p
Scaled Leaf Size	Intercept	0.32	0.12		
	Occupied	0.15	0.16	0.89	0.37
	Season	-0.68	0.15	-4.48	<0.001
	Occupied:Season	0.01	0.21	0.06	0.95
Scaled soil surface temperature	Intercept	-0.61	0.1		
	Occupied	1.22	0.14	8.83	<0.001
	Season	0.05	0.13	0.36	0.72
	Occupied:Season	-0.09	0.18	-0.51	0.61

325   **Figure 1:** Mean values of scaled leaf size and soil surface temperature measured on and

326   around occupied and unoccupied host plants of *L. phlaeas* in early summer and autumn. Points

327   show predicted marginal effects from linear models (Table 4) and bars their 95% confidence

328   intervals.

Relationships between microclimate and host nutritional quality

As expected, in the absence of bare ground host nutritional quality (scaled leaf size) was significantly negatively related to scaled soil surface temperature under plants (β = -0.30 ± 0.08, t = -3.94, df = 313, p < 0.001). At the lowest leaf sizes plants with bare ground were significantly warmer than those without bare ground (β = 0.80 ± 0.11, t = 7.13, df = 313, p < 0.001), and the presence of bare ground (associated with *T. europaea* activity) made the slope of the relationship between host nutritional quality and scaled temperature under plants significantly less negative, almost entirely flattening it (β = 0.28 ± 0.11, t = 2.62, df = 313, p < 0.01) (Figure 2). Results did not differ qualitatively when using predicted nitrogen content as the measure of host nutritional quality (see Table S5).

Figure 2: The modelled relationship between host nutritional quality (measured using the proxy

of leaf size), microclimate and the presence or absence of bare ground. Predictions from linear

model are plotted with their 95% confidence intervals and the underlying data.

Relationships between microclimate and vegetation structure

Variables characterising vegetation structure around *R. acetosa* plants were all significantly

associated with scaled soil surface temperature in the manner expected (Figure 3). Cover of

bare ground (β = 0.02 ± 0.0028, t = 7.15, df = 315, p < 0.001) and cover of litter (β = 0.018 ±

0.0046, t = 3.95, df = 315, p < 0.001) were positively associated with scaled soil surface

temperature, and sward height (β = -0.04 ± 0.0036, t = -10.33, df = 314, p < 0.001) and cover of

vegetation (β = -0.02 ± 0.0037, t = -6.50, df = 315, p < 0.001) were negatively associated.

Individually, sward height explained more variation in microclimate than any other variable (R^2 =

0.25), followed by bare ground (R^2 = 0.14), vegetation cover (R^2 = 0.12) and litter cover (R^2 =

0.05).

Figure 3: Relationships between vegetation structure around host plants and the relative

temperatures measured under them. Predicted trends from linear models are shown with their

95% confidence intervals and the underlying data.

Relationships between vegetation structure and host nutritional quality

Litter cover and sward height were both related to our host nutritional quality proxy (average leaf size) in the manner expected. The association for litter cover was significantly negative (β = -9.14 ± 2.00, t = -4.56, df = 212, p < 0.001) and for sward height positive (but not significantly so; β = 0.272 ± 2.52, t = 0.11, df = 211, p = 0.91). The litter model therefore explained more variation in host nutritional quality than sward height (litter R^2 = 0.09, sward height R^2 = 0.0000055).

DISCUSSION

Insect herbivores are expected to be constrained by both the nutritional quality and the microclimate of their host plants. We predicted (hypothesis 1) that, in open habitats, insects may face a trade-off between these constraints, because plants with high nutritional quality (nitrogen content) are expected to occur in cooler microclimates. In our study of egg-laying patterns of the widespread butterfly *L. phlaeas*, we found strong support for this hypothesis in the form of a significant negative relationship between host plant nitrogen content and microclimate (although this relationship was fundamentally altered if *T. europaea* activity generated bare ground close to host plants). However, we found no evidence that the butterfly's egg-laying decisions shift between seasons to favour host plant nutritional quality when thermal constraints are eased by higher ambient temperatures (our hypotheses 2 and 3). Egg-laying of *L. phlaeas* on *R. acetosa* was primarily determined by microclimate. We found no significant effect of host nutritional quality (proxies of nitrogen content) on the probability of egg-laying, and no difference between a warmer period (in early summer) and a cooler one (in early autumn).

Given that most plants in our sample appeared to be of sub-optimal nutritional quality for *L. phlaeas* (Figure S4) (Langdon, 2024), and evidence that other butterfly species are able to

select host plants on the basis of nutritional quality (Bourn & Thomas, 1993; Letourneau & Fox,

1989; Prudic et al., 2005), it appears that female *L. phlaeas* are missing an opportunity to

maximise larval fitness by laying their eggs on the minority of plants with higher nutritional

quality. Instead, microclimate appears to be the dominant driver of oviposition choice and,

unless there happened to be soil disturbance under a host plant, plants with warmer

microclimates were of lower nutritional quality (as hypothesised).

In our system ovipositing *L. phlaeas* face a choice between high nutritional quality host plants or

warm host plants, adding to previous work showing that egg-laying butterflies may have to

balance nutritional requirements against thermal requirements (Krämer et al., 2012). The

apparent prioritisation of the latter implies that thermal constraints are strong for *L. phlaeas*, in

line with previous work (León-Cortés et al., 2000; Streitberger et al., 2014) and override

nutritional constraints. Many other studies have highlighted the importance of thermal

constraints on habitat use by butterfly species but without examining host plant nutritional

quality (e.g. Curtis & Isaac, 2015; Johansson et al., 2024; Roy & Thomas, 2003; Thomas et al.,

1986; C.D. Thomas et al., 2001; Wilby et al., 2024). Although we do not have data relating *L.*

phlaeas larval performance to temperature, experimental work (Langdon, 2024) suggests that

although most *R. acetosa* plants in our system were of sub-optimal nutritional quality for *L.*

phlaeas (Figure S4), they were still within a range where larval survival was around 80% of the

maximum. Furthermore, by feeding selectively on different plants or plant parts to regulate their

nutrient intake, larvae in our study system might be able to buffer the impacts of low host

nutritional quality. Thus, the fitness consequences of using hosts with low nutritional quality at

our study site are likely to be relatively small for *L. phlaeas*. A further intriguing possibility is that

the fitness costs associated with sub-optimal food quality may also be reduced by high

temperatures. Clissold et al. (2013) found that the efficiency of macronutrient absorption varied

with temperature for *Locusta migratoria* nymphs, and that protein absorption was more efficient

at higher temperatures for nymphs feeding on *Themeda triandra*. Protein is the main pool of

nitrogen in plants, so increased absorption efficiency on warm host plants could offset the cost

of their lower nitrogen levels. Similar work is lacking for Lepdioptera, however, and Clissold et

al. (2013) did not observe the same trend in locusts fed *Triticum aestivium*, suggesting that such

increases in absorption efficiency are unlikely to be universal.

In spite of the far higher ambient temperatures and microclimatic temperatures around *R.*

acetosa in early summer, we found no support for our hypothesis that *L. phlaeas* would shift

oviposition choices to use hosts with higher nutritional quality in relatively cooler microclimates

in early summer compared to autumn. This contrasts with previous work on another lycaenid

butterfly, *Polyommatus bellargus*, in which higher ambient temperatures in early summer

allowed the use of larval host plants growing in cooler microhabitats, doubling habitat availability

(Roy & Thomas, 2003). This difference is difficult to explain and seems unlikely to reflect a

difference in thermal constraints given that *L. phlaeas* occurs north into the Arctic Circle (Kudrna

et al., 2011) while *P. bellargus* reaches the northern edge of its range in southern England,

where it is dependent on conservation management to provide suitably warm microhabitats

(Thomas, 1983).

At our study site, the cooling effect of the tall swards (mean = 23.30 cm, sd = 11.26) may have

meant that only a small proportion of *R. acetosa* plants were thermally suitable for *L. phlaeas*. In

such mesotrophic grasslands, *L. phlaeas* may breed mainly in small habitat islands of disturbed

ground or low fertility soil where host plant microclimates are suitably warm. Even when ambient

temperatures are high, e.g. in early summer, and thermal constraints are expected to be

lessened, there may be little other suitable habitat for the butterfly to use because most host

plants are still too cool. Indeed, we often observed that egg-laying took place on plants growing

on bare ground generated by *T. europaea* disturbance, which was positively associated with the

temperature under host plants. In fact, 55% of plants with *L. phlaeas* eggs had some bare

ground within 25cm of them, compared to only 11% of unoccupied plants. These results

reinforce the importance of soil disturbance for creating microhabitat heterogeneity that impacts

grassland plant and insect communities (e.g. Seifan et al., 2010). Streitberger et al. (2014) also

observed high rates of host plant use by *L. phlaeas* on *T. europaea* earthworks, while

Streitberger & Fartmann (2013) found the same for the hesperiid *Pyrgus malvae*. Soil

disturbance caused by foraging behaviour of *Sus scrofa* (Wild Boar) has similarly been

positively linked to butterfly oviposition choice by *P. malvae* (de Schaetzen et al. 2018) and two

fritillary species (Scherer et al., 2025). The dependence of *L. phlaeas* on small islands of warm

microhabitat may explain why the response to host plant quality that we documented differs

from that found for *P. bellargus* (Roy & Thomas, 2003), where habitats often have uniformly high

thermal suitability as a result of heavy grazing. It also reinforces the importance of continued

conservation management of open habitats to create suitable microclimates for thermally-

constrained species, in spite of the potential benefits of climate change to these species

(O'Connor et al., 2014). In a broader sense, habitat management to create microclimatic

heterogeneity will help to buffer a wide variety of species from the impacts of extreme weather

events (Hayes et al., 2024) and climate change (Suggitt et al., 2018).

We expected that higher nitrogen content in host plants would reflect higher soil nitrogen levels,

which would in turn enhance plant productivity, leading to denser vegetation and lower

temperatures around host plants. We found some support for this mechanism, with strong

associations between measures of vegetation structure and temperature (shorter, more open

vegetation meant higher temperatures) but only weak relationships in the directions expected

between vegetation structure and host nutritional quality – higher litter cover was significantly,

albeit weakly, associated with lower nutritional quality, while higher sward height was positively,

but not significantly, related to host nutritional quality. Despite this, it is difficult to envisage

another mechanism that could alter host nutritional quality and vegetation structure to affect

microclimate simultaneously. Grazing or cutting often does the latter, but the most commonly-

reported plant response to grazing appears to be an increase in nitrogen uptake and availability

in young plant tissues (e.g. Jaramillo & Detling, 1988; Ruess, 1984), which would lead to a

positive correlation between temperature and host nutritional quality, rather than the inverse as

observed here. The rather weak relationships that we detected between host nutritional quality

and vegetation structure may reflect the fact that our measures of vegetation structure, though

commonly used approximations, imperfectly capture the factors driving the link between host

nutritional quality and microclimate. Further work relating a more detailed characterisation of

vegetation structure to host nutritional quality and, crucially, measuring underlying soil nitrogen

levels would help to establish whether this trade-off is indeed a general phenomenon for

grassland insects.

Our results indicate that habitat use by *L. phlaeas* is driven by thermal requirements and not by

nutritional requirements, and egg-laying females may be forced to use host plants of suboptimal

nutritional quality. This provides useful context for laboratory experiments showing strong

impacts of plant nitrogen content on the performance of *L. phlaeas*, with negative implications

for fitness and survival under levels of nitrogen pollution associated with management of

agricultural grasslands (Kurze et al., 2018). Our results suggest that such responses may not

scale up to impacts of nitrogen pollution on *L. phlaeas* populations in the field via changes in

plant nutritional quality, because high-nitrogen plants will be thermally unsuitable. Thus, *L.*

phlaeas and species with similar thermal requirements seem more likely to be negatively

impacted by changes to vegetation structure under nitrogen pollution, than via changes in host

nutritional quality (Nijssen et al., 2017). This aligns with previous work by Klop et al. (2015), who

performed a similar study with the grassland butterfly *Lasiommata megera,* assessing larval

microhabitat requirements in the field and responses to plant nitrogen content under laboratory

conditions. They found only positive effects of fertilisation, and a tendency to use warmer

microhabitats, suggesting that the negative effects of atmospheric nitrogen deposition on *L.*

megera population trends were likely to reflect losses of suitable microhabitats, offsetting any

improvement in host quality. Similarly, Yang et al. (2017) tested the response of larvae of the

moth *Gynaephora alpherakjj* to nitrogen fertilisation of foodplants in alpine grassland and found

that growth rate and feeding rate declined with fertilisation in the field, despite positive impacts

in the laboratory, attributing this difference to the lower temperatures at ground level in fertilised

plots. Interestingly, however, we found that soil disturbance by *T. europaea* disrupted the trade-

off between host nutritional quality and temperature around plants (by creating high

temperatures around plants of high nutritional quality), suggesting that soil disturbance

generated in this manner, or at larger scales by grazing animals, may contribute to maintaining

insect herbivore species of early-successional habitats under nitrogen pollution.

In conclusion, we found that *L. phlaeas* was strongly thermally constrained, and selection of

larval host plants for egg-laying was determined by warm microclimates rather than host

nutritional quality, even during warmer periods of the year. Indeed, these constraints may force

L. phlaeas to use plants of lower nutritional quality, as we observed a negative association

between host nutritional quality and temperatures around plants. We suggest this could be a

general trade-off faced by grassland insect species, driven by underlying variation in soil

nitrogen and its impact on host nutritional quality and vegetation structure, but further

investigation is needed to confirm the mechanisms underlying this interesting pattern. Overall,

this places the strong responses of herbivores, such as *L. phlaeas*, to variation in host

nutritional quality in fertilization experiments into context; under field conditions, thermophilous

insects are more likely to be negatively impacted by nitrogen pollution via changes in vegetation

structure than positively via increased host nutritional quality. As a result, management to

maintain warm microhabitats will be essential to their survival even under a warming climate.

REFERENCES

Ashe-Jepson, E., Hayes, M.P., Hitchcock, G.E., Wingader, K., Turner, E.C. & Bladon, A.J.

(2023) Day-flying lepidoptera larvae have a poorer ability to thermoregulate than adults. Ecology

and Evolution, 13(10), 1–13. https://doi.org/10.1002/ece3.10623

Bobbink, R. (1991) Effects of nutrient enrichment in Dutch chalk grassland. Journal of Applied

Ecology, 28, 28–41. https://www.jstor.org/stable/2404111

Bourn, N.A.D. & Thomas, J.A. (1993) The ecology and conservation of the brown argus butterfly

Aricia agestis in Britain. Biological Conservation, 63(1), 67–74. https://doi.org/10.1016/0006-

3207(93)90075-C

Bourn, N.A.D. & Thomas, J.A. (2002) The challenge of conserving grassland insects at the

margins of their range in Europe. Biological Conservation, 104, 285–292.

https://doi.org/10.1016/S0006-3207(01)00193-8

Carvalheiro, L.G., Biesmeijer, J.C., Franzén, M., Aguirre-Gutiérrez, J., Garibaldi, L.A., Helm, A.,

Michez, D., Pöyry, J., Reemer, M., Schweiger, O., Leon van den, B., WallisDeVries, M.F. &

Kunin, W.E. (2020) Soil eutrophication shaped the composition of pollinator assemblages during

the past century. Ecography, 43, 209–221. https://doi.org/10.1111/ecog.04656

Cease, A.J., Elser, J.J., Ford, C.F., Hao, S., Kang, L. & Harrison, J.F. (2012) Heavy livestock

grazing promotes locust outbreaks by lowering plant nitrogen content. Science, 335(6067), 467–

469. https://doi.org/10.1126/science.1214433

Clissold, F.J., Coggan, N. & Simpson, S.J. (2013) Insect herbivores can choose microclimates

to achieve nutritional homeostasis. Journal of Experimental Biology, 216(11), 2089–2096.

https://doi.org/10.1242/jeb.078782

Clissold, F.J., Sanson, G.D. & Read, J. (2006) The paradoxical effects of nutrient ratios and

supply rates on an outbreaking insect herbivore, the Australian plague locust. Journal of Animal

Ecology, 75(4), 1000–1013. https://doi.org/10.1111/j.1365-2656.2006.01122.x

Curtis, R.J. & Isaac, N.J.B. (2015) The effect of temperature and habitat quality on abundance

of the Glanville fritillary on the Isle of Wight: implications for conservation management in a

warming climate. Journal of Insect Conservation, 19(2), 217–225.

https://doi.org/10.1007/s10841-014-9738-1

de Schaetzen, F., van Langevelde, F. & Wallisdevries, M.F. (2018) The influence of wild boar

(*Sus scrofa*) on microhabitat quality for the endangered butterfly *Pyrgus malvae* in the

Netherlands. Journal of Insect Conservation, 22(1), 51–59. https://doi.org/10.1007/s10841-017-

0037-5

DeMalach, N., Zaady, E. & Kadmon, R. (2017) Light asymmetry explains the effect of nutrient

enrichment on grassland diversity. Ecology Letters, 20(1), 60–69.

https://doi.org/10.1111/ele.12706

Edwards, C.B., Zipkin, E.F., Henry, E.H., Haddad, N.M., Forister, M.L., Burls, K.J., Campbell,

S.P., Crone, E.E., Diffendorfer, J., Douglas, M.R., Drum, R.G. et al. (2025) Rapid butterfly

declines across the United States during the 21st century. Science, 387(6738), 1090–1094.

Ellis, S. (2003) Habitat quality and management for the northern brown argus butterfly *Aricia*

artaxerxes (Lepidoptera: Lycaenidae) in North East England. Biological Conservation, 113(2),

285–294. https://doi.org/10.1016/S0006-3207(02)00376-2

Elser, J.J., Fagan, W.F., Denno, R.F., Dobberfuhl, D.R., Folarin, A., Huberty, A., Interlandi, S.,

Kilham, S.S., McCauley, E., Schulz, K.L., Siemann, E.H. & Sterner, R. W. (2000) Nutritional

constraints in terrestrial and freshwater food webs. Nature, 408(6812), 578–580.

https://doi.org/10.1038/35046058

Fox, R., Dennis, E.B., Brown, A. & Curson, J. (2022) A revised Red List of British butterflies.

Insect Conservation and Diversity 15, 485–495. https://doi.org/10.1111/icad.12582

Fujita, Y., Van Bodegom, P.M. & Witte, J.P.M. (2013) Relationships between nutrient-related

plant traits and combinations of soil N and P fertility measures. PLoS ONE, 8(12).

https://doi.org/10.1371/journal.pone.0083735

Gripenberg, S., Mayhew, P.J., Parnell, M. & Roslin, T. (2010) A meta-analysis of preference-

performance relationships in phytophagous insects. Ecology Letters, 13(3), 383–393.

https://doi.org/10.1111/j.1461-0248.2009.01433.x

Harrel, J. F. (2024). Hmisc: Harrell Miscellaneous. R Package Version 5.1-3. https://cran.r-

project.org/package=Hmisc

Hautier, Y., Niklaus, P. A. & Hector, A. (2009) Competition for light causes plant biodiversity loss

after eutrophication. Science, 324(5927), 636–638. https://doi.org/10.1126/science.1169640

Hayes, M.P., Ashe-Jepson, E., Hitchcock, G.E., Clark, R., Hellon, J., Knock, R.I., Bladon, A.J. &

Turner, E.C. (2024) Heatwave predicts a shady future for insects: impacts of an extreme

weather event on a chalk grassland in Bedfordshire, UK. Journal of Insect Conservation, 28(5),

923–933. https://doi.org/10.1007/s10841-024-00556-5

Hill, G.M., Kawahara, A.Y., Daniels, J.C., Bateman, C.C. & Scheffers, B.R. (2021) Climate

change effects on animal ecology: Butterflies and moths as a case study. Biological Reviews,

96(5), 2113–2126. https://doi.org/10.1111/brv.12746

Jaenike, J. (1978). On optimal oviposition behavior in phytophagous insects. Theoretical

Population Biology, 14(3), 350–356. https://doi.org/10.1016/0040-5809(78)90012-6

Jaramillo, V. J. & Detling, J. K. (1988) Grazing history, defoliation, and competition: Effects on

shortgrass production and nitrogen accumulation. Ecology, 69(5), 1599–1608.

https://doi.org/10.2307/1941657

Johansson, V., Säwenfalk, D.S., Bergman, K.O., Kindvall, O., Askling, J. & Franzén, M. (2024)

Oviposition preferences and larval survival of the marsh fritillary butterfly: The adverse impact of

grazing. Insect Conservation and Diversity, 17(4), 642–650. https://doi.org/10.1111/icad.12734

Klop, E., Omon, B. & Wallisdevries, M.F. (2015) Impact of nitrogen deposition on larval habitats:

the case of the Wall Brown butterfly *Lasiommata megera*. Journal of Insect Conservation, 19(2),

393–402. https://doi.org/10.1007/s10841-014-9748-z

Krämer, B., Kämpf, I., Enderle, J., Poniatowski, D. & Fartmann, T. (2012). Microhabitat selection

in a grassland butterfly: A trade-off between microclimate and food availability. Journal of Insect

Conservation, 16(6), 857–865. https://doi.org/10.1007/s10841-012-9473-4

Kudrna, O., Harpke, A., Lux, K., Pennerstorfer, J., Schweiger, O., Settele, J. & Wiemers, M.

(2011). Distribution atlas of butterflies in Europe. Halle, Germany: Gesellschaft für

Schmetterlingsschutz.

Kurze, S., Heinken, T. & Fartmann, T. (2018) Nitrogen enrichment in host plants increases the

mortality of common Lepidoptera species. Oecologia, 188(4), 1227–1237.

https://doi.org/10.1007/s00442-018-4266-4

Langdon, W.B.V. (2024) How nitrogen availability constrains insect herbivores. PhD thesis,

University of Oxford. https://ora.ox.ac.uk/objects/uuid:ad62c3fd-0e51-40bd-9f65-fe3787e55b71

León-Cortés, J.L., Cowley, M.J.R. & Thomas, C.D. (2000) The distribution and decline of a

widespread butterfly Lycaena phlaeas in a pastoral landscape. Ecological Entomology, 25(3),

285–294. https://doi.org/10.1046/j.1365-2311.2000.00271.x

Letourneau, D.K. & Fox, L.R. (1989) Effects of experimental design and nitrogen on cabbage

butterfly oviposition. Oecologia, 80(2), 211–214. https://doi.org/10.1007/BF00380153

Lüdecke, D. (2018). ggeffects: Tidy Data Frames of Marginal Effects from Regression Models.

Journal of Open Source Software, 3 (26), 772. doi: https://doi.org/10.21105/joss.00772

Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P. & Makowski, D. (2021) performance: An R

package for assessment, comparison and testing of statistical models. Journal of Open Source

Software, 6(60), 3139. https://doi.org/10.21105/joss.03139

Mattson, W.J. (1980) Herbivory in relation to plant nitrogen content. Annual Review of Ecology

and Systematics, 11, 119–161. http://www.jstor.org/stable/2096905

Menzel, A., Yuan, Y., Matiu, M., Sparks, T., Scheifinger, H., Gehrig, R. & Estrella, N. (2020)

Climate change fingerprints in recent European plant phenology. Global Change Biology, 26(4),

2599–2612. https://onlinelibrary.wiley.com/doi/10.1111/gcb.15000

Nijssen, M.E., WallisDeVries, M.F. & Siepel, H. (2017) Pathways for the effects of increased

nitrogen deposition on fauna. Biological Conservation, 212, 423–431.

https://doi.org/10.1016/j.biocon.2017.02.022

O'Connor, R.S., Hails, R.S. & Thomas, J.A. (2014) Accounting for habitat when considering

climate: Has the niche of the Adonis blue butterfly changed in the UK? Oecologia, 174(4), 1463–

1472. https://doi.org/10.1007/s00442-013-2850-1

Oliver, T.H., Hill, J.K., Thomas, C.D., Brereton, T. & Roy, D.B. (2009) Changes in habitat

specificity of species at their climatic range boundaries. Ecology Letters, 12(10), 1091–1102.

https://doi.org/10.1111/j.1461-0248.2009.01367.x

Persson, J., Fink, P., Goto, A., Hood, J. M., Jonas, J. & Kato, S. (2010) To be or not to be what

you eat: Regulation of stoichiometric homeostasis among autotrophs and heterotrophs. Oikos,

119(5), 741–751. https://doi.org/10.1111/j.1600-0706.2009.18545.x

Prudic, K.L., Oliver, J.C. & Bowers, M.D. (2005) Soil nutrient effects on oviposition preference,

larval performance, and chemical defense of a specialist insect herbivore. Oecologia, 143(4),

578–587. https://doi.org/10.1007/s00442-005-0008-5

R Core Team (2024) R: A language and environment for statistical computing. R Foundation for

Statistical Computing.

Rashid, S., Wessely, J., Moser, D., Rumpf, S.B., Kühn, I., Fiedler, K., Hülber, K. & Dullinger, S.,

(2023) Threatened European butterflies concentrate in areas of strong climatic change and

atmospheric deposition pressure. Biological Conservation, 288, 110352.

https://doi.org/10.1016/j.biocon.2023.110352

Raubenheimer, D., Lee, K.P. & Simpson, S.J. (2005) Does Bertrand's rule apply to

macronutrients? Proceedings of the Royal Society B: Biological Sciences, 272(1579), 2429–

2434. https://doi.org/10.1098/rspb.2005.3271

Raubenheimer, D. & Simpson, S.J. (1993) The geometry of compensatory feeding in the locust.

Animal Behaviour, 45(5), 953–964. https://doi.org/10.1006/anbe.1993.1114

Raubenheimer, D. & Simpson, S.J. (2004) Organismal Stoichiometry: Quantifying Non-

Independence Among Food Components. Ecology, 85(5), 1203–1216.

Raubenheimer, D., Simpson, S.J. & Mayntz, D. (2009) Nutrition, ecology and nutritional ecology:

Toward an integrated framework. Functional Ecology, 23(1), 4–16. https://doi.org/10.1111/j.1365-

2435.2009.01522.x

Roth, T., Kohli, L., Rihm, B., Meier, R. & Amrhein, V. (2021) Negative effects of nitrogen

deposition on Swiss butterflies. Conservation Biology, 35(6), 1766–1776.

https://doi.org/10.1111/cobi.13744

Roy, D.B. & Thomas, J.A. (2003) Seasonal variation in the niche, habitat availability and

population fluctuations of a bivoltine thermophilous insect near its range margin. Oecologia,

134, 439–444. https://doi.org/10.1007/s00442-002-1121-3

Ruess, R. W. (1984). Nutrient movement and grazing: Experimental effects of clipping and

nitrogen source on nutrient uptake in *Kyllinga nervosa*. Oikos, 43(2), 183–188.

https://doi.org/10.2307/3544767

Sardans, J., Rivas-Ubach, A. & Peñuelas, J. (2012). The elemental stoichiometry of aquatic and

terrestrial ecosystems and its relationships with organismic lifestyle and ecosystem structure

and function: A review and perspectives. Biogeochemistry, 111, 1–39.

https://doi.org/10.1007/s10533-011-9640-9

Scherer, G., Streanga, B. & Fartmann, T. (2025) Rare butterfly species vitally depend on soil

disturbance by an ecosystem engineer in abandoned calcareous grasslands. Global Ecology

and Conservation, 58, e03451. https://doi.org/10.1016/j.gecco.2025.e03451

Seifan, M., Tielbörger, K., Schloz-Murer, D. & Seifan, T. (2010) Contribution of molehill

disturbances to grassland community composition along a productivity gradient. Acta

Oecologica, 36(6), 569–577. https://doi.org/10.1016/j.actao.2010.08.005

Simpson, S.J., Sibly, R.M., Lee, K.P., Behmer, S.T. & Raubenheimer, D. (2004) Optimal foraging

when regulating intake of multiple nutrients. Animal Behaviour, 68(6), 1299–1311.

https://doi.org/10.1016/j.anbehav.2004.03.003

Sterner, R.W. & Elser, J.J. (2002) Ecological stoichiometry: The biology of elements from

molecules to the biosphere. Princeton, USA: Princeton University Press.

Stewart, K.E.J., Bourn, N.A.D. & Thomas, J.A. (2001). An evaluation of three quick methods

commonly used to assess sward height in ecology. Journal of Applied Ecology, 38(5), 1148–

1154. https://doi.org/10.1046/j.1365-2664.2001.00658.x

Streitberger, M. & Fartmann, T. (2013) Molehills as important larval habitats for the grizzled

skipper, *Pyrgus malvae* (Lepidoptera: Hesperiidae), in calcareous grasslands. European Journal

of Entomology, 110(4), 643–648. https://doi.org/10.14411/eje.2013.087

Streitberger, M., Rose, S., Hermann, G. & Fartmann, T. (2014) The role of a mound-building

ecosystem engineer for a grassland butterfly. Journal of Insect Conservation, 18(4), 745–751.

https://doi.org/10.1007/s10841-014-9670-4

Suggitt, A.J., Wilson, R.J., Isaac, N.J., Beale, C.M., Auffret, A.G., August, T., Bennie, J.J., Crick,

H.Q., Duffield, S., Fox, R. & Hopkins, J.J. (2018) Extinction risk from climate change is reduced

by microclimatic buffering. Nature Climate Change, 8(8), 713–717.

https://doi.org/10.1038/s41558-018-0231-9

Thomas, C.D., Bodsworth, E.J., Wilson, R.J., Simmons, A.D., Davies, Z.G., Musche, M. &

Conradt, L. (2001) Ecological and evolutionary processes at expanding range margins. Nature,

411(6837), 577–581. https://doi.org/10.1038/35079066

Thomas, J.A. (1983) The ecology and conservation of *Lysandra bellargus* (Lepidoptera:

Lycaenidae) in Britain. Journal of Applied Ecology, 20(1), 59–83. https://doi.org/10.4103/1742-

6413.53887

Thomas, J.A. & Lewington, R. (2014) The butterflies of Britain and Ireland (3rd Edition).

Gillingham, UK: British Wildlife Publishing.

Thomas, J.A., Bourn, N.A.D., Clarke, R.T., Stewart, K.E., Simcox, D.J., Pearman, G.S., Curtis,

R. & Goodger, B. (2001) The quality and isolation of habitat patches both determine where

butterflies persist in fragmented landscapes. Proceedings of the Royal Society B: Biological

Sciences, 268(1478), 1791–1796. https://doi.org/10.1098/rspb.2001.1693

Thomas, J.A., Rose, R.J., Clarke, R.T., Thomas, C.D. & Webb, N.R. (1999) Intraspecific

variation in habitat availability among ectothermic animals near their climatic limits and their

centres of range. Functional Ecology, 13(SUPPL. 1), 55–64. https://doi.org/10.1046/j.1365-

2435.1999.00008.x

Thomas, J.A., Simcox, D.J. & Hovestadt, T. (2011) Evidence based conservation of butterflies.

Journal of Insect Conservation, 15(1), 241–258. https://doi.org/10.1007/s10841-010-9341-z

Thomas, J.A., Thomas, C.D., Simcox, D.J. & Clarke, R.T. (1986) Ecology and declining status of

the Silver-Spotted Skipper Butterfly (*Hesperia comma*) in Britain. Journal of Applied Ecology,

23(2), 365–380. https://www.jstor.org/stable/2404023

Throop, H.L. & Lerdau, M.T. (2004) Effects of nitrogen deposition on insect herbivory:

Implications for community and ecosystem processes. Ecosystems, 7(2), 109–133.

https://doi.org/10.1007/s10021-003-0225-x

Van Klink, R., Bowler, D.E., Gongalsky, K.B., Swengel, A.B., Gentile, A. & Chase, J.M. (2020)

Meta-analysis reveals declines in terrestrial but increases in freshwater insect abundances.

Science, 368(6489), 417–420. https://doi.org/10.1126/science.aax9931

van Wingerden, W.K.R.E., Musters, J.C.M. & Maaskamp, F.I.M. (1991) The influence of

temperature on the duration of egg development in West European grasshoppers (Orthoptera:

Acrididae). Oecologia, 87(3), 417–423. https://doi.org/10.1007/BF00634600

Vogels, J.J., Van de Waal, D.B., WallisDeVries, M.F., Van den Burg, A.B., Nijssen, M., Bobbink,

R., Berg, M.P., Olde Venterink, H. & Siepel, H. (2023) Towards a mechanistic understanding of

the impacts of nitrogen deposition on producer–consumer interactions. Biological Reviews, 31.

https://doi.org/10.1111/brv.12972

WallisDeVries, M.F. (2006). Larval habitat quality and its significance for the conservation of

Melitaea cinxia in northwestern Europe. Larvalökologie von Tagfaltern Und Widderchen in

Mitteleuropa. Abhandlungen Aus Dem Westfälischen Museum Für Naturkunde., 68(3–4), 281–

294.

WallisDeVries, M.F. & van Swaay, C.A.M. (2006) Global warming and excess nitrogen may

induce butterfly decline by microclimatic cooling. Global Change Biology, 12(9), 1620–1626.

https://doi.org/10.1111/j.1365-2486.2006.01202.x

Warren, M.S., Maes, D., van Swaay, C.A., Goffart, P., Van Dyck, H., Bourn, N.A., Wynhoff, I.,

Hoare, D. & Ellis, S. (2021) The decline of butterflies in Europe: Problems, significance, and

possible solutions. Proceedings of the National Academy of Sciences, 118(2), p.e2002551117.

https://doi.org/10.1073/pnas.2002551117

White, T.C.R. (1993) The inadequate environment : Nitrogen and the abundance of animals.

Heidelberg, Germany: Springer Berlin.

Wickham, H. (2016) ggplot2: Elegant graphics for data analysis. Springer-Verlag.

https://ggplot2.tidyverse.org

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G.,

Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms,

J., Robinson, D., Seidel, D., Spinu, V. et al. (2019) Welcome to the Tidyverse. Journal of Open

Source Software, 4(43), 1686. https://doi.org/10.21105/joss.01686

Wickham, H., Hester, J. & Bryan, J. (2024) readr: Read rectangular text data. R Package

Version 2.1.5. https://cran.r-project.org/package=readr

Wilby, A., Grubb, L.A., Burrows, J. & Menéndez, R. (2024) Contrasting responses to

microhabitat and temperature determine breeding habitat differentiation between two *Viola*-

feeding butterflies. Ecological Entomology, 49(3), 407–417. https://doi.org/10.1111/een.13312

Wilke, C. (2024) cowplot: Streamlined plot theme and plot annotations for "ggplot2." R Package

Version 1.1.3. https://cran.r-project.org/package=cowplot

Wilson, R.J. & Fox, R. (2021) Insect responses to global change offer signposts for biodiversity

and conservation. Ecological Entomology 46, 699-717. https://doi.org/10.1111/een.12970

Yang, Y., Xi, X., Zhong, X., Eisenhauer, N. & Sun, S. (2017) N addition suppresses the

performance of grassland caterpillars (*Gynaephora alpherakjj*) by decreasing ground

temperature. Ecosphere, 8(3), e01755. https://doi.org/10.1002/ecs2.1755