Elevated *Plasmodium falciparum* sexual conversion in HbAC and HbAS red blood cells without altered IgG/IgM levels to trophozoite and stage I gametocyte-specific antigens

Short title: Elevated *P. falciparum* sexual conversion in HbAC and HbAS RBCs

Yasmina Drissi-El Boukili[1,3], Vera Kühne[1], H. Magloire Natama[4], Pieter Moris[1], Elisabet Tintó-Font[5], Odin Goovaerts[2], Pieter Guetens[1], Ana Moreno-Murillo[1], Yarno Valgaerts[1], T. Edwig Traoré[4], D. Florence Ouedraogo[4], Halidou Tinto[4], Ann Van de Velde[5], Alfred Cortés[6,7], Guy Caljon[3], Anna Rosanas-Urgell[1*]

Department of Biomedical Sciences, Institute of Tropical Medicine, Antwerp, Belgium

Department of Clinical Sciences, Institute of Tropical Medicine, Antwerp, Belgium.

Laboratory of Microbiology, Parasitology and Hygiene (LMPH), Infla-Med Centre of Excellence, University of Antwerp, Belgium.

Unité de Recherche Clinique de Nanoro, Institut de Recherche en Sciences de la Santé, Nanoro, Burkina Faso.

Department of Hematology and Hemostasis, Antwerp University Hospital, Edegem, Belgium.

ISGlobal, Hospital Clinic - Universitat de Barcelona, Barcelona, Catalonia, Spain.

ICREA, Barcelona, Catalonia, Spain.

arosanas@itg.be (*) Corresponding author

AUTHOR CONTRIBUTIONS

Yasmina Drissi-El Boukili: conceptualization, methodology, investigation and laboratory procedures, analysis, writing- original draft, writing- review and editing; Vera Kühne: methodology, analysis, writing- review and editing, substantial revisions and conceptual input; H. Magloire Natama: supervision, field work, writing- review and editing; Pieter Moris: formal

analysis (statistical analysis, model fitting using GLMs and GLMMs); Elisabet Tintó-Font:

methodology (SCAs); Odin Goovaerts: methodology (flow cytometry); Pieter Guetens:

laboratory procedures and analysis (HBBseq and qPCR); Ana Moreno-Murillo and Erin Sauve:

laboratory procedures and analysis (HBBseq); Yarno Valgaerts: laboratory procedures (IgG

and IgM detection by flow cytometry); T. Edwig Traoré, D. Florence Ouedraogo, Halidou

Tinto: field work in Nanoro, Burkina Faso; Ann Van de Velde: clinical evaluation of patients

and coordination of blood sample collection at UZA; Alfred Cortés: supervision, methodology,

writing- review and editing; Guy Caljon, supervision, methodology, writing- review and

editing; Anna Rosanas-Urgell, conceptualization, resources, supervision, funding acquisition,

project administration, methodology, writing- original draft, writing- review and editing,

substantial revisions and conceptual input. All authors read and approved the final manuscript.

ABSTRACT

Malaria transmission relies on the differentiation of *Plasmodium falciparum* asexual parasites

into gametocytes, a process determined by sexual conversion. Mutations in the host

hemoglobin beta (HBB) gene, are known to influence parasite growth, and disease outcome,

yet their impact on sexual conversion remains unclear. We investigated the effect of HBB

mutant genotypes on sexual conversion and humoral immunity in individuals from Nanoro,

Burkina Faso. Sexual conversion rates were assessed with a newly developed *ex vivo* sexual

conversion assay (*ev*SCA) in HBB genotypes HbAA, HbAC, and HbAS, and an *in vitro* SCA

(*iv*SCA) using the NF54-*gexp02-Tom* sexual conversion reporter line cultured in red blood

cells (RBCs) with HbAA, HbAC, HbAS, HbSS, and HbSC genotypes. IgG and IgM responses

against trophozoite- and stage I gametocyte-infected RBC surface antigens were quantified by

flow cytometry in plasma from individuals with HbAA, HbAC, HbAS, HbCC, and HbSC

genotypes. Growth in HBB mutant RBCs significantly increased sexual conversion rates in

both *ex vivo* and *in vitro* assays, while humoral responses did not differ between HBB mutant

and wild-type individuals. These results demonstrate that HBB mutations enhance sexual

conversion independently of IgG- and IgM-mediated immunity, highlighting a host-genetic

factor that may influence malaria transmission potential.

AUTHOR SUMMARY

Transmission of *Plasmodium falciparum* – a malaria causing parasite- depends on the

parasite's ability to form gametocytes, the stages that infect mosquitoes. In many African

regions where malaria is common, hemoglobin variants such as hemoglobin S (the sickle-cell

trait) and C are highly prevalent because they protect against severe disease. However, it

remains unclear whether these variants also influence the parasite's early shift (*i.e.,* sexual

conversion) toward gametocyte development. In our study, we combined *ex vivo* and *in vitro*

sexual conversion assays to measure sexual conversion in red blood cells with different

hemoglobin genotypes. We found that sexual conversion was higher in parasites growing in

red blood cells carrying hemoglobin AC and AS genotypes compared to normal AA

hemoglobin. These findings provide experimental support for earlier epidemiological

observations reporting that individuals with these hemoglobin variants often carry more

gametocytes and are more infectious to mosquitoes. We also measured antibody levels in

individuals with these red blood cells and found no differences in immunity that could explain

the variation in sexual conversion. Our results show that host genetic background can influence

gametocyte development and may shape the infectious reservoir in high-burden settings. This

knowledge is important for designing effective strategies for malaria elimination.

INTRODUCTION

Plasmodium falciparum (*P. falciparum*) causes over half a million deaths annually, mostly in

young children from low- and middle-income countries. Despite the substantial progress in

reducing the global malaria burden during the last decade, this decline has stalled, with some

countries reporting a resurgence in clinical cases (1). Innovative approaches for malaria control

and elimination are thus urgently needed.

Plasmodium infections have exerted strong evolutionary pressure on the human genome, with

hemoglobin mutations providing a well-documented example (2-5). Mutations in the human

HBB gene, such as hemoglobin S (HbS) and hemoglobin C (HbC), result from a substitution

of glutamic acid at position 6 with valine in HbS and lysine in HbC. Individuals can carry

homozygous (HbSS or HbCC), heterozygous (HbAS or HbAC) or compound heterozygous

(HbSC) mutations (6-8). HBB mutations lead to significant changes in red blood cell (RBC)

morphology and function, influencing malaria susceptibility and clinical outcomes (5, 9, 10).

HbS mutations induce hemoglobin polymerization under low oxygen, causing red blood cells

(RBCs) to reshape into sickle forms. HbSS causes ~60-80% RBC sickling (2) with severe

complications including vaso-occlusion, hemolysis, and chronic anemia (11, 12) and high

early-life mortality with estimates ranging from 50-90% among children in low-and middle

income countries (13). On the other hand, HbAS carriers are typically asymptomatic (2), with

fewer sickled RBCs (~40%)(12, 14). HbC mutations cause hemoglobin crystallization without

sickling but increase blood viscosity and shorten RBC life span (15). HbCC and HbAC is

usually a mild and asymptomatic disorder, with HbCC occasionally causing mild chronic

hemolytic anemia in some individuals (2, 7, 15). HbSC carriers display moderate disease

severity and a mixed phenotype with ~50% sickled/crystallized and "pita-shaped" RBCs, (2,

11, 16).

While HbSS and HbCC are relatively rare (~2%), likely due to the high mortality, HbAS and

HbAC can reach frequencies of 10-30% in regions such as Mali and Burkina Faso, which has

been linked to the protection they confer against *P. falciparum* malaria (4, 5, 8, 17-20). HbAC

offers mild protection against severe malaria, while HbAS and HbCC provide strong protection

against both uncomplicated and severe malaria (5, 15, 21-23). The protective mechanisms of

the different HBB mutations are not fully understood (4, 24). HbS infected-RBCs (iRBCs), especially in HbSS, exhibit elevated heme and HO-1−mediated oxidative stress. This causes polymerization, disrupts the actin cytoskeleton, alters iRBCs morphology, and impairs PfEMP-1 iRBCs surface expression (25-27). In HbAS iRBCs, this reduces cytoadherence and sequestration, promoting splenic clearance and controlled parasite infection (28, 29).

Beyond protection from disease, HbS and HbC RBCs may influence the transmission potential of *P. falciparum*. Population studies have reported conflicting evidence regarding whether HbAS influences gametocyte carriage. No significant association has been reported between HbAS and increased gametocyte carriage, though HbAC and HbCC show associations (30). In contrast, other studies found a link between HbAS and gametocyte carriage (31, 32). Few studies have also reported higher infectivity to mosquitoes in individuals carrying HbAS (30, 33), HbAC (30, 34) and HbCC (30) compared to wild-type HbAA iRBCs.

Gametocyte production is initiated by a process called sexual conversion, in which a small proportion (~1%) of asexual *P. falciparum* parasites commit to sexual development at each replication cycle, a proportion referred to as the sexual conversion rate (SCR), and initiate gametocytogenesis (35-37). Sexual commitment is marked by epigenetic activation of the master regulator *pfap2-g* (38, 39), a transcription factor that irreversibly results in sexual conversion and gametocyte development (38, 39). Sexual rings circulate in blood (40-43), while maturation of gametocyte stages I to IV occurs mainly in the bone marrow over ~10 days (44). Mature male and female stage V gametocytes re-enter the peripheral blood (*i.e.,* gametocyte carriage), and are the only parasite stages capable of infecting *Anopheles* mosquitoes. Gametocyte carriage determines human to mosquito transmission and correlates with infectivity to mosquitoes (45-50).

Elevated gametocyte carriage reported in individuals with HbAC and HbAS genotypes could

result from elevated sexual conversion but also prolonged infections, leading to accumulation

of gametocytes in peripheral blood over numerous replication cycles. Several host related

factors have been shown to increase sexual conversion (51), including antimalarial treatment

(52-54), lysophosphatidylcholine depletion (55-57), and reticulocyte-rich environments (58,

59). A terminal investment in transmission through increased sexual conversion (60-63), could

similarly be initiated by the impaired environment in HbS and HbC iRBCs (4, 29, 64-69). To

date, only one study has investigated the link between sexual conversion and HBB genetics,

showing that increased intracellular heme in HbS iRBCs promotes sexual conversion (25).

Another potential driver of elevated sexual conversion in HBB mutant RBCs is the host

immunity against *P. falciparum*. HbS and HbC mutations appear to modulate host immunity

against *P. falciparum* by increasing regulatory cytokines (*e.g.,* IL-10) (68)), upregulating HO-

1 (66), and dampening pro-inflammatory responses (*e.g.,* TNF-α, IFN-γ) (24), which both

protect against severe malaria and enhance parasite clearance. Moreover, HBB mutant RBCs

show elevated antibody levels against specific RBC antigens (24). including increased IgG

against asexual-stage antigens (*e.g.,* AMA1, EBA175, MSP1-3, CSP, and PSE) and RBC

surface antigens such as PfEMP-1 in HbAS individuals across multiple populations (70-73).,

Discrepancies in gametocyte carriage studies highlight the need to investigate upstream

processes such as sexual conversion. Demonstrating that HBB mutant RBCs (*e.g.,* HbAS and

HbAC) enhance *P. falciparum* sexual conversion could improve strategies to interrupt

transmission (74) and identify carriers as key contributors to the infectious reservoir for

targeted transmission-blocking interventions.

In this study, we investigated the impact of host HBB mutant genotypes (HbAA, HbAC and

HbAS, HbSS, HbCC and HbSC) on sexual conversion in *ex vivo* and *in vitro P. falciparum*

cultures (52, 75) and assessed levels of IgG and IgM against trophozoite- and stage I

gametocyte -infected RBCs (Troph-Gam-iRBCs) by flow cytometry.

RESULTS

HbAC and HbAS genotype prevalences in Nanoro, Burkina Faso

Using Sanger Sequencing, we genotyped the HBB gene in participants of a longitudinal cohort

study (*i.e., InHost* study) in Nanoro, Burkina Faso (*n* = 864) and determined the population

frequencies of HbAA, HbAC, HbCC, HbAS, HbSS, and HbSC genotypes (Fig 1). HbAA (wild-

type) was the most frequent (71.5%), followed by HbAC (20%), HbAS (7.7%), HbSC (0.8%),

and HbCC (0.2%); no HbSS individuals were identified.

Fig 1. Genotype frequencies of the HBB gene in the study population. Pie chart depicting the sample sizes (n)
and frequencies (%) of the different HBB genotypes in the study population.

Novel *ex vivo* sexual conversion assays reveal elevated sexual conversion rates in HbAS and HbAC RBCs

162 To investigate the effect of HBB genotypes on *P. falciparum* sexual conversion we measured

163 next cycle SCRs of in host induced *P. falciparum* isolates collected from individuals with

164 different HBB genotypes (HbAA, *n* =14; HbAC, *n* =12; HbAS, *n* = 6) from the Nanoro Health

165 District, Burkina Faso, using an in-house *ex vivo* sexual conversion assay (*ev*SCA), optimized

166 and validated as described in *Supporting information file subsection "Ex vivo sexual*

conversion assays (evSCAs)". SCRs were quantified by immunofluorescent staining (IFAs) detecting the number of parasites stained positive for the Pfs16 protein (stage I gametocytes) (76-78) out of the number of parasites stained positive for DNA/parasite nuclei (*i.e.,* the total parasite population consisting of stage I gametocytes plus schizont stages), expressed as a percentage. The experimental set-up of the *ev*SCAs and representative immunofluorescent assay (IFA) images are shown in Figs 2A and 2F, respectively.

The SCR (median[IQR]) was 2.65% [1.72] in *P. falciparum* isolates from HbAA genotype, 5.05% [2.55] for HbAC and 18.40% [19.70] for HbAS genotypes. SCR increased significantly in HbAC (Beta-binomial GLMM, OR[95%CI] = 1.82 [1.17-2.82], $p = 0.0032**$) and HbAS (Beta-binomial GLMM, OR [95%CI] = 8.16 [4.34-15.37], $p < 0.0001***$) genotypes compared to HbAA (Fig 2B and Table S1). Fold-change in SCRs (median [IQR]) of *P. falciparum* isolates in HbAC and HbAS compared to the mean of wild-type HbAA RBCs were 2.04 [0.85] in HbAC and 7.07 [7.32] in HbAS, with values in HbAS RBCs significantly higher than in HbAC RBCs (Kruskal-Wallis test, overall $p = 0.034*$; Fig 2C). Parasitemias did not significantly differ between mutant HBB RBCs (HbAC and HbAS) and wild-type HbAA RBCs (Kruskal-Wallis test, $p > 0.05$, Fig 2D); and SCRs and initial parasitemia did not correlate (Spearman correlation, $\rho = 0.05$, $p > 0.05$; (Fig S1A). These results suggest that differences in initial parasitemias do not have an impact on the observed elevated SCRs.

The proportion of RBCs infected with stage I gametocytes (extrapolated from proportion of iRBCs in Giemsa-stained slides after MACS enrichment) was highest in cultures with HbAS RBCs, followed by HbAC RBCs, and lowest in wild-type HbAA RBCs (Fig 2E) (same trend observed as in SCRs by HBB genotype, Fig 2B). This confirms that the higher SCRs in mutant HBB genotypes reflect a net increase in gametocyte production rather than a reduction in asexual parasite numbers.

Fig 2. Effect of human hemoglobin beta (HBB) HbAA, HbAC and HbAS genotypes on sexual conversion determined with *ex vivo* sexual conversion assays (*ev*SCAs). (A) Schematic representation of the *ev*SCAs. Parasite isolates from RBCs with different HBB genotypes were cultured *ex vivo* under basal non-inducing (+Choline) conditions. ML10 was added at 11 hpi to block reinvasion. Stage I gametocytes and schizonts were enriched by MACS, and sexual conversion rates were measured by IFA using Pfs16 (sexual stages) and DAPI (total parasites) staining. **(B)** Sexual conversion rates (SCRs) by HBB genotype determined with IFA. Sample size *n* per group: HbAA *n* = 14, HbAC *n* = 12 and HbAS *n* = 6; total *n* = 32. Data presented as the median and IQR of three independent technical IFA replicates. **(C)** Fold-change SCRs in HbAC and HbAS genotypes relative to HbAA. **(D)** Initial parasitemia at 0 hpc (parasites per µl; log10 scale) by HBB genotype determined with *var*ATS qPCR. **(E)** Proportion of asexual and gametocyte stages, after MACS enrichment, within the initial parasitemia by HBB genotype. **(F)** Representative IFA images of parasites in HbAA, HbAC and HbAS RBCs. Top row with DAPI (all parasites; blue arrows: schizonts) and AF488 (white arrows: stage I gametocytes) overlay and bottom row including brightfield (BF) channel to visualize the parasite's hemozoin. Objective: 40×; Scale bar: 5µm. Images were captured with the Revolve (ECHO) camera. **Statistical tests: B,** Beta-binomial generalized linear mixed models (GLMM) with Tukey adjustment for pairwise comparisons; **D,** Kruskal-Wallis test (p-values: bold solid line), followed by the Wilcoxon Rank Sum post-hoc test with Holm correction (*p*-values: normal solid line) when significant. Significant p-values: $p < 0.05$*, $p < 0.01$**, $p < 0.001$***; and non-significant p-values as: ns. **Abbreviations:** Hpc, hours post culture; ML10, cGMP-dependent protein kinase inhibitor; hpi, hours post-invasion; AF, Alexa Fluor; DAPI, DNA stain 4',6-diamidino-2-phenylindole; Pfs16 *antibody: Kindly provided by R. Sauerwein and T. Bousema, Radboud University*; *var*ATS qPCR, *var* gene acidic terminal sequence (*var*ATS) quantitative PCR (qPCR).

In vitro sexual conversion assays with the *P. falciparum* transgenic reporter line NF54-*gexp02-Tom*

To validate findings from the *ev*SCAs, we established an *in vitro* sexual conversion assay (*iv*SCA) based on an assay that was previously published (52) using a *P. falciparum* transgenic reporter line NF54-*gexp02-Tom* (52, 75), which carries a genome-integrated tandem Tomato (tdTom) fluorescent marker under the control of the promoter of the early gametocyte marker gene *gexp02*. In our assay, parasites, were allowed to invade and grow in cryopreserved non-infected RBCs with different HBB genotypes from CRUN and UZA (28 endemic samples: HbAA $n = 5$, HbAC $n = 5$, HbAS $n = 4$, HbSS $n = 12$, HbSC $n = 2$; and six non-endemic HbAA controls, HbAA CTRL; (Fig 3A, Tables 1 and 3).We measured the SCRs by flow cytometry (FC) detecting sexual stages by the expression of the red fluorescent protein TdTom under the promotor of the *gexp02* gene, which is exclusively expressed in sexual stages (75), and the total number of infected RBCs (iRBCs) using the DNA stain Hoechst33342. The *iv*SCA experimental set-up, validated as described in *Supporting information file subsection "In vitro sexual conversion assays (ivSCAs)"*, is illustrated in Fig 3A and representative flow cytometry plots by HBB genotype are shown in Fig S2. Parasite cultures were tightly synchronized by Percoll purification at the schizont stage (~40-44 hpi), followed by the addition of RBCs with different HBB genotypes to allow reinvasion. Subsequently, sorbitol lysis was performed at the ring stage to obtain a narrow developmental window of ~0-5 hpi (Fig 3A). During the assay, choline was either maintained to repress or depleted (-Choline) to induce sexual conversion, following established protocols (55, 75, 79). Choline depletion was performed at ring stage (~0-5 hpi) (Fig 3A). SCRs were determined as the proportion of parasites that developed into gametocytes in the cycle following exposure to RBCs with different HBB genotypes (Generation 1, Fig 3A). Since sexual parasites begin expressing the TdTom reporter at ~10-15 hpi in Generation 1, and SCRs are typically measured at ~25-30 hpi (Generation 1, Fig 3A)

240 (75), and given the observed developmental delays in HbAC, HbAS, HbSS, and HbSC RBCs

241 compared to wild-type HbAA RBCs (Fig 3), we included an additional later timepoint (~35-

242 40 hpi) within the same cycle to ensure robust TdTom detection across all HBB genotypes.

243 **Table 1: Overview of the different experiments with the number of cryopreserved RBC samples by HBB**
244 **genotype.**

Sample by HBB genotype	Experiments				
	1	2	3	4	Total n
HbAA (CRUN)	-	1	3	1	5
HbAC (CRUN)	1	2	1	1	5
HbAS (CRUN)	2	1	1	-	4
HbSS (UZA)	-	2	2	8	12
HbSC (CRUN/UZA)	-	2	-	-	2
Total n° samples	3	8	7	10	28

At ~35-40 hpi (Generation 1) SCRs (median[IQR]) in wild-type HbAA RBC cultures were at

10.8%[0.49] at the threshold of what is considered induced (>10%). SCR was slightly but non-

significantly increased to 13.30%[4.71] for HbAC (Beta-binomial GLM, HbAC vs. HbAA

RBCs: OR[95% CI] = 1.25[0.81−1.92] , $p > 0.05$) and significantly increased to 16.5%[5.64]

for HbAS RBC cultures (HbAS vs. HbAA RBCs: OR[95% CI] = 1.73 [1.12−2.66], $p =$

0.0056**; Table S1). In HbSS the SCRs were significantly lower than in wild-type control

HbAA RBCs (6.96% [3.17], HbSS vs. HbAA (Beta-binomial GLM: OR[95% CI] = 0.64

[0.43−0.95], $p = 0.021$*; Table S1), while in HbSC RBC cultures (SCR of 7.99% [1.83]), there

was no statistically significant difference to HbAA (Beta-binomial GLM, $p > 0.05$, Table S1

and Fig 3B). Similarly, the fold-change in SCRs (median [IQR]) relative to wild-type fresh

HbAA non-endemic RBC cultures showed a trend consistent with the observations above, with

elevated SCRs in HbAC (1.47 [0.66]) and HbAS (2.05 [0.38]) RBCs, while remaining

unchanged (~1) in HbAA (1.23 [0.31]), HbSS (0.84 [0.49]), and HbSC (1.10 [0.25]) RBCs.

However, fold-change differences between HbAA and HBB mutant RBCs were not

statistically significant (Wilcoxon Rank Sum post-hoc test, $p > 0.05$; Fig 3C).

Conversely, at ~25-30 hpi (Generation 1), no differences were found in SCRs between HbAA

and HBB mutant RBCs (Kruskal-Wallis test, $p > 0.05$) (Fig S3A). Compared to ~35-40 hpi

(Generation 1), these results suggest that at ~25-30 hpi (Generation 1) a proportion of parasites

may not yet have reached the stage of TdTom expression. This is supported by the observation

that the SCR (median [IQR]) in HbAC and HbAS increased from 9.41%[5.14] and 15.0%[6.48]

at ~25-30 hpi to 13.3%[4.71] and 16.5%[5.64] at ~35-40 hpi, respectively, within Generation

1 (Fig 3A), indicating that parasites in these RBCs were indeed developmentally delayed.

In cultures in which sexual conversion had been stimulated by choline depletion (-Choline) at

both time points of Generation 1 (Fig 3A), no significant differences in SCRs were observed

between HbAA and HbAC or HbAS RBCs (Table S1 and Figs S3B and S3C). These findings

confirm that maximal SCR induction following choline depletion does not vary across mutant

HBB RBCs compared to HbAA RBCs. The significantly lower SCRs in HbSS and HbSC RBCs

may be explained by the partial arrest and extreme delay in development observed in these

cultures (Fig 3 and Table S2).

Fig 3. Effect of human hemoglobin beta (HBB) HbAA, HbAC and HbAS genotypes on sexual conversion determined with *in vitro* sexual conversion assays (*iv*SCAs). (A) Experimental design of *iv*SCAs using the NF54-*gexp02-Tom P. falciparum* line. Tightly synchronized late-stage schizonts (Generation -1) were allowed to reinvade RBCs of different HBB genotypes, and ring stages (~0-5 hpi) were then cultured under both non-inducing (+Choline) and inducing (-Choline) conditions. Parasite development was assessed by flow cytometry (FC) at timepoints: ~35-40 hpi (Generation 0) and ~25-30 hpi/~35-40 hpi (Generation 1). Sexual conversion rates (SCRs) were assessed FC at ~25-30 hpi and ~35-40 hpi in Generation 1 **(B)** Sexual conversion rates (SCRs) by HBB genotype determined with FC at ~35-40 hpi (Generation 1), in +Choline cultures. Sample size n per group: endemic RBC samples with HbAA $n = 5$, HbAC $n = 5$ and HbAS $n = 4$, and HbSS $n = 12$ and HbSC $n = 2$, total $n = 28$; and non-endemic fresh RBC samples with "HbAA CTRL" $n = 6$. **(C)** Fold-change SCRs in cryopreserved HbAA, HbAC, HbAS, HbSS and HbSC genotypes relative to fresh HbAA RBCs. **(D)** Proportion of asexual and gametocyte stages and total parasitemia by HBB genotype at ~35-40 hpi (Generation 1), in +Choline cultures. **(F)** Reticulocytes percentage by HBB genotype. **Statistical tests:** (B and E), Beta-binomial generalized linear mixed models [GLM] with Bonferroni adjustment for pairwise comparisons; (C) Kruskal-Wallis test with the Wilcoxon Rank Sum post-hoc test with Holm correction. Significant *p*-values are shown as: $p < 0.05^*$, $p < 0.01^{**}$, $p < 0.001^{***}$; non-significant *p*-values as: ns. Overall *p*-values are presented with a bold straight line. **Abbreviations:** Hpi, hours post-invasion; CTRL, control sample (fresh HbAA RBCs).

Differences in SCRs could not be explained by variations in initial parasitemia at ~0-5 hpi

(Generation 0, all cultures were started with a parasitemia below 2.0%), as no significant

differences were found between HBB genotypes (Beta-binomial GLM, $p > 0.05$; Table S1) and

no correlation was observed between SCR and parasitemia (Spearman $\rho = 0.10$, $p > 0.05$; Fig

S1B). Analysis of the proportion of RBCs infected with asexual parasites and stage I

gametocytes by HBB genotype has shown that total gametocytemia (determined by FC at ~35-

40 hpi, Generation 1) was highest in cultures with HbAS RBCs, followed by HbAC RBCs, and

lowest in wild-type HbAA RBCs, while levels in HbSS and HbSC RBCs were similar to or

lower than those in HbAA RBCs (Fig 3D). Total gametocytemia at ~25-30 hpi (Generation 1)

followed the same dynamics (Fig S3E). This confirms that the higher SCRs in HbAS and

HbAC reflect enhanced gametocyte production rather than diminished asexual parasite

numbers.

Since reticulocytes have been proposed to enhance SCRs (58), and individuals with mutant

HBB genotypes often exhibit anemia-associated reticulocytosis, we assessed reticulocyte

percentages across genotypes. Including all HBB groups (HbAA, HbAC, HbAS, HbSS, and

HbSC), we observed significant differences in reticulocyte percentage with the highest median

percentages in HbSS compared to HbAA RBCs (Beta-binomial GLM, OR [95%CI] = 2.56

[1.20−5.44], $p = 0.008**$, Table S1 and Fig 3E). No correlation between reticulocyte

percentage and SCR was found (Spearman $\rho = 0.10$, $p > 0.05$; Fig S3F), reinforcing the

conclusion that higher SCRs are due to intrinsic properties of mutant RBCs rather than

reticulocyte abundance. Additionally, to control for hemolysis during FC experiments, we

assessed RBC counts (RBCs/µl) across HBB genotypes at various timepoints, as shown in

Table S3. No significant differences were observed at any of the time points, indicating that

mutant HBB RBCs did not undergo hemolysis during the course of the experiment (Kruskal-
Wallis test, overall $p > 0.05$).

Red blood cells with mutant HBB genotypes lead to delayed parasite development

P. falciparum growing in HbAS RBCs exhibits developmental delay/arrest before DNA
replication when growing under low oxygen conditions ($\leq 5\%$ O_2, as present in the bone
marrow and spleen, and in standard *P. falciparum* cultures as used for the *iv*SCA).
Developmental delay/arrest in a particular HBB genotype would affect the time at which sexual
conversion can be measured *in vitro.* In NF54-*gexp02-Tom* parasites, grown in HbAA RBCs,
TdTom signal can be detected from ~10-15 hpi until ~96-101 hpi in the cycle after exposure
(Generation 1), and sexual conversion is usually measured at ~25-30 hpi (Generation 1) when
parasites are large rings/early trophozoites (52, 75). If an HBB mutant genotype causes a delay
in development, TdTom might not yet be expressed at ~25-30 hpi (Generation 1).

To determine whether parasite stage development of NF54-*gexp02-Tom* parasites differs
between RBCs with wild-type HbAA and mutant HBB genotypes (HbAC, HbAS, HbSS,
HbSC), we first quantified the percentage of rings, trophozoites and early/late schizonts in a
set of RBC samples with the different HBB genotypes (HbAA, $n = 1$; HbAC, $n = 1$; HbAS, n
$= 2$; HbSS, $n = 2$; HbSC, $n = 2$) using LM (Fig 4A). Parallel analysis of the proportion of late
stages (schizonts) in the same samples by flow cytometry (FC) (gating strategy in Fig 4C)
revealed consistent patterns between the two approaches. We therefore applied FC to the full
set of samples (HbAA, $n = 5$; HbAC, $n = 5$; HbAS, $n = 4$; HbSS, $n = 12$; HbSC, $n = 2$; and
fresh non-endemic HbAA CTRLs, $n = 6$, Fig 4B), enabling accurate detection of genotype-
specific developmental delays.

We found that parasites in HbAA RBCs progressed normally from Generation 0 to 1 with no
apparent delay. In HbAC RBCs, development was slightly delayed (~3 hours delay vs. HbAA

at Generation 0, ~35-40 hpi) but without arrest, whereas in HbAS RBCs a more pronounced

delay was observed (~10 hours delay) and partial developmental arrest in one sample (Fig 4

and Table S2). HbSS and HbSC RBCs presented a similar delay to HbAS (~ 9 and ~16 hours

delay, respectively) with a persistent population of schizonts, trophozoites, and late rings across

all timepoints (Fig 4A), consistent with the high frequency of partially arrested/delayed cultures

(67% in HbSS and 50% in HbSC) (Table S2). The (partial) arrest/extreme delay in these

cultures potentially makes them unsuitable for SCR measurement. Given the observed

developmental delays in all mutant genotypes, measuring sexual conversion at a single

timepoint (~25-30 hpi, Generation 1) may not be accurate. To capture all *gexp02-Tom-*

expressing sexual parasites (75), including those with delayed development, SCRs were

additionally quantified at ~35-40 hpi (Generation 1) (+ 20 hours post-expression of *gexp02*

should account for the 3-15 h delay measured). Detailed analysis is further described in

Supporting information file in sub-section "Assessment of parasite development in RBCs with

different HBB genotypes".

Fig 4. *P. falciparum* parasite development in red blood cells (RBCs) with different HBB genotypes. (A) Microscopy-based quantification of parasite stages expressed as percentages of a total of 100 counted parasites per sample. Y-axis: percentage of parasite stages (%); X-axis: HBB groups. Sample size n per group: HbAA *n* = 1, HbAC *n* = 1 and HbAS *n* = 2, and HbSS *n* = 2 and HbSC *n* =2; total *n* = 8. Timepoints shown are ~35-40 hpi (Generation 0; left graph), ~25-30 hpi (Generation 1; middle graph) and ~35-40 hpi (Generation 1; right graph). **(B)** Dot plots showing the proportion of late stages (*i.e.,* schizont stages) on the total infected RBCs by flow cytometry (Y-axis) by HBB genotype groups (X-axis). Timepoints shown are ~35-40 hpi (Generation 0), ~25-30 hpi (Generation 1) and ~35-40 hpi (Generation 1). Sample size n per group: endemic RBC samples with HbAA *n* = 5, HbAC *n* = 5 and HbAS *n* = 4, and HbSS *n* = 12 and HbSC *n* = 2, total *n* = 28; and non-endemic fresh RBC samples with "HbAA CTRL" *n* = 6. **(C)** Representative flow cytometry plots showing the gating strategy for the detection of late parasite stages. Left: total RBCs gated in the SSC-A versus FSC-A plot. Right: within this population, uninfected RBCs (uRBCs) and infected RBCs (iRBCs) are identified, with iRBCs further gated into early and late parasite stages. **Statistical tests**:(B, right graph) Beta-binomial generalized linear mixed models [GLM] with Bonferroni adjustment for pairwise comparisons; (B, left and middle graph) Kruskal-Wallis test with the Wilcoxon Rank Sum post-hoc test. Significant *p*-values are shown as: $p < 0.05^{*}$, $p < 0.01^{**}$, $p < 0.001^{***}$; non-significant *p*-values as: ns. Overall *p*-values are presented with a bold solid line, and *p*-values after pairwise comparisons with a normal solid line. **Abbreviations:** Hpi, hours post-invasion; SSC-A, Side Scatter-Area; FSC-A, Forward Scatter-Area; CTRL, control sample (fresh HbAA RBCs).

HBB genotypes are uniquely associated with enhanced SCRs compared to other host factors

To assess whether factors other than the mutant HBB genotypes influenced SCRs, we performed a risk analysis of additional variables (*i.e.,* gender, village, ethnicity, age, parasitemias, etc.) stratified by HBB genotype group, in the study population in the *ev*SCAs (n = 32). This showed that only SCRs differ amongst the HBB genotypes HbAA, HbAC and HbAS (Kruskal-Wallis test, overall $p < 0.001$***; Table S4).

For the same aim, we performed a risk analysis of factors in the study population in the *iv*SCAs (n = 34) that may impact sexual conversion rates (SCRs) in addition to the HBB mutant genotypes (Table 2). Our analysis indicated that HBB genotypes and gender are a risk factor for enhanced SCRs (Fisher's exact test, overall $p < 0.01$** and $p < 0.05$*, respectively). The percentage of reticulocytes was negatively associated with SCRs, with the lowest SCR and highest percentage of reticulocytes in HbSS and HbSC RBCs (Spearman $\rho = -0.50$, $p < 0.01$**). The remaining factors did not show any impact on SCRs (Fisher's exact test and Kruskal-Wallis test, $p > 0.05$).

We also explored which factors are potentially associated with HBB genotype (Table S5). We noted that the percentage of reticulocytes is the highest in the HBB mutant genotypes HbSS and HbSC, followed by HbAS, HbAC and HbAA (Kruskal-Wallis, overall $p < 0.001$***). Additional factors such as age, gender, village and ethnicity showed statistically significant differences in their distribution across HBB genotype categories (Fisher's exact test, overall $p < 0.05$), whereas blood groups did not (Fisher's exact test, $p > 0.05$).

Table 2. Risk Analysis of factors for elevated sexual conversion rates (SCRs) at 35-40 hpi (Generation 1), in the study participants within the *in vitro* sexual conversion assays (*n* = 34). Data from *P. falciparum* non-infected samples collected in the Nanoro Department, Burkina Faso and the UZA (Belgium). Statistical tests performed are shown at the bottom of the figure with correspondent *p*-values (with significant *p*-values as: $p < 0.05^*$, $p < 0.01^{**}$, $p < 0.001^{***}$). Abbreviations: HBB, Hemoglobin beta; SCR, Sexual conversion rates; hpi, hours post-invasion; IQR, Interquartile range.

	SCR 35-40 hpi	$p^\$$		
n (%)	28 (100.0)	NA		
HBB Genotype (median (IQR))		<0.01**		
HbAA	10.8 (0.49)			
HbAC	13.30 (4.71)			
HbAS	16.50 (5.64)			
HbSS	6.96 (3.17)			
HbSC	7.99 (1.83)			
ABO blood group (median (IQR))		0.15		
O+	7.57 (7.86)			
A+	8.80 (2.52)			
B+	17.80 (3.48)			
A-	9.34 (2.87)			
B-	10.90 (3.77)			
Gender (median (IQR))		<0.05*		
Female	11.30 (5.04)			
Male	7.57 (4.09)			
Village (median (IQR))		0.30		
Nanoro	11.30 (3.75)			
Nazoanga	15.40 (0.60)			
Soum	0 (0)			
Séguédin	0 (0)			
Ethnicity (median (IQR))		0.40		
Gourounsi	17.80 (0.00)			
Mossi	12.30 (4.14)			
	Median (IQR)	**rho°**	**p°**	
Reticulocytes (percentage)	1.50 (5.63)	-0.50	<0.01**	
Age	27.00 (30.00)	-0.18	0.37	

Significant p-values ($p < 0.05^*$, $p < 0.01^{**}$, $p < 0.001^{***}$).
$Categorical variables test: Fisher's exact test.
$Continuous variables test: Kruskal Wallis test and Post-Hoc analysis with Wilcoxon Rank Sum Post-hoc Test with Holm correction.
° Continuous variables : Spearman correlation test

Comparison of IgG and IgM levels in plasma from different HBB genotypes living in a malaria endemic region by flow cytometry

We analyzed IgG and IgM levels against *P. falciparum* in individuals with different HBB genotypes using flow cytometry, in selected endemic human plasma samples from the *InHost* longitudinal cohort study (see sub-section on *"Materials and methods, Study design"*). Detailed information on assay optimization and validation is provided in the *Supporting information file subsection IgG and IgM detection assays by flow cytometry)*.

For robust ΔMFI values a cut-off of a minimum number of event counts per category was set to 200 events (Figs S4A and S4C). In each assay, positive controls (three acute *P. falciparum-infected* non-endemic plasma from a traveller at D0, D21 and D23) and a negative control ("naïve non-endemic plasma") were analyzed. The "naïve non-endemic plasma" sample consistently did not reach the 200 event counts threshold, confirming its suitability as a negative control. The negative control values served as a baseline for the levels of α-Troph-Gam-iRBCs IgG & IgM and were lower than those of the endemic plasma samples of all HBB genotypes (Fig 5). Endemic plasma samples were considered positive for IgG or IgM when both the event count and the levels of α-Troph-Gam-iRBCs IgG and IgM exceeded those observed in the "naïve non-endemic plasma". All endemic plasma samples exhibited event counts below the threshold of 200 for the levels of total α-Troph-Gam-iRBCs IgM (Hoechst33342+/PerCP-Cy5.5.+) and α-Troph-iRBCs IgM (Hoechst33342+/ tdTom-/PerCP-Cy5.5.+). The number of α-Gam-iRBCs IgG (Hoechst33342+/tdTom+/AF488+) and IgM (Hoechst33342+/tdTom+/PerCP-Cy5.5.+) was below 200 events for all endemic plasma samples (*n* = 61). Thirteen samples (21%) had no detected (*i.e.,* zero) events for α-Gam-iRBCs IgM. Based on these parameters, the MFI could only be evaluated for total and α-Troph-iRBCs IgG in the statistical analyses (Figs S5B and S5D).

Importantly, the events count in α-Troph-Gam-iRBCs and α-Troph-iRBCs IgM of the positive

controls at D21 and D23 exceeded both the event counts observed in the "naïve non-endemic

plasma" and the threshold of 200 events, confirming that our flow cytometry assay can detect

α-Troph-Gam-iRBCs IgM (Fig S5C). The time course of the "acute *P. falciparum*-infected

non-endemic plasma at D0, D21 and D23" generally followed the expected IgG (D0 < D21 <

D23) and IgM (D0 < D21 > D23) dynamics in α-Troph-Gam-iRBCs IgG & IgM after a *P.*

falciparum infection (80, 81).

We found no significant differences in the levels of α-Troph-Gam-iRBCs IgG & IgM (% of

cells/ΔMFI data are only considered robust for α-Troph-Gam-iRBCs IgG, reaching the

minimum threshold of 200 events; Fig 5) (Beta-binomial GLM, $p > 0.05$; Table S1). When

stratifying the Troph-Gam-iRBCs in asexual and sexual iRBCs, there was no significant

difference in the levels of α-asexual and α-sexual iRBCs IgG & IgM in endemic plasma from

individuals carrying different HBB genotypes in either subgroup (Fig 5) (Beta-binomial GLM,

$p > 0.05$; Table S1). We detect low levels or absence of α-Gam-iRBCs IgG & IgM, and low

levels or absence of IgM against Troph-Gam-iRBCs, Troph-iRBCs and Gam-iRBCs in the

plasma of the endemic individuals, as compared to high levels of IgG against Troph-Gam-

iRBCs and Troph-iRBCs (Figs 5 and S5).

The evaluation of variables in the study population ($n = 61$) such as: *P. falciparum* infection

status at the time of sample collection, parasitemia (parasites per µl), presence of gametocytes

in blood, age, gender and village; stratified by HBB genotype (HbAA [$n = 18$], HbAC [$n = 18$],

HbAS [$n = 18$], HbCC [$n = 2$], and HbSC [$n = 5$]) revealed no significant differences between

the categories (Fisher's exact test and Kruskal-Wallis test; $p > 0.05$) (Table S6). In our study

population, there is a homogeneous distribution of individuals across the categorical variables,

with no observed sampling bias that could affect further analysis.

A) IgG+ Troph-Gam-iRBCs (% cells)

B) IgM+ Troph-Gam-iRBCs (% cells)

Fig 5. IgG and IgM mediated humoral immune response against *P. falciparum* in plasma samples from individuals with different human hemoglobin beta (HBB) genotypes. Data from *P. falciparum* infected ($n = 44$) and non-infected ($n = 17$) plasma samples collected at the Clinical Research Unit (CRUN) in Nanoro, Burkina Faso (total $n = 61$). **(A)** Percentage of α-Troph-Gam-iRBCs IgG: total (left), asexual (middle) and sexual (right). **(B)** Percentage α-Troph-Gam-iRBCs IgM: total (left), asexual (middle) sand sexual (right). Positive controls ("*Pf*-infected non-endemic" *i.e.,* three acute *Pf*-infected non-endemic plasma samples from a traveller at D0, D21, and D23) and a negative control *i.e.,* "Naïve non-endemic": naïve plasma from a non-endemic volunteer with HbAA genotype) are shown as individual dots per sample. The Y-axis range differs between graphs for visualization. No significant differences were found (Beta-binomial generalized linear models [GLM] with Bonferroni adjustment for pairwise comparisons, $p > 0.05$). **Abbreviations:** IgG and IgM, Immunoglobulin G and M; Troph-Gam-iRBCs, trophozoite- and gametocyte-infected red blood cells; *Pf, P. falciparum*. D: Day(s).

Further analysis of the levels of α-Troph-Gam-iRBCs IgG & IgM in the study population by "*P. falciparum* infection status", revealed that the levels of α-Troph-Gam-iRBCs IgG & IgM were higher in the *P. falciparum*-infected individuals ($n = 44$, median[IQR] for IgG: 17.90[6.30], and for IgM: 2.03[3.34]) compared to the non-infected individuals ($n = 17$, median[IQR] for IgG: 9.10[9.66], and for IgM: 1.56[1.27]) (Beta-binomial GLM, OR[95%CI] = 1.81 [1.35−2.43], $p = 0.001$** [for IgG]; and OR[95%CI] = 1.51 [1.01−2.23], $p = 0.043$* [for IgM]) (Fig S7). These results confirm the expected IgM dynamics, where IgM levels are

highest during acute or recent infections (*P. falciparum*-infected individuals) compared to past

infections (non-infected individuals) (80, 81).

Stratification analysis of the levels of α-Troph-Gam-iRBCs IgG & IgM by age-group (*i.e.,*

younger age: "1 to 4 years" compared to older age: "5 to 19 years") in *P. falciparum*-infected

compared to non-infected individuals, showed that both *P. falciparum*-infected and non-

infected individuals with younger age had lower levels of α-Troph-Gam-iRBCs IgG (*Pf*-

infected *n* = 22 and median[IQR]: 17.90[7.23]; non-infected *n* = 15 and median[IQR]:

9.30[10.80]) than those with an older age (*P. falciparum*-infected *n* = 22 and median

[IQR]:18.60[5.64]; non-infected *n* = 2 and median[IQR]: 12.60[3.69]) (Welch's ANOVA, *p* =

0.082 [IgG]; Kruskal-Wallis test *p* = 0.38 [IgM]). *P. falciparum*-infected individuals with

younger age had slightly higher levels of α-Troph-Gam-iRBCs IgM (*n* = 22, median [IQR]:

2.19[2.33]) than those with an older age (*n* = 22, median [IQR]: 1.87[3.04]) (Kruskal-Wallis

test, *p* = 0.94). In contrast, the levels of α-Troph-Gam-iRBCs IgM in non-infected individuals

with younger age were slightly lower (*n* = 15, median[IQR]: 1.56[1.12]) than those with an

older age (*n* = 2, median [IQR]: 2.18[0.00]) (Kruskal-Wallis test, *p* = 0.16).

Together these results (the positive controls exceeding the 200 events threshold for α-Troph-

Gam-iRBCs IgM and IgG, the time course dynamic of α-Troph-Gam-iRBCs IgM and IgG in

the acute *P. falciparum*-infected non-endemic plasmas), show that the flow cytometry assay

can accurately detect Troph-iRBCs IgM levels despite the low level or absence in endemic

individuals (Fig 5). The increase of IgG with age (*i.e.,* exposure in endemic individuals) and

decrease of IgM with age gives insight into the reason for the low α-Troph-Gam-iRBCs IgM

levels in the endemic plasma (exposure leads to a build-up of immunity that replaces low-

affinity IgM by an affinity-matured IgG response) (81-85).

DISCUSSION

Our study demonstrates that *P. falciparum* sexual conversion (SC) is significantly influenced by host hemoglobin genotype. Both *ex vivo* and *in vitro* assays showed that parasites growing in RBCs carrying HBB mutations HbAS and HbAC had higher SCRs than those in HbAA. This genotype-dependent modulation of SC was accompanied by a developmental delay in parasites within HbS- and HbC-infected RBCs but was not associated with differences in IgG or IgM responses against trophozoite- and early gametocyte-infected RBC surface antigens. Together, these findings indicate that host RBC intrinsic properties, rather than humoral immunity, underly altered parasite reproductive strategies in HBB mutant backgrounds.

Amoah et al., (2020) (86) analyzed the effect of HBB genotype on SC of field isolates from Ghanaian children aged 6-15 years with uncomplicated malaria, and reported a non-significant trend of decreasing SCRs from HbAA to HbAC and HbAS. The discrepancy with our results is most likely methodological, because quantifying SC in natural infections is challenging as parasites in freshly collected blood include morphologically indistinguishable sexual and asexual rings. The authors used prolonged culture (≥4 days) to distinguish sexual stages from asexual forms, similarly to Usui et al., (2019) (41). The limitation of this approach is the timing mismatch where SCRs are measured as the ratio of stage II gametocytes on day 4 (D4) by the total parasitemia on day 0 (D0). This can underestimate SCR because parasites contributing to the denominator (D0) are counted before culture adaptation affects viability, while the numerator (D4) includes only gametocytes that survive and mature after several days in *ex vivo* conditions, during which parasite loss can occur. Likewise, limitations have been reported when quantifying low SCRs (< 0.3%) in low-density natural infections by LM (41). For instance, at 0.05% parasitemia and 0.3% SCRs, they would be detecting five gametocytes among 30,000 RBCs. This approach provides limited sensitivity and resolution and could explain why up to 24% of samples were reported as gametocyte negative.

We addressed these limitations using an adapted *ev*SCA to quantify next cycle sexual conversion (39). We measure whether in-host exposure in individuals with RBCs with different HBB genotypes leads to stimulation of sexual commitment. We quantify the in-host effect by assessing the subsequent sexual conversion (next cycle, *ex vivo*) as number of stage I gametocytes (gamI) per number of schizonts and gamI (overall parasites) 24 hours post culture (hpc). Quantification is done by immunofluorescence, gamI detected by Pfs16 and overall parasites by DAPI (87). Inclusion of PKG inhibitor ML10 prevented reinvasion, allowing numerator and denominator measurements for the SCR at the same time point, while enrichment using LD columns improved sensitivity. This approach reliably quantified sexual conversion in infections with densities as low as 0.05% (~2,500 parasites/μl) and detected a minimum of 38 stage I gametocytes per sample, with no gametocyte-negative cultures.

We validated *ex vivo* findings using an *iv*SCA with a *gexp02-reporter* line (75), allowing early detection of sexual rings. While our SCR results in *iv*SCA were broadly consistent with those of Flage, B., et al., (2023) (25) for HbAA and HbAS RBCs (median SCRs: 10.8% vs. 10% in HbAA; 16.5% vs. 15% in HbAS), the results are not directly comparable due to clear methodological differences in how SCR was measured. In our assay, SCR is defined and measured as the proportion of *gexp02*-expressing parasites among total parasites at the next cycle conversion (generation 1), while sexual parasites were quantified at generation 1 and total parasitemia at generation 0 in (25), making direct comparison impossible.

We observed a partial developmental delay in parasites infecting HbAS and HbAC RBCs, and a more pronounced arrest or extreme delay in HbSS and HbSC under low-oxygen conditions (5% O_2), both by LM and FC, consistent with the findings of Archer, N.M., et al., (2018) (29). Later time-point measurements (~35-40 hpi) increased SCRs in endemic HbAA, HbAC, and HbAS RBCs compared with ~25-30 hpi, suggesting delayed sexual marker expression and thus underestimation of SCR at the earlier time point due to developmental delay. Of note, parasites

grown in control non-endemic HbAA RBCs showed stable SCRs across both time points. The

observed differences between endemic and non-endemic HbAA RBCs may reflect unmeasured

host genetic factors (*e.g.,* blood group variants or other erythrocytic polymorphisms). In HbSS

and HbSC RBCs, we observe SCRs similar to HbAA and lower than HbAS and HbAC,

consistently across both measurement time points. This contrasts with findings in (25), who

reported markedly higher SCR in HbSS RBCs (35% vs. 6.96% in our study) and no

developmental delay at day 2 of culture, although this was not quantitatively assessed. As noted

above, differences in the timing and approach used to quantify SCR make the results not

comparable between studies.

Under choline-depleted conditions, SCRs in HbSS and HbSC increased modestly (from <10%

to 10-20%) but remained below levels in HbAA, HbAC, and HbAS RBCs (>30-40%). This is

likely due to severe developmental delay causing under-detection of sexually committed rings

at the measured time points rather than inhibited sexual commitment. Overall, our data

indicates that accurately assessing SC in HbSS and HbSC RBCs is challenging with this

approach due to developmental delay. Measuring sexual commitment one cycle earlier, for

instance through *GDV1* activation, might circumvent this issue, although this is experimentally

demanding because *GDV1* is expressed transiently and at low levels. Given the very low

prevalence of HbSS and HbSC genotypes in malaria-endemic populations, the added value of

such complex assays for transmission studies is likely limited.

We evaluated potential confounding factors that were previously to SC (52, 58, 59, 88-92),

including reticulocyte abundance, parasitemia, and ABO blood group. No significant

differences in reticulocyte percentages were observed among HbAA, HbAC, and HbAS RBCs,

and reticulocyte levels did not correlate with SCRs at 35-40 hpi. Similarly, neither baseline

parasitemia nor ABO blood group influenced SCR in either the *ev*SCA or *iv*SCA. Together,

these results indicate that the observed increase in SC is specifically associated with HBB

genotype rather than these other host-related variables.

The increase in SC in RBCs with HBB mutant genotypes *in vitro*, in the absence of plasma and

white blood cells, points to RBC-intrinsic effects rather than host immune responses or soluble

plasma factors. HbS and HbC RBCs exhibit elevated levels of free heme (93), linked to

increased oxidative stress that leads to polymerization of HbS under low oxygen conditions

(94), and delays parasite development (29). The oxidative stress inhibits actin polymerization

in infected HbS and HbC RBCs (27), which alters protein trafficking and PfEMP1 display (66,

69). This reduces cytoadherence (28), and membrane flexibility (95). Oxidative stress also

modifies RBC micro RNAs that can hybridize with parasite mRNAs (65), directly affecting

parasite development and metabolism, and increasing gametogenesis (25). In (25), authors

showed that heme directly induces SC. Taken together, these findings suggest that the effect

of HbAC and HbAS RBCs on SC *in vitro* is likely driven by oxidative stress caused by free

heme within these RBCs, mediated by their specific miRNA profiles. The stronger effect *ex*

vivo, particularly in HbAS (7.07- vs ~2.05- fold *in vitro*), probably reflects additional *in vivo*

factors such as parasite genetic diversity (41, 87), or host factors not captured *in vitro* (87, 96).

To further investigate potential *in vivo* influences, we investigated the humoral immune

response in the different HBB genotypes. Reduced antigen presentation, cytoadherence, and

increased splenic clearance (28) could plausibly lead to an altered immune response in HbAC

and HbAS. However, epidemiological studies examining immunity in individuals with HbAS

and HbAC genotypes have yielded conflicting results. Older children with HbAS have been

shown to be more protected against the establishment of parasitemia, suggesting an acquired

mechanism of protection (67). In contrast, in (97) no differences were found in antibody

responses between individuals with HbAA, HbAS, and HbAC genotypes against their panel of

491 immunoreactive *P. falciparum* antigens. More recent studies have shown that serological

responses tend to be lower in individuals with HbAC and HbAS when focusing on specific surface antigens (98, 99). In a cross-sectional study conducted in Ghana about circulating inflammatory cytokines, has shown that expression levels of CXCL10, TNF-α, CCL2, IL-8, and IL-6 varied by hemoglobin genotype (100).

Given that differences in serological responses appear specific to certain surface antigens, we chose to measure IgG and IgM levels against intact infected RBCs. This approach enables the assessment of immune responses to native surface antigens, minimizing potential artifacts arising from non-native protein folding, glycosylation, or conformation. We focused on trophozoite-infected RBCs (25-30 hpi), as surface antigen presentation (*i.e.,* PfEMP1 expression) begins around 16 hpi and plateaus at approximately 24 hpi (101). Moreover, commitment to SC occurs between 17-37 hpi (39, 52), suggesting that any antibody-mediated induction of sexual commitment would need to occur at this stage.

We compared IgG and IgM responses to trophozoite- and gametocyte-infected RBCs (Troph-Gam-iRBCs) using plasma samples from Burkinabé individuals carrying HbAA, HbAC, HbAS, HbCC, and HbSC genotypes, assessed by flow cytometry. Our analyses revealed no significant differences in IgG or IgM levels against Troph-Gam-iRBCs across genotypes. IgM responses, as well as responses against gametocyte-infected RBCs, were generally low or absent. Therefore, we could not confirm genotype-dependent differences in humoral responses to these parasite stages and consider it unlikely that antibodies directly drive the differential SC observed in these individuals. Nonetheless, this does not exclude the possibility that genotype-specific antibody responses to later parasite stages, particularly during gametocytogenesis, as suggested by recent studies on gametocyte antigenicity (102, 103), could influence transmission potential, either by promoting clearance of developing gametocytes in the human host or by inducing gametocyte sterility within the mosquito vector (104).

Overall, this study robustly demonstrates elevated sexual conversion of *P. falciparum* parasites

in HbAS and HbAC RBCs both *in vitro* and *ex vivo*. These findings further support the view

that *P. falciparum* can adjust its reproductive investment in response to stress-related cues. In

mutant HBB RBCs, elevated levels of free heme can induce developmental delay and thereby

compromise survival within the human host; increased investment in sexual conversion may

thus represent a parasite survival strategy (25, 29, 63).

Previous studies have shown that microenvironmental factors such as LysoPC depletion,

nutrient limitation and exposure to antimalarial drugs can modulate malaria transmission by

changing the rate of parasite's SC. How factors such as free heme, are being translated into a

signal received by epigenetic regulators controlling the sexual commitment gene *ap2-g* or *gdv1*

remains an open question (25, 39).

At the population level, HBB mutant genotypes have been linked to increased gametocyte

carriage and higher mosquito infectivity. In Burkina Faso, it has been reported a two- to four-

fold higher infection rates in HbAS, HbAC, and HbCC carriers (30). Similar trends were

observed in Mali and Cameroon, where HbAC and HbAS individuals carried more

gametocytes and were more infectious to mosquitoes than HbAA (33, 34). These population-

level associations are consistent with our findings, and suggest that increased SC in mutant

HBB RBCs may contribute to enhanced transmission potential.

MATERIALS AND METHODS

659 Details about reagents and tools are accessible in Table S7.

Study design

Sample collection for the present study was embedded in a longitudinal cohort study assessing

the effect of host factors on transmission potential (*InHost* study, ethical advice reference

1261/18 and 19/06/064), conducted in the Nanoro Department, a rural, malaria-hyperendemic

region in Central-West Burkina Faso (Fig 6) (105) [Characterizing the *Plasmodium falciparum*

infectious reservoir asymptomatic asexual and gametocyte stage across off peak and peak

transmission seasons in a rural and high-transmission setting in Burkina Faso: a longitudinal

2-year cohort study, *manuscript in preparation*]. Briefly, 871 study participants of all ages

from four villages (Nanoro, Nazoanga, Séguédin and Soum) were followed-up for two years

(March 2019 - February 2021) through six cross-sectional surveys conducted during both high

and low transmission season to assess asexual and sexual parasite carriage and risk factors.

Active case detection was complemented by passive case detection at peripheral health centers.

673
Fig 6. Map of the study area. The inset map highlights the location of the Nanoro department of Boulkiemdé
Province in Central Burkina Faso, with the black dot indicating the position of the capital, Ouagadougou.

For this study, from all participants of the *InHost* study (*n* = 871), 864 individuals were

genotyped for their human HBB genotype (see section on "Hemoglobin beta sequencing").

Participants were then selected for this sub-cohort based on their genotype. A venous blood

sample was collected from all selected participants, and screened for *P. falciparum* infection.

Samples from *P. falciparum*-infected individuals were used to perform *ex vivo* sexual

conversion assays (*ev*SCAs) and samples from non-infected individuals were preserved to

perform *in vitro* sexual conversion assays (*iv*SCAs). We included all individuals carrying the

HBB genotypes HbAC ($n = 72$), HbAS ($n = 50$), HbCC ($n = 2$), HbSC ($n = 7$), and a random

subset of individuals with the HbAA genotype ($n = 94$) (Fig S6). In total, 225 individuals were

listed (age groups "5 - 19 years" [$n = 166$] and "more than 20 years" [$n = 59$]). Listed

individuals were visited by a nurse at home and were invited to the Clinical Research Unit of

Nanoro (CRUN, Burkina Faso) for screening and enrolment into the study. Population

screening for *P. falciparum* infections and enrolment took place over a two-month period

(August-September 2022).

At CRUN, the eligibility was assessed based on inclusion and exclusion criteria for the *ev*SCAs

(*P. falciparum*-infected individuals) and *iv*SCAs (non-infected individuals). Inclusion criteria

consisted of: being enrolled in the main *InHost* study, a confirmed *P. falciparum* malaria

infection (for the *ev*SCAs) or a confirmed negative *P. falciparum* malaria infection (for the

*iv*SCAs), willing to participate in this study and undergo additional sampling, and signed

informed consent. Exclusion criteria consisted of: mixed infection with other *Plasmodium*

species (only for *P. falciparum*-infected individuals in the *ev*SCAs), antimalarial treatment

within the last two weeks, signs of severe malaria (*e.g.* prostration), severe chronic illness

besides hemoglobinopathies (*e.g.,* cancer, heart disease, kidney or liver disorders, HIV/AIDS,

etc.), and planning to leave the study region during the study period. Of the 225 listed

individuals, 162 were screened at CRUN, of whom 161 participants were selected as eligible

and were included within this sub-cohort (Fig S6). A 15 ml venous blood sample was collected

from each participant. After sample collection, blood samples were screened for *P. falciparum*

infections (see section on "*P. falciparum* diagnosis"). Samples with confirmed *P. falciparum*

infection and parasitemia by light microscopy (LM) $\geq 0.05\%$ (*i.e.* ≥ 2500 parasites/µl) were

processed for *ev*SCAs at CRUN (Table 3). Blood samples negative for *P. falciparum* infection

were processed and preserved for *iv*SCAs conducted at the Institute of Tropical Medicine

(ITM), Antwerp (Table 3, see section on "*iv*SCAs"). *P. falciparum*-infected individuals, as

detected by LM and confirmed with *P. falciparum var* gene acidic terminal sequence (*var*ATS)

quantitative PCR (qPCR), were treated with Artemether-Lumefantrine (AL) following national

guidelines and were asked to return three days later to confirm infection clearance.

Sample size was calculated based on a power analysis assuming an expected effect size of 50%,

a 5% significance level, and 90% power. This sample size was considered sufficient to detect

differences in sexual conversion by HBB genotype, based on effect sizes reported in previous

studies (58). To account for potential variability in HBB genotype groups, a minimum of 8

participants per HBB genotype group was required. To ensure sufficient sampling and

successful assays per HBB genotype group, the sample size was upscaled to a maximum of 15

participants per HBB genotype group.

To increase the number of HbSC and HbSS blood samples for *iv*SCA, additional blood samples

were collected from patients with hemoglobinopathies (HbSS and HbSC) visited at the

University Hospital of Antwerp (UZA), Belgium. A total of 13 adults (aged 19-37 years),

routinely followed at UZA's Hematology Unit, were invited to donate a blood sample with

exclusion criteria of fever. The inclusion period was between October 2023 and April 2024.

Table 3: Number of samples (n) per *P. falciparum* infection status at the time of sample collection and HBB
genetics of the donor, for the *ex vivo* sexual conversion assays (*ev*SCAs), *in vitro* sexual conversion assays
(*iv*SCAs) and Immunoglobulin (Ig) detection assays. Note: *P.falciparum-infected (Pf+)* samples used in the
*ev*SCAs must have a parasitemia $\geq 0.05\%$ (*i.e.,* ≥ 2500 parasites/µl), as the established limit of accurate assessment
of sexual conversion with these assays. Abbreviations: *Pf, P. falciparum*, Hb, hemoglobin; CRUN, Clinical
Research Unit in Nanoro; ITM, Institute of Tropical Medicine in Antwerp; UZA, University Hospital in Antwerp.
729

	HbAA	HbAC	HbAS	HbSC	HbCC	HbSS	Total
EvSCAs							
Pf+ (CRUN, total n)	14	12	6	-	-	-	32
IvSCAs							
Pf- (total n)	5	5	4	2	-	12	28
CRUN (n)	5	5	4	1	-	-	
UZA (n)	-	-	-	1	-	12	
Ig detection							
Pf+ (CRUN)	13	13	13	3	1	-	43
Pf- (CRUN)	5	5	5	2	1	-	18
Total n (*Pf+* and *Pf-*)	18	18	18	5	2	-	61

Plasmodium falciparum diagnosis

P. falciparum malaria diagnosis was done by RDT using the Bioline™ malaria antigen P.f®

test (05FK50, Abbott) (recommended by the National malaria Control Programme (NMCP) in

Burkina Faso), following the manufacturer instructions. In addition, malaria parasites were

detected and quantified by LM following standard protocols at CRUN (106). In brief, venous

blood smears were stained with Giemsa and examined until 200 white blood cells were

counted. Parasitemia was calculated based on the number of asexual parasites per white blood

cells, expressed as asexual parasites per microliter of blood, assuming a white blood cell count

of 8,000 per microliter. The presence of gametocytes and other *Plasmodium* species (*P. ovale*

and P. malariae) was assessed in all positive blood smears (107). Blood samples that tested

positive for *P. falciparum* by both RDT and LM were classified as *P. falciparum*-infected.

Samples that were negative by LM (regardless of the RDT result), were identified as non-

infected (Table 3 and Fig S6). Negative infection with *P. falciparum* was confirmed by *var*ATS

qPCR, using five µl of DNA (extracted with QIAamp DNA Mini Kit, Qiagen Germany), as

previously described (105, 108) (Fig S6). The limit of detection of the *var*ATS qPCR was 0.1

parasite/µl (in our hands), with samples showing Ct > 39.7 considered negative. DNA from the

P. falciparum 3D7 strain served as a positive control, while negative controls included DNA

from non-infected human blood and a no-template control (NTC).

Sample collection and processing

At CRUN, a 15 ml venous blood sample was collected in an Acid Citrate Dextrose (ACD) tube

and the residual blood in the needle was used for *P. falciparum* diagnosis by RDT and LM (see

section on "*P. falciparum* diagnosis"). The sample was then immediately transported to the

laboratory for processing. Samples were centrifuged (1,648 g for 5 minutes) and washed two

times with RPMI to isolate the RBC pellet. RBC pellets from *P. falciparum*-infected blood

samples were immediately put in culture for the *ev*SCAs (see section on "*Ex vivo* sexual

conversion assays (*ev*SCAs)". RBC pellets from non-infected samples were cryopreserved (1:3 dilution in cryopreservation solution containing 3% sorbitol and 0.65% NaCl) and stored directly at -80°C until shipment to ITM, where preserved in liquid nitrogen until use in *iv*SCAs. Two hundred microliters of whole blood were used for DNA extraction and confirmation of *P. falciparum* infection by *var*ATS qPCR (see section on "*P. falciparum* diagnosis). At UZA, 15-20 ml of non-infected venous blood samples were collected in EDTA tubes and immediately transported at 4°C to ITM for processing, done as described at CRUN. The resulting RBC pellets were cryopreserved (1:3 dilution in cryopreservation solution containing 3% sorbitol and 0.65% NaCl) by direct snap-freezing in liquid nitrogen for subsequent *iv*SCAs. Additionally, ABO and Rhesus D blood groups were determined with agglutination tests (BIO-RAD, REF: 801320, 801345, 802032).

Hemoglobin beta sequencing (HBBseq)

In samples collected at CRUN, PCR amplification of the HBB gene (NCBI_Gene:3043) was performed to identify single nucleotide polymorphisms (SNPs) using PCR conditions adapted from (109) and (110). Primer pairs were designed to amplify a 599 bp product (Table S7). PCR reactions were performed in a total volume of 25 µl, containing 2.5 µl of DNA template, 10× Buffer B, 5 mM of each dNTP, 0.3 µM of each primer, 25 mM of $MgCl_2$, and FIREPol® DNA Polymerase (Solis Biodyne, Ref: 01-01-00500). Thermal cycling conditions were as follows: an initial denaturation at 94°C for 6 minutes, followed by 40 cycles of 60.6°C for 30 seconds, and 72°C for 1 minute, with a final extension at 72°C for 10 seconds. After Sanger sequencing, chromatograms were analysed using BioEdit 7.2. (Informer Technologies) to identify the wild-type and the mutant HBB codons (Table 4). In samples collected at UZA, the HBB genotype sequences were determined by capillary hemoglobin electrophoresis at UZA as described elsewhere (111).

Table 4. HBB codons and genotypes identified by Sanger sequencing analysis of the HBB gene (HBB-seq).

HBB allele	Mutation	HBB codon
HbA (wild-type)	-	GAG
HbS (mutant)	c.20A>T = E (Glu)> V (Val)	GTG
HbC (mutant)	c.19G>A = E (Glu) > K (Lys)	AAG
HbSC (double mutant)	HbS mutation (Glu→Val) + HbC mutation (Glu→Lys)	AAG + GTG (112)

Ex vivo sexual conversion assays (*ev*SCAs)

Ex vivo sexual conversion assays in fresh *P. falciparum* field isolates

P. falciparum-infected venous blood samples (15 ml) collected from individuals carrying different HBB genotypes (Table 3; HbAA n = 14, HbAC n = 12, HbAS n = 6; total n = 32) were cultured *ex vivo* to measure sexual conversion rates (SCRs), assay optimization and validation described in *Supporting information file*. Cultures at 3% Hct and 20 ml volume were prepared by mixing 600 µl of *P. falciparum*-infected RBC pellet with 20 ml of RPMI 1640 medium containing 0.5% Albumax II (Gibco, 11021-029) and 2 mM choline chloride (Sigma-Aldrich, C7527) (52, 75). Parasites at the ring stage were incubated using candle jar method under static conditions until ~24 hours post-culture (hpc). At this time, sexually committed rings present in the venous blood sample are expected to have developed into stage I gametocytes expressing Pfs16, while asexual parasites will have progressed to the schizont stage (Fig 2A). During culture, at ~5 hpc, ML10 (80 nM), a cGMP-dependent protein kinase inhibitor, was added to prevent schizont rupture and reinvasion (39, 113). Giemsa-stained smears were examined by LM to confirm that most parasites had reached either stage I gametocytes or schizonts stages. At ~24 hpc, parasites were harvested and purified using MACS LD columns (Miltenyi Biotec). In brief, cultured parasites are centrifuged (1,648 g, 5 min) and pelleted infected-RBCs (iRBCs) were diluted 1:4 with culture medium, loaded onto

an LD-MACS column, and washed four times with 1 ml medium to separate non-infected

RBCs and ring-stage-infected RBCs (first elution). The column was then removed from the

magnet, and stage I gametocytes and schizonts were recovered by gravity elution, followed by

gentle plunger flushing twice with 3 ml medium (second elution). Recovered parasites were

pelleted (1,648 g, 5 min), resuspended 1:2 with non-infected RBCs, and diluted 1:2 in PBS.

From the second elution, four thin blood smears (3 µl each) were prepared and left to air-dry;

three were used for on-site IFAs, and one was Giemsa-stained for LM to verify MACS

enrichment.

Immunofluorescence assays (IFAs)

IFAs were performed at CRUN (Burkina Faso) as previously described (39, 52, 75), with

modifications. Slides were stained in batches of four to eight cultures with three slides per

culture, to optimize processing and staining consistency. Primary antibody mouse anti-Pfs16

(1:400, kindly provided by R. Sauerwein and T. Bousema, Radboud University) (76) and

secondary antibody goat anti-mouse IgG (H+L) Alexa Fluor 488 (1:1000; Abcam, ab150113)

were used, with incubations and washes under static conditions. Slides were mounted with

Aqueous Fluoroshield (Abcam, ab104139) containing DAPI (0.0002%) and stored at 4°C until

shipment to ITM (Belgium). At ITM, slides were visualized within six months of preparation

on an ECHO Revolve epifluorescence microscope (ECHO, San Diego, CA) to determine SCRs

(Fig 2F). Analysis was blinded to HBB genotype. Stage I gametocytes were identified as Pfs16-

positive cells (green) and counted among a minimum of 1,000 DAPI-positive cells (blue) per

slide (stage I gametocytes and schizonts). Slides with fewer than 1,000 DAPI-positive cells

were excluded from further analysis during the first reading. For each culture ($n = 35$), three

of the four prepared slides were stained by IFA and examined by a primary reader. Slides from

three cultures were excluded due to insufficient DAPI-positive cells. From the remaining

cultures ($n = 32$) a random subset of ~10% (4/32), with three IFA slides per culture, was

examined by a second reader to assess inter-reader reliability via intraclass correlation

coefficient (ICC) analysis, based on a two-way random effects model, absolute agreement,

single measures [ICC(2,1) (114)]. For each IFA slide, SCRs were calculated as the proportion

of Pfs16-positive parasites among total DAPI-positive cells. Across the three replicate IFA

slides per culture, the mean SCR (±SD) was computed and used to calculate and report the

median SCR [IQR] by HBB genotype (Figs 4A and 4B), statistical differences were assess as

described for proportions in "Data analysis" section. Fold-change in SCR for HbAC and HbAS

was calculated relative to HbAA by dividing each mutant genotype's median SCR by that of

the wild-type (Fig 2C). Statistical analysis was performed as described in the "Data analysis"

section: Differences in initial parasitemia across HBB genotypes (continuous non-parametric

variables) (Figs 2B-D) and potential confounding effect of initial parasitemia on SCRs (non-

parametric correlation across continuous data) (Fig S1A).

Stable Pfs16 fluorescence was confirmed by the counting of ≥1 stage I gametocyte per 10

fields. For each culture, the three replicate IFA slides were analyzed to quantify SCRs. The

mean, minimum and maximum SCR were calculated across replicates. To assess inter-slide

reproducibility (technical repeatability) of SCR measurements, an ICC (2,1) analysis (two-way

random effects, single measures, consistency) was performed using SCR values from the three

replicate slides per culture (n = 32). Counts of stage I gametocytes and total parasites were

recorded for each IFA slide and summarized as median (Q1-Q3). From 5 out of 32 cultures

(15.6%) specificity controls were performed by staining one slide per culture without primary

antibody (negative controls). Stage V gametocytes (also express Pfs16), identified by their

shape, were excluded from the counting as they are a product of sexual conversion occurring

>10 days prior to the sexual commitment cycle under investigation. Giemsa-stained slides from

each *ev*SCA were examined by LM to estimate parasite purification by MACS and confirm

approximate parasite age/stage to assure parasites were sufficiently mature to express Pfs16

(starts expression at ~20 hpi (39)).

In vitro sexual conversion assays (*iv*SCAs)

In vitro sexual conversion assays (*iv*SCAs) were performed as follows (Fig 3A). NF54-*gexp02-*

Tom parasites (52, 75) were routinely maintained with parasitemias of less than 2 % in O+

RBCs (3% Hct) in RPMI 1640-based medium supplemented with 0.5% Albumax II (Gibco,

ref: 11021-029) and 2 mM choline chloride (+Choline; Sigma-Aldrich, C7527), analogous to

LysoPC supplementation (55, 75, 79). Cultures were incubated at 37°C with shaking conditions

(70 rpm) in a gas mixture (5% O_2, 5.5% CO_2, balance N_2). To minimize bias due to prior culture

duration, all cultures were initiated from parasites of the same cryopreserved batch of an

expanded culture.

To obtain cultures of a well-defined age window, we used Percoll/sorbitol synchronization (Fig

3A). Percoll-purified schizonts (~40-44 hpi, Generation -1) were used to establish fresh

cultures with thawed cryopreserved RBCs (day 3 post thawing) with different HBB genotypes

(exposure, Generation 0), now cultured in static conditions. Thawing of RBCs was performed

using a three-step protocol, sequentially adding three volumes of 3.5% NaCl solution (10× the

RBC volume each time), followed by RPMI washes and centrifugation (1,648 g, 5 min) as per

standard procedure (115). Approximately 5 hours later, when parasites have reinvaded the new

RBCs, cultures were subjected to 5% D-sorbitol lysis (39, 75) to obtain cultures of a defined

0-5 hpi age window. Post-sorbitol initial parasitemia was then measured by flow cytometry,

and parasites were cultured in two parallel subcultures (~200 µl final volume) per sample. One

sub-culture was maintained in +Choline medium (non-inducing sexual conversion conditions),

and the other in choline-free medium (-Choline; inducing conditions). Cultures were

maintained under these respective conditions for two generations (48 hpi, until ~35-40 hpi in

Generation 1, Fig 3A, hpi calculations are based on a 44 hours replication cycle empirically

established for this parasite line in our lab). In each experiment we tested RBCs with at least two different HBB genotypes. Samples with different HBB genotypes were used in the experiments if the recovered RBC pellet volume after thawing was ≥ 30 µl after dilution with RPMI to 50% Hct. From a total of 20 cryopreserved non-infected RBC samples from CRUN that were confirmed to be negative for *P. falciparum* by *var*ATS qPCR, 15 samples had a RBC pellet volume ≥ 30 µl after thawing and dilution in RPMI to 50% Hct (Fig S6). The mean recovered RBC pellet volumes after thawing were: HbAA, 178 µl (range: 50−596 µl); HbAC, 111 µl (range: 30−320 µl); HbAS, 246 µl (range: 64−630 µl); HbSC, 558 µl (range: 35−1080 µl); and HbSS, 600 µl (range: 300−1120 µl). These samples were included within the *iv*SCAs, in addition to non-infected, cryopreserved RBC samples from the UZA HbSS and HbSC (Fig S6). Total numbers of samples tested per HBB genotype and experiment are shown in Table 1. Flow cytometry (FC) analysis was used to measure parasitemia, assess parasite stage development (*i.e.,* proportion of parasites reaching the intraerythrocytic schizont stage or late-stage proportions), and quantify sexual conversion at defined time points during the assay (Fig 3A). Parasitemia was determined at ~0-5 hpi (rings), together with (1) the proportion of schizont stages at ~35-40 hpi in Generation 0 and 1, and (2) sexual conversion rates (SCRs) at ~25-30 hpi (large rings/trophozoites) and ~35-40 hpi (schizonts) in Generation 1 (Fig 3A). The standard timepoint to measure sexual conversion with this parasite line is ~25-30 hpi to ensure detection of all sexually committed parasites, as TdTom-expression under the *gexp02* promoter starts around ~10-15 hpi (75). The ~35-40 hpi timepoint was included at Generation 1 to account for potential delayed stage development in mutant HBB genotypes (see section on "Assessment of parasite stage development in RBCs with different HBB genotypes"). Four independent experiments were conducted, with RBC samples included in each experiment detailed in Table 1. Each experiment included a fresh HbAA blood sample to control the effect of RBC storage after cryopreservation.

Samples for FC analysis (using a CytoFLEX LX flow cytometer, Beckman Coulter) were

prepared by resuspending parasite cultures (1:3) in PBS containing 1% BSA, followed by

staining with Hoechst 33342 (1:500, BD Pharmingen) in a final volume of 100 µl per well.

Samples were then incubated at 37°C for 30 minutes. To measure parasitemia and proportion

of schizont stages, FC analysis (with 100,000 events collected per sample) was set to detect

Hoechst 33342 (350 nm laser; 450/40 filter). RBCs were gated by plotting side scatter area

(SSC-A) versus forward scatter area (FSC-A), with singlets selected using FSC-H versus FSC-

A. Parasites were identified within the singlet gate by Hoechst fluorescence. Hoechst-positive

events were defined as exceeding background levels from unstained negative controls. To

quantify SCRs, Hoechst fluorescence was combined with the fluorescent reporter TdTom (561

nm laser; 610/20 filter) that is integrated in the NF54-*gexp02-Tom* line to distinguish sexual

(Hoechst+/tdTom+) from asexual (Hoechst+/tdTom-) parasites. SCRs were calculated as the

proportion of Hoechst+/tdTom+ events among all Hoechst+ events.

SCRs were expressed as a percentage (median [IQR]) per HBB genotype (Figs 3B and 2A-C).

Fold-changes in SCRs were calculated per experiment (Table 1) by dividing the SCR of each

cryopreserved mutant RBC sample by the SCR of a fresh, non-endemic HbAA RBC control

sample. (Figs 3C and S3D). Statistical analysis was performed as described in the "Data

analysis" section: differences in SCRs and post-sorbitol initial parasitemia across HBB

genotypes tested as described for proportions, differences in fold-change as described for

continuous data across categories (Figs 3B-D and S3A-E).

Reticulocyte percentages were measured for all RBC samples on the same day as Percoll

synchronization (Generation -1, Fig 3A). Briefly, RBC suspensions were mixed 1:1 with

reticulocyte stain (Sigma-Aldrich), incubated 15 min at room temperature, and used to prepare

a thin blood smear for LM estimation of reticulocyte percentage. We determined the ABO and

Rhesus D blood groups in all samples using agglutination tests with Seraclone™ Anti-A, Anti-

B, and Anti-D (RH1) reagents (Bio-Rad).

Assessment of parasite stage development in RBCs with different HBB genotypes

Parasite stage development was evaluated during the *iv*SCAs (Fig 3A) at ~35-40 hpi

(Generation 0) and ~25-30 hpi/~35-40 hpi (Generation 1) using LM and FC. The proportion of

rings, trophozoites and early/late schizonts was quantified at ~35-40 hpi in Generation 0 and

~25-30 hpi/~35-40 hpi in Generation 1 by LM of Giemsa-stained thin blood smears (100

parasite counts per sample; Fig 4A) in a subset of RBC samples (HbAA $n = 1$; HbAC $n = 1$;

HbAS $n = 2$; HbSS $n = 2$; HbSC $n = 2$). FC was used as described earlier to determine the

percentage of "late stages". For this a distinct population of cells were gated with increased

Hoechst 33342 fluorescence (increased DNA content due to multiplication of the nuclei in

schizont stages), within the Hoechst+ cells (Fig 4B). Upon confirming consistent results

between LM and FC after screening the same samples by FC, FC was subsequently used in all

*iv*SCA samples ($n = 28$: HbAA $n = 5$; HbAC $n = 5$; HbAS $n = 4$; HbSS $n = 12$; HbSC $n = 2$;

and fresh non-endemic HbAA CTRLs $n = 6$) to determine the proportion of late-stages (*i.e.,*

schizonts) by HBB genotype (Figs 4B and 4C).

From LM data we calculated: (1) the percentage of rings, trophozoites and early/late schizonts

(Fig 4A), (2) the mean parasite age using the age in hpi (116) of each stage (rings, trophozoites,

schizonts) and the observed proportions; and (3) the developmental delay (in hours) by

subtracting mean parasite age of the mutant genotypes from the wild-type HbAA genotype.

Finally, for each genotype, we compared the developmental delay between Generations 0 and

1 by calculating the difference in mean parasite age at ~35-40 hpi between both generations.

Then, the development of each independent NF54-*gexp02-Tom* culture was classified as:

expected development, *delayed development* or *(partially) arrested/extremely delayed*

development (see detailed analysis in *Supporting information file)*.

Flow cytometry assays to assess IgG and IgM immunity against *P.*
falciparum

To study the IgG and IgM levels against *P. falciparum* in individuals with different HBB

genotypes by flow cytometry, we selected endemic human plasma samples from the *InHost*

longitudinal cohort study. We classified samples as "endemic" or "non-endemic" based on the

country of origin. An "endemic" country has consistent, stable *P. falciparum* transmission,

while a "non-endemic" country has no local transmission, with infections typically imported

(117). As the expected variation and magnitude of potential differences among HBB genotypes

were unknown, we based the sample size on the minimum of 8 individuals per HBB genotype

used in the *ev*SCAs and *iv*SCAs. A maximum of 15-20 individuals per genotype was included

to ensure sufficient sampling. Selection criteria included equal representation of the most

frequent genotype groups (HbAA [$n = 18$], HbAC [$n = 18$], HbAS [$n = 18$], HbCC [$n = 2$], and

HbSC [$n = 5$]), *P. falciparum* infection status with a large proportion of active *P. falciparum*

infections ($n = 10$-15) and a smaller proportion of non-infected individuals ($n = 2$-5), age range

(1-19 years to exclude maternal antibodies), and homogeneous distribution amongst the

ethnicity (all Mossi) and village (Nanoro, Nazoanga, Soum, and Séguédin) groups. Table 3

shows the number of participants per HBB genotype included in the assays. The *P. falciparum*

transgenic line NF54-*gexp02*-Tom (52, 75) was cultured under the same conditions as

described in the *iv*SCAs section and Fig 3A, with the following modifications: (1) parasites

were cultured only in wild-type HbAA fresh RBCs and (2) flow cytometry assays were

performed on early trophozoite/stage I gametocyte-infected RBCs (Troph-Gam-iRBCs) at ~20-

25 hpi (Generation 1), following the protocol reported in (102) with the adaptations described

below.

Human plasma samples were pre-diluted in PBS with 1% Bovine Serum Albumin (BSA, Sigma

Aldrich) and heparin (2,500U; Sigma Aldrich) (PBS-BSA-Hep) at dilutions of 1:5000 (IgG)

and 1:100 (IgM). *In vitro* generated Troph-Gam-iRBCs (25-30hpi; Generation 1) were pre-

diluted to 0.6% Hct in PBS-BSA-Hep. Pre-diluted human plasma samples were incubated at a

1:10 dilution with pre-diluted Troph-Gam-iRBCs, in separate 96-well plates for IgG and IgM

assays for 1.5 hours at room temperature. After 2 washes with PBS-BSA-Hep, secondary

antibodies were added to the Troph-Gam-iRBCs for 30 minutes at 4°C at 0.6% hematocrit.

AlexaFluor488 (AF488) mouse anti-human IgG (Jackson ImmunoResearch Europe) and

PerCP/Cy5.5® mouse anti-human IgM (Abcam) were used at final dilutions of 1:640 and 1:320

in PBS-BSA-Hep for the IgG and IgM assays, respectively. After 2 washes with PBS-BSA-

Hep, the DNA stain Hoechst33342 (BD Pharmingen) was added to the Troph-Gam-iRBCs at

a final dilution of 1:5000 in PBS-BSA-Hep at 0.6% hematocrit for 30 minutes at 37°C. To

verify that the Troph-Gam-iRBCs measured contained exclusively red blood cells (no white

blood cells), each batch pre-diluted Troph-Gam-iRBCs (to 0.6% hematocrit in PBS-BSA-Hep)

were stained with APC anti-human CD236 (Glycophorin C) Recombinant Antibody

(Biolegend®) in triplicate at a 1:40 final dilution (Fig S4C). Immunostained samples were

analyzed on the flow cytometer CytoFLEX LX (Beckman Coulter), collecting 100,000 cells

per culture. We analyzed the median percentage of total, asexual and sexual-infected red blood

cells (Troph-Gam-iRBCs) that were positive for IgG and IgM binding to surface antigens on

the Troph-Gam-iRBCs (*i.e.,* α-Troph-Gam-iRBCs , α- Troph-iRBCs and α-Gam-iRBCs IgG &

IgM). Samples were considered positive for IgG or IgM if the levels of α-Troph-Gam-iRBCs

IgG & IgM exceeded that of a naïve non-endemic plasma from a volunteer (negative control

with HbAA genotype with no prior *P. falciparum* infection, from a non-endemic country [*i.e.,*

Belgium] collected at the clinic of the Institute of Tropical Medicine [ITM] in Antwerp). Three

non-endemic samples from an acute *P. falciparum*-infected traveller collected at the ITM clinic

at day 0, 21 and 23 of *P. falciparum* diagnosis (*i.e.,* acute *P. falciparum-infected* non-endemic

plasma D0, D21 and D23) were used as positive controls. A schematic of the gating strategy

to measure the surface reactivity of asexual (tdTom-) and sexual (tdTom+) Troph-Gam-iRBCs

(Hoechst33342+) to IgG (α-Troph-iRBCs IgG: Hoechst33342+/tdTom-/AF488+ and α-Gam-

iRBCs IgG: Hoechst33342+/tdTom+/ AF488+) or IgM (α-Troph-iRBCs IgM:

Hoechst33342+/tdTom-/PerCP-Cy5.5.+ and α-Gam-iRBCs IgM (Hoechst33342+/

tdTom+/PerCP-Cy5.5.+) in the analyzed endemic human plasma samples is represented in Figs

S4B and S4C.

To determine the optimal plasma dilution factor, a pool of five endemic and hyperimmune

plasma samples from *P. falciparum*-infected individuals in a sub-cohort within the *InHost*

study (*i.e., P. falciparum*-infected hyperimmune endemic plasma pool of symptomatic

individuals with high *P. falciparum* parasitemia ranging from 45,667 to 175,000 parasites per

µl) (53) was titrated for the IgG and IgM assay separately. For the IgM assay, four additional

plasma samples were titrated (*i.e.,* "naïve non-endemic plasma" and "acute *P. falciparum*-

infected non-endemic plasma at D0, D21 and D23") (Fig S4A).

During flow cytometry analysis, α-Troph-Gam-iRBCs IgG (Hoechst33342+/tdTom-

/+/AF488+) and α-Troph-Gam-iRBCs IgM (Hoechst33342+/tdTom-/+/PerCP-Cy5.5.+) levels

for each sample were expressed as total event counts, percentage of positive cells, and median

fluorescence intensity (MFI) per HBB genotype group. The α-Troph-Gam-iRBCs IgG and IgM

levels and signal were compared by HBB genotype group for the total Troph-Gam-iRBCs, and

the asexual or sexual Troph-Gam-iRBCs (Figs 5 and S5). To correct for the background signal

from the non-infected RBCs in the IgG and the IgM channel, the MFI of all samples (including

the positive and the negative controls) was expressed as a ΔMFI. For the asexual and sexual

Troph-Gam-iRBCs, ΔMFI was calculated as ΔMFI (AF488+/PerCP-Cy5.5.+) = ("MFI

IgG+/IgM+ of asexual [tdTom-]/sexual [tdTom+] Troph-iRBCs [Hoechst33342+]") - ("MFI

IgG+/IgM+ of non-infected RBCs [tdTom-/Hoechst 33342-]") within each plasma sample. For

the total Troph-iRBCs, ΔMFI was calculated as ΔMFI (AF488+/PerCP-Cy5.5.+) = ("MFI IgG+

or IgM+ iRBCs [Hoechst33342+]") - ("MFI IgG+ or IgM+ of non-infected RBCs

[Hoechst33342-]" of trophozoite stages incubated with PBS-BSA-Hep, instead of plasma).

Data analysis

1026

All flow cytometry data were analyzed using FlowLogic™ version 8.7 (Inivai

Technologies™). Statistical analyses were performed in RStudio software version

2025.09.1+401(©2025 Posit Software; RRID:SCR_000432), with p-values ($p \leq 0.05$)

considered significant. The statistical analysis type was chosen based on the nature of the

variables: proportions based on counts were assessed using binomial and beta-binomial

generalized linear models (GLM) and mixed models (GLMM) with Bonferroni adjustment for

the contrasts in pairwise comparisons. We chose beta-binomial models due to (possible)

violations of the assumptions of the binomial model (overdispersion), this was borderline

acceptable in "*ev*SCA" (GLMM) but not in "*iv*SCA" (GLM) and "IgG/IgM detection" (GLM).

Random factors were used to account for technical replicates in the GLMM for the "*ev*SCA".

To test for the relevance of other factors we also modelled the initial parasitemia ("*ev*SCA" &

"*iv*SCA"), and reticulocyte percentage ("*iv*SCA"). To address the liberal parametric Wald Z

estimates and confidence intervals, we also computed more conservative Bayesian estimates

and credible intervals using a Markov chain Monte Carlo (MCMC) approach when model

specification and convergence allowed. GLM and GLMM analysis results are reported in Table

S1. When the assumptions of the GLM and GLMM beta-binomial models were violated, we

performed Kruskal-Wallis tests as described below.

Continuous variables were first assessed for normality using the Shapiro-Wilk test to guide the

choice between parametric and non-parametric methods. Continuous non-parametric variables

across categorical groups with ≥2 levels were analyzed using the Kruskal-Wallis test, and when

significant, this was followed by Wilcoxon rank sum post-hoc tests for pairwise comparisons with Holm correction. For normally distributed variables, one-way ANOVA was applied when Bartlett's F-test indicated equal variances; when variances were unequal, Welch's ANOVA was used. Bonferroni post-hoc tests were performed for pairwise comparisons when ANOVA results were significant. Associations between categorical variables were evaluated using Pearson's χ^2 test. Continuous non-parametric variables were correlated using Spearman correlation.

In the *ev*SCAs and the *iv*SCAs, risk analyses stratified by HBB genotype were conducted to examine associations with demographic and clinical variables (*e.g.,* SCRs, gender, age, village, ethnicity, …). An additional risk analysis for factors associated with elevated basal SCRs included the same covariates with the addition of HBB genotype (Table S4 for the *ev*SCAs, and Table S5 for the *iv*SCAs).

Ethical approvals

The *InHost* study, a longitudinal cohort in Burkina Faso, was approved by the ethical review board of the Comité d'Éthique pour la Recherche en Santé (CERS: 2018/10/131) in Burkina Faso, and by the IRB-ITM (1261/18) and IRB-UZA (19/06/064) in Belgium. The amendment for our nested study in the *InHost* study was approved by CERS (N°2022000140/MSHP/ MESRI/CERS), the IRB at ITM (1261/18), and the IRB at UZA (3347) in Belgium. Ethical approval for additional sample collection at the Hematology Unit at UZA was granted by the IRBs at ITM (1538/21) and UZA (3215).

ACKNOWLEDGEMENTS

We are grateful to Robert W Sauerwein and Teun Bousema (Radboud University, The Netherlands) for the anti-Pfs16 monoclonal antibody. We thank Núria Casas-Vila (ISGlobal) for the support and guidance in setting up the *ex vivo* and *in vitro* sexual conversion assays.

REFERENCES

1. World Health Organization (WHO) GMPG. World malaria report 2025: Addressing the threat of antimalarial drug resistance. 2025:213.

2. Hannemann A, Weiss E Fau - Rees DC, Rees Dc Fau - Dalibalta S, Dalibalta S Fau - Ellory JC, Ellory Jc Fau - Gibson JS, Gibson JS. The Properties of Red Blood Cells from Patients Heterozygous for HbS and HbC (HbSC Genotype). Anemia. 2011;2011:248527. (2090-1275 (Electronic)).

3. Haldane JB. Suggestions as to quantitative measurement of rates of evolution. Evolution; international journal of organic evolution. 1949;3(1):51-6.

4. Taylor SM, Cerami C, Fairhurst RM. Hemoglobinopathies: slicing the Gordian knot of Plasmodium falciparum malaria pathogenesis. PLoS Pathogens. 2013;9(5):e1003327.

5. Taylor S, Parobek C, Fairhurst R. Haemoglobinopathies and the clinical epidemiology of malaria: A systematic review and meta-analysis. The Lancet infectious diseases. 2012;12:457-68.

6. Friedman MJ. Erythrocytic mechanism of sickle cell resistance to malaria. Proc Natl Acad Sci U S A. 1978;Apr;75(4):1994-7. (0027-8424 (Print)).

7. Agarwal A, Guindo A Fau - Cissoko Y, Cissoko Y Fau - Taylor JG, Taylor Jg Fau - Coulibaly D, Coulibaly D Fau - Koné A, Koné A Fau - Kayentao K, et al. Hemoglobin C associated with protection from severe malaria in the Dogon of Mali, a West African population with a low prevalence of hemoglobin S. Blood. 2000;96(7)(0006-4971 (Print)):2358–63.

8. Modiano D, Luoni G Fau - Sirima BS, Sirima Bs Fau - Simporé J, Simporé J Fau - Verra F, Verra F Fau - Konaté A, Konaté A Fau - Rastrelli E, et al. Haemoglobin C protects against clinical Plasmodium falciparum malaria. Nature. 2001;414(6861)(0028-0836 (Print)):305–8.

9. Mackinnon MJ, Mwangi Tw Fau - Snow RW, Snow Rw Fau - Marsh K, Marsh K Fau - Williams TN, Williams TN. Heritability of malaria in Africa. PLOS Medicine. 2005; 2(12): e340(1549-1676 (Electronic)).

10. de Mendonça VR, Goncalves Ms Fau - Barral-Netto M, Barral-Netto M. The host genetic diversity in malaria infection. Journal of tropical medicine. 2012;940616(1687-9694 (Electronic)).

11. Lawrence C, Fabry Me Fau - Nagel RL, Nagel RL. The unique red cell heterogeneity of SC disease: crystal formation, dense reticulocytes, and unusual morphology. Blood. 1991;78(8)(0006-4971 (Print)):2104–12.

12. Rees DC, Williams TN, Gladwin MT. Sickle-cell disease. The Lancet. 2010;376(9757):2018-31.

13. Grosse SD, Odame I Fau - Atrash HK, Atrash Hk Fau - Amendah DD, Amendah Dd Fau - Piel FB, Piel Fb Fau - Williams TN, Williams TN. Sickle cell disease in Africa: a neglected cause of early childhood mortality. American journal of preventive medicine. 2011;41(6 Suppl 4)(1873-2607 (Electronic)):S398–S405.

14. Bunn Hf Fau - Noguchi CT, Noguchi Ct Fau - Hofrichter J, Hofrichter J Fau - Schechter GP, Schechter Gp Fau - Schechter AN, Schechter An Fau - Eaton WA, Eaton WA. Molecular and cellular pathogenesis of hemoglobin SC disease. Proceedings of the National Academy of Sciences of the United States of America. 1982;79(23)(0027-8424 (Print)):7527–31.

15. Zaabi. BKSKJEA. Hemoglobin C Disease: Treasure Island (FL): StatPearls Publishing; 2025 Jan-. Last Update: May 29, 2023.

16. Ballas SK, Lewis CN, Noone AM, Krasnow SH, Kamarulzaman E, Burka ER. Clinical, hematological, and biochemical features of Hb SC disease. American Journal of Hematology. 1982;13(1):37-51.

17.   Wellems TE, Fairhurst RM. Malaria-protective traits at odds in Africa? Nature genetics.
2005;37(11):1160-2.
18.   Natama HM, Rovira-Vallbona E, Krit M, Guetens P, Sorgho H, Somé MA, et al.
Genetic variation in the immune system and malaria susceptibility in infants: a nested case-
control study in Nanoro, Burkina Faso. Malaria journal. 2021;20(1)(1475-2875
(Electronic)):94.
19.   Simpore J, Pignatelli S Fau - Barlati S, Barlati S Fau - Musumeci S, Musumeci S.
Modification in the frequency of Hb C and Hb S in Burkina Faso: an influence of migratory
fluxes and improvement of patient health care. Hemoglobin. 2002;26(2)(0363-0269
(Print)):113–20.
20.   Bougouma EC, Tiono AB, Ouédraogo A, Soulama I, Diarra A, Yaro J-B, et al.
Haemoglobin variants and Plasmodium falciparum malaria in children under five years of age
living in a high and seasonal malaria transmission area of Burkina Faso. Malaria Journal.
2012;11(1):154.
21.   May J, Evans Ja Fau - Timmann C, Timmann C Fau - Ehmen C, Ehmen C Fau - Busch
W, Busch W Fau - Thye T, Thye T Fau - Agbenyega T, et al. Hemoglobin variants and disease
manifestations in severe falciparum malaria. JAMA. 2007;297(20)(1538-3598
(Electronic)):2220–6.
22.   Williams TN, Mwangi Tw Fau - Roberts DJ, Roberts Dj Fau - Alexander ND,
Alexander Nd Fau - Weatherall DJ, Weatherall Dj Fau - Wambua S, Wambua S Fau - Kortok
M, et al. An immune basis for malaria protection by the sickle cell trait. PLoS medicine.
2005;2(5); e128(1549-1676 (Electronic)).
23.   Williams TN, Mwangi Tw Fau - Wambua S, Wambua S Fau - Alexander ND,
Alexander Nd Fau - Kortok M, Kortok M Fau - Snow RW, Snow Rw Fau - Marsh K, et al.
Sickle cell trait and the risk of Plasmodium falciparum malaria and other childhood diseases.
The Journal of infectious diseases. 2005;192(1)(0022-1899 (Print)):178–86.
24.   Gong L, Parikh S, Rosenthal PJ, Greenhouse B. Biochemical and immunological
mechanisms by which sickle cell trait protects against malaria. Malaria Journal.
2013;12(1):317.
25.   Flage B, Dent M, Tejero J, Amoah L, Ofori-Acquah S. Heme promotes sexual
conversion of Plasmodium falciparum in human erythrocytes. Frontiers in Malaria. 2023;1.
26.   Vasquez M, Zuniga M, Rodriguez A. Oxidative Stress and Pathogenesis in Malaria.
Frontiers in cellular and infection microbiology. 2021;11(2235-2988 (Electronic)):768182.
27.   Cyrklaff M, Sanchez CP, Kilian N, Bisseye C, Simpore J, Frischknecht F, et al.
Hemoglobins S and C interfere with actin remodeling in Plasmodium falciparum-infected
erythrocytes. Science. 2011;334(6060):1283-6.
28.   Cholera R, Brittain Nj Fau - Gillrie MR, Gillrie Mr Fau - Lopera-Mesa TM, Lopera-
Mesa Tm Fau - Diakité SAS, Diakité Sa Fau - Arie T, Arie T Fau - Krause MA, et al. Impaired
cytoadherence of Plasmodium falciparum-infected erythrocytes containing sickle hemoglobin.
Proceedings of the National Academy of Sciences of the United States of America.
2008;105(3)(1091-6490 (Electronic)):991–6.
29.   Archer NM, Petersen N, Clark MA, Buckee CO, Childs LM, Duraisingh MT.
Resistance to Plasmodium falciparum in sickle cell trait erythrocytes is driven by oxygen-
dependent growth inhibition. Proceedings of the National Academy of Sciences.
2018;115(28):7350-5.
30.   Gouagna LC, Bancone G, Yao F, Yameogo B, Dabire KR, Costantini C, et al. Genetic
variation in human HBB is associated with Plasmodium falciparum transmission. Nature
genetics. 2010;42(4):328-31.

31.   Andolina C, Ramjith J, Rek J, Lanke K, Okoth J, Grignard L, et al. Plasmodium falciparum gametocyte carriage in longitudinally monitored incident infections is associated with duration of infection and human host factors. Scientific Reports. 2023;13(1):7072.

32.   Lawaly YR, Sakuntabhai A, Marrama L, Konate L, Phimpraphi W, Sokhna C, et al. Heritability of the human infectious reservoir of malaria parasites. PloS one. 2010;5(6):e11358-e.

33.   Ngou CM, Bayibéki AN, Abate L, Makinde OS, Feufack-Donfack LB, Sarah-Matio EM, et al. Influence of the sickle cell trait on Plasmodium falciparum infectivity from naturally infected gametocyte carriers. BMC Infectious Diseases. 2023;23(1):317.

34.   Gonçalves BP, Sagara I, Coulibaly M, Wu Y, Assadou MH, Guindo A, et al. Hemoglobin variants shape the distribution of malaria parasites in human populations and their transmission potential. Scientific reports. 2017;7(1):14267-.

35.   Meibalan EM, M. Biology of Malaria Transmission. Cold Spring Harb Perspect Med. 2017;7(3).

36.   Bousema T, Drakeley C. Epidemiology and infectivity of Plasmodium falciparum and Plasmodium vivax gametocytes in relation to malaria control and elimination. Clin Microbiol Rev. 2011;24(2):377-410.

37.   Ngotho P, Soares AB, Hentzschel F, Achcar F, Bertuccini L, Marti M. Revisiting gametocyte biology in malaria parasites. FEMS Microbiology Reviews. 2019;43(4):401-14.

38.   Llorà-Batlle O, Michel-Todó L, Witmer K, Toda H, Fernández-Becerra C, Baum J, et al. Conditional expression of PfAP2-G for controlled massive sexual conversion in Plasmodium falciparum. Science Advances. 2020;6(24):eaaz5057.

39.   Bancells C, Llora-Batlle O, Poran A, Notzel C, Rovira-Graells N, Elemento O, et al. Revisiting the initial steps of sexual development in the malaria parasite Plasmodium falciparum. Nature microbiology. 2019;4(1):144-54.

40.   Farid R, Dixon MW, Tilley L, McCarthy JS. Initiation of gametocytogenesis at very low parasite density in Plasmodium falciparum infection. The Journal of infectious diseases. 2017;215(7):1167-74.

41.   Usui M, Prajapati SK, Ayanful-Torgby R, Acquah FK, Cudjoe E, Kakaney C, et al. Plasmodium falciparum sexual differentiation in malaria patients is associated with host factors and GDV1-dependent genes. Nature Communications. 2019;10(1):2140.

42.   Pelle KG, Oh K, Buchholz K, Narasimhan V, Joice R, Milner DA, et al. Transcriptional profiling defines dynamics of parasite tissue sequestration during malaria infection. Genome Medicine. 2015;7(1):19.

43.   Prajapati SK, Ayanful-Torgby R, Pava Z, Barbeau MC, Acquah FK, Cudjoe E, et al. The transcriptome of circulating sexually committed Plasmodium falciparum ring stage parasites forecasts malaria transmission potential. Nature Communications. 2020;11(1):6159.

44.   Venugopal KA-O, Hentzschel F, Valkiūnas G, Marti MA-O. Plasmodium asexual growth and sexual development in the haematopoietic niche of the host. Nature reviews Microbiology. 2020;18((3)):177–89.

45.   Churcher TS, Bousema T Fau - Walker M, Walker M Fau - Drakeley C, Drakeley C Fau - Schneider P, Schneider P Fau - Ouédraogo AL, Ouédraogo Al Fau - Basáñez M-G, et al. Predicting mosquito infection from Plasmodium falciparum gametocyte density and estimating the reservoir of infection. eLife. 2013;2(2050-084X (Print)):e00626.

46.   Bousema T, Drakeley C. Determinants of Malaria Transmission at the Population Level. . Cold Spring Harbor perspectives in medicine. 2017;7((12)):a025510.

47.   Bradley JA-O, Stone W, Da DF, Morlais I, Dicko A, Cohuet A, et al. Predicting the likelihood and intensity of mosquito infection from sex specific Plasmodium falciparum gametocyte density. LID - 10.7554/eLife.34463 [doi] LID - e34463. (2050-084X (Electronic)).

48. Barry A, Bradley J, Stone W, Guelbeogo MW, Lanke K, Ouedraogo A, et al. Higher
gametocyte production and mosquito infectivity in chronic compared to incident Plasmodium
falciparum infections. Nature Communications. 2021;12(1):2443.

49. Barry A. Mosquitoes infecting children & children infecting mosquitoes in Burkina
Faso: Production and infectivity of malaria transmission stages: Radboud Universiteit
Nijmegen; 2022.

50. Ahmad A, Soumare HM, Camara MM, Jadama L, Gaye PM, Bittaye H, et al. Infectivity
of patent Plasmodium falciparum gametocyte carriers to mosquitoes: establishing capacity to
investigate the infectious reservoir of malaria in a low-transmission setting in The Gambia.
Transactions of the Royal Society of Tropical Medicine and Hygiene. 2021;115((12)):1462–7.

51. Josling GA, Williamson KC, Llinás M. Regulation of Sexual Commitment and
Gametocytogenesis in Malaria Parasites. Annual review of microbiology. 2018;72(1545-3251
(Electronic)):501–19.

52. Portugaliza HP, Miyazaki S, Geurten FJA, Pell C, Rosanas-Urgell A, Janse CJ, et al.
Artemisinin exposure at the ring or trophozoite stage impacts Plasmodium falciparum sexual
conversion differently. eLife. 2020;9:e60058.

53. Portugaliza HP, Natama HM, Guetens P, Rovira-Vallbona E, Somé AM, Millogo A, et
al. Plasmodium falciparum sexual conversion rates can be affected by artemisinin-based
treatment in naturally infected malaria patients. EBioMedicine. 2022;83(2352-3964
(Electronic)):104198.

54. Thommen BT, Passecker A, Buser T, Hitz E, Voss TS, Brancucci NMB. Revisiting the
Effect of Pharmaceuticals on Transmission Stage Formation in the Malaria Parasite
Plasmodium falciparum. Frontiers in cellular and infection microbiology. 2022;12(2235-2988
(Electronic)):802341.

55. Brancucci NMB, Gerdt JP, Wang C, De Niz M, Philip N, Adapa SR, et al.
Lysophosphatidylcholine Regulates Sexual Stage Differentiation in the Human Malaria
Parasite Plasmodium falciparum. Cell. 2017;171(7):1532-44.e15.

56. Neveu G, Beri D, Kafsack BF. Metabolic regulation of sexual commitment in
Plasmodium falciparum. Current opinion in microbiology. 2020;58(1879-0364
(Electronic)):93–8.

57. Harris CT, Tong XA-O, Campelo RA-O, Marreiros IA-O, Vanheer LA-OX, Nahiyaan
N, et al. Sexual differentiation in human malaria parasites is regulated by competition between
phospholipid metabolism and histone methylation. Nature microbiology. 2023;8(7)(2058-5276
(Electronic)):1280–92.

58. Trager W, Gill GS, Lawrence C, Nagel RL. Plasmodium falciparum: enhanced
gametocyte formation in vitro in reticulocyte-rich blood. Experimental parasitology.
1999;91(2):115-8.

59. Birget PLG, Repton C, O'Donnell AJ, Schneider P, Reece SE. Phenotypic plasticity in
reproductive effort: malaria parasites respond to resource availability. Proceedings Biological
sciences. 2017;284(1860).

60. Carter R Fau - Miller LH, Miller LH. Evidence for environmental modulation of
gametocytogenesis in Plasmodium falciparum in continuous culture. Bulletin of the World
Health Organization. 1979;57 Suppl 1(Suppl)(0042-9686 (Print)):37–52.

61. Pollitt LC, Mideo N, Drew DR, Schneider P, Colegrave N, Reece SE. Competition and
the evolution of reproductive restraint in malaria parasites. The American naturalist.
2011;177(3):358-67.

62. Carter LM, Kafsack BF, Llinas M, Mideo N, Pollitt LC, Reece SE. Stress and sex in
malaria parasites: Why does commitment vary? Evol Med Public Health. 2013;2013(1):135-
47.

63. Schneider P, Greischar MA, Birget PLG, Repton C, Mideo N, Reece SE. Adaptive
plasticity in the gametocyte conversion rate of malaria parasites. PLOS Pathogens.
2018;14(11):e1007371.
64. Kaur J, Hora R. '2TM proteins': an antigenically diverse superfamily with variable
functions and export pathways. PeerJ. 2018;6:e4757.
65. Chen S-Y, Wang Y, Telen MJ, Chi J-T. The Genomic Analysis of Erythrocyte
microRNA Expression in Sickle Cell Diseases. PloS one. 2008;3(6):e2360.
66. Fairhurst RM, Baruch Di Fau - Brittain NJ, Brittain Nj Fau - Ostera GR, Ostera Gr Fau
- Wallach JS, Wallach Js Fau - Hoang HL, Hoang Hl Fau - Hayton K, et al. Abnormal display
of PfEMP-1 on erythrocytes carrying haemoglobin C may protect against malaria. Nature.
2005;435(7045)(1476-4687 (Electronic)):1117–21.
67. Gong L, Maiteki-Sebuguzi C Fau - Rosenthal PJ, Rosenthal Pj Fau - Hubbard AE,
Hubbard Ae Fau - Drakeley CJ, Drakeley Cj Fau - Dorsey G, Dorsey G Fau - Greenhouse B,
et al. Evidence for both innate and acquired mechanisms of protection from Plasmodium
falciparum in children with sickle cell trait. Blood. 2012;119(16)(1528-0020
(Electronic)):3808–14.
68. Lelliott PM, McMorran BJ, Foote SJ, Burgio G. The influence of host genetics on
erythrocytes and malaria infection: is there therapeutic potential? Malaria journal.
2015;14(1475-2875 (Electronic)):289.
69. Fairhurst RM, Bess CD, Krause MA. Abnormal PfEMP1/knob display on Plasmodium
falciparum-infected erythrocytes containing hemoglobin variants: fresh insights into malaria
pathogenesis and protection. Microbes and infection. 2012;14(10):851-62.
70. Marsh K, Otoo L Fau - Hayes RJ, Hayes Rj Fau - Carson DC, Carson Dc Fau -
Greenwood BM, Greenwood BM. Antibodies to blood stage antigens of Plasmodium
falciparum in rural Gambians and their relation to protection against infection. Transactions of
the Royal Society of Tropical Medicine and Hygiene. 1989;83(3)(0035-9203 (Print)):293–303.
71. Verra F, Simpore J, Warimwe GM, Tetteh KK, Howard T, Osier FHA, et al.
Haemoglobin C and S Role in Acquired Immunity against Plasmodium falciparum Malaria.
PloS one. 2007;2(10):e978.
72. Diatta AM, Marrama, L., Tall, A., Trape, J. F., Dieye, A., Garraud, O., Mercereau-
Puijalon, O., & Perraut, R. Relationship of binding of immunoglobulin G to Plasmodium
falciparum-infected erythrocytes with parasite endemicity and antibody responses to conserved
antigen in immune individuals. Clinical and diagnostic laboratory immunology.
2004;11(1)(1071-412X (Print)):6–11.
73. Cabrera G, Cot M Fau - Migot-Nabias F, Migot-Nabias F Fau - Kremsner PG, Kremsner
Pg Fau - Deloron P, Deloron P Fau - Luty AJF, Luty AJ. The sickle cell trait is associated with
enhanced immunoglobulin G antibody responses to Plasmodium falciparum variant surface
antigens. The Journal of infectious diseases. 2005;191(10)(0022-1899 (Print)):1631–8.
74. Rabinovich RN, Drakeley C, Djimde AA, Hall BF, Hay SI, Hemingway J, et al.
malERA: An updated research agenda for malaria elimination and eradication. PLoS medicine.
2017;14(11)(1549-1676 (Electronic)):e1002456.
75. Portugaliza HP, Llorà-Batlle O, Rosanas-Urgell A, Cortés A. Reporter lines based on
the gexp02 promoter enable early quantification of sexual conversion rates in the malaria
parasite Plasmodium falciparum. Scientific Reports. 2019;9(1):14595.
76. Moelans IIMD. Pfs16, a potential vaccine candidate against the human malaria parasite
Plasmodium falciparum (Dissertation). PhD Thesis University of Nijmegen. 1995:154 p.
77. Bruce MC, Carter Rn Fau - Nakamura K, Nakamura K Fau - Aikawa M, Aikawa M
Fau - Carter R, Carter R. Cellular location and temporal expression of the Plasmodium
falciparum sexual stage antigen Pfs16. Molecular and biochemical parasitology.
1994;65(1)(0166-6851 (Print)):11–22.

78.   Shabani S, Serbus LR. Pfs16: A Key Parasitophorous Vacuole Membrane Protein
Crucial for Malaria Parasite Development and Transmission. The protein journal.
2025;44(2)(1875-8355 (Electronic)):133–46.
79.   Filarsky M, Fraschka SA, Niederwieser I, Brancucci NMB, Carrington E, Carrió E, et
al. GDV1 induces sexual commitment of malaria parasites by antagonizing HP1-dependent
gene silencing. Science (New York, NY). 2018;359(6381):1259-63.
80.   Boyle MJ, Chan JA, Handayuni I, Reiling L, Feng G, Hilton A, et al. IgM in human
immunity to Plasmodium falciparum malaria. Science Advances. 2019;5(9):eaax4489.
81.   Walker MR, Knudsen AS, Partey FD, Bassi MR, Frank AM, Castberg FA-O, et al.
Acquisition and decay of IgM and IgG responses to merozoite antigens after Plasmodium
falciparum malaria in Ghanaian children. PloS one. 2020;15((12)): e0243943.
82.   Stanisic DI, Fowkes FJ, Koinari M, Javati S, Lin E, Kiniboro B, et al. Acquisition of
antibodies against Plasmodium falciparum merozoites and malaria immunity in young children
and the influence of age, force of infection, and magnitude of response. Infection and
immunity. 2015;83((2)):646–60.
83.   Rogier E, Nace D, Dimbu PR, Wakeman B, Beeson JG, Drakeley C, et al. Antibody
dynamics in children with first or repeat Plasmodium falciparum infections. Frontiers in
medicine. 2022;9(2296-858X (Print)):869028.
84.   Akpogheneta OJ, Duah No Fau - Tetteh KKA, Tetteh Kk Fau - Dunyo S, Dunyo S Fau
- Lanar DE, Lanar De Fau - Pinder M, Pinder M Fau - Conway DJ, et al. Duration of naturally
acquired antibody responses to blood-stage Plasmodium falciparum is age dependent and
antigen specific. Infection and immunity. 2008;76((4)):1748–55. .
85.   Chelimo K, Ofulla Av Fau - Narum DL, Narum Dl Fau - Kazura JW, Kazura Jw Fau -
Lanar DE, Lanar De Fau - John CC, John CC. Antibodies to Plasmodium falciparum antigens
vary by age and antigen in children in a malaria-holoendemic area of Kenya. The Pediatric
infectious disease journal. 2005;24((8)):680–4.
86.   Amoah LE, Acquah FK, Nyarko PB, Cudjoe E, Donu D, Ayanful-Torgby R, et al.
Comparative analysis of asexual and sexual stage Plasmodium falciparum development in
different red blood cell types. Malaria Journal. 2020;19(1):200.
87.   LB. S, Freville A, Voss Till S, Baker David A, Awandare Gordon A, Conway David J.
Plasmodium falciparum Sexual Commitment Rate Variation among Clinical Isolates and
Diverse Laboratory-Adapted Lines. Microbiology Spectrum. 2022;10(6):e02234-22.
88.   Talman AM, Paul, R. E., Sokhna, C. S., Domarle, O., Ariey, F., Trape, J. F., & Robert,
V. Influence of chemotherapy on the Plasmodium gametocyte sex ratio of mice and humans.
The American journal of tropical medicine and hygiene. 2004;71(6):739-44.
89.   Uneke CJ. Plasmodium falciparum malaria and ABO blood group: is there any
relationship? Parasitology research. 2007;100((4)):759–65.
90.   Wolofsky KT, Ayi K Fau - Branch DR, Branch Dr Fau - Hult AK, Hult Ak Fau - Olsson
ML, Olsson Ml Fau - Liles WC, Liles Wc Fau - Cserti-Gazdewich CM, et al. ABO blood
groups influence macrophage-mediated phagocytosis of Plasmodium falciparum-infected
erythrocytes. PLoS pathogens. 2012;8((10)):e1002942.
91.   Afoakwah R, Aubyn E, Prah J, Nwaefuna EK, Boampong JN. Relative Susceptibilities
of ABO Blood Groups to Plasmodium falciparum Malaria in Ghana. Advances in hematology.
2016;2016(1687-9104 (Print)):5368793.
92.   Cserti-Gazdewich C, Musoke C, Dhabangi A, Nakiboneka-Ssenabulya D, Ddungu H,
Nabukeera-Barungi N, et al. ABO In Morbidity and Mortality of Plasmodium Falciparum
malaria. Blood. 2010;116(21):666-.
93.   Sheng K, Shariff M, Hebbel RP. Comparative Oxidation of Hemoglobins A and S.
Blood. 1998;91(9):3467-70.

94. Uzunova VV, Pan W Fau - Galkin O, Galkin O Fau - Vekilov PG, Vekilov PG. Free heme and the polymerization of sickle cell hemoglobin. Biophysical journal. 2010;99((6)):1976–85.

95. Fröhlich B, Jäger J, Lansche C, Sanchez CP, Cyrklaff M, Buchholz B, et al. Hemoglobin S and C affect biomechanical membrane properties of P. falciparum-infected erythrocytes. Communications biology. 2019;2(2399-3642 (Electronic)):311.

96. Smalley ME, Brown J. Plasmodium falciparum gametocytogenesis stimulated by lymphocytes and serum from infected Gambian children. Trans R Soc Trop Med Hyg. 1981;75(2):316-7.

97. Tan X, Traore B Fau - Kayentao K, Kayentao K Fau - Ongoiba A, Ongoiba A Fau - Doumbo S, Doumbo S Fau - Waisberg M, Waisberg M Fau - Doumbo OK, et al. Hemoglobin S and C heterozygosity enhances neither the magnitude nor breadth of antibody responses to a diverse array of Plasmodium falciparum antigens. The Journal of infectious diseases. 2011;204((11)):1750–61.

98. Bailey RD, Lawton JG, Niangaly A, Stucke EM, Bailey JA, Berry AA, et al. Children with hemoglobin C or S trait have low serologic responses to a subset of malaria variant surface antigens. The Journal of infection. 2024;89((4)):106257.

99. Asare KK, Agrah B, Ofori-Acquah FS, Kudzi W, Aryee NA, Amoah LE. Immune responses to P falciparum antibodies in symptomatic malaria patients with variant hemoglobin genotypes in Ghana. BMC Immunology. 2024;25(1):14.

100. Harp KO, Botchway F, Dei-Adomakoh Y, Wilson MD, Hood JL, Adjei AA, et al. Hemoglobin Genotypes Modulate Inflammatory Response to Plasmodium Infection. Frontiers in immunology. 2020;11(1664-3224 (Electronic)):593546.

101. Jensen AA-O, Adams YA-O, Hviid LA-O. Cerebral Plasmodium falciparum malaria: The role of PfEMP1 in its pathogenesis and immunity, and PfEMP1-based vaccines to prevent it. Immunological reviews. 2020;293((1)):230–52.

102. Dantzler KW, Ma S, Ngotho P, Stone WJR, Tao D, Rijpma S, et al. Naturally acquired immunity against immature Plasmodium falciparum gametocytes. Science translational medicine. 2019;11(495).

103. Ngotho P, Press KD, Peedell M, Muasya W, Omondi BR, Otoboh SE, et al. Reversible host cell surface remodelling limits immune recognition and maximizes transmission of Plasmodium falciparum gametocytes. bioRxiv. 2024:2024.04.30.591837.

104. Bousema T, Roeffen W, Meijerink H, Mwerinde H, Mwakalinga S, van Gemert G-J, et al. The Dynamics of Naturally Acquired Immune Responses to Plasmodium falciparum Sexual Stage Antigens Pfs230 & Pfs48/45 in a Low Endemic Area in Tanzania. PloS one. 2010;5(11):e14114.

105. Natama HM, Rovira-Vallbona E, Somé MA, Zango SH, Sorgho H, Guetens P, et al. Malaria incidence and prevalence during the first year of life in Nanoro, Burkina Faso: a birth-cohort study. Malaria Journal. 2018;17(1):163.

106. WHO. WHO. Basic malaria microscopy: part I. Learner's guide. . 2010;2nd ed.

107. Natama HA-OX, Rovira-Vallbona E, Somé MA, Zango SH, Sorgho H, Guetens P, et al. Malaria incidence and prevalence during the first year of life in Nanoro, Burkina Faso: a birth-cohort study. Malaria journal. 2018;17((1)):163.

108. Hofmann N, Mwingira F, Shekalaghe S, Robinson LJ, Mueller I, Felger I. Ultra-sensitive detection of Plasmodium falciparum by amplification of multi-copy subtelomeric targets. PLoS Med. 2015;12(3):e1001788.

109. Wittwer CT, Reed GH, Gundry CN, Vandersteen JG, Pryor RJ. High-Resolution Genotyping by Amplicon Melting Analysis Using LCGreen. Clinical chemistry. 2020;49(6):853-60.

110.   WHO Ma. Methods and techniques for clinical trials on antimalarial drug efficacy:
genotyping to identify parasite populations. Organization IcobMfMVacbtWH, editor.
Amsterdam, The Netherlands2008 2008.
111.   Rasouli Ghahfarokhi SM, Asadi F, Obeidi N. Use of capillary electrophoresis for
detection of hemoglobinopathies in individuals referred to health centers in Masjed-Soleiman.
Iranian Journal of Blood and Cancer. 2017;9:89-92.
112.   da Guarda CC, Yahouédéhou S, Santiago RP, Neres J, Fernandes CFL, Aleluia MM, et
al. Sickle cell disease: A distinction of two most frequent genotypes (HbSS and HbSC). PloS
one. 2020;15((1)):e0228399.
113.   Baker DA-O, Stewart LB, Large JM, Bowyer PW, Ansell KH, Jiménez-Díaz MB, et al.
A potent series targeting the malarial cGMP-dependent protein kinase clears infection and
blocks transmission. Nature communications. 2017;8((1)):430. .
114.   Ten Hove D, Jorgensen, T. D., & van der Ark, L. A. Updated guidelines on selecting
an intraclass correlation coefficient for interrater reliability, with applications to incomplete
observational designs. 2024;29:967-79.
115.   EVIMalaR MA. METHODS IN MALARIA RESEARCH 2013;Sixth Edition
116.   Silamut K, Phu Nh Fau - Whitty C, Whitty C Fau - Turner GD, Turner Gd Fau -
Louwrier K, Louwrier K Fau - Mai NT, Mai Nt Fau - Simpson JA, et al. A quantitative analysis
of the microvascular sequestration of malaria parasites in the human brain. The American
journal of pathology. 1999;155((2)):395–410.
117.   WHO. World malaria report 2024. 2024 11 December 2024.

1435