Transcranial Static Magnetic Stimulation Dissociates the Causal Roles of the Parietal Cortex in Spatial and Temporal Processing

Masakazu Sugimoto[a,b], Ikko Kimura[c], Masamichi J. Hayashi[a,b,*]

[a] Graduate School of Frontier Biosciences, The University of Osaka, Suita, Japan.

[b] Center for Information and Neural Networks (CiNet), Advanced ICT Research Institute, National Institute of Information and Communications Technology, Suita, Japan

[c] Danish Research Centre for Magnetic Resonance, Department of Radiology and Nuclear Medicine, Copenhagen University Hospital - Amager and Hvidovre, Copenhagen, Denmark

[*] Corresponding author

Center for Information and Neural Networks (CiNet), Advanced ICT Research Institute, National Institute of Information and Communications Technology

1-4 Yamadaoka, Suita, 565-0871, Japan

Email: mjhgml@gmail.com (Masamichi J. Hayashi).

Abstract

Accurate time estimation is essential for optimizing our perception and actions. Previous neuroimaging and transcranial magnetic stimulation (TMS) studies have suggested that the right inferior parietal lobule (IPL) and supplementary motor area (SMA) are involved in time perception. However, it remains inconclusive whether the activity in these regions is crucial for time perception, partly due to the possible spread of TMS effects across anatomically connected brain regions. Such a remote effect is less likely to happen with transcranial static magnetic stimulation (tSMS), as the static magnetic field is expected to modulate the firing threshold of neurons rather than directly triggering an action potential. In this study, we aimed to determine the causal relevance of local activities in the right IPL and the SMA for temporal processing using tSMS. 48 human volunteers (26 males and 22 females) participated in the study. We measured duration discrimination thresholds, along with orientation discrimination thresholds as a control, using staircase methods before and during the administration of tSMS over the IPL/SMA. Our results indicated no significant changes in duration discrimination thresholds in either the IPL or SMA conditions. In contrast, we observed an improvement in orientation discrimination thresholds in the IPL condition. This improvement correlated with individual differences in the distance between the scalp and the IPL. Overall, our findings demonstrate a causal involvement of the IPL in orientation processing. The correlation between the effects of tSMS and the scalp-to-target distance suggests that the efficacy of tSMS may be sensitive to the magnetic field strength.

Significance Statement

Accurate time estimation is essential for optimizing our perception and actions. While prior studies using transcranial magnetic stimulation (TMS) investigated a role of inferior parietal lobule (IPL) and supplementary motor area in spatiotemporal processing, the results have been inconclusive because TMS can affect anatomically connected areas. Here, using transcranial static magnetic stimulation (tSMS), we examined the causal relevance of local activities in these areas. Our findings showed that tSMS over the IPL did not alter duration discrimination but, unexpectedly, significantly enhanced orientation discrimination, with the degree of improvement correlating with individual anatomical differences. These results provide new insights into the neural basis of spatial and temporal processing and emphasize the potential of tSMS as a localized neuromodulation technique of interest.

Introduction

Precise estimation of time is essential for adapting to an ever-changing environment and optimizing our actions. For instance, in daily activities such as conversation and dancing to a beat, the ability to accurately estimate and predict the timing of external events enables smooth and effective interactions (Teki et al., 2011; Magyari et al., 2014, 2017; Ross et al., 2016).

Recent functional magnetic resonance imaging (fMRI) studies have shown that multiple brain regions are involved in temporal processing in the range of hundreds of milliseconds to about one second (Wiener et al., 2010b; Coull and Droit-Volet, 2018; Naghibi et al., 2024). A meta-analysis of neuroimaging studies reported that perceptual, subsecond timing tasks engage a distributed network of brain regions, including the supplementary motor area (SMA), inferior parietal lobule (IPL), inferior frontal gyrus (IFG), and basal ganglia (Wiener et al., 2010b). Among these areas, the right IPL and SMA have been highlighted as they seem to contain neural populations tuned for specific durations (Hayashi et al., 2015; Protopapa et al., 2019; Harvey et al., 2020). More recent research has associated activity in the right IPL with individual differences in time discrimination performance (Hayashi et al., 2018) and perceived duration (Hayashi and Ivry, 2020), suggesting its relevance to our subjective experience of time.

Consistent with these neuroimaging findings, several studies have reported that transcranial magnetic stimulation (TMS) applied over the right IPL impairs time discrimination (Bueti et al., 2008; Riemer et al., 2016) or biases perceived durations (Wiener et al., 2010a, 2012). However, it remains unclear whether the modulation of local activity in the area directly affects timing performances or operates through a remote effect, where TMS influences neuronal activity in anatomically connected regions (i.e., remote effect) (Rafique et al., 2015; Bergmann et al., 2021;

Wang et al., 2024). This is of particular concern given that a recent functional connectivity study

reported that the parietal cortex drives duration-selective activities in several brain regions,

including the SMA, visual cortex, IFG, and cerebellum (Protopapa et al., 2023). In line with this

finding, a TMS-EEG study reported that event-related potentials in frontocentral electrodes

around the SMA were modulated by TMS applied over the right IPL and correlated with changes

in timing performance (Wiener et al., 2012). These studies suggest that, even if TMS produces

measurable behavioral effects, it may be premature to attribute these effects to changes in local

activity beneath the TMS coil.

A potentially useful neuromodulation technique for determining the causal role of specific

brain regions is transcranial static magnetic stimulation (tSMS) (Oliviero et al., 2011). tSMS is a

non-invasive brain stimulation method that modulates the firing threshold of neurons in the cortex

by placing a neodymium magnet on the scalp (Sinha et al., 2024). Although the precise

mechanism is not fully understood, it is believed that a static magnetic field of approximately

120–200 mT affects sodium and chloride ion channels, possibly by deforming the lipids in the

cell membrane (Rosen, 2003; Hernando et al., 2020; Sinha et al., 2024). These alternations in ion

influx may lead to inhibitory effects on neural activity within the area exposed to the static

magnetic field. Since tSMS does not trigger action potential of neurons, it may offer advantages

over TMS in producing effects that are relatively localized and with minimal remote influences

on anatomically connected brain regions.

To determine whether the local activities of IPL are crucial for temporal processing, here, we

examined the influence of tSMS over the region on the duration discrimination task. If the area

contains duration-tuned neuronal populations that are crucial for subsecond timing, we predicted

that suppressing the IPL through tSMS should alter duration discrimination performance. In

addition to targeting the IPL, we included stimulation over the SMA as an exploratory feasibility

test for the application of tSMS. This decision was motivated by prior timing studies reporting

limited effects of TMS over the SMA (Del Olmo et al., 2007; F. Giovannelli et al., 2014; Méndez

et al., 2017), potentially because timing-related responses are localized to deeper, medial portions

of the SMA, as demonstrated by previous fMRI studies (Hayashi et al., 2018; Protopapa et al.,

2019). Additionally, we used an orientation discrimination task as a control because a previous

neuroimaging study found that while duration information was decoded from the IPL and, albeit

less clearly, the SMA, orientation information was not evident in these areas (Hayashi et al.,

2018).

Materials and Methods

Participants

A total of 48 healthy, right-handed adult volunteers participated in our experiments (22 females and 26 males; age = 22.1 (mean) ± 1.4 (SD) years). The sample size of 48 participants was determined a priori. Detecting a moderate correlation with adequate statistical power requires a minimum of 46 participants (Bujang and Baharum, 2016). Because our experimental design required the total number of participants to be a multiple of four to allow for counterbalancing, we set the final sample size to 48. This sample size is comparable to, and in some cases larger than, those used in previous tSMS studies targeting the SMA (Pineda-Pardo et al., 2019; Guida et al., 2023).

Participants were recruited through the institutional online recruitment system from July 15, 2021, to April 14, 2023. All participants had normal or corrected-to-normal vision and no history of neurological and psychiatric disorders. This study was conducted in accordance with the Declaration of Helsinki. Participants received detailed written explanations of the experimental procedures and safety measures, and they provided informed consent. The protocol was approved by the institutional ethics and safety committees.

Task and stimuli

Visual stimuli were generated using MATLAB (MathWorks, Natick, MA, USA) and presented on a gamma-corrected CRT monitor (Multiscan G520; Sony Inc., Tokyo, Japan) with a refresh rate of 100 Hz. In each trial, a standard stimulus and a comparison stimulus were sequentially presented at the center of the screen, with an inter-stimulus interval (ISI) of 1.0 s (Fig. 1).

Following the two visual stimuli, a red fixation point was presented to indicate the response period.

The red fixation point remained on the screen until the participant responded. The subsequent trial

began immediately after the response. Participants were instructed to maintain their gaze on the

fixation point throughout the experiments.

In the duration task, the standard and comparison stimuli were black disks (diameter 5.4

degrees), and participants judged whether the second stimulus was shorter or longer than the first.

In the orientation task, the standard and comparison stimuli were sinusoidal Gabor patches (100%

contrast, spatial frequency of 1.6 cycles/degree, Gaussian envelope SD of 2.0 degrees, diameter

~12 degrees). Participants judged whether the second stimulus was rotated clockwise or

counterclockwise relative to the first.

A one-up-three-down staircase procedure was employed to measure participants' thresholds in

both the duration and orientation discrimination tasks. In the duration discrimination task, the

duration of the standard stimulus was set to 0.5 s, while the duration of the comparison stimulus

was 0.5 s + T (initial T = 0.4 s). In the orientation discrimination task, the standard orientation of

the sinusoidal Gabor patches was 45° from vertical while that for the comparison was 45° + T

(initial T = 12°). The stimulus duration for each Gabor patch was 0.06 s. The order of presentation

for the standard and comparison stimuli was randomized across trials.

Until the first three reversals, T was multiplied by 1.5 (increased) when the response was

incorrect and divided by 1.5 (decreased) when the response was correct for both tasks. After

obtaining three reversal points, T was multiplied by 1.3 when the response was incorrect or

divided by 1.3 when the response was correct for three consecutive trials. To mitigate any

potential confounds related to experiment duration (e.g., fatigue), the measurement durations

were fixed at 270 s for the duration task and 215 s for the orientation task based on pilot

experiments. Participants responded by pressing either the "J" key (for shorter or

counterclockwise) or the "K" key (for longer or clockwise) with their right index and middle

fingers, respectively. Participants were explicitly instructed to respond as accurately as possible

and not to count time during the duration task.

Fig. 1. Stimulus sequence. In the duration task (left), two visual stimuli (black disks) were

presented sequentially. Participants judged whether the duration of the second disk was shorter or

longer than that of the first. In the orientation task (right), two Gabor patches were presented

sequentially. Participants judged whether the orientation of the second stimulus was rotated

clockwise or counterclockwise compared to the first. A response was required in every trial during

the response period, indicated by a red fixation point.

Target localization for tSMS

We employed a tSMS, a cylindrical neodymium magnet, with a diameter of 60 mm and a

thickness of 30 mm (MAG 60 r+, Neurek SL., Toledo, Spain). The magnetic field strength was

0.511 T at the surface of the magnet. The right IPL (MNI coordinates x, y, z = 66, -34, 34) and

SMA (x, y, z = 2, 0, 70) were selected as the stimulation sites based on preceding fMRI studies

(Hayashi et al., 2018; Hayashi and Ivry, 2020). To determine the individual's positions of the

stimulation sites, we collected T1-weighted images for each participant with a magnetization-

prepared rapid acquisition with a gradient echo (MPRAGE) sequence using a 3T MRI scanner

(Siemens Inc., Erlangen, Germany), MAGNETOM Prisma (0.9 mm isotropic resolution, TR =

2700 ms, and TE = 3.68 ms) or MAGNETOM Prisma Fit (0.8 mm isotropic resolution, TR =

2500 ms, and TE = 2.18 ms). For each participant, we localized the right IPL and SMA by

converting the MNI coordinates into positions within that individual's structural T1-weighted

image. Using statistical parametric mapping (SPM12) software implemented in MATLAB, we

calculated an affine transformation matrix to align the individual's T1-weighted image with the

standard brain (MNI template). Subsequently, we used the inverse of the transformation on the

neuronavigation system (Brainsight, Rogue Research Inc., Montreal, Quebec, Canada) to

determine the positions of the right IPL and SMA within the participant's structural image.

Experimental Procedure

We conducted two sessions of experiments on different days: one targeting the IPL and the

other targeting the SMA. The interval between sessions ranged from 1 to 27 days. Before each

session, we aligned the MR images with the participant's head and marked the scalp position for

placing the magnet using the Brainsight navigation system. Participants completed 10 practice

trials for both the duration and orientation discrimination tasks.

Experiments consisted of baseline and test blocks, with each block lasting approximately 20

min. In these blocks, participants performed duration and orientation discrimination tasks two

times each (Fig. 2A). Following baseline threshold measurements, which were taken without

applying tSMS, we fixed the magnet over the targeted region using a custom-assembled clamp

with a sliding arm (Slick Co., Ltd., Tokyo, Japan), ensuring that the center of the magnet's bottom

surface aligned with the marked position on the scalp. A chin rest was used to stabilize the

participant's head position. Based on a previous report showing suppression of cortical excitability

following administration of tSMS for 30 min (Dileone et al., 2018; Takamatsu et al., 2021), we

positioned tSMS over the IPL or SMA for 30 min prior to the test block and maintained the

position throughout the test block (Fig. 2B).

Participants were randomly assigned to one of two groups: one group received tSMS over the

IPL in the first session and over the SMA in the second, while the other group was tested in the

opposite order. The task order for the baseline and test blocks was counterbalanced within each

group, with half of the participants performing the duration task followed by the orientation task,

and the other half performing the tasks in reverse order. For each participant, the task order

remained consistent across both sessions.

Fig. 2. Experimental procedure and scalp-to-target distance. (A) Timeline of the experiment.
During both the baseline and test blocks, participants performed duration and orientation tasks,
each repeated two times. tSMS was applied for approximately 50 min, beginning 30 min before
and continuing throughout the 20 min test block. (B) Target regions. We identified the target
regions, right IPL and SMA, based on MNI coordinates from previous studies. The figure

illustrates the target regions on coronal (left) and sagittal (right) views of a standard brain image.

(C) The average distance from the scalp to the IPL and the SMA across participants. The red bar

represents the mean distance to the IPL, while the blue bar represents the mean distance to the

SMA. Error bars indicate the standard error. Gray lines represent individual data. *** $p < 0.001$

Data Analysis

To determine the discrimination thresholds with the staircase procedure, we averaged the last

six reversal points from the collected data (Hayashi et al., 2014). The threshold values were then

averaged between two measurements to establish an individual's discrimination threshold. If the

number of reversal points in one of the two measurements was less than eight, we excluded the

data and used the threshold from the other measurement as the participant's discrimination

threshold.

To assess any potential changes in the baseline threshold across sessions, we compared the

average baseline threshold values between the first and second sessions using two-tailed paired t-

tests ($\alpha = 0.05$), with adjustments for multiple comparisons using the Bonferroni correction. Also,

to evaluate the stability of individual differences in baseline discrimination thresholds, we

calculated Pearson correlation coefficients between the baseline thresholds obtained in the first

and second sessions. To investigate the changes in discrimination thresholds induced by tSMS,

we performed two-tailed one-sample t-tests ($\alpha = 0.05$) on the difference between the test and

baseline thresholds, applying the Bonferroni correction for multiple comparisons across two

target brain regions.

Finally, we examined whether the magnitude of the tSMS effects depends on the distance

between the targeted cortical area and the magnet. We calculated the Pearson correlation

coefficient between the distance from the scalp to the stimulation site (either the IPL or SMA)

and the change in discrimination threshold between the test and baseline blocks. The distance was

measured using the ruler tool in the Brainsight navigation system. We tested the significance of

correlations (two-tailed, $\alpha = 0.05$). Differences in correlation coefficients between tasks were also

tested by bootstrap resampling (10,000 samples) using the Robust Correlation Toolbox

implemented in MATLAB (Pernet et al., 2013).

All statistical analyses, except for the bootstrap resampling, were performed using JASP

(https://jasp-stats.org/). Bayes factors (BF) were calculated for all tests to quantify the relative

evidence supporting the null versus the alternative hypothesis. All Bayesian analyses were also

performed in JASP.

Results

Reliability of the baseline threshold

Among the collected data from the 48 participants, threshold data from eleven measurements in the IPL sessions (four baseline and three test threshold measurements for the duration task, and one baseline and three test thresholds measurements for the orientation task) and six measurements in the SMA sessions (three baseline and two test thresholds measurements for the duration task and one test threshold measurements for the orientation task) were disregarded due to the insufficient number of reversal points. For these blocks, the individual's baseline or test threshold was determined based on the threshold value obtained in the other measurement (see *Data analysis* section in the Materials and Methods).

Preceding to our main analyses, we examined whether the baseline thresholds differed between the first and second sessions. As expected, a paired t-test revealed that the baseline thresholds for duration discrimination were not statistically different ($t(47) = 0.258$, $p = 1.000$, Cohen's $d = 0.037$, BF $= 0.162$) (Fig. 3A, left). In contrast, baseline thresholds for orientation discrimination were significantly lower in the second session compared to the first session ($t(47) = 4.244$, $p < 0.001$, Cohen's $d = 0.613$, BF $= 226.832$) (Fig. 3B, left). This result suggests that, while baseline thresholds were maintained in the duration task, the threshold for the orientation task was improved as the experiment progressed.

We also tested whether the individual differences in baseline thresholds were maintained across sessions. Our correlation analyses revealed that individual differences between the first and the second baseline thresholds were correlated in both duration ($r = 0.437$, $p = 0.002$, BF $= 19.242$) (Fig. 3A, right) and orientation tasks ($r = 0.656$, $p < 0.001$, BF $= 43447.499$) (Fig. 3B,

right). These results indicate that, although the threshold was improved in the second session in

the orientation task, individual differences in the baseline thresholds were maintained across

sessions in both tasks.

Fig. 3. Baseline thresholds. Baseline thresholds for the first (Day 1) and second sessions (Day
2) for the duration task (A, left) and for the orientation task (B, left). Each bar represents the
average discrimination thresholds, with error bars indicating the standard error. The gray lines
indicate the change in individual thresholds between Day 1 and Day 2. ** $p < 0.01$, *** $p < 0.001$,
corrected for multiple comparison. Correlations between thresholds for Day 1 and Day 2 for the
duration task (A, right) and the orientation task (B, right). The black circles represent individual
data. The gray solid line represents the regression line. The black dashed line represents the
diagonal line. *** $p < 0.001$

Effect of tSMS on task performances

Next, we examined whether tSMS over the IPL and SMA altered discrimination thresholds in

duration and orientation tasks. A one-sample t-test for the duration discrimination threshold

revealed no significant difference from the baseline, both in the IPL and SMA conditions (IPL: t

$(47) = -0.716$, $p = 0.954$, Cohen's $d = -0.103$, BF $= 0.200$; SMA: t $(47) = 0.381$, $p = 1.000$, Cohen's

$d = 0.055$, BF $= 0.168$; Fig. 4A). In contrast, the orientation discrimination threshold significantly

decreased when tSMS was applied over the IPL (t $(47) = -2.463$, $p = 0.034$, Cohen's $d = -0.356$,

BF $= 2.371$; Fig. 4B). No such reduction in the orientation discrimination threshold was observed

in either task when tSMS was applied over the SMA (t (47) = -0.393, p = 1.000, Cohen's d = -

0.057, BF = 0.169; Fig. 4B).

Fig. 4. Modulation of discrimination thresholds by tSMS. Differences in the discrimination

threshold between baseline and test phases (Test - Baseline) in the duration (**A**) and the orientation

tasks (**B**) when tSMS was applied over the IPL (red bars) and SMA (blue bars). The gray circles

indicate individual data. Each bar depicts the mean threshold difference, with error bars indicating

the standard error. * p < 0.05, corrected for multiple comparisons.

Effect of magnetic field strength on performance improvement

Since the effective magnetic field strength of tSMS depends on the distance between the

magnet and target, we wondered whether the variability in the effect size of tSMS across

participants is explained by the scalp-to-target distance. First, we examined whether the mean

scalp-to-target distance differed between the two target regions. A paired t-test revealed that the

scalp-to-target distance for the IPL (16.1 ± 0.2 mm) was significantly closer than for the SMA

(24.0 ± 0.2 mm) (t (47) = -25.467, p < 0.001, Cohen's d = -3.676, BF = 7.253 × 10^{25}; Fig. 2C),

indicating that magnetic field strength in the IPL was greater and probably more effective than

that in the SMA.

Next, we tested whether individual differences in the scalp-to-target distance explain the effect

size of tSMS. Our correlation analyses revealed that individual differences in the scalp-to-IPL

distance were positively correlated with the change in orientation discrimination thresholds ($r =$

0.304, $p = 0.036$, BF $= 1.526$; Fig. 5B), whereas such a correlation was absent for the duration

task ($r = -0.029$, $p = 0.846$, BF $= 0.183$; Fig. 5A). A bootstrap analysis further confirmed that the

degree of correlations significantly differed between the two tasks (95% confidence interval (CI)

$= -0.607$–-0.006, $p = 0.046$; Fig. 5C). For the SMA, individual differences in the scalp-to-SMA

distance were not correlated with the degree of threshold modulation in either the duration ($r =$

0.022, $p = 0.883$, BF $= 0.182$; Fig. 5D) or orientation task ($r = 0.016$, $p = 0.917$, BF $= 0.181$; Fig.

5E). Our bootstrap analysis confirmed that the correlations not significantly differed between

these two tasks (95% CI $= -0.396$–0.456, $p = 0.941$; Fig. 5F). These findings together suggest that

the scalp-to-target distance in the IPL, which was shorter than that in the SMA, was predictive of

the degree of improvement in behavioral performances, specifically for the orientation task.

Fig. 5. Correlation between the scalp-to-target distance and the behavioral effect of tSMS.

Correlation between the scalp-to-target distance (**A and B** for IPL, **D and E** for SMA) and the

change in discrimination thresholds (**A and D** for duration task, **B and E** for orientation task).

Gray lines indicate regression lines and circles indicate individual data. Distribution of differences

in correlation coefficients between tasks (i.e., duration - orientation task) for IPL (C) and SMA

(F), estimated by the bootstrap method. The solid line represents the 95% confidence interval, and

the dashed line indicates the value of differences in correlation coefficients between tasks. * p <

0.05

Discussion

Discussion

This study investigated whether the local activities in the IPL and SMA are causally involved

in temporal and spatial processing by measuring duration and orientation discrimination

thresholds, respectively, before and during the application of tSMS. Our within-subject

experiments, which included a relatively large sample size (n = 48), showed no significant

changes in duration discrimination thresholds, irrespective of the target regions. Unexpectedly,

however, we found that orientation discrimination thresholds significantly improved when tSMS

was applied over the IPL (but not SMA). Additionally, the degree of improvement in the

orientation task correlated with individual differences in the scalp-to-target distance of the IPL.

Lack of tSMS effect on duration task

In our experiment, performance in the duration task was not significantly altered by the tSMS

applied over the IPL or SMA. This result may seem surprising given that numerous neuroimaging

studies have reported the involvement of these regions in temporal processing (Coull et al., 2015;

Hayashi et al., 2015, 2018; Protopapa et al., 2019; Hayashi and Ivry, 2020). The absence of a

tSMS effect in our study might be related to the distributed nature of temporal representations in

the brain. A lack of disorders associated with a selective impairment in timing abilities suggests

that time representations might be distributed across various regions (Wiener et al., 2011).

Supporting this idea, previous neuroimaging studies have consistently shown that timing

information is spread across cortical areas, forming a timing network (Hayashi et al., 2018;

Harvey et al., 2020; Cona et al., 2021, 2024; Protopapa et al., 2023). Therefore, we speculate that

disrupting one region of this timing network may not significantly affect timing performance

because the brain can flexibly utilize other reliable sources of timing information within the

network to maintain behavioral performance.

Although our study found no evidence of tSMS effect on time discrimination, previous studies

have reported impairments in timing performance when TMS was applied over the IPL (Bueti et

al., 2008; Wiener et al., 2010a, 2012). This discrepancy may be explained by the combination of

the distributed timing network idea described above and the difference in the degree of focality

between tSMS and TMS methods. TMS is known to influence neuronal activity not only in the

targeted areas but also in anatomically connected regions, which is referred to as the remote effect

(Sack et al., 2007; Hirose et al., 2011; Watanabe et al., 2015; Siebner et al., 2022). In contrast,

tSMS is believed to have a more focal effect due to its mechanism of working through the

deformation of lipids in the cell membrane (Rosen, 2003; Hernando et al., 2020). Therefore, the

varying degrees of focality between TMS and tSMS may account for different behavioral

outcomes. Specifically, while TMS can disrupt timing performance through broader interference

within the timing network, more localized disruption by tSMS might be compensated for by

utilizing timing information from intact areas.

The lack of tSMS effect on the SMA could be explained by the significant distance between

the SMA and the scalp. As demonstrated in the orientation discrimination task (Fig. 5B), the

impact of tSMS on behavioral performance is influenced by the scalp-to-target distance. The

targeted SMA was located on the medial wall of the brain, which may have rendered the magnetic

field too weak to effectively disrupt cortical activity. This notion is consistent with previous TMS

studies that reported no significant effect on time perception when TMS was administered over

the SMA (Del Olmo et al., 2007; Fabio Giovannelli et al., 2014; Méndez et al., 2017). Our findings

suggest a need for future studies to employ neuromodulation techniques that can effectively

stimulate deeper areas (e.g., transcranial focused ultrasound stimulation) to better clarify the role

of the SMA in duration discrimination tasks.

Although the relatively large scalp-to-target distance for the SMA suggests that the effective

magnetic field reaching the cortex may have been weaker, which may potentially attenuating the

impact of tSMS, previous studies have nevertheless demonstrated that tSMS applied over the

SMA can modulate behavioral and physiological measures in the motor domain (Pineda-Pardo et

al., 2019; Pagge et al., 2024). Importantly, these studies employed stimulation procedures broadly

comparable to ours in terms of magnet size, strength, and target definition, making it unlikely that

simple methodological differences alone account for the discrepancy in effectiveness across

studies. Instead, the impact of tSMS on the SMA may be task dependent, reflecting differences in

the neural networks recruited by the task. Specifically, prior work primarily examined motor

processes such as action planning and control, in which behavioral effects may arise through

modulation of interactions within the motor network, including SMA–M1 connectivity (Pineda-

Pardo et al., 2019). In contrast, duration discrimination is thought to rely on widely distributed

timing representations across a broader timing network (Protopapa et al., 2023). Consequently,

perturbation of a single node within this network may be compensated for by intact regions,

thereby reducing the likelihood that measurable behavioral effects emerge. Consistent with this

interpretation, TMS studies have repeatedly reported reliable effects of SMA stimulation on motor

control (Zandbelt et al., 2013; Obeso et al., 2017; Tosun et al., 2017; Berkay et al., 2018), whereas

stimulation of the SMA has often yielded null effects in time perception tasks (Del Olmo et al.,

2007; F. Giovannelli et al., 2014; Méndez et al., 2017). Taken together, these findings suggest that

the behavioral impact of SMA stimulation may depend on the nature of the targeted task and,

consequently, on the networks it engages.

tSMS over the right IPL improved orientation discrimination performance

Orientation discrimination performance was improved when tSMS was applied over the right

IPL. This finding aligns with previous studies showing that TMS over the right IPL significantly

increased reaction times for detecting target orientations (Sack et al., 2007; Sack, 2009),

suggesting that the IPL plays a crucial role in orientation discrimination.

The paradoxical enhancement in performance due to the inhibitory effects of tSMS can be

attributed to increased lateral inhibition mechanisms of orientation-tuned neural populations in

the IPL. In the visual cortex, higher GABA concentration levels have been linked to better

orientation thresholds (Edden et al., 2009), likely due to the enhanced lateral inhibition in

orientation-tuned neurons (Li et al., 2008; Brouwer and Heeger, 2011; Del Rosario et al., 2025).

Notably, a recent fMRI adaptation study highlighted the existence of spatiotopic receptive fields

for orientations in the IPL (Dunkley et al., 2016). Given that another study reported that tSMS

enhances inhibitory signal mediated by GABA receptors (Nojima et al., 2015), we speculate that

our tSMS application might have increased GABA concentration or sensitivity of GABA

receptors associated with orientation channels in the IPL. Consequently, this rise in GABA levels

could have enhanced orientation selectivity, leading to improved orientation discrimination

performance.

Additionally, we found that participants with shorter scalp-to-target distances showed greater

improvements in their orientation discrimination performance. This finding suggests that scalp-

to-target distance could be a useful metric for evaluating the effectiveness of tSMS. For other

neuromodulation techniques, such as TMS, simulating induced electric field in the target region

has been common (Axel Thielscher et al., 2015). Our findings imply that such simulations may

also be beneficial in predicting the effectiveness of tSMS. An important avenue for future research

with tSMS might be to control the dosage across participants based on these simulations, allowing

for an optimal selection of magnetic strength that standardizes the dose or effectiveness of tSMS

across different participants and target regions.

In the present study, the baseline block was conducted first, followed by the test block with the

magnet in place. One might argue that this fixed order could introduce a potential confound, as

task performance may improve over time due to learning effects. However, several lines of

evidence suggest that the improvement observed in our study cannot be fully explained by

learning alone. First, if practice were the primary driver of the observed improvement, similar

effects would be expected regardless of the stimulation site; however, no such improvement was

observed in the SMA condition. Second, by subtracting baseline thresholds within each session,

the influence of learning across sessions was minimized. Finally, individual differences in

threshold improvement observed in the IPL condition were correlated with scalp-to-IPL distance,

which cannot be accounted for by simple practice effects and instead supports the presence of a

physiological impact of tSMS. Taken together, these findings strengthen the interpretation that

the observed enhancement reflects a site-specific physiological effect of tSMS rather than a

learning-related effect.

Limitations

431  A limitation of the current study is that our experimental design, which employed the staircase

432 method, only assessed the modulation of threshold. Consequently, it remains unclear whether

433 tSMS impacted perceived duration and/or orientation (i.e., bias). Previous studies have suggested

434 that the right IPL is associated with the subjective experience of time (Wiener et al., 2010a, 2012;

435 Hayashi and Ivry, 2020), highlighting the necessity of measuring biases in perceived duration.

436 This can be clarified by using discrimination tasks that involve a stable reference unaffected by

437 tSMS, such as a bisection task with an internal reference.

438  Second, since tSMS was applied continuously during the discrimination tasks, the specific

439 processes being modulated (e.g., stimulus encoding, maintenance, comparison) in the orientation

440 task remain unclear. This lack of temporal specificity may not be easily addressed due to the

441 nature of tSMS, necessitating the use of more temporally specific stimulation methods, such as

442 focused ultrasound stimulation.

443  Finally, the absence of a sham condition represents an additional limitation of the present study.

444 The inclusion of a sham magnet condition would allow for more rigorous control of nonspecific

445 influences, including practice-related and placebo effects. Future studies incorporating such a

446 sham condition would help fully dissociate the specific physiological effects of tSMS from these

447 potential confounds.

449 ***Conclusions***

450 The current study aimed to identify the brain regions that are causally involved in temporal and

451 spatial perception using tSMS. The observed improvements in orientation discrimination

452 thresholds induced by tSMS suggest a potential causal involvement of the right IPL in spatial

processing. Furthermore, we demonstrated that individual differences in the distance from the

scalp to the right IPL correlated with the effectiveness of tSMS, underscoring the importance of

considering anatomical differences when predicting and evaluating the efficacy of tSMS.

CRediT authorship contribution statement

Masakazu Sugimoto: Methodology, Software, Validation, Formal analysis, Investigation, Data

Curation, Writing- Original draft preparation, Visualization. **Ikko Kimura:** Methodology,

Writing- Reviewing and Editing. **Masamichi J. Hayashi:** Conceptualization, Methodology,

Software, Writing- Reviewing and Editing, Supervision, Project administration, Funding

acquisition

Acknowledgments

This work was supported by the Japan Society for the Promotion of Science (Grants-in-Aid for

Scientific Research JP21H00315, JP22H01110, and JP23K17649 to M.J.H., and Grant-in-Aid for

JSPS Fellows JP25KJ1802 to M.S.) and the Japan Science and Technology Agency (FOREST

JPMJFR232X to M.J.H.).

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used ChatGPT, Grammarly, and DeepL in order

to improve the readability and language of the manuscript. After using these tools, the authors

reviewed and edited the content as needed and took full responsibility for the content of the

published article.

References

Axel Thielscher, Andre Antunes, Guilherme B Saturnino (2015) Field modeling for transcranial magnetic stimulation: A useful tool to understand the physiological effects of TMS? 2015 37th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC) 222–225.

Bergmann TO, Varatheeswaran R, Hanlon CA, Madsen KH, Thielscher A, Siebner HR (2021) Concurrent TMS-fMRI for causal network perturbation and proof of target engagement. Neuroimage 237:118093.

Berkay D, Eser HY, Sack AT, Çakmak YÖ, Balcı F (2018) The modulatory role of pre-SMA in speed-accuracy tradeoff: A bi-directional TMS study. Neuropsychologia 109:255–261.

Brouwer GJ, Heeger DJ (2011) Cross-orientation suppression in human visual cortex. J Neurophysiol 106:2108–2119.

Bueti D, Bahrami B, Walsh V (2008) Sensory and association cortex in time perception. J Cogn Neurosci 20:1054–1062.

Bujang MA, Baharum N (2016) Sample Size Guideline for Correlation Analysis. World Journal of Social Science Research 3:37.

Cona G, Wiener M, Allegrini F, Scarpazza C (2024) Gradient Organization of Space, Time, and Numbers in the Brain: A Meta-analysis of Neuroimaging Studies. Neuropsychol Rev 34:721–737.

Cona G, Wiener M, Scarpazza C (2021) From ATOM to GradiATOM: Cortical gradients support time and space processing as revealed by a meta-analysis of neuroimaging studies. Neuroimage 224:117407.

Coull JT, Charras P, Donadieu M, Droit-Volet S, Vidal F (2015) SMA selectively codes the active accumulation of temporal, not spatial, magnitude. J Cogn Neurosci 27:2281–2298.

Coull JT, Droit-Volet S (2018) Explicit Understanding of Duration Develops Implicitly through Action. Trends Cogn Sci 22:923–937.

Del Olmo MF, Cheeran B, Koch G, Rothwell JC (2007) Role of the cerebellum in externally paced rhythmic finger movements. J Neurophysiol 98:145–152.

Del Rosario J, Coletta S, Kim SH, Mobille Z, Peelman K, Williams B, Otsuki AJ, Del Castillo Valerio A, Worden K, Blanpain LT, Lovell L, Choi H, Haider B (2025) Lateral inhibition in V1 controls neural and perceptual contrast sensitivity. Nat Neurosci 28:836–847.

Dileone M, Mordillo-Mateos L, Oliviero A, Foffani G (2018) Long-lasting effects of transcranial

static magnetic field stimulation on motor cortex excitability. Brain Stimulation: Basic,
Translational, and Clinical Research in Neuromodulation 11:676–688.

Dunkley BT, Baltaretu B, Crawford JD (2016) Trans-saccadic interactions in human parietal and
occipital cortex during the retention and comparison of object orientation. Cortex 82:263–
276.

Edden RAE, Muthukumaraswamy SD, Freeman TCA, Singh KD (2009) Orientation
discrimination performance is predicted by GABA concentration and gamma oscillation
frequency in human primary visual cortex. Journal of Neuroscience 29:15721–15726.

Giovannelli Fabio, Innocenti I, Rossi S, Borgheresi A, Ragazzoni A, Zaccara G, Viggiano MP,
Cincotta M (2014) Role of the dorsal premotor cortex in rhythmic auditory-motor
entrainment: A perturbational approach by rTMS. Cerebral Cortex 24:1009–1016.

Giovannelli F., Ragazzoni A, Battista D, Tarantino V, Del Sordo E, Marzi T, Zaccara G, Avanzini
G, Viggiano MP, Cincotta M (2014) "..the times they aren't A-Changin''." rTMS does not
affect basic mechanisms of temporal discrimination: A pilot study with ERPs."
Neuroscience 278:302–312.

Guida P, Foffani G, Obeso I (2023) The Supplementary Motor Area and Automatic Cognitive
Control: Lack of Evidence from Two Neuromodulation Techniques. J Cogn Neurosci
35:439–451.

Harvey BM, Dumoulin SO, Fracasso A, Paul JM (2020) A Network of Topographic Maps in
Human Association Cortex Hierarchically Transforms Visual Timing-Selective Responses.
Current Biology 30:1424-1434.e6.

Hayashi MJ, Ditye T, Harada T, Hashiguchi M, Sadato N, Carlson S, Walsh V, Kanai R (2015)
Time Adaptation Shows Duration Selectivity in the Human Parietal Cortex. PLoS Biol
13:e1002262.

Hayashi MJ, Ivry RB (2020) Duration selectivity in right parietal cortex reflects the subjective
experience of time. Journal of Neuroscience 40:7749–7758.

Hayashi MJ, Kantele M, Walsh V, Carlson S, Kanai R (2014) Dissociable Neuroanatomical
Correlates of Subsecond and Suprasecond Time Perception. J Cogn Neurosci 26:1685–1693.

Hayashi MJ, van der Zwaag W, Bueti D, Kanai R (2018) Representations of time in human
frontoparietal cortex. Commun Biol 1:1–10.

Hernando A, Galvez F, García MA, Soto-León V, Alonso-Bonilla C, Aguilar J, Oliviero A (2020)
Effects of Moderate Static Magnetic Field on Neural Systems Is a Non-invasive Mechanical
Stimulation of the Brain Possible Theoretically? Front Neurosci 14:1–9.

Hirose M, Mochizuki H, Groiss SJ, Tanji Y, Nakamura K, Nakatani-Enomoto S, Enomoto H,
Nishizawa M, Ugawa Y (2011) On-line effects of quadripulse transcranial magnetic
stimulation (QPS) on the contralateral hemisphere studied with somatosensory evoked
potentials and near infrared spectroscopy. Exp Brain Res 214:577–586.

Li G, Yang Y, Liang Z, Xia J, Yang Y, Zhou Y (2008) GABA-mediated inhibition correlates with
orientation selectivity in primary visual cortex of cat. Neuroscience 155:914–922.

Magyari L, Bastiaansen MCM, de Ruiter JP, Levinson SC (2014) Early anticipation lies behind
the speed of response in conversation. J Cogn Neurosci 26:2530–2539.

Magyari L, De Ruiter JP, Levinson SC (2017) Temporal Preparation for Speaking in Question-
Answer Sequences. Front Psychol 8:211.

Méndez JC, Rocchi L, Jahanshahi M, Rothwell J, Merchant H (2017) Probing the timing network:
A continuous theta burst stimulation study of temporal categorization. Neuroscience
356:167–175.

Naghibi N, Jahangiri N, Khosrowabadi R, Eickhoff CR, Eickhoff SB, Coull JT, Tahmasian M
(2024) Embodying Time in the Brain: A Multi-Dimensional Neuroimaging Meta-Analysis
of 95 Duration Processing Studies. Neuropsychol Rev 34:277–298.

Nojima I, Koganemaru S, Fukuyama H, Mima T (2015) Static magnetic field can transiently alter
the human intracortical inhibitory system. Clinical Neurophysiology 126:2314–2319.

Obeso I, Wilkinson L, Teo JT, Talelli P, Rothwell JC, Jahanshahi M (2017) Theta burst magnetic
stimulation over the pre-supplementary motor area improves motor inhibition. Brain Stimul
10:944–951.

Oliviero A, Mordillo-Mateos L, Arias P, Panyavin I, Foffani G, Aguilar J (2011) Transcranial
static magnetic field stimulation of the human motor cortex. Journal of Physiology
589:4949–4958.

Pagge C, Caballero-Insaurriaga J, Oliviero A, Foffani G, Ammann C (2024) Transcranial static
magnetic field stimulation of the supplementary motor area decreases corticospinal
excitability in the motor cortex: a pilot study. Sci Rep 14.

Pernet CR, Wilcox R, Rousselet GA (2013) Robust Correlation Analyses: False Positive and
Power Validation Using a New Open Source Matlab Toolbox. Front Psychol 3:606.

Pineda-Pardo JA, Obeso I, Guida P, Dileone M, Strange BA, Obeso JA, Oliviero A, Foffani G
(2019) Static magnetic field stimulation of the supplementary motor area modulates resting-
state activity and motor behavior. Commun Biol 2:397.

Protopapa F, Hayashi MJ, Kulashekhar S, van der Zwaag W, Battistella G, Murray MM, Kanai R,

Bueti D (2019) Chronotopic maps in human supplementary motor area. PLoS Biol
574   17:e3000026.

575   Protopapa F, Kulashekhar S, Hayashi MJ, Kanai R, Bueti D (2023) Effective connectivity in a
576   duration selective cortico-cerebellar network. Sci Rep 13:20674.

577   Rafique SA, Solomon-Harris LM, Steeves JKE (2015) TMS to object cortex affects both object
578   and scene remote networks while TMS to scene cortex only affects scene networks.
579   Neuropsychologia 79:86–96.

580   Riemer M, Rhodes D, Wolbers T (2016) Systematic Underreproduction of Time Is Independent
581   of Judgment Certainty. Neural Plast 2016:1–8.

582   Rosen AD (2003) Mechanism of Action of Moderate-Intensity Static Magnetic Fields on
583   Biological Systems. Cell Biochem Biophys 39:163–174.

584   Ross JM, Iversen JR, Balasubramaniam R (2016) Motor simulation theories of musical beat
585   perception. Neurocase 22:558–565.

586   Sack AT (2009) Parietal cortex and spatial cognition. Behavioural Brain Research 202:153–161.

587   Sack AT, Kohler A, Bestmann S, Linden DEJ, Dechent P, Goebel R, Baudewig J (2007) Imaging
588   the brain activity changes underlying impaired visuospatial judgments: Simultaneous fMRI,
589   TMS, and behavioral studies. Cerebral Cortex 17:2841–2852.

590   Siebner HR et al. (2022) Transcranial magnetic stimulation of the brain: What is stimulated? – A
591   consensus and critical position paper. Clinical Neurophysiology 140:59–97.

592   Sinha A saran, Shibata S, Takamatsu Y, Akita T, Fukuda A, Mima T (2024) Static magnetic field
593   stimulation enhances shunting inhibition via a SLC26 family Cl − channel, inducing
594   intrinsic plasticity. The Journal of Neuroscience 44:e1324222024.

595   Takamatsu Y, Koganemaru S, Watanabe T, Shibata S, Yukawa Y, Minakuchi M, Shimomura R,
596   Mima T (2021) Transcranial static magnetic stimulation over the motor cortex can facilitate
597   the contralateral cortical excitability in human. Sci Rep 11:5370.

598   Teki S, Grube M, Kumar S, Griffiths TD (2011) Distinct neural substrates of duration-based and
599   beat-based auditory timing. Journal of Neuroscience 31:3805–3812.

600   Tosun T, Berkay D, Sack AT, Çakmak Y, Balcı F (2017) Inhibition of pre–supplementary motor
601   area by continuous theta burst stimulation leads to more cautious decision-making and more
602   efficient sensory evidence integration. J Cogn Neurosci 29:1433–1444.

603   Wang JB, Hassan U, Bruss JE, Oya H, Uitermarkt BD, Trapp NT, Gander PE, Howard MA, Keller
604   CJ, Boes AD (2024) Effects of transcranial magnetic stimulation on the human brain
605   recorded with intracranial electrocorticography. Mol Psychiatry 29:1228–1240.

Watanabe T, Hanajima R, Shirota Y, Tsutsumi R, Shimizu T, Hayashi T, Terao Y, Ugawa Y,
Katsura M, Kunimatsu A, Ohtomo K, Hirose S, Miyashita Y, Konishi S (2015) Effects of
rTMS of pre-supplementary motor area on fronto basal ganglia network activity during stop-
signal task. Journal of Neuroscience 35:4813–4823.

Wiener M, Hamilton R, Turkeltaub P, Matell MS, Coslett HB (2010a) Fast forward:
Supramarginal gyrus stimulation alters time measurement. J Cogn Neurosci 22:23–31.

Wiener M, Kliot D, Turkeltaub PE, Hamilton RH, Wolk DA, Coslett HB (2012) Parietal influence
on temporal encoding indexed by simultaneous transcranial magnetic stimulation and
electroencephalography. Journal of Neuroscience 32:12258–12267.

Wiener M, Matell MS, Coslett HB (2011) Multiple Mechanisms for Temporal Processing. Front
Integr Neurosci 5.

Wiener M, Turkeltaub P, Coslett HB (2010b) The image of time: A voxel-wise meta-analysis.
Neuroimage 49:1728–1740.

Zandbelt BB, Bloemendaal M, Hoogendam JM, Kahn RS, Vink M (2013) Transcranial Magnetic
Stimulation and Functional MRI Reveal Cortical and Subcortical Interactions during Stop-
signal Response Inhibition. J Cogn Neurosci 25:157–174.