Inhibition governs stimulus preference encoding in mPFC pyramidal neurons during a social choice

Renad Jabarin[1], Shai Netser[1] and Shlomo Wagner[1,*]

*Lead contact author: shlomow@research.haifa.ac.il

[1]Sagol Department of Neurobiology, Faculty of Natural Sciences, University of Haifa, Haifa, Israel

Summary

Social decision-making requires the brain to continuously evaluate competing options and select appropriate actions. The medial prefrontal cortex (mPFC) has been implicated in social cognition and value-based decision-making, yet the neural mechanisms by which it encodes the relative value of competing social choices remain poorly understood. Here, we employed fiber photometry, optogenetics, and projection-specific recordings in mice performing four distinct binary social discrimination tasks to reveal how mPFC pyramidal neurons encode social choice. We found that mPFC pyramidal neurons exhibit profound inhibition during bouts toward the preferred stimulus. This inhibitory response was specific to transitional bouts, when animals switched between stimuli, and was absent during non-transitional bouts, which occurred during continued engagement with the same stimulus. Notably, this differential activity encoded relative rather than absolute stimulus value, such that the same stimulus evoked opposite neural responses depending on its contextual preference. Accordingly, manipulating the subject's preference between food and social stimuli via food deprivation inverted its inhibition-excitation pattern. Furthermore, negative calcium transients in the mPFC predicted subsequent investigation of the preferred stimulus, suggesting a functional role in guiding behavioral choice. Optogenetic activation of mPFC pyramidal neurons during stimulus investigation produced immediate aversion yet paradoxically biased the stimulated animal to persistently re-engage with the stimulus-associated option through repetitive non-transitional bouts. Finally, projection-specific recordings revealed that mPFC neurons targeting the nucleus accumbens and basolateral amygdala were differentially recruited across tasks, reflecting functional specialization of output pathways. Together, these findings identify inhibition of mPFC pyramidal neurons as a key neural signature for encoding relative stimulus preference, thereby establishing a neural principle by which the mPFC dynamically guides social choice.

Introduction

Social interactions are fundamental to the survival and well-being of many species, including humans. These interactions are composed of a continuous stream of decisions that require an individual to evaluate available social partners, interpret their intentions, and select appropriate actions [1, 2]. Thus, social decision-making is a complex cognitive function that relies on the brain's ability to assign value to different social stimuli and to flexibly adapt behavior in response to a constantly changing social landscape [3]. A breakdown in this critical capacity can lead to profound social deficits, as seen in a variety of neuropsychiatric and neurodevelopmental disorders, most notably autism spectrum disorder (ASD) [4, 5].

The medial prefrontal cortex (mPFC) has emerged as a critical hub in the neural circuitry governing social cognition and value-based decision-making [6, 7]. Extensive research in both humans and animal models has identified the mPFC as a critical regulator of a wide range of social functions, including the recognition of conspecifics, the interpretation of social cues, and the regulation of the approach–avoidance balance [8-10] and the integration of emotional and contextual cues [11]. It has been proposed that the mPFC's function is to learn and represent associations between context and adaptive behavioral responses, a principle that may extend from simple reward-based tasks to the complex, dynamic world of social interaction [12, 13]. Understanding this process is crucial for deciphering how the brain dynamically navigates the complexities of the social world—a process that remains poorly understood

While it is well established that the mPFC encodes context-dependent value in non-social operant tasks involving explicit rewards or costs [14], it remains unclear whether similar principles govern the more fluid and spontaneous nature of social interactions. In natural settings, the 'value' of a social partner is not fixed but rather dynamically shaped by the presence of other individuals, the subject's internal state, and the broader social environment [15, 16]. This concept of relative valuation is a hallmark of flexible and adaptive decision-making. However, a central and unanswered question in social neuroscience is how the mPFC dynamically encodes the relative appeal of social choices, particularly during innate behavior and unconditioned social contexts.

At the circuit level, decision-making is often conceptualized as a competition between neuronal populations that is resolved through a balance of excitatory and inhibitory signaling, ultimately biasing activity in favor of one option over another [17-21]. The

mPFC is thought to implement this process through a 'differential coding' scheme, where

distinct patterns of neural activity represent different types of social information [10, 22].

Despite these advances, a significant gap remains in our understanding of how the

mPFC translates the relative value of social options into a concrete behavioral choice.

While we know that the mPFC is involved, we do not yet have a clear understanding of

the neural code that represents the decision itself, nor do we comprehend how this code

is utilized to guide the initiation, maintenance, and switching of social investigation [23,

24]. Specifically, it is still unknown how the mPFC dynamically encodes the relative

value of competing social and non-social options and how this evolving representation

translates into behavioral choice [25, 26].

Here, we use a battery of four distinct social discrimination tasks in mice, each requiring

a choice between two stimuli of varying motivational value, to show that the mPFC

encodes the relative value of social choices through selective inhibition: pyramidal

neurons are profoundly inhibited during transitional bouts toward the preferred stimulus,

distinguishing it from both the non-preferred stimulus and baseline chamber

investigation. This differential activity was observed specifically during transitional bouts

between stimuli, suggesting a role in the decision to engage with a new option.

Furthermore, we found that negative calcium transients in the mPFC were predictive of

subsequent investigation of the preferred stimulus. Furthermore, distinct mPFC

projections to the NAc and BLA were differentially engaged during social decisions,

indicating that valuation and emotional discrimination are mediated through segregated

downstream circuits. Finally, optogenetic activation of mPFC pyramidal neurons during

stimulus investigation, despite eliciting an immediate avoidance, was sufficient to bias

choice by increasing the preference for the stimulated option via repetitive bouts of

investigation. Together, these findings identify a neural principle underlying social choice

encoding in the mPFC and demonstrate how cortical circuits transform social information

into adaptive decisions.

Results

I. Behavioral characterization of social choice across multiple binary
discrimination tasks

To establish a behavioral framework for interpreting neural activity in the mPFC, we first
assessed how subject mice behave when faced with a binary choice between distinct
stimuli with varying motivational value. To that end, adult males were tested in four
different discrimination tasks that required them to choose between two distinct stimuli.
In the Social Preference (SP) task, subject mice were exposed to a novel age- and sex-
matched conspecific (social stimulus) on one side of the arena and an inanimate object
on the other side (Fig. 1A), and exhibited a higher preference to investigate the social
stimulus over the object, as reflected by their investigation time (Fig. 1B-C) and mean
investigation bout duration (Fig. 1D) for each of the stimuli. In the Sex Preference (SxP)
task, subjects preferred to investigate a female over a male conspecific (Fig. 1F-I). In the
Emotional State Preference for stress (ESPs) task, subjects preferred a stressed over a
naïve conspecific (Fig. 1K-N). Lastly, in the Social vs. Food (SvF) task, subjects
displayed a significant preference for a conspecific over food (Fig. 1P-S). While the four
tasks represent distinct social contexts, they all exhibit a similar level of stimulus
preference, as reflected by the relative discrimination index (RDI; Fig. S1A). Notably, all
four tasks shared one type of stimulus – a novel naïve male, which was the preferred
stimulus in the SP and SvF tasks, but was the non-preferred stimulus in the SxP and
ESPs tasks.

The four tasks also exhibited a similar level of social propensity, as reflected in the total
investigation time (Fig. S1B) and showed a similar distance traveled during each task
(Fig. S1C). Thus, they appear to induce a comparable level of social motivation and
arousal in the subject animals, and differ primarily in the type of choice, dictated by the
identity of the presented stimuli.

We then differentiate between two types of investigation bouts presumably differing in
the type of decision preceding them: transitional vs. non-transitional. Transitional bouts
occur when the subject animal investigates one stimulus following a previous bout with
the other stimulus. As we previously described [27], transitional bouts were more
abundant at the early stage of each of the four tasks, with no apparent correlation with
the investigation time (Fig. 1B,G,L,Q, grey line). They reflect the animal's decision to
stop investigating one stimulus and start investigating the other stimulus, thus
performing a behavioral shift. In contrast, non-transitional bouts occur when the subject

animal investigates one stimulus following a previous bout with the same stimulus,

hence reflecting the sustained tendency of the animal to pursue the same stimulus.

We found that in all four tasks, mice exhibited a greater number of non-transitional bouts

compared to transitional bouts (Fig. S1D). Moreover, we found a significantly higher

number of non-transitional bouts for the preferred stimulus over the non-preferred

stimulus in all tasks, except for the SvF, where only a trend was observed (Fig.

1E,J,O,T). These findings suggest that mice exhibit a consistent tendency to persist in

investigating the preferred stimulus once it has been selected, and that social decision-

making involves both sustained engagement and selective switching between options.

II. Activity of mPFC pyramidal neurons discriminates between transitional and

non-transitional bouts

To examine the activity of mPFC CaMKIIα-expressing pyramidal neurons during the four

social discrimination tasks, we employed fiber photometry, as previously described [28].

For that, we injected the mPFC with AAV viral vectors to express the genetically

encoded calcium indicator GCaMP6f under the control of the CaMKIIα promoter (Fig.

2A). We then recorded calcium signals from subject mice during the experiments shown

in Fig. 1, with each session comprising a 10-minute recording of calcium signals: five

minutes before the introduction of stimuli into stimulus chambers (pre-encounter phase)

and five minutes after it (encounter phase, see timeline in Fig. 2B). Example traces of

normalized calcium signal aligned with the onset of non-transitional and transitional

bouts performed during that session, as well as mean traces for each of the four tasks

are shown in Fig. S2A-H.

We used z-score Analysis to normalize and average all the instantaneous signals

recorded during the two types of behavioral events—namely, transitional and non-

transitional bouts—separately. We then analyzed the averaged z-scored calcium

response for each behavioral event across all sessions of the same task, as previously

described by us [28].

We first determined the responses of mPFC pyramidal neurons during transitional and

non-transitional bouts of investigation conducted toward the two empty stimulus

chambers during the pre-encounter phase of each session. No behavioral preference

was found for one empty chamber over the other in any of the four tasks (Fig. S3A-D).

Still, we found that both non-transitional bouts (in SP, SxP, and ESPs tasks; Fig. S3E-F)

and transitional bouts (in all four tasks; Fig. S3G-H) elicited an immediate increase in the

recorded signal (mainly during -1 to 2s relative to the beginning of the bout), with this

excitatory response being more robust in transitional compared to non-transitional bouts
(Independent t-test- -1to2s: t(45)=-5.35, p<0.001; 3-6s: t(45)=-1.572, p=0.117).
We then analyzed the responses of mPFC pyramidal neurons during the encounter
stage of the various sessions, when subject mice investigated stimulus-containing
chambers. Importantly, we compared these responses with the signals recorded during
transitional and non-transitional bouts toward the empty chambers during the pre-
encounter stage. We found that transitional bouts elicited differential neural responses
between the two stimuli, with the preferred stimulus eliciting a significant inhibition while
the non-preferred one eliciting excitation. Nevertheless, there was no difference between
the excitatory response toward the non-preferred stimulus and the response toward the
empty cages during the pre-encounter stage, while the inhibitory response toward the
preferred stimulus showed a significant difference from the empty chamber response in
all tasks (Fig. 2C-N). In contrast, we found that across all four tasks, non-transitional
bouts during the encounter stage did not show any difference in the response between
the stimuli (Fig. O-R, Fig. S4A-H). These results, which are further apparent when all
four tasks are averaged (Fig. 2S-T), suggest that a differential response between the two
stimuli occur in mPFC pyramidal neurons only during transitional investigation bouts, but
not during non-transitional bouts. They also suggest that it is the preferred stimulus that
elicit an inhibitory response during transitional bouts, while the excitation observed
during non-preferred stimulus investigation is a generic response during chamber
investigation.

III. Context-Dependent Encoding of Relative Stimulus Value in the mPFC
Our results so far show that mPFC CaMKII neurons are differentially responsive to
investigation bouts following a behavioral shift, where the animal leaves one stimulus in
order to investigate the other stimulus, and are less responsive to investigation bouts
that involve repeating the same action consecutively. Yet, this differential response may
be sensitive to the absolute appeal of the stimulus, determined by its identity, or to its
relative appeal, determined by the other option (stimulus) presented in the same context,
rather than by the specific identity of the stimulus itself.
As motioned above, all four tasks used by us shared a common stimulus - a naïve male
conspecific. However, the preference for this stimulus differed between the tasks: it was
the preferred stimulus in the SP and SvF tasks, while in SxP and ESPs tasks it was the
non-preferred stimulus, as shown by the RDI values (Fig. S3A). Despite the identical

nature of this common stimulus across the four tasks, mPFC pyramidal neurons activity
during transitional bouts toward this stimulus depended on whether the stimulus was the
preferred or the non-preferred in that given task (Fig. 3B-C). Specifically, neural activity
during transitional bouts toward the common stimulus was mostly inhibitory in sessions
where the common stimulus was the preferred stimulus (RDI > 0), while in sessions
where it was the non-preferred stimulus (RDI < 0) the neural activity was excitatory (Fig.
3D). This difference was specific for transitional bouts and was not detected during non-
transitional bouts (Fig. 3E-F).

To test directly whether the observed inhibition of mPFC pyramidal neurons in transition
bouts is indeed related to the relative appeal of the stimulus rather than its identity, we
examined how the neural activity of mPFC pyramidal neurons changes when stimuli
identity is held constant but their relative appeal is altered. To that end, we recorded
animals performing the SvF task following 24 hours of food deprivation (Fig. 3G). As
expected, animals exhibited a strong preference for the food over the social stimulus, as
reflected by the time dedicated to investigate each stimulus and the number of non-
transitional bouts (Fig. 3H-I). Accordingly, RDI values were significantly different
between satiety and food deprivation (Fig. 3J). This change in preference was
accompanied by a "flip" in the neural activity pattern observed in the SvF task during
food deprivation, as compared to satiety (Fig. 3K-P). Specifically, transitional bouts
toward the food elicited inhibitory responses, as compared to excitatory responses to the
social (common) stimulus (Fig. 3P). Moreover, the responses during transitional bouts to
the same stimulus, whether food or social, were opposite between satiety, when the
social stimulus was preferred, and food deprivation, where food was the preferred
stimulus (Fig. 3Q-R). These results further demonstrate that the mPFC neurons
differential response during transitional investigation bouts depends on the relative
appeal of the stimulus, with an inhibitory response elicited specifically by the preferred
stimulus.

**The relationship between inhibited activity of mPFC pyramidal neuron activity and
social preference was further assessed** by calculating the probability of social
investigation following significant negative or positive changes in calcium activity (Fig.
S5A-L). This Analysis revealed that negative calcium transients are predictive of the
subjects' behavior in all four tasks, given that they were followed by a statistically
significant increased probability of investigation behavior toward the preferred stimulus,
but not toward the non-preferred stimulus (Fig. 4A-P). Moreover, this increase in

probability during the encounter was significantly higher than that recorded during the pre-encounter stage, suggesting that the negative transients are associated with the investigation of the preferred stimulus. In contrast, no such significant change from baseline or the pre-encounter stage was recorded for positive transients (Fig. 4Q-R, Fig. S6A-H).

Together, these findings suggest that decreased activity of mPFC pyramidal neurons serves a functional role in regulating investigation behavior toward the preferred stimulus and can be predictive of the animal's stimulus investigation behavior.

IV. Optogenetic Stimulation of mPFC Pyramidal Neurons During Stimulus Investigation Modulates Social Choice in a Context-Dependent Manner

To further examine whether pyramidal neurons of the mPFC play a critical role in mediating social choice, we used optogenetic stimulation to activate these neurons during stimulus investigation. For that, we used AAV viral vectors to express the excitatory opsin Channelrhodopsin-2 (ChR2) under the control of the CaMKIIα promoter in the mPFC, followed by optic fiber implantation for blue light delivery (Fig. 5A). Photo-stimulation (7–10 ms pulses at 10 Hz) was delivered specifically during investigation bouts toward a given stimulus (from onset to end of bout, Fig. 5B). Each task was repeated three times under three distinct stimulation protocols: no stimulation, stimulation during bouts with one stimulus, and stimulation during bouts with the other stimulus. Notably, there were no differences in the total investigation time between the four tasks in the "no stimulation" condition, indicating a similar level of social motivation without stimulation (Fig. S7A).

We found that optic stimulation delivered during investigation bouts with the preferred stimulus strengthened the preference toward it in SP, ESPs, and SvF, while merely preserving it in the SxP task, as reflected by the investigation time (Fig. 5C,H,M,R) and RDI (Fig. S7B-E). Conversely, stimulating the mPFC during bouts toward the non-preferred stimulus changes the subject's preference, abolishing it in specifically in the SP task and reversing it to create a preference toward the typically non-preferred stimulus in the SxP, ESPs, and SvF tasks. Thus, stimulating mPFC pyramidal neurons during stimulus investigation seem to enhance the investigation of that stimulus.

We then examined how optic stimulation of mPFC pyramidal neurons affects the structure of stimulus investigation. We first analyzed the time course of all investigation bouts toward a given stimulus and, to our surprise, found that in almost all cases

(besides SvF) the optic stimulation shortens the mean investigation bout rather than
lengthening it (Fig. 5D-E,I-J,N-O,S-T). We then examined the number of transitional and
non-transitional bouts toward each stimulus in all three stimulation conditions. In most
cases we found a significantly greater number of non-transitional bouts toward the
stimulus paired with optic stimulation, while the transitional bouts were affected less
consistently (Fig. 5F,K,P,U; Fig. S7F-M). Accordingly, the ratio between non-transitional
to transition bouts was always significantly higher for the stimulation-paired stimulus
compared to the other stimulus (Fig. 5G,L,Q,V). Moreover, the ratio of non-translational
bouts to all bouts pulled across all tasks wag significantly higher for the stimulation-
paired stimulus compared to the other stimulus or non-stimulation condition (Fig. 5W),
suggesting that optic stimulation elicit a high level of repetitive investigation bouts toward
the stimulation-paired stimulus.
We next tested whether the effect of mPFC pyramidal neuron activation was specific to
social contexts, by delivering optic stimulation during investigation bouts with an empty
chamber. We found that subject animals exhibited a preference for the empty chamber
coupled with optic stimulation (Fig. S8A-I). Thus, it seems as if stimulation of the mPFC
pyramidal neurons exerts a mixed effect on the subject's investigation behavior,
regardless of the stimulus nature: on one hand it shortens the investigation bouts, which
may reflect avoidance toward the stimulus, but on the other hand it increases the
number of non-transitional investigation bouts and made them more repetitive, which
may reflect persistent interest in the stimulus. In order to farther examine this effect, we
looked at the video recordings of the experiments and found that following the
stimulation that was activated at the beginning of an investigation bout, subject animals
rapidly stopped investigating the stimulus, but immediately after the termination of
stimulation they started a new investigation of the same stimulus, a sequence that
repeated itself several times before the animal made a transition and turned to
investigate the other stimulus (Supp. Movie 1-2).
Taken together, the results of the optogenetic stimulation demonstrate that activation of
mPFC pyramidal neurons exerts an immediate aversive effect on social investigation,
prompting the subject to abruptly terminate its ongoing interaction with the stimulus.
However, this manipulation does not alter the motivational drive of the subject to
investigate that stimulus. Consequently, once the optic stimulation is terminated, subject
animals are more likely to resume investigating the same stimulus (i.e., perform a non-
transitional bout) to satisfy their initial motivation for investigating the stimulus, leading to

a counter-intuitive increase in the investigation time for that stimulus (see cartoon in Fig.
5X).
300

V. Functional Dissociation of mPFC Projections' Role in Mediating Social
Choice

So far, we examined the activity of the whole population of mPFC pyramidal neurons.
However, sub-populations of these neurons project to several distinct brain regions [29],
hence may have different principles organizing their activity during a binary choice
between stimuli. Therefore, we examined the activity of two such sub-populations: those
that project to the nucleus accumbens (NAc) and those that project to the basolateral
amygdala (BLA), during the same four tasks used by us so far. To target mPFC neurons
projecting to a given region we injected a retrogade CAV-Cre viral vector mixed with an
AAV vector expressing mCherry (to label the injection site) into either the NAc or the
BLA. We also injected into the mPFC an AAV vector leading to the expression of
GCaMP7 under the human EF1α promoter in a Cre-dependent manner (Fig. 6A-B).
Upon recovery, we used fiber photometry to record calcium signals while subject
animals performed the four tasks, as described above.

In the SP, SxP, and ESPs tasks, subject animals of both groups (mPFC-to-NAc and
mPFC-to-BLA) displayed a similar preference as reported earlier for the CaMKII-
GCaMP-injected animals in SP, SxP, and ESPs tasks (Fig. 6C,E,G,I,K,M). In the SvF
task, however, subject animals from both groups displayed no preference for either
stimulus (Fig. 6O,Q).

We next examined the neural activity of mPFC-to-NAc and mPFC-to-BLA projecting
neurons during non-transitional and transitional bouts in each task. In all four tasks, both
populations of mPFC projecting neurons exhibited no differential responses during non-
transitional bouts to either of the stimuli. Similar to the general population of pyramidal
neurons, differences in neural responses—when present—were only found during
transitional bouts. Yet, these differential responses were task-dependent.

In the SP task, both populations of projecting neurons exhibited a similar pattern of
activity changes during transition bouts, with inhibitory responses to the social stimulus
specifically in transitional bouts (Fig. 6D,F). In the SxP task, neither population showed
any significant response during transitional bouts (Fig. 6H,J). In ESPs, only mPFC-to-
BLA projecting neurons showed inhibitory response during transitional bouts with the

stressed stimulus (Fig. 6L,N). Conversely, In the SvF task only the mPFC-to-NAc

projecting neurons population exhibited inhibitory responses during transitional bouts

with the social stimulus compared to the food (Fig. 6P,R).

Taken together, the mPFC-to-NAc and the mPFC-to-BLA projecting neurons exhibit

distinct pattern of responses during social choice in a task-dependent manner. Although

their engagement varies across tasks, whenever these projections are recruited, their

response profile follows the same consistent pattern observed in the broader CaMKII+

neuronal population: inhibition during transitional bouts toward the preferred stimulus.

Discussion

Social decision-making is a cornerstone of adaptive behavior, requiring the continuous evaluation of social cues and leading to selection of appropriate actions. Here, we provide compelling evidence that the mPFC plays a central role in this process by encoding the relative value of social choices through a context-dependent activity pattern. Our principal finding is that mPFC pyramidal neurons exhibit profound inhibition during transitional bouts toward a preferred stimulus, a response that is not only specific to the preferred option but also predictive of future choices. Such selective inhibition, coupled with a non-specific excitation during investigation of non-preferred stimuli, reveals a complex neural mechanism for social choice that is dynamically updated based on the relative appeal of available options.

Our results show that the mPFC does not encode the absolute identity of a social stimulus, but rather its relative value within a given context, as demonstrated by the "flipping" of the neural response to the same naïve male stimulus, which elicited inhibition when it was the preferred option (in the SP and SvF tasks) but excitation when it was non-preferred (in the SxP and ESPs tasks). These findings are consistent with our previous work on emotion recognition, where we observed inhibition of PrL pyramidal neurons during investigation of emotionally aroused individual, which is preferred over a naïve individual [28]. Context-dependent encoding was further confirmed by the SvF task, where the social stimulus elicited inhibition under satiety conditions (when it was the preferred option), but excitation following 24 hours of food deprivation (when food was the preferred option). **Such modulation of the neural response demonstrates** that the mPFC's encoding of social value is not fixed but rather modulated by the animal's internal motivational state, suggesting that inhibition of pyramidal neuronal activity may be a key mechanism by which the mPFC integrates internal state and external context to guide adaptive social choices. However, the exact neural circuit underlying the inhibitory and excitatory signals we observed remains to be fully elucidated. One possibility is that the inhibition of pyramidal neurons is mediated by local inhibitory interneurons, such as somatostatin (SOM) or parvalbumin (PV) interneurons. Recent work by Scheggia et al. (2020) [30] has demonstrated that SOM interneurons in the prefrontal cortex play a crucial role in discriminating between different affective states[22]. SOM interneurons may also be involved in the inhibitory signal we observed during interactions with preferred stimuli, perhaps by suppressing pyramidal neuronal

activity when the animal commits to a preferred option. Future studies using cell-type-
specific manipulations will be necessary to test this hypothesis.

While previous work has established that the mPFC encodes value in non-social,
operant tasks with explicit rewards [14, 31, 32], our findings demonstrate this principle in
the context of free and spontaneous social interactions- a distinction that highlights a key
advantage of our approach over conditioning paradigms. In operant tasks, animals are
trained to associate specific stimuli with rewards or punishments, and the mPFC's role
may be limited to executing learned associations. In contrast, our paradigms capture the
mPFC's involvement in innate, spontaneous social discrimination, where animals must
continuously evaluate and compare options without explicit training. These differences in
behavioral context may explain why other operant task-based studies focusing on the
mPFC during social approach have reported primarily excitatory responses [7, 16]. The
inhibitory signal we observed appears specific to contexts where animals are making
unconstrained choices between multiple options, as opposed to executing a learned
response to obtain a reward, **suggesting** that the neural mechanisms underlying value
encoding may differ fundamentally between spontaneous and conditioned social
behaviors and underscoring the importance of studying social behavior in multiple
settings.

Additionally, we found that the differential neural responses were primarily observed
during transitional bouts, when the subject animal was switching between stimuli, and
not during non-transitional bouts, which reflect sustained engagement. These findings
suggest that the mPFC is particularly involved in the process of behavioral switching and
reevaluation of choices, rather than in maintaining ongoing behavior. The inhibition
observed during transitional bouts to the preferred stimulus may reflect a neural "reset"
that facilitates the disengagement from the previous (non-preferred) option and the
commitment to the new (preferred) one. The lack of differential activity during non-
transitional bouts may indicate that once a choice is made and the animal is engaged in
sustained investigation, the mPFC's role in value encoding becomes less critical, and
other brain regions may take over the maintenance of the ongoing behavior. The
distinction between transitional and non-transitional bouts is critical for understanding the
dynamic nature of mPFC encoding. It demonstrates that the mPFC is not simply
representing the value of a stimulus in a static manner, but is actively tracking the
animal's behavioral state and updating its encoding based on the animal's previous
decisions. Recent work by Gabriel et al. (2025) [33] supports this notion by showing that

prelimbic cortex population dynamics rapidly generate novel representations during
action selection, with these representations emerging early in learning and predicting
whether animals will take avoidance or exploratory actions. Such dynamic encoding is
essential for adaptive behavior, as it enables the animal to remain flexible and
responsive to environmental changes while maintaining stability once a choice has been
made.

Our optogenetic experiments provide causal evidence for the role of mPFC activity in
biasing social choice. Stimulating mPFC pyramidal neurons during the investigation of a
stimulus enhanced the preference for that stimulus, regardless of its initial value.
Paradoxically, the stimulation itself had an immediate aversive effect, causing the animal
to cease social interaction and move away from the stimulus, which aligns with previous
work showing that optogenetic stimulation of the mPFC can have negative or aversive
effects on behavior, including enhanced fear associations and avoidance [13, 34-36].
However, upon termination of the optic stimulation, animals rapidly and repeatedly re-
engaged with the same stimulus, as reflected in an increased number of non-transitional
bouts. Such pattern of repetitive returns resulted in a net increase in total investigation
time and a robust preference bias toward the stimulated option. Therefore, we propose
that broad excitation in mPFC pyramidal neurons may serve as a "caution" signal,
reflecting a state of heightened vigilance or alertness that, if passing a certain threshold,
drives the animal to cease current behavior and explore alternatives. **The caution
signal interpretation explains** why optogenetic stimulation of mPFC pyramidal neurons
leads to immediate avoidance and shortened bout duration: artificial excitation may have
mimicked a caution signal, prompting the animal to disengage from the social interaction
prematurely. Supporting this interpretation, we observed an excitation effect during
transitional bouts between empty chambers in the pre-encounter period, which was
stronger than the effect found in non-transitional bouts. Transitional bouts, by definition,
involve a shift from one stimulus to another, which may be a moment of heightened
uncertainty or caution as the animal evaluates whether to commit to the new option. In
contrast, non-transitional bouts reflect sustained engagement with a stimulus, a state in
which the animal is naturally less cautious and more committed. By extension, the
profound inhibition we observed during transitional bouts toward the preferred stimulus
may reflect the suppression of this caution signal, effectively creating a "low-caution
state" that permits the animal to commit to and engage in sustained social interaction
with the preferred option. However, future experiments using optogenetic or

chemogenetic inhibition will be crucial for testing whether artificially inducing mPFC
inhibition can create a low-caution state that promotes sustained investigation of any
stimulus, regardless of its initial value.
Our recordings from mPFC neurons projecting to the nucleus accumbens (NAc) and
basolateral amygdala (BLA) indicate that these projection-specific populations exhibit
distinct functional specializations that depend on the behavioral choice context dictated
by the type of stimuli. This finding aligns with a growing body of literature demonstrating
functional specialization of mPFC output pathways. For instance, recent work has shown
that mPFC-NAc projections specifically encode social winning behavior, while mPFC-
BLA projections are more involved in processing negative emotional valence [37, 38].
The distinct responses we observed in mPFC-NAc and mPFC-BLA projecting neurons
may reflect a division of labor, where one pathway signals the value of the chosen option
to reward-related circuits, while the other signals the emotional or social context to
valence-processing regions. Future studies using projection-specific manipulations will
be necessary to determine whether the inhibitory signal we observed is specific to
certain projection pathways and whether selectively manipulating these pathways can
alter social choice.
Several limitations of our study should be acknowledged. First, while we targeted the
mPFC, our viral injection and optic fiber implantation did not allow for a precise
distinction between the prelimbic (PrL) and infralimbic (IL) subregions. Given the known
functional heterogeneity of the mPFC, with the PrL and IL often playing distinct or even
opposing roles in behavior [6, 12], our findings should be interpreted as reflecting the
integrated activity of the mPFC as a functional unit. The PrL has been implicated in
approach behavior and social cognition, while the IL is more often associated with
avoidance and fear extinction [39]. It is possible that the inhibitory and excitatory signals
we observed arise from different subregions, or that the relative contributions of PrL and
IL vary depending on the specific discrimination task. Future studies using more targeted
approaches, such as subregion-specific viral injections or localized optogenetic
manipulations, will be necessary to dissect the specific contributions of the PrL and IL to
social choice. Second, while our fiber photometry recordings revealed a strong
correlation between mPFC inhibition and social choice, this technique does not provide
the single-cell resolution needed to understand the underlying circuit dynamics in detail.
Single-unit recordings or calcium imaging with cellular resolution will be necessary in

future studies to elucidate the precise interplay of excitatory and inhibitory neurons that
gives rise to the observed population-level signals.
Our findings have several broader implications for our understanding of the neural basis
of both typical and atypical social behavior. A hallmark feature of ASD is a disruption in
the balance of excitation and inhibition (E/I balance) in cortical circuits, including the
mPFC [21, 40]. This E/I imbalance has been proposed to underlie many of the core
symptoms of ASD, including social deficits, repetitive behaviors, and sensory
abnormalities. Our findings suggest that a disruption of the mPFC's value encoding
mechanism—specifically, the inhibitory signal that marks commitment to a preferred
social option—could contribute to the social choice deficits observed in ASD. If the
inhibitory signal is weakened or absent, individuals with ASD may have difficulty
committing to social interactions or may fail to appropriately prioritize social stimuli
according to their value. Conversely, if excitatory activity is abnormally elevated, this
could lead to a heightened "caution" signal that promotes avoidance of social
interactions. Indeed, previous work has shown that restoring the balance of excitation
and inhibition in the mPFC can rescue social behavior deficits in mouse models of ASD
[40-42], suggesting that the integrity of mPFC circuits is crucial for normal social
function. Future studies examining whether the context-dependent value encoding
mechanism we observed is disrupted in mouse models of ASD will be important for
understanding the neural basis of social deficits in this disorder and for developing
targeted interventions.
In conclusion, our study provides novel insights into the neural mechanisms of social
choice, demonstrating that the mPFC encodes the relative value of social options
through a context-dependent inhibitory signal that is dynamically updated based on the
subject's behavioral state and motivational context. These findings not only advance our
understanding of the social brain but also provide a new framework for investigating the
neural basis of social deficits in neuropsychiatric disorders.

531
532
533

Declarations

Acknowledgments

We thank Boris Shklyar, Head of the Bio-imaging Unit and Eng. Alex Bizer, the experimental systems engineer of the Faculty of Natural Sciences of the University of Haifa, for their technical help.

Author contributions

R.J.: Formal Analysis, Investigation, Methodology, Validation, Visualization, Writing - original draft, and Writing - review & editing; S.N.: Data curation, Project administration, Software, Validation, Visualization, Writing - original draft, and Writing - review & editing. S.W.: Conceptualization, Funding acquisition, Project administration, Resources, Supervision, Writing - original draft, and Writing - review & editing.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was supported by the Israel Science Foundation (ISF) (grant No. 2220/22), the Ministry of Health of Israel (grant No. 3-19884for PainSociOT), the German Research Foundation (DFG) (SH 752/2-1), the Congressionally Directed Medical Research Programs (CDMRP) (grant No. AR210005), the United States-Israel Binational Science Foundation (grant No. 2019186) and the HORIZON EUROPE European Research Council (ERC-SyG oxytocINspace).

558
559

Supplemental Information

Table S1. An Excel file containing details of all statistical analyses, organized by figure.

Table S2. An Excel file containing processed datasets analyzed and organized by figure.

Tables S3. Contains optogenetic stimulation parameters and data retention rate.

Methods

Resource availability

Further information and requests for resources and reagents should be directed to and will be fulfilled by Shlomo Wagner (shlomow@research.haifa.ac.il).

Materials availability

This study did not generate new unique reagents.

Data and code availability

All the data generated from experiments and the custom codes used to evaluate the conclusions of this paper would be uploaded onto a public repository upon acceptance of the paper.

Main-text figure legends

Figure 1. Behavioral Analysis across four social discrimination tasks.

A. Schematic representation of the SP task.

B. Mean time ($\pm$SEM) dedicated by subject mice (n=49 sessions from 21 animals) for investigating the animal stimulus (dark blue) or the object (light blue), and mean sum of transitional bouts between the two stimuli ($\pm$SEM, gray lines) across the 5-min encounter period of the SP task in (1-min bins).

C. Median (in box plot) value of mean investigation time of i the social stimulus and the object. Each grey line connects data points of the two stimuli from the same session.

D. Median (in box plot) value of mean duration of all investigation bouts toward the social stimulus and the object.

E. Median (in box plot*) value of the number of non-transitional bouts with the social stimulus and the object.

F-J. As in **A-E**, for the SxP task (n=40 sessions).

K-O. As in **A-E**, for the ESPs task (n=30 sessions).

P-T. As in **A-E**, for the SvF task (n=35 sessions).

$p<0.001$, paired samples t-test in **C,* **H***,* **M** and **R**.

*$p<0.05$, **$p<0.01$, ***$p<0.001$, Wilcoxon signed-rank test in **D-E**, **I-J**, **N-O**, and **S-T**.

594   Box plot represents 25 to 75 percentiles of the distribution, while the bold line is the median of the
595   distribution. Whiskers represent the smallest and largest values in the distribution.

597   **Figure 2. mPFC pyramidal neurons display context-dependent calcium dynamics during**
598   **transitional and non-transitional bouts.**

A.   On the left, a schematic representation of the GCaMP6-expressing AAV vector injection
        followed by optic fiber implant in the mPFC. On the right, a fluorescent micrograph of an
        AAV vector-infected mPFC slice, showing GCaMP6-expression (green) and optic fiber
        location (dashed line).

B.   Timeline for a typical session of behavioral testing with fiber photometry recording.

C.   Heat-map (see color code to the right of **F**) of the z-scored calcium signals at the
        beginning (first six seconds) of bout, averaged across all transitional bouts toward the
        stimulus animals, shown for all sessions of the SP task (n=49 sessions, with each row
        corresponding to a single session), using 0.5 s bins. Time' 0' represents the beginning of
        bout.

D-F.   As in **C** for the female (SxP; n=40), stressed (ESPs; n=30), and social (SvF; n=35)
        stimuli, respectively.

G-J.   As in **C** for the non-preferred stimuli across the various tasks.

K.   Superimposed traces of the mean (±SEM) z-scored calcium signals shown in **C and G**,
        averaged across all SP sessions. The dark blue and light blue trace lines represent the z-
        scored calcium signals in transitional bouts to the social stimulus and the object,
        respectively, during the encounter stage of the SP sessions. Dashed black trace is for
        transitional bouts to empty chambers during the pre-encounter stage. Blue-shaded areas
        represent two distinct time windows of the event, which are used for statistical Analysis.
        Asterisks, representing statistical comparisons between the stimuli and across time
        windows, are color-coded according to the type of comparison at the top of the graph.
        Blue and light blue asterisks represent a significant difference in social and object z-
        scored signals, respectively, during the specific time window relative to baseline. Black
        asterisks on dashed lines represent a difference in responses to the two stimuli during
        the same time window. Asterisks on the right side of the graph represent statistical
        differences for social and object z-scored signals in the encounter stage of the session,
        during the 3-6 s time window, relative to the pre-encounter stage, and are color-coded
        according to the stimulus with the significant difference. A significant effect was found for
        stimulus ($p<0.001$), time ($p<0.001$), and their interaction ($p<0.001$) by two-way MM
        ANOVA.

L-N. As in **K**, for transitional bouts in the SxP, ESPS, and SvF tasks, respectively. In all three
        tasks, a significant effect was found for stimulus ($p<0.05, p<0.01, p<0.001$, respectively),

631   time ($p<0.001$), and their interaction ($p<0.01$ in SxP and ESPs; $p<0.001$ in SvF) by two-
632   way MM ANOVA.

O-R.   As in **K**, for non-transitional bouts in the SP, SxP, ESPs, and SvF tasks, respectively. A
634   significant effect was found for time ($p<0.01$ in SP, $p<0.001$ in SxP, $p<0.01$ in ESPs, and
635         $p<0.05$ in SvF) by two-way MM ANOVA.

Q.   Mean trace for the z-scored activity during all transitional bouts from the pooled sessions
of the four tasks during the pre-encounter stage (in black), transitional bouts to the
preferred stimulus (in blue), and transitional bouts to the non-preferred stimulus (in
orange). Please note the difference between the inhibitory response in the trace for
transitional bouts to the preferred stimulus, compared to the traces for transitional bouts
in the pre-encounter stage and to the less-preferred stimulus.

$*p<0.05$, $**p<0.01$, $***p<0.001$, *post-hoc* paired (within stimulus differences in time) or
independent (differences between stimuli within time windows, or differences between pre-
encounter and encounter) t-test with Holm-Šídák correction for multiple comparisons following
detection of main effects by ANOVA.

Figure 3. mPFC pyramidal neurons' activity tracks relative stimulus preference across
648 **contexts and internal motivational states.**

A.   Median relative discrimination index (RDI) for the common stimulus in each of the four
650   tasks. A significant effect was found for task ($p<0.001$) by one-way ANOVA.

B.   Heat-map (see color code to the right of **C**) of the average z-scored calcium signals at
652   the beginning (first six seconds) of transitional bouts to the common stimulus, using 0.5-
653   s bins. Time' 0' represents the beginning of the bout. Each line corresponds to a single
654   session in which the common stimulus was the preferred stimulus (RDI > 0), pooled
655   from all four tasks.

C.   As in **B**, for transitional bouts to the common stimulus in sessions where it was the non-
657   preferred stimulus (RDI<0), pooled from all four tasks.

D.   Superimposed traces of the mean (±SEM) z-scored calcium signals shown in **B-C**,
659   averaged across all sessions. Gray-shaded areas represent three distinct time windows
660   of the event, which were used for the statistical Analysis. A significant effect was found
661   for stimulus ($p<0.001$), time ($p<0.001$), and their interaction ($p<0.001$) by two-way MM
662   ANOVA.

E.   Median z-score of the calcium signal during non-transitional bouts to the common
stimulus in each of the four tasks, divided into two time windows: -1-to-2 s and 3-to-6 s
relative to the bout onset. A significant effect was found for the time effect ($p<0.001$) by
two-way MM ANOVA.

F.   As in **E**, for transitional bouts. A significant effect was found for stimulus ($p<0.001$), time
($p<0.001$), and their interaction ($p<0.001$) by two-way MM ANOVA.

G.   Schematic representation of the SvF task following 24 hours of food deprivation.

H.   Median time dedicated by subject animals to investigate the social stimulus (in purple)
and the food (in brow, n=24 sessions) during the 5-min encounter period of the SvF-food
deprivation task. Each grey line connects data points of the same session.

I.   Median sum of non-transitional bouts to the social stimulus and the food during the 5-
minute encounter period of the task.

J.   Median relative discrimination index (RDI) during the 5-min encounter period of the two
conditions of the SvF tasks (satiety and food deprivation).

K.   Heat-map (see color code to the right of **L**) of the z-scored calcium signals at the
beginning (first six seconds) of non-transitional bouts, averaged across all non-
transitional bouts to the social stimulus during the 5-minute encounter stage of all
sessions of the SvF-food deprivation task.

L.   As in **K**, for non-transitional bouts to the food.

M.   Superimposed traces of the mean (±SEM) z-scored calcium signals shown in **K-L,**
averaged across all SvF-food deprivation sessions. Paige-shaded areas represent two
distinct time windows of the event, which are used for the statistical Analysis. A
significant effect was found for stimulus ($p<0.01$) by two-way MM ANOVA.

N-P.   As in **K-M**, for transitional bouts. A significant effect was found for stimulus ($p<0.05$),
time ($p<0.001$), and their interaction ($p<0.05$) by two-way MM ANOVA.

Q.   Superimposed traces of the mean (±SEM) z-scored calcium signals for transitional bouts
to the social stimulus in the two conditions of the SvF task.

R.   As in **Q**, for transitional bouts to the food.

***$p<0.001$, *post-hoc* pairwise comparisons with Holm-Šídák correction following detection of
main effects by ANOVA in **A**.

*$p<0.05$, **$p<0.01$, ***$p<0.001$, *post-hoc* paired (within stimulus differences in time) or
independent (differences between stimuli within time windows) t-test with Holm-Šídák correction
for multiple comparisons following detection of main effects by ANOVA in **D-F**, **M** and **P**.

*$p<0.05$, **$p<0.01$, independent t-test with Holm-Šídák correction for multiple comparisons in **Q-**
R.

Figure 4. Negative calcium transients in mPFC pyramidal neurons' activity predict social
investigation behavior.

A.   Heat-map (see color code to the right) of the z-scored social stimulus investigation
probability following (4 seconds in 0.2 s bins) the onset of negative change events in the

calcium signal (time '0'), averaged across all negative events within each session, with

each row representing one session of the SP task.

B.   Superimposed traces of the mean (+SEM) social stimulus investigation probability for 4

seconds following the onset (time '0') of negative change events identified in the pre-

encounter stage (black dashed line) and in the encounter stage (dark blue filled line), z-

scored to the probability in the -4 to 0 s window in 0.2 s bins, and averaged across all SP

sessions.

C-D. As in **A-B**, for the object in SP sessions.

E-H. As in **A-D**, for the SxP task.

I-L.   As in **A-D**, for the ESPs task.

M-P. As in **A-D**, for the SvF task.

Q.   Mean traces for the investigation probability of the preferred stimulus (in blue) and the

non-preferred stimulus (in orange) following negative change events in the calcium signal

during the encounter stage of all sessions of the four tasks. Dashed traces represent the

investigation probability of empty chambers in the pre-encounter stage, later used to

house the preferred and non-preferred stimuli. Please note the high probability of

investigating the preferred stimulus in the encounter stage compared to the other traces.

R.   As in **Q**, for positive change events.

*p<0.05, **p<0.01, ***p<0.001, One-sample Wilcoxon signed rank test (within trace differences in

time) or Wilcoxon signed rank test (differences between pre-encounter and encounter).

Figure 5. Optogenetic activation of mPFC pyramidal neurons modulates social

preference.

A.   On the left, A schematic representation of the ChR2-expressing AAV vector injection

followed by optic fiber implantation in the mPFC. On the left, a fluorescent micrograph of

an infected mPFC slice, showing ChR2-mCherry expression (red) and the position of the

optic fiber (dashed line).

B.   A session example for the investigation times of stimulus 1 (in green), stimulus 2 (in red),

and periods of optic stimulation delivered during investigation bouts with stimulus 2 (in

blue), across the 5-min duration of the session. The highlighted gray square represents

an example bout with stimulus 2 in which optic stimulation (10 ms pulses at 10 Hz

frequency) was delivered at the onset of the bout (when the subject's nose touches the

mesh of the stimulus chamber) and terminated at the end of the bout (when the subject

turns away from the stimulus chamber)

N.   Median investigation time of the social (dark blue) and object (light blue) stimuli (n=14),

in SP sessions with no stimulation, or when optogenetic stimulation was consistently

applied during investigation bouts with the social stimulus, or during investigation bouts

with the object. A significant effect was found for stimulus ($p<0.001$), condition ($p<0.05$),
and their interaction ($p<0.001$) by two-way RM ANOVA.

C. Superimposed traces of the normalized mean ($\pm$SEM) probability for investigating the
social stimulus 6 seconds, in 0.2 s bins, following the bout onset (optic stimulation onset)
in SP sessions with no optic stimulation (in gray), and in SP sessions with optic
stimulation delivered during bouts with the social stimulus (in dark blue). The gray-
shaded area represents the time window (0 to 4 s) taken for statistical Analysis.

D. As in D, for the object stimulus in SP sessions with no optic stimulation (in gray), and in
SP sessions with optic stimulation delivered during bouts with the object (in light blue).

E. Median sum of non-transitional bouts to the social stimulus and the object across the
three optic stimulation conditions of the SP task.

F. Median logarithmic ratio of non-transitional/transitional bouts for the social stimulus and
the object across the three optic stimulation conditions of the SP task. The statistics
shown in the figure were calculated from the original data, which was transformed into
logarithmic values for graphical convenience due to the significant variation in the original
data. A significant effect was found for condition by Kruskal-Wallis test for Social
($p<0.001$) and object ($p<0.01$).

H-L. As in **C-G**, for the SxP task.

M-Q. As in **C-G**, for the ESPs task.

R-V. As in **C-G**, for the SvF task.

W. Median ratio of non-transitional/all bouts for the preferred and the non-preferred stimuli,
pulled from all four tasks across the three optic stimulation conditions (no stimulation,
stimulation with the preferred stimulus, and stimulation with the non-preferred stimulus).
Please note the significant increase in the non-transitional/alll bouts ratio for the stimulus
coupled with the optic stimulation.

X. Schematic representation of the behavioral sequence observed during optogenetic
stimulation pyramidal neurons in the mPFC. (1) A subject mouse implanted with an optic
fiber investigates a stimulus mouse positioned behind a wire mesh barrier. Both animals
display engaged social interaction. (2) Blue light stimulation is delivered to the subject
mouse, activating mPFC pyramidal neurons at the onset of the investigation bout with the
stimulus. (3) The subject mouse exhibits an immediate avoidance response and abruptly
ceases interacting with the stimulus. (4) Following the end of investigation bout, and optic
stimulation is terminated. (5) Once the optogenetic stimulation ceases, the subject mouse
displays renewed motivation to investigate the stimulus. The subject mouse then
promptly returns to the mesh barrier to resume investigation of the same stimulus,
demonstrating a non-transitional investigation bout.

p<0.05, **p<0.01, ***p<0.001, *post-hoc* paired t-test with Holm-Šídák correction following
detection of main effects by RM ANOVA in **H**, **M**, and **R**.

p<0.05, **p<0.01, independent test in **D-E**, **I-J,** **N-O,** and **S-T**.

p<0.05, **p<0.01, independent t-test with Holm-Šídák correction for multiple comparisons in **Q-**
R.

p<0.05, **p<0.01, ***p<0.001, Wilcoxon signed rank test with Holm-Šídák correction for multiple
comparisons in **F**, **K**, **P**, and **U**.

p<0.05, **p<0.01, ***p<0.001, *post-hoc* pairwise comparisons with Holm-Šídák correction for
multiple comparisons following detection of main effects by Kruskal-Wallis test in **G**, **L**, **Q**, and **V-**
W.

Figure 6. Projection-specific calcium dynamics in mPFC-to-NAc and mPFC-to-BLA
neurons during transitional bouts.

A. A schematic representation of the CAV_Cre and mCherry expressing AAV vectors
injection into the NAc, and the GCaMP7f AAV vector injection into the mPFC, followed by
optic fiber implantation in the mPFC. On the left, a fluorescent micrograph of an infected
mPFC slice, showing mCherry (red) and GCaMP7f (green) and the position of the optic
fiber (dashed line). On the right, a fluorescent micrograph of an infected NAc slice,
showing mCherry (red) expression.

B. As in **A**, for the injection and expression of GCaMP in mPFC-to-BLA projecting neurons.

C. Median investigation time of the social stimulus and the object (n=27 sessions from 11
animals) during the 5-min encounter period of SP sessions recorded from animals with
GCaMP expression in mPFC-to-NAc neuronal cells.

D. Superimposed traces of the mean (±SEM) z-scored calcium signals for transitional bouts
with the social stimulus (dark blue) and object (light blue), averaged across all SP
sessions, and aligned to the bout onset (time '0'). Blue-shaded areas represent two
distinct time windows of the event, which were used for the statistical Analysis. A
significant effect was found for stimulus ($p<0.01$) and time x stimulus interaction ($p<0.01$)
by two-way MM ANOVA.

E-F. As **B-D**, for SP sessions recorded from animals with GCaMP expression in mPFC-to-BLA
neuronal cells (n=25 sessions from 12 animals).

G-J. As in **C-F**, for the SxP task ($n_{mPFC-to-NAc}=19$; $n_{mPFC-to-BLA}=31$).

K-N. As in **C-F**, for the ESPs task ($n_{mPFC-to-NAc}=22$; $n_{mPFC-to-BLA}=27$).

O-R. As in **C-F**, for the SvF task ($n_{mPFC-to-NAc}=18$; $n_{mPFC-to-BLA}=32$).

p<0.05, ***p<0.001, paired t-test in **C**, **E**, **G**, **I**, **K,** and **M**.

STAR Methods

Method details

Experimental model and subject details

Animals

C57BL/6J: C57BL/6J naïve adult (8-18 week-old) male and female mice (Envigo RMS, Israel) were used either as subjects or stimuli.

Maintenance: Animals were housed in groups of 2-5 sex-matched mice per cage. All behavioral sessions were conducted while subjects were group-housed, with the exception of the SP task in CaMKIIα–GCaMP6f–injected mice, which were recorded after two to three weeks of individual housing to enhance social motivation. All animals were housed at the animal facility of the University of Haifa under veterinary supervision, in a 12 h light/12 h dark cycle (lights on at 9 PM), with *ad libitum* access to food (standard chow diet, Envigo RMS, Israel) and water.

Institutional review board

All experiments were performed according to the National Institutes of Health guide for the care and use of laboratory animals and approved by the Institutional Animal Care and Use Committee of the University of Haifa.

Experiments

Behavioral experiments were conducted in the dark phase of the dark/light cycle in a sound attenuated chamber (Faraday cage), under dim red light.

Experimental setups

The experimental setup [43] consisted of a white Plexiglas arena (37 X 22 X 35 cm) placed in the middle of an acoustic cabinet (60 X 65 X 80 cm). Two Plexiglas triangular chambers (12 cm isosceles, 35 cm height), into which an animal or object (plastic toy) stimulus could be introduced, were placed in two randomly selected opposite corners of the arena. A metal mesh (12 X 6 cm, 1 X 1 cm holes) placed at the bottom of the

triangular chamber allowed direct interaction with the stimulus through the mesh. A high-
quality monochromatic camera (Flea3 USB3, Flir for optogenetic experiments or
Grasshopper3 for fiber photometry recordings) equipped with a wide-angle lens was
placed at the top of the acoustic chamber and connected to a computer, enabling a clear
view and recording (~30 frames/s) of subject behavior using commercial software
(FlyCapture2 or SpinView, FLIR).

Behavioral paradigms

All behavioral paradigms started with 15 minutes of habituation in which subject animals
were moved into the experimental arena with two empty stimuli chambers. Throughout
this time, social stimuli were placed in their chambers near the acoustic cabinet for
acclimation. After habituation, the chambers containing the relevant stimuli were
diagonally placed at opposite ends of the arena in a random fashion. Each task was
conducted for 5 minutes, and differed mainly in the stimuli presented to the subjects-

Social preference (SP): Subject mice were introduced to either a social (naïve, novel,
age- and sex-matched conspecific) and an object (mouse-size lego block).

Sex Preference (SxP): subject mice were presented with a novel male on one side, and
a novel female conspecific on the other.

Emotional State Preference for stress stimuli (ESPs) - subject mice were introduced to a
naive stimulus animal and to another stimulus animal that had been constrained for 15
min in a 50 ml tube pierced with multiple holes for ventilation. Each "stressed" stimulus
animal was used as a stimulus for one session only.

Social vs. food (SvF, in satiety): Subject mice were introduced to a novel, naïve, social
stimulus, and to food pellets composed of standard chow, under satiety conditions (*ad*
libitum access to food and water prior to the task). Food pellets were placed inside a
dedicated chamber fitted with a metal mesh with small holes (0.5 × 0.5 cm) that
prevented subjects from grabbing or consuming the food during testing, thereby
minimizing distraction and maintaining engagement with the experimental task.

Social vs. food (SvF, in food deprivation)- subject mice were food deprived, with full
access to water for 24 hours before the task. In the task, subjects were introduced to a
novel, naïve, social stimulus, and to food pellets composed of standard chow.

Stereotactic surgery for viral injection and optic fiber implantation

Surgery: Mice were anesthetized and analagized with an intraperitoneal injection of a mixture of ketamine, domitor, and the painkiller meloxicam (0.13 mg/g, 0.01 mg/g, and 0.01 mg/g, respectively). Anesthesia levels were monitored by testing toe pinch reflexes and maintained using isoflurane (0.2%; SomnoSuite, Kent Scientific Corporation). Body temperature was kept constant at approximately 37°C using a closed-loop custom-made temperature controller connected to a temperature probe and a heating pad placed under the animal. Ophthalmic ointment (Duratears, Alcon, Couvreur, NV) was applied to maintain eye moisture. Anesthetized animals were fixed in a stereotaxic apparatus (Kopf), with the head flat. The head was shaved, the scalp was disinfected, and the skin was removed to expose the skull. The skull was then leveled using bregma–lambda measurements.

The region of interest was then marked, and a hole (unilateral, right hemisphere) was drilled for viral injection and optic fiber implantation. Additional holes were drilled for supporting screws. Viral injection was performed as previously described in detail (https://star-protocols.cell.com/protocols/1283) using a glass capillary filled with the virus ssAAV-1/2-mCaMK2a-GCaMP6f-WPRE-SV40p(A) for fiber photometry recordings or ssAAV-1/2-mCaMK2α-hChR2(H134R)_mCherry-WPRE-SV40p(A) for excitatory optogenetic stimulation, was slowly lowered into the mPFC (AP, +2 mm; ML, -0.35 mm; DV, +2.0 mm) and left in place for 5 min prior to injection and 10 min following injection to prevent retraction of the virus. For targeting mPFC projecting neurons to the NAc and the BLA, CAV2-Cre (Canine Adenovirus Type 2 expressing Cre recombinase) in combination of 1:5 ratio with ssAAV-1/2-hSyn1-hChR2(H134R)_mCherry-WPRE-hGHp(A) for marking the injection site were injected into either the NAc (AP, +1.42 mm; ML, -1.2 mm; DV, +4.4 mm) or the BLA (AP, -1.43 mm; ML, -3.0 mm; DV, +4.8 mm). ssAAV-1/2-hEF1a-dlox-jGCaMP7f(rev)-dlox-WPRE-bGHp(A) was injected into the mPFC (AP, +2 mm; ML, -0.35 mm; DV, +2.0 mm). The Cav_cre virus was obtained from the Plateforme de Vectorologie de Montpellier (PVM), Institut de Génomique Fonctionnelle, CNRS, University of Montpellier, France. All other viruses were purchased from the Viral Vector Facility (VVF), Institute of Pharmacology & Toxicology, University of Zurich, Switzerland.

A total of 300 nl of the relevant virus was then delivered by manual application using a 50 ml syringe connected to the glass capillary through a plastic tube for applying air pressure (BRAND, disposable BLAUBRAND micropipettes, intraMark, 5 μl). Following

viral injection, an optic fiber (Doric lenses, 200 µm, NA 0.66, 3 mm long, zirconia ferrule,

flat-fiber tip) was inserted into the mPFC with the exact coordinates as the viral injection,

and placed 200 μm above the injection depth. Screws and optic fiber were fixed to the

skull plate using dental cement (UNIFAST, GC America).

After the surgery, antisedan (0.1 mL/10g body weight) was administered subcutaneously

to wake the mice from anesthesia. The mice were injected with meloxicam (5%, 0.01

mg/g) and baytril (5%, 0.03 ml/10g) to relieve pain and prevent infections for three days

following surgery. Behavioral testing with optogenetic stimulation or fiber photometry

was conducted three weeks after surgery to allow optimal viral expression. Following

surgery, all subjects were housed with one or two cage mates.

In vivo fiber photometry recordings

Test: All fiber photometry recordings were performed using the apparatus described

above. The optic fiber of the implanted mouse was connected to the optical patch cords

via a sleeve connector (Doric) under brief isoflurane anesthesia. The wired animal was

then allowed to habituate in the behavior setup for 15 min. The recording session started

with an additional 5-minute baseline (pre-encounter) period with empty chambers.

Thereafter, the empty chambers were replaced with chambers containing the

appropriate stimuli for a 5-minute test. Following the test, the stimulus-containing

chambers were again replaced with empty ones for an additional 5 minutes. Each

subject was recorded in two different task sessions per day, with at least 40 minutes

between sessions for the same animal.

Fiber photometry recording and synchronization: Calcium signals were recorded using

one of two fiber photometry recording setups composed of the RZ10x system of Tucker

Davis Technologies and an optical path by Doric, which includes 600 µm mono-fiber

optic patch cords connected to either a four ports Fluorescence MiniCube

(FMC4_IE(400-410)_E(460-490)_F(500-550)_S) for one-color recordings or a six ports

Fluorescence MiniCube (FMC6_IE(400-410)_E1(460-490)_F1(500-540)_E2(555-

570)_F2(580-680)_S) for two-colors recordings. Optic paths were connected using a 200

µm optical patch cord with a fiber-optic rotary joint (FRJ_1×1_PT_200/230/LWMJ-

0.57_1m_FCM_0.15m_FCM) that was connected to the recorded animal. A camera

(Grasshopper3 USB3, FLIR) was placed on top of the arena and connected to the digital

port of the RZ10✗ system and configured to deliver strobes for every frame acquired at a

frame rate of 30 frames/sec. These were later used to synchronize the video frames with

the calcium signal. TDT Synapse software was used to record the GCaMP6f/7f signal

channel (excitation at 465 nm, modulated at 330 Hz), the isosbestic control channel (405

nm, modulated at 210 Hz), and the digital channel receiving the camera strobes, with a

sampling rate of ~$10^{17.3}$ Hz. LED currents were adjusted to return the voltage to

between 0 and 200 mV for each signal, were offset by 5 mA, and were demodulated

using a 6 Hz low-pass frequency filter. Light intensity levels for the 465-nm and 405-nm

channels were verified by measuring output power with an external power meter

(PM100D, Thorlabs) every 2–3 recording days.

***In vivo* optogenetic stimulation**

<u>Test:</u> In all optogenetic experiments (SP, SxP, ESPs, SvF–Satiety, and SvF–Food

Deprivation), subject mice (n = 14) were connected to the stimulation apparatus and

allowed to habituate for 20 min in an arena containing empty chambers. After

habituation, the empty chambers were replaced with the appropriate stimuli, and

interactions were recorded for 5 min. In control experiments with empty chambers, new

empty chambers were introduced into the arena at this stage.

Each subject was tested three times in the SP, SxP, ESPs, and SvF tasks, each session

employing a different stimulation condition: (1) no stimulation, (2) stimulation delivered

during investigation bouts with one of the two presented stimuli, and (3) stimulation

delivered during investigation bouts with the other stimulus. Optic stimulation during an

investigation bout was manually delivered and defined as light delivery from the onset of

nose contact with the mesh of a stimulus chamber until the subject's head turned away

from it.

In the SvF–Food Deprivation and empty-chambers tasks, each mouse was tested

twice—once without stimulation and once with optic stimulation delivered during

investigation bouts with either the social stimulus (SvF–Food Deprivation) or one of the

empty chambers.

Each day, subject mice participated in either a single experiment with optogenetic

stimulation or in two sequential experiments, consisting of a non-stimulated session

followed 20 minutes later by a stimulated session. In the SvF–Food Deprivation task,

both conditions were conducted on the same day, with the non-stimulated session

always preceding the stimulated session, to avoid repeated food-deprivation

manipulations. All other tasks were conducted in a randomized order.

<u>Optogenetic stimulation:</u> Subject mice were lightly anesthetized with isoflurane and

connected to the laser (473 nm, model FTEC2471-M75YY0, Blue Sky Research) via a

fiber-optic rotary joint attached to 200 µm core multimode fiber patch cables connecting

the animal to the light source for optic stimulation delivery (Thorlabs, Fiber Optic Rotary

Joint Patch Cables for Optogenetics, product group #6865). Animals were given 20 min

to recover from any residual effects of the brief isoflurane anesthesia before testing, until

they showed full mobility. In all experiments, optogenetic stimulation consisted of

continuous light pulses delivered at 10 Hz throughout the entire duration of each

investigation bout with a given stimulus. The initial photo-stimulation protocol used 10

ms pulses at a light power of 8.23 mW. However, these parameters induced seizure-like

behavior in a subset of animals, which we attributed to the high level of ChR2 expression

in the mPFC. This phenomenon has been previously reported in optogenetic studies of

cortical regions, where brief optogenetic stimuli can elicit seizures that increase in

severity with repeated stimulation (Cela et al., 2019; Osawa et al., 2013). To ensure that

experimental outcomes reflected controlled neuronal modulation rather than pathological

network states, we systematically optimized the stimulation parameters by first reducing

light power from 8.23 mW to no lower than 4.65 mW, and subsequently shortening pulse

duration from 10 ms to 7 ms, while maintaining the 10 Hz frequency (see Tables S3A-B

for detailed timeline and voltage-light power calibration). The minimal used parameters

of 4.65 mW and 7 ms pulses are well within the established range for effective ChR2

activation in vivo. Using a computational model based on direct measurements in

mammalian brain tissue [44] (https://web.stanford.edu/group/dlab/cgi-

bin/graph/chart.php), we calculated that 4.65mW of 473nm light delivered through a

200µm fiber (NA 0.66) produces an irradiance of approximately 4.99mW/mm² at 250µm

distance from the fiber tip (the maximal distance between the fiber tip and the virus

injection site), and 1.26mW/mm^2 at a distance of 500µm which is the highest penetration

depth of blue light stimulation in rodent brain tissues [36]. These values exceed the

generally accepted threshold of 1 mW/mm² required for suprathreshold activation of

ChR2-expressing neurons [44-46], while remaining within the recommended safe

operating range of less than 10-20 mW/mm² at target cells to minimize tissue heating

[36]. Furthermore, our pulse duration of 7 ms is well above the opening rate time

constant of ChR2 (τ ≈ 1.21 ms), ensuring sufficient channel opening to trigger action

potentials reliably [3-Lin 2011]. Optical stimulation was manually delivered via a Master8

Channel Programmable Pulse Stimulator (A.M.P.I). Sessions in which severe seizures

prevented completion of behavioral testing were excluded from analysis. Overall, 12 out

of 150 stimulation sessions (8%) were excluded due to severe seizures, yielding a 92%

data retention rate across all animals (Table S3C). Importantly, even at the optimized

parameters (4.65 mW, 7 ms pulses, 10 Hz), animals continued to exhibit the intended

behavioral response pattern, characterized by abrupt termination of investigation bouts

upon stimulation onset, followed by immediate re-engagement with the same stimulus

once stimulation ceased, confirming that the optimized parameters were sufficient to

produce the desired modulatory effect on mPFC pyramidal neuron activity. Importantly,

even at the lowest parameters (4.65 mW, 7 ms pulses, 10 Hz), animals continued to

exhibit the intended behavioral response pattern, characterized by the abrupt termination

of investigation bouts upon stimulation onset, followed by immediate re-engagement with

the same stimulus once stimulation ceased (see Supp. Movie 3).

1026

***Post-mortem* histological Analysis of viral injections and optic-fiber placement**

Each implanted mouse was anesthetized with isoflurane (induction: 3%, 0.5%-0.8%

maintenance in 200 mL/min of air), attached to the stereotaxic device, and the optic fiber

implant was gently extracted from its head. The mouse was then perfused with

phosphate-buffered saline (PBS) and fixed using a 4% paraformaldehyde (PFA, Sigma)

solution. The brains were harvested and placed in PFA (4%) for 48 hours, followed by

sectioning into 50-μm slices in the horizontal axis using a VT1200s Leica sliding

vibratome. The slices were collected onto microscope slides and stained with DAPI

(Fluoroshield with DAPI, histology mounting medium, Merck). Images of subject brain

slices were acquired for verification of the viral expression placement and optic fiber path

within the mPFC using an epifluorescence microscope (Ti2 eclipse, Nikon) equipped

with a blue filter for DAPI staining, FITC for detecting GCaMP6f/7f-expressing cells, and

TRITC for detecting mCherry-expressing cells.

1040

1041   Quantification and statistical analysis

1042   All processed data are detailed in Table S2, organized by the various figures.

Tracking software and behavioral analyses

All video recordings were analyzed using *TrackRodent* (https://github.com/shainetser/TrackRodent) [47]. The *BlackMouseWiredBodyBased* algorithm was applied to sessions involving tethered animals in both fiber photometry and optogenetic experiments. Behavioral analyses were performed as previously described [48].

Parameters for behavioral quantification

All behavioral parameters reported in this study were automatically extracted using the TrackRodent software, which continuously tracks the animal's position and investigation of the stimuli chambers throughout the experiment. Behavioral events were identified and quantified using predefined algorithms and rule-based classifications embedded in the software. The Relative Discrimination Index (RDI) was calculated as the difference in investigation time between the two stimuli—specifically, investigation of the preferred stimulus minus the investigation of the non-preferred stimulus—normalized by the total investigation time toward both stimuli. For the naïve male stimulus (the common stimulus across all four tasks), the RDI was calculated as the investigation time for the common stimulus (independent of preference) minus the investigation time for the other stimulus, divided by the sum of both.

Investigation bouts were defined as events beginning when the animal's head (or body center, depending on the selected algorithm in TrackRodent software) made contact with the metal mesh of the stimulus chamber and ending when the animal disengaged from the mesh. Each bout was subsequently categorized as either non-transitional or transitional. Non-transitional bouts were those preceded by a bout directed toward the same stimulus, whereas transitional bouts were those preceded by a bout directed toward the alternate stimulus. A minimum interval of >0.5 s between successive bouts was required to classify bouts as distinct. Behavioral events of interest detected by TrackRodent were then aligned with the calcium imaging data to enable precise synchronization between neural activity and behavior.

Synchronization of physiological data to behavioral events

We used a custom-made code called DeepPhenotyping to visualize and synchronize the different signals and behavioral events, as previously described [49, 50]. For the calcium signal with each event type ("Transitional bouts" and "Non-transitional bouts"), the

software considered the relevant sample rate and aligned the signal to the beginning of
the event. For calculating the subjects' investigation probability of a given stimulus
between different optic stimulation conditions, the investigation time for the relevant
stimulus was synchronized to the onset of all bouts (regardless of type) which is also the
onset of the optical stimulation in conditions where optic stimulation was delivered.
Derived investigation probabilities were then normalized to the maximum (divided by
P_{max} of 1) and used directly for statistical Analysis.

1083

Calcium signal data analysis
Calcium signals were analyzed using DeepPhenotyping. We first fitted the 405 nm
channel onto the 465 nm channel to de-trend signal bleaching and any movement
artifacts, according to the manufacturer's protocol(https://github.com/tjd2002/tjd-shared-
code/blob/master/matlab/photometry/FP_normalize.m). Next, the de-trended signal was
aligned to the video recording using the timestamps recorded by the digital port of the
RZ10x system. We aligned the de-trended signal to each bout type (transition\non-
transition) event and normalized it using the z-score (0.5 s bins), with the pre-event
period of -4 s to -1 s serving as a baseline.
As for calculating the population average signal amplitude over time, we first calculated
the AUC for each session in time bins of five seconds (using the original detrended
signal which is samples at 1017.3Hz) and then averaged the signals across all sessions.
For calculating the probability of social investigation following significant negative or
positive changes in calcium activity, we first segmented the GCaMP signal into 0.2-s
time bins. For each bin, neural activity was z-scored within an 8-s window (–4 s to +4 s
relative to the bin onset) using the preceding 4 s (–4 s to 0 s) as baseline. The mean z-
score during the 4 s following each bin was then compared to baseline activity. Time
bins showing a significant deviation ($|Z| \geq 1.96$) were classified as either positive or
negative calcium events. Following the detection of a positive or negative event, a 6-s
interval was imposed before the next sampling bin to ensure that detected events were
distinct and did not reflect the same change in the recorded signal at adjacent time
points. All identified positive and negative events were subsequently aligned with the
animal's investigation time of a given stimulus, and the probability of investigation was
calculated in 8-s time windows centered on the event onset. The mean probability during
the –4 to 0 s interval served as the baseline for z-scoring post-event (0 to 4 s)

probabilities. z-scored probabilities for positive and negative events were then averaged

across all events pooled within each session. The area under the curve (AUC) in the 0-

to4s time window was computed to quantify differences in investigation probability

following positive versus negative events, as well as to assess deviations from baseline

(0) within each event type.

Statistical Analysis

All statistical results and parameters are presented in Table S1, organized according to

the corresponding figures. Sample sizes for all behavioral experiments were determined

based on previously published power calculations [48]. Statistical analyses were

performed using SPSS v27.0 (IBM) and GraphPad Prism v10.1.2 (GraphPad Software).

The Kolmogorov–Smirnov and Shapiro–Wilk tests were used to assess the normality of

the dependent variables. For normally distributed data, one-sample t-tests were used to

compare group means to a fixed reference value (0 in this case). Two-tailed paired t-

tests were used to compare conditions or stimuli within the same group, and two-tailed

independent t-tests were used to compare variables between distinct groups. When

normality assumptions were violated, nonparametric alternatives were applied. The

Wilcoxon signed-rank test was used to compare group medians to a set value (0) or

between conditions within the same group, and the Mann–Whitney U test was used to

compare conditions between groups. For comparisons involving multiple factors,

analyses of variance (ANOVA) were conducted according to the experimental design.

Mixed-model (MM) ANOVA was applied when data included both within-subject

(repeated-measures) and between-subject factors. This model included one random

effect (subject ID), one within-subject effect, one between-subject effect, and their

interaction. Two-way between-subjects ANOVA was used for comparisons between

groups involving two independent factors, whereas two-way repeated-measures (RM)

ANOVA was used for comparisons within groups across multiple conditions or variables.

The latter model included one random effect (subject ID), two within-subject effects, and

their interaction. When the assumption of sphericity required for ANOVA was violated,

the Greenhouse–Geisser correction was applied. Whenever significant main effects or

interactions were detected, post hoc Student's t-tests with Holm–Šidák correction were

conducted. For repeated-measures data violating normality assumptions, Friedman's

nonparametric test was applied, followed by post hoc Dunn's paired comparisons. For

between-group comparisons involving more than two levels, the Kruskal–Wallis test was

used, followed by Dunn's multiple-comparisons post hoc tests. Statistical significance was set at $p < 0.05$.

References

1. Chen, P. and W. Hong, *Neural Circuit Mechanisms of Social Behavior.* Neuron, 2018. **98**(1): p. 16-30.
2. Gangopadhyay, P., et al., *Prefrontal–amygdala circuits in social decision-making.* Nature neuroscience, 2021. **24**(1): p. 5-18.
3. Kietzman, H.W. and S.L. Gourley, *How social information impacts action in rodents and humans: the role of the prefrontal cortex and its connections.* Neuroscience & Biobehavioral Reviews, 2023. **147**: p. 105075.
4. Cao, W., H. Li, and J. Luo, *Prefrontal cortical circuits in social behaviors: an overview.* Journal of Zhejiang University-SCIENCE B, 2024. **25**(11): p. 941-955.
5. Bicks, L.K., et al., *Prefrontal Cortex and Social Cognition in Mouse and Man.* Front Psychol, 2015. **6**: p. 1805.
6. Ko, J., *Neuroanatomical Substrates of Rodent Social Behavior: The Medial Prefrontal Cortex and Its Projection Patterns.* Front Neural Circuits, 2017. **11**: p. 41.
7. Lee, E., et al., *Enhanced neuronal activity in the medial prefrontal cortex during social approach behavior.* Journal of Neuroscience, 2016. **36**(26): p. 6926-6936.
8. Yashima, J., T. Uekita, and T. Sakamoto, *The prelimbic cortex but not the anterior cingulate cortex plays an important role in social recognition and social investigation in mice.* PloS one, 2023. **18**(4): p. e0284666.
9. Kuga, N., et al., *Prefrontal-amygdalar oscillations related to social behavior in mice.* eLife, 2022. **11**: p. e78428.
10. Li, H., et al., *Brain circuits that regulate social behavior.* Molecular Psychiatry, 2025: p. 1-17.
11. van den Bos, W. and B. Güroğlu, *The role of the ventral medial prefrontal cortex in social decision making.* Journal of Neuroscience, 2009. **29**(24): p. 7631-7632.
12. Euston, D.R., A.J. Gruber, and B.L. McNaughton, *The role of medial prefrontal cortex in memory and decision making.* Neuron, 2012. **76**(6): p. 1057-1070.
13. Levy, D.R., et al., *Dynamics of social representation in the mouse prefrontal cortex.* Nat Neurosci, 2019. **22**(12): p. 2013-2022.
14. Glanzberg, J.T., et al., *Individual differences in prelimbic neural representation of food and cocaine seeking.* Cell reports, 2024. **43**(12).
15. Frost, N.A., et al., *Context-invariant socioemotional encoding by prefrontal ensembles.* Nature Communications, 2025. **16**(1): p. 5455.
16. Isaac, J., et al., *Sex differences in neural representations of social and nonsocial reward in the medial prefrontal cortex.* Nature Communications, 2024. **15**(1): p. 8018.
17. Lam, N.H., et al., *Effects of altered excitation-inhibition balance on decision making in a cortical circuit model.* Journal of Neuroscience, 2022. **42**(6): p. 1035-1053.

18. Tan, T., et al., *Neural circuits and activity dynamics underlying sex-specific*
effects of chronic social isolation stress. Cell Rep, 2021. **34**(12): p. 108874.
19. Huang, W.-C., Y. Chen, and D.T. Page, *Hyperconnectivity of prefrontal cortex to*
amygdala projections in a mouse model of macrocephaly/autism syndrome.
Nature communications, 2016. **7**(1): p. 13421.
20. Liu, Y., et al., *A molecularly defined mPFC-BLA circuit specifically regulates*
social novelty preference. Science Advances, 2025. **11**(17): p. eadt9008.
21. Yizhar, O., et al., *Neocortical excitation/inhibition balance in information*
processing and social dysfunction. Nature, 2011. **477**(7363): p. 171-178.
22. Kingsbury, L. and W. Hong, *A multi-brain framework for social interaction.* Trends
in neurosciences, 2020. **43**(9): p. 651-666.
23. Klune, C.B., B. Jin, and L.A. DeNardo, *Linking mPFC circuit maturation to the*
developmental regulation of emotional memory and cognitive flexibility. Elife,
2021. **10**: p. e64567.
24. Ebstein, R.P., et al., *Genetics of human social behavior.* Neuron, 2010. **65**(6): p.
831-844.
25. McGraw, L.A. and L.J. Young, *The prairie vole: an emerging model organism for*
understanding the social brain. Trends in neurosciences, 2010. **33**(2): p. 103-
109.
26. Carlén, M., *What constitutes the prefrontal cortex?* Science, 2017. **358**(6362): p.
478-482.
27. Netser, S., et al., *Distinct dynamics of social motivation drive differential social*
behavior in laboratory rat and mouse strains. Nat Commun, 2020. **11**(1): p. 5908.
28. Jabarin, R., et al., *Distinct prelimbic cortex neuronal responses to emotional*
states of others drive emotion recognition in adult mice. Current Biology, 2025.
35(5): p. 994-1011. e8.
29. Anastasiades, P.G. and A.G. Carter, *Circuit organization of the rodent medial*
prefrontal cortex. Trends in neurosciences, 2021. **44**(7): p. 550-563.
30. Scheggia, D., et al., *Somatostatin interneurons in the prefrontal cortex control*
affective state discrimination in mice. Nature neuroscience, 2020. **23**(1): p. 47-60.
31. Sul, J.H., et al., *Distinct roles of rodent orbitofrontal and medial prefrontal cortex*
in decision making. Neuron, 2010. **66**(3): p. 449-460.
32. Wang, Q., et al., *Distributed value representation in the medial prefrontal cortex*
during intertemporal choices. Journal of Neuroscience, 2014. **34**(22): p. 7522-
7530.
33. Gabriel, C.J., et al., *Transformations in prefrontal ensemble activity underlying*
rapid threat avoidance learning. Current Biology, 2025. **35**(5): p. 1128-1136. e4.
34. Klavir, O., et al., *Manipulating fear associations via optogenetic modulation of*
amygdala inputs to prefrontal cortex. Nature neuroscience, 2017. **20**(6): p. 836-
844.
35. Yizhar, O., *Optogenetic insights into social behavior function.* Biological
psychiatry, 2012. **71**(12): p. 1075-1080.
36. Yizhar, O., et al., *Optogenetics in neural systems.* Neuron, 2011. **71**(1): p. 9-34.
37. Choi, T.-Y., et al., *Distinct prefrontal projection activity and transcriptional state*
conversely orchestrate social competition and hierarchy. Neuron, 2024. **112**(4):
p. 611-627. e8.
38. Huang, W.-C., et al., *Social behavior is modulated by valence-encoding mPFC-*
amygdala sub-circuitry. Cell reports, 2020. **32**(2).
39. Giustino, T.F. and S. Maren, *The role of the medial prefrontal cortex in the*
conditioning and extinction of fear. Frontiers in behavioral neuroscience, 2015. **9**:
p. 298.

40. Selimbeyoglu, A., et al., *Modulation of prefrontal cortex excitation/inhibition*
balance rescues social behavior in CNTNAP2-deficient mice. Science
translational medicine, 2017. **9**(401): p. eaah6733.
41. Luo, Y., et al., *Reduced excitatory activity in the developing mPFC mediates a*
PVH-to-PVL transition and impaired social cognition in autism spectrum
disorders. Translational Psychiatry, 2024. **14**(1): p. 325.
42. Ferguson, B.R. and W.-J. Gao, *PV interneurons: critical regulators of E/I balance*
for prefrontal cortex-dependent behavior and psychiatric disorders. Frontiers in
neural circuits, 2018. **12**: p. 37.
43. Netser, S., et al., *A novel system for tracking social preference dynamics in mice*
reveals sex-and strain-specific characteristics. Molecular autism, 2017. **8**(1): p.
53.
44. Aravanis, A.M., et al., *An optical neural interface: in vivo control of rodent motor*
cortex with integrated fiberoptic and optogenetic technology. Journal of neural
engineering, 2007. **4**(3): p. S143.
45. Cela, E., et al., *An optogenetic kindling model of neocortical epilepsy.* Scientific
Reports, 2019. **9**(1): p. 5236.
46. Britt, J.P., R.A. McDevitt, and A. Bonci, *Use of channelrhodopsin for activation of*
CNS neurons. Current protocols in neuroscience, 2012. **58**(1): p. 2.16. 1-2.16.
19.
47. Netser, S., et al., *A novel system for tracking social preference dynamics in mice*
reveals sex- and strain-specific characteristics. Mol Autism, 2017. **8**: p. 53.
48. Netser, S., et al., *A system for tracking the dynamics of social preference*
behavior in small rodents. Journal of Visualized Experiments (JoVE), 2019(153):
p. e60336.
49. Barbier, M., et al., *Altered Neural Activity in the Mesoaccumbens Pathway*
Underlies Impaired Social Reward Processing in Shank3-Deficient Rats.
Advanced Science, 2025. **12**(17): p. 2414813.
50. Mohapatra, A.N., et al., *Impaired emotion recognition in Cntnap2-deficient mice*
is associated with hyper-synchronous prefrontal cortex neuronal activity.
Molecular Psychiatry, 2025. **30**(4): p. 1440-1452.

1268

1269

Transitional bouts

Non-transitional bouts

Social Preference

Social

A

B *** ***

Object

C

D *

Sex Preference

Female

E

F ** *

Male

G

H **

Emotional State Preference-stress

Stressed

I

J ** **

Naive

K

L

Social vs. Food

Social

M

N *** *

Food

O

P

Q Negative changes

Mean Z-score probability

Time (s)

R Positive changes

Mean Z-score probability

Time (s)

Preferred_Pre
Non-preferred_Pre
Preferred_Enc
Non-preferred_Enc

A

promoter
mCaMKIIa — hChR2_mCherry

mPFC

M2
Cg1
PrL
IL
MO   VO

Optic fiber

DAPI
mCherry
1000μm

B

10 Hz, 10ms pulses

Beginning of bout → End of bout

475nm

Laser On   Laser Off

Optic stimulation
Stimulus 2 inv.
Stimulus 1 inv.

1   2   3   4

Time (min)

Social Preference

C Investigation Time (s) — No stim. with Social with Object — Social / Object — *** ***

D Normalized probability vs Time (s) — No stim. / with Social — **

E Normalized probability vs Time (s) — No stim. / with Object — *

F Number of non-trans. bouts — Social / Object — No stim. with Social with Object — ** * ** / * **

G Log10 (Non-trans.\Trans. ratio) — Social / Object — *** *** / * **

Sex Preference

H Investigation Time (s) — No stim. with Female with Male — Female / Male — ** ** ***

I Normalized probability vs Time (s) — No stim. / with Female — *

J Normalized probability vs Time (s) — No stim. / with Male

K Number of non-trans. bouts — Female / Male — No stim. with Female with Male — ** ** / ** **

L Log10 (Non-trans.\Trans. ratio) — Female / Male — ** *** / *** ***

Emotional State Preference-stress

M Investigation Time (s) — No stim. with Stressed with Naive — Stressed / Naive — * *** *

N Normalized probability vs Time (s) — No stim. / with Stressed — *

O Normalized probability vs Time (s) — No stim. / with Naive

P Number of non-trans. bouts — Stressed / Naive — No stim. with Stressed with Naive — ** ** / ** **

Q Log10 (Non-trans.\Trans. ratio) — Stressed / Naive — *** *** / *** ***

Social vs. Food

R Investigation Time (s) — No stim. with Social with Food — Social / Food — ** *** ***

S Normalized probability vs Time (s) — No stim. / with Social

T Normalized probability vs Time (s) — No stim. / with Food — *

U Number of non-trans. bouts — Social / Food — No stim. with Social with Food — *** ** / *** **

V Log10 (Non-trans.\Trans. ratio) — Social / Food — *** *** / * **

W Non-trans\All bouts ratio — Preferred / Non-preferred — No stim. with preferred with non-preferred — *** *** *** / *** *** ***

X

Start of an investigation bout

Delivery of optogenetic stimulation — ON

Abrupt termination of interaction — 3

Termination of optic stimulation — OFF — 4

Motivation to invesitigate stimulus persists

Return to stimulus ''Non-transitional bout'' — 5