Autolamellasomes: An Autophagy-Dependent Parallel Pathway for Bulk ER Degradation and Lysosome Biogenesis

Di Lu[1*], Rouxuan Zhang[2], Wanjing Shi[1], Delin Zhan[1], Yuntian Yang[1], Xulin Sun[1], Han Zhang[1], Ying Li[3], Xueming Li[2] and Li Yu[1*]

[1]The State Key Laboratory of Membrane Biology, Tsinghua University-Peking University Joint Center for Life Sciences, Beijing Frontier Research Center for Biological Structure, School of Life Sciences, Tsinghua University, Beijing 100084, China.

[2] The State Key Laboratory of Membrane Biology, Tsinghua University-Peking University Joint Center for Life Sciences, Beijing Frontier Research Center for Biological Structure, MOE Key Laboratory of Protein Sciences, School of Life Sciences, Tsinghua University, Beijing 100084, China.

[3]Technology Center for Protein Sciences, School of Life Sciences, Tsinghua University, Beijing, China.

* Correspondence: Li Yu, liyulab@mail.tsinghua.edu.cn;

Di Lu, ld2023@mail.tsinghua.edu.cn

Abstract

Autophagy maintains cellular homeostasis through lysosomal degradation of cytoplasmic components, yet how prolonged autophagy activation reshapes organelle architecture remains poorly understood. Here we identify a previously unrecognized form of endoplasmic reticulum (ER) remodeling induced by chronic mTOR inhibition. We show that sustained autophagy triggers the formation of multilamellar ER-derived structures, termed autolamellasomes, which mediate bulk ER degradation. Unlike canonical ER-phagy, autolamellosome biogenesis requires the core autophagy machinery but is independent of all known ER-phagy receptors. Using cryo-electron tomography, correlative light–electron microscopy, and *in vitro* reconstitution, we

demonstrate that autolamellasomes arise through autophagy-dependent assembly and compaction of fragmented ER membranes into concentric stacks subsequently engulfed by lysosomes. Autolamellosomes form constitutively at low levels across diverse cell types, accumulate in senescent cells and fibroblasts from Hutchinson–Gilford progeria syndrome (HGPS), and thus represent an aging-associated pathway linking sustained mTOR suppression to ER and lysosomal homeostasis. This work not only resolves the long-standing origin of intra-lysosomal membrane whorls but also establishes a cell-free system for dissecting autophagy-driven membrane remodeling, revealing a conserved degradative route coupling nutrient sensing to membrane turnover, lipid catabolism, and aging.

Key words

Autolamellasome, Autophagy, ER, Lysosome, mTOR

Introduction

Autophagy is an evolutionarily conserved catabolic process essential for maintaining cellular homeostasis in eukaryotes[1-4]. This lysosomal degradation pathway ensures the continuous turnover of cytoplasmic components—ranging from proteins and lipids to entire organelles—thereby supplying nutrients during starvation and removing damaged or superfluous structures[5-7]. The best-characterized form, macroautophagy (hereafter referred to as autophagy), entails the de novo formation of a double-membrane vesicle known as the autophagosome, which engulfs cytoplasmic cargo and delivers it to lysosomes for degradation[8-10].

Originally regarded as a non-selective "bulk" degradation process activated by nutrient deprivation, autophagy is now recognized as a highly selective and precisely regulated system[11-14]. Distinct selective autophagy pathways—including ER-phagy, mitophagy, xenophagy, and aggrephagy—target specific substrates such as damaged mitochondria,

intracellular pathogens, or protein aggregates[15]. This selectivity is conferred by cargo

receptors containing LC3-interacting region (LIR) motifs that bridge designated cargo

to the Atg8/LC3 family proteins decorating the phagophore[16-19].

A paradigmatic example of such precision is endoplasmic reticulum autophagy (ER-

phagy), a process that controls ER size, morphology, and function through the

lysosomal turnover of ER subdomains[20]. The ER forms a dynamic network responsible

for protein synthesis, lipid metabolism, and calcium storage; its homeostasis is critical

for cell survival and stress adaptation[21,22]. ER-phagy can occur via several

morphologically distinct routes: (1) macro-ER-phagy, in which ER fragments are

sequestered by autophagosomes[23]; (2) micro-ER-phagy, involving direct lysosomal

invagination and engulfment of ER regions[24]; (3) ER-to-lysosome–associated

degradation (ERLAD), where ER-derived vesicles directly fuse with endolysosomes;

and most importantly, receptor-mediated ER-phagy[25,26].

The initiation and specificity of receptor-mediated ER-phagy are governed by a family

of ER-phagy receptors—transmembrane or membrane-associated proteins that

recognize ER subdomains and recruit the autophagy machinery through conserved LIR

motifs[27]. Beyond serving as tethers, many receptors such as FAM134B[28] and RTN3L[29]

actively remodel the ER membrane through reticulon homology domains (RHDs) that

induce curvature and fragmentation, while intrinsically disordered regions (IDRs)

provide flexible linkers between ribosome-studded ER and the growing phagophore[30-

32]. Distinct receptors thus confer subdomain selectivity: FAM134B preferentially

targets ER sheets, whereas RTN3L and ATL3[33] act on tubules. Others, such as CCPG1,

also recruit upstream autophagy regulators (e.g., FIP200), directly initiating local

autophagosome formation at the ER surface[34,35]. Together, these mechanisms ensure

the spatial and temporal precision of receptor-mediated ER turnover.

A morphological hallmark of lysosomes is their dense, lamellar content composed of

intraluminal vesicles and membrane whorls[36]. Under various physiological or

pathological conditions, lamellated membrane structures—including membrane

whorls[37], myelin figures, and zebra bodies[38]—have long been observed within

lysosomes in diverse contexts such as autophagy, drug-induced phospholipidosis[39], and

lysosomal storage disorders like Fabry disease[38,40,41]. These concentric, multilamellar

stacks are thought to arise when membrane-rich autophagic cargo resists lysosomal

degradation, leading to lipid reorganization into multilamellar arrays[42].

Pharmacological or genetic disruption of lipid catabolism similarly promotes such

lamellation. First described by Rozenszajn et al. in 1969[43], these structures remain a

morphological hallmark of lysosomes, and their increased abundance is a sign of

lysosomal dysfunction. However, the biophysical mechanisms underlying their

initiation and persistence remain elusive.

In yeast, ER stress triggers the formation of large multilamellar ER whorls that are

eliminated via a micro-ER-phagy route distinct from canonical autophagy[24]. This

process involves direct invagination of the vacuolar membrane, depends on the ESCRT

machinery[44] and the ER-localized Nem1-Spo7 phosphatase complex, but proceeds

independently of ATG genes and classical ER-phagy receptors[45]. In mammalian cells,

severe ER stress similarly induces the formation of massive, concentric ER whorls—

up to 10 μm in diameter—through a PERK- and COPII-dependent pathway. These

structures function to alleviate ER stress by concentrating PERK and sequestering

translocons[46], and their formation and clearance are likewise independent of macro- and

micro-autophagy.

Here, we identify a distinct form of ER remodeling that emerges under chronic mTOR

inhibition—a condition that persistently activates autophagy[47]. We show that prolonged

mTOR suppression triggers the formation of extensive multilamellar ER-derived

structures, which we term autolamellasomes. Unlike conventional double-membrane

autophagosomes, autolamellasomes are composed of densely packed, concentric ER

membranes that are subsequently engulfed by lysosomes for degradation. Genetic and biochemical analyses reveal that their formation requires the core autophagy machinery but proceeds independently of all known ER-phagy receptors, defining a unique, autophagy-dependent yet receptor-independent ER remodeling and degradation pathway. Using cell-free reconstitution, we demonstrate that autolamellasomes arise from the autophagy-driven assembly and compaction of fragmented ER membranes. They are detectable under basal conditions but markedly accumulate in senescent cells and in fibroblasts derived from Hutchinson–Gilford progeria syndrome (HGPS) patients, consistent with chronic mTOR inhibition during aging. Temporally, autolamellosome formation follows conventional autophagosome-mediated cytosolic degradation and facilitates the bulk catabolism of lipid-rich ER membranes, sustaining long-term energy balance under nutrient stress.

Together, these findings define autolamellasomes as a parallel structure to autophagosomes that mediates bulk ER degradation under prolonged mTOR inhibition, linking mTOR signaling to membrane homeostasis and cellular aging.

Results

Prolonged mTOR Inhibition Induces the Formation of "Free Whorls"

Prolonged inhibition of mTOR by Torin1 led to the accumulation of numerous whorl-like membrane structures within autolysosomes (Fig. 1A). It is well-established that mTOR inhibition induces autophagy, characterized by the transient appearance of double-membraned autophagosomes. However, transmission electron microscopy (TEM) analysis revealed that 24 hours after Torin1 treatment, typical autophagosomes were rarely observed. Instead, cells contained abundant autolysosomes filled with densely packed membrane whorls with size of ~ 500 nm-2 μm, Intriguingly, we also observed multilamellar whorls composed of stacked double membranes that were not enclosed by a limiting membrane with similar size with whorls inside autolysosome. Because these structures resided outside autolysosomes, we refer to them as "free

whorls" (Fig. 1B).

Free whorls were frequently observed in close proximity to the endoplasmic reticulum

(ER). Notably, at the time points when free whorls appeared, conventional double-

membraned autophagosomes were nearly undetectable. Time-course TEM analysis

revealed that free whorls began to emerge approximately 4 hours after starvation and

progressively accumulated with prolonged treatment (Fig.1C). The number of

autolysosomes or lysosomes containing whorls increased with a similar trajectory

(Fig.1C), and the number of internal whorls within each whorl-bearing lysosome rose

correspondingly (Fig.1D). Similarly, inhibition of mTOR by other means—including

rapamycin treatment and nutrient deprivation—also induced prominent whorl

formation (Fig.2Q-2R). Moreover, this phenomenon was consistently observed across

all tested cell lines derived from multiple species, although the abundance of free

whorls and whorl-containing lysosomes varied among different cell types (Fig.1H,

Extended Fig.1A-1B).

Because chemical fixation can potentially introduce artifacts, we next employed cryo-

electron tomography (cryo-ET), which preserves native membrane ultrastructure with

high fidelity. Cryo-ET unambiguously confirmed the presence of free whorls and

revealed that each consisted of multiple concentric double membranes with a luminal

spacing of ~15 nm (Fig.1E-1F). Three-dimensional FIB-SEM further demonstrated that

each whorl forms a spherical structure (Fig.1G).

Together, these findings indicate that prolonged mTOR inhibition triggers the formation

of multilamellar free whorls, which subsequently appear to be engulfed by

autolysosomes or lysosomes, revealing an unexpected ultrastructural consequence of

sustained mTOR suppression.

The Free Whorls Originate from the Endoplasmic Reticulum

The frequent association of free whorls with the endoplasmic reticulum (ER) suggested that they may arise from the ER. To test this hypothesis, we first labeled live cells with ER-Tracker and observed that prolonged mTOR inhibition transformed the ER morphology from a reticular network into punctate structures (Fig.2A). Co-labeling with LAMP1-YFP and mCherry-REEP5 revealed the appearance of LAMP1-YFP$^-$/mCherry-REEP5$^+$ puncta, as well as the translocation of mCherry-REEP5 signal into LAMP1-YFP$^+$ lysosomes or autolysosomes (Fig.2C). Three-dimensional confocal imaging confirmed the existence of these REEP5-positive puncta (Fig.2D).

To exclude potential artifacts caused by the large fluorescent tag, we generated EGFP- or Halo-tagged REEP5 constructs and observed comparable puncta formation, confirming that the phenomenon is tag-independent (Extended Fig.2A, 2C). Quantitative analysis showed that nearly all cells undergoing prolonged mTOR inhibition developed ER puncta (Fig.2B, 2E-2F, Extended Fig.2B, 2D-2E). When other ER markers (mCherry-SEC61B, Calnexin-Halo, and ER-DsRed) were used, the proportion of cells with puncta was similar to that observed with REEP5 (Extended Fig.2F-2H).

To rule out artifacts arising from overexpression, we next examined endogenous ER proteins by immunostaining with antibodies against LAMP1 and either SEC61B or REEP5. Consistently, SEC61B- and REEP5-positive puncta were detected both inside and outside autolysosomes/lysosomes, supporting the conclusion that these structures are not overexpression artifacts (Fig.2G). Notably, the frequency and Pearson correlation coefficients in immunostained samples were lower than in overexpression systems, implying that ER protein overexpression can enhance puncta formation to some extent (Fig.2H-2K).

We next examined the ultrastructure of these puncta using APEX labeling for electron

microscopy. Because APEX2-catalyzed reactions generate electron-dense DAB precipitates, this method enables precise localization of tagged proteins[48]. APEX2-tagged SEC61B and REEP5 both exhibited dark, electron-dense labeling on the whorls (Fig.2L, Extended Fig.2M). Similarly, correlative light and electron microscopy (CLEM) confirmed colocalization of REEP5 signal with the whorls, providing direct evidence that the whorls originate from the ER (Fig.2M).

To capture the dynamics of ER puncta formation, we performed live-cell time-lapse imaging. The fraction of cells containing REEP5 puncta increased sharply within 5 hours after mTOR inhibition, accompanied by a parallel rise in the Pearson correlation coefficient with LAMP1-YFP (Fig.2N, Extended Fig.2I-2J). These observations indicate that ER puncta form rapidly and are subsequently delivered into autolysosomes for degradation.

Consistent with this, Western blot analysis revealed progressive degradation of ER proteins following prolonged mTOR inhibition. After 24 hours, most ER-resident proteins were degraded to varying degrees, with the notable exception of TEX264, which remained largely stable (Fig.2O). Five independent experiments confirmed this degradation pattern (Fig.2P). Label-free quantitative mass spectrometry produced similar results (Fig.2Q), although TEX264 appeared reduced in this dataset—likely reflecting normalization differences, as Western blots were normalized to actin, whereas MS data were normalized to total protein, leading to an apparent increase in actin and relative decrease in TEX264 in the Torin1-treated group.

Time-course analysis further revealed distinct degradation kinetics among ER proteins. The autophagy substrate p62 was degraded within 4 hours of mTOR inhibition, whereas ER proteins displayed diverse temporal profiles. For instance, TEX264 was degraded at early time points but its expression rebounded by 12 hours, suggesting enhanced transcription or translation. In contrast, ATL3 and REEP5 levels remained stable until

12 hours but declined sharply by 24 hours, while other ER proteins showed gradual

decreases (Fig.2R-2S).

Together, these data indicate that prolonged mTOR inhibition induces the formation of

ER-derived puncta, which are subsequently engulfed and degraded by

autolysosomes/lysosomes. The distinct kinetics and selectivity of degradation suggest

that multiple mechanisms—including transcriptional regulation, selective ER-phagy,

and puncta formation—contribute to the turnover of ER components under sustained

mTOR inhibition.

Formation of Whorls Depends on the Core Autophagy Machinery but Not ER-

Phagy Receptors

We next investigated the genetic requirements governing whorl formation. Given their

frequent association with ER membranes, we first examined whether known ER-phagy

receptors are required. Deletion of the major and widely recognized ER-phagy

receptors—including FAM134B[28], SEC62[49], TEX264[32,50], ATL3[33], and CCPG1[35]—did

not affect whorl formation. To further exclude the involvement of canonical ER-phagy

receptors, we generated a cell line lacking all five known receptors (FAM134B, ATL3,

SEC62, TEX264, and CCPG1), hereafter referred to as ERphR-KO cells. Across three

independently derived clones, simultaneous loss of these receptors had minimal impact

on whorl formation. We then asked whether genes implicated in the generation of large

ER whorls (~10 μm) under ER stress conditions might contribute. However, knockout

of these genes, or of key transcriptional regulators of the unfolded protein response

(UPR), similarly had no detectable effect on whorl formation (Fig.3A).

We next tested genes involved in general membrane dynamics based on functional

prediction. Among these, knockout of NPC2 modestly reduced whorl formation,

whereas deletion of PI4KIIA, VPS13C, or SCAP had no discernible effect. In striking

contrast, disruption of core autophagy genes markedly impaired whorl biogenesis. Deletion of ATG2 (ATG2A/2B DKO), ATG7, ATG9A, or ATG8 almost completely abolished whorl formation, while loss of ATG5, ATG14, ATG13, ATG16L, or ATG3 produced moderate reductions. Knockout of Beclin-1 or WIPI2 had a weak effect, and deletion of ULK1 or p62 showed no significant impact (Fig.3A). Together, these results demonstrate that the core autophagy machinery—but not canonical ER-phagy receptors—is essential for whorl formation.

We next assessed how disruption of these pathways affects ER protein turnover. Two complementary assays were employed. First, we monitored cleavage of mCherry-REEP5, a well-established reporter for ER-phagy. Knockout of autophagy core components—including ATG2, ATG5, and ATG7—completely blocked mCherry-REEP5 cleavage (Fig.3B-3C). In contrast, deletion of individual ER-phagy receptors (FAM134B, SEC62, ATL3) had no significant effect, while TEX264 and CCPG1 knockouts modestly reduced cleavage efficiency (Extended Fig.3A-3C). Second, we examined endogenous ER protein degradation. Core autophagy gene deletion broadly prevented degradation of multiple ER proteins, whereas ERphR-KO and individual ER-phagy receptor knockout cells exhibited normal degradation (Fig.3B, 3D-3E, Extended Fig.3A, 3D-3E). Collectively, these findings indicate that degradation of ER components under prolonged mTOR inhibition depends primarily on the core autophagy machinery, rather than on specific ER-phagy receptors.

We next analyzed ER morphology in cells lacking essential autophagy genes. ER translocation into autolysosomes and autolysosomal whorls were abolished in ATG2, ATG5, ATG7 and ATG8-hexa knockout cells (Fig.3F-3I, Extended Fig.3F-3G, 3M-3P), whereas ERphR-KO did not affect whorl formation (Fig.3J-3K, Extended Fig.3H-3I). Consistent with the genetic data, ER puncta in wild-type cells were positive for LC3 (Extended Fig.3J-3L), and notably, they exhibited strong ATG9A signal—distinct from conventional autophagosomes, which typically show weak ATG9A labeling[51,52]

(Fig.3L-3M). Triple labeling with BFP-LC3, mCherry-REEP5, and LAMP1-YFP revealed two classes of puncta: (i) REEP5$^+$/LC3$^+$/LAMP1$^-$ puncta representing free whorls, and (ii) REEP5$^+$/LC3$^+$/LAMP1$^+$ puncta corresponding to whorls enclosed within autolysosomes (Fig.3N).

APEX2 labeling confirmed these observations at the ultrastructural level, revealing LC3-positive autophagosomes, multilamellar whorls, and whorls enclosed within autolysosomes (Fig.3O-3P). Together, these results establish that core autophagy components are indispensable for whorl formation and turnover, whereas canonical ER-phagy receptors play only a minor role.

Based on these findings, we designate these structures "autolamellasomes", reflecting both their dependence on autophagy machinery and their distinctive multilamellar morphology. This nomenclature also distinguishes them from classical ER whorls generated under ER-stress conditions.

Autolamellasomes Form via Autophagy-Dependent Assembly of ER Fragments

We next sought to elucidate the cellular mechanism underlying autolamellasome formation. Reasoning that expanding the ER might enhance autolamellasome generation, we treated cells with oleic acid (OA), which is known to induce ER membrane expansion[53,54], followed by prolonged mTOR inhibition (Fig.4A). Unexpectedly, OA treatment alone caused the appearance of large ER patches in both wild-type and autophagy-deficient (ATG2, ATG5, and ATG7-deficient) cells (Fig.4B). Fluorescence recovery after photobleaching (FRAP) experiments showed rapid recovery of mCherry-REEP5 signal within these patches, indicating lateral diffusion of ER fragments across patch boundaries—a behavior reminiscent of liquid–liquid phase separation (Extended Fig.4A-4B).

When prolonged mTOR inhibition was applied, wild-type cells readily formed autolamellasomes (Fig.4B). Time-lapse imaging revealed that autolamellasomes emerged from ER patches and subsequently translocated into autolysosomes (Extended Fig.4C). In contrast, in autophagy-deficient cells, ER patches enlarged over time but failed to generate autolamellasomes. Interestingly, two distinct patterns were observed among the autophagy-deficient cell types: in ATG2- and ATG7-deficient cells, autolamellasome formation was completely blocked, whereas ATG5-deficient cells exhibited a moderate level of formation. OA treatment induced ER patch formation in most wild-type and ATG2-deficient cells, but only in a subset of ATG7- or ATG5-deficient cells. However, upon prolonged mTOR inhibition, nearly all autophagy-deficient cells displayed enlarged ER patches (Fig.4C-4D). These observations suggest that while ATG5 and ATG7 may negatively modulate OA-induced ER patch formation, the core autophagy machinery is required to convert ER patches into autolamellasomes.

To further characterize the nature of ER patches, we performed correlative light and electron microscopy (CLEM) in wild-type cells. CLEM revealed that these patches consist of clustered ER fragments (Fig.4E), consistent with their rapid fluorescence recovery after photobleaching. TEM analysis showed abundant autolamellasomes either free in the cytoplasm or enclosed within autolysosomes in wild-type cells, whereas autophagy-deficient cells accumulated disordered ER fragments instead (Fig.4F-4G). APEX2 labeling of SEC61B confirmed that in wild-type cells, autolamellasomes emerged directly from ER fragment clusters, reflecting the assembly of ordered multilamellar structures from disordered ER components. In contrast, no autolamellasome structures were detected in autophagy-deficient cells (Fig.4H-4I).

Together, these data demonstrate that autolamellasomes arise through the autophagy-dependent assembly of ER fragments, with the core autophagy machinery orchestrating their transformation from disordered ER clusters into organized multilamellar structures.

Basal Autolamellasome Formation Occurs Widely *In Vitro* and *In Vivo*

One of the defining ultrastructural features of lysosomes is the presence of internal membrane whorls. This observation prompted us to ask whether these whorls might originate from basal-level autolamellasome formation followed by lysosomal engulfment. To test this hypothesis, we performed TEM and fluorescence microscopy analyses across a wide range of immortalized cell lines, primary cultured cells, mouse zygotes, and multiple mouse organs.

We found that both free autolamellasomes and autolamellasomes enclosed within lysosomes are widely present under basal conditions. The abundance of these structures varied considerably among cell types and tissues (Fig.5A-5I, Extended Fig.5A-5G). For example, the number of autolamellasomes in U2OS cells was comparable to that observed in NRK cells after prolonged mTOR inhibition, while mouse zygotes displayed predominantly free autolamellasomes (Fig.5E-5F). This diversity highlights the dynamic basal turnover of ER-derived membranes across physiological contexts.

Interestingly, we later realized that our use of NRK cells for mTOR inhibition experiments was entirely serendipitous—chosen without prior knowledge that NRK cells exhibit the lowest basal level of autolamellasomes among the lines examined. This fortunate coincidence provided an unusually clean baseline for visualizing and quantifying induced autolamellasome formation.

We next examined whether basal autolamellasome formation depends on autophagy core genes. Under nutrient-rich conditions, ATG7-deficient cells showed a marked reduction in autolamellasome number, though the effect was less dramatic than under prolonged mTOR inhibition (Fig.5J-5L). This suggests that while autophagy contributes substantially, additional mechanisms may also promote low-level

autolamellasome formation in an ATG7-independent manner.

Because chloroquine (CQ) inhibits lysosomal degradation and, in some contexts, autophagosome-lysosome fusion[55], we next assessed its effect on autolamellasome formation. CQ treatment robustly induced autolamellasome formation, as confirmed by ER Tracker staining (Fig.5M-5N), TEM (Fig.5O-5P), cryo-tomography (Fig.5Q), and APEX2 labeling of SEC61B and LC3 (Fig.5R). In wild-type cells, CQ caused massive translocation of ER material into lysosomes, whereas in ATG2-, ATG5-, and ATG7- deficient cells, this translocation was strongly reduced. In these autophagy-deficient cells, lysosomes lacked intralysosomal whorls (Fig.5S-5W).

Together, these findings resolve a longstanding question in lysosome biology: the origin of intralysosomal whorls. Our results demonstrate that these whorls arise from basal autolamellasome formation and lysosomal incorporation, a constitutive process present across diverse cell types and organisms. The serendipitous use of NRK cells, with their exceptionally low basal activity, provided a natural "zero point" that made this discovery possible.

In Vitro Reconstitution of Autolamellasome and Autophagosome Budding

We next sought to establish a cell-free system capable of reconstituting autolamellasome budding *in vitro* in a manner dependent on autophagy core components. The assay was conceptually adapted from the classic COPII-budding assay developed by the Schekman and Novick laboratories, which uses permeabilized (semi-intact) cells to study vesicle formation[56,57]. However, instead of collecting released vesicles, we monitored the formation of autolamellasomes within permeabilized cells by transmission electron microscopy (TEM).

In this assay, cells were first pretreated with oleic acid (OA) for 24 hours to expand the

ER network. They were then permeabilized with digitonin in reconstitution buffer, supplemented with an S150 cytosolic fraction, ATP, and GTP, followed by incubation at 30 °C for 6-8 hours. Control samples received reconstitution buffer and digitonin alone (Fig.6A-6B). To ensure complete permeabilization and cell death, we stained samples with Apotracker, propidium iodide (PI), and Cy5. Apotracker labeling confirmed phosphatidylserine exposure on the plasma membrane outer leaflet, Cy5 positivity indicated membrane permeabilization, and PI nuclear staining verified loss of nuclear membrane integrity—together confirming that all treated cells were fully permeabilized and non-viable (Extended Fig.6A-6D, Extended Fig.6-6O).

To assess lysosomal integrity, we used Magic Red and Lysotracker staining. Both signals were abolished following digitonin treatment, indicating lysosomal disruption. Besides, microtubule integrity was also destroyed—a condition expected to facilitate the observation of free autolamellasomes (Extended Fig.6E-6H, 6J-6M).

TEM analysis revealed striking differences between control and reconstituted samples. In control preparations, internal membrane structures were extensively disrupted. In contrast, samples containing cytosol and nucleotides displayed numerous autolamellasomes at various stages of formation, as well as autophagosomes (Fig.6C-6E). Compared with TEM analysis of intact cells, the proportion of free autolamellasomes was substantially increased, suggesting efficient reconstitution of their formation within permeabilized cells.

We next examined the dependence on ATG7. Using ATG7-deficient cells as the membrane source, addition of wild-type S150 cytosol partially rescued autolamellasome formation to ~60 % of wild-type efficiency. Surprisingly, cytosol derived from ATG7-deficient cells still supported low-level formation—about one-third of wild-type levels—indicating that while ATG7 is indispensable for autolamellasome formation *in vivo*, limited assembly can occur *in vitro* (Fig.6F-6H). Together, these

results establish a foundational cell-free system that captures the essential process of

autolamellasome and autophagosome biogenesis. This *in vitro* approach reveals a core

mechanistic principle: the formation of these structures is driven by soluble cytosolic

factors acting upon permeabilized cellular membranes. Furthermore, the system

demonstrates that while the core component ATG7 is a major accelerator of

autolamellasome formation, the process retains a basal, ATG7-independent capacity.

This suggests the existence of either alternative, low-efficiency pathways or a residual

self-organizing potential within the membrane itself that can be co-opted by the cytosol

to initiate this form of membrane remodeling.

Induction of Autolamellasomes in Senescent Cells and Cells from Premature-

Aging Patients

Lysosomal activity is known to decline with age in both animals and humans[58]. To test

whether autolamellasomes accumulate in senescent cells, we employed two well-

established models of cellular senescence: doxorubicin-induced DNA damage[59] and

tert-butyl hydroperoxide (TBHP)-induced oxidative stress[60]. Senescence was

confirmed by SA–β-galactosidase staining, which showed robust activity in both

models (Fig.7A-C, Extended Fig.7B-7D).

Transmission electron microscopy (TEM) and fluorescence imaging revealed a

dramatic accumulation of autolamellasomes in senescent cells, whereas autophagy-

deficient cells failed to form autolamellasomes and instead accumulated large, irregular

ER patches (Fig.7F-7L, Extended Fig.7A, Extended Fig.7G-7M). Interestingly, mTOR

activity was largely suppressed in both senescence models (Fig.7D-E, Extended Fig.7E-

7F). Moreover, treatment with MHY1485—a small-molecule activator of mTOR—

effectively prevented ER translocation into autolysosomes (Fig.7M-7N). Collectively,

these findings mechanistically explain enhanced autolamellasome formation under

mTOR inhibition, suggesting that their accumulation in senescent cells results from

both reduced lysosomal degradation capacity and active induction via mTOR inhibition.

To further explore this phenomenon in the context of genetic premature aging, we

analyzed human dermal fibroblasts (HDFs) derived from Hutchinson-Gilford Progeria

Syndrome (HGPS) patients[61]—a rare laminopathy that causes severe premature aging

in children. Compared with HDFs from healthy individuals, HGPS fibroblasts

displayed markedly increased autolamellasome accumulation (Fig.7O-7R) and

reduction of mTOR activity (Fig.7S-7T). HGPS arises from a mutation in the LMNA

gene, which encodes lamin A; this mutation produces a truncated, farnesylated form

known as progerin, leading to nuclear envelope deformation and global cellular stress[62].

To directly test the role of progerin, we established a doxycycline-inducible progerin-

expressing cell line. Induction of progerin expression led to robust formation of

autolamellasomes, whereas this response was completely abolished in autophagy-

deficient cells, confirming that autolamellasome formation is autophagy-dependent

(Fig.7U-7W, Extended Fig.7N).

Together, these findings position autolamellasome accumulation as a conserved cellular

signature of aging, emerging at the intersection of declining lysosomal function and

persistent ER stress. Rather than a passive breakdown, this process represents an active,

autophagy-dependent remodeling pathway, triggered by the near-universal age-

associated suppression of mTOR. This pathway thus links core hallmarks of aging—

dysfunctional organelles and nutrient signaling collapse—into a unified, structural

adaptation.

Discussion

This study uncovers a previously unrecognized form of ER remodeling that operates

under conditions of chronic mTOR inhibition. We identify autolamellasomes,

multilamellar ER-derived structures that arise through the core autophagy machinery

but independently of canonical ER-phagy receptors. Unlike conventional autophagosomes, autolamellasomes mediate bulk ER degradation via autophagy-dependent assembly and lysosomal incorporation of compacted ER fragments. Their presence across multiple cell types, and their accumulation in senescent and HGPS fibroblasts, reveal a conserved, aging-associated remodeling pathway linking sustained mTOR suppression to ER and lysosomal homeostasis. These findings redefine how long-term autophagy activation reshapes cellular architecture, providing a structural and mechanistic framework that connects nutrient-sensing, membrane dynamics, and aging.

Beyond revealing a new mode of ER turnover, our findings also resolve a long-standing question in cell biology—the origin of the membrane whorls long observed within lysosomes. These enigmatic multilamellar structures, variably termed myelin figures or zebra bodies, have been described for decades in electron microscopy studies but lacked a defined biogenic pathway. Our data demonstrate that these intralysosomal whorls originate from basal autolamellosome formation, followed by lysosomal incorporation. Importantly, autolamellasomes are not restricted to experimental mTOR inhibition but are constitutively generated at low levels across a wide range of cell types and tissues, from cultured mammalian cells to mouse embryos. This basal activity likely represents a housekeeping mechanism that continuously remodels the ER and contributes to the steady-state membrane flux through lysosomes. Thus, the identification of autolamellasomes provides a unifying explanation for the biogenesis of lysosomal membrane whorls and establishes a physiological link between ER dynamics, lysosome structure, and long-term membrane homeostasis.

The autolamellosome pathway is mechanistically distinct from previously described ER-derived whorls in both yeast and mammalian systems. In *Saccharomyces cerevisiae*, ER whorls form rapidly under acute ER stress and are degraded through a specialized micro-ER-phagy route. This process involves direct invagination of the vacuolar

membrane, followed by ESCRT-mediated scission to internalize the whorl. It proceeds independently of the core ATG machinery and canonical ER-phagy receptors, relying instead on vacuolar sorting components and the ER-localized Nem1-Spo7 phosphatase complex, which regulates phosphatidic acid turnover and membrane curvature. In contrast, the ER-stress-induced whorls in mammalian cells—large, concentric stacks up to several micrometers in diameter—arise through a PERK- and COPII-dependent remodeling program that serves an adaptive function: concentrating PERK and sequestering translocons to transiently dampen protein import during stress. These structures are not degraded via autophagy, but rather disassembled upon stress resolution, and their biogenesis is tightly coupled to the unfolded protein response (UPR) signaling network. By contrast, autolamellosomes emerge under sustained mTOR inhibition and depend on the core autophagy machinery but not ER-phagy receptors or UPR signaling. They represent a degradative remodeling route that links nutrient sensing to long-term membrane recycling, distinguishing them from both the ESCRT-dependent ER whorl degradation in yeast and PERK-driven ER whorl formation in mammalian.

A key advance of this study is the establishment of a cell-free reconstitution system that faithfully recapitulates autolamellasome formation *in vitro*. This approach demonstrates that multilamellar ER structures can self-organize from cellular membranes in response to soluble cytosolic factors and energy cofactors, revealing that the core autophagy machinery is sufficient to drive their assembly. The significance of this finding extends far beyond the specific phenomenon described here: cell-free reconstitution has historically been pivotal in cell biology, transforming abstract cellular processes into experimentally tractable biochemical reactions. Seminal examples include the reconstitution of COPII- and COPI-coated vesicle budding from the ER and Golgi membranes—discoveries that defined the molecular logic of vesicular trafficking. By establishing a comparable system for autolamellosome biogenesis, our work brings this newly identified pathway into the same mechanistic framework, enabling its dissection at molecular resolution. This cell-free system provides not only proof of principle that

autolamellosome formation is a bona fide, self-organizing process but also a powerful platform for future biochemical and structural reconstitution of autophagy-driven membrane remodeling.

Despite elucidating the existence and mechanistic basis of autolamellosome formation, our study has several important limitations that point to future directions. The specific molecular components that distinguish this pathway from canonical autophagy remain unknown. Although the core autophagy machinery is essential, additional factors likely confer the multilamellar morphology and ER specificity of autolamellosomes. A comprehensive whole-genome CRISPR or siRNA screen will be critical to identify such unique regulators, including membrane-shaping or lipid-metabolizing proteins. Moreover, the physiological significance of autolamellosomes requires validation *in vivo*. Developing animal models that allow controlled modulation of autolamellosome formation will enable investigation of their roles in organ function, stress adaptation, and organismal physiology. Given their capacity for extensive membrane remodeling and lipid catabolism, autolamellosomes may serve as an adaptive mechanism for energy recycling during prolonged nutrient limitation. Their pronounced accumulation in senescent and progeroid cells further suggests that this pathway participates in the cellular response to aging-associated mTOR suppression and lysosomal decline. Because chronic mTOR dysregulation, ER stress, and impaired lipid homeostasis are hallmarks of diverse human disorders—including metabolic diseases, neurodegeneration, and cancer—autolamellosome formation may represent a broader protective or maladaptive remodeling response in these contexts. Future studies integrating unbiased genetic screening, *in vitro* reconstitution, disease modeling, and physiological analyses will be essential to define how autolamellosomes contribute to membrane homeostasis, metabolic resilience, aging, and disease pathogenesis.

Figures and figure legends

Figure 1 Prolonged mTOR inhibition induces the formation of "free whorls".

(A) Representative TEM images of NRK cells treated with or without 1 µM Torin1 for

24 h. Red arrows indicate electron-dense whorls. ER: the endoplasmic reticulum; mt:

mitochondria; LY: lysosome; Golgi: Golgi apparatus. Scale bars: 5 µm (overview) or

500 nm (magnified view).

(B) Representative TEM images of autophagosome (AP), free whorls (FW) at different stages and lysosome/autolysosome containing with whorls (Whorled lysosome). NRK cells were treated with 1 μM Torin1 for a time series. Scale bars, 200 nm.

(C) TEM images from (B) were quantified for the amount of each kind of structure (FW: free whorls, intraLY-W: intralysosomal whorls, AP: autophagosome) in 15 μm^2 area. For each time point, more than 38 cells were imaged. Data are reported as mean ± SEM.

(D) TEM images from (B) were quantified for the number of whorls contained in each lysosome/autolysosome. For each time point, more than 70 lysosomes/ autolysosomes were quantified. Data are presented as violin plots showing the full distribution, with the mean indicated by a triangle.

(E) Representative cryo-ET (cryo-electron tomography) images and 3D rendered images of free whorl (FW) and lysosome/autolysosome containing with whorls (Whorled lysosome) in NRK cells treated with 1 μM Torin1 for 24 h. L1-L3 indicated different layers of the structure. Scale bars, 100 nm.

(F) Lumen diameter of the whorls from (E) were quantified. Data are reported as mean ± SEM. n=34 for FW group, n=48 for intraLY-W group. ns, not significant. P values were calculated using a two-tailed, unpaired t test. Asterisks indicated the lumens of whorls.

(G) Representative FIB (focused ion beam) images of free whorl (FW) and lysosome/autolysosome containing with whorls (Whorled lysosome). L1-L5 indicated different layers of the structure.

(H) Quantification of free whorls (FW) and intralysosomal whorls (intraLY-W) from TEM images of multiple cell lines treated with 1 μM Torin1 for 24 h. M, mock; T, Torin1. Sample sizes (n) were as follows: L929 (M: n=40; T: n=68), MEF (M: n=78; T: n=56), 4T1 (M: n=54; T: n=34), THP1 (M: n=43; T: n=46), HepG2 (M: n=74; T: n=54), 293FT (M: n=32; T: n=29), MGC803 (M: n=35; T: n=38), COS7 (M: n=64; T: n=49), and S2 (M: n=25; T: n=31).

607 **Figure 2 Whorls originate from the Endoplasmic Reticulum.**

608 (A) Representative confocal images of NRK cells treated with or without 1 µM Torin1

609 for 24 h. NRK cells were labeled by ER Tracker. Arrows indicate ER puncta. Scale

610 bars, 5 µm.

611 (B) Quantification of the ratio of cells shown in (A) contained with ER puncta. More

than 230 cells were analyzed per group in each experiment (n=3 experiments). Error bars indicate mean ± SEM. ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(C) Representative SIM images of NRK cells stably co-expressing LAMP1-YFP and mCherry-REEP5 treated with or without 1 μM Torin1 for 24 h. White arrows: free-standing ER puncta, yellow arrows: ER puncta inside lysosomes. Scale bars, 5 μm.

(D) Representative z-stack image and its 3D rendered image of cells from (C). Scale bars, 5 μm.

(E-F) Quantification of the cells shown in (B). (E) Ratio of cells with mCherry-REEP5 labeled-ER puncta. (F) Pearson correlation coefficient of LAMP1-YFP and mCherry-REEP5 signals. More than 135 cells were analyzed per group in each experiment (n=3 experiments). Error bars indicate mean ± SEM. ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(G) Endogenous immunostaining of NRK cells treated with or without 1 μM Torin1 for 24 h. Cells were stained with anti-LAMP1/anti-SEC61B and anti-LAMP1/anti-REEP5 antibodies. White arrows: free-standing ER puncta, yellow arrows: ER puncta inside lysosomes. Scale bars, 5 μm.

(H-K) Quantification of the cells shown in (G). (H) Ratio of cells with SEC61B labeled-ER puncta. (I) Pearson correlation coefficient of LAMP1 and SEC61B signals. (J) Ratio of cells with REEP5 labeled-ER puncta. (K) Pearson correlation coefficient of LAMP1 and REEP5 signals. More than 180 cells were analyzed per condition. Error bars indicate mean ± SEM. *, p<0.05, ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(L) Representative APEX2-TEM images of free whorls in NRK cells stably expressing APEX2-EGFP-SEC61B or APEX2-EGFP-REEP5 treated with Torin1. Scale bar, 200 nm.

(M) CLEM data of NRK cells stably co-expressing LAMP1-YFP and mCherry-REEP5 treated with Torin1. The marked region in the fluorescent graph stacked with the electron microscope graph was enlarged and demonstrated on the right. Scale bars

denoted 10 μm for the stacked graph, 2 μm for the electron microscope graph.

(N) Time-lapse live cell images of NRK cells stably co-expressing LAMP1-mStaygold

and HaloTag-REEP5 treated with Torin1. Here shows representative snapshots at

indicated time points. Scale bar, 5 μm.

(O) Western blot results of NRK cells treated with or without 1 μM Torin1 for 24 h.

Cell lysates were subjected to SDS-PAGE and immunoblotted with the indicated

antibodies. Blots are representative of n=5 independent biological replicates.

(P) Quantification of integrated band intensities from (O). Individual data points

represent relative protein levels in the Torin1-treated group normalized to the untreated

control and loading control Actin. n=5 independent biological repeats. Data are

presented as mean ± SEM.

(Q) Heatmap of relative protein abundances measured by label-free mass spectrometry

in NRK cells treated with or without 1 μM Torin1 for 24 h. Values were normalized to

the mock-1 group. n=3 independent biological repeats. Color scale indicates fold

change relative to mock-1 group.

(R) Western blot results of NRK cells treated with 1 μM Torin1 in a time series. Cell

lysates were subjected to SDS-PAGE and immunoblotted with the indicated antibodies.

Blots are representative of n=4 independent biological replicates.

(S) Quantification of integrated band intensities from (R). Individual data points

represent relative protein levels in the time point normalized to 0 h and loading control

Actin. n=4 independent biological repeats. Data are presented as mean ± SEM.

(T) Schematic diagram illustrating that mTOR inhibition stimulates parallel formation

of autophagosomes and free whorls.

Figure 3 Formation of whorls depends on the core autophagy machinery but not

ER-phagy receptors.

(A) Table of knockout genes and their effects on whorls formation. *ERphR*: ER phagy

receptor penta KO. ↓ slightly decrease，↓↓ moderated decease, ↓↓↓ almost disappear.

(B) Western blot analysis of endogenous proteins and mCherry cleavage in WT and

autophagy-deficient NRK cells. WT, *Atg2* KO, *Atg5* KO, *Atg7* KO NRK cells were treated with or without 1 μM Torin1 for 24 h. Cell lysates were subjected to SDS-PAGE and immunoblotted with the indicated antibodies. Asterisks denote cleaved mCherry fragments generated upon autophagic degradation. Actin served as a loading control. Blots are representative of n=3 independent biological replicates.

(C) Quantification of mCherry-REEP5 cleavage. The percentage of cleaved mCherry fragments relative to full-length mCherry-REEP5 was quantified from the Western blot shown in (B). Data are presented as mean ± SEM from n=3 independent biological replicates.

(D) Western blot analysis of *ERphR* KO (ER-phagy receptor penta-KO) clones. Wild-type (WT) and three independent *ERphR* KO NRK cell clones (#9, #19, #26) were treated with or without 1 μM Torin1 for 24 h. Cell lysates were subjected to SDS-PAGE and immunoblotted with the indicated antibodies. Blots are representative of n=3 independent biological replicates.

(E) Quantification of integrated band intensities from (D). Individual data points represent relative protein levels in the Torin1-treated group normalized to the untreated control and loading control Actin. n=3 independent biological repeats. Data are presented as mean ± SEM.

(F) Wild-type (WT) and autophagy-deficient NRK cells (*Atg2* KO, *Atg5* KO, or *Atg7* KO, as indicated) stably co-expressing LAMP1-YFP and mCherry-REEP5 were treated with 1 μM Torin1 for 24 h. Cells were imaged by Structured Illumination Microscopy (SIM). Scale bars, 5 μm.

(G) Pearson correlation coefficient of LAMP1 and REEP5 signals of the cells shown in (F). More than 300 cells were analyzed for each group (n=3 independent biological repeats). Error bars indicate mean ± SEM. ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(H) Representative TEM images of WT and autophagy-deficient (*Atg2* KO, *Atg5* KO, or *Atg7* KO, as indicated) NRK cells treated with 1 μM Torin1 for 24 h. Scale bars, 200 nm.

(I) TEM images from (H) were quantified for the amount of each kind of structure (AL: autolamellasome, LYAL: lysosome/autolysosome containing with autolamellasome) in 15 μm^2 area. More than 37 cells were imaged for each group. Data are reported as mean ± SEM.

(J) Representative TEM images of WT NRK and *ERphR* KO clones treated with 1 μM Torin1 for 24 h. Scale bars, 200 nm.

(K) TEM images from (J) were quantified for the amount of each kind of structure (AL: autolamellasome, LYAL: lysosome/autolysosome containing with autolamellasome) in 15 μm^2 area. More than 43 cells were imaged for each group. Data are reported as mean ± SEM.

(L) Representative confocal images of NRK cells stably co-expressing EGFP-ATG9A and mCherry-REEP5 treated with or without 1 μM Torin1 for 24 h. Scale bar, 5 μm.

(M) Pearson correlation coefficient of ATG9A and REEP5 signals of the cells shown in (L). More than 150 cells were analyzed for each group. Error bars indicate mean ± SEM. ****, $p < 0.0001$. P values were calculated using a two-tailed, unpaired t test.

(N) Representative confocal images of NRK cells stably co-expressing BFP-LC3, mCherry-REEP5 and LAMP1-YFP treated with Torin1. (i) Representative images of BFP-LC3$^+$/mCherry-REEP5$^+$/LAMP1-YFP$^-$ puncta, corresponding to LC3 labeled free ER. (ii) Representative images of BFP-LC3$^+$/mCherry-REEP5$^+$/LAMP1-YFP$^+$ ER puncta, indicating ER in lysosomes.

(O) Representative APEX2-TEM images of cells stably expressing APEX2-mCherry-LC3 NRK cells treated with 1 μM Torin1. APEX2 labeled autophagosome (AP), the endoplasmic reticulum (ER), autolamellasome (AL) are shown here as indicated. Scale bars, 200 nm.

(P) NRK cells stably expressing APEX2-mCherry-LC3 were treated with 1 μM Torin1 and processed for APEX2-TEM at the indicated time points. Red arrows mark LC3-positive structures. Scale bars, 500 nm.

(Q) Schematic diagram illustrating that autophagy genes (ATGs) are key regulators of autolamellasome formation.

Figure 4 Autolamellasome formation via autophagy-dependent assembly of ER fragments

(A) Schematic diagram of the sequential drug treatment protocol in NRK cells. NRK cells were primed with 200 μM oleic acid (OA) for the initial incubation period, followed by direct supplementation with 1 μM Torin1 for further treatment (OAT).

(B) Wild-type (WT) and autophagy-deficient NRK cells (*Atg2* KO, *Atg5* KO, or *Atg7* KO, as indicated) stably co-expressing LAMP1-YFP and mCherry-REEP5 were treated with 200 µM oleic acid (OA) for 24 h, or OA followed by 1 µM Torin1. Cells were imaged by confocal microscopy. Arrows indicate ER puncta while asterisks indicate ER patches. Scale bar, 5 µm.

(C-D) Quantification of cells from (B). (C) Statistics of the ratio of cells with ER puncta. (D) Statistics of the ratio of cells with ER patches. More than 70 cells were analyzed for each group (n=3 independent biological repeats). Error bars indicate mean ± SEM.

(E) CLEM data of NRK-*Atg7*-KO cells stably expressing LAMP1-YFP/mCherry-REEP5 treated with OAT. The marked region was enlarged and demonstrated on the right. Scale bars denoted 5 µm for the fluorescence graph, 1 µm for the electron microscope graph.

(F) Representative TEM images of WT and autophagy-deficient NRK cells treated OAT. Scale bar, 500 nm.

(G) TEM images from (F) were quantified for the amount of each kind of structure (AL: autolamellasome, LYAL: lysosome/autolysosome containing with autolamellasome) in 15 µm^2 area. More than 60 cells were imaged for each group. Data are reported as mean ± SEM.

(H) Representative APEX2-TEM images of WT and autophagy-deficient NRK cells stably expressing APEX2-EGFP-SEC61B and treated with OAT. Red dashed lines indicate clusters of ER fragments. Red arrow indicates autolamellasome, yellow arrow indicates ER fragments. Scale bars, 500 nm.

(I) TEM images from (H) were quantified for APEX2-postive structures (AL: autolamellasome, LYAL: lysosome/autolysosome containing with autolamellasome) in 35 µm^2 area. More than 50 cells were imaged for each group. Data are reported as mean ± SEM.

(J) Schematic diagram illustrating oleic acid (OA)-induced clustering of ER fragments and subsequent Torin1-driven, autophagy-dependent autolamellasome formation.

Figure 5 Widespread basal autolamellasome formation *in vitro* and *in vivo*.

(A) Quantification of AL (autolamellasome) and LYAL (lysosome/autolysosome containing with autolamellasome) number per 15 μm² area from TEM images for each indicated cell line. More than 30 cells were analyzed per group. Data are reported as mean ± SEM.

(B) Representative TEM images of AL (autolamellasome) and LYAL (lysosome/autolysosome containing with autolamellasome) in HeLa cells. Scale bars, 200 nm.

(C) HeLa cells were co-stained with anti-LAMP1/anti-SEC61B antibodies. White arrow: free-standing ER puncta, yellow arrow: ER puncta inside lysosomes. Scale bars: 5 μm (main image) and 2 μm (magnified view).

(D) Representative cryo-ET images and 3D rendered images of AL (autolamellasome) and LYAL (lysosome/autolysosome containing with autolamellasome) in HeLa cells. L1-L3 indicated different layers of the structure. Scale bars, 100 nm.

(E) Representative TEM images of AL (autolamellasome) and LYAL (lysosome/autolysosome containing with autolamellasome) in mouse zygotes. Scale bars, 200 nm.

(F) Quantification of AL (autolamellasome) and LYAL (lysosome/autolysosome containing with autolamellasome) number per 35 μm^2 area from (E). Mouse zygotes were collected at 3, 7, and 12 hours and imaged. n = 3 (z3h), n = 3 (z7h), and n = 2 (z12h). Data are reported as mean ± SEM.

(G) Quantification of AL (autolamellasome) and LYAL (lysosome/autolysosome containing with autolamellasome) number per 65 μm^2 area from TEM images for each indicated primary cell. More than 35 cells were analyzed per group. Data are reported as mean ± SEM.

(H) Representative TEM images of AL (autolamellasome) and LYAL (lysosome/autolysosome containing with autolamellasome) in primary MEF cells. Scale bars, 200 nm.

(I) Primary MEF cells were co-stained with anti-LAMP1/anti-REEP5 antibodies. White arrow: free-standing ER puncta, yellow arrow: ER puncta inside lysosomes. Scale bars: 10 μm (main image) and 2 μm (magnified view).

(J) Representative TEM images of WT and ATG7 KO HeLa cells. Scale bars, 500 nm.

(K) Validation of knockout efficiency of ATG7 KO HeLa cells via Western blot.

(L) Quantification of AL (autolamellasome) and LYAL (lysosome/autolysosome

containing with autolamellasome) number per 35 μm^2 area from (J). More than 50 cells were analyzed per group. Data are reported as mean ± SEM.

(M) Representative confocal images of NRK cells treated with or without 100 nM chloroquine (CQ) for 18 h. Arrow indicates ER Tracker-labeled puncta. Scale bar, 5 μm.

(N) Quantification of the number of ER puncta per cell in cells shown in (M). More than 120 cells were analyzed per group. Error bars indicate mean ± SEM. ****, $p < 0.0001$. P values were calculated using a two-tailed, unpaired t test.

(O) Representative TEM images of NRK cells treated with 100 nM chloroquine (CQ) for 18 h. Scale bar, 200 nm.

(P) Quantification of AL (autolamellasome) and LYAL (lysosome/autolysosome containing with autolamellasome) number per 15 μm^2 area from (O). More than 65 cells were analyzed per group. Data are reported as mean ± SEM.

(Q) Representative cryo-ET (cryo-electron tomography) images and 3D rendered images of autolamellasome (AL) and lysosome/autolysosome containing with autolamellasome (LYAL) in NRK cells treated with 100 nM chloroquine (CQ) for 18 h. L1-L3 indicated different layers of the structure. Scale bars, 100 nm.

(R) Representative APEX2-TEM images of autolamellasomes in NRK cells stably expressing APEX2-EGFP-SEC61B or APEX2-mCherry-LC3. Cells were treated with 100 nM chloroquine (CQ) for 18 h. Scale bars, 200 nm.

(S) Representative SIM images of WT and autophagy-deficient NRK cells stably co-expressing LAMP1-YFP and mCherry-REEP5. Cells were treated with 100 nM CQ for 18 h. Scale bars, 5 μm.

(T-U) Quantification of the cells shown in (S). (T) Pearson correlation coefficient of LAMP1 and REEP5 signals. (U) Ratio of cells with ER puncta. More than 140 cells were analyzed in each group for each experiment (n=3 independent biological repeats). Error bars indicate mean ± SEM. ****, $p < 0.0001$. P values were calculated using a two-tailed, unpaired t test.

(V) Representative TEM images of WT and autophagy-deficient NRK cells stably co-

826   expressing LAMP1-YFP/mCherry-REEP5. Cells were treated with 100 nM CQ for 18

827   h. Blue arrow: autolamellasomes. Red arrows: lysosomes. Scale bar, 500 nm.

828   (W) Quantification of AL (autolamellasome) and LYAL (lysosome/autolysosome

829   containing with autolamellasome) number per 35 μm^2 area from (V). More than 40 cells

830   were analyzed per group. Data are reported as mean ± SEM.

Figure 6 *In vitro* reconstitution of autolamellasome.

(A) Schematic diagram illustrating the process of *in vitro* reconstitution of autolamellasome. NRK cells were primmed with 200 µM OA for 20-24 h, followed by incubation in reconstitution buffer containing 40 µg/mL digitonin (mock group) or supplemented with S150, ATP and GTP (all group) for 6-8 h at 30 °C.

(B) Table of various protocols for *in vitro* autolamellasome reconstitution and their

efficiencies.

(C) Representative TEM images of membrane structures in mock group and all group

from protocol A. Red arrows indicated autollamellasome-like structures. Scale bar=5

μm for large view. Scale bar=500 nm for magnified view.

(D) Representative TEM images of autollamellasomes, autollamellasome precursors

and autophagosomes in all group. Scale bars, 200 nm.

(E) Quantification of sealed mature autolamellasomes (Mature) and unsealed

autolamellasome precursors (Precursor) in the *in vitro* reconstitution system using

different protocols. Protocol A: n=3 independent biological repeats, >35 fields of view

per group per experiment. Protocols B, C, and D: >30, >50, and >40 fields of view per

group, respectively. Data are presented as mean ± SEM.

(F) Representative TEM images of membrane structures reconstituted with *Atg7*

knockout membranes (protocol A). Conditions: I, reconstitution buffer containing 40

μg/mL digitonin; II: buffer (reconstitution buffer containing 40 μg/mL digitonin)

supplemented with *Atg7* knockout S150, ATP, and GTP; III: buffer supplemented with

WT S150, ATP, and GTP. Red arrows indicate autolamellasomes. Boxed regions are

magnified and demonstrated on the right. Scale bar, 500 nm.

(G) Western blot analysis of extracted S150 from WT and *Atg7* knockout NRK cells.

(H) Quantification of sealed mature autollamellasomes (Mature) and unsealed

autollamellasome precursors from the *in vitro* reconstitution system in (F). More than

45 fields of view were analyzed per group. Data are presented as mean ± SEM.

Figure 7 Autolamellasomes increased in senescent cells and progeroid cells.

(A) Schematic diagram illustrating the process of Doxorubicin-induced cellular

senescence.

(B) β-galactosidase staining results of WT and autophagy-deficient NRK cells. Scale

bar, 100 pixels.

(C) Quantification of β-galactosidase-positive cell ratios from (B). More than 45 cells were analyzed per group. Data are presented as mean ± SEM. ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(D) Western blot results of NRK cells treated with or without 400 nM Doxorubicin (DOXO) for 2 days. Cell lysates were subjected to SDS-PAGE and immunoblotted with the indicated antibodies. GAPDH served as a loading control. Blots are representative of n=3 independent biological replicates.

(E) Quantification of phospho-p70 S6K (T389) band intensities from (D). Individual data points represent values normalized to total p70 S6K and expressed relative to the untreated control. n=3 independent biological repeats. Data are presented as mean ± SEM.

(F) Representative confocal (upper layer) and SIM images (lower layer) of Doxorubicin (DOXO)-treated WT and autophagy-deficient NRK cells. Arrows indicate ER puncta; asterisks indicate ER patches. Scale bars: 10 μm (confocal), 5 μm (SIM).

(G-I) Statistics of cells from (F). (G) Pearson correlation coefficient of LAMP1 and REEP5 signals. (H) Ratio of cells with ER puncta. (I) Ratio of cells with ER patches. More than 49 cells were analyzed in each group for each experiment (n=3 independent biological repeats). Data are presented as mean ± SEM. ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(J) Representative TEM images of Doxorubicin (DOXO)-treated WT and autophagy-deficient NRK cells. Red arrows indicate autolamellasomes (AL), yellow arrow indicates lysosomes containing with autolamellasomes (LYAL). Scale bar, 200 nm.

(K) TEM images from (J) were quantified for the amount of each kind of structure (AL: autolamellasome, LYAL: lysosome/autolysosome containing with autolamellasome) in 15 μm^2 area. More than 39 cells were imaged for each group. Data are reported as mean ± SEM.

(L) Quantification of frequency of existence of ER patch per 65 μm^2 area from TEM images in (J).

(M) Representative confocal images of NRK cells treated with or without 400 nM

Doxorubicin (DOXO), 400 nM DOXO together with 10 μM MHY1485 for 2 days.

Scale bars, 5 μm.

(N) Quantification of the Pearson correlation coefficient of LAMP1 and REEP5 signals

from (M). More than 130 cells were analyzed in each group for each experiment (n=3

independent biological repeats). Data are presented as mean ± SEM. ****, p<0.0001.

P values were calculated using a two-tailed, unpaired t test.

(O) Representative confocal images of ER Tracker-labeled HDF (human dermal

fibroblasts) from normal donors and HGPS (Hutchinson-Gilford progeria syndrome)

patients. Scale bars, 10 μm.

(P) Quantification of the ratio of cells with ER puncta from (O). Normal HDF from 2

healthy donors and HGPS HDF form 2 HGPS patients were acquired. More than 225

cells were analyzed for each group. Data are reported as mean ± SEM. ****, p<0.0001.

P values were calculated using a two-tailed, unpaired t test.

(Q) Representative TEM images of Normal HDF and HGPS HDF. Red arrow indicates

autolamellasome (AL), yellow arrows indicate lysosome/autolysosome containing with

autolamellasomes (LYAL). Scale bars, 200 nm.

(R) TEM images from (Q) were quantified for the amount of each kind of structure (AL:

autolamellasome, LYAL: lysosome/autolysosome containing with autolamellasome) in

15 μm^2 area. More than 65 areas were imaged for each group. Data are reported as mean

± SEM.

(S) Western blot results of indicated HDF. Cell lysates were subjected to SDS-PAGE

and immunoblotted with the indicated antibodies. GAPDH served as a loading control.

(T) Phospho-p70 S6K (T389) and phospho-p70 S6K (T229) band intensities from (S)

were normalized to total p70 S6K and expressed relative to Normal-1.

(U) Schematic diagram illustrating doxycycline (Dox) inducible expression of EGFP-

Progerin in NRK cells.

(V) Representative TEM images of AL (autolamellasome) and LYAL

(lysosome/autolysosome containing with autolamellasome) in doxycycline (Dox)-

induced EGFP-Progerin NRK cells. Scale bars, 200 nm.

(W) Quantification of AL (autolamellasome) and LYAL (lysosome/autolysosome

containing autolamellasomes) number per 15 μm^2 area from TEM images from (V).

WT and autophagy-deficient NRK cells were treated with doxycycline (Dox) for 0, 5,

and 10 days (D0, D5, D10). More than 60 areas were imaged for each group. Data are

reported as mean ± SEM.

Extended figures and figure legends

Extended Data Fig. 1. Free whorl formation is a widespread phenomenon.

(A) Representative TEM images of 293FT cells treated with or without 1 μM Torin1

for 24 h. Scale bar, 500 nm.

(B) Representative TEM images of free whorls in various cells treated with 1 μM

Torin1 for 24 h. Scale bars, 200 nm.

Extended Data Fig. 2. Whorls originate from the endoplasmic reticulum.

(A) Representative confocal images of NRK cells stably expressing EGFP-REEP5
treated with or without 1 μM Torin1 for 24 h. Arrows indicate EGFP-REEP5 labeled-
ER puncta. Scale bar, 5 μm.

(B) Quantification of the ratio of cells shown in (A) contained with ER puncta. More

than 280 cells were analyzed per group in each experiment (n=3 experiments). Error bars indicate mean ± SEM. ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(C) Representative confocal images of NRK cells stably expressing LAMP1-mStaygold/HaloTag-REEP5 treated with or without 1 μM Torin1 for 24 h. Scale bar, 5 μm.

(D-E) Quantification of the cells shown in (C). (D) Pearson correlation coefficient of LAMP1-mStaygold and HaloTag-REEP5 signals. (E) Ratio of cells with HaloTag-REEP5 labeled-ER puncta. More than 280 cells were analyzed per group in each experiment (n=3 experiments). Error bars indicate mean ± SEM. ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(F) Representative confocal images of NRK cells stably expressing LAMP1-mStaygold/Calnexin-HaloTag, LAMP1-YFP/mCherry-SEC61B, LAMP1-YFP/ER-DsRed treated with or without 1 μM Torin1 for 24 h. Scale bar, 5 μm.

(G-H) Quantification of the cells shown in (F). (G) Pearson correlation coefficient of LAMP1 and ER proteins. (H) Ratio of cells with ER puncta. Sample sizes (n) were n>90 (Calnexin), n>100 (SEC61B), and n>50 (ER-DsRed) cells per group in each experiment (n=3 experiments). Error bars indicate mean ± SEM. ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(I-J) Quantification of NRK cells stably co-expressing LAMP1-YFP and mCherry-REEP5 treated with 1 μM Torin1 in a time series. (I) Pearson correlation coefficient of LAMP1 and REEP5 signals. (J) Ratio of cells with ER puncta. More than 85 cells were analyzed in each time point. Error bars indicate mean ± SEM. ns, not significant, **, p<0.01, ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(K-L) NRK cells stably expressing LAMP1-YFP and mCherry-REEP5 were treated with Torin1 at the indicated concentrations for 9 h. (K) Pearson correlation coefficient of LAMP1 and REEP5 signals. (L) Ratio of cells exhibiting ER puncta. More than 160 cells were analyzed per dose. Data are presented as mean ± SEM. ns, not significant, *, p<0.05, ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(M) APEX2-TEM images of cells stably expressing APEX2-EGFP-SEC61B or APEX2-EGFP-REEP5. In H_2O_2-treated groups, APEX2 catalyzes DAB polymerization, generating electron-dense structures that reveal protein localization. In the absence of H_2O_2, cells exhibit low electron contrast. Scale bars, 500 nm.

(N) Representative SIM images of NRK cells stably co-expressing LAMP1-YFP/mCherry-REEP5. Cells were treated with 1 µM rapamycin or serum-starved (FBS withdrawal) for 24 h, alongside untreated controls. White arrows: free-standing ER puncta, yellow arrows: ER puncta inside lysosomes. Scale bars, 5 µm.

(O-P) Quantification of the cells shown in (N). (O) Pearson correlation coefficient of LAMP1 and REEP5 signals. (P) Ratio of cells with ER puncta. More than 180 cells were analyzed in each group. Error bars indicate mean ± SEM. ****, $p<0.0001$. P values were calculated using a two-tailed, unpaired t test.

(Q) Representative TEM images of free whorls (FW) and intralysosomal whorls (intraLY-W) in NRK cells treated with 1 µM rapamycin or serum-starved (FBS withdrawal) for 24 h. Scale bar, 200 nm.

(R) TEM images from (Q) were quantified for the amount of each kind of structure (FW: free whorls, intraLY-W: intralysosomal whorls) in 15 μm^2 area. More than 70 cells were imaged for each group. Data are reported as mean ± SEM.

Extended Data Fig. 3. Autolamellasome formation depends on autophagy core machinery but not ER phagy receptors.

(A) Western blot analysis of endogenous proteins and mCherry cleavage in WT and ER phagy receptor KO NRK cells. WT, *Fam134b* KO, *Sec62* KO, *Tex264* KO, *Atl3* KO and *Ccpg1* KO NRK cells were treated with or without 1 µM Torin1 for 24 h. Cell lysates were subjected to SDS-PAGE and immunoblotted with the indicated antibodies. Asterisks denote cleaved mCherry fragments generated upon autophagic degradation. Actin served as a loading control. Blots are representative of n=3 independent biological replicates.

(B-C) Quantification of mCherry-REEP5 cleavage. The percentage of cleaved mCherry fragments relative to full-length mCherry-REEP5 was quantified from the Western blot

shown in (A). Data are presented as mean ± SEM from n=3 independent biological replicates.

(D) Western blot analysis of endogenous proteins in WT, *Atg7* KO, *Fam134b* KO and *Tex264* KO NRK cells treated with or without 1 μM Torin1 for 24 h. Cell lysates were subjected to SDS-PAGE and immunoblotted with the indicated antibodies. Actin served as a loading control. Blots are representative of n=3 independent biological replicates.

(E) Quantification of integrated band intensities from (D). Individual data points represent relative protein levels in the Torin1-treated group normalized to the untreated control and loading control Actin. n=3 independent biological repeats. Data are presented as mean ± SEM.

(F) Representative confocal images of WT and autophagy-deficient KO (*Atg2* KO, *Atg5* KO, *Atg7* KO) NRK cells treated with 1 μM Torin1 for 24 h. NRK cells were labeled by ER Tracker. Arrows indicate ER puncta. Scale bars, 5 μm.

(G) Quantification of the ratio of cells shown in (F) contained with ER puncta. More than 250 cells were analyzed per group in each experiment (n=3 experiments). Error bars indicate mean ± SEM. ****, $p < 0.0001$. P values were calculated using a two-tailed, unpaired t test.

(H) Representative confocal images of *ERphR* KO NRK cells treated with 1 μM Torin1 for 24 h. NRK cells are labeled by ER Tracker. Arrows indicate ER puncta. Scale bars, 5 μm.

(I) Quantification of the ratio of cells shown in (H) contained with ER puncta. More than 89 cells were analyzed per group. Error bars indicate mean ± SEM. ns, not significant. P values were calculated using a two-tailed, unpaired t test.

(J) Representative confocal images of NRK cells stably co-expressing EGFP-LC3 and mCherry-REEP5 treated with or without 1 μM Torin1 for 24 h. Scale bars, 5 μm.

(K) Pearson correlation coefficient of LC3 and REEP5 signals of the cells shown in (J). More than 100 cells were analyzed for each group in each experiment (n=3 independent experiments). Error bars indicate mean ± SEM. ****, $p < 0.0001$. P values were

calculated using a two-tailed, unpaired t test.

(L) Time-lapse live cell images of NRK cells stably co-expressing EGFP-LC3 and

mCherry-REEP5 treated with Torin1. Here shows representative snapshots at indicated

time points. Scale bar, 5 µm.

(M) WT and *Atg8* hexa KO HeLa cells stably co-expressing LAMP1-YFP and

mCherry-REEP5 were treated with or without 1 µM Torin1 for 24 h. More than 30 cells

were analyzed for each group. Scale bars, 5 µm.

(N) Pearson correlation coefficient of LAMP1 and REEP5 signals of the cells shown in

(M). More than 30 cells were analyzed for each group. Error bars indicate mean ± SEM.

**, $p < 0.01$, ns, not significant. P values were calculated using a two-tailed, unpaired t

test.

(O) Representative TEM images of WT and *Atg8* hexa KO HeLa cells treated with or

without 1 µM Torin1 for 24 h. Red arrow indicates autolamellasome (AL), yellow

arrows indicate lysosome/autolysosome containing with autolamellasomes (LYAL).

Scale bars, 500 nm.

(P) TEM images from (O) were quantified for the amount of each kind of structure (AL:

autolamellasome, LYAL: lysosome/autolysosome containing with autolamellasome) in

15 µm^2 area. More than 45 cells were imaged for each group. Data are reported as mean

± SEM.

(Q) Verification of knockout cells. Western blot results, quantitative RT-PCR and fpkm

values are shown here.

1055

1056

Extended Data Fig. 4. Autophagy-dependent assembly of autolamellasome from ER fragments.

(A) FRAP analysis of OAT-treated NRK-*Atg7*-KO cells stably expressing mCherry-REEP5. The boxed regions are magnified on the right. Dashed circles indicate the bleached region. Time intervals are indicated relative to bleaching. Scale bars: 5 μm (main image) and 1 μm (magnified view).

(B) Normalized mean fluorescence intensity of the bleached region over time. More

than 20 cells were analyzed. Data are presented as mean ± SEM. Intensity values were

normalized to pre-bleach levels.

(C) Time-lapse live cell images of NRK cells stably co-expressing LAMP1-YFP and

mCherry-REEP5 treated with OAT. Here shows representative snapshots at indicated

time points. Arrows indicate ER puncta while asterisks indicate ER patches. Scale bar,

5 μm.

(D) Western blot results of NRK cells treated with 200 μM OA 24 h followed by 1 μM

Torin1 in a time series. Cell lysates were subjected to SDS-PAGE and immunoblotted

with the indicated antibodies. Blots are representative of n=3 independent biological

replicates.

(E) Quantification of integrated band intensities from (D). Individual data points

represent relative protein levels in the time point normalized to 0 h and loading control

Actin. n=3 independent biological repeats. Data are presented as mean ± SEM.

1077

Extended Data Fig. 5. Basal formation of autolamellasomes in cell lines and

primary cells.

(A) Representative TEM images of AL (autolamellasome) and LYAL

(lysosome/autolysosome containing with autolamellasome) in indicated cell lines.

Scale bars, 200 nm.

(B) Indicated cell lines were co-stained with anti-LAMP1 and anti-SEC61B antibodies.

White arrows: free-standing ER puncta, yellow arrows: ER puncta inside lysosomes.

Scale bars: 5 μm (main image) and 2 μm (magnified view).

(C) Statistic of cells from (B) for the ratio of cells with SEC61B puncta. More than 38

cells were analyzed for each group. Data are presented as mean ± SEM.

(D) Representative TEM images of whole mouse zygote (scale bar=10 μm) and

enlarged view (scale bar=200 nm).

(E) Indicated primary cells were co-stained with anti-LAMP1 and anti-REEP5

antibodies. White arrows: free-standing ER puncta, yellow arrows: ER puncta inside

lysosomes. Scale bars: 10 μm (main image) and 2 μm (magnified view).

(F) Statistic of cells from (E) for the ratio of cells with REEP5 puncta. More than 60

cells were analyzed for each group. Data are presented as mean ± SEM.

(G) Representative TEM images of AL (autolamellasome) and LYAL

(lysosome/autolysosome containing with autolamellasome) in indicated primary cells.

Scale bars, 200 nm.

1098

Extended Data Fig. 6. *In vitro* reconstitution of autolamellasome.

(A) Representative confocal images of cells following protocol A. Cell death was verified using Apotracker and propidium iodide (PI). Scale bar, 10 μm.

(B) Quantification of the ratio of cells double-positive for Apotracker and PI from (A). More than 150 cells were analyzed per group. Data are presented as mean ± SEM. ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(C) Representative confocal images of cells following protocol A. Cell permeability was verified using Cy5. Scale bar, 10 μm.

(D) Quantification of the ratio of Cy5-positive cells in (C). More than 85 cells were analyzed per group in each experiment (n=3 independent biological repeats). Data are presented as mean ± SEM. ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(E) Representative confocal images of cells following protocol A. Lysosome integrity was verified using Magic Red. Scale bar, 10 μm.

(F) Quantification of intact lysosomes per cell from (E). More than 40 cells were analyzed per group. Data are presented as mean ± SEM. ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(G) Representative confocal images of cells following protocol A. Microtubule integrity was verified using SirTubulin. Scale bar, 10 μm.

(H) Quantification of microtubule mean intensity per cell in (G). More than 37 cells were analyzed per group. Data are presented as mean ± SEM. ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(I) Negative staining-TEM images of S150 and P150. Scale bar, 1 μm.

(J) Schematic diagram illustrating the process of *in vitro* reconstitution (Protocol D). OA primmed NRK cells were permeabilized in reconstitution buffer containing 40 μg/mL digitonin for 30 min, followed by incubation in reconstitution buffer (mock group) or buffer supplemented with S150, ATP and GTP (all group) for 6-8 h at 30 ℃.

(K) Representative confocal images of cells following protocol D. NRK cells were treated with or without 40 μg/mL digitonin for 30 min. Lysosome integrity was verified using Lysotracker and Magic Red. Scale bars, 10 μm.

(L-M) Quantification of mean intensity of Lysotracker signals (L) or Magic Red signals (M) from cells in (K). More than 34 cells were analyzed per group for (L) and more than 45 cells were analyzed per group for (M). Data are presented as mean ± SEM. ****, p<0.0001. P values were calculated using a two-tailed, unpaired t test.

(N) Cell death in NRK cells following protocol D were verified using Apotracker and PI staining and imaged by confocal microscope. Scale bar, 10 μm.

(O) Quantification of the ratio of cells positive for both Apotracker and PI from (N).

1136   More than 100 cells were analyzed per group. Data are presented as mean $\pm$ SEM. ****,

1137 $p<0.0001$. P values were calculated using a two-tailed, unpaired t test.

1138

Tet on EGFP-Progerin

1140

Extended Data Fig. 7. Increased basal autolamellasome formation in senescent

cells and cells from premature aging patients.

(A) Representative TEM images of DOXO-induced senescent WT and autophagy-

deficient NRK cells. White dashed lines indicate accumulation of whorls. Black dashed

lines indicate clusters of ER fragments. Red arrows indicate autolamellasomes (AL)

and yellow arrows indicate lysosome/autolysosome containing with autolamellasome

(LYAL). Scale bars: 5 μm (large view), 500 nm (magnified view).

(B) Schematic diagram illustrating the process of TBHP-induced cellular senescence.

(C) β-galactosidase staining results of WT and autophagy-deficient (*Atg2* KO and *Atg7*

KO) NRK cells after TBHP treatment. *Atg5* KO NRK cells died shortly after treatment.

Scale bar, 100 pixels.

(D) Quantification of β-galactosidase-positive cell ratios from (C). More than 40 cells

were analyzed per group. Data are presented as mean ± SEM. ns, not significant. P

values were calculated using a two-tailed, unpaired t test.

(E) Western blot results of NRK cells treated with or without TBHP as indicated in (B).

Cell lysates were subjected to SDS-PAGE and immunoblotted with the indicated

antibodies. GAPDH served as a loading control. Blots are representative of n=3

independent biological replicates.

(F) Quantification of phospho-p70 S6K (T389) band intensities from (E). Individual

data points represent values normalized to total p70 S6K and expressed relative to the

untreated control. n=3 independent biological repeats. Data are presented as mean ±

SEM.

(G) Representative confocal (left layer) and SIM images (right layer) of TBHP-treated

WT and autophagy-deficient NRK cells. Arrows indicate ER puncta; asterisks indicate

ER patches. Scale bars denote 10 μm for confocal images and 5 μm for SIM images.

(H-J) Statistics of cells from (G). (H) Pearson correlation coefficient of LAMP1 and

REEP5 signals. (I) Ratio of cells with ER puncta. (J) Ratio of cells with ER patches.

More than 65 cells were analyzed in each group for each experiment (n=3 independent

biological repeats). Data are presented as mean ± SEM. ****, p<0.0001. P values were

calculated using a two-tailed, unpaired t test.

(K) Representative TEM images of TBHP-treated WT and autophagy-deficient NRK

cells. Blue arrows indicate electron dense structures, red arrows indicate

autolamellasome (AL), and yellow arrow indicates lysosome/autolysosome containing

with autolamellasomes (LYAL). Black dashed lines indicate clusters of ER fragments.

Boxed regions are magnified on the right. Scale bars, 2 μm for upper images and 1 μm

for upper images.

(L) TEM images from (K) were quantified for the amount of each kind of structure (AL:

autolamellasome, LYAL: lysosome/autolysosome containing with autolamellasome) in

15 μm^2 area. More than 40 cells were imaged for each group. Data are reported as mean

± SEM.

(M) Quantification of frequency of existence of ER patch per 65 μm^2 area from TEM

images in (K).

(N) Representative TEM images of WT and autophagy-deficient NRK cells expressing

doxycycline (Dox)-inducible EGFP-Progerin, induced for 0, 5, or 10 days (D0, D5,

D10).   Red arrows indicate autolamellasome (LY), and yellow arrow indicates

lysosome/autolysosome containing with autolamellasomes (LYAL). Scale bars, 500

nm.

1188

1189   **Supplementary videos and information**

1190

1191   **Supplementary video 1. Continuous formation of ER puncta and their**

1192   **translocation into lysosomes/autolysosomes during Torin1 treatment.** Time-lapse

1193   live cell imaging of NRK cells stably co-expressing LAMP1-mStaygold and HaloTag-

1194   REEP5 treated with 1 µM Torin1**.** Cells were imaged under Olympus FV4000 confocal

1195   microscope**.** Scale bar, 10 µm.

1196

1197   **Supplementary video 2. Continuous formation of ER puncta and their**

1198   **colocalization with LC3 during Torin1 treatment.** Time-lapse live cell imaging of

1199   NRK cells stably co-expressing BFP-LC3 and mCherry-REEP5 treated with 1 µM

1200   Torin1 starting at 4 h and imaged on an Olympus FV4000 confocal microscope**.** Scale

1201   bar, 10 µm.

1202

1203   **Supplementary video 3. FRAP analysis of ER patch.** *Atg7* KO NRK cells stably

1204   expressing mCherry-REEP5 were primed with 200 µM oleic acid for 24 h, then treated

1205   with 1 µM Torin1 for 20 h. Cells were photobleached and imaged on a Nikon AX R

1206   microscope. Scale bar, 10 µm.

1207

1208   **Supplementary video 4. Continuous transition from ER patches to ER puncta**

1209   **during Torin1 treatment.** Time-lapse live cell imaging of NRK cells stably co-

1210   expressing Lamp1-YFP and mCherry-REEP5. Cells were pretreated with 200 µM oleic

acid for 24 h, then treated with 1 µM Torin1 and imaged on an Olympus FV4000 confocal microscope. Scale bar, 10 µm.

Resource availability

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Li Yu. All data supporting the findings of this study are available from the corresponding author upon reasonable request.

Acknowledgements

This research was supported by Tsinghua University Dushi Program (grant no. 20251080019), the Ministry of Science and Technology of the People's Republic of China (grant no. 2024YFA1307301，2024YFF1502900), the National Natural Science Foundation of China (grant no. 32030023，92354306, and 32330025), Tsinghua-Toyota Joint Research Fund (grant no. 20233930058), and Scientific and Technological Innovation Project of China Academy of Chinese Medical Sciences (grant no.CI2023C024YL). We sincerely show our gratefulness to the State Key Laboratory of Membrane Biology for confocal microscopy imaging and facility support. We are grateful to Yan Yang, Chenguang Zhao, Yanfei Hu, Qian Li from the Cell Biology Facility affiliated with the Center of Biomedical Analysis, Tsinghua University, for technical assistance and equipment support. We thank SLSTU-Nikon Biological Imaging Center for microscopy imaging and facility support. We thank Dr. Dachuan Zhang (Tsinghua University, Dr. Li Yu's lab) for providing NRK-*Atf4*-KO, NRK-*Atf6*-KO, NRK-*Xbp1*-KO cells. We thank Dr. Jiayu He (Xinjiang Medical University, Dr. Na Mi's lab) for providing NRK-*Atg14*-KO cells. We thank Dr. Du Feng from Guangzhou Medical University for providing HeLa-*ATG7*-KO cells and its wildtype HeLa cells. We thank Dr. Michael Lazarou from Monash University for providing HeLa-*ATG8*-hexa KO cells and its wildtype HeLa cells. We thank Dr. Xiangyang Wang from Dr. Chuanmao Zhang's lab (Peking University, Kunming University of Science and Technology) for providing human dermal fibroblast (HDF) cells. We thank Dr. Zheng

Jiang from Dr. Anming Meng's lab (Tsinghua University) for providing mouse zygotes. We thank Dr. Wenfeng Fu from Dr. Dong Li's lab (Tsinghua University) for providing the LAMP1-mStaygold and the Calnexin-HaloTag plasmids. We thank Dr. Xiong Ji from Peking University for proving the pMK243 (Tet-OsTIR1-PURO) plasmid.

Author contributions

L.Y. and D.L. conceived and designed the study. L.Y. supervised the project and wrote the manuscript. D.L. performed cell biology, molecular biology, biochemistry and EM experiments. R.Z. performed cryo-EM experiments. W.S., D.Z. and Y.Y. participated in generation of knockout cells. X.S. and H.Z. isolated primary cells from mouse tissues. All authors reviewed and approved the final manuscript.

Declaration of interests

L.Y. is the scientific founder of Migrasome therapeutics Ltd. All other authors declare no conflicts of interest.

Materials and Methods

All research adhered to the guidelines from the Academic Integrity and Ethics committee of Tsinghua University.

Reagents

Torin1 (MCE, HY-13003), ER-Tracker™ Blue-White DPX (ThermoFisher, E12353), 3,3′-Diaminobenzidine tetrahydrochloride hydrate (DAB tetrahydrochloride hydrate, Sigma-Aldrich, D5637), Hydrogen peroxide (Sigma-Aldrich, 88597), Rapamycin (MCE, HY-10219), Sodium oleate (Sigma-Aldrich, O7501), Chloroquine (MCE, HY-17589A); ATP (ThermoFisher, R0441); GTP (Sangon Biotech, A620332-0250); Creatine Phosphate 2Na (Lablead, 0271); Creatine Kinase (Yeasen, 14502ES10); Apotracker Green (BioLegend, 427401); Propidium iodide Staining Solution (Yeasen, 40710ES03); Cy5 (Solarbio, S1053); Lysotracker (Invitrogen, L12492); Magic Red

(Abcam, ab270772)); SirTubulin (Cytoskeleton, Inc., CY-SC002); Doxorubicin

hydrochloride (MCE, HY-15142); TBHP (tert-Butyl hydroperoxide solution, Aladdin,

B106035), G418 sulfate (Amresco, E859), Hygromycin B (Roche, 10843555001),

Puromycin (ThermoFisher, A1113803), Doxycycline (Sigma-Aldrich, D3072),

Vigofect (Vigorous, T001), Senescence β-Galactosidase Staining Kit (Beyotime,

C0602), Collagenase I (Solarbio, C8140), Collagenase IV (Yeasen, 40510ES76),

DNaseI (Aladdin, D106200), Liberase (Sigma-Aldrich, 05401127001).

1276

Antibodies

RTN3 (Proteintech Cat# 12055-2-AP, RRID:AB_2301357); FAM134B (Proteintech

Cat# 21537-1-AP, RRID:AB_2878879); TEX264 (Proteintech Cat# 25818-1-AP,

RRID:AB_2880272); ATL3 (Proteintech Cat# 16921-1-AP, RRID:AB_2290228);

SEC62 (Abcam Cat# ab140644); REEP5 (Proteintech Cat# 14643-1-AP,

RRID:AB_2923648); REEP5 (Proteintech Cat# 68119-1-Ig, RRID: AB_2178440);

SEC61B (Abcam Cat# ab15576, RRID:AB_301985); Calnexin (Abcam Cat# ab22595,

RRID:AB_2069006); p62 (SQSTM1) (MBL International Cat# PM045,

RRID:AB_1279301); GAPDH (Proteintech Cat# 60004-1-Ig, RRID:AB_2107436);

LAMP1 (Sigma-Aldrich Cat# L1418, RRID:AB_477157); LAMP1 (Enzo Life

Sciences Cat# ADI-VAM-EN001, RRID:AB_10630197); mCherry-Tag (Easybio Cat#

BE2026, RRID:AB_2936383); ATF4 (Proteintech Cat# 10835-1-AP,

RRID:AB_2058600); SEC22B (Synaptic Systems Cat# 186 003, RRID:AB_993020);

SAR1B (Proteintech Cat# 15453-1-AP, RRID:AB_2183985); SEC23A (Abcam Cat#

ab137582); ATG5 (Cell Signaling Technology Cat# 12994, RRID:AB_2630393);

ATG7 (Cell Signaling Technology Cat# 8558, RRID:AB_10831194); ATG9A (Abcam

Cat# ab108338, RRID:AB_10863880); ATG14 (Cell Signaling Technology Cat# 96752,

RRID:AB_2737056); ATG16L (Cell Signaling Technology Cat# 8089,

RRID:AB_10950320); ATG13 (Cell Signaling Technology Cat# 13273,

RRID:AB_2798169); Beclin-1 (Cell Signaling Technology Cat# 3495,

RRID:AB_1903911); ULK1 (Cell Signaling Technology Cat# 8054,

RRID:AB_11178668); FIP200 (Cell Signaling Technology Cat# 12436, RRID:AB_2797913); WIPI2 (abcepta Cat# AP13314a, RRID:AB_11136530); ATG3 (Cell Signaling Technology Cat# 3415, RRID:AB_2059244); PI4KIIA (Proteintech Cat# 15318-1-AP, RRID:AB_2268225); VPS13C (Proteintech Cat# 28676-1-AP, RRID:AB_2881191); SCAP (Abcam Cat# EPR26242-60); p70(S6K) (Cell Signaling Technology Cat# 2708, RRID:AB_390722); Phospho-p70(S6K) (Thr389) Antibody (Cell Signaling Technology Cat# 9206, RRID:AB_330944); Phospho-p70 (S6K) (Thr389) Antibody (Proteintech Cat# 28735-1-AP, RRID:AB_2918197); Phospho-p70(S6K) (Thr229) Antibody (Proteintech Cat# 81592-1-RR, RRID:AB_2935429); Goat Anti-Rabbit IgG (H+L) (Jackson Cat#111-035-003, RRID:AB_2313567); Goat Anti-Mouse IgG (H+L) (Jackson Cat#115-035-003, RRID:AB_10015289); Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 546 (ThermoFisher Cat#A-11035, RRID:AB_2534093); Goat anti-mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 488 (ThermoFisher, Cat#A-11029, RRID:AB_2534093).

Overexpression plasmids and stable cell lines

mCherry-REEP5, HaloTag-REEP5, mCherry-SEC61B, BFP-LC3 were constructed on the PiggyBac transposon vector (RRID:Addgene_210330; RRID:Addgene_210332). LAMP1-YFP was constructed on the pEYFP-N1 vector. ER-DsRed was purchased from Addgene (DsRed2-ER-5, RRID:Addgene_55836). EGFP-REEP5, APEX2-EGFP-REEP5, APEX2-EGFP-SEC61B, EGFP-LC3, were constructed on the pEGFP-C1 vector. APEX2-mCherry-LC3 was constructed on the pmCherry-C1 vector. EGFP-Progerin was constructed on the pMK243 vector (Tet-OsTIR1-PURO) (this plasmid is a kind gift from Dr. Chuanmao Zhang's lab). LAMP1-mStaygold and Calnexin-HaloTag were kind gifts from Dr. Dong Li's lab (Tsinghua University). Stable NRK cell lines expressing the following fluorescent fusion proteins were generated via electroporation, antibiotic selection (hygromycin, G418 sulfate, and puromycin), and FACS sorting: LAMP1-YFP/mCherry-REEP5, APEX2-EGFP-REEP5, APEX2-EGFP-

SEC61B, EGFP-REEP5, LAMP1-YFP/HaloTag-REEP5, LAMP1-mStaygold/Calnexin-HaloTag, LAMP1-YFP/mCherry-SEC61B, LAMP1-YFP/ER-DsRed, BFP-LC3/LAMP1-YFP/mCherry-REEP5, EGFP-LC3/mCherry-REEP5, and APEX2-mCherry-LC3.

Generation of KO cell lines

All the knockout cell lines were generated with CRISPR-Cas9 system. Genes of interest were deleted using a modified PX458 plasmid or PX459 plasmid (provided by Wei Guo from Zhejiang University, Haining, China) that contains two single guide RNAs (sgRNAs) coupled with Cas9 nuclease. The sgRNAs were designed at the website of Broad Institute (https://portals.broadinstitute.org/gppx/crispick/public). Cells transfected with PX458 were seeded into 96-well plates by fluorescence-activated cell sorter selection of EGFP signal expression. Cells transfected with PX459 were selected with puromycin for 7 days and then seeded into 96-well plates by FACS. Genotyping PCR, Western blotting and qPCR were applied for knockout verification.

The targeted genome sequence pairs (two sites slicing) are listed below:

Fam134b-gRNA1: 5'-GCCACTGCATTGCAGAATCA-3'

Fam134b-gRNA2: 5'-CAGAAGAAACGTGAGAGATC-3'

Sec62-gRNA1: 5'-ACAGTTGAATCGAAGATACT-3'

Sec62-gRNA2: 5'-AAATGGGCAAAGGCCAAGAA-3'

Tex264-gRNA1: 5'-GGAGGCTCCTTACCGTCTGG-3'

Tex264-gRNA2: 5'-AGGTGGATGAGCTCAGGTGA-3'

Atl3-gRNA1: 5'-GCTCTGACCCAGAAACTACT-3'

Atl3-gRNA2: 5'-TGAGCTAGAAGAGAGAGCCT-3'

Ccpg1-gRNA1: 5'-CATGCTAATAGCAATCACCA-3'

Ccpg1-gRNA2: 5'-ATTAAAGATGACCTTAACCT-3'

Sar1b-gRNA1: 5'-ATATTTGACTGGATTTACAG-3'

Sar1b -gRNA2: 5'-TCCTTGGATTGGATAATGCC-3'

Sec23a-gRNA1: 5'-AGGACAGGCTCGTACTGAAT-3'

Sec23a-gRNA2: 5'-GATATCCCAGCATATGTAGG-3'

Atg5-gRNA1: 5'-AAGAGTCAGCTATTTGACGC-3'

Atg5-gRNA2: 5'-TCCGTGCAAGGATGCAGTTG-3'

Atg7-gRNA1: 5'-GAACGAGTATCGCCTGGACG-3'

Atg7-gRNA2: 5'-TGACGCCTTCAGTTCGACAC-3'

Atg9a-gRNA1: 5'-AGGCCGAGTACAAACGTGGA-3'

Atg9a-gRNA2: 5'-GAGCTGGGCAAACTCGTCCC-3'

Atg16l1-gRNA1: 5'-AAAAGCACGACGTACCAAAT-3'

Atg16l1-gRNA2: 5'-AGAAGCCAATCGCCTTAATG-3'

Atg13-gRNA1: 5'-GTCGAGCCTGGACAATCACT-3'

Atg13-gRNA2: 5'-GATCTCACTCAAGACTTCTG-3'

Beclin1-gRNA1: 5'-CCTGGACCGAGTGACCATTC-3'

Beclin1-gRNA2: 5'-GGAAGAGGCTAACTCAGGAG-3'

Ulk1-gRNA1: 5'-ATCATGTCCCAGCACTACGA-3'

Ulk1-gRNA2: 5'-GTGGTGGAAAAATTCATCTG-3'

Wipi2-gRNA1: 5'-ACTGCTACTTGGCGTACCCA-3'

Wipi2-gRNA2: 5'-GCTCTATATACACAACATCC-3'

Atg3-gRNA1: 5'-GTAGACACGTACCATAACAC-3'

Atg3-gRNA2: 5'-AGGGGAAGAATTGAAAGTGA-3'

Pi4k2a-gRNA1: 5'-TTAGGGTTAAGGTTCCCGTA-3'

Pi4k2a -gRNA2: 5'-ATTGACCGAGTAAAGTCCAG-3'

Vps13c-gRNA1: 5'-TTAGAACACTGGTACGTCAC-3'

Vps13c-gRNA2: 5'-TGGTTAGAAATGCACCATAC-3'

Npc2-gRNA1: 5'-CAGGAATAGGGAAGTAGACT-3'

Npc2-gRNA2: 5'-AAAGGTGACGTTGACACTGT-3'

Scap-gRNA1: 5'-TCTGTCCCGAGCATTCCAAC-3'

Scap-gRNA2: 5'-ACCTACCCGCCATTGAGTGT-3'

Srebf1-gRNA1: 5'-AGGCTTGAGTACCCCAGCAC-3'

Srebf1-gRNA2: 5'-AGCATGCCTCGGCTATGTGA-3'

NRK-*Perk*-KO, NRK-*Sec22b*-KO, NRK-*Atg2*-KO, NRK-*p62*-KO, MEF-*Fip200*-KO,

HeLa-*ATG8*-hexa KO cells were reported before[46,63-66]. NRK-*Atf4*-KO, NRK-*Atf6*-KO,

NRK-*Xbp1*-KO cells were kind gifts from Dr. Dachuan Zhang (Tsinghua University, Li

Yu's lab), NRK-*Atg14*-KO cell line was a kind gift from Dr. Jiayu He (Xinjiang Medical

University, Dr. Na Mi's lab).

1390

Cell culture

S2 cells were cultured at 28 °C and 5% CO_2 in Schneider's Drosophila medium

supplemented with 10% fetal bovine serum, 1% penicillin-streptomycin and 1%

Glutamax. MDA-MB-231 cells were cultured at 37 °C and 5% CO_2 in RPMI-1640 high

glucose medium supplemented with 10% fetal bovine serum, 1% penicillin-

streptomycin and 1% Glutamax. THP1 cells were cultured at 37 °C and 5% CO_2 in

RPMI-1640 high glucose medium supplemented with 10% fetal bovine serum, 1%

penicillin-streptomycin, 1% Glutamax and 100 ng/mL PMA. The isolated fascia

fibroblasts (FFB) and the isolated mouse endometrial stromal cells (MESC) were

cultured at 37 °C and 5% CO_2 in DMEM/F-12 medium supplemented with 10% fetal

bovine serum, 1% penicillin-streptomycin and 1% Glutamax. The isolated hepatocytes

were cultured at 37 °C and 5% CO_2 in Williams E Medium supplemented with 5% fetal

bovine serum, 1% penicillin-streptomycin and 1% Glutamax. Other cells were cultured

at 37 °C and 5% CO_2 in DMEM high glucose medium supplemented with 10% fetal

bovine serum, 1% penicillin-streptomycin and 1% Glutamax.

1406

Immunofluorescence and live cell imaging

For immunofluorescence, cells were grown on confocal dishes and fixed by 4%

paraformaldehyde (PFA) at room temperature for 15 min. Then, samples were blocked

and permeabilized in PBS containing 0.03% saponin and 10% FBS for 30 minutes at

room temperature. Following was incubation with primary antibodies diluted in PBS

containing 10% FBS overnight at 4°C. After washing three times with PBS, cells were

incubated with secondary antibodies diluted in PBS containing 10% FBS for 1 h at

room temperature in the dark. After final washes with PBS, the samples were imaged

with light microscopes.

For live cell imaging, cells were seeded on 3.5mm confocal dishes and monitored with

microscopes equipped with a Chamlide environmental incubator system, which

maintained the cells at 37 °C and 5% CO_2. Confocal images were acquired with

Olympus (FV3000, FV4000) and Nikon (A1HD25, AX R). SIM images were acquired

with a Nikon N-SIM microscope equipped with a 100×oil-immersion objective (NA

1.49). Images were reconstructed and analyzed with cellSens software (Olympus), NIS-

Elements AR software (Nikon) or ImageJ software (NIH). Z-stack images were

reconstructed with Imaris software.

1424

Western blotting and intensities analysis

Cells were solubilized in lysis buffer (2.5% SDS, 50 mM Tris-HCl, pH 7.4) and

denatured at 95°C for 15 min. Following protein quantification with the BCA assay kit

(Thermo Fisher, 23227), samples were separated on precast SDS-PAGE gradient gels

(Lablead; Epizyme) and electroblotted onto PVDF membranes (Millipore, ISEQ00010).

The membranes were blocked in TBST (20 mM Tris-HCl, pH 7.6, 137 mM NaCl and

0.2% Tween-20) containing 5% milk for 1 h at room temperature. Primary antibodies

were diluted in Solution I (TOYOBO, NKB-201). After incubation with primary

antibodies at 4 °C overnight, the membranes were washed with TBST and then

incubated with horseradish peroxidase (HRP)-conjugated secondary antibodies diluted

in antibody solution buffer (NCM Biotech, WB100D) at room temperature for 1 h. After

final washes with TBST, the membranes were detected and quantified with ChemiDoc

Touch Imaging System (Bio-Rad, ChemiDoc MP, RRID:SCR_021693) using Westar

ETAC (Cyanagen).

Band intensities were quantified using either ImageLab software (Bio-Rad) or ImageJ

software (NIH). For ImageLab software (Bio-Rad), lanes were automatically detected

and manually adjusted for alignment. Background subtraction was applied using the

rolling disk method. Band volumes (integrated intensity) were measured and

normalized to the corresponding loading control (e.g., Actin or GAPDH) from the same lane. For ImageJ software (NIH), images were converted to 8-bit grayscale format, and background subtraction was performed using the "Rolling ball radius" method. Proteins of interest were selected using rectangular selections, and lane boundaries were manually defined. The integrated density values were measured and normalized to the corresponding loading control from the same blot. Data from at least three independent biological replicates were analyzed and expressed as fold change relative to the control group.

Quantitative real-time PCR

Total RNA was extracted with MolPure® Flash Cell/Tissue Total RNA Kit (Yeasen, 19221ES50). cDNA was synthesized using a ReverTra Ace qPCR RT Kit (TOYOBO, FSQ-101). Quantitative real-time PCR was performed using 2× T5 Fast qPCR Mix (SYBR Green I) (TSINGKE Biological Technology, TSE202). Primer sequences are provided below:

Ccpg1-1F: 5'-GTCACTGGCGTCTGGAGC-3'

Ccpg1-1R: 5'-TCAGACATTTTTCAGGTCCGAAG-3'

Ccpg1-2F: 5'-CGCGAGCAGGTTGTTGTT-3'

Ccpg1-2R: 5'-TTCAGACATTTTTCAGGTCCGA-3'

Npc2-F: 5'-CCTGAGCCTGACGGTTGTAA-3'

Npc2-R: 5'-CACCACCAGTTTTAGAGAGGG-3'

Srebf1-F: 5'-CTTGACCGACATCGAAGACAT-3'

Srebf1-R: 5'-CCAGCATAGGGGGCATCAAA-3'

Label-free Mass Spectrum

For protein extraction, cells were lysed in 10 volumes of ice-cold lysis buffer (1% sodium deoxycholate, 10 mM Tris-HCl, pH 8.5), immediately boiled at 100 °C for 10 min to inactivate proteases, and sonicated on a pre-chilled (4 °C) non-contact ultrasonicator (30 s on/30 s off, 15 cycles). Lysates were cleared by brief centrifugation,

and supernatants were stored at -80 °C until analysis. Protein concentration was determined by BCA assay. Label-free quantitative proteomic analysis was performed by Protein Chemistry and Proteomics Platform (Technology Center for Protein Sciences, Tsinghua University) using three biological replicates.

TEM

Cells were cultured on 3.5 mm confocal dishes and fixed in 2.5% glutaraldehyde (GA) (Legene). The cells were washed with 0.1 M PB buffer (pH 7.2) three times, and post-fixed in 1% osmium tetroxide (Ted Pella)/1.5% K3[Fe(CN)6] in distilled water for 30 min at 4 °C. After three washes with distilled water, the samples were stained with 1% uranyl acetate overnight at 4°C in the dark. After three washes with distilled water, the cells were dehydrated in a graded ethanol series (50%, 70%, 80%, 90%, 100%, 100%, 100%, for 2 min each on ice, then 100%, 5 min at RT). The cells were infiltrated in Pon 812 (SPI) resin (50%, 67%, 75% for 30 min each, then 100% for 1 h and 100% overnight at RT), and then polymerized at 60 °C for 48 h. Ultrathin sections of 70 nm were cut by ultramicrotome (Leica EM UC7). After staining with uranyl acetate and lead citrate, sections were observed under 120 kV transmission electron microscope (Hitachi, HT-7800).

CLEM

Cells grown on gridded glass-bottom dishes (Cellvis, D35-14-1.5GI) were fixed with 4% PFA and immediately imaged with confocal microscope to document the cells of interest at different magnifications. After that, the cells were post-fixed with 2.5% GA and prepared for TEM. The steps for preparing ultrathin sections were the same as standard TEM sample preparation protocols.

APEX2-TEM

Cells stably expressing APEX2-EGFP-REEP5 (or APEX2-EGFP-SEC61B, APEX2-mCherry-LC3) were cultured on 3.5 mm confocal dishes and fixed with 2.5% GA. All

subsequent steps were performed on ice within 1 h of fixation. After three washes with 0.1M PB buffer (PH 7.2), cells were incubated in a freshly prepared solution of 0.5 mg/mL (1.4 mM) DAB tetrahydrochloride (Sigma, D5637) in 0.1M PB buffer (PH 7.2) for 20 min. Next, a fresh solution containing 0.5 mg/mL (1.4 mM) DAB and 0.03% (v/v) H_2O_2 was used for the enzymatic reaction. After reaction, the cells were washed three times with 0.1M PB buffer (PH 7.2). Post-fixation staining was performed with 2% osmium tetroxide for 5 min. Cells were then rinsed three times in distilled water, and placed in 2% aqueous uranyl acetate (Electron Microscopy Sciences) overnight. Subsequent steps followed standard TEM sample preparation protocols.

FIB-SEM

The resin block was trimmed to expose cells at the surface, then coated with a ~20-nm gold layer followed by a ~600-nm platinum layer applied perpendicular to the block face. Imaging was performed on a Helios NanoLab G3 UC scanning electron microscope in back-scattered electron (BSE) mode using TLD (through-the-lens) and ICD (in-column) detectors. Each milling cycle used a 30 kV, 0.79 nA ion beam to remove a 10-nm-thick surface layer, and images were acquired with a 2 kV, 0.4 nA electron beam at 8 μs/pixel dwell time.

Sample Vitrification

For sample preparation, a cell suspension with a concentration of approximately 3×10^6 cells/mL was used. 200-mesh copper grids with a continuous carbon film (EMCN, BZ1011125a) were rendered hydrophilic by glow-discharging in ambient air for 25 seconds at 15 mA using a Pelco EasiGlow glow-discharge system. The vitrification process was carried out using a Leica EM GP2 automatic plunge-freezer, operated within an environmental chamber maintained at 25°C and 75% relative humidity to minimize sample evaporation.

A volume of 3 μL of the suspended cell solution was applied to the carbon-coated side of the glow-discharged grid. To achieve a thin, uniform layer of cells suitable for

imaging, 2 µL of buffer was applied to the reverse side of the grid. The grid was blotted from the reverse side for 4-6 s and immediately plunged into liquid ethane cooled by liquid nitrogen, resulting in the vitrification of the sample. Grids were subsequently stored in liquid nitrogen until required for further processing.

Cryo-lamella preparation

Lamellae were prepared from the vitrified cells using a multi-step milling procedure analogous to established protocols[67] with a Helios dual-beam cryo-focused ion beam/scanning electron microscope (cryo-FIB/SEM) (Thermo Fisher Scientific). To balance milling speed with precision and minimize beam-induced sample damage during the milling process, the Ga^{2+} ion beam current at an accelerating voltage of 30 kV was progressively reduced from 0.79 nA for rough milling to 24 pA for final polishing. This procedure yielded lamellae with a final thickness between 100 to 200 nm.

Cryo-electron tomography data collection

Lamellae were imaged on a 300 kV Titan Krios electron microscope (Thermo Fisher Scientific) equipped with a Gatan post-column energy filter. Images were recorded at a magnification of ×53,000, resulting in a calibrated pixel size of 2.4 Å. Tilt-series were collected using the bidirectional tilt scheme over a range of ±60°, with a 2° angular increment. A target defocus of approximately -5 µm was applied, and the total cumulative electron dose for each tilt-series was limited to 120 $e^-/Å^2$.

Tomogram reconstruction

Following data acquisition, the raw movie frames were processed to correct for specimen drift and beam-induced motion using MotionCor2[68]. The resulting tilt-series were subsequently aligned using patch tracking routine and reconstructed into tomograms by weighted back-projection, employing the IMOD software suite[69]. To enhance the signal-to-noise ratio for visualization and subsequent segmentation, the

1559 final tomograms were four-fold binned, and a SIRT-like filter was applied.

1560

Tomogram segmentation and visualization

Three-dimensional segmentation of cellular features within the reconstructed

tomograms was performed using Amira (Thermo Fisher Scientific). A semi-automated

workflow was employed to generate accurate 3D models of membranes and other

structures of interest, which combined automated feature recognition with subsequent

manual refinement. Features such as membranes were first identified using threshold-

based tools within the software. Each segmented feature was then manually inspected,

corrected, and annotated, to ensure anatomical accuracy and remove any artifacts from

the automated process. The finalized 3D models were then rendered within Amira for

visualization and analysis.

1571

FRAP

Fluorescence recovery after photobleaching (FRAP) experiments were performed using

a Nikon AX R confocal microscope equipped with a 100×oil-immersion objective.

After acquiring five prebleach frames (1s intervals), a circular region of interest (ROI;

1 μm diameter) was photobleached with the 488 nm laser at 100% power for 5 s.

Postbleach recovery was recorded at 1-s intervals for 60 s. Mean fluorescence intensity

within the bleached ROI was quantified at each time point, background-subtracted

using an extracellular ROI, corrected for acquisition photobleaching by normalization

to an unbleached reference ROI in the same cell, and expressed relative to the mean

prebleach intensity (set as 100%). Recovery curves were plotted as mean ± SEM

from >20 cells per condition using GraphPad Prism 10 software.

1583

Isolation of primary cells

In this project, we isolated several primary cells from mouse tissues, including lung

fibroblasts (LFB), fascia fibroblasts (FFB), dermal fibroblasts (DFB), hepatocytes,

mouse embryonic fibroblasts (MEF) and mouse endometrial stromal cells (MESC),

following previously described protocols[70-74].

Human dermal fibroblasts (HDF) from healthy donors and HGPS patients were

obtained from Dr. Chuanmao Zhang's lab, as previously reported[61].

1591

***In vitro* reconstitution**

S150 cell extract preparation: Cells were harvested and washed with cold PBS, and

resuspended in 2× cytosol buffer (50 mM Hepes-KOH, 50 mM KOAc, 2.5 mM MgCl2,

250 mM sucrose, PH 7.4) containing protease inhibitors. Lysis was performed by

mechanical disruption (25G needle or homogenizer) until >80% efficiency was

confirmed by typan blue (Solarbio Life Sciences, C0040) staining. The lysate was

centrifuged sequentially at 1,000×g for 10 min and 120,000×g for 20 min at 4°C. The

supernatant was then ultracentrifuged at 150,000×g for 1 h at 4°C. The final S150

supernatant was acquired, protein concentration was determined by BCA assay (~10

mg/mL), and aliquots were flash-frozen in liquid nitrogen and stored at -80°C.

We have tried several protocols for *in vitro* reconstitution, the most efficient is described

below. After incubation with 200 μM oleic acid for 20 h, cells grown on 3.5 mm

confocal dishes were treated with reaction buffer (25 mM Hepes-KOH, 25 mM KOAc,

1.25 mM MgCl2, 125 mM sucrose, PH 7.4) containing 40 μg/mL digitonin and protease

inhibitors alone (mock group), or additionally supplemented with 50% S150 and an

energy system (1 mM ATP, 0.5 mM GTP, 20 mM creatine phosphate, 0.2 mg/mL

creatine kinase) (all group). Both groups were incubated at 30°C for 6-8 h to allow

reconstitution, then fixed with 2.5% GA for TEM analysis. Cell death was confirmed

using Apotracker and PI, and plasma membrane permeabilization was assessed with

Cy5.

1612

Negative staining TEM

A droplet of S150 or resuspended P150 was deposited onto copper grids for 5 minutes.

Excess sample was blotted with filter paper. The grid was promptly washed with a

droplet of double distilled water (ddH2O), stained with 1% uranyl acid for one minute,

and washed twice with droplets of ddH$_2$O. Residual water was blotted using filter paper,

and the grid was air-dried before imaging by transmission electron microscopy (TEM).

1619

Statistical analysis

Normality was assessed using a Kolmogorov-Smirnov test. Statistical analyses were

conducted with GraphPad Prism 10 using an unpaired two-tailed Student's t-test.

Sample sizes were determined empirically without a priori statistical calculations, and

all acquired data were retained for analysis. Error bars indicate mean ± SEM. *P<0.05,

**P<0.01, ***P<0.001, ****P<0.0001, ns, not significant.

1626

1627

References

1. Tsukada, M., and Ohsumi, Y. (1993). Isolation and characterization of autophagy-defective
mutants of Saccharomyces cerevisiae. FEBS Lett *333*, 169-174. 10.1016/0014-
5793(93)80398-e.

2. Takeshige, K., Baba, M., Tsuboi, S., Noda, T., and Ohsumi, Y. (1992). Autophagy in yeast
demonstrated with proteinase-deficient mutants and conditions for its induction. J Cell
Biol *119*, 301-311. 10.1083/jcb.119.2.301.

3. Mizushima, N., Noda, T., Yoshimori, T., Tanaka, Y., Ishii, T., George, M.D., Klionsky, D.J.,
Ohsumi, M., and Ohsumi, Y. (1998). A protein conjugation system essential for autophagy.
Nature *395*, 395-398. 10.1038/26506.

4. Kametaka, S., Matsuura, A., Wada, Y., and Ohsumi, Y. (1996). Structural and functional
analyses of APG5, a gene involved in autophagy in yeast. Gene *178*, 139-143.
10.1016/0378-1119(96)00354-x.

5. Deretic, V., Saitoh, T., and Akira, S. (2013). Autophagy in infection, inflammation and
immunity. Nat Rev Immunol *13*, 722-737. 10.1038/nri3532.

6. Mizushima, N., and Komatsu, M. (2011). Autophagy: renovation of cells and tissues. Cell
147, 728-741. 10.1016/j.cell.2011.10.026.

7. Deretic, V., and Klionsky, D.J. (2008). How cells clean house. Sci Am *298*, 74-81.
10.1038/scientificamerican0508-74.

8. Klionsky, D.J., and Emr, S.D. (2000). Autophagy as a regulated pathway of cellular
degradation. Science *290*, 1717-1721. 10.1126/science.290.5497.1717.

9. Gómez-Sánchez, R., Tooze, S.A., and Reggiori, F. (2021). Membrane supply and
remodeling during autophagosome biogenesis. Curr Opin Cell Biol *71*, 112-119.
10.1016/j.ceb.2021.02.001.

10. Nishimura, T., and Tooze, S.A. (2020). Emerging roles of ATG proteins and membrane
lipids in autophagosome formation. Cell Discov *6*, 32. 10.1038/s41421-020-0161-3.

11. Kirkin, V., McEwan, D.G., Novak, I., and Dikic, I. (2009). A role for ubiquitin in selective

1655   autophagy. Mol Cell *34*, 259-269. 10.1016/j.molcel.2009.04.026.

1656   12.   Vargas, J.N.S., Hamasaki, M., Kawabata, T., Youle, R.J., and Yoshimori, T. (2023). The
1657   mechanisms and roles of selective autophagy in mammals. Nat Rev Mol Cell Biol *24*, 167-
1658   185. 10.1038/s41580-022-00542-2.

1659   13.   Youle, R.J., and Narendra, D.P. (2011). Mechanisms of mitophagy. Nat Rev Mol Cell Biol
1660   *12*, 9-14. 10.1038/nrm3028.

1661   14.   Hong, Z., Pedersen, N.M., Wang, L., Torgersen, M.L., Stenmark, H., and Raiborg, C. (2017).
1662   PtdIns3P controls mTORC1 signaling through lysosomal positioning. J Cell Biol *216*, 4217-
1663   4233. 10.1083/jcb.201611073.

1664   15.   Li, W., He, P., Huang, Y., Li, Y.F., Lu, J., Li, M., Kurihara, H., Luo, Z., Meng, T., Onishi, M., et
1665   al. (2021). Selective autophagy of intracellular organelles: recent research advances.
1666   Theranostics *11*, 222-256. 10.7150/thno.49860.

1667   16.   Pankiv, S., Clausen, T.H., Lamark, T., Brech, A., Bruun, J.A., Outzen, H., Øvervatn, A., Bjørkøy,
1668   G., and Johansen, T. (2007). p62/SQSTM1 binds directly to Atg8/LC3 to facilitate
1669   degradation of ubiquitinated protein aggregates by autophagy. J Biol Chem *282*, 24131-
1670   24145. 10.1074/jbc.M702824200.

1671   17.   Birgisdottir Å, B., Lamark, T., and Johansen, T. (2013). The LIR motif - crucial for selective
1672   autophagy. J Cell Sci *126*, 3237-3247. 10.1242/jcs.126128.

1673   18.   Noda, N.N., Ohsumi, Y., and Inagaki, F. (2010). Atg8-family interacting motif crucial for
1674   selective autophagy. FEBS Lett *584*, 1379-1385. 10.1016/j.febslet.2010.01.018.

1675   19.   Lamark, T., and Johansen, T. (2021). Mechanisms of Selective Autophagy. Annu Rev Cell
1676   Dev Biol *37*, 143-169. 10.1146/annurev-cellbio-120219-035530.

1677   20.   Grumati, P., Dikic, I., and Stolz, A. (2018). ER-phagy at a glance. J Cell Sci *131*.
1678   10.1242/jcs.217364.

1679   21.   Bravo, R., Parra, V., Gatica, D., Rodriguez, A.E., Torrealba, N., Paredes, F., Wang, Z.V.,
1680   Zorzano, A., Hill, J.A., Jaimovich, E., et al. (2013). Endoplasmic reticulum and the unfolded
1681   protein response: dynamics and metabolic integration. Int Rev Cell Mol Biol *301*, 215-290.
1682   10.1016/b978-0-12-407704-1.00005-1.

1683   22.   Schwarz, D.S., and Blower, M.D. (2016). The endoplasmic reticulum: structure, function
1684   and response to cellular signaling. Cell Mol Life Sci *73*, 79-94. 10.1007/s00018-015-2052-
1685   6.

1686   23.   Stolz, A., and Grumati, P. (2019). The various shades of ER-phagy. Febs j *286*, 4642-4649.
1687   10.1111/febs.15031.

1688   24.   Schuck, S., Gallagher, C.M., and Walter, P. (2014). ER-phagy mediates selective
1689   degradation of endoplasmic reticulum independently of the core autophagy machinery.
1690   J Cell Sci *127*, 4078-4088. 10.1242/jcs.154716.

1691   25.   Fregno, I., and Molinari, M. (2019). Proteasomal and lysosomal clearance of faulty
1692   secretory proteins: ER-associated degradation (ERAD) and ER-to-lysosome-associated
1693   degradation (ERLAD) pathways. Crit Rev Biochem Mol Biol *54*, 153-163.
1694   10.1080/10409238.2019.1610351.

1695   26.   Chino, H., and Mizushima, N. (2020). ER-Phagy: Quality Control and Turnover of
1696   Endoplasmic Reticulum. Trends Cell Biol *30*, 384-398. 10.1016/j.tcb.2020.02.001.

1697   27.   Chino, H., and Mizushima, N. (2023). ER-Phagy: Quality and Quantity Control of the
1698   Endoplasmic Reticulum by Autophagy. Cold Spring Harb Perspect Biol *15*.

10.1101/cshperspect.a041256.

28.   Khaminets, A., Heinrich, T., Mari, M., Grumati, P., Huebner, A.K., Akutsu, M., Liebmann, L.,
Stolz, A., Nietzsche, S., Koch, N., et al. (2015). Regulation of endoplasmic reticulum
turnover by selective autophagy. Nature *522*, 354-358. 10.1038/nature14498.

29.   Grumati, P., Morozzi, G., Hölper, S., Mari, M., Harwardt, M.I.E., Yan, R., Müller, S., Reggiori,
F., Heilemann, M., and Dikic, I. (2017). Full length RTN3 regulates turnover of tubular
endoplasmic reticulum via selective autophagy. Elife *6*. 10.7554/eLife.25555.

30.   Rudinskiy, M., Galli, C., Raimondi, A., and Molinari, M. (2025). The intrinsically disordered
regions of organellophagy receptors are interchangeable and control organelle
fragmentation, ER-phagy and mitophagy flux. Nat Cell Biol *27*, 1431-1447.
10.1038/s41556-025-01728-4.

31.   Mo, J., Chen, J., and Zhang, B. (2020). Critical roles of FAM134B in ER-phagy and diseases.
Cell Death Dis *11*, 983. 10.1038/s41419-020-03195-1.

32.   Chino, H., Hatta, T., Natsume, T., and Mizushima, N. (2019). Intrinsically Disordered Protein
TEX264 Mediates ER-phagy. Mol Cell *74*, 909-921.e906. 10.1016/j.molcel.2019.03.033.

33.   Chen, Q., Xiao, Y., Chai, P., Zheng, P., Teng, J., and Chen, J. (2019). ATL3 Is a Tubular ER-
Phagy Receptor for GABARAP-Mediated Selective Autophagy. Curr Biol *29*, 846-855.e846.
10.1016/j.cub.2019.01.041.

34.   Mochida, K., and Nakatogawa, H. (2022). ER-phagy: selective autophagy of the
endoplasmic reticulum. EMBO Rep *23*, e55192. 10.15252/embr.202255192.

35.   Smith, M.D., Harley, M.E., Kemp, A.J., Wills, J., Lee, M., Arends, M., von Kriegsheim, A.,
Behrends, C., and Wilkinson, S. (2018). CCPG1 Is a Non-canonical Autophagy Cargo
Receptor Essential for ER-Phagy and Pancreatic ER Proteostasis. Dev Cell *44*, 217-
232.e211. 10.1016/j.devcel.2017.11.024.

36.   Pollard, T.D., Earnshaw, W. C., Lippincott-Schwartz, J., and Johnson, G. T. . (2016). Cell
Biology: Third Edition. (Elsevier).

37.   Klumperman, J., and Raposo, G. (2014). The complex ultrastructure of the endolysosomal
system. Cold Spring Harb Perspect Biol *6*, a016857. 10.1101/cshperspect.a016857.

38.   Germain, D.P. (2010). Fabry disease. Orphanet J Rare Dis *5*, 30. 10.1186/1750-1172-5-30.

39.   Wu, S.Z., Liang, X., Geng, J., Zhang, M.B., Xie, N., and Su, X.Y. (2019). Hydroxychloroquine-
induced renal phospholipidosis resembling Fabry disease in undifferentiated connective
tissue disease: A case report. World J Clin Cases *7*, 4377-4383. 10.12998/wjcc.v7.i24.4377.

40.   Raposo, G., Marks, M.S., and Cutler, D.F. (2007). Lysosome-related organelles: driving
post-Golgi compartments into specialisation. Curr Opin Cell Biol *19*, 394-401.
10.1016/j.ceb.2007.05.001.

41.   Faigle, W., Raposo, G., Tenza, D., Pinet, V., Vogt, A.B., Kropshofer, H., Fischer, A., de Saint-
Basile, G., and Amigorena, S. (1998). Deficient peptide loading and MHC class II
endosomal sorting in a human genetic immunodeficiency disease: the Chediak-Higashi
syndrome. J Cell Biol *141*, 1121-1134. 10.1083/jcb.141.5.1121.

42.   Platt, F.M., Boland, B., and van der Spoel, A.C. (2012). The cell biology of disease: lysosomal
storage disorders: the cellular impact of lysosomal dysfunction. J Cell Biol *199*, 723-734.
10.1083/jcb.201208152.

43.   Rozenszajn, L.A., Epstein, Y., and Fischer, D. (1969). Myelin figures in blood cells, lysosomes,
and lipidoses. Am J Clin Pathol *52*, 473-476. 10.1093/ajcp/52.4.473.

44.   Hurley, J.H., Coyne, A.N., Miączyńska, M., and Stenmark, H. (2025). The expanding
repertoire of ESCRT functions in cell biology and disease. Nature *642*, 877-888.
10.1038/s41586-025-08950-y.

45.   Schäfer, J.A., Schessner, J.P., Bircham, P.W., Tsuji, T., Funaya, C., Pajonk, O., Schaeff, K.,
Ruffini, G., Papagiannidis, D., Knop, M., et al. (2020). ESCRT machinery mediates selective
microautophagy of endoplasmic reticulum in yeast. Embo j *39*, e102586.
10.15252/embj.2019102586.

46.   Xu, F., Du, W., Zou, Q., Wang, Y., Zhang, X., Xing, X., Li, Y., Zhang, D., Wang, H., Zhang, W.,
et al. (2021). COPII mitigates ER stress by promoting formation of ER whorls. Cell Res *31*,
141-156. 10.1038/s41422-020-00416-2.

47.   Mizushima, N., Levine, B., Cuervo, A.M., and Klionsky, D.J. (2008). Autophagy fights disease
through cellular self-digestion. Nature *451*, 1069-1075. 10.1038/nature06639.

48.   Lam, S.S., Martell, J.D., Kamer, K.J., Deerinck, T.J., Ellisman, M.H., Mootha, V.K., and Ting,
A.Y. (2015). Directed evolution of APEX2 for electron microscopy and proximity labeling.
Nat Methods *12*, 51-54. 10.1038/nmeth.3179.

49.   Fumagalli, F., Noack, J., Bergmann, T.J., Cebollero, E., Pisoni, G.B., Fasana, E., Fregno, I.,
Galli, C., Loi, M., Soldà, T., et al. (2016). Translocon component Sec62 acts in endoplasmic
reticulum turnover during stress recovery. Nat Cell Biol *18*, 1173-1184. 10.1038/ncb3423.

50.   An, H., Ordureau, A., Paulo, J.A., Shoemaker, C.J., Denic, V., and Harper, J.W. (2019).
TEX264 Is an Endoplasmic Reticulum-Resident ATG8-Interacting Protein Critical for ER
Remodeling during Nutrient Stress. Mol Cell *74*, 891-908.e810.
10.1016/j.molcel.2019.03.034.

51.   Orsi, A., Razi, M., Dooley, H.C., Robinson, D., Weston, A.E., Collinson, L.M., and Tooze, S.A.
(2012). Dynamic and transient interactions of Atg9 with autophagosomes, but not
membrane integration, are required for autophagy. Mol Biol Cell *23*, 1860-1873.
10.1091/mbc.E11-09-0746.

52.   Judith, D., and Tooze, S.A. (2019). ATG9A supplies PtdIns4P to the autophagosome
initiation site. Autophagy *15*, 1660-1661. 10.1080/15548627.2019.1632124.

53.   Suzuki, E., Matsuda, T., Kawamoto, T., Takahashi, H., Mieda, Y., Matsuura, Y., Takai, T.,
Kanno, A., Koyanagi-Kimura, M., Asahara, S.I., et al. (2018). Docosahexaenoic Acid
Reduces Palmitic Acid-Induced Endoplasmic Reticulum Stress in Pancreatic B Cells. Kobe
J Med Sci *64*, E43-e55.

54.   Schuck, S., Prinz, W.A., Thorn, K.S., Voss, C., and Walter, P. (2009). Membrane expansion
alleviates endoplasmic reticulum stress independently of the unfolded protein response.
J Cell Biol *187*, 525-536. 10.1083/jcb.200907074.

55.   Fedele, A.O., and Proud, C.G. (2020). Chloroquine and bafilomycin A mimic lysosomal
storage disorders and impair mTORC1 signalling. Biosci Rep *40*. 10.1042/bsr20200905.

56.   Ruohola, H., Kabcenell, A.K., and Ferro-Novick, S. (1989). Reconstitution of transport from
the ER to the Golgi complex in yeast using microsomes and permeabilized yeast cells.
Methods Cell Biol *31*, 143-154. 10.1016/s0091-679x(08)61606-4.

57.   Baker, D., Hicke, L., Rexach, M., Schleyer, M., and Schekman, R. (1988). Reconstitution of
SEC gene product-dependent intercompartmental protein transport. Cell *54*, 335-344.
10.1016/0092-8674(88)90196-1.

58.   López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M., and Kroemer, G. (2023). Hallmarks

1787   of aging: An expanding universe. Cell *186*, 243-278. 10.1016/j.cell.2022.11.001.

59. Chang, B.D., Xuan, Y., Broude, E.V., Zhu, H., Schott, B., Fang, J., and Roninson, I.B. (1999). Role of p53 and p21waf1/cip1 in senescence-like terminal proliferation arrest induced in human tumor cells by chemotherapeutic drugs. Oncogene *18*, 4808-4818. 10.1038/sj.onc.1203078.

60. Chen, Q., and Ames, B.N. (1994). Senescence-like growth arrest induced by hydrogen peroxide in human diploid fibroblast F65 cells. Proc Natl Acad Sci U S A *91*, 4130-4134. 10.1073/pnas.91.10.4130.

61. Wang, X., Ma, L., Lu, D., Zhao, G., Ren, H., Lin, Q., Jia, M., Huang, F., Wang, S., Xu, Z., et al. (2024). Nuclear envelope budding inhibition slows down progerin-induced aging process. Proc Natl Acad Sci U S A *121*, e2321378121. 10.1073/pnas.2321378121.

62. Eriksson, M., Brown, W.T., Gordon, L.B., Glynn, M.W., Singer, J., Scott, L., Erdos, M.R., Robbins, C.M., Moses, T.Y., Berglund, P., et al. (2003). Recurrent de novo point mutations in lamin A cause Hutchinson-Gilford progeria syndrome. Nature *423*, 293-298. 10.1038/nature01629.

63. Sun, D., Wu, R., Zheng, J., Li, P., and Yu, L. (2018). Polyubiquitin chain-induced p62 phase separation drives autophagic cargo segregation. Cell Res *28*, 405-415. 10.1038/s41422-018-0017-7.

64. Sho, T., Li, Y., Jiao, H., and Yu, L. (2024). Migratory autolysosome disposal mitigates lysosome damage. J Cell Biol *223*. 10.1083/jcb.202403195.

65. Nguyen, T.N., Padman, B.S., Usher, J., Oorschot, V., Ramm, G., and Lazarou, M. (2016). Atg8 family LC3/GABARAP proteins are crucial for autophagosome-lysosome fusion but not autophagosome formation during PINK1/Parkin mitophagy and starvation. J Cell Biol *215*, 857-874. 10.1083/jcb.201607039.

66. Ren, J., Liang, R., Wang, W., Zhang, D., Yu, L., and Feng, W. (2020). Multi-site-mediated entwining of the linear WIR-motif around WIPI β-propellers for autophagy. Nat Commun *11*, 2702. 10.1038/s41467-020-16523-y.

67. Wagner, F.R., Watanabe, R., Schampers, R., Singh, D., Persoon, H., Schaffer, M., Fruhstorfer, P., Plitzko, J., and Villa, E. (2020). Preparing samples from whole cells using focused-ion-beam milling for cryo-electron tomography. Nat Protoc *15*, 2041-2070. 10.1038/s41596-020-0320-x.

68. Zheng, S.Q., Palovcak, E., Armache, J.P., Verba, K.A., Cheng, Y., and Agard, D.A. (2017). MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. Nat Methods *14*, 331-332. 10.1038/nmeth.4193.

69. Mastronarde, D.N., and Held, S.R. (2017). Automated tilt series alignment and tomographic reconstruction in IMOD. J Struct Biol *197*, 102-113. 10.1016/j.jsb.2016.07.011.

70. Edelman, B.L., and Redente, E.F. (2018). Isolation and Characterization of Mouse Fibroblasts. Methods Mol Biol *1809*, 59-67. 10.1007/978-1-4939-8570-8_5.

71. Correa-Gallegos, D., Ye, H., Dasgupta, B., Sardogan, A., Kadri, S., Kandi, R., Dai, R., Lin, Y., Kopplin, R., Shenai, D.S., et al. (2023). CD201(+) fascia progenitors choreograph injury repair. Nature *623*, 792-802. 10.1038/s41586-023-06725-x.

72. Xu, Z., Chen, D., Hu, Y., Jiang, K., Huang, H., Du, Y., Wu, W., Wang, J., Sui, J., Wang, W., et al. (2022). Anatomically distinct fibroblast subsets determine skin autoimmune patterns.

Nature *601*, 118-124. 10.1038/s41586-021-04221-8.

73. Charni-Natan, M., and Goldstein, I. (2020). Protocol for Primary Mouse Hepatocyte
Isolation. STAR Protoc *1*, 100086. 10.1016/j.xpro.2020.100086.

74. Rizo, J.A., Spencer, T.E., and Kelleher, A.M. (2024). Protocol for the establishment and
characterization of an endometrial-derived epithelial organoid and stromal cell co-
culture system. STAR Protoc *5*, 102894. 10.1016/j.xpro.2024.102894.

1837